



HAL
open science

Uranyl-chelating peptides to help understanding uranium toxicity at a molecular level

Fanny Laporte, Matthieu Stark, Colette Lebrun, Vicky Gathu, Pier Lorenzo Solari, Gaëlle Creff, Jérôme Roques, Christophe Den Auwer, Claude Vidaud, Pascale Delangle

► To cite this version:

Fanny Laporte, Matthieu Stark, Colette Lebrun, Vicky Gathu, Pier Lorenzo Solari, et al.. Uranyl-chelating peptides to help understanding uranium toxicity at a molecular level. *BIO Web of Conferences*, 2019, 14, pp.06005. 10.1051/bioconf/20191406005 . hal-02447953

HAL Id: hal-02447953

<https://hal.science/hal-02447953>

Submitted on 4 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Uranyl-chelating peptides to help understanding uranium toxicity at a molecular level

Fanny Laporte,¹ Matthieu Stark,¹ Colette Lebrun,¹ Stéphane Oros,¹ Nathalie Sisommay,¹ Vicky Gathu,¹ Pier Lorenzo Solari,⁵ Gaëlle Creff,³ Jérôme Roques,⁴ Christophe Den Auwer,³ Claude Vidaud² and Pascale Delangle^{1,*}

¹INAC-SyMMES, Univ. Grenoble Alpes, CEA, CNRS, 38000 Grenoble, France

²BIAM, CEA, CNRS, Aix-Marseille Université, 13108 Saint-Paul-lez-Durance, France

³Institut de Chimie de Nice, Université Côte d'Azur, 06108 Nice, France

⁴Institut de Physique Nucléaire d'Orsay, CNRS-IN2P3, Univ. Paris-Sud, Univ. Paris-Saclay, France


⁵Synchrotron SOLEIL, L'orme des Merisiers, Saint-Aubin, 91192 Gif-sur-Yvette, France

Context

Uranium is a natural element widely found in the environment, due to both natural occurrence in mineral ores or in sea water and industrial applications. The production of nuclear energy uses enriched uranium in ²³⁵U for nuclear fission. Despite its ubiquitous distribution, uranium has no essential role in living organisms and presents radiological and chemical toxicities. Despite significant recent advances in the field, there is still a serious lack of knowledge about the molecular interactions responsible for uranium toxicity. The underlying mechanisms need to be unravelled to predict the effect of uranium on living organisms and also to help in designing efficient detoxification agents, to be used in case of dirty bombs or accidental release of uranium in the environment.

Design of biomimetic peptides

At the CIBEST lab, we propose a biomimetic approach to help understanding the interactions of metals with biological molecules, relevant to their regulation and toxicity.[1, 2] This strategy was applied for uranium, which most stable form *in vivo* is the uranyl cation, UO₂²⁺, which prefers four to six oxygen donors in its equatorial plane, perpendicular to O-U-O bonds. Short peptide sequences are used as simple models of metal-binding sites in proteins to get structural and thermodynamic data, which are not accessible directly with large biological molecules. Cyclic constrained peptide scaffolds that orient metal-coordinating groups found in proteins toward the equatorial plane of the uranyl cation revealed to be highly effective uranyl-binding peptides.[3]



* Corresponding author: pascale.delangle@cea.fr

Results

A combination of synthesis, analytical chemistry, X-Absorption Spectroscopy and theoretical calculations allowed us to correlate the amino acid sequence to the stability of the uranyl complexes. Uranium-coordinating amino acids with side-chains containing oxygen donors, such as aspartic acids, glutamic acids and phosphoserines were inserted in the cyclic peptide sequences. The largest uranyl complexes stabilities were obtained when four coordinating amino acids were converging to the uranyl equatorial plane.[3-5] Importantly, phosphate groups significantly enhance the affinity of the biomimetic peptides for uranyl, which strongly supports the importance of phosphoamino acids in uranyl binding in proteins. Therefore, our biomimetic approach reinforces the relevance of considering phosphoproteins such as osteopontin as potential uranyl targets in vivo.[6-8]

Acknowledgements

This work was supported by the toxicology and NRBC programs of CEA and the LabEx ARCANÉ (Grant ANR-11-LABX-0003-01).

References

1. O. Sénèque, P. Rousselot-Pailley, A.M. Pujol, D. Boturyn, S. Crouzy, O. Proux, A. Manceau, C. Lebrun, P. Delangle, *Inorg. Chem.*, **57**, 2705 (2018)
2. A.S. Jullien, C. Gateau, C. Lebrun, P. Delangle, *Inorg. Chem.*, **54**, 2339 (2015)
3. C. Lebrun, M. Starck, V. Gathu, Y. Chenavier, P. Delangle, *Chem. Eur. J.*, **20**, 16566 (2014)
4. M. Starck, F. A. Laporte, S. Oros, N. Sisommay, V. Gathu, P. L. Solari, G. Creff, J. Roques, C. Den Auwer, C. Lebrun, P. Delangle, *Chem. Eur. J.*, **23**, 5288 (2017)
5. M. Starck, N. Sisommay, F. A. Laporte, S. Oros, C. Lebrun, P. Delangle, *Inorg. Chem.*, **24**, 11557 (2015)
6. L. Qi, C. Basset, O. Averseng, E. Quemeneur, A. Hagege, C. Vidaud, *Metallomics*, **6**, 166 (2014)
7. T. N. S. Huynh, C. Vidaud, A. Hagege, *Metallomics*, **8**, 1185 (2016).
8. F. Laporte, PhD thesis, Univ. Grenoble Alpes (2017).