



HAL
open science

Chebyshev's bias in dihedral and generalized quaternion Galois groups

Alexandre Bailleul

► **To cite this version:**

Alexandre Bailleul. Chebyshev's bias in dihedral and generalized quaternion Galois groups. 2020.
hal-02453465v1

HAL Id: hal-02453465

<https://hal.science/hal-02453465v1>

Preprint submitted on 23 Jan 2020 (v1), last revised 9 Jun 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chebyshev's bias in dihedral and generalized quaternion Galois groups

Alexandre Bailleul

Abstract

We study the inequities in the distribution of Frobenius elements in Galois extensions of the rational numbers with Galois groups that are either dihedral D_{2^n} or (generalized) quaternion \mathbb{H}_{2^n} of two-power order. In the spirit of recent work of Fiorilli and Jouve [5], we study, under natural hypotheses, some families of such extensions, in a horizontal aspect, where the degree is fixed, and in a vertical aspect, where the degree goes to infinity. Our main contribution uncovers in families of extensions a phenomenon, for which Ng gave numerical evidence in [18] : real zeros of Artin L -functions sometimes have a radical influence on the distribution of Frobenius elements.

Introduction

The prime number theorem in arithmetic progressions states that if $q \geq 1$ is an integer and a is prime to q , then

$$\pi(x, q, a) := |\{p \leq x \mid p = a \pmod{q}\}| \underset{x \rightarrow +\infty}{\sim} \frac{1}{\varphi(q)} \text{Li}(x),$$

where Li is the logarithmic integral. Even though for a and b coprime to q , $\pi(x, q, a)$ and $\pi(x, q, b)$ have the same asymptotic value, it may happen that one count could be larger than the other most of the time. This is Chebyshev's bias : there seems to be more primes congruent to $3 \pmod{4}$ than $1 \pmod{4}$ in generic intervals $[2, x]$.

These so-called "prime number races" have been studied extensively by Rubinstein and Sarnak in [21]. They managed to explain conditionally Chebyshev's bias : under natural hypotheses which we will consider later, the inequality $\pi(x, 4, 3) > \pi(x, 4, 1)$ holds "99% of the time" (this will be explained rigorously in a later paragraph). For a general modulus q , they have shown a bias towards non-quadratic residues mod q against quadratic residues mod q .

In [18], Ng, following a suggestion made in [21], extended Rubinstein and Sarnak's framework to conjugacy classes of the Galois group G of a Galois extension L/K of number fields, in the context of the Chebotarev density theorem. Recall that if C is such a conjugacy class, then

$$\pi(x, C, L/K) := \left| \left\{ \mathfrak{p} \triangleleft \mathcal{O}_K \text{ unramified} \mid N(\mathfrak{p}) \leq x, \left(\frac{\mathfrak{p}}{L/K} \right) = C \right\} \right| \underset{x \rightarrow +\infty}{\sim} \frac{|C|}{|G|} \text{Li}(x),$$

where $N(\mathfrak{p})$ is the norm of the prime ideal \mathfrak{p} and $\left(\frac{\mathfrak{p}}{L/K}\right)$ is the Artin symbol, or Frobenius conjugacy class, at \mathfrak{p} . Taking $L = \mathbb{Q}(\zeta_q)$, where ζ_q is a primitive q -th root of unity in \mathbb{C} , and $K = \mathbb{Q}$, we get the prime number theorem in arithmetic progressions and one recovers the setting of [21]. We say we study *Chebotarev biases* when comparing the behaviours of prime ideal counting functions $\pi(x, C_1, L/K)$ and $\pi(x, C_2, L/K)$ for distinct conjugacy classes C_1, C_2 of $\text{Gal}(L/K)$.

In recent work, Fiorilli and Jouve ([5]) have shown some Chebotarev races to be extremely biased (*i.e.* as for the original case where one compares $\pi(x, 4, 3)$ and $\pi(x, 4, 1)$, the underlying density is close to 1) by using large deviations principles. In the opposite direction, they managed to show central limit theorem behaviours which correspond to moderately biased Chebotarev races (*i.e.* the underlying density is close to $\frac{1}{2}$). In both cases, the asymptotic results appear as we let the conductors go to infinity. In [5], the theoretical results are applied to some families of number field extensions with Galois groups dihedral of order $2p$ (p an odd prime), quasidihedral, symmetric or Frobenius of order $p(p-1)$.

In this paper, we exploit work of Fröhlich on root numbers of quaternion extensions and we perform class field theoretic constructions to study, in the context of [5], Chebyshev's bias in some families of number field extensions with Galois groups of 2-power order : dihedral or (generalized) quaternion. Our main contributions highlight the role of central zeros of Artin L -functions in the study of this bias. They come as the result of the two following points of view. We first work in a "horizontal aspect", in which the Galois group is fixed (up to isomorphism). In this context, thanks to a result of Fröhlich (Theorem 3.3), we construct two different kinds of families of number fields which have quaternion Galois groups over \mathbb{Q} , but in which the existence or not of a central zero for the corresponding Dedekind zeta function leads to opposite biases (Theorem 3.5). Second, in the "vertical aspect", we build arbitrarily high degree towers of number field extensions with Galois groups dihedral or quaternion of 2-power order, and exhibit different kinds of behaviours in both families (Theorems 4.15 and 4.16). We obtain extreme bias, moderate bias and unbiased races (the underlying density is plainly equal to $\frac{1}{2}$). We emphasize that some of these behaviours are directly linked to the existence or not of a central zero for the corresponding Dedekind zeta functions in the quaternion case. This provides the first theoretical construction that confirms numerical evidence obtained by Ng. Finally, we are able to prove partial monotonicity phenomena in the evolution of the bias inside the tower itself (Theorem 4.17).

Outline of the paper

In Section 1 we introduce the preliminary results and conjectures needed to study Chebyshev's bias in number field extensions, and state abridged versions of our main results. In Section 2.1 we recall the definition of the Artin conductor of a complex character of the Galois group of a number field extension, and relate it to the study of Chebyshev's bias in the extension considered. Then in Section 2.2, we study the character-theoretic properties of dihedral and generalized quaternion groups of 2-power order. In Section 3 we focus on the "horizontal aspect" of our study, in which the Galois group is fixed to be a quaternion group of order 8. In Sections 4.1 and 4.2 we give bounds on moments of the random variables governing Chebyshev's bias in Galois extensions with Galois groups dihedral and generalized quaternion groups of 2-power order. In Section 4.3 we construct such extensions with

controlled ramification, and in Section 4.4 we give estimates on Chebyshev's bias in those extensions.

Acknowledgments

The author would like to thank Florent Jouve for his doctoral advising and for suggesting to study quaternion extensions of \mathbb{Q} in the context of Chebyshev's bias, Daniel Fiorilli for fruitful discussions on the subject, Philippe Cassou-Noguès for introducing him to the paper [6], which was the starting point of this work, Gerhard Niklasch for technical help in bounding discriminants in the proof of Proposition 4.9, and Alain Debreil for providing L^AT_EX code for the lattice of subgroups in Section 3.

1 Chebotarev biases in Galois groups of number fields

1.1 Notations and recollection on Artin characters

We recall the following standard notations.

In this section and sections 1.2 and 1.4, we let L/K be a Galois number field extension with Galois group G . Let C_1 and C_2 be distinct conjugacy classes of G . We would like to give a notion of measure to the set

$$\mathcal{P}_{L/K, C_1, C_2} := \{x \geq 2 \mid \pi(x, C_1, L/K) > \pi(x, C_2, L/K)\},$$

where C_1 and C_2 are conjugacy classes of G . A first guess would be to use the natural density, defined as the limit as x tends to $+\infty$ of

$$\frac{|\mathcal{P}_{L/K, C_1, C_2} \cap [0, x]|}{x},$$

where $|\cdot|$ denotes the Lebesgue measure. Unfortunately, it has been shown that this limit does not exist, even in the case of prime number races [12]. Therefore, we use the notion of logarithmic density, better-suited to the study of Chebyshev's bias as was first observed in [26].

Definition 1.1. *Let A be a Borel set in \mathbb{R} . Its logarithmic density is, when it exists,*

$$\delta(A) := \lim_{X \rightarrow +\infty} \frac{1}{X} \int_2^X \mathbf{1}_A(e^t) dt.$$

If C_1 and C_2 are conjugacy classes of G , we write $\delta(L/K, C_1, C_2)$ for $\delta(\mathcal{P}_{L/K, C_1, C_2})$. We say the race between C_1 and C_2 is unbiased if $\delta(L/K, C_1, C_2) = \frac{1}{2}$, that it is biased towards C_1 if $\delta(L/K, C_1, C_2) > \frac{1}{2}$ and biased towards C_2 if $\delta(L/K, C_1, C_2) < \frac{1}{2}$.

Ng has shown in [18] that this logarithmic density always exists under some natural hypotheses (see section 1.3). Our goal will be to give estimates for such densities. Following [5], we introduce some quantities related to the conjugacy classes under consideration, but also to the irreducible characters of G .

Definition 1.2. Let C be a conjugacy class of G . Then we define

$$C^{1/2} := \{g \in G \mid g^2 \in C\}$$

and

$$z(C) := 2 \sum_{\chi \neq \chi_0} \chi(C) \operatorname{ord}_{s=1/2} L(s, \chi, L/K),$$

where the sum is taken over all the non trivial irreducible characters χ of G , and $s \mapsto L(s, \chi, L/K)$ is the Artin L -function associated to χ .

A certain class of characters of G plays a particular role in our study, the class of symplectic characters. We introduce them in the following definition.

Definition 1.3. Let χ be a character of G . We define its Frobenius-Schur index by

$$\varepsilon_2(\chi) := \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$

We also define $\operatorname{Irr}(G)$ to be the set of irreducible characters of G , and $\operatorname{Irr}^{\operatorname{real}}(G)$ to be the set of real-valued irreducible characters of G .

The Frobenius-Schur index of an irreducible (complex) character of G determines if such a character can be afforded by a real-valued representation or not, thanks to the following well-known result from character theory.

Theorem 1.4 ([9] p.58). Let χ be an irreducible complex character of G . Then only one of the following three statements holds :

- i) $\varepsilon_2(\chi) = 0$, in this case χ is not real-valued. We say that χ is unitary.
- ii) $\varepsilon_2(\chi) = 1$, in this case χ is real-valued and can be afforded by a real-valued representation of G . We say that χ is orthogonal.
- iii) $\varepsilon_2(\chi) = -1$, in this case χ is real-valued and cannot be afforded by a real-valued representation of G . We say that χ is symplectic.

It is expected that symplectic characters are exactly the irreducible characters which can yield real zeros for their associated Artin L -function (see conjecture LI below).

Finally, we recall that if χ is an irreducible character of G , then its Artin L -function satisfies a functional equation ([16, p.28]) of the form

$$\Lambda(s, \chi, L/K) = W(\chi, L/K) \Lambda(1-s, \bar{\chi}, L/K),$$

where $W(\chi, L/K)$ is a complex number of modulus 1, called the root number of χ , and $s \mapsto \Lambda(s, \chi, L/K)$ is the completed Artin L -function associated to χ , which is the product of $s \mapsto L(s, \chi, L/K)$ with Gamma factors coming from archimedean places. Unless there is an ambiguity in the extension considered, we will usually write $W(\chi)$ for $W(\chi, L/K)$. The root number satisfies $\overline{W(\chi)} = W(\bar{\chi})$. In particular if χ is real-valued then $W(\chi) = \pm 1$. We have the following important way to detect central zeros of Artin L -functions.

Proposition 1.5. *If $\chi \in \text{Irr}^{\text{real}}(G)$ with $W(\chi) = -1$, then $L\left(\frac{1}{2}, \chi, L/K\right) = 0$.*

Proof. Since χ is real-valued, evaluating the functional equation at $\frac{1}{2}$ we find

$$\Lambda\left(\frac{1}{2}, \chi\right) = -\Lambda\left(\frac{1}{2}, \chi\right),$$

i.e. $\Lambda\left(\frac{1}{2}, \chi\right) = 0$. Since the Gamma function never vanishes on \mathbb{C} , this implies that $L\left(\frac{1}{2}, \chi, L/K\right) = 0$. \square

It is expected that the converse also holds for real-valued characters when the base field is \mathbb{Q} , see conjecture LI⁺ below.

1.2 Conjectures

As in [5], we consider natural conjectures on the distribution of zeros and poles of Artin L -functions, some of which are generalizations of conjectures used in the work of Rubinstein and Sarnak. Recall that L/K is a Galois extension of number fields.

Conjecture 1.6 (Artin's conjecture). *If χ is a non-trivial irreducible character of G , then $s \mapsto L(s, \chi, L/K)$ is entire.*

Conjecture 1.7 (GRH). *If χ is an irreducible character of G , then the non-trivial zeros of $s \mapsto L(s, \chi, L/K)$ have real part $\frac{1}{2}$.*

Conjecture 1.8 (LI⁻). *Let L_0/\mathbb{Q} be the Galois closure of L/\mathbb{Q} . Then the multiset of imaginary parts of zeros*

$$\Gamma_{L_0/\mathbb{Q}} := \bigcup_{\chi \in \text{Irr}(\text{Gal}(L_0/\mathbb{Q}))} \left\{ \gamma > 0 \mid L\left(\frac{1}{2} + i\gamma, \chi, L_0/\mathbb{Q}\right) = 0 \right\}$$

is linearly independent over \mathbb{Q} .

Conjecture 1.9 (LI). *LI⁻ is true and if $\chi \neq \chi_0$ is a unitary or orthogonal character of $\text{Gal}(L_0/\mathbb{Q})$ (see Theorem 1.4) then $L\left(\frac{1}{2}, \chi, L_0/\mathbb{Q}\right) \neq 0$. If χ is a symplectic character of $\text{Gal}(L_0/\mathbb{Q})$ then $\text{ord}_{s=1/2} L(s, \chi, L_0/\mathbb{Q})$ is bounded by some absolute constant M_0 .*

Conjecture 1.10 (LI⁺). *LI is true and if χ is a symplectic irreducible character of $\text{Gal}(L_0/\mathbb{Q})$ then $\text{ord}_{s=1/2} L(s, \chi, L_0/\mathbb{Q}) = \frac{1-W(\chi)}{2}$.*

Let us make a few comments about these conjectures.

- First, Artin's conjecture needs to be assumed in order to be able to prove explicit formulas for number field analogs of Chebyshev's ψ function, involving sums over zeros of Artin L -functions, and discarding the existence of possible poles. In the cases we will consider, namely when G is a dihedral group or generalized quaternion group, Artin's conjecture is known to be true, because such groups are supersolvable, which means they have a normal series

$$\{1\} = G_1 \triangleleft G_2 \triangleleft \dots \triangleleft G_k = G$$

where each quotient G_{i+1}/G_i is cyclic for $1 \leq i \leq k-1$. This implies that their irreducible characters are induced by those of abelian subgroups. By inductive properties of Artin L -functions, we are reduced to knowing Artin's conjecture in the case of characters of abelian Galois groups, but this is exactly one of the consequences of class field theory, together with work of Hecke on the L -functions bearing his name.

- The Generalized Riemann Hypothesis (GRH) needs to be assumed so that, for conjugacy classes C_1 and C_2 of G ,

$$\frac{|C_1|}{|G|} \pi(x, C_1, L/K) - \frac{|C_2|}{|G|} \pi(x, C_2, L/K)$$

is oscillating with amplitude of size roughly \sqrt{x} . This is central in the analysis of Chebyshev's bias. One can still show the existence of a limiting logarithmic distribution for a convenient renormalization of $\frac{|C_1|}{|G|} \pi(x, C_1, L/K) - \frac{|C_2|}{|G|} \pi(x, C_2, L/K)$ without assuming GRH (see [4]), though it depends on the supremum of real parts of non-trivial zeros of the Artin L -functions considered.

- The hypothesis LI⁺ actually contains two statements. The most obvious one is the linear independence of the positive imaginary parts of zeros of Artin L -functions. This hypothesis appears because, in order to understand Chebyshev's bias using explicit formulas for prime counting functions, one needs good information on the joint distribution of the values of $e^{i\gamma_1 y}, e^{i\gamma_2 y}, \dots$ in \mathbb{S}^1 , where $\gamma_1, \gamma_2, \dots$ denotes the aforementioned positive imaginary parts of zeros, and y varies in \mathbb{R} . Again, weaker hypotheses have been used to show the existence of a limiting logarithmic distribution ([4]) and results on the bias. Another aspect of LI we highlight is that it is stated for L -functions over \mathbb{Q} instead of L -functions over K . Indeed, those L -functions over K factorize as products of L -functions relative to L_0/\mathbb{Q} , so linear independence is typically false. The last aspect addressed by LI is about the multiplicity of the zeros of Artin L -functions over \mathbb{Q} . One should expect those Artin L -functions to be "primitive" in the sense of [22], and those should not satisfy any kind of non-trivial algebraic relations, except for their respective functional equations. Instead of the simplicity of the zeros, we could have opted for an hypothesis about the boundedness of the multiplicities of such zeros (called BM in [5]), which would have been enough for a few intermediate results of this paper.
- The assumptions on the order of vanishing at $\frac{1}{2}$ in LI and LI⁺ appear because the quantity $z(C)$ defined above is involved in the mean of the random variables attached to the limiting distribution governing the bias. Examples of symplectic characters with root numbers -1 were first given by Armitage ([1]) and Serre (unpublished). In particular, they yield Artin L -functions vanishing at $1/2$ because of Proposition 1.5. As was pointed out by the author of the present paper to the authors of [5] while both papers were under preliminary form, using one of the aforementioned examples, the conjecture "Artin L -functions attached to orthogonal or unitary characters do not vanish at $1/2$ " could not hold over general number fields K . Indeed, consider Serre's example, as described in [18, section 5.3.3] : let $L = \mathbb{Q}(\theta)$, where $\theta = \sqrt{\frac{5+\sqrt{5}}{2} \frac{41+\sqrt{5 \cdot 41}}{2}}$. It is a Galois extension of \mathbb{Q} with Galois group isomorphic to the quaternion group

\mathbb{H}_8 (see section 2). The root number of its non-abelian (symplectic) character ψ is -1 , so $L(1/2, \psi, L/\mathbb{Q}) = 0$. Now consider the subfield $K = \mathbb{Q}(\sqrt{5}) \subset L$. We have $\text{Gal}(L/K) \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ which does not admit any irreducible symplectic character. But we have the classical factorization

$$\begin{aligned}\zeta_L(s) &= P_1(s)L(s, \psi, L/\mathbb{Q})^2 \\ &= P_2(s),\end{aligned}$$

where P_1 is the product of the Artin L -functions attached to the four abelian irreducible characters of $\text{Gal}(L/\mathbb{Q})$, and P_2 is the product of the Artin L -functions attached to the four irreducible characters of $\text{Gal}(L/K)$. Since $L(1/2, \psi, L/\mathbb{Q}) = 0$, at least one of those Artin L -functions, attached to a non-symplectic irreducible character, must vanish at $1/2$. This shows that we cannot expect easy non-vanishing statements at the central point in the relative case L/K when $K \neq \mathbb{Q}$, and that such a statement should involve the way irreducible characters of G are "induced" to Artin characters over \mathbb{Q} .

1.3 Main results

Our first result is in a horizontal aspect : we exhibit moderately biased races in families of extensions of \mathbb{Q} with fixed Galois group isomorphic to \mathbb{H}_8 , the usual quaternion group of order 8.

Theorem A. *Assume GRH and LI⁺. For any function f such that $f(n) \xrightarrow[n \rightarrow +\infty]{} +\infty$, there exist two families $(K_d)_d$ and $(L_d)_d$ of number fields, indexed by square-free integers satisfying $d > 1$ and $d \equiv 1 \pmod{4}$, such that for any d in the index set :*

- i) $\mathbb{Q}(\sqrt{d}) \subset K_d \cap L_d$.
- ii) $\text{Gal}(K_d/\mathbb{Q}) \simeq \text{Gal}(L_d/\mathbb{Q}) \simeq \mathbb{H}_8$.
- iii) $0 < \frac{1}{2} - \delta(K_d/\mathbb{Q}, C_1, C_{-1}) \ll \frac{1}{f(d)}$, where C_1 and C_{-1} denote the conjugacy classes of 1 and -1 in \mathbb{H}_8 .
- iv) $0 < \delta(L_d/\mathbb{Q}, C_1, C_{-1}) - \frac{1}{2} \ll \frac{1}{f(d)}$.

The most important feature of this result is that we are able to exhibit two families in which the biases are opposite to each other, one is $< \frac{1}{2}$ while the other is $> \frac{1}{2}$, the difference coming from the existence of a central zero for the corresponding Dedekind zeta function in the first case, and the absence of such a zero in the second case. Moreover, the prime number races considered can be specified to be as moderately biased as possible, in the sense that the logarithmic densities can be made arbitrarily close to $\frac{1}{2}$ when d grows, while ensuring that $\mathbb{Q}(\sqrt{d}) \subset K_d \cap L_d$. In fact the choice of f allows one to have arbitrarily large variances for the random variables $X(K_d/\mathbb{Q}, C_1, C_{-1})$ and $X(L_d/\mathbb{Q}, C_1, C_{-1})$ governing the biases (see Theorem 1.11).

The second part of our work is in a vertical aspect : we build families of extensions of \mathbb{Q} with Galois groups dihedral and quaternion of 2-power order, in which we are able to show extreme biases, moderate biases and absence of bias.

Theorem B. *Assume GRH and LI⁺. There exist absolute constants $c_1, c_2, c_3 > 0$ and a family $(\mathcal{D}_n)_n$ of number fields such that for any $n \geq 3$, the following hold :*

- i) $\text{Gal}(\mathcal{D}_n/\mathbb{Q}) \simeq D_{2^{n-1}} = \langle r, s \mid r^{2^{n-1}} = s^2 = 1, srs^{-1} = r^{-1} \rangle$.
- ii) $c_1 \exp(-c_2 2^n) < \delta(\mathcal{D}_n/\mathbb{Q}, C_1, C_{-1}) < \exp\left(-c_3 \frac{2^n}{n}\right)$ where C_a denotes the conjugacy class of a , and -1 denotes $r^{2^{n-2}} \in D_{2^{n-1}}$.
- iii) $0 < \frac{1}{2} - \delta(\mathcal{D}_n/\mathbb{Q}, C_{-1}, C_s) \ll \frac{1}{2^{n/3}}$.
- iv) $\delta(\mathcal{D}_n/\mathbb{Q}, C_{r^k}, C_s) = \delta(\mathcal{D}_n/\mathbb{Q}, C_{r^k}, C_{rs}) = \frac{1}{2}$ when $1 \leq k \leq 2^{n-2} - 1$ is odd.

We chose a sample of possible couples of conjugacy classes in the above statement. For an exhaustive treatment of conjugacy classes, see Theorem 4.15. In [5], Fiorilli and Jouve, relying on a construction of Klüners, only deal with dihedral groups of order $2p$ with p varying in the set of odd primes. The following result deals with generalized quaternion Galois groups.

Theorem C. *Assume GRH and LI⁺. There exist absolute constants $c_1, c_2, c_3 > 0$ and two families $(\mathcal{Q}_n^+)_n$ and $(\mathcal{Q}_n^-)_n$ of number fields such that for any $n \geq 3$, the following hold :*

- i) $\text{Gal}(\mathcal{Q}_n^\pm/\mathbb{Q}) \simeq \mathbb{H}_{2^n} = \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle$, and the root number of each symplectic character of $\text{Gal}(\mathcal{Q}_n^\pm/\mathbb{Q})$ is the same (we denote it by $W_{\mathcal{Q}_n^\pm}$). Moreover, $W_{\mathcal{Q}_n^+} = 1$ and $W_{\mathcal{Q}_n^-} = -1$.
- ii) $c_1 \exp(-c_2 2^n) < \left| \frac{1+W_{\mathcal{Q}_n^\pm}}{2} - \delta(\mathcal{Q}_n^\pm/\mathbb{Q}, C_1, C_{-1}) \right| < \exp\left(-c_3 \frac{2^n}{n}\right)$, where C_a denotes the conjugacy class of a , and -1 denotes $x^{2^{n-2}}$.
- iii) $\delta(\mathcal{Q}_n^+/\mathbb{Q}, C_1, C_{x^k}) = \frac{1}{2}$ when $1 \leq k \leq 2^{n-2} - 1$ is even and $c_1 \exp(-c_2 8^n) < \delta(\mathcal{Q}_n^-/\mathbb{Q}, C_1, C_{x^k}) < \exp\left(-c_3 \frac{2^n}{n}\right)$ for $1 \leq k \leq 2^{n-2} - 1$.
- iv) $0 < \frac{1}{2} - \delta(\mathcal{Q}_n^+/\mathbb{Q}, C_1, C_y) \ll \frac{1}{2^{n/3}}$ and $c_1 \exp(-c_2 2^n) < \delta(\mathcal{Q}_n^-/\mathbb{Q}, C_1, C_y) < \exp\left(-c_3 \frac{2^n}{n}\right)$.

Again, we chose to focus on a sample of meaningful cases of bias estimates in ii), iii) and iv), but more have been computed and are stated in Theorem 4.16. The new and remarkable feature of this theorem is that it highlights the role played by the existence or not of a central zero for the corresponding Dedekind zeta function. The presence (or the inexistence) of such a zero leads to : extreme biases of opposite signs in ii), extreme biases or no biases in iii), extreme bias or moderate bias in iv).

Finally, our last contribution is to show that we can observe a monotonicity phenomenon in the evolution of Chebyshev's bias in subextensions of dihedral extensions of \mathbb{Q} .

Theorem D. *Assume GRH and LI⁺. Consider the number fields $(\mathcal{D}_n)_n$ as in Theorem B. For every $n \geq 3$, there are dihedral subfields $\mathcal{D}_n^{(i)} \subset \mathcal{D}_n$, $3 \leq i \leq n$, decreasing for inclusion*

with i , such that for any $\varepsilon > 0$ and any sufficiently large n , for $3 \leq i < j \leq n$ such that $i \leq n^{\frac{1+\varepsilon}{2}}$ and $j \geq n^{\frac{1+3\varepsilon}{2}}$, we have

$$\delta(\mathcal{D}_n/\mathcal{D}_n^{(j)}, C_1^{(j)}, C_{-1}^{(j)}) < \delta(\mathcal{D}_n/\mathcal{D}_n^{(i)}, C_1^{(i)}, C_{-1}^{(i)}),$$

where $C_1^{(k)}$ and $C_{-1}^{(k)}$ denote the conjugacy classes of 1 and -1 in $\text{Gal}(\mathcal{D}_n/\mathcal{D}_n^{(k)})$.

This statement means that, in the subextensions of $\mathcal{D}_n/\mathcal{D}_n^{(i)}$, Chebyshev's bias is more extreme at the bottom of the tower than at the top. A more general result including the quaternion fields \mathcal{Q}_n^\pm of Theorem C is stated as Theorem 4.17. The proof relies on a new large deviation bound for the values of Chebyshev's bias in families of number field extensions (see Theorem 4.14).

1.4 The probabilistic approach to Chebotarev biases

Let us recall the setting of [5] together with useful results. As before, L/K is a Galois extension of number fields with group G . We will assume L/\mathbb{Q} is Galois, with Galois group G^+ , so that G is a subgroup of G^+ . This will be the case in our applications. If C is a conjugacy class of G , then we denote by C^+ the conjugacy class generated by C in G^+ , *i.e.* $C^+ := \bigcup_{a \in G^+} aCa^{-1}$. In what follows, if F/E is a Galois extension of number fields and $\chi \in \text{Irr}(\text{Gal}(F/E))$, a summation over $\gamma_\chi > 0$ means a summation over the corresponding zero multiset

$$\Gamma_{F/E, \chi} := \{\gamma > 0 \mid L(1/2 + i\gamma, \chi, F/E) = 0\}.$$

We also write $\Gamma_{F/E} := \bigcup_{\chi \in \text{Irr}(\text{Gal}(F/E))} \Gamma_{F/E, \chi}$ and $\Gamma_{F/E}^{\text{real}} := \bigcup_{\chi \in \text{Irr}^{\text{real}}(\text{Gal}(F/E))} \Gamma_{F/E, \chi}$.

We summarize in the next statement the main result giving a probabilistic interpretation of the logarithmic densities we are studying.

Theorem 1.11 ([5], Proposition 3.18 and Lemma 3.20). *Assume Artin's conjecture, GRH and LI. For any $\gamma \in \Gamma_{L/\mathbb{Q}}$, we introduce the random variable $X_\gamma = \Re(Z_\gamma)$, where $(Z_\gamma)_\gamma$ is a family of independent random variables uniform on the unit circle. Then for any conjugacy classes C_1 and C_2 of G , we have*

$$\delta(L/K, C_1, C_2) = \mathbb{P}(X(L/K, C_1, C_2) > 0)$$

where

$$X(L/K, C_1, C_2) := \frac{|C_2^{1/2}|}{|C_2|} - \frac{|C_1^{1/2}|}{|C_1|} + z(C_2) - z(C_1) + 2 \sum_{\lambda \in \text{Irr}(G^+)} |\lambda(C_2^+) - \lambda(C_1^+)| \sum_{\gamma_\lambda > 0} \frac{X_{\gamma_\lambda}}{\sqrt{\frac{1}{4} + \gamma_\lambda^2}},$$

unless $C_1^+ = C_2^+$.

Moreover, under the previous condition, we have

$$\mathbb{E}(X(L/K, C_1, C_2)) = \frac{|C_2^{1/2}|}{|C_2|} - \frac{|C_1^{1/2}|}{|C_1|} + z(C_2) - z(C_1)$$

and

$$\text{Var}(X(L/K, C_1, C_2)) = 2 \sum_{\lambda \in \text{Irr}(G^+)} |\lambda(C_1^+) - \lambda(C_2^+)|^2 \sum_{\gamma_\lambda > 0} \frac{1}{\frac{1}{4} + \gamma_\lambda^2}.$$

Theorem 1.11 is not the exact transcription of [5, Proposition 3.18], but it is the same, specialized to the case of the class function $\frac{|G|}{|C_1|}\mathbf{1}_{C_1} - \frac{|G|}{|C_2|}\mathbf{1}_{C_2}$, under the hypothesis LI^+ : each zero in the sum giving $\text{Var}(X(L/K, C_1, C_2))$ is a zero of exactly one function $s \mapsto L(s, \chi, L/\mathbb{Q})$, with multiplicity one. In other words, the union

$$\Gamma_{L/\mathbb{Q}} = \bigcup_{\chi \in \text{Irr}(G^+)} \Gamma_{L/\mathbb{Q}, \chi}$$

is a disjoint union, and each of its elements appears only once in the sums indexed by $\lambda \in \text{Irr}(G^+)$. Note that the expression of $\text{Var}(X(L/K, C_1, C_2))$ involves zeros of L -functions attached to characters of G^+ , not of G . This is because our linear independence and simplicity hypothesis LI^+ was stated over \mathbb{Q} . If $C_1^+ = C_2^+$ then it is easy to see that $\frac{|C_2^{1/2}|}{|C_2|} - \frac{|C_1^{1/2}|}{|C_1|} + z(C_2) - z(C_1) = 0$, since C_1 and C_2 are conjugated in G^+ , and the approach to Chebyshev's bias initiated by Rubinstein and Sarnak [21] and generalized by Ng [18], breaks down. In fact, one can show that $\pi(x, C_1, L/K) = \pi(x, C_2, L/K)$ for any $x \geq 2$ in this case.

The random variables at play are symmetric about their mean m . This is because their characteristic functions are products of Bessel J_0 functions, which are even, multiplied by $t \mapsto e^{imt}$ ([18, Theorem 5.2.1]). Therefore, it is easily seen that the corresponding logarithmic density δ will be $< \frac{1}{2}$, $> \frac{1}{2}$ or $= \frac{1}{2}$ according to whether $m < 0$, $m > 0$ or $m = 0$, respectively. They also do not have any atoms, as was shown in [4, Theorem 2.2].

We introduce one more quantity in order to state the main results of [5] on Chebyshev's bias in families of number fields.

Definition 1.12. *Let C_1 and C_2 be conjugacy classes of G such that $C_1^+ \neq C_2^+$. Assuming Artin's conjecture, GRH and LI^+ , the bias factor of the race between C_1 and C_2 is*

$$B(L/K, C_1, C_2) := \frac{\mathbb{E}(X(L/K, C_1, C_2))}{\sqrt{\text{Var}(X(L/K, C_1, C_2))}}.$$

Note that the above variance is non-zero since $C_1^+ \neq C_2^+$.

The next theorem is a result giving extremely biased races.

Theorem 1.13 ([5, Proposition 5.3]). *Assume GRH, LI^+ and Artin's Conjecture. Then there exists an absolute constant $c_3 > 0$ such that for any conjugacy classes C_1, C_2 of G satisfying $C_1^+ \neq C_2^+$ and $B(L/K, C_1, C_2) > 0$, we have*

$$1 - \delta(L/K, C_1, C_2) < \exp(-c_3 B(L/K, C_1, C_2)^2).$$

Remark. This theorem is based on a large deviation result due to Montgomery and Odlyzko (see Theorem 4.13 below). The idea is that if the quantity B is positive and large, then the mean of the associated random variable X is large compared to its variance, and the distribution of the random variable X is very concentrated around its (positive) mean, and therefore

it takes positive values with high probability.

We are actually able to provide lower bounds in the context of Theorem 1.13, but not as uniform as the previous upper bounds (see Theorem 4.14). This new bound will allow us to exhibit a monotonicity phenomenon in the values of Chebyshev's bias in certain towers of number field extensions (see Theorem [?Mono]).

The second result can be seen as a converse to the above theorem. We can roughly state it as saying that if the quantity B is small, then the race will be moderately biased, *i.e.* the logarithmic density δ will be close to $\frac{1}{2}$.

Theorem 1.14 ([5], Theorem 5.10). *Assume GRH, LI⁻ and Artin's Conjecture. Then for any conjugacy classes C_1, C_2 of G such that $C_1^+ \neq C_2^+$, if $|B(L/K, C_1, C_2)|$ is small enough then we have*

$$\delta(L/K, C_1, C_2) = \frac{1}{2} + \frac{B(L/K, C_1, C_2)}{\sqrt{2\pi}} + O\left(B(L/K, C_1, C_2)^3 + \text{Var}(X(L/K, C_1, C_2))^{-1/3}\right).$$

Remarks.

- i) This result is a consequence of a central limit behaviour for the random variable $X(L/K, C_1, C_2)$ (explaining the $\sqrt{2\pi}$ factor), provided its variance is large enough, which can be established thanks to bounds on Bessel J_0 functions which appear in the characteristic functions of $X(L/K, C_1, C_2)$.
- ii) If $\mathbb{E}(X(L/K, C_1, C_2))$ is bounded and $B(L/K, C_1, C_2)$ approaches zero then the main term besides $\frac{1}{2}$ is $\text{Var}(X(L/K, C_1, C_2))^{-1/3}$.

2 Recollection on ramification and on dihedral and quaternion groups of 2-power order

2.1 Artin conductors

Let L/K be a Galois extension of number fields with Galois group G and C_1, C_2 be distinct conjugacy classes of G . In [5], bounds for the variance of $X(L/K, C_1, C_2)$ are given in terms of the local ramification data of the extension L/K . More precisely, a link between this variance and the Artin conductors of the irreducible characters of G is established.

Recall that if \mathfrak{p} is a prime ideal of \mathcal{O}_K and \mathfrak{P} is a prime ideal in \mathcal{O}_L above \mathfrak{p} , the inertia subgroup $\mathcal{I}(\mathfrak{P}/\mathfrak{p}) := \{g \in G \mid \forall x \in \mathcal{O}_L, g(x) = x \bmod \mathfrak{P}\} \subset G$ admits a filtration $(G_i(\mathfrak{P}/\mathfrak{p}))_{i \geq 0}$ defined as follows : for integers $i \geq 0$, define

$$G_i(\mathfrak{P}/\mathfrak{p}) := \{g \in G \mid \forall x \in \mathcal{O}_L, g(x) = x \bmod \mathfrak{P}^{i+1}\} \subset G.$$

Obviously, $G_0(\mathfrak{P}/\mathfrak{p}) = \mathcal{I}(\mathfrak{P}/\mathfrak{p})$ and the subgroups $G_i(\mathfrak{P}, \mathfrak{p})$ of G only depend on \mathfrak{p} up to conjugacy, so in the sequel we drop the dependency on \mathfrak{P} and denote them by $G_i(\mathfrak{p})$. It is also known that the elements of this filtration are eventually trivial. We note that if L/K

is tamely ramified, then $G_i(\mathfrak{p})$ is trivial for $i \geq 1$ ([23, Chapitre IV §2 Corollaire 3]). If $\rho : G \rightarrow \mathbf{GL}(V)$ is a complex representation, we obtain complex representations of all the ramification subgroups $G_i(\mathfrak{p})$ by restriction. If χ is the character ρ , we define

$$n(\chi, \mathfrak{p}) := \sum_{i \geq 0} \frac{|G_i(\mathfrak{p})|}{|G_0(\mathfrak{p})|} \operatorname{codim}(V^{G_i(\mathfrak{p})}),$$

which only depends on \mathfrak{p} and not on \mathfrak{P} since the $G_i(\mathfrak{P}/\mathfrak{p})$ are conjugates in G and so the dimension of their invariant subspaces are all the same. This sum is finite since the $G_i(\mathfrak{p})$ are eventually trivial, but it is also known that this sum is an integer ([23, Chapitre VI §2 Corollaire 2]), so we can finally define the Artin conductor of the character χ by

$$\mathfrak{f}(L/K, \chi) := \prod_{\mathfrak{p}} \mathfrak{p}^{n(\chi, \mathfrak{p})}.$$

This ideal of \mathcal{O}_K is well-defined since it is easy to see that for \mathfrak{p} unramified in L we have $n(\chi, \mathfrak{p}) = 0$, so that there are only finitely many prime ideals actually contributing to the product. We recall the important conductor-discriminant formula ([23, Chapitre VI §3 Corollaire 2]).

Theorem 2.1. *We have*

$$D_{L/K} = \prod_{\chi \in \operatorname{Irr}(G)} \mathfrak{f}(L/K, \chi)^{\chi(1)},$$

where $D_{L/K}$ is the relative discriminant of L/K .

The following quantity appears in the functional equation of the Artin L -functions associated to a character χ .

Definition 2.2 ([16] p.28). *For any irreducible character χ of G , define*

$$A(\chi) := |d_K|^{\chi(1)} N_{K/\mathbb{Q}}(\mathfrak{f}(L/K, \chi)),$$

where d_K is the absolute discriminant of K .

Lemma 2.3 ([5], Lemma 4.3). *Assume Artin's conjecture. Then for any irreducible character χ of G , we have*

$$B_0(\chi) := \sum_{\gamma_\chi \neq 0} \frac{1}{\frac{1}{4} + \gamma_\chi^2} \asymp \log A(\chi).$$

Recall that the sums $B_0(\chi)$, for non-trivial irreducible characters χ of G , appear in the variances in Theorem 1.11. That explains why we will be interested in bounding the quantities $\log A(\chi)$.

2.2 Character theory of dihedral and quaternion groups of 2-power order

We obtain our main contributions in the setting of Galois extensions of number fields with Galois groups dihedral or quaternion of 2-power order. The goal of this section is to recollect some character-theoretic facts about these groups which will be used throughout sections 3 and 4. Let us first recall the classical presentations of those groups.

Definition 2.4. Let $n \geq 3$ be an integer. The dihedral group of order 2^n is

$$D_{2^{n-1}} := \langle r, s \mid r^{2^{n-1}} = s^2 = 1, srs^{-1} = r^{-1} \rangle.$$

The generalized quaternion group of order 2^n is

$$\mathbb{H}_{2^n} := \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle.$$

The group \mathbb{H}_8 is the usual quaternion group of order 8, with elements usually denoted i, j, k satisfying

$$i^2 = j^2 = k^2 = ijk = -1, (-1)^2 = 1.$$

This notation will be used in section 3. In $D_{2^{n-1}}$, we will denote by -1 the element $r^{2^{n-2}}$, and in \mathbb{H}_{2^n} we will also denote by -1 the element $x^{2^{n-2}} = y^2$. Both of those elements are of order 2 and generate the center of the respective group they belong to.

These groups are special instances of metacyclic groups (*i.e.* groups admitting a cyclic normal subgroup with cyclic quotient). As a consequence, computations are easily carried in such groups and the conjugacy classes are easily identified ([24]).

Lemma 2.5. Let $n \geq 3$ be an integer. Then $D_{2^{n-1}} = \langle r, s \mid r^{2^{n-1}} = s^2 = 1, srs^{-1} = r^{-1} \rangle$ has the following $2^{n-2} + 3$ conjugacy classes :

- i) The trivial conjugacy class $\{1\}$, denoted by C_1 .
- ii) The conjugacy class $\{-1\}$, denoted by C_{-1} .
- iii) Pairs of powers $\{r^k, r^{-k}\}$ for $1 \leq k \leq 2^{n-2} - 1$, denoted by $C_{r,k}$.
- iv) The conjugacy class $\{r^k s \mid 0 \leq k \leq 2^{n-2} - 1 \text{ even}\}$ of s , denoted by C_s .
- v) The conjugacy class $\{r^k s \mid 0 \leq k \leq 2^{n-1} - 1 \text{ odd}\}$ of rs , denoted by C_{rs} .

Lemma 2.6. Let $n \geq 3$ be an integer. Then $\mathbb{H}_{2^n} = \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle$ has the following $2^{n-2} + 3$ conjugacy classes :

- i) The trivial conjugacy class $\{1\}$, denoted by C_1 .
- ii) The conjugacy class $\{-1\}$, denoted by C_{-1} .
- iii) Pairs of powers $\{x^k, x^{-k}\}$ for $1 \leq k \leq 2^{n-2} - 1$, denoted by C_{x^k} .
- iv) The conjugacy class $\{x^k y \mid 0 \leq k \leq 2^{n-1} - 1 \text{ even}\}$ of y , denoted by C_y .
- v) The conjugacy class $\{x^k y \mid 0 \leq k \leq 2^{n-1} - 1 \text{ odd}\}$ of xy , denoted by C_{xy} .

In particular, both $D_{2^{n-1}}$ and \mathbb{H}_{2^n} have $2^{n-2} + 3$ isomorphism classes of irreducible complex representations. We refer the reader to [9] for classical facts about the representation theory of finite groups. It is a well-known fact that even though those two groups are not isomorphic, they have the same character table.

Lemma 2.7. Let $n \geq 3$ be an integer and let $D_{2^{n-1}} = \langle r, s \mid r^{2^{n-1}} = s^2 = 1, srs^{-1} = r^{-1} \rangle$. Let also $\zeta_n = \exp\left(\frac{2i\pi}{2^{n-1}}\right) \in \mathbb{C}$. Then the following homomorphisms are representatives of the $2^{n-2} + 3$ isomorphism classes of irreducible complex representations of $D_{2^{n-1}}$:

i) Four abelian representations, coming from the abelianization of $D_{2^{n-1}}$, isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, given as follows :

$$\chi_0 : r, s \mapsto 1 \quad \chi_1 : r \mapsto 1, s \mapsto -1 \quad \chi_2 : r \mapsto -1, s \mapsto 1 \quad \chi_3 : r \mapsto -1, s \mapsto -1.$$

ii) For $1 \leq j \leq 2^{n-2} - 1$, a degree 2 representation given by :

$$r \mapsto \begin{pmatrix} \zeta_n^j & 0 \\ 0 & \zeta_n^{-j} \end{pmatrix}, \quad s \mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

with character denoted by ψ_j .

The character table of $D_{2^{n-1}}$ is

	C_1	C_{-1}	$C_{r^k}, 1 \leq k \leq 2^{n-2} - 1$	C_s	C_{rs}
χ_0	1	1	1	1	1
χ_1	1	1	1	-1	-1
χ_2	1	1	$(-1)^k$	1	-1
χ_3	1	1	$(-1)^k$	-1	1
$\psi_j, 1 \leq j \leq 2^{n-2} - 1$	2	$(-2)^j$	$\zeta_n^{jk} + \zeta_n^{-jk} = 2 \cos\left(\frac{jk\pi}{2^{n-2}}\right)$	0	0

Lemma 2.8. Let $n \geq 3$ be an integer and let $\mathbb{H}_{2^n} = \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle$. Let also $\zeta_n = \exp\left(\frac{2i\pi}{2^{n-1}}\right) \in \mathbb{C}$. Then the following homomorphisms are representatives of the $2^{n-2} + 3$ isomorphism classes of irreducible complex representations of \mathbb{H}_{2^n} :

i) Four abelian representations, coming from the abelianization of \mathbb{H}_{2^n} , isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, given as follows :

$$\chi_0 : x, y \mapsto 1 \quad \chi_1 : x \mapsto 1, y \mapsto -1 \quad \chi_2 : x \mapsto -1, y \mapsto 1 \quad \chi_3 : x \mapsto -1, y \mapsto -1.$$

ii) For $1 \leq j \leq 2^{n-2} - 1$, a degree 2 representation given by :

$$x \mapsto \begin{pmatrix} \zeta_n^j & 0 \\ 0 & \zeta_n^{-j} \end{pmatrix}, \quad y \mapsto \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

with character denoted by ψ_j .

The character table of \mathbb{H}_{2^n} is

	C_1	C_{-1}	$C_{x^k}, 1 \leq k \leq 2^{n-2} - 1$	C_y	C_{xy}
χ_0	1	1	1	1	1
χ_1	1	1	1	-1	-1
χ_2	1	1	$(-1)^k$	1	-1
χ_3	1	1	$(-1)^k$	-1	1
$\psi_j, 1 \leq j \leq 2^{n-2} - 1$	2	$(-2)^j$	$\zeta_n^{jk} + \zeta_n^{-jk} = 2 \cos\left(\frac{jk\pi}{2^{n-2}}\right)$	0	0

We remark that all the irreducible characters of $D_{2^{n-1}}$ and \mathbb{H}_{2^n} are real-valued (which does not necessarily mean they are afforded by real-valued representations, see Lemma 2.10) and they have bounded degrees (at most 2) with respect to n .

Lemma 2.9. *With the same notations as in Lemma 2.7, the irreducible character ψ_j of $D_{2^{n-1}}$, for $1 \leq j \leq 2^{n-2} - 1$, is faithful if and only if j is odd. The same holds for the irreducible characters of \mathbb{H}_{2^n} as in Lemma 2.8.*

Proof. Recall that a character χ of a group G is faithful if the corresponding representation is faithful (that is, injective). This happens if and only if $\chi(g) \neq \chi(1)$ for any $g \in G \setminus \{1\}$.

Obviously, if j is even then $\psi_j(-1) = 2 = \psi_j(1)$ so ψ_j is not faithful. Conversely, if j is odd, then $\chi(r^k)$ is never equal to 2, for $1 \leq k \leq 2^{n-2} - 1$. Indeed, since j is odd, the 2-adic valuation of jk is the same as that of k , which is $< n - 2$, therefore $\frac{jk}{2^{n-1}}$ is not an integer, and $\frac{jk\pi}{2^{n-2}} = \frac{2jk\pi}{2^{n-1}}$ is not an integer multiple of 2π . \square

Lemma 2.10. *Every irreducible character of $D_{2^{n-1}}$ is orthogonal, and with the same notations as in Lemma 2.8, the irreducible character ψ_j of \mathbb{H}_{2^n} , for $1 \leq j \leq 2^{n-2} - 1$, is symplectic if and only if j is odd.*

Proof. Clearly, χ_0, χ_1, χ_2 and χ_3 are afforded by real representations so, they are orthogonal (even in the case of \mathbb{H}_{2^n}). Consider now a non-abelian character ψ_j , $1 \leq j \leq 2^{n-2} - 1$, of $D_{2^{n-1}}$. Then by definition

$$\varepsilon_2(\psi_j) = \frac{1}{2^n} \sum_{g \in D_{2^{n-1}}} \psi_j(g^2).$$

Since each $r^k s$, with $0 \leq k \leq 2^{n-1} - 1$, satisfies $(r^k s)^2 = 1$ and $1^2 = 1$, the sum on the right-hand side contains at least $2^{n-1} + 1$ terms equal to $\psi_j(1) = 2$. Since this is more than half the number of elements in $D_{2^{n-1}}$ and each $\psi_j(g)$ is smaller than 2 in absolute value, the reverse triangular inequality implies that $\varepsilon_2(\chi) > 0$. By Theorem 1.4 it must be 1, and ψ_j is orthogonal.

Now let ψ_j , $1 \leq j \leq 2^{n-2} - 1$, be a non-abelian irreducible character of \mathbb{H}_{2^n} . The squares in \mathbb{H}_{2^n} are precisely the even powers of x . Indeed, since $y^{-1} = -y$, we have, for any $k \in \{0, \dots, 2^{n-1} - 1\}$,

$$(x^k y)^2 = -x^k y x^k y^{-1} = -1.$$

From this, and denoting $r(h)$ the number of square roots of $h \in \mathbb{H}_{2^n}$, we see that $r(-1) = 2^{n-1} + 2$ while $r(x^{2k}) = 2$ for $0 \leq k \leq 2^{n-3} - 1$. Now if j is odd, then

$$\begin{aligned} \varepsilon_2(\psi_j) &= \frac{1}{2^n} \left(4 - 2(2^{n-1} + 2) + 4 \sum_{k=1}^{2^{n-3}-1} \cos\left(\frac{2jk\pi}{2^{n-2}}\right) \right) \\ &= -1 + \frac{1}{2^{n-2}} \sum_{k=1}^{2^{n-3}-1} \cos\left(\frac{2jk\pi}{2^{n-2}}\right). \end{aligned}$$

It is easily seen from the triangular inequality that this quantity has to be negative. As before, this shows that $\varepsilon_2(\psi_j) = -1$.

Conversely, if j is even, the reverse triangular inequality implies that

$$\varepsilon_2(\psi_j) = \frac{1}{2^n} \left(4 + 2(2^{n-1} + 2) + 4 \sum_{k=1}^{2^{n-3}-1} \cos \left(\frac{2jk\pi}{2^{n-2}} \right) \right) > 0,$$

which implies that $\varepsilon_2(\psi_j) = 1$. \square

Let us remark that for $n \geq 4$, the group $\mathbb{H}_{2^n} = \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle$ contains $\mathbb{H}_{2^{n-1}}$ as a subgroup generated by $\{x^2, y\}$, and inductively, contains $\mathbb{H}_{2^i} = \langle x^{2^{n-i}}, y \rangle$ for every $3 \leq i \leq n$. Similarly, for $n \geq 3$, $D_{2^{n-2}}$ appears as a subgroup of $D_{2^{n-1}} = \langle r, s \mid r^{2^{n-1}} = s^2 = 1, sr s^{-1} = r^{-1} \rangle$ generated by $\langle r^2, s \rangle$. In particular when L/K is a Galois extension of number field extension with group \mathbb{H}_{2^n} or $D_{2^{n-1}}$, it contains subextensions with Galois groups \mathbb{H}_{2^i} or $D_{2^{i-1}}$ for every $3 \leq i \leq n$. To study Chebyshev's bias in such subextensions, we will need to know how irreducible characters are induced from such \mathbb{H}_{2^i} to \mathbb{H}_{2^n} , and from $D_{2^{i-1}}$ to $D_{2^{n-1}}$.

Lemma 2.11. *With the same notations as in Lemma 2.8, denote by $\psi_k^{(i)}$, for $3 \leq i \leq n$, $1 \leq k \leq 2^{i-2} - 1$, the character of the subgroup \mathbb{H}_{2^i} of \mathbb{H}_{2^n} associated to the representation*

$$x^{2^{n-i}} \mapsto \begin{pmatrix} \zeta_i^k & 0 \\ 0 & \zeta_i^{-k} \end{pmatrix}, \quad y \mapsto \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

and denote by $\chi_0^{(i)}, \chi_1^{(i)}, \chi_2^{(i)}$ and $\chi_3^{(i)}$ the four abelian characters of \mathbb{H}_{2^i} corresponding to the notations of Lemma 2.8. For $k \in \{1, \dots, 2^{i-2} - 1\}$, one has

$$\text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \psi_k^{(i)} = \sum_{\substack{1 \leq l \leq 2^{n-2}-1 \\ l \equiv \pm k \pmod{2^{i-1}}}} \psi_l.$$

Also,

$$\text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \chi_0^{(i)} = \text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \chi_1^{(i)} = \chi_0 + \chi_1 + \chi_2 + \chi_3 + \sum_{\substack{1 \leq j \leq 2^{n-2}-1 \\ j \equiv 0 \pmod{2^{i-1}}}} \psi_j$$

and

$$\text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \chi_2^{(i)} = \text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \chi_3^{(i)} = \sum_{\substack{1 \leq j \leq 2^{n-2}-1 \\ j \equiv 0 \pmod{2^{i-2}}, j \not\equiv 0 \pmod{2^{i-1}}}} \psi_j.$$

Those equalities also hold when \mathbb{H}_{2^n} and \mathbb{H}_{2^i} are replaced by $D_{2^{n-1}}$ and $D_{2^{i-1}}$ respectively.

Proof. Let $1 \leq j \leq 2^{n-2} - 1$ be odd. We use the Frobenius reciprocity formula and the character table in Lemma 2.8 to compute

$$\begin{aligned} \langle \text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \psi_k^{(i)}, \psi_j \rangle_{\mathbb{H}_{2^n}} &= \langle \psi_k^{(i)}, \psi_{j|\mathbb{H}_{2^i}} \rangle_{\mathbb{H}_{2^i}} = \frac{1}{2^i} \sum_{g \in \mathbb{H}_{2^i}} \psi_k^{(i)}(g) \overline{\psi_j(g)} \\ &= \frac{1}{2^i} \left(2 \cdot 2 + (-2) \cdot (-2) + 2 \sum_{1 \leq l \leq 2^{i-2}-1} \psi_k^{(i)}(x^{2^{n-i}}) \overline{\psi_j(x^{2^{n-i}})} \right) \\ &= \frac{1}{2^i} \left(8 + 2 \sum_{1 \leq l \leq 2^{i-2}-1} (\zeta_i^{kl} + \zeta_i^{-kl}) (\zeta_n^{jl2^{n-i}} + \zeta_n^{-jl2^{n-i}}) \right). \end{aligned}$$

Since $\zeta_n^{2^m} = \zeta_{n-2^m}$ for any $0 \leq m \leq n-1$, we find

$$\begin{aligned} \langle \text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \psi_k^{(i)}, \psi_j \rangle_{\mathbb{H}_{2^n}} &= \frac{1}{2^i} \left(8 + 2 \sum_{1 \leq l \leq 2^{i-2}-1} (\zeta_i^{kl} + \zeta_i^{-kl}) (\zeta_i^{jl} + \zeta_i^{-jl}) \right) \\ &= \frac{1}{2^i} \left(8 + 2 \sum_{1 \leq l \leq 2^{i-2}-1} (\zeta_i^{(k+j)l} + \zeta_i^{-(k+j)l} + \zeta_i^{(k-j)l} + \zeta_i^{-(k-j)l}) \right) \\ &= \frac{1}{2^i} \left(8 + 2 \sum_{\substack{1 \leq l \leq 2^{i-1}-1 \\ l \neq 2^{i-2}}} (\zeta_i^{(k+j)l} + \zeta_i^{(k-j)l}) \right). \end{aligned}$$

Now, if $k+j = 0 \pmod{2^{i-1}}$ then $k-j \neq 0 \pmod{2^{i-1}}$, otherwise we would get $2k = 0 \pmod{2^{i-1}}$, *i.e.* $k = 0 \pmod{2^{i-2}}$, which cannot be because $1 \leq k \leq 2^{i-2}-1$. Therefore, the geometric progressions missing two terms sum to

$$\begin{aligned} \sum_{\substack{1 \leq l \leq 2^{i-1}-1 \\ l \neq 2^{i-2}}} (\zeta_i^{(k+j)l} + \zeta_i^{(k-j)l}) &= 2^{i-1} - 2 + \left(\frac{1 - \zeta_i^{(k-j)2^{i-1}}}{1 - \zeta_i^{(k-j)}} - \zeta_i^{(k-j)2^{i-2}} - 1 \right) \\ &= 2^{i-1} - 4. \end{aligned}$$

Similarly, if $k-j = 0 \pmod{2^{i-1}}$, then $k+j \neq 0 \pmod{2^{i-1}}$, and we find

$$\sum_{\substack{1 \leq l \leq 2^{i-1}-1 \\ l \neq 2^{i-2}}} (\zeta_i^{(k+j)l} + \zeta_i^{(k-j)l}) = 2^{i-1} - 4.$$

Finally, if $k+j \neq 0 \pmod{2^{i-1}}$ and $k-j \neq 0 \pmod{2^{i-1}}$, we find

$$\sum_{\substack{1 \leq l \leq 2^{i-1}-1 \\ l \neq 2^{i-2}}} (\zeta_i^{(k+j)l} + \zeta_i^{(k-j)l}) = -4.$$

To sum up, we have found

$$\langle \text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \psi_k^{(i)}, \psi_j \rangle_{\mathbb{H}_{2^n}} = \begin{cases} 1 & \text{if } k+j = 0 \pmod{2^{i-1}} \text{ or } k-j = 0 \pmod{2^{i-1}} \\ 0 & \text{otherwise} \end{cases}.$$

We have found 2^{n-i} irreducible components of $\text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \psi_k^{(i)}$ of degree 2. Since $\text{Ind}_{\mathbb{H}_{2^i}}^{\mathbb{H}_{2^n}} \psi_k^{(i)}(1) = [\mathbb{H}_{2^n} : \mathbb{H}_{2^i}] \psi_k^{(i)}(1) = 2^{n-i+1}$, these are the only ones.

The other induced characters are computed in a similar manner, using the Frobenius reciprocity formula. The fact that \mathbb{H}_{2^n} and $D_{2^{n-1}}$ have the same character table with conjugacy classes indexed similarly with respect to their generators imply that the same computations work in the $D_{2^{n-1}}$ case. \square

Corollary 2.12. *Let $3 \leq i \leq n-1$. If $\chi \in \text{Irr}(\mathbb{H}_{2^i})$ then for any $\chi' \in \text{Irr}(\mathbb{H}_{2^i}) \setminus \{\chi\}$, inducing from \mathbb{H}_{2^i} to \mathbb{H}_{2^n} we have $(\chi, \chi') = 1$ unless $\deg(\chi) = 1$. This also holds when \mathbb{H}_{2^n} and \mathbb{H}_{2^i} are replaced by $D_{2^{n-1}}$ and $D_{2^{i-1}}$ respectively.*

The above conditions will allow us to obtain a lower bound for $1 - \delta$ for relative number field extensions (i.e. with base field different from \mathbb{Q}) with Galois groups \mathbb{H}_{2^i} and $\mathcal{D}_{2^{i-1}}$ using Theorem 4.14.

The following lemma will be used to compute moments in Section 4.

Lemma 2.13. *For $i \geq 3$ and $1 \leq k \leq 2^{i-2} - 1$,*

$$\sum_{\substack{j=1 \\ j \text{ odd}}}^{2^{i-2}-1} (\zeta_i^{jk} + \zeta_i^{-jk}) = 0.$$

Proof. This is clear for $i = 3$, so assume $i \geq 4$. This sum can be rewritten

$$\sum_{j=1}^{2^{i-2}-1} (\zeta_i^{jk} + \zeta_i^{-jk}) - \sum_{h=1}^{2^{i-3}-1} (\zeta_i^{2hk} + \zeta_i^{-2hk}).$$

As in the previous proof, we see that the first sum is simply

$$\sum_{j=0}^{2^{i-1}-1} \zeta_i^{jk} - \zeta_i^0 - \zeta_i^{k2^{i-2}} = -1 - (-1)^k$$

since $0 < k < 2^{i-1}$ and so $\zeta_i^k \neq 1$. The second sum

$$\sum_{h=1}^{2^{i-3}-1} (\zeta_i^{2hk} + \zeta_i^{-2hk})$$

is being dealt with similarly, because $\zeta_i^2 = \zeta_{i-1}$ and $0 < k < 2^{i-2}$, so it also equals $-1 - (-1)^k$. Thus the two sums cancel each other. \square


3 Extensions of \mathbb{Q} of group \mathbb{H}_8 : horizontal Chebotarev biases

In this section, we will depart temporarily from the notations we used in section 2, and we will use the more familiar i, j, k notations for elements of \mathbb{H}_8 . Our goal is to prove Theorem A. We will be constructing two distinct families of number fields with Galois group over \mathbb{Q} isomorphic to \mathbb{H}_8 and with opposite biases along each family, due to the existence or not of a central zero for the corresponding Dedekind zeta functions.

Recall that $\mathbb{H}_8 = \{\pm 1, \pm i, \pm j, \pm k\}$ is the quaternion group of order 8. Its elements all commute with -1 and satisfy the following relations

$$i^2 = j^2 = k^2 = -1, ij = k, ji = -k, jk = i, kj = -i, ki = j \text{ and } ik = -j.$$

Here is the lattice of subgroups of \mathbb{H}_8 (see [3])



The character table of \mathbb{H}_8 is

	C_1	C_{-1}	C_i	C_j	C_k
χ_0	1	1	1	1	1
χ_i	1	1	1	-1	-1
χ_j	1	1	-1	1	-1
χ_k	1	1	-1	-1	1
ψ	2	-2	0	0	0

where C_x denotes the conjugacy class of $x \in \mathbb{H}_8$.

Let K/\mathbb{Q} be a Galois extension, with Galois group G isomorphic to \mathbb{H}_8 . Assuming LI, we can compute the values of the different means and variances, according to the formulas given in Theorem 1.11 (since

$$\mathbb{E}(X(K/\mathbb{Q}, C_1, C_2)) = -\mathbb{E}(X(K/\mathbb{Q}, C_2, C_1)) \text{ and } \text{Var}(X(K/\mathbb{Q}, C_1, C_2)) = \text{Var}(X(K/\mathbb{Q}, C_2, C_1)),$$

there are only $\binom{5}{2} = 10$ races to consider).

Proposition 3.1. *Assume LI for the extension K/\mathbb{Q} . Let $\mathfrak{o} = \text{ord}_{s=\frac{1}{2}} L(s, \psi, K/\mathbb{Q})$, where ψ is the non-abelian character of \mathbb{H}_8 in the above character table. Then one has*

a	b	$\mathbb{E}(X(K/\mathbb{Q}, C_a, C_b))$	$\text{Var}(X(K/\mathbb{Q}, C_a, C_b))$
1	-1	$4(1 - 2\mathfrak{o})$	$16B_0(\psi)$
1	i, j, k	$-2(1 + 2\mathfrak{o})$	$4 \sum_{\chi \neq \chi_b} B_0(\chi)$
-1	i, j, k	$2(2\mathfrak{o} - 3)$	$4 \sum_{\chi \neq \chi_b} B_0(\chi)$
i, j, k	i, j, k	0	$4B_0(\chi_a) + 4B_0(\chi_b)$

where B_0 is defined in Lemma 2.3.

Remarks. Let us make a few comments on these values :

- i) The presence or not of a zero at $\frac{1}{2}$ for $s \mapsto L(s, \psi, K/\mathbb{Q})$ changes the sign $\mathbb{E}(X(K/\mathbb{Q}, C_1, C_{-1}))$.
- ii) The race between C_1 and C_b , for $b \in \{i, j, k\}$, is always biased towards C_b . This was expected since 1 is a square in G , and b is not.
- iii) The presence or not of a zero with multiplicity at least 2 at $\frac{1}{2}$ for $s \mapsto L(s, \psi, K/\mathbb{Q})$ changes the sign $\mathbb{E}(X(K/\mathbb{Q}, C_{-1}, C_b))$, for $b \in \{i, j, k\}$.
- iv) There is no bias in the race between C_a and C_b , for $a \in \{i, j, k\}$ and $b \in \{i, j, k\} \setminus \{a\}$.

Since we will be using the hypothesis LI^+ , under which the order of vanishing \mathfrak{o} can only be 0 or 1, and we want to exhibit a change of leading conjugacy class in Chebotarev races, we will now focus on the conjugacy classes C_1 and C_{-1} . When K/\mathbb{Q} is tamely ramified, we can deduce more precise bounds for the variances.

Proposition 3.2. *Assume LI and that K/\mathbb{Q} is tamely ramified. Then*

$$\text{Var}(X(K/\mathbb{Q}, C_1, C_{-1})) \asymp \log |d_K|.$$

Proof. Since \mathbb{K}/\mathbb{Q} tame, the filtration of the inertia subgroup (see section 2.1) at any prime number ramified in K only has length 1, *i.e.* for any prime p and $i \geq 1$, $|G_i(p)| = 1$. In particular, for any non-trivial character χ of G and any prime number p , one has

$$n(\chi, p) = \text{codim } V^{\mathcal{I}(p)},$$

where V is the space of the representation affording χ .

Using the definition, we find for any prime p ramified in K (so that $\mathcal{I}(p) = G_0(p) \neq \{1\}$),

$$n(\chi_i, p) = \begin{cases} 1 & \text{if } \mathcal{I}(p) = \langle j \rangle \text{ or } \mathcal{I}(p) = \langle k \rangle \\ 0 & \text{otherwise} \end{cases},$$

$$n(\chi_j, p) = \begin{cases} 1 & \text{if } \mathcal{I}(p) = \langle i \rangle \text{ or } \mathcal{I}(p) = \langle k \rangle \\ 0 & \text{otherwise} \end{cases},$$

$$n(\chi_k, p) = \begin{cases} 1 & \text{if } \mathcal{I}(p) = \langle i \rangle \text{ or } \mathcal{I}(p) = \langle j \rangle \\ 0 & \text{otherwise} \end{cases}$$

and

$$n(\psi, p) = 2.$$

This yields

$$\mathfrak{f}(K/\mathbb{Q}, \chi_i) = \prod_{\substack{p|d_K \\ \mathcal{I}(p)=\langle j \rangle}} p \times \prod_{\substack{p|d_K \\ \mathcal{I}(p)=\langle k \rangle}} p,$$

$$\mathfrak{f}(K/\mathbb{Q}, \chi_j) = \prod_{\substack{p|d_K \\ \mathcal{I}(p)=\langle i \rangle}} p \times \prod_{\substack{p|d_K \\ \mathcal{I}(p)=\langle k \rangle}} p,$$

$$\mathfrak{f}(K/\mathbb{Q}, \chi_k) = \prod_{\substack{p|d_K \\ \mathcal{I}(p)=\langle i \rangle}} p \times \prod_{\substack{p|d_K \\ \mathcal{I}(p)=\langle j \rangle}} p$$

and

$$f(K/\mathbb{Q}, \psi) = \prod_{p|d_K} p^2.$$

Since the base field is \mathbb{Q} , we have $A(\chi) = f(K/\mathbb{Q}, \chi)$ for any character χ of G , and the conductor-discriminant formula (Theorem 2.1) thus gives

$$|d_K| = \prod_{\substack{p|d_K \\ e_p=8}} p^4 \prod_{\substack{p|d_K \\ e_p=4}} p^6 \prod_{\substack{p|d_K \\ e_p=2}} p^4,$$

where e_p is the ramification index of the ramified prime p . In particular, we have

$$A(\psi)^2 \leq d_K \leq A(\psi)^3,$$

so that

$$\log A(\psi) \asymp \log d_K.$$

Finally, Theorem 1.11 and Lemma 2.3 yield

$$\text{Var}(X(K/\mathbb{Q}, C_1, C_{-1})) \asymp B_0(\psi) \asymp \log A(\psi) \asymp \log |d_K|.$$

□

The fact that a bound such as Proposition 3.2 holds follows from the fact that the character ψ is faithful, because this implies that each ramified prime appears with positive exponent in $A(\chi) = f(K/\mathbb{Q}, \psi)$. This observation will be used again in our study of biases in towers of extensions (see Proposition 4.4).

We will use the following theorem of Fröhlich to construct the quaternion extensions of \mathbb{Q} we will be interested in :

Theorem 3.3 ([6]). *Let d be a square-free integer, $d > 1$ and $d \equiv 1 \pmod{4}$. Let \mathfrak{R} be a finite set of primes unramified in $\mathbb{Q}(\sqrt{d})$ and not containing 2. Then there exist infinitely many tamely ramified extensions K/\mathbb{Q} and L/\mathbb{Q} such that :*

- i) $\text{Gal}(K/\mathbb{Q}) \simeq \text{Gal}(L/\mathbb{Q}) \simeq \mathbb{H}_8$.*
- ii) $\mathbb{Q}(\sqrt{d}) \subset K \cap L$.*
- iii) Every prime in \mathfrak{R} is ramified in K and L .*
- iv) $W(\psi, K/\mathbb{Q}) = -1$ and $W(\psi, L/\mathbb{Q}) = 1$.*

Remark. Theorem 3.3 is in fact a weak version of Fröhlich's theorem in [6]. Actually, one can specify any finite number of unramified primes in K and L as well (the ramification must be compatible with the fact that $\mathbb{Q}(\sqrt{d}) \subset K \cap L$), and one can also ask for K and L to be totally real or totally imaginary.

Corollary 3.4. *For any square-free integer $d > 1$ with $d \equiv 1 \pmod{4}$ and any prime p , there exist infinitely many tamely ramified extensions K/\mathbb{Q} and L/\mathbb{Q} such that :*

i) $\text{Gal}(K/\mathbb{Q}) \simeq \text{Gal}(L/\mathbb{Q}) \simeq \mathbb{H}_8$.

ii) $\mathbb{Q}(\sqrt{d}) \subset K \cap L$.

iii) p is ramified in K and L .

iv) $L\left(\frac{1}{2}, \psi, K/\mathbb{Q}\right) = 0$ and if LI^+ holds then $L\left(\frac{1}{2}, \psi, L/\mathbb{Q}\right) \neq 0$.

Proof. The existence of the number fields K follows from Theorem 3.3 without the assumption of LI^+ : we consider tamely ramified extensions K/\mathbb{Q} with Galois group \mathbb{H}_8 , ramified at p , containing $\mathbb{Q}(\sqrt{d})$ and such that $W(\psi) = -1$. Proposition 1.5 shows that $L(1/2, \psi, K/\mathbb{Q}) = 0$.

The existence of the number fields L follows similarly because, assuming LI^+ , $W(\psi) = 1$ implies that $s \mapsto L(s, \psi, L/\mathbb{Q})$ does not vanish at $\frac{1}{2}$. \square

We are now ready to prove Theorem A, which we recall for convenience.

Theorem 3.5. *Assume GRH and LI^+ . For any function f such that $f(n) \xrightarrow{n \rightarrow +\infty} +\infty$, there exist two infinite families $(K_d)_d$ and $(L_d)_d$ of Galois extensions of \mathbb{Q} , indexed by square-free integers satisfying $d > 1$ and $d \equiv 1 \pmod{4}$, such that for any d in the index set :*

i) $\mathbb{Q}(\sqrt{d}) \subset K_d \cap L_d$.

ii) $\text{Gal}(K_d/\mathbb{Q}) \simeq \text{Gal}(L_d/\mathbb{Q}) \simeq \mathbb{H}_8$.

iii) $0 < \frac{1}{2} - \delta(K_d/\mathbb{Q}, C_1, C_{-1}) \ll \frac{1}{f(d)}$.

iv) $0 < \delta(L_d/\mathbb{Q}, C_1, C_{-1}) - \frac{1}{2} \ll \frac{1}{f(d)}$.

Proof. For any square-free $d > 1$ with $d \equiv 1 \pmod{4}$, we choose a field K_d given in the first part of Corollary 3.4, ramified at the smallest prime larger than $e^{f(d)^3}$. Since K_d/\mathbb{Q} is tamely ramified and we are assuming LI^+ (so that $\mathfrak{o} = 1$), the computations made in Proposition 3.2 yield

$$B(X(K_d/\mathbb{Q}, C_1, C_{-1})) = \frac{\mathbb{E}(X(K/\mathbb{Q}, C_1, C_{-1}))}{\sqrt{\text{Var}(X(K/\mathbb{Q}, C_1, C_{-1}))}} = \frac{1 - 2\mathfrak{o}}{\sqrt{2}B_0(\psi)^{1/2}} = -\frac{1}{\sqrt{2}B_0(\psi)^{1/2}}.$$

In particular, $\mathbb{E}(X(K/\mathbb{Q}, C_1, C_{-1}))$ is negative, because of the existence of a real zero of $s \mapsto L(s, \psi, K_d/\mathbb{Q})$, and this implies that $\delta(K_d/\mathbb{Q}, C_1, C_{-1}) < \frac{1}{2}$. Lemma 2.3 yields

$$B_0(\psi) \asymp \log A(\psi),$$

which satisfies (as remarked in the proof of Proposition 3.2)

$$\log A(\psi) \asymp \log |d_{K_d}|,$$

since ψ is faithful. Moreover, since K_d/\mathbb{Q} is ramified at the smallest prime larger than $e^{f(d)^3}$, we have

$$\log |d_{K_d}| \gg \log \left(e^{f(d)^3} \right) = f(d)^3.$$

Therefore,

$$|B(X(K_d/\mathbb{Q}, C_1, C_{-1}))| \ll \frac{1}{f(d)^{3/2}}.$$

Similarly, we have

$$\text{Var}(X(K/\mathbb{Q}, C_1, C_{-1}))^{-1} \ll \frac{1}{f(d)^3}.$$

Since Artin's conjecture holds for extensions with Galois group \mathbb{H}_8 (recall that this group is supersolvable, and that Artin's conjecture is known in that case), Theorem 1.14 implies that

$$0 < \frac{1}{2} - \delta(K_d/\mathbb{Q}, C_1, C_{-1}) \ll \frac{1}{f(d)}.$$

The construction of L_d follows along the same lines, by choosing fields as in the second part of Corollary 3.4. In that case, the mean is positive because there is no real zero of $s \mapsto L(s, \psi, L_d/\mathbb{Q})$, so $\delta(K_d/\mathbb{Q}, C_1, C_{-1}) > \frac{1}{2}$, and the estimates are the same. \square

Remark. If we had a statement analogous to Theorem 3.3 in which we could specify if $s \mapsto L(s, \psi, K/\mathbb{Q})$ vanishes, or not, at $\frac{1}{2}$ with multiplicity at least two, then we would be able to obtain a result similar to Theorem A for the race between C_{-1} and C_b , for $b \in \{i, j, k\}$, *i.e.* moderately biased races with biases of opposite signs, bounded in absolute value by $\frac{1}{f(d)}$, for any function f going to infinity. Such a result would of course contradict LI^+ .

4 Chebyshev's bias in towers

We now consider the vertical aspect of our problem. Our goal is to prove Theorems B, C and D. Instead of working with a family of extensions of \mathbb{Q} with fixed Galois group, we want to study different Galois extensions of \mathbb{Q} with Galois groups dihedral of order a power of two or generalized quaternion of increasing sizes. We also want to compare Chebyshev's bias in subextensions. For this to make sense, we need to make a few observations first.

For $n \geq 4$, $\mathbb{H}_{2^{n-1}}$ appears as a (normal) subgroup of \mathbb{H}_{2^n} . Indeed, if $\mathbb{H}_{2^n} = \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle$, then it is easy to see that $\langle x^2, x^k y \rangle$ is a normal subgroup of \mathbb{H}_{2^n} isomorphic to $\mathbb{H}_{2^{n-1}}$, for any $0 \leq k \leq 2^{n-2} - 1$. Therefore, if K/\mathbb{Q} is a number field extension with $\text{Gal}(K/\mathbb{Q}) \simeq \mathbb{H}_{2^n}$ then there exist number fields $K \supset K_3 \supset K_4 \supset \cdots \supset K_{n-1} \supset K_n = \mathbb{Q}$ with $\text{Gal}(K/K_i) \simeq \mathbb{H}_{2^i}$ for all $3 \leq i \leq n$, namely the subfields fixed by $\langle x^{2^{n-i}}, y \rangle$. In the following, we will always assume that the quaternion subgroups have been chosen so that $\text{Gal}(K/K_i)$ is generated by $x^{2^{n-i}}$ and y , for $3 \leq i \leq n-1$. Similarly, if $n \geq 3$ then $D_{2^{n-1}}$ is a normal subgroup of $D_{2^n} = \langle r, s \mid r^{2^{n-1}} = s^2, sr s^{-1} = r^{-1} \rangle$, generated for example by r^2 and s . Those observations will allow us to compare Chebyshev's bias in subextensions (see Theorem 4.17).

4.1 Moments in generalized quaternion Galois groups

In this section, we compute bounds on the moments of the random variables attached to the different Chebotarev races in number field extensions with Galois group generalized quaternion. They will be used in the proof of Theorem C.

We will use the following fact about root numbers of symplectic characters of generalized quaternion groups.

Theorem 4.1 ([7], Theorem 3). *Let N/L be a tamely ramified extension of number fields, with Galois group generalized quaternion. Then the root numbers of the symplectic irreducible characters of $\text{Gal}(N/L)$ are all equal.*

Now, let $n \geq 3$, and assume that K/\mathbb{Q} is a Galois extension with Galois group

$$G := \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle \simeq \mathbb{H}_{2^n},$$

and let $\mathbb{Q} = K_n \subset \cdots \subset K_3 \subset K$ be intermediate number fields as before, that is

$$G_i := \langle x^{2^{n-i}}, y \rangle \simeq \mathbb{H}_{2^i}$$

and

$$K_i := K^{G_i},$$

for $3 \leq i \leq n$. We will denote $x^{2^{n-i}}$ by x_i so that $G_i = \langle x_i, y \rangle$. Conjugacy classes in G_i will be denoted, according to our notations in Lemma 2.6, by $C_1^{(i)}, C_{-1}^{(i)}, C_{x_i^k}^{(i)}, C_y^{(i)}$ and $C_{x_i y}^{(i)}$. Note that each G_i is supersolvable, so Artin's conjecture is known to hold for Artin L -functions attached to irreducible characters of G_i .

Recall that if C is a conjugacy class of G_i , then C^+ is the conjugacy class $\bigcup_{g \in G} gCg^{-1}$ of $G_i^+ = G$. The following lemma is immediate.

Lemma 4.2. *Let $i \in \{3, \dots, n-1\}$. Using the notations of section 2, we have*

- i) $C_1^{(i)+} = C_1$.
- ii) $C_{-1}^{(i)+} = C_{-1}$.
- iii) For any $1 \leq k \leq 2^{i-2} - 1$, $C_{x_i^k}^{(i)+} = C_{x_i^k}$.
- iv) $C_y^{(i)+} = C_y$.
- v) For any $1 \leq k \leq 2^{i-2} - 1$, $C_{x_i y}^{(i)+} = C_y$.

In particular, if C_1 and C_2 are distinct conjugacy classes of G_i , then $C_1^+ \neq C_2^+$, unless, $\{C_1, C_2\} = \{C_y^{(i)}, C_{x_i y}^{(i)}\}$.

When K/\mathbb{Q} is tamely ramified, we denote by W_{K/K_i} the root number of any of the symplectic characters of $\text{Gal}(K/K_i)$. We first relate the root numbers of symplectic characters of G to those of each G_i , and the orders of vanishing at $1/2$ of the corresponding Artin L -functions.

Proposition 4.3. *Assume LI^+ and assume K/\mathbb{Q} is tamely ramified. Then for any $3 \leq i \leq n$, we have $W_{K/K_i} = W_K$, where $W_k = W_{k/\mathbb{Q}}$ is the root number of any symplectic character of G . Moreover, if $W_K = -1$ then only the symplectic characters of G_i have their Artin L -function vanish at $1/2$, and the order of vanishing is 2^{n-i} .*

Proof. Clearly, K/K_i is tamely ramified so our definition of W_{K/K_i} makes sense using Theorem 4.1. For any $3 \leq j \leq n$, consider the classical decomposition

$$\begin{aligned}
\zeta_K(s) &= L(s, \chi_0, K/K) \\
&= L(s, \text{Ind}_{\{1\}}^{G_j} \chi_0, K/K_j) \\
&= L(s, \text{reg}_{G_j}, K/K_j) \\
&= L(s, \sum_{\chi \in \text{Irr}(G_j)} \chi(1)\chi, K/K_j) \\
&= \prod_{\chi \in \text{Irr}(G_j)} L(s, \chi, K/K_j)^{\chi(1)},
\end{aligned}$$

where reg_{G_j} is the character of the regular representation of G_j .

If $W_K = 1$, consider the previous factorization with $j = n$. By LI^+ , none of the factors vanish at $1/2$, and so ζ_K does not vanish at $1/2$. Using now the decomposition with $j = i$, we see that $L(1/2, \chi, K/K_i) \neq 0$ for each irreducible character χ of G_i . In particular, $W_{K/K_i} \neq -1$, i.e. $W_{K/K_i} = 1$.

Conversely, if $W_K = -1$, then ζ_K vanishes at $1/2$ to order 2^{n-2} . Indeed, under LI^+ , the only factors that vanish at $1/2$ are the $L(s, \chi, K/\mathbb{Q})^{\chi(1)}$ for $\chi = \chi_0$ or χ symplectic. There are 2^{n-3} symplectic characters, all vanishing at $1/2$ to order one (again, because of LI^+ , and the fact all such characters have root number -1) and satisfying $\chi(1) = 2$. Moreover, $L(s, \chi_0, K/\mathbb{Q}) = \zeta(s)$ is the classical Riemann ζ function, which is known not to vanish at $1/2$. Using Lemma 2.11, we see that for $1 \leq k \leq 2^{i-2} - 1$ odd we have

$$\begin{aligned}
L(s, \psi_k^{(i)}, K/K_i) &= L(s, \text{Ind}_{G_i}^G \psi_k^{(i)}, K/\mathbb{Q}) \\
&= L(s, \sum_{\substack{0 \leq l \leq 2^{n-2}-1 \\ l \equiv \pm k \pmod{2^{i-1}}} \psi_l, K/\mathbb{Q}) \\
&= \prod_{\substack{0 \leq l \leq 2^{n-2}-1 \\ l \equiv \pm k \pmod{2^{i-1}}} L(s, \psi_l, K/\mathbb{Q}).
\end{aligned}$$

Since each $L(s, \psi_l, K/\mathbb{Q})$, with $1 \leq l \leq 2^{n-2} - 1$ odd, vanishes at $1/2$ to order one, we see that each $L(s, \psi_k^{(i)}, K/K_i)$ vanishes at $1/2$ as well, to order 2^{n-i} , for $1 \leq k \leq 2^{i-2} - 1$ odd. In particular, $W_{K/K_i} = -1$. Moreover, the 2^{i-3} symplectic characters of G_i contribute to the vanishing of ζ_K at $1/2$ to order $2 \times 2^{i-3} \times 2^{n-i} = 2^{n-2}$, so no other irreducible character of G_i have its Artin L -function vanishing at $1/2$. \square

In Proposition 4.4, 4.5 and 4.6, we give bounds on the variances and compute the means of the random variables attached to the Chebotarev races in the extensions K/K_i .

Proposition 4.4. *Assume GRH and LI^- . Then for any $3 \leq i \leq n$ and any distinct conjugacy classes C_1, C_2 of G_i such that $\{C_1, C_2\} \neq \{C_y^{(i)}, C_{xiy}^{(i)}\}$,*

$$\text{Var}(X(K/K_i, C_1, C_2)) \ll \log |d_K|.$$

Moreover if K/\mathbb{Q} is tamely ramified then for any $3 \leq i \leq n$,

$$\text{Var}(X(K/K_i, C_1, C_2)) \gg \frac{\log |d_K|}{4^n},$$

and if $\{C_1, C_2\} \cap \{C_{x_i^k}^{(i)} \mid 1 \leq k \leq 2^{i-2} - 1\} = \emptyset$ this can be improved to

$$\text{Var}(X(K/K_i, C_1, C_2)) \gg \log |d_K|.$$

Proof. By Lemma 4.2, the condition $\{C_1, C_2\} \neq \{C_y^{(i)}, C_{x_i y}^{(i)}\}$ ensures that $C_1^+ \neq C_2^+$. Recall that

$$\text{Var}(X(K/K_i, C_1, C_2)) = \sum_{\chi \in \text{Irr}(G^+)} |\chi(C_1^+) - \chi(C_2^+)|^2 B_0(\chi)$$

by Theorem 1.11, with

$$B_0(\chi) \asymp \log A(\chi)$$

by Lemma 2.3. In particular, we see using Lemma 4.2 that our estimates on $\text{Var}(X(K/K_i, C_1, C_2))$ do not depend on K_i , so it is enough to prove them in the case $i = n$, that is $K_i = \mathbb{Q}$.

Since the values taken by the irreducible characters of G are bounded in absolute value uniformly in n , we get

$$\text{Var}(X(K/\mathbb{Q}, C_1, C_2)) \ll \sum_{\chi \in \text{Irr}(G)} \log A(\chi) \leq \sum_{\chi \in \text{Irr}(G)} \chi(1) \log A(\chi).$$

Recall that for any irreducible character χ of G ,

$$A(\chi) = \mathfrak{f}(K/\mathbb{Q}, \chi)$$

is the Artin conductor of χ , so the conductor-discriminant formula (Theorem 2.1) yields

$$\begin{aligned} \sum_{\chi \in \text{Irr}(G)} \chi(1) \log A(\chi) &= \sum_{\chi} \chi(1) \mathfrak{f}(K/\mathbb{Q}, \chi) \\ &= \log |d_K|. \end{aligned}$$

This proves the stated upper bound.

To prove the first lower bound, assume K/\mathbb{Q} is tamely ramified. If χ is an irreducible symplectic character of G then χ is faithful by Lemma 2.9. This implies that there are no invariant vectors for the representation of character χ . Moreover, since K/\mathbb{Q} is tamely ramified, the ramification groups of index ≥ 2 are trivial for any prime p ramified in K , and for any such prime we find $n(\chi, p) = 2$ (see section 2.1 for the definition of $n(\chi, p)$). In particular,

$$A(\chi) = \mathfrak{f}(K/\mathbb{Q}, \chi) = \prod_{p|d_K} p^2.$$

Now, since K/\mathbb{Q} is tamely ramified, each prime \mathfrak{p} of K above a ramified prime number p appears in the factorization of $\partial_{K/\mathbb{Q}}$, the different of K/\mathbb{Q} , with exponent $e_{\mathfrak{p}} - 1$ where $e_{\mathfrak{p}}$ denotes the corresponding ramification index ([17, Theorem 2.6]). Since the discriminant of K/\mathbb{Q} is the K/\mathbb{Q} -norm of the different, this yields

$$|d_K| = \prod_{\mathfrak{p}|\partial_{K/\mathbb{Q}}} N_{K/\mathbb{Q}}(\mathfrak{p})^{e_{\mathfrak{p}}-1} = \prod_{p|d_K} p^{(e_p-1)f_p g_p},$$

where f_p (respectively g_p) denotes the residual degree of (respectively the number of primes above) the prime p . In particular, this exponent is less than $e_p f_p g_p = [K : \mathbb{Q}] = 2^n$. Therefore we have

$$\begin{aligned} \prod_{p|d_K} p^{2^n} &\geq \prod_{p|d_K} p^{(e_p-1)f_p g_p} \\ &= |d_K|. \end{aligned}$$

Recall there are exactly 2^{n-3} symplectic characters of G , so we find

$$\begin{aligned} \prod_{\chi \text{ symplectic}} A(\chi) &= \prod_{p|d_K} p^{2^{n-2}} \\ &\geq |d_K|^{1/4}, \end{aligned}$$

and we find

$$\sum_{\chi \text{ symplectic}} \log A(\chi) \gg \log |d_K|.$$

Finally, a simple Taylor expansion shows that the minimum value of $|\chi(C_1) - \chi(C_2)|$ when χ varies in the set of symplectic irreducible characters of G and $\{C_1, C_2\} \neq \{C_y, C_{xy}\}$ is $\gg \frac{1}{2^n}$, so we conclude that

$$\text{Var}(X(K/\mathbb{Q}, C_1, C_2)) \gg \frac{1}{4^n} \sum_{\chi \text{ symplectic}} \log A(\chi) \gg \frac{\log |d_K|}{4^n}.$$

As for the last lower bound, if $\{C_1, C_2\} \cap \{C_{x^k} \mid 1 \leq k \leq 2^{n-2} - 1\} = \emptyset$ then we actually have $|\chi(C_1) - \chi(C_2)| \gg 1$ for χ irreducible symplectic, so

$$\text{Var}(X(K/\mathbb{Q}, C_1, C_2)) \gg \sum_{\chi \text{ symplectic}} \log A(\chi) \gg \log |d_K|.$$

□

Proposition 4.5. *Assume GRH and LI⁺. If K/\mathbb{Q} is tamely ramified then for any $3 \leq i \leq n$,*

$$\mathbb{E}(X(K/K_i, C_1^{(i)}, C_{-1}^{(i)})) = -2^{n-1}(1 - W_K) + 2^{i-1},$$

where $W_k = W_{k/\mathbb{Q}}$ is the root number of any symplectic character of G .

Proof. From Theorem 1.11 we have

$$\mathbb{E}(X(K/K_i, C_1^{(i)}, C_{-1}^{(i)})) = \frac{|(C_{-1}^{(i)})^{1/2}|}{|C_{-1}|} - \frac{|(C_1^{(i)})^{1/2}|}{|C_1|} + 2 \sum_{\chi \neq \chi_0} (\chi(C_{-1}^{(i)}) - \chi(C_1^{(i)})) \text{ord}_{s=1/2} L(s, \chi, K/K_i).$$

Since we are assuming LI⁺, Proposition 4.3 shows that only symplectic characters, i.e. the $\psi_j^{(i)}$ with $1 \leq j \leq 2^{i-2} - 1$ odd, contribute to the sum. By the same proposition, their root

numbers are all equal to W_K and the order of vanishing of their Artin L -functions is 0 or 2^{n-i} , so

$$2 \sum_{\chi \neq \chi_0} (\chi(C_{-1}^{(i)}) - \chi(C_1^{(i)})) \text{ord}_{s=1/2} L(s, \chi, K/K_i) = \begin{cases} 0 & \text{if } W_K = 1 \\ 2 \sum_{\chi \text{ symplectic}} (-4) \cdot 2^{n-i} = -2^n & \text{if } W_K = -1 \end{cases} .$$

In the proof of Lemma 2.10, we have observed that -1 has $2^{i-1} + 2$ square roots in $\text{Gal}(K/K_i)$, while 1 has 2 . This yields

$$\mathbb{E}(X(K/K_i, C_1^{(i)}, C_{-1}^{(i)})) = \begin{cases} 2^{i-1} & \text{if } W_K = 1 \\ 2^{i-1} - 2^n & \text{if } W_K = -1 \end{cases} .$$

□

For the sake of completeness, we give below an exhaustive list of $\mathbb{E}(X(K/K_i, C_1, C_2))$ for each $3 \leq i \leq n$ and for all possible choices of distinct conjugacy classes C_1, C_2 of G_i .

Proposition 4.6. *Assume GRH and LI⁺. If K/\mathbb{Q} is tamely ramified then for any $3 \leq i \leq n$,*

C_1	C_2	$\mathbb{E}(X(K/K_i, C_1, C_2))$	Conditions on the classes
$C_1^{(i)}$	$C_{-1}^{(i)}$	$-2^{n-1}(1 - W_K) + 2^{i-1}$	none
$C_1^{(i)}$	$C_{x_i^k}^{(i)}$	$-2^{n-2}(1 - W_K) + (-1)^k - 1$	$1 \leq k \leq 2^{i-2} - 1$
$C_1^{(i)}$	$C_y^{(i)}/C_{x_i y}^{(i)}$	$-2^{n-2}(1 - W_K) - 2$	none
$C_{-1}^{(i)}$	$C_{x_i^k}^{(i)}$	$2^{n-2}(1 - W_K) - 1 - (-1)^k - 2^{i-1}$	$1 \leq k \leq 2^{i-2} - 1$
$C_{-1}^{(i)}$	$C_y^{(i)}/C_{x_i y}^{(i)}$	$2^{n-2}(1 - W_K) - 2 - 2^{i-1}$	none
$C_{x_i^k}^{(i)}$	$C_{x_i^l}^{(i)}$	$(-1)^l - (-1)^k$	$1 \leq k, l \leq 2^{i-2} - 1, k \neq l$
$C_{x_i^k}^{(i)}$	$C_y^{(i)}/C_{x_i y}^{(i)}$	$(-1)^{k+1} - 1$	$1 \leq k \leq 2^{i-2} - 1$
$C_y^{(i)}$	$C_{x_i y}^{(i)}$	0	none

Proof. The first row was computed in Proposition 4.5. Each of the sums

$$\sum_{\chi \neq \chi_0} \chi(C_{x_i^k}^{(i)}) \text{ord}_{s=1/2} L(s, \chi, K/K_i)$$

for $1 \leq k \leq 2^{i-2} - 1$ is zero, because Proposition 4.3 shows they only involve symplectic characters, for which $\text{ord}_{s=1/2} L(s, \chi, K/K_i)$ does not depend on χ , and those sums reduce to

$$\text{ord}_{s=1/2} L(s, \psi_1^{(i)}, K/K_i) \sum_{\substack{j=1 \\ j \text{ odd}}}^{2^{i-2}-1} (\zeta_i^{jk} + \zeta_i^{-jk}),$$

which is zero by Lemma 2.13. Therefore using Theorem 1.11, we see that the remaining computations only involves counting square roots in $G_i \simeq \mathbb{H}_{2^i}$, and, when one of the conjugacy classes is $C_1^{(i)}$ or $C_{-1}^{(i)}$, the fact there are 2^{i-3} irreducible symplectic characters of G_i with L -functions vanishing at $1/2$ to order $2^{n-i-1}(1 - W_K)$. The last row is clear because

the elements of $C_y^{(i)}$ and $C_{x_i y}^{(i)}$ have no square root in G_i and symplectic characters vanish on those conjugacy classes. \square

The table of 4.6 that any Chebotarev race involving the classes $C_1^{(i)}$ or $C_{-1}^{(i)}$ is influenced by the root number of symplectic characters of G .

4.2 Moments in dihedral Galois groups of order a power of two

We now turn to the case of number field extensions with dihedral Galois group of 2-power order. The same methods as in the previous section are applied to get bounds on the moments of the random variables attached to the different Chebotarev races in such extensions. Those bounds will be used to prove Theorem B.

Let $n \geq 2$ and let L/\mathbb{Q} be a Galois extension with Galois group $D_{2^{n-1}} = \langle r, s \mid r^{2^{n-1}} = s^2 = 1, srs^{-1} = r^{-1} \rangle$. Let $\mathbb{Q} = L_n \subset L_{n-1} \subset \dots \subset L_3 \subset L$ be the subextensions such that $G_i := \langle r^{2^{n-i}}, s \rangle$ and $L_i = L^{G_i}$. We will denote $r^{2^{n-i}}$ by r_i . Conjugacy classes in G_i will be denoted, following our notations in Lemma 2.5, by $C_1^{(i)}, C_{-1}^{(i)}, C_{r_i^k}^{(i)}, C_s^{(i)}$ and $C_{r_i s}^{(i)}$. The estimates from the previous section are proved similarly in the dihedral case, so we state them without proof. Since each irreducible character of $D_{2^{n-1}}$ is orthogonal by Lemma 2.10, and since we will be working under the hypothesis LI, no considerations on root numbers are involved.

Proposition 4.7. *Assume GRH and LI $^+$. Then for any $2 \leq i \leq n$ and any distinct conjugacy classes C_1, C_2 of G_i such that $\{C_1, C_2\} \neq \{C_s^{(i)}, C_{r_i s}^{(i)}\}$,*

$$\text{Var}(X(L/L_i, C_1, C_2)) \ll \log |d_L|.$$

Moreover if L/\mathbb{Q} is tamely ramified then for any $2 \leq i \leq n$,

$$\text{Var}(X(L/L_i, C_1, C_2)) \gg \frac{\log |d_L|}{4^n},$$

and if $\{C_1, C_2\} \cap \{C_{r_i^k}^{(i)} \mid 1 \leq k \leq 2^{i-2} - 1\} = \emptyset$ this can be improved to

$$\text{Var}(X(L/L_i, C_1, C_2)) \gg \log |d_L|.$$

Proposition 4.8. *Assume GRH and LI. For any $3 \leq i \leq n$,*

C_1	C_2	$\mathbb{E}(X(L/L_i, C_1, C_2))$	Conditions on the classes
$C_1^{(i)}$	$C_{-1}^{(i)}$	$-2^{i-1} + 1$	none
$C_1^{(i)}$	$C_{r_i^k}^{(i)}$	$-2^{i-1} + (-1)^k$	$1 \leq k \leq 2^{i-2} - 1$
$C_1^{(i)}$	$C_s^{(i)}/C_{r_i s}^{(i)}$	$-2^{i-1} - 1$	none
$C_{-1}^{(i)}$	$C_{r_i^k}^{(i)}$	$(-1)^k - 1$	$1 \leq k \leq 2^{i-2} - 1$
$C_{-1}^{(i)}$	$C_s^{(i)}/C_{r_i s}^{(i)}$	-2	none
$C_{r_i^k}^{(i)}$	$C_{r_i^l}^{(i)}$	$(-1)^l - (-1)^k$	$1 \leq k, l \leq 2^{i-2} - 1, k \neq l$
$C_{r_i^k}^{(i)}$	$C_s^{(i)}/C_{r_i s}^{(i)}$	$(-1)^{k+1} - 1$	$1 \leq k \leq 2^{i-2} - 1$
$C_s^{(i)}$	$C_{r_i s}^{(i)}$	0	none

4.3 Construction of the towers

We now construct towers of dihedral and generalized quaternion extensions of \mathbb{Q} which are tamely ramified, and with controlled discriminants, properties which will enable us to apply effectively the results from sections 4.1 and 4.2.

Proposition 4.9. *Assume GRH for the Dedekind zeta functions of the number fields $\mathbb{Q}(\sqrt[5 \cdot 2^{n-1}]{\varepsilon}, \mu_{5 \cdot 2^{n-1}})$, where μ_k denotes a primitive k -th root of unity. There exists a sequence $(\mathcal{D}_n)_{n \geq 3}$ of number fields such that for any $n \geq 3$:*

- i) *The extension \mathcal{D}_n/\mathbb{Q} is tamely ramified and Galois with Galois group isomorphic to the dihedral group $D_{2^{n-1}}$ of order 2^n .*
- ii) $\mathbb{Q}(\sqrt{5}) \subset \mathcal{D}_n$.
- iii) $2^n \ll \log |d_{\mathcal{D}_n/\mathbb{Q}}| \ll n2^n$.

Proof. Fix $K = \mathbb{Q}(\sqrt{5})$, let $n \geq 3$ and let $\varepsilon = \frac{3+\sqrt{5}}{2}$ be the fundamental totally positive unit of K . Using class field theory, it is shown in the proof of [20, Theorem 3.2] that there exists a number field \mathcal{D}_n which has Galois group over \mathbb{Q} isomorphic to $D_{2^{n-1}}$, containing K and such that only 5 and another odd prime p ramify in \mathcal{D}_n . The prime number p is chosen so that p splits in the extension $M_n := \mathbb{Q}(\sqrt[5 \cdot 2^{n-1}]{\varepsilon}, \mu_{5 \cdot 2^{n-1}})$.

Using the bound for the least prime ideal in the Chebotarev density theorem stated in [13, (1.2)], we can find such a p satisfying

$$p \ll (\log |d_{M_n}|)^2.$$

Now, M_n is obtained in at most $2n$ steps of adjoining square roots of algebraic units, starting from the field $\mathbb{Q}(\sqrt[5]{\varepsilon}, \mu_5)$. If M/L is one of those steps, we have $M = L(\sqrt{\eta})$, where η is a unit of L , and the relative discriminant $D_{M/L}$ has to divide the discriminant of $X^2 - \eta$, which is $4\eta\mathcal{O}_L = 4\mathcal{O}_L$. Since

$$|d_M| = N_{L/\mathbb{Q}}(D_{M/L})|d_L|^{[M:L]} \leq 4^{[L:\mathbb{Q}]}|d_L|^2,$$

we find

$$|d_M|^{\frac{1}{[M:\mathbb{Q}]}} \leq 2|d_L|^{\frac{1}{[L:\mathbb{Q}]}}.$$

Using this iteratively on the (at most) $2n$ steps, we find that

$$|d_{M_n}|^{\frac{1}{[M_n:\mathbb{Q}]}} \ll 2^{2n}.$$

Using the same reasoning, we see that

$$[M_n : \mathbb{Q}] \ll 4^n,$$

so finally we can choose the prime p so that

$$p \ll (n[M_n : \mathbb{Q}])^2 \ll n^2 16^n.$$

As in the proof of Proposition 4.4,

$$|d_{\mathcal{D}_n}| = 5^{(e_5-1)f_5g_5} \times p^{(e_p-1)f_pg_p} \leq (5p)^{2^n},$$

where e_q (respectively f_q, g_q) denotes the ramification index of the prime number q (respectively the residual degree of q , the number of primes above q). This shows that

$$\log |d_{\mathcal{D}_n}| \ll 2^n \log p \ll n2^n.$$

The lower bound on the discriminant simply comes from Minkowski's bound ([14] p.120)

$$|d_{\mathcal{D}_n}| \geq \frac{2^{n2^n}}{(2^n)!} \left(\frac{\pi}{4}\right)^{2^{n-1}}.$$

Since $[\mathcal{D}_n : \mathbb{Q}] = 2^n$ and $d_{\mathcal{D}_n}$ is odd, \mathcal{D}_n/\mathbb{Q} is tamely ramified. □

Remark. The quadratic field $\mathbb{Q}(\sqrt{5})$ plays no particular role in our construction. For our purpose, we could replace the integer 5 by any positive square-free integer d which is 1 mod 4.

Proposition 4.10. *Assume GRH. There exists two sequences $(\mathcal{Q}_n^+)_{n \geq 3}$ and $(\mathcal{Q}_n^-)_{n \geq 3}$ of number fields such that for any $n \geq 3$:*

- i) *The extension $\mathcal{Q}_n^\pm/\mathbb{Q}$ is tamely ramified and Galois with Galois group isomorphic to the generalized quaternion group \mathbb{H}_{2^n} .*
- ii) *$\mathbb{Q}(\sqrt{5}) \subset \mathcal{Q}_n^\pm$.*
- iii) *$2^n \ll \log |d_{\mathcal{Q}_n^\pm/\mathbb{Q}}| \ll n2^n$.*
- iv) *$W_{\mathcal{Q}_n^+} = 1$ and $W_{\mathcal{Q}_n^-} = -1$.*

Proof. We use the same notations as in the proof of Proposition 4.9 : we let $K = \mathbb{Q}(\sqrt{5})$ and $\varepsilon = \frac{3+\sqrt{5}}{2}$ be the fundamental totally positive unit of K . We use again a theorem of Fröhlich [8, Chapter V, Proposition 3.1]. It states that for $e \in \{-1, 1\}$, the set of prime numbers p such that there exists a number field $N[p]$ such that the only ramified primes in $N[p]$ are 5 and p , satisfying $\text{Gal}(N[p]/\mathbb{Q}) \simeq \mathbb{H}_8$, $\mathbb{Q}(\sqrt{5}) \subset N[p]$ and $W_{N[p]} = e$ have positive density. It is shown by translating the last two conditions into an arithmetic condition on p , and then using Chebotarev's density theorem. The arithmetic condition amounts to prescribing the Frobenius conjugacy class of p in $\text{Gal}(M'_n/\mathbb{Q})$, where $M'_n = K(\mu_{2^{n-1}}, \sqrt[n]{\varepsilon})$, with $\mu_{2^{n-1}}$ a primitive 2^{n-1} -th root of unity. As in the proof of Proposition 4.9, we apply, under GRH, the bound on the least prime ideal in Chebotarev's density theorem of [13] to choose

$$p \ll (\log |d_{M'_n}|)^2 \ll n^2 16^n.$$

Call \mathcal{Q}_n^\pm the number field constructed this way (the superscript \pm indicating which root number was prescribed). Since only 5 and p ramify in \mathcal{Q}_n^\pm , we get as in the proof of Proposition 4.9

$$\log |d_{\mathcal{Q}_n^\pm/\mathbb{Q}}| \ll n2^n.$$

The lower bound on the discriminant follows once again from Minkowski's bound, and tame ramification follows from the fact that $d_{\mathbb{Q}_n^\pm}$ is odd. \square

Remark. If we do not want to specify the root numbers in our quaternion extensions of \mathbb{Q} , we could use the methods of [2] to construct extensions of \mathbb{Q} satisfying i), ii) and iii) using the extensions \mathcal{D}_n/\mathbb{Q} of Proposition 4.9. Indeed, it is shown in [2] that, if Q/\mathbb{Q} is a tamely ramified extension with Galois group \mathbb{H}_8 and $\mathbb{Q}(\sqrt{5}) \subset Q$ (again, the number 5 has no particular significance), then the composite field $Q\mathcal{D}_n$ contains a subfield Q_n with Galois group \mathbb{H}_{2^n} over \mathbb{Q} , and the upper bound on $\log |d_{\mathcal{D}_n}|$ then implies a similar bound on $\log |d_{Q_n}|$. Tame ramification follows from the tame ramification of Q/\mathbb{Q} and \mathcal{D}_n/\mathbb{Q} , and the lower bound on the discriminant from Minkowski's bound.

4.4 A large deviation result for Chebyshev's bias

In this section we prove a lower bound in the context of Theorem 1.13. We will make use of the following bounds on sums of zeros of Artin L -functions. The first one is the so-called Riemann-Von Mangoldt formula, stated in [10], in the particular case of Artin L -functions, for which the conductor of $s \mapsto L(s, \chi, L/K)$ is simply $A(\chi)$ and its analytic conductor is bounded by $A(\chi)(|s| + 4)^{[K:\mathbb{Q}]\chi(1)}$ (see [10, §5.13]).

Lemma 4.11. *Let χ be an irreducible character of G , and for any $T > 0$, define $N(T, \chi)$ to be the number of zeros $\rho = \beta + i\gamma$ of $s \mapsto L(s, \chi, L/K)$ with $0 \leq \beta \leq 1$ and $0 < \gamma \leq T$. Then we have*

$$N(T, \chi) = \frac{T}{2\pi} \log \left(A(\chi) \left(\frac{T}{2\pi e} \right)^{[K:\mathbb{Q}]\chi(1)} \right) + O \left(\log \left(A(\chi)(T + 4)^{[K:\mathbb{Q}]\chi(1)} \right) \right).$$

Proof. This is essentially [10, Theorem 5.8], except that we are counting zeros $\beta + i\gamma$ with $0 < \gamma \leq T$ and not $|\gamma| \leq T$. Our main term is half the one in [10, Theorem 5.8], in which the count is made by adding the contributions of zeros $\beta + i\gamma$ with $\gamma \geq 0$ for $s \mapsto L(s, \chi, L/K)$ and for $s \mapsto L(s, \bar{\chi}, L/K)$. It is also stated in the proof that the number of real zeros is taken into account in the error term. \square

Lemma 4.12 ([5], Lemma 5.4). *Assume Artin's conjecture. Then for any irreducible character χ of G and for $T \geq 1$, we have*

$$\sum_{0 < \gamma_\chi \leq T} \frac{1}{\sqrt{\frac{1}{4} + \gamma_\chi^2}} = \frac{\log T}{2\pi} \log \left(A(\chi) \left(\frac{T^{1/2}}{2\pi e} \right)^{[K:\mathbb{Q}]\chi(1)} \right) + O \left(\log \left(A(\chi)(T + 4)^{[K:\mathbb{Q}]\chi(1)} \right) \right).$$

In order to state our improvement of Theorem 1.13, we first recall the large deviation result of Montgomery and Odlyzko which was used to derive Theorem 1.13, and will be used to prove Theorem 4.14.

Theorem 4.13 ([15], Theorem 2). *Let $(W_n)_{n \geq 1}$ be a family of independent real random variables such that*

i) For all $n \geq 1$, $\mathbb{E}(W_n) = 0$.

ii) For all $n \geq 1$, $|W_n| \leq 1$ a.s.

iii) There exists $c > 0$ such that for all $n \geq 1$, $\mathbb{E}(W_n^2) > c$.

Let $(r_n)_n$ be a real sequence decreasing to zero such that $\sum_{n \geq 1} r_n^2 < +\infty$ and let $W = \sum_{n \geq 1} r_n W_n$.

Let $V \geq 0$ and $\alpha > 0$. If $\sum_{r_n \geq \alpha} r_n \leq \frac{V}{2}$ then

$$\mathbb{P}(W \geq V) \leq \exp\left(-\frac{1}{16}V^2\left(\sum_{r_n < \alpha} r_n^2\right)^{-1}\right),$$

If $\sum_{r_n \geq \alpha} r_n \geq 2V$ then

$$\mathbb{P}(W \geq V) \geq a_1 \exp\left(-a_2V^2\left(\sum_{r_n < \alpha} r_n^2\right)^{-1}\right),$$

where a_1 and a_2 are positive constants depending only on c .

The following theorem provides a lower bound for $1 - \delta$, which was not obtained in [5], in the context of Theorem 1.13. The proof proceeds by taking into account distinct characters of G whose corresponding induced characters to G^+ are not orthogonal, which in general leads to complications in estimating Chebyshev's bias. This is because, in Theorem 1.11, the means are expressed using values of characters of G , while the variances involve values of characters of G^+ .

Before stating the result, we set a few notations for readability. If $\chi \in \text{Irr}(G)$ and $\lambda \in \text{Irr}(G^+)$ we will write $\lambda \mid \chi$ if $\langle \lambda, \text{Ind}_G^{G^+} \chi \rangle \neq 0$, that is if λ is a component of the character $\text{Ind}_G^{G^+} \chi$ of G^+ , and $\lambda \nmid \chi$ otherwise. For $\chi, \chi' \in \text{Irr}(G)$, we write $(\chi, \chi') = 1$ when $\langle \text{Ind}_G^{G^+} \chi, \text{Ind}_G^{G^+} \chi' \rangle = 0$, i.e. when $\lambda \mid \chi$ implies $\lambda \nmid \chi'$.

Theorem 4.14. *Assume GRH, LI and Artin's Conjecture. Let*

$$S = \{\chi \in \text{Irr}(G) \mid \forall \chi' \in \text{Irr}(G) \setminus \{\chi\}, (\chi, \chi') = 1\}$$

and $R = \text{Irr}(G) \setminus (S \cup \{\chi_0\})$. Let $b_1 := \max_{\chi \in R} \chi(1)$, $b_2 := |R|$ and $M = \max_{\chi \in R} \text{ord}_{s=1/2} L(s, \chi, L/K) + 1$. There exist absolute constants $c_1, c_2 > 0$ such that for any conjugacy classes C_1, C_2 of G satisfying $C_1^+ \neq C_2^+$ and $B(L/K, C_1, C_2) > 0$, we have

$$c_1 \exp\left(-c_2 Q(C_1, C_2) B(L/K, C_1, C_2)^2\right) \leq 1 - \delta(L/K, C_1, C_2)$$

where $Q(C_1, C_2) = \max\left(e^C \sqrt{\frac{M b_1 b_2}{\lambda^*(1) b_3(C_1, C_2)}}, C^{\frac{b_3(C_1, C_2)}{b_4(C_1, C_2)}}, C\right)$ for some absolute constant $C > 0$

and where $\lambda^* \in \text{Irr}(G^+)$ is such that $|\lambda^*(C_2^+) - \lambda^*(C_1^+)| = \max_{\lambda \in \text{Irr}(G^+)} |\lambda(C_2^+) - \lambda(C_1^+)| =: b_3(C_1, C_2) > 0$ and $b_4(C_1, C_2) = \min_{\lambda \in \text{Irr}(G^+), \lambda(C_2^+) \neq \lambda(C_1^+)} |\lambda(C_2^+) - \lambda(C_1^+)| > 0$.

Proof. We introduce the random variable $W = X(L/K, C_1, C_2) - \mathbb{E}(X(L/K, C_1, C_2))$. It obviously satisfies the hypotheses of Theorem 4.13 with $(r_n)_n$ the ordered sequence of non-zero $\frac{2|\lambda(C_2^+) - \lambda(C_1^+)|}{\sqrt{\frac{1}{4} + \gamma_\lambda^2}}$, λ varying in $\text{Irr}(G^+)$ and γ_λ in $\Gamma_{L/\mathbb{Q}, \lambda}$, and $(W_n)_n$ the accordingly ordered sequence of X_γ . Indeed, the series

$$\sum_{\gamma_\lambda > 0} \frac{1}{\frac{1}{4} + \gamma_\lambda^2}$$

converges for any $\lambda \in \text{Irr}(G^+)$.

We set $V = \mathbb{E}(X(L/K, C_1, C_2))$ and look for $\alpha > 0$ such that $\sum_{r_n \geq \alpha} r_n \geq 2V$. By Theorem 1.11, we have

$$\begin{aligned} \mathbb{E}(X(L/K, C_1, C_2)) &= \frac{|C_2^{1/2}|}{|C_2|} - \frac{|C_1^{1/2}|}{|C_1|} + z(C_2) - z(C_1) \\ &= \sum_{\chi \in \text{Irr}(G)} (\chi(C_2) - \chi(C_1))(\varepsilon_2(\chi) + 2 \text{ord}_{s=1/2} L(s, \chi, L/K)). \end{aligned}$$

If $\chi \in S$, then for some $\lambda \in \text{Irr}(G^+)$ we have, by a quick computation using the Frobenius reciprocity formula,

$$\begin{aligned} |\lambda(C_2^+) - \lambda(C_1^+)| &= |\lambda|_G(C_2) - \lambda|_G(C_1)| = \left| \sum_{\chi' \in \text{Irr}(G)} (\chi'(C_2) - \chi'(C_1)) \langle \lambda|_G, \chi' \rangle \right| \\ &= |\chi(C_2) - \chi(C_1)| \langle \lambda, \text{Ind}_G^{G^+} \chi \rangle. \end{aligned}$$

From the factorisation

$$L(s, \chi, L/K) = \prod_{\lambda \in \text{Irr}(G^+)} L(s, \lambda, L/\mathbb{Q})^{\langle \lambda, \text{Ind}_G^{G^+} \chi \rangle},$$

we see that

$$\text{ord}_{s=1/2} L(s, \chi, L/K) = \sum_{\lambda \in \text{Irr}(G^+)} \langle \lambda, \text{Ind}_G^{G^+} \chi \rangle \text{ord}_{s=1/2} L(s, \lambda, L/\mathbb{Q}) \leq M_0 \sum_{\lambda \in \text{Irr}(G^+)} \langle \lambda, \text{Ind}_G^{G^+} \chi \rangle$$

(recall that M_0 comes from the hypothesis L1). Therefore,

$$\begin{aligned} \sum_{\chi \in S} (\chi(C_2) - \chi(C_1))(\varepsilon_2(\chi) + 2 \text{ord}_{s=1/2} L(s, \chi, L/K)) &\leq \sum_{\chi \in S} |\chi(C_2) - \chi(C_1)| (1 + 2M_0 \sum_{\lambda | \chi} \langle \lambda, \text{Ind}_G^{G^+} \chi \rangle) \\ &= (1 + 2M_0) \sum_{\lambda \in S^+} |\lambda(C_2^+) - \lambda(C_1^+)| \end{aligned}$$

where $S^+ = \{\lambda \in \text{Irr}(G^+) \mid \exists \chi \in S, \lambda \mid \chi\}$. We now see that it is enough to have

$$\sum_{0 < \gamma_\lambda \leq T_0(\lambda)} \frac{1}{\sqrt{\frac{1}{4} + \gamma_\lambda^2}} \geq (2 + 4M_0)$$

for any $\lambda \in S^+$ and some $T_0(\lambda) > 0$, to bound from above the contribution of characters of S in $2V$. It remains to bound from above the sum

$$2 \sum_{\chi \in R} (\chi(C_2) - \chi(C_1)) (\varepsilon(\chi) + 2 \operatorname{ord}_{s=1/2} L(s, \chi, L/K)) \leq 4M \sum_{\chi \in R} |\chi(C_2) - \chi(C_1)|.$$

By definition of b_1 and b_2 , this last sum is $\leq 8Mb_1b_2$, so we choose $T_0(\lambda^*) > 0$ such that

$$\sum_{0 < \gamma_{\lambda^*} \leq T_0(\lambda^*)} \frac{1}{\sqrt{\frac{1}{4} + \gamma_{\lambda^*}^2}} \geq \frac{8Mb_1b_2}{b_3(C_1, C_2)}.$$

To do so, we use Lemma 4.12 (for the extension L/\mathbb{Q}) to get

$$\sum_{0 < \gamma_{\lambda} \leq T} \frac{1}{\sqrt{\frac{1}{4} + \gamma_{\lambda}^2}} = \frac{\log T}{2\pi} \log \left(A(\lambda) \left(\frac{T^{1/2}}{2\pi e} \right)^{\lambda(1)} \right) + O \left(\log \left(A(\lambda) (T+4)^{\lambda(1)} \right) \right)$$

for any $\lambda \in \operatorname{Irr}(G^+)$. A simple computation shows that it is enough to choose $T_0(\lambda^*) = \max \left((2\pi e)^4, e \sqrt{\frac{64\pi M b_1 b_2}{\lambda^*(1) b_3(C_1, C_2)}}, e^{16\pi C'} \right)$ to ensure

$$\sum_{0 < \gamma_{\lambda^*} < T_0(\lambda^*)} \frac{1}{\sqrt{\frac{1}{4} + \gamma_{\lambda^*}^2}} \geq \frac{8Mb_1b_2}{b_3(C_1, C_2)},$$

where $C' > 0$ is such that

$$\left| \sum_{0 < \gamma_{\lambda^*} \leq T} \frac{1}{\sqrt{\frac{1}{4} + \gamma_{\lambda^*}^2}} - \frac{\log T}{2\pi} \log \left(A(\lambda^*) \left(\frac{T^{1/2}}{2\pi e} \right)^{\lambda^*(1)} \right) \right| \leq C' \log \left(A(\lambda^*) (T+4)^{\lambda^*(1)} \right),$$

while we have

$$\sum_{0 < \gamma_{\lambda} \leq A} \frac{1}{\sqrt{\frac{1}{4} + \gamma_{\lambda}^2}} \geq (2 + 4M_0)$$

for any $\lambda \in \operatorname{Irr}(G^+)$ and some absolute constant $A > 0$, so we set $T_0(\lambda) = A$ for every $\lambda \neq \lambda^*$.

We now turn to the choice of $\alpha \geq 0$ such that

$$\sum_{r_n \geq \alpha} r_n \geq 2V.$$

It is enough to have

$$\sum_{r_n \geq \alpha} r_n \geq \sum_{\lambda \in \operatorname{Irr}(G^+)} \sum_{0 < \gamma_{\lambda} \leq T_0(\lambda)} \frac{2|\lambda(C_2^+) - \lambda(C_1^+)|}{\sqrt{\frac{1}{4} + \gamma_{\lambda}^2}},$$

so we require that

$$0 < \gamma_{\lambda} \leq T_0(\lambda) \Rightarrow \gamma_{\lambda} \leq \sqrt{\frac{4|\lambda(C_2^+) - \lambda(C_1^+)|^2}{\alpha^2} - \frac{1}{4}} =: R_{\alpha, \lambda}.$$

We choose α to be the minimal (positive) value of $\frac{2|\lambda(C_2^+) - \lambda(C_1^+)|}{\sqrt{\frac{1}{4} + T_0(\lambda)^2}}$, attained for some $\lambda_m \in \text{Irr}(G^+)$ (recall that we discarded the characters $\lambda \in \text{Irr}(G^+)$ such that $\lambda(C_2^+) = \lambda(C_1^+)$ in our definition of the r_n 's). For this choice of α we therefore have $\sum_{r_n \geq \alpha} r_n \geq 2V$.

Theorem 4.13 now yields

$$\mathbb{P}(W \geq V) \geq a_1 \exp \left(-a_2 V^2 \left(\sum_{r_n < \alpha} r_n^2 \right)^{-1} \right).$$

We have

$$\begin{aligned} \sum_{r_n < \alpha} r_n^2 &= \sum_{\lambda \in \text{Irr}(G^+)} \sum_{\gamma_\lambda > R_{\alpha, \lambda}} \frac{4|\lambda(C_2^+) - \lambda(C_1^+)|^2}{\frac{1}{4} + \gamma_\lambda^2} \\ &\geq \sum_{\lambda \in \text{Irr}(G^+)} \sum_{2R_{\alpha, \lambda} \geq \gamma_\lambda > R_{\alpha, \lambda}} \frac{4|\lambda(C_2^+) - \lambda(C_1^+)|^2}{\frac{1}{4} + \gamma_\lambda^2} \\ &\geq \sum_{\lambda \in \text{Irr}(G^+)} \frac{4|\lambda(C_2^+) - \lambda(C_1^+)|^2}{\frac{1}{4} + 4R_{\alpha, \lambda}^2} (N(2R_{\alpha, \lambda}, \lambda) - N(R_{\alpha, \lambda}, \lambda)) \end{aligned}$$

As in the proof of [5, Lemma 4.3] we see, using Lemma 4.12, that

$$N(2R_{\alpha, \lambda}, \lambda) - N(R_{\alpha, \lambda}, \lambda) \gg R_{\alpha, \lambda} \log A(\lambda).$$

Thus,

$$\sum_{r_n < \alpha} r_n^2 \gg \frac{1}{\max_{\lambda \in \text{Irr}(G^+)} R_{\alpha, \lambda}} \sum_{\lambda \in \text{Irr}(G^+)} |\lambda(C_2^+) - \lambda(C_1^+)|^2 \log A(\lambda).$$

Combining Theorem 1.11 and Lemma 2.3, we finally get

$$\sum_{r_n < \alpha} r_n^2 \gg \frac{1}{\max_{\lambda \in \text{Irr}(G^+)} R_{\alpha, \lambda}} \text{Var}(X(L/K, C_1, C_2)).$$

We can now conclude that

$$\mathbb{P}(W \geq V) \geq a_1 \exp \left(-a_3 \left(\max_{\lambda \in \text{Irr}(G^+)} R_{\alpha, \lambda} \right) B(L/K, C_1, C_2)^2 \right)$$

for some absolute constant $a_3 > 0$. But, recalling the choice we made for α ,

$$\max_{\lambda \in \text{Irr}(G^+)} R_{\alpha, \lambda} = \sqrt{\left(\frac{1}{4} + T_0(\lambda_m)^2 \right) \frac{|\lambda^*(C_2^+) - \lambda^*(C_1^+)|^2}{|\lambda_m(C_2^+) - \lambda_m(C_1^+)|^2}}.$$

To determine the size of this quantity, we consider two cases : either $\lambda_m = \lambda^*$, in which case we find $\max_{\lambda \in \text{Irr}(G^+)} R_{\alpha, \lambda} \asymp T_0(\lambda^*) \asymp e^C \sqrt{\frac{Mb_1b_2}{\lambda^*(1)b_3(C_1, C_2)}}$, or $\lambda_m \neq \lambda^*$, in which case $\max_{\lambda \in \text{Irr}(G^+)} R_{\alpha, \lambda} \asymp \frac{b_3(C_1, C_2)}{b_4(C_1, C_2)}$ since $T_0(\lambda_m)$ was chosen as an absolute constant.

It simply remains to note that, by symmetry of W about 0, we have

$$\mathbb{P}(W \geq V) = \mathbb{P}(W \leq -V) = 1 - \mathbb{P}(X(L/K, C_1, C_2) > 0) = 1 - \delta(L/K, C_1, C_2).$$

□

Note that when $K = \mathbb{Q}$ in the previous theorem, we have $R = \emptyset$ so that $Q(C_1, C_2)$ can be taken to be $C \left(\frac{b_3(C_1, C_2)}{b_4(C_1, C_2)} + 1 \right)$, where $C > 0$ is an absolute constant. In our applications (Theorems 4.15 and 4.16), the quantities b_1, b_2, M and $\frac{b_3(C_1, C_2)}{b_4(C_1, C_2)}$ will be bounded.

4.5 Estimates on the bias in the towers

Using the number fields constructed in Proposition 4.9 and Proposition 4.10, we can now state the following result, a more exhaustive version of Theorem B.

Theorem 4.15. *Assume GRH and LI⁺ for the number fields \mathcal{D}_n of Proposition 4.9. There exist absolute constants $c_1, c_2, c_3 > 0$ such that for any $n \geq 3$, the following hold :*

C_a	C_b	Estimate on $\delta(\mathcal{D}_n/\mathbb{Q}, C_1, C_2)$	Condition on the classes
C_1	C_{-1}, C_s, C_{rs}	$c_1 \exp(-c_2 2^n) < \delta(\mathcal{D}_n/\mathbb{Q}, C_1, C_b) < \exp\left(-c_3 \frac{2^n}{n}\right)$	none
C_1	C_{r^k}	$c_1 \exp(-c_3 8^n) < \delta(\mathcal{D}_n/\mathbb{Q}, C_1, C_{r^k}) < \exp\left(-c_4 \frac{2^n}{n}\right)$	none
C_{-1}	C_{r^k}	$\frac{1}{2}$	k even
C_{-1}	C_s, C_{rs}	$0 < \frac{1}{2} - \delta(\mathcal{D}_n/\mathbb{Q}, C_{-1}, C_b) \ll \frac{1}{2^{n/3}}$	none
C_{r^k}	C_{r^l}	$\frac{1}{2}$	$k = l \pmod{2}$
C_{r^k}	C_s, C_{rs}	$\frac{1}{2}$	k odd
C_s	C_{rs}	$\frac{1}{2}$	none

Proof. For the first part of the theorem, the bounds obtained in Proposition 4.7 (for the variance), Proposition 4.8 (for the mean) and Proposition 4.9 (for the discriminant) show that

$$\frac{2^n}{n} \ll B(\mathcal{D}_n/\mathbb{Q}, C_1, C_b)^2 = \frac{\mathbb{E}(X(\mathcal{D}_n/\mathbb{Q}, C_1, C_b))^2}{\text{Var}(X(\mathcal{D}_n/\mathbb{Q}, C_1, C_b))} \ll 2^n.$$

We combine this estimate with Theorems 1.13 and 4.14 (and the fact $\mathbb{E}(X(\mathcal{D}_n/\mathbb{Q}, C_1, C_b)) < 0$) to get

$$c_1 \exp(-c_2 2^n) < \delta(\mathcal{D}_n/\mathbb{Q}, C_1, C_{-1}) < \exp\left(-c_3 \frac{2^n}{n}\right),$$

for some absolute constants $c_1, c_2, c_3 > 0$ (which may differ from the constants of Theorems 1.13 and 4.14, but by an absolute factor).

The proof is similar for $\delta(\mathcal{D}_n/\mathbb{Q}, C_1, C_{r^k})$, the only difference being the lower bound

$$\text{Var}(X(\mathcal{D}_n/\mathbb{Q}, C_1, C_{r^k})) \gg \frac{\log |d_{\mathcal{D}_n}|}{4^n} \gg \frac{1}{2^n}$$

from Proposition 4.7 and 4.9. To deal with $\delta(\mathcal{D}_n/\mathbb{Q}, C_{-1}, C_b)$, $b = s$ or $b = rs$, we simply apply Theorem 1.14, together with Proposition 4.7.

Finally, each case in which $\delta(C_1, C_2, L/\mathbb{Q}) = \frac{1}{2}$ comes from the fact that $\mathbb{E}(X(C_1, C_2, L/\mathbb{Q})) = 0$. \square

Remark. We could not produce estimates for $\delta(\mathcal{D}_n/\mathbb{Q}, C_{r^k}, C_b)$ for $b = -1$ (when k is odd), $b = r^l$ (when k and l do not have the same parity) and $b = s$ or $b = rs$ (when k is even) because we do not have good enough bounds on $\text{Var}(X(\mathcal{D}_n/\mathbb{Q}, C_{r^k}, C_b))$ to conclude that $B(\mathcal{D}_n/\mathbb{Q}, C_{r^k}, C_b)$ is large or not.

We now turn our attention to the extensions $\mathcal{Q}_n^\pm/\mathbb{Q}$ built in Proposition 4.10. The proof of the next theorem is the same as for Theorem 4.15, except that the value of $W_{\mathcal{Q}_n^\pm}$ determines in some cases the class towards which there is a bias. This is a more exhaustive version of Theorem C.

Theorem 4.16. *Assume GRH and LI⁺ for the number fields \mathcal{Q}_n^\pm of Proposition 4.10. There exist absolute constants $c_1, c_2, c_3 > 0$ such that for any $n \geq 3$, denoting \mathcal{Q}_n^\pm by \mathcal{Q}_n , the following hold :*

C_a	C_b	Estimate on $\delta(\mathcal{Q}_n/\mathbb{Q}, C_a, C_b)$	Conditions
C_{-1}	C_1	$c_1 \exp(-c_2 2^n) < \left \frac{1-W_{\mathcal{Q}_n}}{2} - \delta(\mathcal{Q}_n/\mathbb{Q}, C_{-1}, C_b) \right < \exp\left(-c_3 \frac{2^n}{n}\right)$	none
C_{-1}	C_y, C_{xy}	$c_1 \exp(-c_2 2^n) < \delta(\mathcal{Q}_n/\mathbb{Q}, C_{-1}, C_b) < \exp\left(-c_3 \frac{2^n}{n}\right)$	$W_{\mathcal{Q}_n} = 1$
C_{-1}	C_y, C_{xy}	$0 < \frac{1}{2} - \delta(\mathcal{Q}_n/\mathbb{Q}, C_{-1}, C_b) \ll \frac{1}{2^{n/3}}$	$W_{\mathcal{Q}_n} = -1$
C_{-1}	C_{x^k}	$c_1 \exp(-c_2 8^n) < \delta(\mathcal{Q}_n/\mathbb{Q}, C_{-1}, C_{x^k}) < \exp\left(-c_3 \frac{2^n}{n}\right)$	$W_{\mathcal{Q}_n} = 1$
C_{-1}	C_{x^k}	$\delta(\mathcal{Q}_n/\mathbb{Q}, C_{-1}, C_{x^k}) = \frac{1}{2}$	$W_{\mathcal{Q}_n} = -1$ and k odd
C_1	C_{x^k}	$\frac{1}{2}$	$W_{\mathcal{Q}_n} = 1$ and k even
C_1	C_{x^k}	$c_1 \exp(-c_2 8^n) < \delta(\mathcal{Q}_n/\mathbb{Q}, C_1, C_{x^k}) < \exp\left(-c_3 \frac{2^n}{n}\right)$	$W_{\mathcal{Q}_n} = -1$
C_1	C_y, C_{xy}	$0 < \frac{1}{2} - \delta(\mathcal{Q}_n/\mathbb{Q}, C_1, C_b) \ll \frac{1}{2^{n/3}}$	$W_{\mathcal{Q}_n} = 1$
C_1	C_y, C_{xy}	$c_1 \exp(-c_2 2^n) < \delta(\mathcal{Q}_n/\mathbb{Q}, C_1, C_b) < \exp\left(-c_3 \frac{2^n}{n}\right)$	$W_{\mathcal{Q}_n} = -1$
C_{x^k}	C_{x^l}	$\frac{1}{2}$	$k = l \pmod{2}$
C_{x^k}	C_y, C_{xy}	$\frac{1}{2}$	k odd
C_y	C_{xy}	$\frac{1}{2}$	none

Remark. As in Theorem 4.15, we could not produce bounds for $\delta(\mathcal{Q}_n/\mathbb{Q}, C_{x^k}, C_b)$ for $b = 1$ (when k is odd and $W_{\mathcal{Q}_n} = 1$), $b = -1$ (when k is even and $W_{\mathcal{Q}_n} = -1$), $b = x^l$ (when k and l do not have the same parity) and $b = y$ or $b = xy$ (when k is even).

We now prove a more general version of Theorem D : we are able to observe monotonicity in the evolution of the bias in the subextensions of \mathcal{D}_n/\mathbb{Q} and $\mathcal{Q}_n^+/\mathbb{Q}$.

Theorem 4.17. *Assume GRH and LI⁺. For any $n \geq 3$ and $3 \leq i \leq n$, let $\mathcal{D}_n^{(i)} = \mathcal{D}_n^{(r_i, s)}$ as in section 4.2, and $(\mathcal{Q}_n^+)^{(i)} = (\mathcal{Q}_n^+)^{(x_i, y)}$ as in section 4.1. Then for any $\varepsilon > 0$ and any sufficiently large n , for $3 \leq i < j \leq n$ such that $i \leq n \frac{1+\varepsilon}{2}$ and $j \geq n \left(\frac{1+3\varepsilon}{2}\right)$, we have*

$$\delta(\mathcal{D}_n/\mathcal{D}_n^{(j)}, C_1^{(j)}, C_{-1}^{(j)}) < \delta(\mathcal{D}_n/\mathcal{D}_n^{(i)}, C_1^{(i)}, C_{-1}^{(i)}),$$

$$1 - \delta(\mathcal{Q}_n^+ / (\mathcal{Q}_n^+)^{(j)}, C_1^{(j)}, C_{-1}^{(j)}) < 1 - \delta(\mathcal{Q}_n^+ / (\mathcal{Q}_n^+)^{(i)}, C_1^{(i)}, C_{-1}^{(i)})$$

and

$$\delta(\mathcal{Q}_n^- / (\mathcal{Q}_n^-)^{(j)}, C_1^{(j)}, C_{-1}^{(j)}) < \delta(\mathcal{Q}_n^- / (\mathcal{Q}_n^-)^{(i)}, C_1^{(i)}, C_{-1}^{(i)}).$$

Proof. We combine the bounds of Propositions 4.7, 4.8 and 4.9, with Theorems 1.13 and 4.14. With notations from Theorem 4.14, it is easy to see from Corollary 2.12 that $R = \{\chi_2, \chi_3\}$ so that $b_1 = b_2 = 2$, $M = 1$, and $\frac{b_3(C_1, C_{-1})}{b_4(C_1, C_{-1})} = 1$. We thus find

$$c_1 \exp(-c_2 2^{2i-n}) < \delta(\mathcal{D}_n / \mathcal{D}_n^{(i)}, C_1^{(i)}, C_{-1}^{(i)}) < \exp\left(-c_3 \frac{2^{2i-n}}{n}\right)$$

for any $3 \leq i \leq n$ and for some absolute $c_1, c_2, c_3 > 0$.

In order to have $\delta(\mathcal{D}_n / \mathcal{D}_n^{(j)}, C_1^{(j)}, C_{-1}^{(j)}) < \delta(\mathcal{D}_n / \mathcal{D}_n^{(i)}, C_1^{(i)}, C_{-1}^{(i)})$, it is therefore enough to have

$$\exp\left(-c_3 \frac{2^{2j-n}}{n}\right) < c_1 \exp(-c_2 2^{2i-n}).$$

This is equivalent to

$$c_2 2^{2i} < c_3 \frac{2^{2j}}{n} + 2^n \log(c_1).$$

Now if $j \geq n \left(\frac{1+3\varepsilon}{2}\right)$ and n is large enough then $c_3 \frac{2^{2j}}{n} \geq c_3 \frac{2^{n(1+3\varepsilon)}}{n} > 2^{n(1+2\varepsilon)}$ while if $i \leq n \left(\frac{1}{2} + \varepsilon\right)$ and n is large enough we have $c_2 2^{2i} + 2^n \log(1/c_1) \leq 2^{n(1+2\varepsilon)}$ and the desired inequality holds.

The proof is similar in the case of \mathcal{Q}_n^\pm by using the bounds of Proposition 4.4, Proposition 4.6 and Proposition 4.10 with Theorems 1.13 and 4.14. \square

Remark. A similar proof can be applied to the other extremely biased Chebotarev races from Theorems 4.15 and 4.16, which yields similar monotonicity results.

References

- [1] J. V. Armitage, *Zeta functions with a zero at $s = \frac{1}{2}$* , *Invent. Math.* **15** (1972), 199–205.
- [2] Pierre Damey and Jacques Martinet, *Plongement d’une extension quadratique dans une extension quaternionienne*, *J. Reine Angew. Math.* **262/263** (1973), 323–338. Collection of articles dedicated to Helmut Hasse on his seventy-fifth birthday.
- [3] Alain Debreil, *Groupes finis et treillis de leurs sous-groupes*, *Mathématiques en devenir*, Calvage et Mounet, 2016.
- [4] Lucile Devin, *Chebyshev’s bias for analytic L-functions*, *Mathematical proceedings of the cambridge philosophical society*, 2019, pp. 1–38.
- [5] Daniel Fiorilli and Florent Jouve, *Distribution of Frobenius elements in families of Galois extensions* (January 15, 2020), available at <http://arxiv.org/abs/2001.05428v1>.
- [6] A. Fröhlich, *Artin root numbers and normal integral bases for quaternion fields*, *Invent. Math.* **17** (1972), 143–166.
- [7] ———, *The Galois module structure of algebraic integer rings in fields with generalised quaternion group* (1974), 81–86. *Bull. Soc. Math. France Mém.* 27.
- [8] ———, *Galois module structure of algebraic integers*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, vol. 1, Springer-Verlag, Berlin, 1983.
- [9] I. Martin Isaacs, *Character theory of finite groups*, Dover Publications, Inc., New York, 1994. Corrected reprint of the 1976 original [Academic Press, New York; MR0460423 (57 #417)].
- [10] Henryk Iwaniec and Emmanuel Kowalski, *Analytic number theory*, *American Mathematical Society Colloquium Publications*, vol. 53, American Mathematical Society, Providence, RI, 2004.
- [11] Christian U. Jensen, Arne Ledet, and Noriko Yui, *Generic polynomials*, *Mathematical Sciences Research Institute Publications*, vol. 45, Cambridge University Press, Cambridge, 2002. Constructive aspects of the inverse Galois problem.
- [12] Jerzy Kaczorowski, *On the distribution of primes (mod 4)*, *Analysis* **15** (1995), no. 2, 159–171.
- [13] J. C. Lagarias, H. L. Montgomery, and A. M. Odlyzko, *A bound for the least prime ideal in the Chebotarev density theorem*, *Invent. Math.* **54** (1979), no. 3, 271–296.
- [14] Serge Lang, *Algebraic number theory*, Second, *Graduate Texts in Mathematics*, vol. 110, Springer-Verlag, New York, 1994.
- [15] Hugh L. Montgomery and Andrew M. Odlyzko, *Large deviations of sums of independent random variables*, *Acta Arith.* **49** (1988), no. 4, 427–434.
- [16] M. Ram Murty and V. Kumar Murty, *Non-vanishing of L-functions and applications*, *Modern Birkhäuser Classics*, Birkhäuser/Springer Basel AG, Basel, 1997. [2011 reprint of the 1997 original] [MR1482805].
- [17] Jürgen Neukirch, *Algebraic number theory*, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 322, Springer-Verlag, Berlin, 1999. Translated from the 1992 German original and with a note by Norbert Schappacher, With a foreword by G. Harder.
- [18] Nathan Christopher Ng, *Limiting distributions and zeros of artin l-functions*, Ph.D. Thesis, 2000.
- [19] A. M. Odlyzko, *Bounds for discriminants and related estimates for class numbers, regulators and zeros of zeta functions: a survey of recent results*, *Sém. Théor. Nombres Bordeaux (2)* **2** (1990), no. 1, 119–141.
- [20] Bernat Plans, *On the minimal number of ramified primes in some solvable extensions of \mathbb{Q}* , *Pacific J. Math.* **215** (2004), no. 2, 381–391.
- [21] Michael Rubinstein and Peter Sarnak, *Chebyshev’s bias*, *Experiment. Math.* **3** (1994), no. 3, 173–197.

- [22] Zeév Rudnick and Peter Sarnak, *Zeros of principal L-functions and random matrix theory*, Duke Math. J. **81** (1996), no. 2, 269–322. A celebration of John F. Nash, Jr.
- [23] Jean-Pierre Serre, *Corps locaux*, Hermann, 1980.
- [24] Dane Skabelund, *Character Tables of Metacyclic Groups*, Ph.D. Thesis, 2013.
- [25] Hiraku Tôyama, *A note on the different of the composed field*, Kodai mathematical seminar reports, 1955, pp. 43–44.
- [26] Aurel Wintner, *On the distribution function of the remainder term of the prime number theorem*, Amer. J. Math. **63** (1941), 233–248.

Alexandre Bailleul, UNIV. BORDEAUX, IMB, UMR 5251, F 33405, TALENCE, FRANCE
Email address: alexandre.bailleul@math.u-bordeaux.fr