



A new operational classification of the Conoidea (Gastropoda)

P. Bouchet, Yu I Kantor, A. Sysoev, N. Puillandre

► To cite this version:

P. Bouchet, Yu I Kantor, A. Sysoev, N. Puillandre. A new operational classification of the Conoidea (Gastropoda). *Journal of Molluscan Studies*, 2011, 77 (3), pp.273-308. 10.1093/mollus/eyr017 . hal-02458082

HAL Id: hal-02458082

<https://hal.science/hal-02458082>

Submitted on 28 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A NEW OPERATIONAL CLASSIFICATION OF THE CONOIDEA (GASTROPODA)

P. BOUCHET¹, Yu. I. KANTOR², A. SYSOEV³ AND N. PUILLANDRE¹

¹ *Museum National d'Histoire Naturelle, Departement Systematique et Evolution, UMR 7138, 55, Rue Buffon, 75231 Paris, France*

² *A.N. Severtsov Institute of Ecology and Evolution of Russian Academy of Sciences, Leninski Prospekt 33, Moscow 119071, Russia*

³ *Zoological Museum of Moscow State University, Bolshaya Nikitskaya str. 6, Moscow 1253009, Russia*

RUNNING HEAD: CLASSIFICATION OF CONOIDEA

(Received 14 October 2010; accepted 29 March 2011)

ABSTRACT

A new genus-level classification of the Conoidea is presented, based on the molecular phylogeny of Puillandre *et al.* in the accompanying paper. Fourteen lineages are recognized and ranked as families to facilitate continuity in the treatment of the names Conidae (for ‘cones’) and Terebridae in their traditional usage. The hitherto polyphyletic ‘Turridae’ is now resolved as 13 monophyletic families, in which the 358 currently recognized genera and subgenera are placed, or tentatively allocated: Conorbidae (2 (sub)genera), Borsoniidae (34), Clathurellidae (21), Mitromorphidae (8), Mangeliidae (60), Raphitomidae (71), Cochlespiridae (9), Drilliidae (34), Pseudomelatomidae (= Crassispiridae) (59), Clavatulidae (14), Horaiclavidae new family (28), Turridae s. s. (16) and Strictispiridae (2). A diagnosis with description of the shell and radulae is provided for each of these families.

INTRODUCTION

The history of classification of the Conoidea can be divided in two main periods. During the first period, from H. & A. Adams (1853), Bellardi (1875), Fischer (1887) and Cossmann (1896) to the 1980s (e.g. Powell, 1942, 1966; Morrison, 1965; McLean, 1971), classifications were based solely on the morphology of the ‘hard parts’ (shell and radula). Contradictions were commonplace and none of these classifications satisfied the malacological community, even if some of them were adopted by other authors (eg. Kilburn, 1983, 1985, 1986, 1988, 1991, 1992, 1993, 1994, 1995; Chang, 1995, 2001). Actually, only the Conidae (cone snails) and Terebridae (auger snails) were confidently recognized as well-defined groups, all the other constituents being lumped in the artificial group ‘turrids’, more or less separated into different subfamilies. Indeed, numerous cases of homoplasy for the morpho-anatomical characters used in these pre-1990s classifications are known (Taylor, 1994; Kantor, Medinskaya & Taylor, 1997; Fedosov & Kantor, 2008; Holford *et al.*, 2009), thus rendering earlier classifications questionable. Protoconch characters, in particular, have been much used and much misunderstood (Bouchet, 1990). Inevitably, different authors emphasized different types of characters (radula, foregut anatomy, shape of the shell, protoconch): when one taxonomist recognized a group based on protoconch similarities and ignored the radula morphology, another one stated that the similarity of the protoconch was the result of convergence, and split the group into as many groups as there were radula types. This contributed to the conoideans earning their reputation of ‘taxonomic nightmare’.

In 1993, based on anatomical characters (and not just the radula), Taylor, Kantor and Sysoev proposed a completely revised classification that recognized six families. One of their major results was that *Conus*, hitherto a genus in its own monotypic family Conidae, became one of many genera in an expanded family Conidae, which also now contained genera formerly placed in ‘Turridae’, thus confirming Thiele’s (1929) intuitive classification. The Taylor *et al.* (1993) classification formed the basis of the system published by Ponder & Bouchet (in Bouchet & Rocroi, 2005; Fig. 1A). Most recently, a new shell-and-radula classification of the Toxoglossa, with emphasis on *Conus* and other ‘cone-like’ genera, has been proposed by Tucker & Tenorio (2009; Fig. 1C), who recognized two superfamilies: the Turroidea, grouping together the Turridae of Taylor *et al.* (1993), plus the Terebridae, Drilliidae, Pseudomelatomidae and Strictispiridae, and the Conoidea, corresponding to the Conidae of Taylor *et al.* (1993) except for several of their constituents having been raised to family level. Based on a dataset of 57 genera and molecular sequences of fragments of one mitochondrial (COI) and three nuclear (28S, 18S and H3) genes, Puillandre *et al.* (2008) published the first molecular phylogeny of Conoidea. Even though ranks differed, most of the lineages defined by Taylor *et al.* (1993) were also retrieved in this molecular approach. The new molecular phylogeny of Puillandre *et al.* (in press) is based on a dataset of 102 conoidean genera (87 ‘turrids’, 5 cones and 10 terebrids) and sequences of three gene fragments (COI, 12S rRNA and 16S rRNA). Because of the high congruency between the classification based on anatomical characters and the molecular tree obtained, we are inclined to think that our understanding of the phylogeny of the Conoidea has now reached a stable position, and that a new classification is warranted. The classification presented below (Fig. 1B) aims to transform the molecular phylogeny into an operational classification by: (1) presenting revised morphological diagnoses for the now redefined families of Conoidea; (2) allocating (sometimes tentatively) all Recent genera recognized or used in current literature to the corresponding family.

Figure
1

MATERIAL AND METHODS

Genus-group names have been allocated to the newly defined families based on the following criteria, in order of decreasing robustness:

- (1) 226 genera were assigned to a family based on shell characters, and phenetic resemblance to those genera with radula and/or molecular characters available; those are marked¹.
- (2) 102 genera were classified on the basis of radula morphology (both ours and published data), and congruence between radula and molecular characters for those genera that were sequenced; these are marked².
- (3) 98 genera were classified in a family on the basis of the molecular data; these are marked³.
- (4) 63 genera and subgenera do not fall easily into the morphological groups resulting from the preceding steps, but they have tentatively been assigned to a family as a working hypothesis; these genera are preceded by a question mark.
- (5) 172 subgenera and/or synonyms are also listed; subgenera are in parentheses, synonyms in square brackets.

Our aim was to provide an exhaustive list of conoidean genus-group names based on Recent type species or reasonably recognized as Recent in the literature. Genera present exclusively in the fossil record are not included. In very few cases, obviously erroneous attribution of Recent species to entirely fossil genera also excluded these genera from our listing. Junior homonyms and *nomina nuda* are also not included because they are unavailable for nomenclatural purposes. When the status of a nominal genus is uncertain, we adopted a ‘valid until synonymized’ approach; that is, a name is regarded valid (irrespective of our opinion) if it has not been synonymized in literature. The names of synonyms are placed after corresponding valid names (the synonyms applied to the valid subgenus name are positioned after the subgenus name). All the genus and subgenus names are also listed in alphabetic order in Appendix 1.

Radulae

To illustrate the range of radular types in the molecularly defined clades, the radulae were extracted as far as possible from the specimens used for the molecular analyses or from conspecific specimens (after rehydration when soft parts had been dried), cleaned with diluted bleach, rinsed in distilled water, mounted on stubs, air dried, coated with gold-palladium, and investigated with a JEOL JSM 840A scanning electron microscope. Some of the previously photographed radulae (examined by YuIK and/or J.D. Taylor) were additionally illustrated. For radular descriptions we mostly followed the terminology accepted and discussed in Kantor & Taylor, 2000). In radular formulae the parentheses indicate partial or complete fusion of lateral and rachidian teeth (for more details see Kantor, 2006).

Ranking and nomenclature

Although the names Conoidea and Toxoglossa are used interchangeably in the taxonomic literature, we have avoided the name Toxoglossa because (1) it is not typified and cannot be used for a family-group name, and (2) many of the included taxa do not have a toxoglossate radula. Within the Conoidea, ranking of the clades was determined by a conservative approach, thus retaining the names Terebridae and Conidae s. s. (the latter including only the cone snails *Profundiconus*, *Californiconus*, *Conasprella*, *Conus* and *Taranteconus*) in their accustomed usage at family rank. An alternative would have been to recognize only two families, Conidae *sensu* Taylor *et al.* (1993) and Turridae (including the Turridae, Terebridae, Drillidae and Pseudomelatomidae), with the resulting inconvenience that Conidae s. s. and Terebridae, although monophyletic, would lose their traditional usage and the vast associated

literature dealing with these names that largely ignores the Taylor *et al.* (1993) and Bouchet & Rocroi (2005) classifications.

Fourteen clades of rank equivalent to Conidae *s. s.* and Terebridae are recognizable from the molecular phylogenetic tree (Puillandre *et al.*, in press). Forty-three family-group names within Conoidea are nomenclaturally available (Appendix 2), of which five are based on a genus with a fossil type species (Andoniinae Vera-Pelaez, 2002, Cryptoconinae Cossmann, 1896, Hemiconidae Tucker & Tenorio, 2009, Johnwyattidae Serna, 1979, Siphopsinae Le Renard, 1995). These five cannot be applied to a molecular clade and will not be used in our classification. Names were applied to clades based on the position of their type genus in the tree. If more than one family-group name was applicable, the valid name was determined by priority. If the type genus of a nominal family name was not sequenced, application of the name was determined by reference to the morphologically most similar genus used in the analysis.

Because the molecular taxon sampling is still too patchy for such levels, we have abstained from extending the classification below family level (i.e. subfamilies, tribes), even when some molecular clades obviously match previously recognized ‘subfamilies’ (e.g. Califoniconiidae, Oenopotinae, Zemaciinae, Zonulispirinae). They are all included in a family (Appendix 2), without precluding their usefulness and potential taxonomic validity.

Illustration of voucher material

Each family (except the Conidae and Terebridae, already extensively covered and illustrated in other recent works, e.g. Röckel, Korn & Kohn, 1995; Terryn, 2007; Tucker & Tenorio, 2009) was illustrated by one or several shells, radulae and protoconchs, covering the morphological variability of the group. As far as possible, specimens used for the molecular analyses were used for illustration. However, since a substantial part of the morphological variability was not covered by our dataset, shells and radulae of other specimens are also illustrated.

RESULTS

SUPERFAMILY CONOIDEA FLEMING, 1822

Family Conidae Fleming, 1822

[= Conilithidae Tucker & Tenorio, 2009, **new synonym**; = Taranteconidae Tucker & Tenorio, 2009, **new synonym**]

Diagnosis: Shell medium-sized to large or very large, normally 20–50 mm, up to 170 mm high, conical or biconical, with narrow aperture and short siphonal canal. Shell with considerable internal remodelling due to inner wall resorption. Spiral sculpture usually developed, axial sculpture absent or in form of shoulder tuberculation. Anal sinus on subsutural ramp, shallow to moderately deep. Operculum present, small, with terminal nucleus. Radula of marginal hypodermic teeth, generally harpoon-shaped, barbed at tip, often with complex inner structure of folds and serration, base small and swollen, tooth canal opening (sub)terminally, rarely laterally. Subradular membrane vestigial. Teeth can be attached to the membrane by long or very long flexible ligament. Tooth wall forms several overlapping layers.

Remarks: in Conidae and the other families that are included in this major clade (family Conidae *sensu* Taylor *et al.*, 1993), the radula consists only of marginal teeth that are usually

enrolled with completely overlapping edges (hypodermic). Teeth at their formation in the radular sac are already enrolled and they are attached to the radular membrane only by their bases, sometimes through a long flexible ligament. In the ‘turrids’ with enrolled teeth, these are attached to the membrane along most of its length (Kantor & Taylor, 2000). A molecular phylogeny of the Conidae is currently in preparation (C. Meyer, personal communication).

Included genera:

- Africonus* Petuch, 1975¹
- Asprella* Schaufuss, 1869¹
 - [= *Embrikena* Iredale, 1937]
 - [= *Fumiconus* da Motta, 1991]
 - [= *Graphiconus* da Motta, 1991]
- Astroconus* Tucker & Tenorio, 2009¹
- Bathyconus* Tucker & Tenorio, 2009¹
- Calamiconus* Tucker & Tenorio, 2009¹
- Calibanus* da Motta, 1991¹
- Californiconus* Tucker & Tenorio, 2009³
- Chelyconus* Mörch, 1852¹
- Conasprella* Thiele, 1929³
- Conaspelloides* Tucker & Tenorio, 2009¹
- Conus* Linnaeus, 1758³
 - [= *Coronaxis* Swainson, 1840]
 - [= *Cucullites* Herrmannsen, 1847]
 - [= *Cucullus* Röding, 1798]
- Cylindrus* Batsch, 1789¹ (For authorship of *Cylindrus*, see Dubois & Bour (2010))
 - [= *Cylinder* Montfort, 1810]
 - [= *Cylindrella* Swainson, 1840]
- Dalliconus* Tucker & Tenorio, 2009¹
- Darioconus* Iredale, 1930¹
 - [= *Erythroconus* da Motta, 1991]
 - [= *Regiconus* Iredale, 1930]
- Dauciconus* Cotton, 1945¹
 - [= *Cariboconus* Petuch, 1991]
- Dendroconus* Swainson, 1840¹
 - [= *Clebula* Iredale, 1930]
- Ductoconus* da Motta, 1991¹
- Dyraspis* Iredale, 1949¹
- Endemoconus* Iredale, 1931¹
 - [= *Kemasprella* Powell, 1958]
 - [= *Mamiconus* Cotton & Godfrey, 1932]
- Eremiconus* Tucker & Tenorio, 2009¹
- Eugeniconus* da Motta, 1991¹
- Floraconus* Iredale, 1930¹
- Fulgiconus* da Motta, 1991¹
 - [= *Thoraconus* da Motta, 1991]
- Fusiconus* da Motta, 1991¹
- Gastridium* Modeer, 1793¹
 - [= *Rollus* Montfort, 1810]
 - [= *Tuliparia* Swainson, 1840]
- [= *Utriculus* Schumacher, 1817]
- Genuanoconus* Tucker & Tenorio, 2009¹
- Gladioconus* Tucker & Tenorio, 2009¹
- Globiconus* Tucker & Tenorio, 2009¹
- Gradiconus* da Motta, 1991¹
- Harmoniconus* da Motta, 1991¹
- Hermes* Montfort, 1810¹
 - [= *Theliconus* Swainson, 1840]
- Jaspidiconus* Petuch, 2003¹
- Kalloconus* da Motta, 1991¹
- Kenyonia* Brazier, 1896¹
- Ketyconus* da Motta, 1991¹
- Kioconus* da Motta, 1991¹
 - [= *Ongococonus* da Motta, 1991]
- Kohniconus* Tucker & Tenorio, 2009¹
- Kuradoconus* Shikama & Habe, 1968¹
- Lamniconus* da Motta, 1991¹
- Lautoconus* Monterosato, 1923¹
- Leporiconus* Iredale, 1930¹
- Leptoconus* Swainson, 1840¹
- Lilliconus* Raybaudi Massilia, 1994¹
- Lithoconus* Mörch, 1852¹
 - [= *Tesselliconus* da Motta, 1991]
- Lividoconus* Wils, 1970¹
 - [= *Splinoconus* da Motta, 1991]
- Miliariconus* Tucker & Tenorio, 2009¹
- Monteiroconus* da Motta, 1991¹
- Nataliconus* Tucker & Tenorio, 2009¹
- Parviconus* Cotton & Godfrey, 1932¹
- Perplexiconus* Tucker & Tenorio, 2009¹
- Phasmococonus* Mörch, 1852¹
- Pionoconus* Mörch, 1852¹
 - [= *Heroconus* da Motta, 1991]
 - [= *Socioconus* da Motta, 1991]
 - [= *Strioconus* Thiele, 1929]
- Plicaustraconus* Moolenbeek 2008¹
- Profundiconus* Kuroda, 1956³
 - [= *Lizaconus* da Motta, 1991]
- Protoconus* da Motta, 1991¹ [invalid name]
 - [= *Seminoleconus* Petuch, 2003]
- Protostrioconus* Tucker & Tenorio, 2009¹
- Pseudoconorbis* Tucker & Tenorio, 2009¹

- Pseudolilliconus* Tucker & Tenorio, 2009¹
Pseudonoduloconus Tucker & Tenorio, 2009¹
Puncticulis Swainson, 1840¹
Purpuriconus da Motta, 1991¹
[= *Magelliconus* da Motta, 1991]
Pyruconus Olsson, 1967¹
Quasiconus Tucker & Tenorio, 2009¹
Rhizoconus Mörch, 1852¹
Rhombiconus Tucker & Tenorio, 2009¹
[= *Rhombus* Montfort, 1810]
Rolaniconus Tucker & Tenorio, 2009¹
Sciteconus da Motta, 1991¹
Lindaconus Petuch, 2002¹
[= *Calusaconus* Petuch, 2003]
[= *Spuriconus* Petuch, 2003]
- Stellaconus* Tucker & Tenorio, 2009¹
Stephanoconus Mörch, 1852¹
Strategoconus da Motta, 1991¹
Taranteconus Azuma, 1972³
[= *Cornutoconus* Suzuki, 1972]
Textilia Swainson, 1840¹
Trovaconus Tucker & Tenorio, 2009¹
Turriconus Shikama & Habe, 1968¹
Varioconus da Motta, 1991¹
Viminiconus Tucker & Tenorio, 2009¹
Virgiconus Cotton, 1945¹
Virroconus Iredale, 1930¹
Vituliconus da Motta, 1991¹
Ximeniconus Emerson & Old, 1962¹
Yedoconus Tucker & Tenorio, 2009¹

Conorbidae de Gregorio, 1880
(Figs 2O, 22C)

Diagnosis (from Tucker & Tenorio, 2009): “Radular tooth: anterior fold is usually present; basal spur is directed toward the apex of the tooth or parallel with the tooth base; the waist, base and C-fold are absent; terminating cusps, serratins and accessory process are also absent. Shell characters: the interior of the shell is extensively remodelled including the columellar region; nodules are absent but cords may be present. Shells can be squatly conical to elongated or biconical.”

Remarks: The type genus of the family is *Conorbis* Swainson, 1840, based on an Eocene type species. The only Recent species that was believed to belong to *Conorbis*, *Conus coromandelicus* E.A. Smith, 1894, was made the type of the new genus *Pseudoconorbis* Tucker & Tenorio, 2009, and included by them in their family Conilithidae (herein in Conidae). In the molecular phylogeny of Puillandre *et al.* (in press), what we have identified as ‘*Benthofascis*’ constitutes an independent lineage (i.e. not included in any other family-ranked clade), characterized by a long branch. This certainly explains why its position is not the same in the MLA (sister group of the Borsoniidae) and in the BA (included in the Borsoniidae). *Benthofascis lozoueti* (the sequenced species) differs from other species of the genus in the absence of inner shell wall resorption (N. Puillandre, personal observations), but phylogenetic and taxonomic value of this character remains unverified. The problem can be resolved by sequencing other species of *Benthofascis*, which have inner wall resorption. Until then we prefer to keep *Benthofascis lozoueti* within this genus and to apply the name Conorbidae to this lineage.

Included genera:
Artemidiconus da Motta, 1991¹

Benthofascis Iredale, 1936³

Borsoniidae A. Bellardi, 1875
[= Zemaciinae Sysoev, 2003, new synonym]
(Figs 2A-N, 3)

Figure
2

Figure
3

Diagnosis: Shell small to large (5-80 mm), fusiform to biconic, sometimes with strong to obsolete columellar pleats. Sculpture usually well developed, axial ribs sometimes obsolete to absent. Siphonal canal short to moderately long. Anal sinus on subsutural ramp, deep. Protoconch when multispiral with up to 5 whorls, initially smooth and then with arcuate axial riblets, when paucispiral up to 2 smooth whorls. Operculum with terminal nucleus, fully developed to missing. Radula of hypodermic marginal teeth that usually have a weakly developed solid basal part, often attached to the ligament (marked by an arrow on Fig. 3 C). Tooth canal opening (sub)terminally or, sometimes, laterally. At their tip teeth can have weak to rather strong barb(s) (*Genota*, Fig. 19A). Overlapping of the tooth edges is weak (Fig. 19C). In *Zemacies* the radula is completely absent.

Remarks: This is a rather heterogeneous group. Obviously, it is not fully resolved as it is based on molecular data and comprises rather conchologically different clades. This could be explained by the fact that many taxa of this group are among most ancient of conoideans, known since the Paleocene (*Zemacies*, *Borsonia*, *Tomopleura*) or Eocene (*Bathytoma*, *Genota*, *Microdrillia*). The loss of apomorphies by mutation can be more important for old taxa (Puillandre *et al.*, in press).

The names *Borsoniinae* Bellardi, 1875, and *Pseudotominae* Bellardi, 1875, were established simultaneously. As First Revisers, under Art. 24 of the ICZN Code, we here give precedence to the former over the latter.

Included genera:

- | | |
|--|---|
| <i>Aphanitoma</i> Bellardi, 1875 ¹ | ? <i>Glyptaesopus</i> Pilsbry & Olsson, 1941 ² |
| <i>Asthenotoma</i> Harris & Burrows, 1891 ¹ | <i>Heteroturris</i> Powell, 1967 ¹ |
| <i>Austroturris</i> Laseron, 1954 ¹ | <i>Maoritomella</i> Powell, 1942 ² |
| ? <i>Apaturris</i> Iredale, 1917 ¹ | [= <i>Narraweena</i> Laseron, 1954] |
| <i>Bathytoma</i> Harris & Burrows, 1891 ³ | <i>Microdrillia</i> Casey, 1903 ³ |
| (<i>Micantapex</i> Iredale, 1936) | [= <i>Acropota</i> Nordsieck, 1977] ² |
| (<i>Parabathytoma</i> Shuto, 1961) | <i>Ophiodermella</i> Bartsch, 1944 ² |
| (<i>Riuguhdrillia</i> Oyama, 1951) | <i>Paraborsonia</i> Pilsbry, 1922 ¹ |
| ? <i>Belaterricula</i> Powell, 1951 ² | <i>Phenatoma</i> Finlay, 1924 ² |
| <i>Borsonella</i> Dall, 1908 ² | <i>Pulsarella</i> Laseron, 1954 ² |
| <i>Borsonellopsis</i> McLean, 1971 ¹ | <i>Retidrillia</i> McLean, 2000 ¹ |
| <i>Borsonia</i> Bellardi, 1839 ³ | <i>Suavodrillia</i> Dall, 1908 ² |
| (<i>Boettgeriola</i> Wenz, 1943) | <i>Tomopleura</i> Casey, 1904 ³ |
| <i>Cordieria</i> Rouault, 1848 ¹ | <i>Tropidoturris</i> Kilburn, 1986 ² |
| <i>Darbya</i> Bartsch, 1944 ¹ | ? <i>Typhlodaphne</i> Powell, 1951 ¹ |
| <i>Diptychophlia</i> Berry, 1964 ¹ | <i>Typhlomangelia</i> G.O. Sars, 1878 ³ |
| <i>Drilliola</i> Cossmann, 1903 ² | [= <i>Viridoturris</i> Powell, 1964] ² |
| <i>Filodrillia</i> Hedley, 1922 ¹ | [= <i>Vexithara</i> Finlay, 1926] |
| <i>Genota</i> H. Adams & A. Adams, 1853 ³ | <i>Zemacies</i> Finlay, 1926 ³ |
| [= <i>Genotia</i> Fischer, 1883] | |

Figure
4

Clathurellidae H. Adams & A. Adams, 1858
(Figs 4A-E, 5)

Figure
5

Diagnosis: Shell small to medium-sized (5-40, usually 10-20mm), fusiform to broadly fusiform, subsutural ramp usually unclearly separated. Sculpture mostly strong, usually cancellate, subsutural fold lacking. Shell surface often microgranular. Apertural armature often well developed, in the form of pleats and denticles on both inner and outer lips; no true

columellar pleats. Siphonal canal well expressed, moderately long. Anal sinus on subsutural ramp, deep, often (sub)tubular. Protoconch typically multispiral, up to six whorls, generally smooth with keeled last whorls; when paucispiral, protoconch usually smooth or spirally striated, sometimes with remaining carination in last portion. Operculum always absent. Radula of hypodermic marginal teeth that usually have relatively small solid basal part (Fig. 5). Distinct ligaments not found. At their tip, teeth can have a weak barb.

Included genera:

- Arista* Hedley, 1922¹
- ?*Adanaclava* Bartsch, 1950¹
- ?*Clathromangelia* Monterosato, 1884¹
- Clathurella* Carpenter, 1857³
- Comarmondia* Monterosato, 1844¹
[= *Bellatula* Strand, 1929]
- Corinnaeturris* Bouchet & Warén, 1980²
- Crockerella* Hertlein & Strong, 1951²
- Etrema* Hedley, 1918³
[= *Iraqetrema* Dance & Eames, 1966]
(*Etremopa* Oyama, 1953)
(*Etremopsis* Powell, 1942)

- Euclathurella* Woodring, 1928¹
- Glyphostoma* Gabb, 1872²
(*Euglyphostoma* Woodring, 1970)
- (*Glyphostomopsis*) Bartsh, 1934
- Lienardia* Jousseaume, 1884¹
(*Thetidos* Hedley, 1899)
- Nannodiella* Dall, 1919³
- Paraclathurella* Boettger, 1895¹
- Pseudoetrema* Oyama, 1953¹
- Strombinoturris* Hertlein & Strong, 1951²
- Turrella* Laseron, 1954¹

Mitromorphidae Casey, 1904, new rank
(Figs 4F-H, 6)

Diagnosis: Shell small to medium-sized, 3-30 mm, usually 5-10 mm high, biconic, of mitriform shape. Sculpture rather smooth, with dominant spiral elements. Aperture narrow, with or without 1-3 columellar pleats, sometimes with denticles within. Siphonal canal short or indistinct. Anal sinus from indistinct to rather shallow indentation on weakly pronounced subsutural ramp. Protoconch multispiral or paucispiral, up to 4.5 smooth whorls. No operculum. Radula of hypodermic, marginal, relatively short, awl-shaped teeth with large swollen solid basal part (Fig. 6). Distinct ligaments present, short. Tooth canal opening subterminally or laterally. At their tip, teeth can have a weak barb (Fig. 6 C).

Remarks: Mitrolumninae was established as a substitute name for Diptychomitrinae. Mitromorphidae and Mitrolumninae were published the same year but Mitromorphidae (19 May 1904) has priority over Mitrolumninae (31 August 1904).

Included genera:

- Anarithma* Iredale, 1918¹
- Arielia* Shasky, 1961¹
(*Vexiariela* Shuto, 1983)
- Lovellona* Iredale, 1917³
- Maorimorpha* Powell, 1939¹
- Mitrellatoma* Powell, 1942¹
- Mitromorpha* Carpenter, 1865³
- [= *Cymakra* Gardner, 1937]
- [= *Helenella* Casey, 1904]
- [= ?*Itia* Marwick, 1931]
- [= *Mitrtihara* Hedley, 1922]
- [= (*Mitrolunna* Bucquoy, Dautzenberg & Dollfus, 1883)]
- Scrinium* Hedley, 1922¹

Mangeliidae P. Fischer, 1883, new rank
[= Oenopotinae Bogdanov, 1987 new synonym]
(Figs 7, 8)

Figure
6

Figure
7

Figure
8

Diagnosis: Shell small to medium size, 3-30 mm, usually 6-12 mm, oval to low- or high-fusiform, usually with comparatively low spire, often with a shoulder angulation. Spiral and axial sculpture both well developed. Shell surface often bearing microsculpture of spirally aligned granules. Subsutural ramp usually not separated sculpturally. Anal sinus on subsutural ramp, shallow to rather deep, rarely tubular. Aperture normally not constrained with strong outgrowths or callus pads, rarely denticulate. Siphon rather short to moderately long. Protoconch typically multispiral, of up to five whorls, with axially ribbed protoconch II; spiral cords on protoconch II present or absent. When paucispiral, protoconch usually spirally lirate. Operculum present, with terminal nucleus (Oenopotinae), or normally absent. Radula of marginal teeth of very variable morphology. Teeth can be from semi-enrolled (Fig. 8 F-G) to true hypodermic. Frequently side projections around the base are present and there is large irregularly shaped ‘root’ projecting from the base. Barbs may be present, from small to very large (Fig. C) or absent (Fig. 8B). Tooth canal opening laterally.

Included genera:

- | | |
|--|---|
| <i>Acmaturris</i> Woodring, 1928 ¹ | <i>Kurtzia</i> Bartsch, 1944 ² |
| <i>Agathotoma</i> Cossmann, 1899 ² | <i>Kurtziella</i> Dall, 1918 ² |
| <i>Anticlinura</i> Thile, 1934 ³
[= <i>Clinuromella</i> Beets, 1943] | <i>Kurtzina</i> Bartsch, 1944 ² |
| <i>Apispiralia</i> Laseron, 1954 ¹ | <i>Leiocithara</i> Hedley, 1922 ¹ |
| <i>Apitua</i> Laseron, 1954 ¹ | <i>Liracraea</i> Odhner, 1924 ² |
| <i>Bactrocythara</i> Woodring, 1928 ¹ | ? <i>Macteola</i> Hedley, 1918 ¹ |
| <i>Bela</i> Gray, 1847 ²
[= <i>Ginnania</i> Monterosato, 1884] | <i>Mangelia</i> Risso, 1826 ²
[= <i>Cyrtocythara</i> Nordsieck, 1977] |
| [= <i>Smithiella</i> Monterosato, 1890] | [= <i>Cytharella</i> Monterosato, 1875] |
| <i>Bellacythara</i> McLean, 1971 ² | [= <i>Lyromangelia</i> Nordsieck, 1977] |
| <i>Benthomangelia</i> Thiele, 1925 ³ | [= <i>Mangiliella</i> Bucqoy, Dautzenberg & Dollfus, 1883] |
| <i>Brachycythara</i> Woodring, 1928 ¹ | [= <i>Rissomangelia</i> Monterosato, 1917] |
| ? <i>Cacodaphnella</i> Pilsbry & Lowe, 1932 ¹ | [= <i>Rugocythara</i> Nordsieck, 1977] |
| <i>Citharomangelia</i> Kilburn, 1992 ² | [= <i>Villiersiella</i> Monterosato, 1884] |
| <i>Cryoturris</i> Woodring, 1828 ¹ | <i>Marita</i> Hedley, 1922 ¹ |
| <i>Curtitoma</i> Bartsch, 1941 ²
[= <i>Nematoma</i> Bartsch, 1941] | <i>Mitraguraleus</i> Laseron, 1954 ¹ |
| [= <i>Widalli</i> Bogdanov, 1986] | ? <i>Neoguraleus</i> Powell, 1939 ² |
| <i>Cytharopsis</i> A. Adams, 1865 ¹ | ? <i>Notocytharella</i> Hertlein & Strong, 1955 ¹ |
| <i>Eucithara</i> Fischer, 1883 ³
[= ? <i>Cythara</i> Schumacher, 1817] | <i>Obesotoma</i> Bartsch, 1941 ² |
| ? <i>Euryentmema</i> Woodring, 1928 ¹ | <i>Oenopota</i> Mörch, 1852 ³
(<i>Nodotoma</i> Bartsch, 1941) |
| <i>Fehria</i> van Aartsen, 1988 ¹ | <i>Oenopotella</i> Sysoev, 1988 ¹ |
| <i>Genotina</i> Vera-Peláez, 2004 ¹ | <i>Papilocithara</i> Kilburn, 1992 ¹ |
| <i>Gingicithara</i> Kilburn, 1992 ² | ? <i>Paraspirotropis</i> Sysoev & Kantor, 1984 ² |
| <i>Glyphoturris</i> Woodring, 1928 ¹ | <i>Perimangelia</i> McLean, 2000 ¹ |
| <i>Granotoma</i> Bartsch, 1941 ² | <i>Platocythara</i> Woodring, 1928 ¹ |
| <i>Granoturris</i> Fargo, 1953 ¹ | ? <i>Propebela</i> Iredale, 1918 ²
[= <i>Antiguraleus</i> Powell, 1942] |
| <i>Guraleus</i> Hedley, 1918 ²
(<i>Euguraleus</i> Cotton, 1947) | [= ? <i>Belalora</i> Powell, 1951] |
| <i>Hemicythara</i> Kuroda & Oyama in Kuroda,
Habe & Oyama, 1971 ¹ | [= <i>Cestoma</i> Bartsch, 1941] |
| <i>Heterocithara</i> Hedley, 1922 ³ | [= <i>Funitoma</i> Bartsch, 1941] |
| <i>Ithycythara</i> Woodring, 1928 ² | [= ? <i>Lorabela</i> Powell, 1951] |
| | [= <i>Turritomella</i> Bartsch, 1941] |
| | (<i>Canetoma</i> Bartsch, 1941) |

Pseudorhaphitoma Boettger, 1895²
Pyrgocythara Woodring, 1928²
Rubellatoma Bartsch & Rehder, 1939¹
Saccharoturris Woodring, 1928¹
Stellatoma Bartsch & Rehder, 1939¹
Suturocythara García, 2008¹

Tenaturris Woodring, 1928²
Toxicochlespira Sysoev & Kantor, 1990³
?Venustoma Bartsch, 1941¹
Vitjazinella Sysoev, 1988¹
?Vitricythara Fargo, 1953¹

Raphitomidae A. Bellardi, 1875

(Figs 9, 10)

Figure
9

Figure
10

Diagnosis: Shell very variable in shape and size, from buccinoid to ovate or to elongate-fusiform or high-cylindrical and from about 2 to over 140 mm high. Sculpture variously developed: from nearly smooth shell to well developed spiral and axial elements, subsutural ramp, when separated from rest of the whorl, usually smooth or with traces of anal sinus growth. Apertural armature rarely well expressed, inner lip usually smooth. Anal sinus either subsutural, shaped as a ‘reversed-L’, or on subsutural ramp, asymmetric, very shallow to rather deep, typically not constrained but sometimes even almost tubular. Siphon short to rather long. Protoconch typically planktotrophic, multispiral, of 2.5-6.5 whorls, protoconch I often spirally striated, protoconch II with diagonally cancellated sculpture. There are many variations of the basic pattern of the protoconch II sculpture, though a combination of spiral and axial elements is retained, with the latter usually more pronounced, but protoconchs with a spiral keel and axial pillars (Fig. 23) depart from this pattern. Paucispiral protoconch with predominantly spiral striation. No operculum. Radula of hypodermic marginal teeth of very variable morphology. Teeth can be relatively very long (to 12.8% of shell length in *Daphnella cladara*, Fig. 10E) to very short and reduced (0.13% of shell length in *Spergo fusiformis*, Fig. 10H). Solid base of the teeth from very small (Fig. 10E, G) to rather large (Fig. 10B, F), sometimes side projections around the base are present (Fig. 10F). Barb(s) present or absent. At least in one species cockscomb-shaped structure is present near the tooth tip (Fig. 10G). Tooth canal opening laterally or subapically. Reduction of radular apparatus and other structures of the anterior foregut is frequent grading to complete absence.

Remarks: This is the largest and most variable taxon in the Conoidea, as concerns the number of species, with the largest vertical range (intertidal to hadal depths).

Included genera:

Abyssobela Kantor & Sysoev, 1986¹
?Abyssothauma Sysoev, 1996¹
Acamptodaphne Shuto, 1971¹
Acanthodaphne Bonfitto & Morassi, 2006¹
Aliceia Dautzenberg & Fischer, 1897¹
Asperdaphne Hedley, 1922¹
 (Aspertilla Powell, 1944)
Austrodaphnella Laseron, 1954¹
?Austropusilla Laseron, 1954¹
 (Metaclathurella Shuto, 1983)
Bathybela Kobelt, 1905²
 [= *Bathypota* Nordsieck, 1968]
Buccinaria Kittl, 1887²
 [= *Otomella* Bartsch, 1933]
 [= *Pionotoma* Kuroda, 1952]

Cenodagreutes E.H. Smith, 1967¹
Clathurina Melvill, 1917¹
Clinura Bellardi, 1875¹
Cryptodaphne Powell, 1942¹
Daphnella Hinds, 1844³
 [= *Eudaphnella* Bartsch, 1933]
 [= *Paradaphne* Laseron, 1954]
 (*Diaugasma* Melvill, 1917)
 (*Hemidaphne* Hedley, 1918)
Eubela Dall, 1889¹
Eucyclotoma Boettger, 1895³
 [= *Turhyssa* Dall, 1924]
?Exomilus Hedley, 1918¹
Famelica Bouchet & Warén, 1980¹
Favriella Hornung, 1920¹

- Glyphostomoides* Shuto, 1983³
Gymnobela Verrill, 1884³
[= *Majox* Nordsieck, 1968]
[= *Watsonaria* Nordsieck, 1968]
(*Theta* Clarke, 1959)
Hemilienardia Boettger, 1895³
Isodaphne Laseron, 1954¹
Kermia Olivier, 1915³
Kuroshiodaphne Shuto, 1965¹
Leiosyrinx Bouchet & Sysoev, 2001²
Lusitanops Nordsieck, 1968¹
Magnella Dittmer, 1960¹
Microdaphne McLean, 1971¹
Microgenia Laseron, 1954¹
Mioawateria Vella, 1954¹
Neopleurotomoides Shuto, 1971¹
Nepotilla Hedley, 1918²
Pagodidaphne Shuto, 1983¹
Paramontana Laseron, 1954²
Philbertia Monterosato, 1884²
[= *Leufroyia* Monterosato, 1884]
[= *Lineotoma* Nordsieck, 1977]
[= *Peratotoma* Harris & Burrows, 1891]
Phymorhynchus Dall, 1908²
Pleurotomella Verrill, 1873³
[= *Azorilla* Nordsieck, 1968]
[= *Azorita* Nordsieck, 1968]
[= *Anomalotomella* Powell, 1966]
[= *Fusidaphne* Laseron, 1954]
[= *Tasmadaphne* Laseron, 1954]
(*Cyrtoides* Nordsieck, 1968)
Pontiothauma E.A. Smith, 1895²
- Pseudodaphnella* Boettger, 1895²
(*Qii* Zhang, 1995)
Raphitoma Bellardi, 1848³
Rimosodaphnella Cossmann, 1915³
Rocroithys Sysoev & Bouchet, 2001²
Spergo Dall, 1895²
(*Speoides* Kuroda & Habe in Habe, 1962)
Stilla Finlay, 1926¹
Taranidaphne Morassi & Bonfitto, 2001¹
Taranis Jeffreys, 1867³
[= ?*Allo* Lamy, 1934]
[= ?*Feliciella* Lamy, 1934]
[= *Fenestrosyrinx* Finlay, 1926]
?*Teleochilus* Harris, 1897¹
[= *Litachilus* Powell, 1944]
Teretia Norman, 1888²
Teretiopsis Kantor & Sysoev, 1989³
Thatcheria Angas, 1877³
[= *Cochliococonus* Yokoyama, 1928]
?*Thatcheriasyrinx* Powell, 1969¹
Thatcherina Vera-Peláez, 1998¹
?*Thesbia* Jeffreys, 1867²
Tritonoturris Dall, 1924³
Truncadaphne McLean, 1971¹
Tuskaroria Sysoev, 1988¹
Typhlosyrinx Thiele, 1925²
Veprecula Melvill, 1917³
[= *Mordica* Dall, 1924]
Vepridaphne Shuto, 1983¹
Xanthodaphne Powell, 1942²
Zenepos Finlay, 1928¹

Figure
11

Cochlespiridae Powell, 1942
(Figs 11A-D, 12)

Figure
12

Diagnosis: Shell of moderate size, about 20-30 mm, up to 100 mm high, high-pagodiform to fusiform, with a tall spire and usually a long siphonal canal. Axial sculpture poorly developed or absent, subsutural ramp usually smooth. Anal sinus deep, on subsutural ramp. Protoconch paucispiral, smooth. Operculum with terminal nucleus. Radula 1-0-R-0-1. Rachidian broad, subrectangular or arched, with a single rather large cusp (Fig. 12), rarely absent (some *Aforia*). Marginal teeth duplex, with well developed accessory limb.

Remarks: This is a clade with poor congruence between molecular and shell characters. The contents of most genera need revision and the family limits remain uncertain.

Sibogasyrinx was proposed as a subgenus of *Leucosyrinx* for two species characterized by a low position of the peripheral angle. During our examination of radulae of species provisionally attributed to *Leucosyrinx*, two distinct radulae types were found, differing in the presence of the rachidian, without apparent congruence with shell morphology. In *S. pyramidalis*, the type species of *Sibogasyrinx*, as well as in the sequenced specimen (Fig.

11B), the radula is characterized by the presence of a well developed rachidian. Although the sequenced specimen is conchologically closer to typical *Leucosyrinx*, the molecular sequence suggests affinities to *Cochlespira*.

Included genera:

- ?*Aforia* Dall, 1889²
[= *Irenosyrinx* Dall, 1908]
[= *Danilacarina* Bozzetti, 1997 syn. nov.]
(?*Abyssaforia* Sysoev & Kantor, 1987)
(?*Dallaforia* Sysoev & Kantor, 1987)
(?*Steiraxis* Dall, 1896)
?*Apiotoma* Cossmann, 1889¹

- ?*Clavosurcula* Schepman, 1913¹
Cochlespira Conrad, 1865³
[= *Ancistrosyrinx* Dall, 1881]
[= *Coronasyrinx* Powell, 1944]
[= *Pagodasyrinx* Shuto, 1969]
?*Nihonia* MacNeil, 1961¹
Sibogasyrinx Powell, 1969³

Drilliidae Olsson, 1964
(Figs 11F-L, 13)

Diagnosis: Shell small to medium-sized, usually 15–25 mm, up to 50 mm high, with a rather high spire and usually truncated base. Spiral sculpture often obsolete. Anal sinus on subsutural ramp, deep, (sub)symmetrical, sometimes tubular. Protoconch usually paucispiral (up to 2 whorls), smooth or abapically carinate. Operculum with terminal nucleus. Radular formula 1-1-R-1-1, rarely 1-1-0-1-1, central tooth small, from narrow unicuspis (Fig. 13B) to subrectangular with additional cusps (Fig. 13C), rarely reduced to completely absent. Lateral teeth broad, pectinate, and arched, marginal teeth from simple flat and sharply pointed (Fig. 13 A) to duplex with slightly thickened edges and to loosely enrolled with the small barb near the tip (*Imaclava*, Fig. 13D).

Remarks: The genera *Cruziturridula* and *Fusiturridula* form an unsupported group that is sister to Drillidae. In the studied *Cruziturridula arcuata* (Reeve, 1843) the very characteristic radula differs from the Drillidae: 1-0-0-0-1. Marginal teeth are loosely enrolled with little overlap of the edges, with two barbs on the tip and a tongue-shape extension at the base (Fig. 13E). The type species of *Fusiturridula*, *Turris fusinella* Dall, 1908, is different from what is currently conceived as belonging to that genus (e.g. Williams, 2006), but those species are similar to *Cruziturridula sensu auctt.* The only data on the radula was provided by Powell (1966: 31), who stated (without mentioning the species): “Radula of marginals only, wishbone-type, but long and narrow”. Although *Cruziturridula* and *Fusiturridula* definitely do not belong in Drillidae and may represent a new family, they are here provisionally placed in Drillidae for lack of a better alternative.

Included genera:

- ?*Acinodrillia* Kilburn, 1988¹
Agladrillia Woodring, 1928³
Bellaspira Conrad, 1868¹
Calliclava McLean, 1971³
Cerodrillia Bartsch & Rehder, 1939³
(*Lissodrillia* Bartsch and Rehder, 1943)
Clavus Monfort, 1810³
[= *Clavicantha* Swainson, 1840]
[= *Eldridgea* Bartsch, 1934]
[= *Tylotia* Melvill, 1917]
Clathrodrillia Dall, 1918³

- Conopleura* Hinds, 1844³
Crassopleura Monterosato, 1884¹
?*Cruziturridula* Marks, 1951³
Cymatosyrinx Dall, 1889²
Douglassia Bartsch, 1934¹
Drillia Gray, 1838³
Elaeocyma Dall, 1918²

Globidrillia Woodring, 1928²
Imaclava Bartsch, 1944³
Iredalea Oliver, 1915³
 [= *Brephodrillia* Pilsbry & Lowe, 1932]
Kylix Dall, 1919²
Leptadrillia Woodring, 1928²
Neodrillia Bartsch, 1943²
Orrmaesia Kilburn, 1988²

?Paracuneus Laseron, 1954¹
Plagiostropha Melvill, 1927²
Splendrillia Hedley, 1922³
 (*Hauturua* Powell, 1942)
Spirotropis G.O. Sars, 1878²
?Stenodrillia Korobkov, 1955¹
Syntomodrillia Woodring, 1928²
Tylotiella Habe, 1964²

Figure
14

Pseudomelatomidae Morrison, 1965 (Figs 14-16)

Figure
15

Diagnosis: Shell small to rather large, 15 to 100 mm high, claviform to fusiform. Spiral and axial sculpture generally well developed, often strong. Subsutural fold often present. Anal sinus on subsutural ramp, usually moderately deep to very deep, often constrained by callus rendering anal sinus subtubular. Protoconch usually paucispiral, sometimes multispiral, with up to 3 whorls, smooth or sometimes axially or spirally sculptured on later whorls. Operculum with terminal nucleus.

Radula very variable, 4 types are recorded:

1-(1-R-1)-1 – *Comitas* type (includes also *Knefastia* and *Antiplanes*). The central formation is rather variable in degree of development of the rachidian tooth and fusion of 3 teeth. In some *Comitas* (Fig. 15C-D) the central formation looks like single well defined tooth [as in *C. murrawolga* (Garrard, 1961)], while in *C. onokeana vivens* Dell, 1956, *Knefastia* and *Antiplanes*, the rachidian is totally reduced and the formation appears as two poorly developed paired plates (reduced laterals) (Fig. 15B). Marginal teeth in *Comitas* and *Knefastia* flat, broadly oval, with thickened edges and teeth tips and without pronounced accessory limb. In *Antiplanes* marginal teeth narrow elongate, with well developed accessory limb.

1-1-0-1-1 – *Crassiclava* type. Differs from the *Comitas* type by the better defined laterals (Fig. 15E, F) that are low, arcuate and sharply curved towards the midline of the ribbon.

1-0-R-0-1 – *Pseudomelatoma* type (includes also *Hormospira* and *Tiariturris*, Fig. 15A). Rachidian with strong cusp and subrectangular base (contrary to the other types with a central formation formed by fused lateral and rachidian). Despite neither ontogeny nor folding of the radula have been examined, we tentatively treat this structure as a true rachidian. Marginal teeth simple, solid and strongly curved, attached to the membrane by rather a narrow base and free along most of their length.

1-0-0-0-1 – Most genera of the family. Marginal teeth elongated, narrow, flat, with thickened edge (e. g. *Funa*, *Carinodrillia*, Fig. 16C), or trough-shaped in transverse section, sometimes with small barb near the tip (*Cheungbeia*, Fig. 16E), which may become semi-enrolled (e.g. *Pyrgospira*, *Pilsbryspira*, *Zonulispira*, Fig. 16B), to nearly hollow, where limbs overlap at significant length of the teeth (*Ptychobela*, Fig. 16D).

Remarks: Anatomically, Pseudomelatomidae is the most variable family of Conoidea. Most genera were formerly included in the subfamily Crassispirinae, but the nomenclaturally valid name for this clade is Pseudomelatomidae. Its constituents includes several taxa that were previously recognized as separate (sub)families: Zonulispirinae, characterized by semi-enrolled marginal radular teeth (a character found in several branches of the clade), and Pseudomelatominae, defined on the basis of the very characteristic solid marginal teeth and strongly developed rachidian.

Figure
16

The genus *Leucosyrinx* has long been a convenient genus for placement of turreted-fusiform species, mostly from deep water of the Atlantic and Indo-Pacific. Currently it is a mixture of species of probably different taxonomic position, and the membership of the genus needs revision. The type species from the northern Atlantic, *Leucosyrinx verrilli* (Dall, 1881), has a radula consisting of only duplex, rather robust marginal teeth (Powell, 1966: text fig. B12), a radula type also found by us in Indo-Pacific species (Fig. 16A). A second type of radula is found in *Sibogasyrinx*, originally described as a subgenus of *Leucosyrinx*, and which clusters in the molecular tree with the Cochlespiridae (see under that family). In our study, true *Leucosyrinx* appears to be sister to the Pseudomelatomidae, but this relationship has poor support. As a working hypothesis, we tentatively include *Leucosyrinx* in the Pseudomelatomidae.

Included genera:

- ?*Abyssocomitas* Sysoev & Kantor, 1986 ²
- Aguilaria* Taylor & Wells, 1994 ²
- ?*Anticomitas* Powell, 1942 ¹
- Antimelatoma* Powell, 1942 ¹
- Antiplanes* Dall, 1902 ²
 - [= *Rectiplanes* Bartsch, 1944]
 - [= *Rectisulcus* Habe, 1958]
- Austrotoma* Finlay, 1924 ¹
- ?*Benthodaphne* Oyama, 1962 ¹
- Burchia* Bartsch, 1944 ²
- Buridrilla* Olsson, 1942 ¹
- Calcatodrilla* Kilburn, 1988 ²
- Carinodrilla* Dall, 1919 ³
- ?*Carinoturris* Bartsch, 1944 ²
- Cheungbeia* Taylor & Wells, 1994 ³
- Comitas* Finlay, 1926 ³
- Compsodrilla* Woodring, 1928 ²
 - (*Mammillaedrilla* Kuroda and Oyama in Kuroda, Habe & Oyama, 1971)
- ?*Conorbela* Powell, 1951 ²
- Conticosta* Laseron, 1954 ¹
- Crassiclava* McLean, 1971 ²
- Crassispira* Swainson, 1840 ³
 - (*Crassispirella* Bartsch & Rehder, 1939)
- Cretaspira* Kuroda & Oyama in Kuroda, Habe & Oyama, 1971 ¹
- Dallspira* Bartsch, 1950 ²
- Doxospira* McLean, 1971 ²
- Eplodeira* Hedley, 1918 ²
 - [= *Epidirona* Hedley, 1931]
- Funa* Kilburn, 1988 ³
- Gibbaspira* McLean, 1971 ²
- Glossispira* McLean, 1971 ²
- Hindsiclava* Hertlein & Strong, 1955 ³
- [= *Turrigemma* Berry, 1958]
- Hormospira* Berry, 1958 ²
- Inquisitor* Hedley, 1918 ³
- Knefastia* Dall, 1919 ³
- ?*Kurilohadalia* Sysoev & Kantor, 1986 ²
- Kurodadrilla* Azuma, 1975 ¹
- ?*Leucosyrinx* Dall 1889 ³
- Lioglyphostoma* Woodring, 1928 ²
- Lioglyphostomella* Shuto, 1970 ¹
- Maesiella* McLean, 1971 ²
- Megasurcula* Casey, 1904 ²
- ?*Meggittia* Ray, 1977 ¹
- Miraclathurella* Woodring, 1928 ²
- ?*Monilispira* Bartsch & Rehder, 1939 ²
- Naudedrilla* Kilburn, 1988 ²
- Otitoma* Jousseaume, 1898 ³
 - [= *Thelecytharella* Shuto, 1969]
- ?*Paracomitas* Powell, 1942 ¹
- Pilsbryspira* McLean, 1971 ³
 - (*Nymphispira* McLean, 1971)
- Plicisyrix* Sysoev & Kantor, 1986 ²
- Pseudomelatoma* Dall, 1918 ³
- ?*Pseudotaranis* McLean, 1995 ²
- Ptychobela* Thiele, 1925 ³
 - [= ?*Brachytoma* Swainson, 1840]
- ?*Pyrgospira* McLean, 1971 ³
- ?*Rhodopetoma* Bartsch, 1944 ²
- Sediliopsis* Petuch, 1988 ¹
- ?*Shutonia* Van der Bijl, 1993 ¹
- Striospira* Bartsch, 1950 ²
- Tiariturris* Berry, 1958 ³
- Viridrilla* Bartsch, 1943 ¹
 - [= *Viridrillina* Bartsch, 1943]
- Zonulispira* Bartsch, 1950 ³

Figure
17

Figure
18

Clavatulidae Gray, 1853

(Figs 17A-F, 18)

Diagnosis: Shell medium-sized to rather large (usually 15-30 mm, up to 85 mm high), broad-fusiform to turreted-fusiform, with high spire and usually moderately long siphonal canal. Subsutural ramp usually well developed, with very shallow to rather deep anal sinus situated on its lower part or shifted abapically, with apex at almost a peripheral position. Sculpture variously developed, from almost smooth shell surface to well developed axial ribs and spiral cords. Protoconch only known as paucispiral, up to ca. 2.5 smooth whorls. Operculum with medio-lateral nucleus. Radular formula 1-(1-R-1)-1. The central formation is composed of very thin, broad, plate-like lateral teeth and a small, but sometimes strong, narrow rachidian. The central formation is variously developed, sometimes appearing as a pronounced tooth (Fig. 18B), sometimes clearly consisting of three elements (*Clionella sinuata*, see Taylor *et al.*, 1993; Fig. 18A, B), to its nearly complete reduction (Fig. 18C). Marginal teeth usually duplex, with sharp-edged major limb and a deep socket where an accessory limb is inserted, often with angulation distal to the socket (arrow in Fig. 18E). In *Toxiclionella* the marginal teeth are hypodermic, loosely enrolled, attached along their length to radular membrane, having two barbs at the tip and a tooth canal opening subterminally (Fig. 18D).

Remarks: The genus *Gemmuloborsonia* represents a sister group to the (Clavatulidae + Horaiclavidae) clade, but this node is not well supported. Because it resembles much more the Clavatulidae than the Horaiclavidae in terms of shell and radular characters (Fig. 18F), *Gemmuloborsonia* is provisionally included in the former family. If further studies support that hypothesis, then the diagnosis of the Clavatulidae should be amended to account for the presence of weak columellar pleats and multispiral Turridae-type protoconch present in *Gemmuloborsonia*.

Included genera:

- | | |
|--|---|
| <i>Benthoclionella</i> Kilburn, 1974 ² | (<i>Tomellana</i> Wenz, 1943) |
| <i>Caliendrula</i> Kilburn, 1985 ¹ | <i>Pusionella</i> Gray, 1847 ³ |
| <i>Clavatula</i> Lamarck, 1801 ³ | [= <i>Netrum</i> Philippi, 1850] |
| <i>Clionella</i> Gray, 1847 ² | <i>Scaevatula</i> Gofas, 1989 ¹ |
| ? <i>Fusiturris</i> Thiele, 1929 ¹ | <i>Toxiclionella</i> Powell, 1966 ² |
| [= <i>Tyrrhenoturris</i> Coen, 1929] | <i>Trachydrillida</i> Nolf & Swinnen, 2010 ¹ |
| ? <i>Gemmuloborsonia</i> Shuto, 1989 ³ | <i>Turricula</i> Schumacher, 1817 ³ |
| <i>Makiyamaia</i> Kuroda in MacNeil, 1961 ¹ | [= <i>Surcula</i> H. & A. Adams, 1853] |
| <i>Perrona</i> Schumacher, 1817 ³ | |

Figure
19

Horaiclavidae new family
(Figs 17 G-R, 19)

Type genus: *Horaiclavus* Oyama, 1954.

Diagnosis: Shell generally small, 5-25 mm, usually 7-15 mm high, shortly claviform, with relatively low spire and a short, truncated, poorly differentiated siphonal canal. Subsutural ramp usually poorly differentiated. Axial sculpture almost always present, usually as strong sinuate ribs. Spiral sculpture normally weak or obsolete, often with glossy shell surface. Anal sinus on subsutural slope, weak to moderately deep, often constrained by callus. Protoconch of up to 3.5 medially carinate but otherwise smooth whorls when multispiral, but usually paucispiral and smooth. Operculum with terminal nucleus. Radular formula: 1-0-0-0-1. Marginal teeth duplex, with lanceolate major limb and usually narrow accessory limb, which is inserted in a shallow socket. Major limb often with angulation lateral to the place of

accessory limb insertion. Rarely (some *Paradrillia*, cf. Kilburn, 1988, Fig. 17; *Inkinga*, Fig. 19B) the additional limb is of similar size to the major limb and the teeth become trough-shaped in transverse section, with a collar near the base (arrows in Fig. 19B). In several species of *Horaiclavus* the radular apparatus is absent.

Remarks: This family shares many characters with Pseudomelatomidae, conchologically differing by a small stout shell with short siphonal canal and usually poorly developed spiral sculpture.

Radulae also are rather similar to many representatives of Pseudomelatomidae and no clear cut distinctions were found. Genera currently included in Horaiclavidae have been usually included in the Crassispiridae (= Pseudomelatomidae), and the clear molecular-based division between the two clades seems to be not so clearly reflected in shell-based distinction. Therefore, the generic composition of the family is somewhat provisional and needs confirmation by further molecular data and/or a detailed analysis of conchological and radular characters.

Included genera:

Anacithara Hedley, 1922³
Anguloclavus Shuto, 1983³
?Aoteadrillia Powell, 1942²
?Austrocarina Laseron, 1954¹
Astrodrillia Hedley, 1918¹
 (*Regidrillia* Powell, 1942)
Buchema Corea, 1934²
Carinapex Dall, 1924³
Ceritoturris Dall, 1924³
Coronacomitas Shuto, 1983¹
Cypharoclavus Kuroda & Oyama in
 Kuroda, Habe & Oyama, 1971¹
?Darrylia García, 2008¹
Graciliclava Shuto, 1983¹
Haedropleura Bucquoy, Dautzenberg &
 Dollfus, 1883²
Horaiclavus Oyama, 1954³
Inodrillia Bartsch, 1943¹

[= *Inodrillara* Bartsch, 1943]
 [= *Inodrillina* Bartsch, 1943]
Inkinga Kilburn, 1988¹
Iwaaoa Kuroda, 1940¹
?Marshallena Allan, 1927²
 [= *Sugitanitoma* Kuroda, 1959]
Mauidrillia Powell, 1942²
?Micropleurotoma Thiele, 1929²
Naskia Sysoev & Ivanov, 1985¹
Numa Kilburn, 1988²
Paradrillia Makiyama, 1940³
 [= *Vexitomina* Powell, 1942]
 [= *Alticlavatula* MacNeil, 1960]
?Pseudexomilus Powell, 1944¹
Psittacodrillia Kilburn, 1988¹
Striatoguraleus Kilburn, 1994¹
?Thelecythara Woodring, 1928¹

Figure
20

Figure
21

Turridae H. Adams & A. Adams, 1853
(Figs 20, 21)

Diagnosis: shell of medium to large size (usually 20-30 mm, up to 110 mm high), short- to high-fusiform, usually with a high spire and a long (rarely short and truncated) siphonal canal. Axial sculpture weak or absent. Anal sinus on whorl periphery. Protoconch typically multispiral, up to 6 whorls, protoconch I smooth, protoconch II with arcuate axial riblets; reduced paucispiral protoconch usually smooth, may have arcuate axial riblets. Operculum fully developed, with terminal nucleus. Radular formula typically 1-(1:R:1)-1. Small and narrow rachidian and plate-like laterals are fused together, together constituting a central formation of different development (Kantor, 2006), varying from a well defined broad central tooth (Fig. 20B) to a tooth clearly formed of three elements (Fig. 20A) through gradual reduction of rachidian and/or laterals to complete absence (Fig. 20D). Marginals duplex, of variable morphology, from broadly oval and flattened (Fig. 21E, F) with nearly equally

developed limbs to awl-shaped and divided only in basal part (Fig. 21A). In most cases the major limb is large and knife-shaped, while accessory limb is dorsal and more weakly developed (Fig. 21). In *Iotyrris*, marginal teeth have equally developed limbs that form a shallow broad trough (Fig. 21D).

Remarks: This group is well defined by its usually narrowly fusiform shell with obsolete axial sculpture and peripheral anal sinus. However, the genus *Lucerapex*, although fully conforming conchologically, occupies a position on the tree (sister group to Turridae + Terebridae) that excludes it from the Turridae. It is nevertheless tentatively included here in Turridae.

Included genera:

<i>Cryptogemma</i> Dall, 1918 ²	[= <i>Oxytropa</i> Glibert, 1955]
<i>Decollidrilla</i> Habe & Ito, 1965 ¹	<i>Ptychosyrinx</i> Thiele, 1925 ³
<i>Epidirella</i> Iredale, 1913 ¹	[= <i>Bathybermudia</i> Haas, 1949]
[= <i>Austrogemmula</i> Laseron, 1954]	<i>Turridrupa</i> Hedley, 1922 ³
<i>Gemmula</i> Weinkauff, 1875 ³	<i>Turris</i> Batsch, 1789 ³ (For authorship of Turris, see Dubois & Bour (2010))
[= <i>Eugemmula</i> Iredale, 1931]	[= <i>Pleurotoma</i> Lamarck, 1799]
(<i>Pinguigemmula</i> MacNeil, 1960)	[= <i>Pleurotomus</i> Blainville, 1810]
<i>Iotyrris</i> Medinskaya & Syssoev, 2001 ³	(<i>Annulaturris</i> Powell, 1966)
<i>Kuroshoturris</i> Shuto, 1961 ¹	<i>Unedogemmula</i> MacNeil, 1961 ²
<i>Lophiotoma</i> Casey, 1904 ³	<i>Xenuroturris</i> Iredale, 1929 ³
[= <i>Lophioturris</i> Powell, 1964]	[= <i>Clamturris</i> Iredale, 1931]
? <i>Lucerapex</i> Iredale, 1936 ³	
<i>Polystira</i> Woodring, 1928 ³	

Figure
22

Strictspiridae McLean, 1971
(Figs 11E, 22A, B)

Diagnosis: Shell medium-sized, to 20 mm, claviform. Spiral and axial sculpture well developed, shoulder with a marked subsutural fold. Anal sinus deep, laterally directed. Parietal callus well developed. Protoconch paucispiral in all species examined, smooth. Operculum leaf-shaped, with terminal nucleus. Radula consisting of a pair of solid awl-shaped marginal teeth that have a prominent flange, located above the base of the tooth and firmly attached to the radular membrane.

Remarks: We did not obtain any material of Strictspiridae for molecular analysis and the position of the family group remains unclear. The shell resembles that of Pseudomelatomidae, and the shape of the radular marginal teeth is also somewhat similar to *Pseudomelatoma*, *Hormospira* and *Tiariturris*, but it differs in the absence of the rachidian. Among the examined species of *Strictspira*, *S. paxillus* is characterised by rather unusual characters, such as the absence of the venom gland together with very large and powerful odontophore (Kantor & Taylor, 1994). Until molecular data are available, we conservatively treat this family as valid, following Taylor *et al.* (1993).

Included genera:

<i>Strictspira</i> McLean, 1971 ²	<i>Cleospira</i> McLean, 1971 ²
--	--

Terebridae Mörch, 1852

Diagnosis: Shell medium-sized to large, 8-270 mm, usually 30-100 mm high, auger-shaped, with high to very high multiwhorled spire and flattened shell profile, aperture relatively small. Siphonal canal short, anal sinus not pronounced. Protoconch with up to 5 smooth whorls when multispiral. Radular formula 1-0-0-0-1 but radular apparatus absent in many species. Marginal teeth range from solid and curved to hypodermic, with or without small barb at the tip. Hypodermic teeth without solid bases. In some species (e.g. *Impages hectica*) the walls of the teeth are penetrated by numerous holes.

Included genera (After Bratcher & Cernohorsky, 1987; Terryn, 2007; Terryn & Holford, 2008):

<i>Cinguloterebra</i> Oyama, 1961 ³	[= <i>Decorihastula</i> Oyama, 1961]
<i>Clathroterebra</i> Oyama, 1961 ³	<i>Oxymeris</i> Dall, 1903 ³
<i>Duplicaria</i> Dall, 1908 ¹	[= <i>Nototerebra</i> Cotton, 1947]
[= <i>Diplomeriza</i> Dall, 1919]	<i>Pellifronia</i> Terryn & Holford, 2008 ¹
[= <i>Myurellisca</i> Bartsch, 1923]	<i>Perirhoe</i> Dall, 1908 ¹
[= <i>Pervicacia</i> Iredale, 1924]	[= <i>Abretiella</i> Bartsch, 1923]
<i>Euryta</i> H. & A. Adams, 1853 ¹	[= <i>Dimidacus</i> Iredale, 1929]
<i>Euterebra</i> Cotton & Godfrey, 1932 ³	<i>Pristiterebra</i> Taki & Oyama, 1954 ¹
[= <i>Gradaterebra</i> Cotton & Godfrey, 1932]	[= <i>Laeviacus</i> Oyama, 1961]
[= <i>Partecosta</i> Dance & Eames, 1966]	<i>Strioterebrum</i> Sacco, 1891 ³
<i>Granuliterebra</i> Oyama, 1961 ¹	[= <i>Brevimyurella</i> Oyama, 1961]
<i>Hastula</i> H. & A. Adams, 1853 ³	[= <i>Punctoterebra</i> Bartsch, 1923]
[= <i>Acuminia</i> Dall, 1908]	<i>Terebra</i> Bruguière, 1789 ³
[= <i>Egentelaria</i> Rehder, 1980]	[= <i>Myurellina</i> Bartsch, 1923]
[= <i>Hastulina</i> Oyama, 1961]	[= <i>Panaterebra</i> Olsson, 1967]
<i>Hastulopsis</i> Oyama, 1961 ³	[= <i>Paraterebra</i> Woodring, 1928]
<i>Impages</i> E.A. Smith, 1873 ¹	[= <i>Subula</i> Schumacher, 1817]
<i>Mazatlania</i> Dall, 1900 ¹	[= <i>Terebrum</i> Montfort, 1910]
<i>Microtrypetes</i> Pilsbry & Lowe, 1932 ¹	<i>Terenolla</i> Iredale, 1929 ³
<i>Myurella</i> Hinds, 1845 ³	<i>Triplostephanus</i> Dall, 1908 ¹

Figure
23

ACKNOWLEDGEMENTS

The authors thank B. A. Marshall, M. G. Harasewych, R. N. Kilburn, M. J. Tenorio, J. K. Tucker and D. Tippett for constructive comments and help to complete the lists of genera and subgenera. A. E. Fedosov, J. D. Taylor, J. McLean, M.G. Harasewych and D. L. Tippett helped with shell and radula pictures. J. A. Todd provided specimens for dissection and photography. Barbara Buge curated and photographed the voucher specimens in MNHN.

REFERENCES

- ADAMS, H. & ADAMS, A. 1853. *The genera of recent Mollusca*. John Van Voorts, London.
- BELLARDI, L. 1875. Novae pleurotomidarum Pedimonti et Liguria fossilium: dispositionis prodromus. *Bullettino della Societa Malacologica Italiana*, **1**: 16-24.
- BOUCHET, P. 1990. Turrid genera and mode of development: the use and abuse of protoconch morphology. *Malacologia*, **32**: 69-77.
- BOUCHET, P. & ROCROI, J.-P. 2005. Classification and nomenclator of Gastropod families. *Malacologia*, **47**: 1-397.

- BRATCHER, T. & CERNOHORSKY, W.O. 1987. *Living terebras of the world*. American Malacologists, Inc, New York.
- CHANG, C.-K. 1995. Reevaluation of the classification of Turridae. *Bulletin of Malacology, Taiwan*, **19**: 49.
- CHANG, C.-K. 2001. *Small turrids of Taiwan*. Thorsson, W. M., Taiwan.
- COSSMANN, M. 1896 *Essais de Paléoconchologie comparée, 2ème livraison*. Paris.
- DUBOIS, A. & BOUR, R. 2010. The distinction between family-series and class-series nomina in zoological nomenclature, with emphasis on the nomina created by Batsch (1788, 1789) and on the higher nomenclature of turtles. *Bonn Zoological Bulletin*, **57**: 149-171.
- FEDOSOV, A. & KANTOR, Y. 2008. Toxoglossan gastropods of the subfamily Crassispirinae (Turridae) lacking a radula, and a discussion of the status of the subfamily Zemaciinae. *Journal of Molluscan Studies*, **74**: 27-35.
- FISCHER, P. 1887. *Manuel de conchyliologie et de paléontologie conchyliologique*. F. Savy, Paris.
- HOLFORD, M., PUILLANDRE, N., TERRYN, Y., CRUAUD, C., OLIVERA, B.M. & BOUCHET, P. 2009. Evolution of the Toxoglossa venom apparatus as inferred by molecular phylogeny of the Terebridae. *Molecular Biology and Evolution*, **26**: 15-25.
- KANTOR, Y. & TAYLOR, J.D. 1994. The foregut anatomy of *Strictispira paxillus*. *Journal of Molluscan Studies*, **60**: 343-346.
- KANTOR, Y. & TAYLOR, J.D. 2000. Formation of marginal radular teeth in Conoidea (Neogastropoda) and the evolution of the hypodermic envenomation mechanism. *Journal of Zoology*: 251-262.
- KANTOR, Y.I. 2006. On the morphology and homology of the "central tooth" in the radula of Turrinae (Conoidea: Turridae). *Ruthenica*, **16**: 47-52.
- KANTOR, Y.I., MEDINSKAYA, A.I. & TAYLOR, J.D. 1997. Foregut anatomy and relationships of the Crassispirinae (Gastropoda, Conoidea). *Bulletin of the Natural History Museum of London (Zoology)*, **63**: 55-92.
- KILBURN, R.N. 1983. Turridae (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 1. Subfamily Turrinae. *Annals of the Natal Museum*, **25**: 549-585.
- KILBURN, R.N. 1985. Turridae (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 2. Subfamily Clavatulinae. *Annals of the Natal Museum*, **26**: 417-470.
- KILBURN, R.N. 1986. Turridae (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 3. Subfamily Borsoniinae. *Annals of the Natal Museum*, **27**: 633-720.
- KILBURN, R.N. 1988. Turridae (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 4. Subfamily Drilliinae, Crassispirinae and Strictispirinae. *Annals of the Natal Museum*, **29**: 167-320.
- KILBURN, R.N. 1991. Turridae (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 5. Subfamily Taraninae. *Annals of the Natal Museum*, **32**: 325-339.
- KILBURN, R.N. 1992. Turridae (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 6. Subfamily Mangeliinae, section 1. *Annals of the Natal Museum*, **33**: 461-575.
- KILBURN, R.N. 1993. Turridae (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 6. Subfamily Mangeliinae, section 2. *Annals of the Natal Museum*, **34**: 317-367.
- KILBURN, R.N. 1994. Turridae [s.l.] (Mollusca: Gastropoda) of southern Africa and Mozambique. Part 7. Subfamily Crassispirinae, section 2. *Annals of the Natal Museum*, **35**: 177-228.
- KILBURN, R.N. 1995. Turridae (s.l.) of southern Africa and Mozambique (Mollusca: Gastropoda: Conoidea) Part 8. Conidae: subfamily Mangeliinae, section 3. *Annals of the Natal Museum*, **36**: 261-269.

- MCLEAN, J.H. 1971. A revised classification of the family Turridae, with the proposal of new subfamilies, genera and subgenera from the Eastern pacific. *Veliger*, **14**: 114-130.
- MORRISON, J.P.E. 1965. On the families of Turridae. In: *The Thirty-First Annual Meeting of the American Malacological Union*, pp. 1-2. New-York.
- POWELL, A.W.B. 1942. The New-Zealand Recent and fossil Mollusca of the family Turridae. With general notes on turrid nomenclature and systematics. *Bulletin of the Auckland. Institute and Museum*, **2**: 1-192.
- POWELL, A.W.B. 1966. The molluscan families Speightiidae and Turridae. An evaluation of the valid taxa, both Recent and fossil, with lists of characteristics species. *Bulletin of the Auckland. Institute and Museum*, **5**: 5-184.
- PUILLANDRE, N., SAMADI, S., BOISSELIER, M.C., SYSOEV, A.V., KANTOR, Y.I., CRUAUD, C., COULOUX, A. & BOUCHET, P. 2008. Starting to unravel the toxoglossan knot: molecular phylogeny of the “turrids” (Neogastropoda: Conoidea). *Molecular Phylogenetics and Evolution*, **47**: 1122-1134.
- RÖCKEL, D., KORN, W. & KOHN, A.J. 1995. *Manual of the living Conidae. Vol. I, Indo-Pacific*. Christa Hemmen Verlag, Wiesbaden.
- TAYLOR, J.D. 1994. Foregut anatomy of the larger species of Turrinae, Clavatulinae and Crassispirinae (Gastropoda: Conoidea) from Hong-Kong. In: *Third International Workshop on the Malacofauna of Hong-Kong and Southern China* (Morton, B., ed.), pp. 195-213. Hong-Kong University Press, Hong-Kong.
- TAYLOR, J.D., KANTOR, Y.I. & SYSOEV, A.V. 1993. Foregut anatomy, feedings mechanisms and classification of the Conoidea (= Toxoglossa)(Gastropoda). *Bulletin of the Natural History Museum of London (Zoology)*, **59**: 125-170.
- TERRYN, Y. 2007. *A collectors guide to Recent Terebridae (Mollusca: Neogastropoda)*. ConchBooks, Hackenheim.
- TERRYN, Y. & HOLFORD, M. 2008. The Terebridae of the Vanuatu Archipelago with a revision of the genus *Granuliterebra* Oyama 1961. *Visaya, Supplement 3*:
- THIELE, J. 1929. *Handbuch der systematischen weichtierkunde*. G. Fischer, Jena.
- TUCKER, J.K. & TENORIO, M.J. 2009. *Systematic classification of Recent and fossil conoidean gastropods*. Conchbooks, Hackenheim, Germany.
- WILLIAMS, M. 2006. *Shallow-water Turridae of Florida and the Caribbean*. M. Williams, Tallevast, Florida.

Figure captions:

Figure 1. Comparison of the last two conoidean classifications with the new classification proposed in this article. **A.** Taylor *et al.* (1993) and Bouchet & Rocroi (2005). (¹ Elevated at the familial rank in Bouchet & Rocroi 2005. ² Absent in Taylor *et al.*, 1993). **B.** Proposed classification. **C.** Tucker & Tenorio (2009). † fossil taxa

Figures 2-21. Abbreviation and symbols: SL, shell length; * , sequenced species; **, sequenced specimen.

Figure 2. Shells.

A-N. Borsoniidae.

- A.** *Typhlomangelia* sp.**, MNHN IM200717931, Solomon Islands, SALOMON 2, st. CP2269, 07°45.1'S, 156°56.3'E, 768-890 m, SL 14.2 mm.
- B.** Borsoniidae gen. 1**, MNHN IM200717911, Philippines, PANGLAO 2005, st. CP2333, 9°38.2'N, 123°43.5'E, 584-596 m, SL 17.1 mm (radula: Fig. 3 D).
- C.** *Darbya lira* Bartsch, 1934, USNM 810565, off Puerto Rico, SL 19.4 mm.
- D.** *Borsonia* sp.**, MNHN IM200717932, Solomon Islands, SALOMON 2, st. CP2197, 08°24.4'S, 159°22.5'E, 897-1057 m, male, SL 24.6 mm (radula: Fig. 3B).
- E.** *Bathytmoma neocaledonica* Puillandre *et al.*, 2010**, paratype, MNHN IM200717857, Coral Sea, EBISCO, st. CP2551, SL 44 mm (radula: Fig. 3C).
- F.** *Zemacies excelsa* Sysoev & Bouchet, 2001**, MNHN IM200911056, New Caledonia, Musorstrom 4, st. DW226, 22°47'S, 167°22'E, 395m, SL 38 mm.
- G.** *Tomopleura reevii* (C. B. Adams, 1850)**, MNHN IM200717875, Philippines, PANGLAO 2004, st. T26, 09°43.3'N, 123°48.8'E, 123-135 m, SL 7.7 mm.
- H.** *Microdrillia cf. optima* (Thiele, 1925)**, MNHN IM200717887, Philippines, PANGLAO 2004, st. T36, 09°29.3'N, 123°51.5'E, 95-128 m, SL unknown (broken).
- I.** *Borsonellopsis chrysothermis* (Dall, 1919), SIO (Scripps Institution of Oceanography) M2530, San Diego Trough, off San Diego Co., California, 1244 m, SL 19.3 mm.
- J.** *Belaterricula turrita* (Strebel, 1908), USNM 897559, South Georgia Island, 54.3°S, 35.6°W, 238-247 m, SL 49.8 mm.
- K.** *Glyptaesopus oldroydi* (Arnold, 1903). 10 m, off Punta Abreojos, Baja California, Mexico, SL 8.3 mm.
- L.** *Tropidoturris fossata fossata* (Sowerby, 1903), holotype, SAMC A346, off Cape Vidal, Zululand, 146-182 m, SL 20.4 mm.
- M.** *Genota mitriformis* (Wood, 1828)**, MNHN IM200742293, Angola, 30-50m, SL 30 mm (radula: Fig. 3 A).
- N.** *Ophiodermella inermis* (Hinds, 1843), USNM 56092, W coast of North America, SL 40.7 mm.

O: Conorbidae.

- O.** *Benthofascis lozoueti* Sysoev & Bouchet, 2001**, MNHN IM200742331, New Caledonia, NORFOLK 2, st. DW2147, 22°50'S, 167°16'E, 496 m, SL 16.5 mm.

Abbreviation and symbols: SL, shell length; * , sequenced species; **, sequenced specimen.

Photo credits: B. Buge (A, B, D, E, G, H), M.G. Harasewych and D. Tippett (C, J, N), R.N. Kilburn (L), J. McLean (I, K).

Figure 3. Radulae. Borsoniidae.

- A.** *Genota mitriformis* (Wood, 1828), MNHN IM200742293 (shell: Fig. 2M).
- B.** *Borsonia* sp., MNHN IM200717932 (shell: Fig. 2D).
- C.** *Bathytmoma neocaledonica* Puillandre *et al.*, 2010, MNHN IM200717857 (shell: Fig. 2E).

D. Borsoniidae gen. 1, MNHN IM200717911 (shell: Fig. 2B).

Figure 4. Shells.

A-E. Clathurellidae.

A. *Nannodiella ravella* (Hedley, 1922)*, MNHN IM200742350, Philippines, PANGLAO 2004, st. T9, 09°33.5'N, 123°49.5'E, 97-120 m. SL 5.8 mm (radula: Fig. 5A).

B. *Comarmondia gracilis* (Montagu, 1803), MNHN, Le Brusc, Cap Sicié, Provence, France, 40-100 m, SL 23 mm.

C. *Etrema cf. tenera* (Hedley, 1899)**, MNHN IM200717869, Philippines, PANGLAO 2004, st. S21, 09°41.7'N, 123°50.9'E, 4-12 m, SL unknown (broken).

D. *Glyphostoma rostrata*. Sysoev & Bouchet, 2001, MNHN, New Caledonia, BATHUS 4, st. DW896, 20°16'S, 163°52'E, 315-350 m, SL 21.5 mm.

E. *Lienardia nigrotincta* (Montrouzier, 1872)*, MNHN, Touho, New Caledonia, 20°45.2'S, 165°16.3'E, intertidal, SL 6.9 mm (radula: Fig. 5B).

F. *Strombinoturris crockeri* Hertlein & Strong, 1951, LACM 747-37, off Isabel Island, Mexico, 18-33 m, SL 47.5 mm (courtesy of J. McLean).

G-I: Mitromorphidae.

G. *Lovellona atramentosa* (Reeve, 1849)*, MNHN, Philippines, PANGLAO 2004, st. M2, 09°32.8'N, 123°45.9'E, 0-2 m, SL 9.0 mm (radula: Fig. 6A).

H. *Mitromorpha metula* (Hinds, 1843)*, MNHN IM200742339, Philippines, PANGLAO 2004, st. B8, 09°37.1'N, 123°46.1'E, 3 m, SL 3.1 mm (radula: Fig. 6B).

I. *Anarithma* sp.*, MNHN, Philippines, PANGLAO 2004, st. S5, 09°37.1'N, 123°46.1'E, 2-4 m, SL 6.9 mm (radula: Fig. 6C).

Abbreviation and symbols: SL, shell length; *, sequenced species; **, sequenced specimen.

Photo credits: B. Buge (D, F, G, K), M.G. Harasewych and D. Tippett (H), P. Maestrati (E, M), J. McLean (O).

Figure 5. Radulae. Clathurellidae.

A. *Nannodiella ravella* (Hedley, 1922), MNHN IM200742350 (shell: Fig. 4A).

B. *Lienardia nigrotincta* (Montrouzier, 1872), MNHN (shell: Fig. 4E).

C. *Lienardia jusseaumei* (Hervier, 1896), MNHN, Philippines, PANGLAO 2004, st. B7, 9°35.9'N, 123°51.8'E, 4-30 m.

Figure 6. Radulae. Mitromorphidae.

A. *Lovellona atramentosa* (Reeve, 1849), MNHN (shell: Fig. 4G).

B. *Mitromorpha metula* (Hinds, 1843), MNHN IM200742339 (shell: Fig. 4H).

C. *Anarithma* sp., MNHN (shell: Fig. 4 I).

Figure 7. Shells. Mangeliidae.

A. *Bela clarae* Penas & Rolan, 2008, paratype, MNHN Moll 21396, Spain, Barcelona, Siges, Vallcarca, 45-60 m, SL 7 mm.

B. *Macteola segesta* (Duclos in Chenu, 1850), MNHN, LIFOU 2000, st. 1425, 20°46.8'S, 167°07.2'E, 4-5 m, SL 4.6 mm.

C. *Benthomangelia cf. trophonoidea* (Schepman, 1913)**, MNHN IM200717835, Vanuatu, BOA1, st. CP2462, 16°38'S, 167°57'E, 618-641 m, SL 21 mm (radula: Fig. 8A).

D. *Citharomangelia* sp., MNHN, Island Futuna, MUSORSTOM 7, st. DW504, 14°20'S, 178°04'E, 300-390 m, SL 15.4 mm.

E. *Anticlinura* sp.**, MNHN IM200742513, Solomon Islands, SALOMON 2, st. CP2182, 08°47.0'S, 159°37.9'E, 762-1060 m, SL 17.3 mm (radula: Fig. 8E).

- F.** *Eucithara* cf. *coronata* (Hinds, 1843)**, MNHN IM200717900, Philippines, PANGLAO 2004, st. B8, 09°37.1'N, 123°46.1'E, 3 m, SL 8.7 mm (radula: Fig. 8D).
- G.** *Leiocithara* sp., MNHN, New Caledonia, BATHUS 2, st. DW719, 22°48'S, 167°16'E, 444-445 m, SL 5.4 mm.
- H.** *Oenopota uschakovi* Bogdanov, 1985, holotype, ZIN 37974/1, Iturup Island, Kurile Islands, 44°24.2'N, 147°09'01"E. 160 m, SL 10.3 mm.
- I.** *Genotina adamii* (Bozzetti, 1994), MNHN, LIFOU 2000, st. DW1650, 20°54.15'S, 167°01.7'E, 120-150 m, SL 20.0 mm.
- J.** *Vitjazinella multicostata* Sysoev, 1988, Paratype, ZMMU 22356, Izu-Bonin Trench, 33°18'N, 149°46'E., 6096 m, SL 7.2 mm.
- K.** *Oenopotella ultraabyssalis ultraabyssalis* Sysoev, 1988, holotype, ZMMU Lc-22681, Kurile-Kamchatka trench, 49°29' N, 158°41' E, 7210-7230 m, SL 12.8 mm.
- L.** *Pseudorhaphitoma* sp., Philippines, Balicasag Island, lumun-lumun net, 35-40m deep, SL 19.7 mm
- M.** *Paraspirotropis simplicissima* (Dall, 1907), ZIN, off Shikotan I., S Kurile Islands, 1450-1530 m, SL 28.0 mm.
- N.** *Heterocithara* sp.**, MNHN IM200717884, Philippines, PANGLAO 2004, st. L46, 09°30.9'N, 123°41.2'E, 90-110 m, SL 3.4 mm.
- O.** *Toxicochlespira pagoda* Sysoev & Kantor, 1990**, MNHN IM200717925, Solomon Islands, SALOMON 2, st. CP2227, 6°37.2'N, 156°12.7'E, 508-522 m, SL 18.0 mm (radula: Fig. 8B).
- Abbreviation and symbols: SL, shell length; *, sequenced species; **, sequenced specimen. Photo credits: B. Buge (F, O), P. Maestrati (B, D, G, I, L).

Figure 8. Radulae. Mangeliidae.

- A.** *Benthomangelia trophonoidea* (Schepman, 1913), MNHN IM200717835 (shell: Fig. 7C).
- B.** *Toxicochlespira pagoda* Sysoev & Kantor, 1990, MNHN IM200717925 (shell: Fig. 7L).
- C.** Mangeliinae gen. 2. MNHN IM200910331, Philippines, PANGLAO 2004, st. S26, 9°41.50'N, 123°51.00'E, 21m, SL unknown (broken).
- D.** *Eucithara* cf. *coronata* (Hinds, 1843), MNHN IM200717900 (shell: Fig. 7F).
- E.** *Anticlinura* sp., MNHN IM200742513 (shell: Fig. 7E).
- F.** Mangeliidae gen. sp., MNHN, Philippines, PANGLAO 2004, st. T26, 9°43.3'N, 123°48.8'E, 123-135 m.
- G.** *Mangelia powisiana* (Dautzenberg, 1887). Plymouth, England, after Taylor *et al.* (1993).

Figure 9. Shells. Raphitomidae.

- A.** *Hemilienardia calcicincta* (Melvill & Standen, 1895)*, Philippines, PANGLAO 2004, st. B14, 09°38.5'N, 123°49.2'E, 2-4 m SL 4.6 mm.
- B.** *Hemilienardia malleti* (Recluz, 1852), MNHN, New Caledonia, MONTROUZIER, st. 1245, 20°45.2'S, 165°16.3'E, intertidal, SL 4.4 mm.
- C.** *Rimosodaphnella* sp.**, MNHN IM200717836, Vanuatu, BOA1, st. CP2462, 16°37.5'S, 167°57.4'E, 618-641 m. SL 15.1 mm.
- D.** *Veprecula* sp.**, MNHN IM200717883, Philippines, PANGLAO 2004, st. L46, 09°30.9'N, 123°41.2'E, 90-110 m, SL 3.4 mm.
- E.** *Glyphostomoides* sp.**, MNHN IM200717892, Philippines, PANGLAO 2004, st. T39, 09°30'N, 123°50.4'E, 100-138 m, SL 7.0 mm.
- F.** *Aliceia* sp., MNHN, Wallis Islands, MUSORSTOM 7, st. DW604, 13°21'S, 176°08'W, 415-420 m, SL 4.0 mm.
- G.** *Thatcheriasyrinx orientis* (Melvill, 1904), MNHN, New Caledonia, 20°49'S, 165°19'E, 105-110 m, SL 3.6 mm.

- H.** *Tritonoturris cumingii* (Powys in Powys and Sowerby, 1835), MNHN, New Caledonia, SL 20 mm.
- I.** *Rocroithys niveus* Sysoev & Bouchet, 2001, holotype, MNHN, Vanuatu, 15°08'S 167°12'E, 312-314 m, SL 25.5 mm.
- J.** *Pleurotomella* sp.**, MNHN IM200717848, Coral Sea, EBISCO, st. DW2625, 20°05'S, 160°19'E, 627-741 m, SL 20.6 mm.
- K.** *Taranis* sp.**, MNHN IM200742296, Philippines, AURORA 07, st. CP2749, 15°56.6'N, 121°50.2'E, 473 m, SL 6 mm. A07 1717.
- L.** *Nepotilla* sp., MNHN, New Caledonia, SMIB8, st. DW146-147, 24°55'S, 168°22', 508-532 m, SL 1.5 mm.
- M.** *Eucyclotoma cymatodes* (Hervier, 1899)**, MNHN IM200717903, Philippines, PANGLAO 2004, st. S12, 09°29.4'N, 123°56.0'E, 6-8 m, SL 5.0 mm.
- N.** *Buccinaria pendula* Bouchet & Sysoev, 1997, MNHN, New Caledonia, BATHUS 1, st. DE697, 20°34'S, 164°58'E, 570-650m, SL 7.2 mm.
- O.** *Microdaphne morrisoni* Rehder, 1980, MNHN, New Caledonia, SL 4.5 mm.
- P.** *Cryptodaphne* sp., MNHN, New Caledonia, MUSORSTOM 4, st. DW150, 19°07'S, 163°22'E, 110m, SL 11.2 mm.
- Q.** *Kermia aureotincta* (Hervier, 1897)**, MNHN IM200717878, Philippines, PANGLAO 2004, st. B25, 09°29'N, 123°56.1'E, 16 m, SL 4.5 mm.
- R.** *Teretiopsis cf. hyalina* Sysoev & Bouchet, 2001**, MNHN IM200717845, Coral Sea, EBISCO, st. CP2651, 21°29'S, 162°36'E, 883-957 m, SL 22.8 mm.
- S.** *Daphnella itonis* Sysoev & Bouchet, 2001, holotype, MNHN, New Caledonia, 22°48'S 167°16'E, 444-445 m, SL 66.0 mm.
- T.** *Phymorhynchus major* Warèn & Bouchet, 2001, holotype, MNHN, East Pacific Rise, Site Barbecue, 09°50'N, 104°17'W, 2505 m, SL 72.0 mm.
- U.** *Gymnobela* sp.**, MNHN IM200717841, Coral Sea, EBISCO, st. CP2648, 21°32'S, 162°30'E, 750-458 m, SL 25.7 mm.
- V.** *Thatcheria mirabilis* Angas, 1877**, MNHN IM200717924, Solomon Islands, SALOMON 2, st. CP2184, 08°16.9'S, 159°59.7'E, 464-523 m, SL 32.5 mm.
- Abbreviation and symbols: SL, shell length; *, sequenced species; **, sequenced specimen. Photo credits: B. Buge (C-E, J, Q, R, U, V), P. Maestrati (F-G, L, P).

Figure 10. Radulae. Raphitomidae.

- A.** *Buccinaria pendula* Bouchet & Sysoev, 1997, MNHN, New Caledonia, BATHUS 4, st. CP948, 20°33'S, 164°57'E, 533-610m.
- B.** *Daphnella pulvisculus* Chino, 2006, MNHN, New Caledonia, BATHUS 4, st. DW927, 18°56'S, 163°22'E, 444-452m.
- C.** *Gymnobela yoshidai* Kuroda & Habe in Habe, 1962, MNHN, Norfolk ridge, NORFOLK 2, st. DW2058, 24°40'S, 168°40'E, 591-1032m.
- D.** *Kermia irretita* (Hedley, 1899), MNHN, New Caledonia, LIFOU 2000, st. 1419, 20°55.6'S, 167°04.5'E, 0-5m.
- E.** *Daphnella cladara* Sysoev & Bouchet, 2001, MNHN, Norfolk Ridge, LITHIST, st. CP09, 24°53'S, 168°22'E, 518-540m.
- F.** *Daphnella mitrellaformis* (Nomura, 1940), MNHN, New Caledonia, LIFOU 2000, st. DW1649, 20°54'S, 167°01'E, 150-200m.
- G.** *Miowateria* sp., MNHN, Fidji, MUSORSTOM 10, st. CP1354, 17°43'S, 178°55'E, 959-963m.
- H.** *Spergo fusiformis* (Kuroda & Habe in Habe, 1962), MNHN, Tongatapu, Tonga, BORDAU 2, st. CP1566, 21°02'S, 175°18'W, 530-531m.

Figure 11. Shells.

A-D. Cochlespiridae.

A. *Nihonia maxima* Sysoev, 1997, holotype, MNHN, Indonesia, 09°23'S 131°09'E, 275 m - 246 m, SL 128 mm.

B. *Sibogasyrinx* sp.**, MNHN IM200717701, Vanuatu, BOA1, st. CP2432, 14°59.7'S, 166°55.0'E, SL 46.0 mm (radula: Fig. 12B, C).

C. *Aforia multisprialis* Dell, 1990, USNM 870023, N of Livingston Island, South Shetland Islands, 62°S, 61.1°W, 1437 m, SL 45.6 mm (courtesy of USNM)

D. *Cochlespira pulchella* (Schepman, 1913)**, MNHN IM200717920, PANGLAO 2005, st. CP2340, 09°29.4'N, 123°44.4'E, SL 22.0 mm.

E. Strictispiridae.

E. *Strictispira solida* (C.B.Adams, 1850), USNM 900425, E side of Marquesas Keys, Florida Keys, SL 17.5 mm (courtesy of M.G. Harasewych, D. Tippett).

F-L. Drillidae.

F. *Plagiostropha caledoniensis* Wells, 1995, S New Caledonia, SL 12.2 mm.

G. *Imaclava unimaculata* (Sowerby I, 1834)* subadult, det. J.A. Todd, NHMUK MOEA 20100529 Gulf of Panama, JTD-00-46, 08°31.37'N, 79°05.79'W, 24-25 m , SL 25.5 mm.

H. *Clavus canalicularis* (Roeding, 1798)*, MNHN IM200742345, Philippines, PANGLAO 2004, st. B11, 09°29.4'N, 123°56.0'E, 2-4 m, SL 19 mm.

I. *Cruziturricula arcuata* (Reeve, 1843)* subadult, det. J. A. Todd, YK; NHMUK 20100542, Gulf of Panama, JTD-00-34, 08°26.24'N, 79°09.14'W, 66-68 m, SL = 17.6 mm (radula: Fig. 13E).

J. *Conopleura striata* Hinds, 1844*, New Caledonia, MONTROUZIER, st. 1311, 20°40.4'S, 164°14.9'E, 10-60 m, SL 15.6 mm.

K. *Iredalea pupoidea* (H. Adams, 1872)*, MNHN, New Caledonia, MONTROUZIER, st. 1240, 20°46.5'S, 165°14'E-165°15'E, 2m, SL 6.2 mm.

L. *Splendrillia* sp.**, MNHN IM200717847, Coral Sea, EBISCO, st. DW2617, 20°06'S, 160°22'E, 427-505 m, SL 16.4 mm (radula: Fig. 13A, B).

Abbreviation and symbols: SL, shell length; *, sequenced species; **, sequenced specimen. Photo credits: B. Buge (B-D, F), A. Fedosov (I, G), M.G. Harasewych and D. Tippett (C, E),

Figure 12. Radulae. Cochlespiridae.

A. *Cochlespira radiata* (Dall, 1889), MNHN, SE Brazil, after Kantor & Taylor, 2000.

B-C. *Sibogasyrinx* sp., MNHN IM200717701 (shell: Fig. 11B).

Figure 13. Radulae. Drillidae.

A-B. *Splendrillia* sp., MNHN IM200717847 (shell: Fig. 11F).

C. *Clavus exasperatus* (Reeve, 1843), MNHN, New Caledonia, LIFOU 2000, st. 1420, 20°47.7'S, 167°09.35'E, 4-5m.

D. *Imaclava pilsbryi* (Bartsch, 1950), after Kantor & Taylor, 2000.

E. *Cruziturricula arcuata* (Reeve, 1843) **, NHMUK MOEA 20100541, Gulf of Panama, JTD-00-34, 08°26.24'N, 79°09.14'W, 66-68 m (shell: Fig. 11I).

Symbol: **, sequenced specimen.

Figure 14. Shells. Pseudomelatomidae.

A. *Comitas* sp. **, MNHN IM200717918, Philippines, PANGLAO 2005, st. CP2388, 09°26.9'N, 123°34.5'E, 762-786 m, SL 113.8 mm (radula: Fig. 15C, D).

B. *Knefastia tuberculifera* (Broderip & Sowerby, 1829)*, NHMUK MOEA 20100534, Gulf of Panama, JTD-00-18, 08°19.50'N, 78°47.71'W, 25-32 m. SL 36.6 mm.

- C.** *Otitoma* sp.**, MNHN IM200717905, Philippines, PANGLAO 2005, st. CP2348, 09°26.6'N, 123°52.5'E, 196-216 m, SL 17.2 mm.
- D.** *Pyrgospira aenone* (Dall, 1919)**, det. J. A. Todd, YK, NHMUK MOEA 20100539, Gulf of Panama, JTD-00-18, 08°19.50'N, 78°47.71'W, 25-32 m. SL 36.2 mm.
- E.** *Inquisitor* sp.**, MNHN IM200717851, Coral Sea, EBISCO, st. DW2625, 20°05'N, 160°19'E, 627-741 m, SL 32.2 mm (radula: Fig. 16F).
- F.** *Funa incerta* (Smith, 1877)**, NHMUK MOEA 20100553, coll. B. Morton, st. 70, off southern Hong Kong, SL 42.8 mm.
- G.** *Leucosyrinx* sp.**, MNHN IM200717846, Coral Sea, EBISCO, st. CP2600, 19°38'S, 158°46'E, 603-630 m, SL 27.1 mm (radula: Fig. 16A).
- H.** *Hindsiclava alesidota* (Dall, 1889)**, det. J. A. Todd. NHMUK MOEA 20100524, Lower Florida Keys, JTD-01-15, 24°33.47'N, 81°07.72'W, 117-148 m. SL 37.9 mm.
- I.** *Conorbela antarctica* (Strebel, 1908) USNM 894465, Zavodovski Island, South Sandwich Islands, 56.27°S, 27.5°W, 208-375 m, SL 23.6 mm (courtesy USNM).
- J.** *Carinodrillia dichroa* Pilsbry & Lowe, 1932*, det. J. A. Todd. NHMUK MOEA 20100531 Gulf of Panama, JTD-00-18, 08°19.50'N, 78°47.71'W, 25-32 m SL 23.0 mm (radula: Fig. 16C).
- K.** *Cheungbeia robusta* (Hinds, 1839)*, NHMUK MOEA 20100558, coll. B. Morton, st. 70, off southern Hong Kong, South China Sea, SL 21.2 mm (radula: Fig. 16E).
- L.** *Plicisyrinx vitjazi* Sysoev, Kantor, 1986, holotype, ZMMU Lc-22382, E of northern Honshu, Japan, 40°55.2'N, 144°53.3'E, 3880-4000 m, SL 13.9 mm.
- M.** *Maesiella maesae* McLean & Poorman, 1971, LACM 1513 holotype, Guaymas, Sonora, Mexico SL 9.2 mm.
- N.** *Kurilohadalia elongata* Sysoev & Kantor, 1986, holotype, ZMMU Lc-22401, Kurile-Kamchatka trench, 43°55' N, 149°47' E, 8080-8160 m, SL 12.5 mm.
- O.** *Crassispira quadrilirata* (E. A. Smith, 1882)**, MNHN IM200717755, Philippines, PANGLAO 2004, st. L46, 09°30.9'N, 123°41.2'E, 90-110 m, SL 14.0 mm.
- P.** *Epideira sibogae* (Schepman, 1913), MNHN, Vanuatu, MUSORSTOM 8, st. DW1097, 15°05'S, 167°11'E, 281-288 m, SL 24.1 mm.
- Q.** *Tiariturris spectabilis* Berry, 1958**, det. J. A. Todd, NHMUK MOEA 20100540, Gulf of Panama, JTD-00-34, 08°26.24'N, 79 09.14'W, 66-68 m SL 48.7 mm (radula: Fig. 15A).
- R.** *Pseudomelatoma penicillata* Carpenter, 1864, USNM 130590, California, SL 28.1 mm.
- S.** *Megasurcula carpenteriana* (Gabb, 1865), USNM 523713, California, SL 84.1 mm. (courtesy of M.G. Harasewych, D. Tippett).
- T.** *Antiplanes sanctioannis* (E. A. Smith, 1875), ZMMU Lc-17096, off northern Kurile Islands, 49°40.9'N, 154°41'E, 142 m, SL 37.2 mm.
- U.** *Naudedrillia* sp., New Caledonia, VAUBAN, st. 3, 22°17'S, 167 °12'E, 390 m, SL 19.0 mm.
- V.** *Zonulispira* sp. *, NHMUK MOEA 20100537, det. J. A. Todd, Gulf of Panama, JTD-00-18, 08°19.50'N, 78°47.71'W, 25-32 m SL.21.2 mm (radula: Fig. 16B).
- Abbreviation and symbols: SL, shell length; *, sequenced species; **, sequenced specimen. Photo credits: B. Buge (A, G), A. Fedosov (B-F, H, J, K, Q), M.G. Harasewych and D. Tippett (R, S), P. Maestrati (U), J. McLean (M).

Figure 15. Radulae Pseudomelatomidae.

- A.** *Tiariturris spectabilis* Berry, 1958**, NHMUK MOEA 20100540, Gulf of Panama, JTD-00-34, 08°26.24'N, 79 09.14'W, 66-68 m (shell: Fig. 14Q).
- B.** *Comitas onokeana vivens* Dell, 1956, MNHN, New Caledonia, MONTROUZIER, st. 1269, after Kantor & Taylor, 2000.
- C-D.** *Comitas* sp., MNHN IM200717918 (shell: Fig. 14A).

E-F. *Crassiclava turricula* (Sowerby, 1834). Off Nacascola, West side of Bahia Culebra, Costa Rica, after Kantor *et al.*, 1997.

Figure 16. Radulae. Pseudomelatomidae.

- A.** *Leucosyrinx* sp., MNHN IM200717846 (shell: Fig. 14 G).
B. *Zonulispira* sp.**, NHMUK MOEA 20100536, Gulf of Panama, JTD-00-18, 08°19.50'N, 78°47.71'W, 25-32 m (shell: Fig. 14V).
C. *Carinodrillia dichroa* Pilsbry & Lowe, 1932**, NHMUK MOEA 20100530, Gulf of Panama, JTD-00-18, 08°19.50'N, 78°47.71'W, 25-32 m. (shell: Fig. 1 J).
D. *Ptychobela suturalis* (Gray, 1838)**, det. J. A. Todd, YK, NHMUK MOEA 20100560, coll. B. Morton, off southern Hong Kong, Sta. 71.
E. *Cheungbeia robusta* (Hinds, 1839)*, NHMUK MOEA 20100557, coll. B. Morton, off southern Hong Kong, Sta. 70. (shell: Fig. 14K).
F. *Inquisitor* sp., MNHN IM200717851 (shell: Fig. 14E).
Symbols: * , sequenced species; **, sequenced specimen.

Figure 17. Shells.

A-F. Clavatulidae.

- A.** *Gemmuloborsonia colorata* (Sysoev & Bouchet, 2001)**, MNHN IM200717849, Coral Sea, EBISCO, st. DW2619, 20°06'S, 160°23'E, 490-550 m, SL 45.5 mm (radula: Fig. 18F).
B. *Pusionella compacta* Strebel, 1914**, MNHN IM200717830, Cacuaco, Bengo Prov., Angola, 5-10 m, SL 24.1 mm (radula: Fig. 18C).
C. *Clavatula xanteni* Nolf & Verstraeten, 2006**, MNHN IM200717829, Ilha de Luanda, Luanda Prov., Angola, 40-60 m, SL 20.5 mm (radula: Fig. 18E).
D. *Perrona subspirata* (von Martens, 1902)**, MNHN IM200717833, Moçâmedes, Namibe Prov., Angola, 5-10 m, SL 24.9 mm.
E. *Turridula javana* (Linnaeus, 1767), S India, SL 63 mm.
F. *Fusiturris undatiruga* (Bivona, 1838), Sierra Leone, 60-80 m, SL 43 mm.

G-R: Horaiclavidae.

- G.** *Marshallena philippinarum* (Watson, 1882), MNHN Philippines, PANGLAO 2005, st. CP2334, 09°37.5'N, 123°40.2'E, 606-631 m, SL 26 mm.
H. *Horaiclavus splendidus* (A. Adams, 1867)**, MNHN IM200717840, Coral Sea, EBISCO, st. DW2631, 21°03'S, 160°44'E, 372-404 m, SL 18.0 mm (radula: Fig. 19D).
I. *Ceritoturris pupiformis* (Smith, 1884)**, MNHN IM200717888, Philippines, PANGLAO 2004, st. T36, 09°29.3'N, 123°51.5'E, 95-128 m, SL 3.4 mm (radula: Fig. 19C).
J. *Inkinga* sp., New Caledonia, NORFOLK 1, st. DW1737, 22°52'S, 167°12'E, 343-400 m, SL 11.1 mm. (radula: Fig. 19B).
K. Horaiclavidae gen. 1**, MNHN IM200742501, Solomon islands, SALOMON 2, st. CP2219, 07°58.3'S, 157°34.4'E, 650-836 m, SL 10.1 mm.
L. *Buchema interpleura* (Dall & Simpson, 1901), USNM 900896, Lemon Cay, San Blas Islands, Panama, Caribbean Sea, 2 m, SL 10.0 mm.
M. *Carinapex minutissima* (Garrett, 1873), MNHN, New Caledonia, LIFOU 2000, st. 1457, 20°46.8'S, 167°02.75'E, 5-10 m, SL 3.1 mm.
N. *Paradrillia* sp.**, MNHN IM200742475, Philippines, PANGLAO 2005, st. CP2396 09°36.3'N, 123°42.0'E, 609-673 m, SL 11.5 mm (radula: Fig. 19A).
O. *Haedropleura septangularis* MNHN, Finistère, France, 20-30 m, SL 11 mm.
P. *Anguloclavus* sp.**, MNHN IM200717908, PANGLAO 2005, st. CP2332, 09°38.8'N, 123°45.9'E, SL 19.0 mm.

Q. *Iwaoa* sp., New-Caledonia, MUSORSTOM6, st. DW489, 20°48'S, 167°06'E, 700 m, SL 13.5 mm.

R. *Anacithara themeropsis* (Melvill & Standen, 1896), MNHN, LIFOU 2000, st. 1427, 20°47.6'S, 167°10.2'E, 10 m, SL 4.5 mm.

Abbreviation and symbols: SL, shell length; *, sequenced species; **, sequenced specimen. Photo credits: B. Buge (B-D, H, I, P, O), M.G. Harasewych and D. Tippett (L), P. Maestrati (J, Q, R).

Figure 18. Radulae. Clavatulidae.

A-B. *Turricula nelliae* (E.A. Smith, 1877)**, NHMUK MOEA 20100551, coll. D.G. Reid, Danang, Vietnam.

C. *Pusionella compacta* Strebel, 1914, MNHN IM200717830 (shell: Fig. 17B).

D. *Toxiclionella tumida* (Sowerby, 1870), South Africa, after Kantor, Taylor (2000).

E. *Clavatula xanteni* Nolf & Verstraeten, 2006, MNHN IM200717829 (shell: Fig. 17C).

F. *Gemmuloborsonia colorata* (Sysoev & Bouchet, 2001), MNHN IM200717849 (shell: Fig. 17A).

Symbol: **, sequenced specimen.

Figure 19. Radulae. Horaiclavidae.

A. *Paradrillia* sp., MNHN IM200742475 (shell: Fig. 17N).

B. *Inkinga* sp., MNHN (shell: Fig. 17 J).

C. *Ceritoturris pupiformis* (Smith, 1884)**, MNHN IM200717888 (shell: Fig. 17I).

D. *Horaiclavus splendidus* (Adams, A., 1867), MNHN IM200717840 (shell: Fig. 17H).

Symbol: **, sequenced specimen.

Figure 20. Shells. Turridae.

A. *Xenuroturris legitima* Iredale, 1929**, MNHN IM200717684, Vanuatu, SANTO 2006, st. DR087, 15°38.5'S, 167°15.1'E, 13 m, SL 57.0 mm (radula: Fig. 21C).

B. *Iotyrris cingulifera* (Lamarck, 1822)**, MNHN IM200717685, Vanuatu, SANTO 2006, st. FS84, 15°33.6'S, 167°16.6'E, 8-9 m, SL 15.5 mm (radula: Fig. 21D).

C. *Decollidrillia nigra* Habe & Ito, 1965, ZMMU uncataloged, southern Kurile Islands, SL 12.8 mm.

D. *Lophiotoma acuta* (Perry, 1811)**, MNHN IM200717860, Philippines, PANGLAO 2004, st. R44, 09°33.3'N, 123°43.9'E, 2 m, SL 44.0 mm.

E. *Turridrupa acutigemmatata* (E. A. Smith, 1877), New Caledonia, 46 m, SL 26.5 mm.

F. *Turris babylonia* (Linnaeus, 1758)**, MNHN IM200717754, Philippines, PANGLAO 2004, st. R42, 09°37.1'N, 123°52.6'E, 8-22 m, SL 79.4 mm.

G. *Gemmula rarimaculata* (Kuroda & Oyama, 1971)**, MNHN IM200717838, Coral Sea, EBISCO, st. DW2533, 22°18'S, 159°28'E, 360-370 m, SL 13.7 mm.

H. *Ptychosyrinx chilensis* Berry, 1968, USNM 870005, S of Coquimbo, Chile, 31.1° S, 71.8° W, 179-187 m, SL 21.1 mm (courtesy of USNM)

I. *Lucerapex cf. casearia* (Hedley & Petterd, 1906)**, MNHN IM200742448, Philippines, PANGLAO 2005, st. CP2363, 09°06.0'N, 123°25.0'E, 437-439 m, SL 21.0 mm (radula: Fig. 21 E).

J. *Cryptogemma cornues* (Okutani, 1966), ZIN 58809/1, off Shikotan Id., Kurile Islands, 1450-1530 m, 12.3 mm.

Abbreviation and symbol: SL, shell length; **, sequenced specimen.

Figure 21. Radulae. Turridae.

- A.** *Turridrupa* cf. *armillata* (Reeve, 1845), MNHN IM200740773, Coral Sea, EBISCO, st. DW2607, 19°33'S, 158°40'E, 400-413m.
- B.** *Gemmula kieneri* (Doumet, 1840), MNHN, Vanuatu, MUSORSTOM 8, st. CP1123, 15°07'S, 166°55'E, 262-352m.
- C.** *Xenuroturris legitima* Iredale, 1929, MNHN IM200717684 (shell: Fig. 20A).
- D.** *Iotyrris cingulifera* (Lamarck, 1822), MNHN IM200717685 (shell: Fig. 20B).
- E.** *Lucerapex* cf. *casearia* (Hedley & Petterd, 1906), MNHN IM200742448 (shell: Fig. 20I).
- F.** *Turris crispa* (Lamarck, 1816), MNHN, Ile Ouen-Baie du Prony, st. 80, 22°31'S, 166°28'E, 33m, SS 39 mm.

Figure 22. Radulae.

- A, B.** Strictspiridae. *Strictispira paxillus* (Reeve, 1845), British Virgin Islands, ANSP. A. teeth detached from the membrane and showing the large median flange; B. Two teeth, showing the large median flange attached to the membrane (courtesy of J.D. Taylor, NHM).
- C.** Conorbidae. *Benthofascis lozoueti* Sysoev & Bouchet, 2001, MNHN IM200742331 (shell: Fig. 2O).

Figure 23. Protoconchs. Only multispiral protonchs are illustrated.

- A.** Drillidae. *Iredalea* sp. (= *aureolinea*), MNHN, New-Caledonia, LIFOU 2000, st. 1441, 20°46.4'S, 167°02.0'E, 20m.
- B.** Pseudomelatomidae. *Inquisitor* cf. *nodicostatus* Kilburn, 1988, MNHN, Nw-Caledonia, MUSORSTOM 6, st. DW391, 20°47'S, 167°06'E, 390m.
- C.** Clavatulidae. *Clavatula strebeli* Knudsen, 1952, MNHN, Ilha de Luanda, Angola, 40-60m.
- D.** Horaiclavidae. *Ceritoturris pupiformis* (Smith, 1884)*, MNHN, New-Caledonia, MONTROUZIER, st.1249, 20°49'S, 165°19'E, 80-140m.
- E.** Turridae. *Gemmula* sp.**, MNHN IM200740766, Coral Sea, EBISCO, DW2625, 20°05'S, 160°20'E, 627-741m.
- F.** Borsoniidae. *Microdrillia fastosa* (Hedley, 1907), MNHN, New-Caledonia, SMIB 8, st. DW146, 24°55'S, 168°22'E, 514-522m.
- G, H.** Clathurellidae.
- G.** *Clathurella curtisiana* (Hedley, 1922), MNHN, New-Caledonia, MONTROUZIER, st.1250, 20°46.7'S, 165°13.7'E, 3-6m.
- H.** *Nannodiella acricula* (Hedley, 1922), MNHN, New-Caledonia, MONTROUZIER, st.1261, 20°46'S, 165°15'E, 45-56m.
- I.** Mitromorphidae. *Mitromorpha metula*** (Hinds, 1843), MNHN IM200717898, Philippines, PANGLAO 2004, st. B8, 9°37.1'N, 123°46.1'E, 3m.
- J-L.** Mangeliidae.
- J.** *Benthomangelia* sp., MNHN, Fidji, BORDAU 1, st. CP1407, 16°40'S, 179°39'E, 499-529m.
- K.** *Neoguraleus* sp., MNHN, New Caledonia, CONCALIS, st. DW2982, 18°02'S, 163°04'E, 320-337m.
- L.** *Pyrgocythara notabilis* (Smith, 1888), MNHN, New Caledonia, LIFOU 2000, st. 1429, 20°47.5'S, 167°07.1'E, 8-18m.
- M-Q.** Raphitomidae.
- M.** *Neopleurotomoides* sp., MNHN, New- Caledonia, BIOCAL, st. CP75, 22°19'S, 167°23'E.
- N.** *Famelica* sp., MNHN, New- Caledonia, BATHUS 4, st. DW914, 18°49'S, 163°15'E, 600-616m.
- O.** *Tritonoturris cumingi* (Powys, 1835), MNHN, Touho, New Caledonia, st. 1357, 22°30.2'S, 166°26.4'E, 22-35m.

P. *Veprecula* sp., MNHN, New Caledonia, MONTROUZIER, st. 1306, 20°39.1'S, 164°12.4'E, 11-13m.

Q. *Pleurotomella packardi* Verrill, 1872, MNHN, INCAL, Porcupine abyssal plain, st. WS10, 47°27'N, 09°40'W, 4354m.

Symbol: **, sequenced specimen.

Appendix 1: Index of genera and subgenera. Genus-group taxa removed from Conoidea are preceded by an asterisk.

<i>Abretiella</i>	Terebridae	<i>Belalora</i>	?Mangeliidae
<i>Abyssaforia</i>	?Cochlespiridae	<i>Belaterricula</i>	?Borsoniidae
<i>Abyssobelaa</i>	Raphitomidae	<i>Bellacythara</i>	Mangeliidae
<i>Abyssocomitas</i>	?Pseudomelatomidae	<i>Bellaspira</i>	Drilliidae
<i>Abyssothauma</i>	?Raphitomidae	<i>Bellatula</i>	Clathurellidae
<i>Acamptodaphne</i>	Raphitomidae	* <i>Belomitria</i>	Buccinidae
<i>Acanthodaphne</i>	Raphitomidae	<i>Benthoclionella</i>	Clavatulidae
<i>Acinodrillia</i>	?Drilliidae	<i>Benthodaphne</i>	?Pseudomelatomidae
<i>Acmaturris</i>	Mangeliidae	<i>Benthofascis</i>	Conorbidae
<i>Acrista</i>	Clathurellidae	<i>Benthomangelia</i>	Mangeliidae
<i>Acropota</i>	Borsoniidae	<i>Boettgeriola</i>	Borsoniidae
<i>Acuminia</i>	Terebridae	<i>Borsonella</i>	Borsoniidae
<i>Adanaclava</i>	?Clathurellidae	<i>Borsonellopsis</i>	Borsoniidae
<i>Aforia</i>	?Cochlespiridae	<i>Borsonia</i>	Borsoniidae
<i>Africonus</i>	Conidae	<i>Brachycythara</i>	Mangeliidae
<i>Agathotoma</i>	Mangeliidae	<i>Brachytoma</i>	?Pseudomelatomidae
<i>Agladrillia</i>	Drilliidae	<i>Brephodrillia</i>	Drilliidae
<i>Aguilaria</i>	Pseudomelatomidae	<i>Brevimyurella</i>	Terebridae
<i>Aliceia</i>	Raphitomidae	<i>Buccinaria</i>	Raphitomidae
<i>Allo</i>	?Raphitomidae	<i>Buchema</i>	Horaiclavidae
<i>Alticlavatula</i>	Horaiclavidae	<i>Burchia</i>	Pseudomelatomidae
<i>Anacithara</i>	Horaiclavidae	<i>Buridrillia</i>	Pseudomelatomidae
<i>Anarithma</i>	Mitromorphidae	<i>Cacodaphnella</i>	?Mangeliidae
<i>Ancistrosyrinx</i>	Cochlespiridae	<i>Calamiconus</i>	Conidae
<i>Anguloclavus</i>	Horaiclavidae	<i>Calcatodrillia</i>	Pseudomelatomidae
<i>Annulaturris</i>	Turridae	<i>Calibanus</i>	Conidae
<i>Anomalotomella</i>	Raphitomidae	<i>Caliedrula</i>	Clavatulidae
<i>Anticlinura</i>	Mangeliidae	<i>Californiconus</i>	Conidae
<i>Anticomitas</i>	?Pseudomelatomidae	<i>Calliclava</i>	Drilliidae
<i>Antiguraleus</i>	Mangeliidae	<i>Calusaconus</i>	Conidae
<i>Antimelatoma</i>	Pseudomelatomidae	<i>Canetoma</i>	Mangeliidae
* <i>Antimitra</i>	Buccinidae	<i>Caribocoonus</i>	Conidae
<i>Antiplanes</i>	Pseudomelatomidae	<i>Carinapex</i>	Horaiclavidae
<i>Aoteadrillia</i>	?Horaiclavidae	<i>Carinodrillia</i>	Pseudomelatomidae
<i>Apaturris</i>	?Borsoniidae	<i>Carinoturris</i>	?Pseudomelatomidae
<i>Aphanitoma</i>	Borsoniidae	<i>Cenodagreutes</i>	Raphitomidae
<i>Apiotoma</i>	?Cochlespiridae	<i>Ceritoturris</i>	Horaiclavidae
<i>Apispiralia</i>	Mangeliidae	<i>Cerodrillia</i>	Drilliidae
<i>Apitua</i>	Mangeliidae	<i>Cestoma</i>	Mangeliidae
<i>Arielia</i>	Mitromorphidae	<i>Chelyconus</i>	Conidae
<i>Artemidiconus</i>	Conorbidae	<i>Cheungbeia</i>	Pseudomelatomidae
<i>Asperdaphne</i>	Raphitomidae	<i>Cinguloterebra</i>	Terebridae
<i>Aspertilla</i>	Raphitomidae	<i>Citharomangelia</i>	Mangeliidae
<i>Asprella</i>	Conidae	<i>Clamturris</i>	Turridae
<i>Asthenotoma</i>	Borsoniidae	<i>Clathrodrillia</i>	Drilliidae
<i>Astrocarina</i>	?Horaiclavidae	<i>Clathromangelia</i>	?Clathurellidae
<i>Astroconus</i>	Conidae	<i>Clathroterebra</i>	Terebridae
<i>Austrodaphnella</i>	Raphitomidae	<i>Clathurella</i>	Clathurellidae
<i>Austrodrillia</i>	Horaiclavidae	<i>Clathurina</i>	Raphitomidae
<i>Austrogemmula</i>	Turridae	<i>Clavatula</i>	Clavatulidae
<i>Austropusilla</i>	?Raphitomidae	<i>Clavicantha</i>	Drilliidae
<i>Austrotoma</i>	Raphitomidae	<i>Clavosurcula</i>	?Cochlespiridae
<i>Austroturris</i>	Borsoniidae	<i>Clavus</i>	Drilliidae
<i>Azorilla</i>	Raphitomidae	<i>Clebula</i>	Conidae
<i>Azorita</i>	Raphitomidae	<i>Cleospira</i>	Strictispiridae
<i>Bactrocycla</i>	Mangeliidae	<i>Clinura</i>	Raphitomidae
<i>Bathybela</i>	Raphitomidae	<i>Clinuromella</i>	Mangeliidae
<i>Bathybermudia</i>	Turridae	<i>Clionella</i>	Clavatulidae
* <i>Bathyclionella</i>	Buccinidae	<i>Cochlespira</i>	Cochlespiridae
<i>Bathyconus</i>	Conidae	<i>Cochlioconus</i>	Raphitomidae
<i>Bathypota</i>	Raphitomidae	<i>Comarmondia</i>	Clathurellidae
<i>Bathytoma</i>	Borsoniidae	<i>Comitas</i>	Pseudomelatomidae
<i>Bela</i>	Mangeliidae	<i>Compsodrillia</i>	Pseudomelatomidae

<i>Conasprella</i>	Conidae	<i>Epidirella</i>	Turridae
<i>Conasprelloides</i>	Conidae	<i>Epidirona</i>	Pseudomelatomidae
<i>Conopleura</i>	Drilliidae	<i>Eremiconus</i>	Conidae
<i>Conorbela</i>	?Pseudomelatomidae	<i>Erythroconus</i>	Conidae
<i>Conticosta</i>	Pseudomelatomidae	<i>Etrema</i>	Clathurellidae
<i>Conus</i>	Conidae	<i>Etremopa</i>	Clathurellidae
<i>Cordieria</i>	Borsoniidae	<i>Etremopsis</i>	Clathurellidae
<i>Corinnaeturris</i>	Clathurellidae	<i>Eubela</i>	Raphitomidae
<i>Cornutoconus</i>	Conidae	<i>Eucithara</i>	Mangeliidae
<i>Coronacomitas</i>	Horaiclavidae	<i>Euclathurella</i>	Clathurellidae
<i>Coronasyrinx</i>	Cochlespiridae	<i>Eucyclotoma</i>	Raphitomidae
<i>Coronaxis</i>	Conidae	<i>Eudaphnella</i>	Raphitomidae
<i>Crassiclava</i>	Pseudomelatomidae	<i>Eugemmula</i>	Turridae
<i>Crassispira</i>	Pseudomelatomidae	<i>Eugeniconus</i>	Conidae
<i>Crassispirella</i>	Pseudomelatomidae	<i>Euglyphostoma</i>	Clathurellidae
<i>Crassopleura</i>	Drilliidae	<i>Euguraleus</i>	Mangeliidae
<i>Cretaspira</i>	Pseudomelatomidae	<i>Eumetadrillia</i>	Drilliidae
<i>Crockerella</i>	Clathurellidae	<i>Euryentemema</i>	?Mangeliidae
<i>Cruziturricula</i>	?Drilliidae	<i>Euryta</i>	Terebridae
<i>Cryoturris</i>	Mangeliidae	<i>Euterebra</i>	Terebridae
<i>Cryptodaphne</i>	Raphitomidae	<i>Exomilus</i>	?Raphitomidae
<i>Cryptogemma</i>	Turridae	<i>Famelica</i>	Raphitomidae
* <i>Cryptomitra</i>	Buccinidae	<i>Favriella</i>	Raphitomidae
<i>Cucullites</i>	Conidae	<i>Fehria</i>	Mangeliidae
<i>Cucullus</i>	Conidae	<i>Feliciella</i>	?Raphitomidae
<i>Curtitoma</i>	Mangeliidae	<i>Fenestrosyrinx</i>	Raphitomidae
<i>Cylinder</i>	Conidae	<i>Fenimorea</i>	Drilliidae
<i>Cylindrella</i>	Conidae	<i>Filodrillia</i>	Borsoniidae
<i>Cylindrus</i>	Conidae	<i>Floraconus</i>	Conidae
<i>Cymakra</i>	Mitromorphidae	<i>Fulgiconus</i>	Conidae
<i>Cymatosyrinx</i>	Drilliidae	<i>Fumiconus</i>	Conidae
<i>Cyrtocythara</i>	Mangeliidae	<i>Funa</i>	Pseudomelatomidae
<i>Cyrtoides</i>	Raphitomidae	<i>Funitoma</i>	Mangeliidae
<i>Cythara</i>	?Mangeliidae	<i>Fusiconus</i>	Conidae
<i>Cytharella</i>	Mangeliidae	<i>Fusidaphne</i>	?Raphitomidae
<i>Cytharoclavus</i>	Horaiclavidae	<i>Fusisyrinx</i>	?Drilliidae
<i>Cytharopsis</i>	Mangeliidae	<i>Fusiturricula</i>	?Drilliidae
<i>Dallaforia</i>	?Cochlespiridae	<i>Fusiturris</i>	?Clavatulidae
<i>Dalliconus</i>	Conidae	<i>Gastridium</i>	Conidae
<i>Dallspira</i>	Pseudomelatomidae	<i>Gemmula</i>	Turridae
<i>Danilacarina</i>	Cochlespiridae	<i>Gemmuloborsonia</i>	?Clavatulidae
<i>Daphnella</i>	Raphitomidae	<i>Genota</i>	Borsoniidae
* <i>Daphnellopsis</i>	Muricidae	<i>Genotia</i>	Borsoniidae
<i>Darbya</i>	Borsoniidae	<i>Genotina</i>	Mangeliidae
<i>Darioconus</i>	Conidae	<i>Genuanoconus</i>	Conidae
<i>Darrylia</i>	?Horaiclavidae	<i>Gibbaspira</i>	Pseudomelatomidae
<i>Dauciconus</i>	Conidae	<i>Gingicithara</i>	Mangeliidae
<i>Decollidrilla</i>	Turridae	<i>Ginnania</i>	Mangeliidae
<i>Decorihastula</i>	Terebridae	<i>Gladioconus</i>	Conidae
<i>Dendroconus</i>	Conidae	<i>Globiconus</i>	Conidae
<i>Diaugasma</i>	Raphitomidae	<i>Globidrillia</i>	Drilliidae
<i>Dimidacus</i>	Terebridae	<i>Glossispira</i>	Pseudomelatomidae
<i>Diplomeriza</i>	Terebridae	<i>Glyphostoma</i>	Clathurellidae
<i>Diptychophlia</i>	Borsoniidae	<i>Glyphostomoides</i>	Raphitomidae
<i>Douglassia</i>	Drilliidae	<i>Glyphostomopsis</i>	Clathurellidae
<i>Doxospira</i>	Pseudomelatomidae	<i>Glyphoturris</i>	Mangeliidae
<i>Drillia</i>	Drilliidae	<i>Glyptaesopus</i>	?Borsoniidae
<i>Drilliola</i>	Borsoniidae	<i>Graciliclava</i>	Horaiclavidae
<i>Ductoconus</i>	Conidae	<i>Gradaterebra</i>	Terebridae
<i>Duplicaria</i>	Terebridae	<i>Gradiconus</i>	Conidae
<i>Dyraspis</i>	Conidae	<i>Granotoma</i>	Mangeliidae
<i>Egentelaria</i>	Terebridae	<i>Granoturris</i>	Mangeliidae
<i>Elaeocyma</i>	Drilliidae	<i>Granuliterebra</i>	Terebridae
<i>Eldridgea</i>	Drilliidae	<i>Graphiconus</i>	Conidae
<i>Embrikena</i>	Conidae	<i>Guraleus</i>	Mangeliidae
<i>Endemoconus</i>	Conidae	<i>Gymnobela</i>	Raphitomidae
<i>Epideira</i>	Pseudomelatomidae	<i>Haedropleura</i>	Horaiclavidae

<i>Harmoniconus</i>	Conidae	<i>Lissodrillia</i>	Drilliidae
<i>Hastula</i>	Terebridae	<i>Litachilus</i>	Raphitomidae
<i>Hastulina</i>	Terebridae	<i>Lithoconus</i>	Conidae
<i>Hastulopsis</i>	Terebridae	<i>Lividoconus</i>	Conidae
<i>Hauturua</i>	Drilliidae	<i>Lizaconus</i>	Conidae
<i>Helenella</i>	Mitromorphidae	<i>Lophiotoma</i>	Turridae
<i>Hemicythara</i>	Mangeliidae	<i>Lophioturris</i>	Turridae
<i>Hemidaphne</i>	Raphitomidae	<i>Lorabela</i>	?Mangeliidae
<i>Hemilenardia</i>	Raphitomidae	<i>Lovellona</i>	Mitromorphidae
<i>Hermes</i>	Conidae	<i>Lucerapex</i>	?Turridae
<i>Heroconus</i>	Conidae	<i>Lusitanops</i>	Raphitomidae
<i>Heterocithara</i>	Mangeliidae	<i>Lyromangelia</i>	Mangeliidae
<i>Heteroturris</i>	Borsoniidae	<i>Macteola</i>	?Mangeliidae
<i>Hindsclava</i>	Pseudomelatomidae	<i>Maesiella</i>	Pseudomelatomidae
<i>Horaiclavus</i>	Horaiclavidae	<i>Magelliconus</i>	Conidae
<i>Hormospira</i>	Pseudomelatomidae	<i>Magnella</i>	Raphitomidae
<i>Imaclava</i>	Drilliidae	<i>Majox</i>	Raphitomidae
<i>Impages</i>	Terebridae	<i>Makiyamaia</i>	Clavatulidae
<i>Inkinga</i>	Horaiclavidae	<i>Mamiconus</i>	Conidae
<i>Inodrillara</i>	Horaiclavidae	<i>Mammillaedrillia</i>	Pseudomelatomidae
<i>Inodrillia</i>	Horaiclavidae	<i>Mangelia</i>	Mangeliidae
<i>Inodrillina</i>	Horaiclavidae	<i>Mangiliella</i>	Mangeliidae
<i>Inquisitor</i>	Pseudomelatomidae	<i>Maorimorpha</i>	Mitromorphidae
<i>Iotyrris</i>	Turridae	<i>Maoritomella</i>	Borsoniidae
<i>Iraquetrema</i>	Clathurellidae	<i>Marita</i>	Mangeliidae
<i>Iredalea</i>	Drilliidae	<i>Marshallena</i>	?Horaiclavidae
<i>Irenosyrinx</i>	Cochlespiridae	<i>Mauidrillia</i>	Horaiclavidae
<i>Isodaphne</i>	Raphitomidae	<i>Mazatlania</i>	Terebridae
<i>Ithycythara</i>	Mangeliidae	<i>Megasurcula</i>	Pseudomelatomidae
<i>Itia</i>	?Mitromorphidae	<i>Meggittia</i>	?Pseudomelatomidae
<i>Iwaoa</i>	Horaiclavidae	<i>Metaclathurella</i>	Raphitomidae
<i>Jaspidiconus</i>	Conidae	<i>Micantapex</i>	Borsoniidae
* <i>Jumala</i>	Buccinidae	<i>Microdaphne</i>	Raphitomidae
<i>Kalloconus</i>	Conidae	<i>Microdrillia</i>	Borsoniidae
<i>Kenyonia</i>	Conidae	<i>Microgenia</i>	Raphitomidae
<i>Kermasperella</i>	Conidae	<i>Micropleurotoma</i>	?Horaiclavidae
<i>Kermia</i>	Raphitomidae	<i>Microtrypetes</i>	Terebridae
<i>Ketyconus</i>	Conidae	<i>Miliariconus</i>	Conidae
<i>Kioconus</i>	Conidae	<i>Mioawateria</i>	Raphitomidae
<i>Knefastia</i>	Pseudomelatomidae	<i>Miraclathurella</i>	Pseudomelatomidae
<i>Kohniconus</i>	Conidae	<i>Mitraguraleus</i>	Mangeliidae
<i>Kuradoconus</i>	Conidae	<i>Mitrellatoma</i>	Mitromorphidae
<i>Kurilohadalia</i>	?Pseudomelatomidae	<i>Mitrithara</i>	Mitromorphidae
<i>Kurodadrillia</i>	Pseudomelatomidae	<i>Mitrolumna</i>	Mitromorphidae
<i>Kuroshiodaphne</i>	Raphitomidae	<i>Mitromorpha</i>	Mitromorphidae
<i>Kuroshioturris</i>	Turridae	<i>Monilispira</i>	Pseudomelatomidae
<i>Kurtzia</i>	Mangeliidae	<i>Monteiroconus</i>	Conidae
<i>Kurtziella</i>	Mangeliidae	<i>Mordica</i>	Raphitomidae
<i>Kurtzina</i>	Mangeliidae	<i>Myurella</i>	Terebridae
<i>Kylix</i>	Drilliidae	<i>Myurellina</i>	Terebridae
<i>Laeviacus</i>	Terebridae	<i>Myurellisca</i>	Terebridae
<i>Lamniconus</i>	Conidae	<i>Nannodiella</i>	Clathurellidae
<i>Lautoconus</i>	Conidae	<i>Narraweena</i>	Borsoniidae
<i>Leiocithara</i>	Mangeliidae	<i>Naskia</i>	Horaiclavidae
<i>Leiosyrinx</i>	Raphitomidae	<i>Nataliconus</i>	Conidae
<i>Leporiconus</i>	Conidae	<i>Naude drillia</i>	Pseudomelatomidae
<i>Leptadrillia</i>	Drilliidae	<i>Nematoma</i>	Mangeliidae
<i>Leptoconus</i>	Conidae	<i>Neodrillia</i>	Drilliidae
<i>Leucosyrinx</i>	?Pseudomelatomidae	<i>Neoguraleus</i>	?Mangeliidae
<i>Leufroyia</i>	Raphitomidae	<i>Neopleurotomoides</i>	Raphitomidae
<i>Lienardia</i>	Clathurellidae	<i>Nepotilla</i>	Raphitomidae
<i>Lilliconus</i>	Conidae	<i>Netrum</i>	Clavatulidae
<i>Lindaconus</i>	Conidae	<i>Nihonia</i>	?Cochlespiridae
<i>Lineotoma</i>	Raphitomidae	<i>Nodotoma</i>	Mangeliidae
<i>Lioglyphostoma</i>	Pseudomelatomidae	<i>Notocytharella</i>	?Mangeliidae
<i>Lioglyphostomella</i>	Pseudomelatomidae	<i>Nototerebra</i>	Terebridae
<i>Liracraea</i>	Mangeliidae	<i>Nquma</i>	Horaiclavidae

<i>Nymphispira</i>	Pseudomelatomidae	<i>Ptychosyrinx</i>	Turridae
<i>Obesotoma</i>	Mangeliidae	<i>Pulsarella</i>	Borsoniidae
<i>Oenopota</i>	Mangeliidae	<i>Puncticulis</i>	Conidae
<i>Oenopotella</i>	Mangeliidae	<i>Punctoterebra</i>	Terebridae
<i>Ongoconus</i>	Conidae	<i>Purpuriconus</i>	Conidae
<i>Ootiomella</i>	Raphitomidae	<i>Pusionella</i>	Clavatulidae
<i>Ophiodermella</i>	Borsoniidae	<i>Pyrgocythara</i>	Mangeliidae
<i>Orrmaesia</i>	Drilliidae	<i>Pyrgospira</i>	Pseudomelatomidae
<i>Otitoma</i>	Pseudomelatomidae	<i>Pyruconus</i>	Conidae
<i>Oxymeris</i>	Terebridae	<i>Qii</i>	Raphitomidae
<i>Oxytropa</i>	Turridae	<i>Quasiconus</i>	Conidae
<i>Pagodasyrinx</i>	Cochlespiridae	<i>Raphitoma</i>	Raphitomidae
<i>Pagodidaphne</i>	Raphitomidae	<i>Rectiplanes</i>	Pseudomelatomidae
<i>Panaterebra</i>	Terebridae	<i>Rectisulcus</i>	Pseudomelatomidae
<i>Papilocithara</i>	Mangeliidae	<i>Regiconus</i>	Conidae
<i>Parabathyтома</i>	Borsoniidae	<i>Regidrillia</i>	Horaclavidae
<i>Paraborsonia</i>	Borsoniidae	<i>Retidrillia</i>	Borsoniidae
<i>Paraclathurella</i>	Clathurellidae	<i>Rhizoconus</i>	Conidae
<i>Paracomitas</i>	?Pseudomelatomidae	<i>Rhodopetoma</i>	?Pseudomelatomidae
<i>Paracuneus</i>	?Drilliidae	<i>Rhombiconus</i>	Conidae
<i>Paradaphne</i>	Raphitomidae	<i>Rhombus</i>	Conidae
<i>Paradrillia</i>	Horaclavidae	<i>Rimosodaphnella</i>	Raphitomidae
<i>Paramontana</i>	Raphitomidae	<i>Rissomangelia</i>	Mangeliidae
<i>Parasprioptropis</i>	?Mangeliidae	<i>Riuguhdillia</i>	Borsoniidae
<i>Paraterebra</i>	Terebridae	<i>Rocroithys</i>	Raphitomidae
<i>Partecosta</i>	Terebridae	<i>Rolaniconus</i>	Conidae
<i>Parviconus</i>	Conidae	<i>Rollus</i>	Conidae
<i>Pellifronia</i>	Terebridae	<i>Rubellatoma</i>	Mangeliidae
<i>Peraotoma</i>	Raphitomidae	<i>Rugocythara</i>	Mangeliidae
<i>Perimangelia</i>	Mangeliidae	<i>Saccharoturris</i>	Mangeliidae
<i>Perirhoe</i>	Terebridae	<i>Scaeavatula</i>	Clavatulidae
<i>Perplexiconus</i>	Conidae	<i>Sciteconus</i>	Conidae
<i>Perrona</i>	Clavatulidae	<i>Scrinium</i>	Mitromorphidae
<i>Pervicacia</i>	Terebridae	<i>Sediliopsis</i>	Pseudomelatomidae
<i>Phasmocoonus</i>	Conidae	<i>Seminoleconus</i>	Conidae
<i>Phenatoma</i>	Borsoniidae	<i>Shutonia</i>	?Pseudomelatomidae
<i>Philbertia</i>	Raphitomidae	<i>Sibogasyrinx</i>	Cochlespiridae
<i>Phymorhynchus</i>	Raphitomidae	<i>Smithiella</i>	Mangeliidae
<i>Pilsbryspira</i>	Pseudomelatomidae	<i>Socioconus</i>	Conidae
<i>Pinguigemmula</i>	Turridae	<i>Speoides</i>	Raphitomidae
<i>Pionoconus</i>	Conidae	<i>Spergo</i>	Raphitomidae
<i>Pionotoma</i>	Raphitomidae	<i>Spirotropis</i>	Drilliidae
<i>Plagiostropha</i>	Drilliidae	<i>Splendrilla</i>	Drilliidae
<i>Platycythara</i>	Mangeliidae	<i>Spinocoonus</i>	Conidae
<i>Pleurotoma</i>	Turridae	<i>Spuriconus</i>	Conidae
<i>Pleurotomella</i>	Raphitomidae	<i>Steiraxis</i>	?Cochlespiridae
<i>Pleurotomus</i>	Turridae	* <i>Steironepion</i>	Columbellidae
<i>Plicaustraconus</i>	Conidae	<i>Stellaconus</i>	Conidae
<i>Plicisyrix</i>	Pseudomelatomidae	<i>Stellatoma</i>	Mangeliidae
<i>Polystira</i>	Turridae	<i>Stenodrillia</i>	?Drilliidae
<i>Pontiothauma</i>	Raphitomidae	<i>Stephanoconus</i>	Conidae
<i>Pristiterebra</i>	Terebridae	<i>Stilla</i>	Raphitomidae
<i>Profundiconus</i>	Conidae	<i>Strategoconus</i>	Conidae
<i>Propebela</i>	Mangeliidae	<i>Striatoguraleus</i>	Horaclavidae
<i>Protoconus</i>	Conidae	<i>Strictispira</i>	Strictispiridae
<i>Protostrioconus</i>	Conidae	<i>Strioconus</i>	Conidae
<i>Pseudexomilus</i>	?Horaclavidae	<i>Striospira</i>	Pseudomelatomidae
<i>Pseudoconorbis</i>	Conidae	<i>Strioterebrum</i>	Terebridae
<i>Pseudodaphnella</i>	Raphitomidae	<i>Strombinoturris</i>	Clathurellidae
<i>Pseudoetrema</i>	Clathurellidae	<i>Suavodrillia</i>	Borsoniidae
<i>Pseudolilliconus</i>	Conidae	<i>Subula</i>	Terebridae
<i>Pseudomelatoma</i>	Pseudomelatomidae	<i>Sugitanitoma</i>	Horaclavidae
<i>Pseudonodulocoonus</i>	Conidae	<i>Surcula</i>	Clavatulidae
<i>Pseudorhaphitoma</i>	Mangeliidae	* <i>Surculina</i>	Turbinellidae
<i>Pseudotaranis</i>	?Pseudomelatomidae	<i>Suturocythara</i>	Mangeliidae
<i>Psittacodrillia</i>	Horaclavidae	<i>Syntomodrillia</i>	Drilliidae
<i>Ptychobela</i>	Pseudomelatomidae	<i>Taranidaphne</i>	Raphitomidae

<i>Taranis</i>	Raphitomidae	* <i>Turrijaumelia</i>	Columbellidae
<i>Taranteconus</i>	Conidae	<i>Turris</i>	Turridae
<i>Tasmadaphne</i>	Raphitomidae	<i>Turritomella</i>	Mangeliidae
<i>Teleochilus</i>	?Raphitomidae	<i>Tuskaroria</i>	Raphitomidae
<i>Tenaturris</i>	Mangeliidae	<i>Tylotia</i>	Drilliidae
<i>Terebra</i>	Terebridae	<i>Tylotiella</i>	Drilliidae
<i>Terebrum</i>	Terebridae	<i>Typhlodaphne</i>	?Borsoniidae
<i>Terenolla</i>	Terebridae	<i>Typhlomangelia</i>	Borsoniidae
<i>Teretia</i>	Raphitomidae	<i>Typhlosyrinx</i>	Raphitomidae
<i>Teretiopsis</i>	Raphitomidae	<i>Tyrrhenoturris</i>	Clavatulidae
<i>Tesseliconus</i>	Conidae	<i>Unedogemmula</i>	Turridae
<i>Textilia</i>	Conidae	<i>Utriculus</i>	Conidae
<i>Thatcheria</i>	Raphitomidae	<i>Varioconus</i>	Conidae
<i>Thatcheriasyrinx</i>	?Raphitomidae	<i>Venustoma</i>	?Mangeliidae
<i>Thatcherina</i>	Raphitomidae	<i>Veprecula</i>	Raphitomidae
<i>Thelecythara</i>	Horaiclavidae	<i>Vepridaphne</i>	Raphitomidae
<i>Thelecytharella</i>	Pseudomelatomidae	<i>Vexiariela</i>	Mitromorphidae
<i>Theliconus</i>	Conidae	<i>Vexithara</i>	Borsoniidae
<i>Thesbia</i>	?Raphitomidae	<i>Vexitomina</i>	Horaiclavidae
<i>Theta</i>	Raphitomidae	<i>Villiersiella</i>	Mangeliidae
<i>Thetidos</i>	Clathurellidae	<i>Viminiconus</i>	Conidae
<i>Thoraconus</i>	Conidae	<i>Virgiconus</i>	Conidae
<i>Tiariturris</i>	Pseudomelatomidae	<i>Viridoturris</i>	Borsoniidae
<i>Tomellana</i>	Clavatulidae	<i>Viridrillia</i>	Pseudomelatomidae
<i>Tomopleura</i>	Borsoniidae	<i>Viridrillina</i>	Pseudomelatomidae
<i>Toxiclionella</i>	Clavatulidae	<i>Virroconus</i>	Conidae
<i>Toxicochlespira</i>	Mangeliidae	<i>Vitjazinella</i>	Mangeliidae
<i>Trachydrillida</i>	Clavatulidae	<i>Vitricythara</i>	Mangeliidae
<i>Triplostephanus</i>	Terebridae	<i>Vituliconus</i>	Conidae
<i>Tritonoturris</i>	Raphitomidae	<i>Watsonaria</i>	Raphitomidae
<i>Tropidoturris</i>	Borsoniidae	<i>Widalli</i>	Mangeliidae
<i>Trovaconus</i>	Conidae	<i>Xanthodaphne</i>	Raphitomidae
<i>Truncadaphne</i>	Raphitomidae	<i>Xenuroturris</i>	Turridae
<i>Tuliparia</i>	Conidae	<i>Ximeniconus</i>	Conidae
<i>Turrella</i>	Clathurellidae	<i>Yeddoconus</i>	Conidae
<i>Turhyssa</i>	Raphitomidae	<i>Zemacies</i>	Borsoniidae
<i>Turriconus</i>	Conidae	<i>Zenepos</i>	Raphitomidae
<i>Turricula</i>	Clavatulidae	* <i>Zetekia</i>	Columbellidae
<i>Turridrupa</i>	Turridae	<i>Zonulispira</i>	Pseudomelatomidae
<i>Turrigemma</i>	Pseudomelatomidae		

Appendix 2: List of all the family and subfamily names available for the Conoidea (see Bouchet and Rocroi, 2005 for details). The status of names including only fossil genera is not discussed.

Acidae Gray, 1853 (nomenclaturally invalid)
Andoniinae Vera-Pelaez, 2002 (fossil)
Belinae Bellardi, 1875 (synonym of Mangeliidae)
Borsoniidae Bellardi, 1875
Brachytominae Thiele, 1929 (nomen dubium)
Californiconinae Tucker & Tenorio, 2009 (synonym of Conidae)
Clathurellidae H. Adams & A. Adams, 1858
Clavatulidae Gray, 1853
Clavinae Casey 1904 (nomenclaturally invalid)
Clionellidae Stimpson, 1865 (synonym of Clavatulidae)
Cochlespiridae Powell, 1942
Conidae Flemming, 1922
Conilithidae Tucker & Tenorio, 2009 (synonym of Conidae)
Conorbidae de Gregorio 1880
Conulinae Rafinesque, 1815 (nomenclaturally invalid)
Crassispirinae McLean, 1971 (synonym of Pseudomelatomidae)
Cryptoconinae Cossmann, 1896 (fossil)
Cytharinae Thiele, 1929 (synonym of Mangeliidae)
Daphnellinae Casey, 1904 (synonym of Raphitomidae)
Defranciinae Gray, 1853 (nomenclaturally invalid)
Diptychomitrinae Bellardi, 1888 (synonym of Mitromorphidae)
Drilliidae Olson, 1964
Hemiconidae Tucker & Tenorio, 2009 (fossil)
Horaiclavidae fam. nov.
Johnwyattidae Serna, 1979 (fossil)
Lorinae Thiele, 1925 *sensu* opinion 666 (synonym of Clathurellidae)
Lorinae Thiele, 1925 *sensu* Thiele (synonym of Mangeliidae)
Mangeliidae Fisher, 1883
Melatomidae Gill, 1871 (synonym of Clavatulidae)
Mitrolumnidae Sacco, 1904 (synonym of Mitromorphidae)
Mitromorphidae Casey, 1904
Oenopotinae Bogdanov, 1987 (synonym of Mangeliidae)
Pervicaciinae Rudman, 1969 (synonym of Terebridae)
Pleurotominae Gray, 1838 (synonym of Turridae)
Pleurotomellinae Nordsieck, 1968 (synonym of Raphitomidae)
Pseudomelatomidae Morrison, 1965
Pseudotominae Bellardi, 1875 (synonym of Clathurellidae)
Puncticuliinae Tucker & Tenorio, 2009 (synonym of Conidae)
Pusionellinae Gray, 1853 (synonym of Clavatulidae)
Raphitomidae Bellardi, 1875
Siphopsinae Le Renard, 1995 (fossil)
Speightidae Powell, 1942 (fossil; doubtfully Conoidea)
Strictispiridae McLean, 1971
Taraninae Casey, 1904 (synonym of Raphitomidae)
Taranteconidae Tucker & Tenorio, 2009 (synonym of Conidae)
Terebridae Mörch, 1852

Thatcheriidae Powell, 1942 (synonym of Raphitomidae)
Turriculinae Powell, 1942 (synonym of Clavatulidae)
Turridae H. Adams & A. Adams, 1853
Zemaciinae Sysoev, 2003 (synonym of Borsoniidae)
Zonulispirinae McLean 1971 (synonym of Pseudomelatomidae)

ATaylor, et al., 1993 /
Bouchet & Rocroi 2005**B**

This paper

C

Tucker & Tenorio 2009

CONOIDEACLAVATULIDAE¹

DRILLIIDAE

PSEUDOMELATOMIDAE

STRICTISPIRIDAE

TEREBRIDAE

TURRIDAE

Crassispirinae

Zonulispirinae

Cochlespirinae

Turrinae

Zemaciinae²

CONIDAE

Clathurellinae

Coninae

Conorbinae

Raphitominae

Mangeliinae

Oenopotinae

CONOIDEA

CLAVATULIDAE

DRILLIIDAE

PSEUDOMELATOMIDAE

HORAICLAVIDAE

STRICTISPIRIDAE

TEREBRIDAE

TURRIDAE

COCHLESPIRIDAE

BORSONIIDAE

MITROMORPHIDAE

CLATHURELLIDAE

CONIDAE

CONORBIDAE

RAPHITOMIDAE

MANGELIIDAE

TURROIDEA

CLAVATULIDAE

DRILLIIDAE

PSEUDOMELATOMIDAE

STRICTISPIRIDAE

TEREBRIDAE

TURRIDAE

Crassispirinae

Zonulispirinae

Cochlespirinae

Turrinae

Zemaciinae²**CONOIDEA**

BORSONIIDAE

CLATHURELLIDAE

Mitromorphinae

Clathurellinae

†CRYPTOCONIDAE

CONIDAE

Coninae

Puncticuliinae

CONILITHIDAE

Conilithinae

Californiconinae

†HEMICONIDAE


TARANTECONIDAE

CONORBIDAE

RAPHITOMIDAE





Raphitominae


Mangeliinae






Borsoniidae

Conorbidae











A










B









C





Mangeliidae


















































Clavatulidae









Horaiclavidae










Turridae





Strictispiridae



Conorbiidae

