



HAL
open science

Evolution of nutrient acquisition: when space matters

Sébastien Barot, Stefan Bornhofen, Simon Boudsocq, Xavier Raynaud,
Nicolas Loeuille

► **To cite this version:**

Sébastien Barot, Stefan Bornhofen, Simon Boudsocq, Xavier Raynaud, Nicolas Loeuille. Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 2016, 30 (2), pp.283-294. 10.1111/1365-2435.12494 . hal-02458157

HAL Id: hal-02458157

<https://hal.science/hal-02458157v1>

Submitted on 28 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evolution of nutrient acquisition: when space matters

Journal:	<i>Functional Ecology</i>
Manuscript ID:	FE-2015-00010.R1
Manuscript Type:	Standard Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Barot, Sébastien; IRD, IEES-P Bornhofen, Stefan; EISTI, Boudsocq, Simon; INRA, Eco & Sols Raynaud, Xavier; UPMC, IEES-P Loeuille, Nicolas; UPMC, IEES-P
Key-words:	nutrient acquisition, nutrient conservation, spatial model, local vs. regional competition, evolution of altruism, plant traits, eco-evolutionary dynamics, nutrient enrichment

SCHOLARONE™
Manuscripts

view

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

1 Evolution of nutrient acquisition: when space matters

2 Sébastien Barot¹, Stefan Bornhofen², Simon Boudsocq³, Xavier

3 Raynaud⁴ and Nicolas Loeuille⁵

4 ¹IRD, IEES-P, UMR 7618 (CNRS, INRA, UPMC, IRD), 46 Rue d'Ulm, F-75230 Paris

5 Cedex 05, France; ²Ecole Internationale des Sciences du Traitement de l'Information

6 (EISTI), avenue du Parc, 95011 Cergy-Pontoise Cedex, France; ³INRA, Eco & Sols,

7 UMR 1222, Centre INRA de Montpellier, 2 place Pierre Viala, 34060 Montpellier Cedex

8 01, France; ⁴UPMC, IEES-P, UMR 7618, 46 Rue d'Ulm, F-75230 Paris Cedex 05,

9 France; ⁵UPMC, IEES-P, 7 quai St.-Bernard, F-75252 Paris Cedex 05, France

10 Summary

11 **1.** Evolution of nutrient acquisition by plants should depend on two forces: local competition is
12 based on the capacity to exploit the local nutrient resource, and regional competition is based on
13 the capacity to occupy the whole landscape through seed production and dispersal.

14 **2.** We build a spatially-explicit simulation model where a limiting nutrient is recycled in each
15 local patch of a lattice by individual plants. The model includes both local and regional
16 competition.

17 **3.** Heterogeneity in nutrient availability and dispersal limitation mitigate the effect of
18 competition for the local nutrient resource and allow the evolution of lower rates of nutrient
19 uptake. Our spatially explicit model suggests that evolution in richer ecosystems selects
20 "expensive" strategies (high acquisition, low conservation of resources) compared to poor
21 ecosystems.

22 4. Low rates of nutrient acquisition can be considered as a form of altruism because they leave
23 more resource available for other individuals. Our model thus suggests that the influence of
24 spatial processes on the evolution of altruism is pervasive and is linked to key aspects of
25 ecosystem functioning.

26 5. Because our model both includes regional and local competition, evolution does not minimize
27 the availability of mineral nutrient, though evolution or species replacement is often thought to
28 minimize the availability of nutrient. Taken together, our work confirms that the interplay
29 between local and regional competition is critical for the evolution of plant nutrient strategies
30 and its effect on ecosystem properties.

31 **Key words:** nutrient acquisition, nutrient conservation, spatial model, local vs. regional
32 competition, evolution of altruism, spatial heterogeneity, dispersal, plant traits, eco-evolutionary
33 dynamics, nutrient enrichment.

34 Introduction

35 Spatial heterogeneity in the availability of mineral nutrients and the spatial structure of plant
36 populations must be taken into account to predict the evolution of plant strategies to acquire
37 mineral nutrients because: (1) Individual growth and reproduction depend on the local
38 availability of the mineral resource. (2) A mutant can establish locally only competing with
39 locally growing individuals and not with the whole resident population. (3) The ability of a
40 mutant to establish locally depends on the modifications it may impose to the local nutrient
41 availability even if its density is negligible at the population scale, i.e. the local success of a
42 mutant depends on positive or negative feedbacks it builds with its local environment. These

43 points emphasize local processes, but these processes only take place because the mineral
44 resource and plant individuals are spatialized.

45 Studies of adaptive plant strategies have already pointed out the importance of nutrient
46 heterogeneity and the ecological and evolutionary influences of two scales of competition
47 (Loreau 1998; Barot *et al.* 2014). (1) Within-cycle competition (local competition throughout
48 this paper) arises between strategies/individuals that are locally competing for the same nutrient
49 resource (thus sharing the same material cycle). (2) Between-cycle competition (regional
50 competition throughout this paper) arises between strategies/individuals growing spatially-apart,
51 therefore not directly competing for the same local resources (not sharing the same material
52 cycle) but competing regionally for space and access to the regional pool of nutrient. While local
53 competition is based on the capacity to exploit the local nutrient resource (which for example
54 involves the investment into a dense root system), regional competition is based on density-
55 dependence and the capacity to occupy the whole landscape (e.g. the whole surface of a plot, all
56 local patches of a spatial model) through the production of seeds and their dispersal. Thus, these
57 two scales of competition may lead to conflicting selection pressures because the strategy (i.e.
58 the resource allocation) maximizing the exploitation of a local resource and its accumulation into
59 the biomass is likely to be different from the strategy maximizing the number of individuals and
60 their distribution over the whole landscape (Loreau 1998; Barot *et al.* 2014). This is likely to
61 impede the minimization of nutrient availability predicted by the R* theory (Tilman 1982). Most
62 evolutionary models of nutrient cycling are mean-field models (Loeuille, Loreau & Ferrière
63 2002; Menge, Levin & Hedin 2008; Boudsocq, Barot & Loeuille 2011) and only predict the
64 evolutionary consequences of local competition. On the one hand, some models have addressed
65 the evolution of nutrient cycling related traits in a spatial context (de Mazancourt, Loreau &

66 Dieckmann 2001; Loeuille & Leibold 2008). However, these studies do not directly aim at
67 assessing the respective influence of local and regional competition. On the other hand, many
68 models address the interplay between local and regional competition but without addressing
69 explicitly evolutionary issues (e.g. Tilman 1994; Gravel *et al.* 2010) (see also a review
70 Fagerström & Westoby 1997).

71 At least two types of process should influence the respective importance of local and
72 regional competition on the evolution of nutrient acquisition by plants. First, the primary
73 producer capacity to colonize new patches through dispersal influences the intensity of local
74 competition between local mutants and the resident population through density-dependant
75 effects. Second, lateral fluxes of nutrient between patches may decrease the heterogeneity in
76 nutrient availability. Hence, both dispersal and nutrient fluxes between patches mitigate the
77 effect of regional competition (Loeuille & Leibold 2008). Through these processes, the whole
78 nutrient resource tends to be shared at the regional scale between all individuals and between the
79 various nutrient strategies they may display. This should increase the strength of selective
80 pressures for the uptake of local resources and should thus push towards the evolution of higher
81 rates of nutrient acquisition. Taken together, a mean-field model should lead to the evolution of
82 higher rates of mineral nutrient acquisition than a spatial model and, in a spatial model,
83 increasing seed dispersal and homogenizing fluxes of mineral nutrients should lead to the
84 evolution of higher rates of nutrient acquisition.

85 To assess the influence of local and regional competition and the impact of seed dispersal
86 and homogenizing nutrient fluxes on the evolution of nutrient acquisition we developed a
87 spatially-explicit model where primary producers compete locally for a heterogeneously
88 distributed limiting nutrient and regionally through seed production and their dispersal. Our

89 model is based on two previously published models: (1) a mean-field analytical model
90 (Boudsocq, Barot & Loeuille 2011) that predicts the evolution of nutrient acquisition, (2) a
91 spatially-explicit and individual-based model (Barot *et al.* 2014) that predicts the evolution of
92 plant capacity to control the mineralization of their litter. In the non-spatial model (Boudsocq,
93 Barot & Loeuille 2011), the fitness of a mutant only depends on the immediate balance between
94 acquisition and turnover of nutrient and this balance does not depend on inputs of nutrients to the
95 ecosystem, nutrient losses from the ecosystem or mineralization. Thus the mutant fitness cannot
96 depend on its capacity to modify its local environment. In contrast, in our spatial model, the
97 fitness of a mutant depends, through the building of local feedback loops and through
98 heterogeneity in nutrient availability, on complex interactions between all nutrient fluxes and the
99 mutant strategy for nutrient acquisition.

100 Taken together, the characteristics of our model should allow the evolution of the
101 diversity of plant strategies observed in different environments (Grime 2001; Craine 2009) and,
102 in particular, to conservative (low rate of nutrient uptake and efficient conservation of the
103 nutrient within the biomass) and acquisitive plants (high rate of nutrient uptake and poor
104 conservation of the nutrient within the biomass). Using this model we test the following
105 hypotheses: (1) The evolution of nutrient acquisition strategies depends on the relative strength
106 of local versus regional competition and spatial processes that influence this strength: dispersal
107 and lateral nutrient fluxes. More specifically, we expect increasing dispersal and lateral fluxes of
108 nutrients to select for higher rates of nutrient uptake. (2) The evolution of nutrient acquisition
109 depends on the nutrient richness of the ecosystem and all fluxes influencing this richness.
110 Namely, a gradient of nutrient enrichment leads to the evolution of Stress-tolerators with low
111 rates of nutrient uptake at the nutrient-poor end of the gradient and Competitors with high rates

112 of nutrient uptake at the other end of the gradient (Chapin III 1980; Grime 2001; Loeuille &
113 Loreau 2004). (3) Ecosystem properties should be impacted by the evolution of nutrient
114 acquisition. It has been proposed that local resource competition should allow the species that
115 minimizes resource availability to exclude all other species and this should maximize the
116 biomass of primary producer and primary production (Tilman 1982; Loreau 1998). On the
117 contrary regional competition should impede this minimization of the availability of mineral
118 nutrients (Loreau 1998).

119 **Description of the Model**

120 MODEL STRUCTURE

121 The model (see Fig. S1-A and Fig. S1-B and the list and description of parameters in Table 1) is
122 a spatial and individual-based version of a mean-field model of recycling of a limiting mineral
123 nutrient (Boudsocq, Barot & Loeuille 2011) between plants (P), dead organic matter (D) and
124 mineral nutrient (N). The limiting nutrient can virtually be any of the mineral nutrients that limit
125 primary production (generally nitrogen or phosphorus). Note that this previous model was not
126 individual-based. Here, the spatial features and the dynamics of individuals are modelled as in
127 Barot et al.'s model (Barot *et al.* 2014). Sessile terrestrial plants interact locally with soil nutrient
128 cycling and disperse through space by seed production. Our model thus couples a nutrient
129 cycling model with a spatial demographic model.

130 Space is described as a lattice of contiguous patches, each containing two abiotic nutrient
131 compartments (D , N) and one nutrient compartment for each locally growing plant individual (P_i ,
132 Fig. S1-A). Nutrient cycling and individual plant growth is modelled through the integration of a
133 system of differential equations in each patch. There is one equation for each plant individual so

134 that demographic events lead to the suppression (death) and addition (birth) of equations. We
135 assume that one time step corresponds to one year so that demographic events occur at the end of
136 each year whereas nutrient cycling is calculated on a sub-year basis through the integration of the
137 differential equations.

138 The model takes into account three types of ecological processes that are detailed next
139 section: (1) within-patch nutrient dynamics, (2) between-patch nutrient dynamics, corresponding
140 to fluxes of mineral nutrient that tend to homogenize resource availability and (3) demography of
141 plants. In order to assess the importance of space in the observed model dynamics, we compare
142 simulations with a 400 patch lattice to others where the lattice has only one larger patch (non-
143 spatial model). Individual plants growing in the same patch are competing for mineral nutrient
144 within this patch, as implemented in many spatial and individual-based models of plant growth
145 and competition (Czárán & Bartha 1992; Berger *et al.* 2008). There is a single source of
146 heterogeneity: the interaction between nutrient cycling and plant demography, and the
147 underlying stochasticity. More mineral nutrient is transiently available in local patches that
148 contain fewer individual plants or a lower total biomass of plants.

149 The size of the patches has to be chosen appropriately, so that they only host a relatively
150 low number of plant individuals that belong to the same competitive neighbourhood. More
151 precisely, this number of individuals and patch size are determined by inputs of mineral nutrient
152 and organic matter (see below). For example, patches of one square meter (as implemented in
153 our simulation, see below) would be relevant to model small and short-lived herbaceous plants
154 for which about 50 individuals may be thought as locally competing. If plants live longer,
155 individuals may grow larger (e.g. perennial bunch grasses) and only a few individuals compete

156 for the same nutrient resource in each one-square-meter patch. The different parameters of the
 157 model and their default values are given in Table 1.

158 ECOLOGICAL DYNAMICS

159 *Within-patch nutrient dynamics*: In each patch (subscript j), the limiting nutrient is recycled
 160 within the patch (through the absorption of mineral nutrient, the turnover of the biomass of the
 161 primary producer, and the mineralization of the resulting dead organic matter) and each patch
 162 receives inputs and loses nutrients. In contrast to the original non-spatial model, each patch of
 163 the lattice can be occupied by several individuals (index i , nutrient content $P_{j,i}$) that compete for
 164 the local pool of mineral nutrient (thus the number of differential equations for each patch varies
 165 with the number of individuals growing in the patch). We assume that plants are identical for all
 166 parameters except their rate of nutrient uptake ($u_{Nj,i}$) and their rate of nutrient turnover to the
 167 ecosystem ($d_{Pj,i}$). There is a rate of mineralization (m_D) and a rate of nitrogen fixation (f_P , if the
 168 limiting nutrient is nitrogen and if the plant is able to fix nitrogen). Nutrient inputs in its organic
 169 matter form (R_D) and in its mineral form (R_N) are independent of the size of the nutrient pools.
 170 Nutrients diffuse out of the ecosystem through fixed rates, respectively l_P , l_D , and l_N for the P, D
 171 and N compartments respectively. l_P denotes losses of nutrients through fires in terrestrial
 172 ecosystems. Dead organic matter is lost through erosion and leaching (l_D). Mineral nutrients are
 173 lost through leaching and denitrification (l_N). For each patch of the lattice, the equations thus
 174 become:

$$175 \quad \frac{dP_{j,i}}{dt} = u_{Nj,i} N_j P_{j,i} - (d_{Pj,i} + l_P - f_P) P_{j,i} \quad (\text{for each individual } i \text{ of local patch } j),$$

$$176 \quad \frac{dD_j}{dt} = \sum_i d_{Pj,i} P_{j,i} + R_D - (m_D + l_D) D_j,$$

$$177 \quad \frac{dN_j}{dt} = m_D D_j + R_N - \left(\sum_i u_{N_j,i} P_{j,i} + l_N \right) N_j + H (\bar{N} - N_j). \text{ (Eq 1)}$$

178 *Between-patch nutrient dynamics.* These dynamics correspond to processes leading to an
 179 homogenization of nutrient content between patches. To do so, the quantity $H (\bar{N} - N_j)$ is
 180 added to the derivative of the N_j value of each patch. H varies between 0 and $+\infty$ and \bar{N} is the
 181 mean nutrient availability calculated over all patches of the lattice. Nutrient availability tends to
 182 homogenize at the regional scale when H increases (all local N_j values are equal to \bar{N}), i.e.
 183 higher H values correspond to a higher rate of nutrient mixing among patches. When H increases
 184 the mineral resource is more and more shared between all individuals of the population and
 185 competition for nutrient acquisition is no longer purely local, i.e. between individuals growing in
 186 the same patch. Hence, studying the effect of H constitutes a theoretical experiment to determine
 187 the influence of the way the mineral resource is shared at the population scale.

188 *Demography.* Discrete demographic events, i.e. mortality and reproduction, occur at the
 189 end of every year (Fig. S1-B). While $d_{P_j,i}$ denotes the rate of nutrient turnover of the primary
 190 producer because of the death of parts of the primary producer (death of roots and leaves,
 191 exudate production), μ denotes individual mortality (Table 1). The amount of nutrient contained
 192 in dead individuals is transferred to the local dead organic matter compartment (D_j).

193 At the end of each year, individuals in the lattice may produce seeds that are dispersed on
 194 the lattice. Parameter ρ is the percentage of the stock of nutrient contained in each individual at
 195 the end of the year that is allocated to seed production. γ is the quantity of nutrient in a seed. The
 196 number of seeds produced by the individual i (in patch j) is thus the largest previous integer of
 197 $P_{j,i} \rho / \gamma$. Seed dispersal is modelled by moving each seed and creating a new individual with a

198 nutrient content γ in another patch. We choose a random angle (between 0 and 360°) to
 199 determine the direction of dispersal and consider that the dispersal distance from the centre of the
 200 patch of the mother primary producer follows a centred normal distribution (mean=0) with
 201 standard deviation σ (if the distance is negative 180° is added to the original angle): when σ
 202 increases dispersal distance tends to increase. There is no habitat choice. For simplification,
 203 production of seed starts at age 1 year for individuals that are large enough at this age
 204 ($P_{j,i} \rho > \gamma$). Note that individuals falling under a threshold nutrient stock (i.e. the nutrient content
 205 of a seed) are also considered to die.

206 EVOLUTIONARY DYNAMICS

207 As in the original non-spatial model (Boudsocq, Barot & Loeuille 2011), we study the evolution
 208 of a trait s that can be considered as the investment of the primary producer into the acquisition
 209 of mineral nutrient. The trait s can be either negative or positive. When s increases, both the
 210 investment into nutrient acquisition and nutrient turnover increase. For any individual i of trait
 211 $s_{j,i}$, the nutrient uptake rate and the nutrient turnover of the primary producer are linked via the
 212 following trade-off equations:

$$213 \quad u_{Nj,i} = u_{N0} e^{bs_{j,i}},$$

$$214 \quad d_{Pj,i} = d_{P0} e^{cs_{j,i}}.$$

215 This trade-off is justified by several complementary mechanisms (see details in the Discussion).
 216 For example, the more a plant invests into the development of its root system, the uptake and
 217 assimilation of nutrients, the less resource are allocated to maintenance and defence functions.
 218 Indeed, the existence of trade-offs between nutrient uptake and anti-herbivore defences have

219 been documented (Herms & Mattson 1992; Mauricio 1998). This mechanism should increase the
220 turnover of the plant biomass and losses of mineral nutrients contained in this biomass.

221 The relation between b and c (real-valued positive parameters) determines the strength of
222 the trade-off: if $b < c$, the trade-off function displaying $d_{Pj,i}$ as a function of $u_{Nj,i}$ is convex
223 (curving up); otherwise if $b > c$ the trade-off function is concave (curving down); if $b=c$ the
224 nutrient turnover increases linearly with the uptake capacity. Taken together, b determines the
225 benefit of the investment into nutrient uptake (s), while c determines its cost. Because of this
226 trade-off, s can be viewed either as a strategy of nutrient capture or a strategy of nutrient
227 turnover. Since the model is not analytically tractable, numerical simulations have to be run.
228 Each new seed has a probability p_m of being a mutant for s , in which case the size of the mutation
229 is drawn from a centred normal distribution with a standard deviation m .

230 The capacity of a strategy to persist or to invade a resident strategy, and thus evolutionary
231 dynamics, depend (1) on the effect of the investment into nutrient acquisition on individual
232 growth in local patches, (2) the capacity to disperse from patch to patch through seed production.
233 Thus selection occurs via the growth of individuals during each time step (the growth of each
234 individual depends on its strategy for nutrient acquisition), and via demography at the end of
235 each time step: the size of individuals having a non-optimal uptake strategy decreases and reach
236 the threshold under which they die, i.e. the size of a seed (see above in the section about
237 demography), and larger individuals produce more seeds.

238 IMPLEMENTATION

239 A simulation platform has been implemented in the Java programming language (1.5). The
240 integration of the differential equations is based on the classical fourth-order Runge-Kutta

241 method with an integration step $\Delta t = 0.1$ year. In our simulations H varies between 0 (no
242 homogenization) and 10 (strong homogenization). Fig. S6 shows that $H=10$ leads to a null
243 standard deviation of nutrient availability (perfect homogenization of the mineral nutrient
244 availability).

245 Parameters chosen for nutrient cycling are adapted from Woodmansee et al. (1981) who
246 describe nitrogen cycling in a temperate grassland (see Table 1). Unless stated in figure captions
247 (main text and the supplementary material) we keep these parameter values. We consider one
248 square meter wide patches. Unless stated, the primary producer is considered to be a short-lived
249 plant. For mutations we always used $p_m = 10^{-4}$ and $m = 0.05$. The grid size is always 20 X 20
250 patches except in one occasion (hereafter one-patch model) where only one patch is modelled but
251 increasing accordingly input of mineral nutrients and organic matter (X400) to support the same
252 total biomass as in the 20X20 patch model. This corresponds to a shift between 400 one square
253 meter patches to one single 400 m² patch. The lattice is wrapped on a torus to handle seed
254 dispersal. Variables following normal distribution (dispersal distance and mutation size) are
255 simulated using the polar method (Marsaglia & Bray 1964).

256 Simulations are initialized with a single seed in each patch and a fixed amount of mineral
257 nutrient (20 dg m⁻²). 0 is always chosen as a starting value for s . Preliminary analyses have
258 shown that results are not sensitive to these initial conditions in terms of ecological (N^* , D^* , P^*)
259 and evolutionary equilibriums (s^*), provided that these conditions allow for the initial survival of
260 the plant species.

261 Results (Fig. 1, 2) are presented using locally-weighted polynomial regressions to
262 compute trend lines. All figures were produced using R 2.10 (2010). In the main text we focus on
263 the evolutionary effects of inputs of mineral nutrient to the ecosystem. Fig. S3 displays results

264 concerning the effects of the rate of mineral nutrient losses from the ecosystem and the rate of
265 dead organic matter mineralization that also influence the availability of mineral nutrient.

266 **Results**

267 EVOLUTIONARY DYNAMICS

268 The simulation model leads to the same types of evolutionary outcomes as the analytical model
269 of Boudsocq et al. (2011): (i) a stable evolutionary equilibrium (Continuously Stable Strategy or
270 CSS, Eshel 1983), (ii) a tragedy-of-the common scenario (Hardin 1968) where the primary
271 producer compartment asymptotically decreases, while uptake and productivity increase,
272 eventually leading to the primary producer extinction, and (iii) an accumulation of nutrient in all
273 compartments due to an efficient recycling (inputs of nutrient are higher than outputs). In this
274 last case, the model predicts the infinite accumulation of the limiting nutrient, which is not
275 realistic but denotes cases where evolution drives the primary producer to switch to an other
276 limiting factor. Evolutionary branching does not occur.

277 In particular, when the benefit of the investment in nutrient acquisition is low in
278 comparison to its cost ($b < c$) and when inputs of nutrient to the primary producer compartment
279 (nitrogen fixation) are equal or lower to its rate of mineral nutrient turnover ($f_p \leq l_p$),
280 simulations lead to a CSS we note s^* . Simulations with identical parameter values always lead
281 qualitatively to the same results, but the inherent stochasticity of the model (mortality, seed
282 dispersal and mutations) may lead to small differences in the s^* observed at the end of a
283 simulation. All the results displayed hereafter correspond to CSS cases because of the parameter
284 values we choose ($f_p = l_p = 0$). To insure the meaningfulness of the results we always run
285 simulations long enough to reach the evolutionary equilibrium (400000 time steps while 100000

286 time steps are generally enough to reach the CSS). Fig. S2 displays an example of evolutionary
287 dynamics leading to an evolutionary equilibrium together with the corresponding pairwise-
288 invisibility plot (Geritz *et al.* 1998) that proves that this is a case of CSS. Such dynamics have
289 been simulated for each point of Fig. 1 and 2.

290 Besides, for any set of ecological parameters leading to a CSS and for any s value, the
291 ecological dynamics leads to equilibrium values for P (sum of P_i of all individuals in a local
292 patch), N and D when their values are averaged over all patches of the model (see also Fig. S2).
293 We use the same notation (*) for these ecological equilibriums (P^* , N^* ...) at the evolutionary
294 equilibrium (CSS, s^*).

295 CAPACITY TO ACQUIRE NUTRIENTS AT THE EVOLUTIONARY EQUILIBRIUM

296 Consistent with our predictions, the homogenization of mineral nutrient and the dispersal
297 capacity of the primary producer increase the evolved primary producer capacity for nutrient
298 acquisition (Fig. 1).

299 The evolved capacity for nutrient acquisition (s^*) increases with inputs of mineral
300 nutrient to the ecosystem (R_N) whatever the trade-off shape (Fig. 2 A), the individual mortality
301 (Fig. 2 B), the intensity of mineral nutrient homogenization (Fig. 2 C), or the dispersal capacity
302 (Fig. 2 D). The less restrictive the trade-off (increasing b values), the higher the evolved s^*
303 values. For increasing individual mortality rates the amplitude of the effect of nutrient inputs on
304 the evolved nutrient acquisition becomes higher. As in Figure 1, homogenization of mineral
305 nutrient and seed dispersal increases the evolved capacity for mineral nutrient acquisition. This
306 effect of nutrient homogenization disappears for high inputs of mineral nutrients. In other words,
307 the evolved capacity for nutrient acquisition always increases with nutrient inputs but this effect
308 is weaker when nutrient homogenization increases. The non-spatial model (only one patch) leads

309 to a much higher value ($s^*=0.77$, Fig. 2 C) of the capacity for nutrient acquisition than the
310 maximum rate of nutrient homogenization and a very efficient seed dispersal ($s^*=0.27$, Fig. 1).
311 Perfect homogenization brought by the non-spatial model (one patch) leads to a higher evolved
312 capacity for nutrient acquisition than the mere homogenization ($H=10$) of mineral nutrient
313 availability (Fig. 2 C). In the case of perfect homogenization, inputs of nutrient no longer impact
314 the evolved capacity for nutrient acquisition.

315 Increasing the mineralization rate and decreasing the rate of mineral nutrient loss from
316 the ecosystem have the same effect on the evolution of nutrient acquisition as increasing nutrient
317 inputs (Fig. S3).

318 ECOSYSTEM PROPERTIES AT THE EVOLUTIONARY EQUILIBRIUM

319 Figure 3 describes the ecological effect of the investment into nutrient acquisition (s) and
320 nutrient inputs (R_N) on the average availability of mineral nutrient and the mean size of the
321 primary producer compartment (for each combination of R_N and s , the model is run till an
322 ecological equilibrium is reached but without allowing evolution). There is always a unique s
323 value that minimizes the mean N , or maximizes the mean P , for each R_N value. However, the
324 values that minimize N and maximize P are always distinct. On the same figure is plotted the
325 outcome of the evolution of s for each R_N value. Evolution neither minimizes N nor optimizes P .
326 Similarly, evolution does not maximize primary production (results not shown). The difference
327 between the evolved N^* value and the lowest possible N^* value increases with nutrient
328 enrichment (Fig. 3, see also confirmation in Fig. S4).

329 **Discussion**330 SPATIALIZATION AND LOW HOMOGENIZING FLUXES ALLOW FOR THE
331 EVOLUTION OF ALTRUIST NUTRIENT STRATEGIES

332 Increasing dispersal and lateral fluxes of nutrient simultaneously homogenizes the availability of
333 the mineral resource and increases the evolved capacity for nutrient acquisition at the CSS (s^*).
334 This happens because these two processes mitigate the influence of regional competition and
335 increase the evolutionary influence of local competition for a shared mineral resource, which
336 tends to push towards less altruist strategies, i.e. towards the evolution of higher uptake rates of
337 this resource. This interpretation is detailed below and is supported by three converging lines of
338 researches: (1) models and experiments showing that heterogeneity in resource availability and
339 spatial structuration of populations promote the evolution of altruism (Lion & van Baalen 2008;
340 Bachmann *et al.* 2013), (2) the framework of the tragedy of the common that shows that
341 competition for a common resource pushes towards the evolution of higher rates of consumption
342 of this resource (Rankin, Bargum & Kokko 2007), (3) researches suggesting that a low rate of
343 resource consumption can be considered as altruist (Pfeiffer, Schuster & Bonhoeffer 2001; Kreft
344 2004; Burtsev & Turchin 2006).

345 It is logical that increasing dispersal and increasing resource homogeneity lead to s^*
346 values closer to the ones obtained for the non-spatial model (only one patch) where only local
347 competition occurs. The non-spatial model, resource homogenization, and efficient dispersal lead
348 to high s^* values because competition for a shared resource favours strategies that increase the
349 acquisition of the resource (Rankin, Bargum & Kokko 2007). In the non-spatial model, the
350 resource is fully shared by all individuals, while with homogenizing fluxes of nutrient the
351 resource is only partially shared. In the same vein, when dispersal ability increases, plants with

352 alternative competing strategies are more likely to compete for the same local resource in the
353 same patches. In the non-spatial model and in the fully-homogenized version of the model the
354 mineral resource is fully-shared by all individuals, but the former case leads to much higher s^*
355 values. Indeed, in the non-spatial model, only local competition for the nutrient acts as a
356 selective pressure unaltered by any spatial process while, in the homogenized models, mutants
357 are locally selected by local competition but still have to invade the whole population, which
358 requires producing seeds and dispersing them, thus involving regional competition.

359 Our evolutionary results reveal the essence of the tragedy of commons (Rankin, Bargum
360 & Kokko 2007): competition for a shared nutrient resource drives the evolution towards higher
361 rates of resource use thereby forbidding the maximization of collective properties (e.g.
362 population biomass). Such a scenario has already been pointed out for roots and soil resources
363 (Gersani *et al.* 2001; Craine 2006). A low rate of resource or social good acquisition is
364 considered as a form of altruism since it leaves more resource available for other individuals
365 (Burtsev & Turchin 2006; Rankin, Bargum & Kokko 2007). This follows classical definitions of
366 altruism (Hamilton 1963; West, Griffin & Gardner 2007): having a low rate of nutrient uptake
367 deprives a plant of an available resource, and can thus be considered as costly, while this benefits
368 to other plants for which more resource is available. As in all cases of evolution of an altruist
369 trait, the obvious cost of the altruist trait must be counterbalanced by indirect fitness benefits
370 (West, Griffin & Gardner 2007). In our model, such indirect benefits emerge from the interaction
371 between spatial variations in resource availability and dispersal processes. As such, our results
372 are consistent with other observations of the emergence of altruism due to spatial structure (Kerr
373 *et al.* 2006; Bachmann *et al.* 2013).

374 Our results can therefore be interpreted in terms of evolution of altruism and are
375 consistent with the established finding that it is promoted by heterogeneity, low dispersal and
376 spatial isolation of strategies (Lion & van Baalen 2008). This is also consistent with the fact that
377 low dispersal increases the impact of kin selection and facilitates group selection (Lion & van
378 Baalen 2008) that are also known to foster the evolution of altruism (Nowak 2006). Indeed, in
379 our model, the mixture of regional and local competition allows for the apparition of selection at
380 the scale of local patches. Evolution of altruism has often been studied from the point of view of
381 social interactions (Ferrière & Michod 1995). Our results and others on the evolution of bacteria
382 (Pfeiffer, Schuster & Bonhoeffer 2001), predators (Goodnight *et al.* 2008) and parasites
383 (Haraguchi & Sasaki 2000) suggest that the influence of spatial processes on the evolution of
384 altruism is pervasive and that many key aspects of ecosystem functioning should be analysed
385 through this prism.

386 NUTRIENT ENRICHMENT SELECTS FOR HIGHER RATES OF RESOURCE 387 ACQUISITION

388 In our spatial model, increasing nutrient inputs to the ecosystem (Fig. 2), increasing the
389 mineralization rate and decreasing the rate of nutrient loss from the ecosystem (Fig. S3) lead to
390 the evolution of higher rates of nutrient acquisition. These three parameters influence three
391 features in interactions along the eco-evolutionary dynamics implemented in the model: the
392 mean availability of mineral nutrient, the variability in this availability and the number of
393 individual plants in each patch. Two arguments support the important role played by
394 heterogeneity in nutrient availability and the number of individuals in a patch: First, in the mean-
395 field analytical model (Boudsocq, Barot & Loeuille 2011) and in the non-spatial version of our
396 simulation model (only one patch), the evolutionary effect of resource enrichment disappears. In

397 our spatial model, individuals with a higher rate of nutrient uptake benefit more from this
398 strategy, on the average, if the heterogeneity in nutrient availability is low. Indeed, in a patch
399 with a low nutrient availability a higher rate of nutrient uptake leads for an individual to a low
400 increase in the nutrient acquired by this mutant that nevertheless pays a high cost for this
401 increase in terms of nutrient turnover because of the trade-off. Second, we have shown that
402 homogenizing fluxes affect the evolution of the investment into nutrient acquisition because they
403 modulate the respective influence of local and regional competition, which is an other way to
404 state that they modulate the number of individuals that are locally competing for the same
405 resource. Thus, the evolved investment into nutrient acquisition should increase with the mean
406 number of individuals in each patch, i.e. in each competitive neighbourhood, as with
407 homogenizing fluxes.

408 This general rationale is supported by complementary results. An important difference
409 between the spatial and non-spatial models is that spatial models allow for heterogeneity in
410 nutrient availability. In our simulation model this heterogeneity is sustained by the stochasticity
411 in plant demography. For example, if several individuals die at the same time in a patch, more
412 mineral nutrient is likely to be locally available (more organic matter to be mineralized and less
413 biomass to take up mineral nutrients). Thus, the local dynamics of nutrient pools are a permanent
414 succession of transient dynamics (see Fig. S5). This allows an increase in nutrient inputs (R_N), an
415 increase in the mineralization rate (m_D) and a decrease in the rate of nutrient loss from the
416 ecosystem (l_N) to modify the average mineral nutrient availability and the spatial and temporal
417 variability in this availability (see Fig. S6 for nutrient inputs). On the contrary, in the analytical
418 non-spatial model, nutrient availability is kept at an equilibrium that does not depend on nutrient
419 inputs, mineralization or the rate of mineral nutrient loss from the ecosystem (see formula for the

420 ecological equilibrium, Boudsocq, Barot & Loeuille 2011). Fig. S6 shows that at the ecological
421 scale (the acquisition of mineral nutrient is not allowed to evolve) the variability in the
422 availability of mineral nutrient always decreases with inputs of mineral nutrient while the mean
423 availability of mineral nutrient may increase or decrease with these inputs. Moreover, at the
424 evolutionary equilibrium, the number of individuals by patch also always increases with nutrient
425 inputs (see Fig. S7).

426 While it is beyond the objective of this article to fully disentangle the ecological and
427 evolutionary feedbacks between local resource availability, its heterogeneity and the local
428 density of plants, a way to study these feedbacks is to disrupt them. Fig. S8 shows that
429 constraining the maximum number of individuals to one individual reverses the effect of nutrient
430 inputs: increasing nutrient inputs decreases the evolved investment into nutrient acquisition. This
431 suggests that: (1) When a maximum of one individual by patch is allowed, increasing nutrient
432 inputs leads to the evolution of lower rates of nutrient uptake because more resource becomes
433 available for a single individual and because lower rates of uptake allow absorbing large quantity
434 of nutrient. (2) When the number of individuals by patch is allowed to increase due to an
435 increase in nutrient inputs (see S6 and S7), higher uptake rates evolve precisely because the
436 number of individuals sharing the local resource increases. These results, the comparison
437 between the spatial and non-spatial models and the effects of homogenizing fluxes confirm that
438 heterogeneity in nutrient availability and local density in competitors determine the evolutionary
439 effects of nutrient enrichment. This general interpretation is parsimonious: the effects of nutrient
440 enrichment, seed dispersal, and heterogeneity in nutrient availability would all be linked to the
441 way these processes modulate the balance between local and regional competition and the
442 number of individuals sharing the same local mineral resource. When nutrient inputs increase,

443 the number of locally competing individuals increases, which leads to a stronger evolutionary
444 influence of local competition, and a stronger selection for nutrient acquisition, as when seed
445 dispersal or homogenizing fluxes of nutrient increase. In both cases (homogenizing fluxes and
446 nutrient enrichment), the local pools of mineral nutrient are more shared or shared among more
447 individuals.

448 EVOLUTION OF ACQUISITIVE AND CONSERVATIVE PLANTS

449 One of our main conclusions is that, whatever the underlying mechanisms, when nutrient
450 resource is not fully shared by all competitors and is heterogeneously distributed, nutrient
451 enrichment drives the evolution towards higher rates of nutrient acquisition. Because of the
452 trade-off between nutrient acquisition and nutrient turnover in primary producers, this also
453 means that strategies more conservative for nutrient evolve in nutrient poor ecosystems.
454 Moreover, our rate of nutrient acquisition is analogous to a relative growth rate (rescaled by the
455 availability of nutrient). Consequently, our results are consistent with Grime's theory about the
456 evolution of terrestrial plants in nutrient-rich and nutrient-poor ecosystems: Competitors (i.e.
457 acquisitive plants) have evolved high relative growth rates in nutrient-rich ecosystems, while
458 Stress-tolerators (i.e. conservative plants) have evolved lower relative growth rates and lower
459 nutrient turnovers in nutrient-poor ecosystems (Grime 1977; Grime 2001).

460 Our model is too simple to predict specific traits of low- and high- nutrient plants such as
461 shoot root-ratio, root length or defences against herbivores. Similarly, it does not take into
462 account the interactions between competition for light (i.e. carbon) and nutrient (see above).
463 However, the generality of our model uncovers fundamental aspects of the evolutionary
464 consequences of competition for a limiting but recycled resource: (1) The trade-off we
465 implement is the most fundamental one concerning the management of a resource by an

466 organism. How efficiently does the organism capture the resource? How long is it able to
467 conserve it? (2) The existence of this trade-off that constrains the evolution of acquisitive and
468 conservative plants is supported by several documented mechanisms. First, the investment into
469 nutrient acquisition may reduce the investment into defences against herbivores and protective
470 structures (Herms & Mattson 1992). Note that this suggests that our results also give an
471 evolutionary support for plants in nutrient-poor environments to be more defended against
472 herbivores (Coley, Bryant & Chapin 1985). Second, plants that absorb large amounts of mineral
473 nutrient tend to have higher nutrient concentrations in their biomass, which decreases the
474 longevity of roots and leaves and leads to stronger herbivore attacks (Eissenstat *et al.* 2000; Silla
475 & Escudero 2004; Endara & Coley 2011). Third, increasing the capacity to uptake mineral
476 nutrients requires increasing the biomass of roots and especially of thin roots that have a short
477 life-span. These three mechanisms should increase the turnover of plant biomass and the
478 turnover of the nutrients contained in this biomass. Taken together, our results give some
479 evolutionary support for the existence of a root economic spectrum (Reich & Cornelissen 2014)
480 that parallels the leaf-economic spectrum (Wright *et al.* 2004).

481 We have shown that the evolution of nutrient acquisition depends on individual mortality:
482 the evolutionary effect of nutrient enrichment depends on longevity, and the higher the longevity
483 the less sensitive the evolved investment into nutrient acquisition to nutrient availability. The full
484 interpretation of this result goes beyond the objectives of this article, but this confirms that
485 nutrient acquisition strategies should interact with life-history traits (e.g. individual mortality,
486 investment into reproduction, age at first reproduction, seed size, dispersal...). Together, they
487 modulate the influence of local and regional competition and influence plant capacity to
488 conserve mineral nutrients (e.g. long-lived plants conserve mineral nutrients longer than

489 annuals). Our framework could thus allow testing further the evolutionary background of plant
490 functional classifications that also involve plant life-histories (Grime 2001; Craine 2009). This
491 would require studying the joint evolution of functional and demographic traits and should lead
492 to the evolution of ruderals –the third strategy of Grime's CSR theory– that have not appeared in
493 our present model (see for example in Bornhofen, Barot & Lattaud 2011).

494 IMPACT OF EVOLUTION ON MINERAL NUTRIENT AVAILABILITY

495 That evolution does not maximize the biomass of the primary producer is caused, as in
496 the original mean field analytical model (Boudsocq, Barot & Loeuille 2011), by the basic
497 functioning of selection: selection is based on the per capita growth rate of mutants and not on
498 the ultimate biomass or primary productivity. The non-minimization of the evolved average
499 nutrient availability (N^*) contradicts both the non-spatial evolutionary model (Boudsocq, Barot
500 & Loeuille 2011) and Tilman's R^* theory (Tilman 1982). This influential theory suggests that
501 the ecological dynamics of species replacement due to competition for a limiting single resource
502 should lead to the exclusion of all species but the one that reduces the most the mineral resource
503 availability at equilibrium. Our new result must stem from the combination of local and regional
504 competition and the impeding of the minimization of nutrient availability by regional
505 competition (Loreau 1998). Indeed, local competition is only based on plant capacity to absorb
506 mineral nutrient while regional competition requires plant individuals to be large enough and to
507 be able to invest enough resource into seed production and seed dispersal. Local competition
508 thus tends to select for high rates of nutrient uptake while regional competition tends to select, in
509 a density-dependent way, for lower rates of nutrient uptake allowing plants to be larger and to
510 invest more into seed production. Besides, homogenization of the availability of the mineral
511 resource and an efficient dispersal increase the influence of competition for the mineral resource

512 because these processes drive this resource to be more evenly distributed among individual
513 plants and the strategies they represent. Therefore, homogenization of the resource and seed
514 dispersal decrease the difference between the evolved availability of mineral nutrient and the
515 lowest possible availability of mineral nutrient (Fig. S4), this difference being caused by regional
516 competition. As aquatic systems are supposedly more homogeneous than terrestrial systems, our
517 results might explain why most conclusive positive tests of the R* theory have been achieved in
518 aquatic systems (Miller *et al.* 2005) and very few, apart from Tilman's initial papers (Tilman &
519 Wedin 1991; Wedin & Tilman 1993), in terrestrial systems.

520 CONCLUSION

521 Our results about the evolution of nutrient acquisition in a spatial context and former results
522 about the evolution of plant capacity to control mineralization (Barot *et al.* 2014) support each
523 other and are complementary. Since mineralization does not impact individual fitness directly,
524 but only through a modification of the availability of the mineral resource, higher mineralization
525 cannot evolve at all in a non-spatial model. On the contrary, nutrient acquisition directly impacts
526 individual fitness so that spatial processes only modify the evolution of nutrient acquisition
527 predicted by the non-spatial model (Boudsocq, Barot & Loeuille 2011). However, in both cases
528 the relative influence of local and regional competition determines the evolutionary dynamics
529 and leads to comparable results: (1) In Barot *et al.* (2014) homogenization (through seed
530 dispersal or homogenizing fluxes of nutrient) leads to the evolution of lower mineralization rate
531 while in the present work homogenization leads to the evolution of higher rates of nutrient
532 uptake. (2) In Barot *et al.* (2014) nutrient enrichment leads to lower mineralization rates while in
533 the present work it leads to higher rates of nutrient uptake. (3) Both in Barot *et al.* (2014) and the
534 present work, the mixture of local and regional competition does not lead through evolution to

535 the maximization of the plant biomass or the minimization of the availability of the mineral
536 nutrient. We contend here that a low rate of nutrient uptake can be considered as an altruist
537 strategy because it leaves more mineral nutrient available to other plants (see above and a
538 classical definition, Hamilton 1963). Considering a high rate of mineralization as an altruist
539 strategy is probably even more obvious because it directly increases the availability of the
540 mineral resource for all plants of the local patch, while the plant that increases mineralization is
541 the only one to bear the cost of this increase in mineralization (the plant losses more nutrient).
542 With this joint interpretation of low rates of nutrient uptake and high rates of mineralization as
543 altruist strategies the results of the present work and the model by Barot et al. (2014) are fully
544 consistent. The higher the influence of local competition the less altruist the evolved strategy: the
545 evolved rate of mineral nutrient acquisition becomes higher and the evolved decomposition rate
546 lower. Similarly, an increase in nutrient inputs pushes plants towards less altruist strategies
547 probably because an increase in nutrient inputs tends to increase the local densities in
548 competitors (see above).

549 Starting from a very simple and general analytical model based on plant biomass we have
550 added some realistic features of plant competition: interaction between local and regional
551 competition and some simple features of plant demography. The resulting model remains general
552 and theoretical. Some features of the model may appear as unrealistic, but our work has paved
553 the way to the implementation of many other influential mechanisms to test the robustness of our
554 results: (1) plasticity in the investment into nutrient acquisition depending on nutrient availability
555 and local density of competitors (Cahill *et al.* 2010), (2) more refine functions for nutrient uptake
556 and local competition (see for example results in Raynaud & Leadley 2004), (3) implementation
557 of more realistic lateral nutrient fluxes corresponding to particular mechanisms (e.g. impact of

558 herbivores) and interactions between lateral fluxes of mineral nutrient and dead organic matter
559 (Gravel *et al.* 2010), (4) more realistic life-cycles to represent particular plant types (e.g. size
560 dependent survival, age at first reproduction, different seed sizes), (5) coevolution between
561 nutrient acquisition and life-history traits such as the investment into seed production or dispersal
562 (Craine 2009; Suzuki & Kimura 2011), (6) different shapes of trade-off (de Mazancourt &
563 Dieckmann 2004) between nutrient acquisition and nutrient turnover. In particular, while our
564 model always leads to the selection of a single nutrient uptake strategy, point (2) and the use of
565 an uptake function increasing with biomass with a downward curvature (Rastetter & Ågren
566 2002) should allow the coexistence of various strategies and may thus lead to the diversification
567 of the strategy of nutrient acquisition through evolutionary branching (Geritz *et al.* 1998).

568 In conclusion, the merit of our study is to link demographic processes, functional
569 processes, evolution and ecosystem properties, which still constitutes a very open research
570 avenue (Fussmann, Loreau & Abrams 2007). This allows us to link theories that are usually
571 developed independently, e.g. theories about the evolution of altruism and evolutionary impact
572 of spatial processes (Lion & van Baalen 2008) vs. the existence of distinct plant strategies
573 depending on resource availability (Grime 2001). In particular, while ecosystem ecology is not
574 prone to evolutionary interpretations, we suggest that many aspects of ecosystem functioning are
575 linked to the evolution of altruism.

576 **Acknowledgements**

577 SB and XR have been funded by the ANR project “3worlds” of the program “Intensive
578 Calculations and Simulations”. We thank E. Thébault for her relevant comments. We thank two
579 anonymous reviewers for their precious suggestions.

580 **References**

- 581 Bachmann, H., Fischlechner, M., Rabbers, I., Barfa, N., dos Santos, F.B., Molenaar, D. &
582 Teusink, B. (2013) Availability of public goods shapes the evolution of competing metabolic
583 strategies. *Proceedings of the National Academy of Sciences of the USA* 442, 75-78.
- 584 Barot, S., Bornhofen, S., Loeuille, N., Perveen, N., Shahzad, T. & Fontaine, S. (2014) Nutrient
585 enrichment and local competition influence the evolution of plant mineralization strategy, a
586 modelling approach. *Journal of Ecology*, 102, 357-366.
- 587 Berger, U., Piou, C., Schiffers, K. & Grimm, V. (2008) Competition among plants: concepts,
588 individual-based modelling approaches, and a proposal for a future research strategy.
589 *Perspectives in Plant Ecology, Evolution and Systematics*, 9, 121-135.
- 590 Bornhofen, S., Barot, S. & Lattaud, C. (2011) The evolution of CSR life-history strategies in a
591 plant model with explicit physiology and architecture. *Ecological Modelling*, 222, 1-10.
- 592 Boudsocq, S., Barot, S. & Loeuille, N. (2011) Evolution of nutrient acquisition: when adaptation
593 fills the gap between contrasting ecological theories. *Proc. Royal. Soc London B*, 278, 449-
594 457.
- 595 Burtsev, M. & Turchin, P. (2006) Evolution of cooperative strategies from first principles.
596 *Nature*, 440, 1041-1044.
- 597 Cahill, J.F., McNickle, G.G., Haag, J.J., Lamb, E.G., Nyanumba, S.M. & Clair, C.C.S. (2010)
598 Plants integrate information about nutrients and neighbors. *Science*, 328, 1657-1657.
- 599 Chapin III, F.S. (1980) The mineral nutrition of wild plants. *Annual Reviews on Ecology and*
600 *Systematics*, 11, 233-260.
- 601 Craine, J.M. (2006) Competition for nutrients and optimal root allocation. *Plant and Soil*, 285,
602 171-185.

- 603 Craine, J.M. (2009) Resource strategies of wild plants. Princeton University Press, Princeton,
604 New Jersey.
- 605 Czárán, T. & Bartha, S. (1992) Spatiotemporal dynamic models of plant populations and
606 communities. *Trends in Ecology and Evolution*, 7, 38-42.
- 607 de Mazancourt, C. & Dieckmann, U. (2004) Trade-off geometries and frequency-dependent
608 selection. *American Naturalist*, 164, 765-778.
- 609 de Mazancourt, C., Loreau, M. & Dieckmann, U. (2001) Can the evolution of plant defense lead
610 to plant-herbivore mutualism? *American Naturalist*, 109-123.
- 611 Eissenstat, D.M., Wells, C.E., Yanai, R.D. & Whitbeck, J.L. (2000) Building roots in a changing
612 environment: implications for root longevity. *New Phytologist*, 147, 33-42.
- 613 Endara, M.-J. & Coley, P.D. (2011) The resource availability hypothesis revisited: a meta-
614 analysis. *Functional Ecology*, 25, 389-398.
- 615 Eshel, I. (1983) Evolutionary and continuous stability. *Journal of Theoretical Biology*, 103, 99-
616 111.
- 617 Fagerström, T. & Westoby, M. (1997) Population dynamics in sessile organisms: some general
618 results from three seemingly different theory-lineages. *Oikos*, 80, 588-594.
- 619 Ferrière, R. & Michod, R.E. (1995) Invading wave of cooperation in a spatial iterated prisoner's
620 dilemma. *B. Proceedings of the Royal Society of London B*, 259, 77-83.
- 621 Fussmann, G.F., Loreau, M. & Abrams, P.A. (2007) Eco-evolutionary dynamics of communities
622 and ecosystems. *Functional Ecology*, 21, 465-477.
- 623 Geritz, S.A.H., Kisdi, E., Meszéna, G. & Metz, J.A.J. (1998) Evolutionarily singular strategies
624 and the adaptive growth and branching of the evolutionary tree. *Evolutionary Ecology*, 12,
625 35-57.

- 626 Gersani, M., Brown, J.S., O'Brien, E.E., Maina, G.M. & Abramsky, Z. (2001) Tragedy of the
627 commons as a result of root competition. *Journal of Ecology*, 89, 660-669.
- 628 Goodnight, C., Rauch, E., Sayama, H., De Aguiar, M.A.M., Baranger, M. & Bar-Yam, Y. (2008)
629 Evolution in spatial predator-prey models and the "Prudent predator": The inadequacy of
630 steady-state organism fitness and the concept of individual and group selection. *Complexity*,
631 13, 23-44.
- 632 Gravel, D., Guichard, F., Loreau, M. & Mouquet, N. (2010) Source and sink dynamics in meta-
633 ecosystems. *Ecology*, 91, 2172-2184.
- 634 Grime, J.P. (1977) Evidence for the existence of three primary strategies in plants and its
635 relevance to ecological and evolutionary theory. *American Naturalist*, 111, 1169-1194.
- 636 Grime, J.P. (2001) Plant strategies, vegetation processes, and ecosystem properties. Wiley, John
637 Wiley, New York, USA.
- 638 Hamilton, W.D. (1963) The evolution of altruistic behavior. *American Naturalist*, 97, 354-356
- 639 Haraguchi, Y. & Sasaki, A. (2000) The evolution of parasite virulence and transmission rate in a
640 spatially structured population. *Journal of Theoretical Biology*, 203, 85-96.
- 641 Hardin, G. (1968) Tragedy of commons. *Science*, 162, 1243-&.
- 642 Herms, D.A. & Mattson, W.J. (1992) The dilemma of plants: to grow or defend. *Quarterly*
643 *Review of Biology*, 67, 283-335.
- 644 Kerr, B., Neuhauser, C., Bohannan, B.J. & Dean, A.M. (2006) Local migration promotes
645 competitive restraint in a host-pathogen 'tragedy of the commons'. *Nature*, 442, 75-78.
- 646 Kreft, J.U. (2004) Biofilms promote altruism. *Microbiology-Sgm*, 150, 2751-2760.
- 647 Lion, S. & van Baalen, M. (2008) Self-structuring in spatial evolutionary ecology. *Ecology*
648 *Letters*, 11, 277-295.

- 649 Loeuille, N. & Leibold, M.A. (2008) Ecological consequences of evolution in plant defenses in a
650 metacommunity. *Theoretical Population Biology*, 74, 34-45.
- 651 Loeuille, N. & Loreau, M. (2004) Nutrient enrichment and food chains: can evolution buffer top-
652 down control? *Theoretical Population Biology*, 65, 285-298.
- 653 Loeuille, N., Loreau, M. & Ferrière, R. (2002) Consequences of plant-herbivore coevolution on
654 the dynamics and functioning of ecosystems. *Journal of Theoretical Biology*, 217, 369-381.
- 655 Loreau, M. (1998) Ecosystem development explained by competition within and between
656 material cycles. *Proceedings of the Royal Society of London B*, 265, 33-38.
- 657 Marsaglia, G. & Bray, T.A. (1964) A Convenient method for generating normal variables. *SIAM*
658 *Review*, 6, 260-264.
- 659 Mauricio, R. (1998) Costs of resistance to natural enemies in field populations of the annual
660 plant *Arabidopsis thaliana*. *American Naturalist*, 151, 20-28.
- 661 Menge, D.N.L., Levin, S.A. & Hedin, L.O. (2008) Evolutionary tradeoffs can select against
662 nitrogen fixation and thereby maintain nitrogen limitation. *Proceedings of the National*
663 *Academy of Sciences of the United States of America*, 105, 1573-1578.
- 664 Miller, T.E., Burns, J.H., Munguia, P., Walters, E.L., Kneitel, J.M., Richards, P.M., Mouquet, N.
665 & Buckley, H.L. (2005) A critical review of twenty years' use of the resource-ratio theory.
666 *American Naturalist*, 165, 439-448.
- 667 Nowak, M.A. (2006) Five rules for the evolution of cooperation. *Nature*, 314, 1560-1563.
- 668 Pfeiffer, T., Schuster, S. & Bonhoeffer, S. (2001) Cooperation and competition in the evolution
669 of ATP-producing pathways. *Science*, 292, 504-507.
- 670 Rankin, D.J., Bargum, K. & Kokko, H. (2007) The tragedy of the commons in evolutionary
671 biology. *Trends in Ecology & Evolution*, 22, 643-651.

- 672 Rastetter, E.B. & Ågren, G.I. (2002) Changes in individual allometry can lead to species
673 coexistence without niche separation. *Ecosystems*, 5, 789-801.
- 674 Raynaud, X. & Leadley, P.W. (2004) Soil characteristics play a key role in modeling nutrient
675 competition in plant communities. *Ecology*, 85, 2200-2214.
- 676 Reich, P.B. & Cornelissen, H. (2014) The world-wide 'fast-slow' plant economics spectrum: a
677 traits manifesto. *Journal of Ecology*, 102, 275-301.
- 678 Silla, F. & Escudero, A. (2004) Nitrogen-use efficiency: trade-offs between N productivity and
679 mean residence time at organ, plant and population levels. *Functional Ecology*, 18, 511-521.
- 680 Suzuki, S. & Kimura, H. (2011) Oscillatory dynamics in the coevolution of cooperation and
681 mobility. *Journal of theoretical biology*, 287, 42-47.
- 682 team, R.d.c. (2010) R: A language and environment for statistical computing. R Foundation for
683 Statistical Computing, Vienna, Austria.
- 684 Tilman, D. (1982) Resource competition and community structure. Princeton University Press,
685 Princeton, New Jersey.
- 686 Tilman, D. (1994) Competition and biodiversity in spatially structured habitats. *Ecology*, 75, 2-
687 16.
- 688 Tilman, D. & Wedin, D. (1991) Dynamics of nitrogen competition between successional grasses.
689 *Ecology*, 72, 1038-1049.
- 690 Wedin, D. & Tilman, D. (1993) Competition among grasses along a nitrogen gradient: initial
691 conditions and mechanisms of competition. *Ecological Monographs*, 63, 199-229.
- 692 West, S.A., Griffin, A.S. & Gardner, A. (2007) Social semantics: altruism, cooperation,
693 mutualism, strong reciprocity and group selection. *Journal of Evolutionary Biology*, 20, 415-
694 432.

695 Woodmansee, R.G., Vallis, I. & Mott, J.J. (1981) Grassland nitrogen. Terrestrial nitrogen cycles
696 (eds F.E. Clark & T. Rosswall), pp. 443-462. Ecological Bulletins, Stockholm.

697 Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-Bares,
698 J., Chapin, T., Cornelissen, J.H.C., Diemer, M., Flexas, J., Garnier, E., Groom, P.K., Gulias,
699 J., Hikosaka, K., Lamont, B.B., Lee, T., Lee, W., Lusk, C., Midgley, J.J., Navas, M.L.,
700 Niinemets, U., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V.I.,
701 Roumet, C., Thomas, S.C., Tjoelker, M.G., Veneklaas, E.J. & Villar, R. (2004) The
702 worldwide leaf economics spectrum. *Nature*, 428, 821-827.

703

704

705

706

707 **Supporting information**708 **Figure S1.** General graphical description of the spatial model709 **Figure S2.** Example of evolutionary dynamics710 **Figure S3.** Effect of the mineralization rate (m_D) and the rate of mineral nutrient loss from the
711 ecosystem (l_N) on the evolution of nutrient acquisition (s^*)712 **Figure S4.** Joint effect of the capacity for nutrient acquisition (s) and the inputs of mineral
713 nutrient (R_N) on the availability of mineral nutrient at the ecological equilibrium (N^*) for four
714 combinations of dispersal and lateral nutrient fluxes715 **Figure S5.** Examples of ecological dynamics in one patch showing temporal variations in the
716 availability of mineral nutrient and the number of plant individuals717 **Figure S6.** Ecological effects of nutrient inputs on the mean and standard deviation of the
718 equilibrium nutrient availability and the number of plant individuals in each patch719 **Figure S7.** Evolutionary effect of inputs of mineral nutrient (R_N) on the local density of plants720 **Figure S8.** Effect of the inputs of mineral nutrient (R_N) on the evolution of nutrient acquisition
721 (s^*) when only one individual is allowed to survive in each patch

722 Table 1: Model parameters.

Parameter type	Symbol	Default values	Definition
Nutrient recycling within a local patch	d_{P0}	0.275 yr ⁻¹	Initial rate of plant nutrient turnover through root and leaf turnover
	m_D	0.0766 yr ⁻¹	Mineralization of dead organic mater
	u_{N0}	1.43 dg ⁻¹ m ² yr ⁻¹	Initial uptake of mineral nutrient
Inputs of nutrient	R_D	2 dg m ⁻² yr ⁻¹	Mineral atmospheric deposition
	R_N	1.8 dg m ⁻² yr ⁻¹	Organic atmospheric deposition
	f_P	0 yr ⁻¹	Symbiotic nitrogen fixation
Losses of nutrient	l_P	$l_P, 0$ yr ⁻¹	Loss of nutrient from the ecosystem because of fires or human exportation of biomasses
	l_D	0.038 yr ⁻¹	Loss of dissolved organic matter or litter from the ecosystem
	l_N	0.05 yr ⁻¹	Nutrient leaching from soils, denitrification
Nutrient fluxes between patches	H	0 yr ⁻¹	Homogenizing lateral fluxes of nutrient
Demographic parameters	μ	0.5	Individual mortality
	γ	0.3 g	Nutrient content of a seed
	ρ	0.1	Percentage of the nutrient stock of plant individuals allocated to seed production
	σ	2 m	Standard deviation of the centred normal distribution that determines seed dispersal distances
Trade-off parameters	c	4	Costs of the investment into nutrient acquisition
	b	3.5	Benefits of the investment into nutrient acquisition

723

34

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

724

For Peer Review

Version preprint

725 Figure 1. Effect of the spatial homogenization of the mineral nutrient resource (H) on the
726 evolved capacity for nutrient acquisition (s^*) for different dispersal capacities (σ). A point is
727 missing for $\sigma=0.5$ and $R_N=0$ because these parameters lead to a non-viable population. See Table
728 1 for parameter values.

729


730 Figure 2. Effect of the inputs of mineral nutrients (R_N) on the evolved capacity for nutrient
731 acquisition (s^*), as a function of four cofactors: (A) the trade-off parameter b . (B) individual
732 mortality μ , (C) homogenization of the availability of the mineral resource H (in this case “1
733 patch” denotes non-spatial simulations with only one patch), (D) efficiency of dispersal (σ). See
734 Table 1 for parameter values.

735

736 Figure 3. Contour plots of the joint effect of the capacity for nutrient acquisition (s) and the
737 inputs of mineral nutrients into the ecosystem (R_N) on (A) the mean availability of mineral
738 nutrient (N) and (B) the mean size of the primary producer compartment (P). The thick solid
739 lines denote the position of (A) the minimum N value, (B) the maximum P value for each R_N
740 value. The dots and the dashed lines denote the evolved capacity for nutrient acquisition (s^*) for
741 each R_N value. See Table 1 for parameter values.

742

743 Fig 1




744

745

746

Peer Review

747 Fig 2




748

749

Version preprint

750 Fig 3

751



752

753

754

755

756


Fig. S1. General description of the spatial model

Figure S1-A describes the spatial setting of the model and nutrient cycling while Figure S1-B describes the scheduling and implementation of the model.

Fig. S1-A first describes (central patch of the model) the nutrient cycling model (see Table 1 and text for detailed explanations) that allows the local competition between plant individuals ($P_{j,1}, \dots, P_{j,n}$) sharing the same material cycle in a local patch (subscript j) of the ecosystem, i.e. a local patch of the model. Second, Fig. S1-A shows that the model also allows for regional competition between individuals growing in different patches, thus not sharing the same local mineral resource: individuals die, reproduce and disperse seeds between local patches. Finally, the model also mimics lateral fluxes of mineral nutrient which allows modulating the heterogeneity in nutrient availability and the way nutrients are shared between the individuals of the population.

Fig. S1-B shows that during each year the system of differential equations is integrated in each cell to simulate the growth of each plant individual in interaction with local mineral resources. Discrete demographic events happen at the end of each year and are implemented through the suppression and addition of differential equations.

Figure S1-A




view

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

Figure S1-B



Peer Review






Version preprint

Fig. S2. Example of evolutionary dynamics

A: Pairwise-invasibility plot (PIP). Each black dot corresponds to a simulation that led to the successful invasion of a resident population at its ecological equilibrium (s value on the x axis) by a mutant (s value on the y axis) starting at a very low density. In these invasion simulations, s values are not allowed to evolve. S values for the resident populations and mutants are explored systematically with a step of 0.025 so that each PIP is based on 161X161 invasion simulations. For resident strategy of nutrient uptake below s^* invasion can only be successful for mutants with higher s values than the resident s (black area above the diagonal), while the reverse is true for resident strategies above s^* . Evolution through successive replacement by new mutants therefore pushes the strategy for nutrient uptake towards s^* . Since, no strategy can invade a resident strategy at s^* (white areas above and below the diagonal at s^*) evolution stops at s^* an evolutionary type of singularity called Continuously Stable Strategy (Geritz et al. 1998).

B: Evolutionary dynamics. An example of evolutionary dynamics is displayed. It confirms the result suggested by the pairwise-invasibility plot: evolution drives the strategy for nutrient uptake towards the CSS s^* . Each dot corresponds to a different mutant. The larger and the darker a dot, the higher the biomass of the corresponding mutant.

C, B, D: changes in the size of the three ecological compartments through evolution. d_{p0} , 0.275 yr^{-1} ; m_D , 0.0766 yr^{-1} ; u_{NO} , 1.43 $\text{dg}^{-1} \text{m}^2 \text{yr}^{-1}$; R_D , 2 $\text{dg m}^{-2} \text{yr}^{-1}$; R_N , 6 $\text{dg m}^{-2} \text{yr}^{-1}$; l_P , 0 yr^{-1} ; l_D , 0.038 yr^{-1} ; l_N , 0.05 yr^{-1} ; f_P , 0 yr^{-1} ; b , 4; c , 3.

A Pairwise-invasibility plot**B Evolutionary dynamics****C Primary producer****D Dead organic matter****E Mineral nutrient**

Geritz, S. A. H., E. Kisdi, G. Meszéna, and J. A. J. Metz. 1998. Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. *Evolutionary Ecology* 12:35-57.

Comment citer ce document :


Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

Fig. S3. Effect of the mineralization rate (m_D) and the rate of mineral nutrient loss (I_N) on the evolution of nutrient acquisition (s^*).


Effect of the mineralization rate (m_D) on the evolution of nutrient acquisition (s^*) with different cofactors: (A) the trade-off parameter b . (B) individual mortality μ , (C) the intensity of homogenisation H (in this case “1 patch” denotes non-spatial simulations with only one patch), (D) the efficiency of dispersal σ .

For Peer Review


A Interaction with the trade-off




B Interaction with mortality



C Interaction with homogenization




D Interaction with dispersal




Version preprint

Effect of the rate of mineral nutrient loss (l_N) on the evolution of nutrient acquisition (s^*) with different cofactors: (A) the trade-off parameter b , (B) individual mortality μ , (C) the intensity of homogenisation H (in this case 1 patch denotes non-spatial simulations with only one patch), (D) the efficiency of dispersal σ .


A Interaction with the trade-off




B Interaction with mortality



C Interaction with homogenization



D Interaction with dispersal



Comment citer ce document :





Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

Fig. S4. Joint effect of the capacity for nutrient acquisition (s) and the inputs of mineral nutrient (R_N) on the availability of mineral nutrient at the ecological equilibrium (N) for four combinations of dispersal and lateral nutrient fluxes.

Contour plot of the joint effect of the capacity for nutrient acquisition (s) and the inputs of mineral nutrients into the ecosystem (R_N) on the availability of mineral nutrient (N) in four cases: dispersal is either low ($\sigma=0.5$) or high ($\sigma=10$), and homogenization is either low ($H=0$) or high ($H=10$). The thick solid lines denote the position of the minimum N value for each R_N value. The dots and the dashed lines denote the evolved capacity for nutrient acquisition (s^*) for each R_N value.

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494



Comment citer ce document :


Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

Fig. S5. Examples of ecological dynamics in one local patch showing temporal variations in the availability of mineral nutrient

The ecological dynamics in one patch of the 400-patch lattice is displayed: the availability of mineral nutrient (in black, N) and the number of individuals (in red, $\#$). The investment into the acquisition of mineral nutrient is not allowed to evolve ($s=0$). Results are displayed for the period 9600-9800 years after the start of the simulation so that it can be considered that a steady-state has been reached. Each panel corresponds to different conditions: A-B-C corresponds to an increase in inputs of mineral nutrients; D and E correspond to an intermediate value of nutrient inputs. For D nutrient availability is homogenized ($H=10$), while for E the capacity of seed dispersal is very low ($\sigma=0.5$).

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494



Comment citer ce document :







Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

Fig. S6. Ecological effects of nutrient inputs on the mean and standard deviation of the equilibrium nutrient availability and the number of plant individuals in each patch

Ecological effect of nutrient enrichment (increase in inputs of mineral nutrients) (1) on the mean and standard deviation (sd) of the availability of mineral nutrient (panels A, B, C), (2) on the mean and standard deviation of the number of individuals by patch (#, panels D, E, F). Means and standard deviations are computed over all the patches of the simulated lattice (20X 20 patches) at the end of 10000 years of ecological simulation. The investment into mineral nutrient acquisition is not allowed to evolve ($s=0$) and it can be considered that an ecological steady state has been reached. Different situations are modeled: (A and D) no homogenization of mineral nutrient and high seed dispersal ability; (B and E) homogenization of mineral nutrient; (C and F) low seed dispersal. Missing points for low dispersal are due to the fact that, in this case, for low nutrient inputs populations of primary producers get extinct.

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494


A, $\sigma=2, H=0$ D, $\sigma=2, H=0$ B, homogenization of nutrient, $\sigma=2, H=10$ E, homogenization of nutrient, $\sigma=2, H=10$ C, low dispersal, $\sigma=0.5, H=0$ F, low dispersal, $\sigma=0.5, H=0$ 

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

Fig. S7. Evolutionary effect of inputs of mineral nutrient (R_N) on the local density of plants

Effect of inputs of mineral nutrient (R_N) on the local crowding (mean number of individuals by local patch) with different covariables: (A) the trade-off parameter b . (B) individual mortality μ , (C) the intensity of homogenisation H , (D) the efficiency of dispersal σ . The investment into nutrient acquisition (s) is allowed to evolve and has reached an equilibrium.



Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494


For Peer Review

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494

Fig. S8. Effect of the inputs of mineral nutrient (R_N) on the evolution of nutrient acquisition (s^*) when only one individual is allowed to survive in each local patch

Effect of the inputs of mineral nutrient to the ecosystem (R_N) on the evolution of nutrient acquisition (s^*) in the case where only one individual is allowed to survive in each local patch and with different cofactors: (A) the trade-off parameter b , (B) individual mortality μ , (C) the intensity of homogenisation H , (D) the efficiency of dispersal σ .



Comment citer ce document :

For Peer Review

Comment citer ce document :

Barot, S., Bornhofen, S., Boudsocq, S., Raynaud, X., Loeuille, N. (2016). Evolution of nutrient acquisition: when space matters. *Functional Ecology*, 30 (2), 283-294. , DOI : 10.1111/1365-2435.12494