

Highlighting and Reducing the Impact of Negative Aging Stereotypes During Older Adults' Cognitive Testing

Marie Mazerolle, Isabelle Regner, François Rigalleau, Pascal Huguet

► To cite this version:

Marie Mazerolle, Isabelle Regner, François Rigalleau, Pascal Huguet. Highlighting and Reducing the Impact of Negative Aging Stereotypes During Older Adults' Cognitive Testing. Journal of visualized experiments: JoVE, 2020, 155, pp.e59922. 10.3791/59922 . hal-02458675

HAL Id: hal-02458675 https://hal.science/hal-02458675

Submitted on 28 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- 1 TITLE:
- 2 Highlighting and Reducing the Impact of Negative Aging Stereotypes During Older Adults'
- 3 Cognitive Testing
- 4

5 **AUTHORS AND AFFILIATIONS:**

- 6 Marie Mazerolle¹, Isabelle Régner², François Rigalleau³, Pascal Huguet⁴
- 7
- ¹Department and Laboratory of Psychology, MSHE, University of Burgundy Franche-Comté,
 Besançon, France
- 10 ²Aix-Marseille University, CNRS, LPC, Marseille, France
- ³Department of Psychology, University of Poitiers, CNRS, Poitiers, France
- ⁴University of Clermont Auvergne, CNRS, LAPSCO, Clermont-Ferrand, France
- 13
- 14 Corresponding author:
- 15 Isabelle Régner (isabelle.regner@univ-amu.fr)
- 16
- 17 Email addresses of co-authors:
- 18 Marie Mazerolle (marie.mazerolle@univ-fcomte.fr)
- 19 François Rigalleau (francois.rigalleau@univ-poitiers.fr)
- 20 Pascal Huguet (pascal.huguet@uca.fr)
- 21

22 **KEYWORDS:**

behavior, stereotype threat, aging, memory, neuropsychological testing, dementia, socialcognition

25

26 **SUMMARY:**

Here, we present a protocol designed to show how negative aging stereotypes can impair memory performance of older adults during cognitive testing and how to reduce this deleterious effect. This method can help older people to perform at an optimal level during testing in both lab studies and clinical settings.

31

32 ABSTRACT:

33 As life expectancy increases, aging has become a major health challenge, resulting in a huge 34 effort to better discriminate between normal and pathological cognitive decline. It is thus 35 essential that cognitive tests and their administration are as fair as possible. However, an 36 important source of bias during cognitive testing comes from negative aging stereotypes that 37 can impair the memory performance of older adults and inflate age differences on cognitive 38 tasks. The fear of confirming negative aging stereotypes creates an extra pressure among older 39 adults which interferes with their intellectual functioning and leads them to perform below 40 their true abilities. Here, we present a protocol that highlights simple but efficient interventions 41 to alleviate this age-based stereotype threat effect. The first study showed that simply 42 informing older participants about the presence of younger participants (threat condition) led 43 older adults to underperform on a standardized memory test compared with younger 44 participants, and that this performance difference was eliminated when the test was presented

as age-fair (reduced-threat condition). The second study replicated these findings on short
 cognitive tests used to screen for predementia in clinical settings and showed that teaching
 older adults about stereotype threat inoculated them against its effects. These results provide
 useful recommendations about how to improve older adults' memory assessment both in lab

- 49 studies and in clinical settings.
- 50

51 **INTRODUCTION:**

52 A growing field of laboratory research in social cognition conducted in the healthy population 53 has demonstrated that members of groups whose abilities are negatively stereotyped typically 54 underperform when the negative stereotypes are made relevant to the performance at hand, a 55 phenomenon called stereotype threat (ST). In addition to the normal anxiety associated with 56 taking cognitive tests, the fear of confirming negative stereotypes creates extra pressure that 57 may interfere with cognitive functioning and lead to perform below one's abilities^{1,2}. Many 58 findings demonstrate that negative aging stereotypes (e.g., the culturally shared beliefs that 59 aging inescapably causes severe cognitive decline and diseases such as Alzheimer's disease 60 [AD]) contribute, at least in part, to the differences classically observed in the healthy population between younger and older adults in memory tasks³⁻⁵. Without denying the impact 61 62 of aging on cognitive functioning, research clearly demonstrates that age-related stereotypes 63 are powerful enough to artificially decrease older adults' performance on memory tests.

64

65 Detrimental age-based ST effects are readily observable and fairly easy to produce with instructional manipulations⁶, such as simply emphasizing the memory component of the test⁷⁻⁹, 66 67 highlighting differences in performance between young and older adults^{10,11}, or implicitly activating negative aging stereotypes^{12,13}. Given the results obtained in laboratory studies, it is 68 69 very likely that negative aging stereotypes also permeate, at least implicitly, the standard 70 neuropsychological testing settings during the screening for pre-dementia. Indeed, because of 71 the lengthening of life expectancy, more and more people are concerned with the possibility of 72 getting AD or other forms of dementia. Importantly, false-positive errors are quite frequent in 73 the diagnosis of prodromal state of AD¹⁴, which could be explained, at least in part, by 74 transitory impaired performances in older adults due to the age-based ST phenomenon¹⁵.

75

76 For these reasons, it is important to provide efficient methods to deactivate the influence of 77 negative aging stereotypes and thereby help older people to perform at their maximum during 78 memory assessment in general and during neuropsychological testing specifically. Some 79 methods, such as de-emphasizing the memory component of the test (e.g., characterizing the 80 task as a vocabulary test), have already proved efficient to eliminate ST effects in older adults on explicit memory tests taken in the context of lab studies^{7,9,16}. However, such instructions are 81 82 not compatible with the ecological clinical context of the neuropsychological testing, in which 83 older adults come to have their memory abilities assessed. The goal of our articles is to present 84 two methods likely to alleviate age-based ST effects among older adults, either in the lab or in 85 the clinical contexts. The first one, especially suited for the lab context, consists in telling older 86 adults that performance on the ongoing memory tests usually does not differ between younger 87 and older adults (i.e., age-fair instructions). The second method, which can be implemented in 88 both lab and clinical contexts, consists in explaining to older adults (or patients) the negative

89 90 91	impact of aging stereotypes, which may help them reappraise the situation, reduce evaluative pressure, and feel less threatened during the testing.
92	PROTOCOL:
93	The present research was carried out in accordance with French standards, each participant
94	provided informed consent and procedures were consistent with the guidelines of the
95	American Psychological Association.
96	
97	1. Highlighting and reducing age-based stereotype threat on a lab memory test
98	
99	1.1. Participants screening
100	
101	1.1.1. Recruit participants in the desired age range (e.g., younger participants: 18–35 years old;
102	older participants: 60-85 years old) for a study on general mental abilities and/or the impact of
103	emotions on various cognitive tasks and questionnaires.
104	
105	1.1.2. Ensure that participants are native speakers, have normal or corrected-to-normal vision
106	and hearing, are free of neurological and psychiatric disorders, and have no current depression
107	(Geriatric Depression Scale ¹⁷) nor abnormal anxiety (State-Trait Anxiety Inventory ¹⁸).
108 109	1.1.3. Screen older participants for cognitive impairment using the Mini-Mental State
1109	Examination (MMSE) ¹⁹ . Include older adults who meet or exceed a cutoff score corresponding
111	to his/her age and educational level ²⁰ , and who live at home.
112	to his/her age and educational level , and who live at nome.
112	1.2. Conditions and experimental design
114	
115	1.2.1. Test each participant individually at home or in local community associations' premises
116	by an experimenter trained at neuropsychological testing.
117	
118	1.2.2. Run the study as a single session, but inform participants there are two separate studies.
119	
120	1.2.3. Use the "first study" as a baseline measure of participants' working memory capacity. In
121	the "first study", minimize evaluative pressure, informing participants they will perform a
122	cognitive task currently under elaboration. Encourage participants to do their very best ²¹ .
123	
124	1.2.4. Use the "second study" to evaluate the impact of stereotype threat on participants'
125	working memory capacity. That is, inform participants that they are going to perform a fully
126	validated and diagnostic test of memory capacity.
127	
128	1.2.4.1. Randomly assign participants to one of two conditions (threat versus reduced-threat
129	conditions).
130	NOTE: Soo Figure 1 for a visual depiction of the presedures full details on this research
131	NOTE: See Figure 1 for a visual depiction of the procedure; full details on this research were
132	published in Mazerolle et al. ²² .

133

134 1.2.4.2. In the "threat condition", simply tell participant that both younger and older adults areparticipating in the study.

136

1.2.4.3. In the "reduced threat condition", tell participants that both younger and older adults
are participating in the study but add that younger and older adults usually obtain the same
performances on the ongoing test.

- 140
- 141 1.3. Memory tests and procedure142
- 1.3.1. Use a validated memory test that is typically used in cognitive lab studies. Because ST
 effects typically occur on difficult tasks, use a difficult test, such as the reading span test²³.
- 145
 146 1.3.2. Use two alternative versions of the reading span test (one for the "first study" and the
 147 other one for the "second" study), with different sentences but match them in number of
 148 words and in length, frequency, and number of syllables of the last word.
- 149

152

- 150 1.3.3. Use a reading span test that contains 42 sentences: 12 series of two to five sentences151 with 3 series per length.
- 153 1.3.4. Present one sentence at a time and instruct participant to read it, at his/her own pace,154 while memorizing the last word of the sentence.
- 155
- 156 1.3.5. Ask participant not to interrupt his reading and not to pause between two sentences.
- 157
- 158 1.3.6. Have participant to recall the words immediately after he/she finished reading the last159 sentence of the block considered.
- 160
- 1.3.7. Do not require participants' recall being serial but constrain them not to start by recalling
 the final word of the last sentence read. Do not limit the recall time and invite participants to
 find as many words as possible.
- 164
- 165 1.3.8. Do not inform participants of their errors, and do not give them a second chance to 166 answer.
- 167
- 168 1.3.9. Compute the final score of each participant using the mean proportion of correctly169 recalled words across the 12 series.
- 170
- 171 2. Highlighting and reducing age-based stereotype threat on short cognitive tests used to172 screen for predementia in clinical settings
- 173
- 174 2.1. Participants screening
- 175

176 2.1.1. Recruit older participants aged 60 to 85 for a study on general mental abilities and/or the 177 impact of emotions on various cognitive tasks and questionnaires.

178

179 2.1.2. Ensure that participants live at home, have no history of significant trauma nor chronic 180 illness, have normal or corrected-to-normal vision and hearing, have no current depression 181 (Geriatric Depression Scale¹⁷) nor abnormal anxiety (State-Trait Anxiety Inventory¹⁸).

- 182
- 183 2.2. Conditions and experimental design
- 184

185 2.2.1. Test each participant individually at home or in local community associations' premises 186 by an experimenter trained at neuropsychological testing.

187

188 2.2.2. Have participants taking two short cognitive tests used to screen for predementia (or 189 mild cognitive impairment [MCI]): the MMSE¹⁹ and the Montreal Cognitive Assessment 190 (MoCA²⁴) in a counterbalanced order.

191

192 2.2.3. Randomly assign participants to one of the two order (MMSE then MoCA or MoCA then 193 MMSE).

195 2.2.4. Before taking the first test, randomly assign participants to one of two threat conditions 196 (threat versus reduced-threat conditions).

197

194

198 2.2.5. In the "threat condition", inform participants that both younger and older adults are 199 taking part in the study about memory capacity.

200

201 2.2.6. In the "reduced threat condition", inform participants that both younger and older adults 202 are taking part in the study about memory capacity but add that younger and older adults 203 typically obtain the same performances on the ongoing tests.

204

205 2.2.7. Before taking the second test, use an educational intervention to debrief all participants 206 (regardless of the threat condition they were previously assigned) about age-based stereotype 207 threat. Tell participants that "Numerous studies demonstrated that negative age-related 208 beliefs, especially about memory capacity decline, trigger anxiety, which in turn prevents older 209 adults from performing optimally. Actually, it is well-known that performance on memory tests 210 may not accurately reflect people's true abilities, because performance can be affected by the 211 testing context. In other words, people's negative reputation in an ability domain may impair 212 their performance on tests that assess this stereotyped ability, for example when older adults 213 complete a memory test. Therefore, if you feel anxious during the subsequent memory test, 214 this does not mean that you are unable to complete the tests. It rather probably means that 215 your feelings are affected by the general negative reputation of older adults regarding 216 memory."

217

218 NOTE: See Figure 2 for a visual depiction of the educational intervention; full details on this 219 research were published in Mazerolle et al.²⁵.

- 220
- 221 2.3. Short cognitive neuropsychological tests and procedure
- 222

2.3.1. Administrate the 8 subtests of the MMSE¹⁹ (orientation, registration, attention and 223 224 calculation, recall, language, copying) to evaluate memory, orientation to time and place, 225 attention and executive functioning, language and visuospatial abilities. Follow Folstein, 226 Folstein and McHugh's administration guidelines¹⁹ and compute the final score adding the scores of every subtest. Obtain a final score rated on 30-point scale. 227

228

229 2.3.2. Administrate the 8 subtests of the MoCA²⁴ (visuospatial/executive, naming, memory, 230 attention, language, abstraction, delayed recall, orientation) to evaluate memory, orientation 231 to time and place, attention and executive functioning, language and visuospatial abilities. 232 Follow administration guidelines available at www.mocatest.org and compute the final score 233 adding the scores of every subtest. Obtain a final score rated on 30-point scale.

234

235 **REPRESENTATIVE RESULTS:**

236 We addressed the hypothesis that stereotype threat impairs the working memory performance 237 of older adults and that this effect can be reduced or eliminated by a simple instruction. The 238 expected interaction between age group and threat instructions was significant, F(1, 214) =239 4.85, p < 0.03, $\eta_p^2 = 0.02$, and is depicted in **Figure 3**. In the threat condition, older participants underperformed (mean [M] = 0.73, standard error [SE] = 0.01) relative to younger participants 240 241 $(M = 0.78, SE = 0.01), F(1, 214) = 12.87, p < 0.001, \eta_p^2 = 0.06$, whereas the two age groups 242 performed equally well in the reduced-threat condition (F < 1). Moreover, older participants in 243 the reduced-threat condition obtained a better working memory score (M = 0.77, SE = 0.01) relative to those in the threat condition (M = 0.73, SE = 0.01), F(1, 214) = 9.42, p < 0.002, $\eta_p^2 =$ 244 245 0.04. The performance of younger participants did not differ across the two conditions (threat 246 condition: M = 0.78, SE = 0.01; reduced-threat condition: M = 0.78, SE = 0.01; F < 1). More details about this research can be found in Mazerolle et al.²². These findings highlight the 247 248 deleterious impact of very subtle cues in the environment (e.g., mentioning the presence of 249 younger adults in a study about memory) on older adults' performance on a laboratory test. 250 Because stereotype threat seems to be powerful enough to create significant differences 251 between younger and older adults, we went further and tested its impact on brief cognitive 252 tests classically used by primary care clinicians for the diagnosis of predementia.

253

254 We tested the impact of stereotype threat on older adults' performance on two brief cognitive 255 tests used to screen for predementia states: the MoCA and the MMSE. As shown in Figure 4 256 (left panel), during test 1 (before debriefing), participants' performance was higher in the 257 reduced-threat condition (M = 28.50; SE = 0.47) than in the threat condition (M = 26.40; SE = 0.43) on the MMSE, F(1, 76) = 12.506, p = 0.001, $\eta_p^2 = 0.141$. The same pattern occurred on 258 259 MoCA, with participants underperforming in the threat condition (M = 24.80; SE = 0.53) as 260 compared to the reduced-threat condition (M = 27.45; SE = 0.53), F(1, 76) = 10.153, p = 0.002, n_{p}^{2} = 0.118). These results show that the performances of older adults on short cognitive tests 261 262 are susceptible to stereotype threat effects.

263

264 Because these tests are used to screen for predementia state, we also examined the proportion 265 of participants meeting the clinical criteria of MCI on MoCA and MMSE (using a cutoff at 26 for 266 both tests). During test 1, stereotype threat was powerful enough on the MoCA to result in 50% 267 of our sample performing under the cutoff score of 26/30 in the threat condition, versus 15% in 268 the reduced-threat condition (p = 0.041, Fischer's exact test). The same pattern occurred on 269 MMSE, with 30% of the participants scoring below 26 in the threat condition, against 5% in 270 reduced-threat condition (p = 0.038, Fisher's exact test). More information about this study can 271 be found in Mazerolle et al.²⁵.

272

273 We also hypothesized that an educational intervention (debriefing) consisting in informing 274 older adults about the stereotype threat phenomenon and alleviating the evaluative pressure 275 of the test, would reduce the impact of negative aging stereotypes on their performance. As 276 shown in Figure 4 (right panel), after the educational intervention (debriefing), older adults 277 performed equally well on test 2 (be it MoCA or MMSE), regardless of the administration 278 condition of their previous test 1, threat or reduced-threat. In other words, lower performance 279 of participants at test 1 due to threat instructions was restored at test 2 thanks to the 280 educational intervention. More details about these findings can be found in Mazerolle et al.²⁵.

282 **FIGURE LEGENDS:**

283

286

281

Figure 1: Visual depiction of the procedure to test age-based stereotype threat on a lab memory test.

Figure 2: Visual depiction of the educational intervention to reduce age-based stereotypethreat effects.

289

Figure 3: Reading span score (adjusted for covariates) as a function of instructions condition
 and age group. Error bars indicate standard errors of the mean. This figure has been modified
 from Mazerolle et al.²².

293

297

Figure 4: MMSE and MoCA scores as a function of threat conditions before debriefing and after debriefing participants about the impact of stereotype threat. Error bars represent standard errors of the mean. This figure has been modified from Mazerolle et al.²⁵.

298 **DISCUSSION:**

299 The present studies show that stereotype threat, a neglected source of stress in many testing 300 situations, can lead older adults to perform below their true abilities on memory tests. The 301 method presented here highlights the crucial importance of the instructions given to 302 participants and patients before testing memory. Simply mentioning that younger adults are 303 taking part in the study (without mentioning any expected age-related differences in 304 performance) is sufficient to inflate by 40% (MMSE and MoCA averaged) the number of older 305 adults meeting clinical criteria for predementia on short cognitive tests. The present findings 306 also showed that presenting the memory tests as age fair or explaining to older adults the 307 negative impact of aging stereotypes on their performances are two efficient strategies to help

them resist to age-based stereotype threat. Taken together, these results highlight the critical importance of taking into account stereotype treat effects when assessing older adults' memory using short cognitive tests, especially given the current pressure on general practitioners to participate in the screening for predementia^{26,27}. It is also important for experimental studies about cognitive aging because short cognitive tests are used to exclude participants who are suspected to show predementia symptoms only because they scored below a cutoff.

315

316 Some of the critical aspects of the method deserve particular attention. A typical mistake when 317 one discovers stereotype threat research²⁸ consists in assuming that a stereotype threat 318 condition requires the implementation of an extra pressure that does not exist in conventional 319 real-life testing, and that standard real-life testing instructions can be used to operationalize a 320 no-stereotype threat control condition. Actually, it is quite the contrary: The standard real-life 321 testing settings are likely to induce stereotype threat, either implicitly or explicitly, because of 322 the words used to present the tests and/or any environmental cues that are related to negative 323 aging stereotypes (e.g., the presence of flyers or posters on Alzheimer's disease or other 324 dementia). It is much more difficult to find a way to eliminate stereotype threat in a testing 325 situation. Therefore, any modification to the instructions that have been found efficient to 326 reduce stereotype threat should be carefully considered to avoid the belief that one is 327 comparing a threat and a reduced-threat condition while, in fact, one is comparing two 328 stereotype threat conditions.

329

330 Although the present findings were obtained on lab memory tests and short cognitive tests, 331 they raise the question of whether age-based stereotype threat effects can also affect older 332 patients' performances on complete neuropsychological batteries used in memory clinics for 333 the diagnosis of MCI, the prodromal stage of AD. In our view, it seems very likely that age-based 334 stereotype threat effects contribute, at least in part, to the 53% of false positive errors 335 observed in the diagnosis of MCI. Without denying that aging may be associated with cognitive 336 decline and neurodegenerative diseases such as MCI or AD for many people, our findings 337 suggest paying special attention to the influence of negative aging stereotypes that have been 338 largely overlooked in the context of neuropsychological testing.

339

Several individual factors can make older adults more or less vulnerable to age-based stereotype threat effects. As indicated by a recent review⁵, these moderators include older adults' level of education, physical and psychological health, subjective age, stigma consciousness, domain identification (i.e., importance attached to memory), and memory selfefficacy. Future research, especially those conducted within the clinical setting where a diagnosis is requested, should thus consider some of these factors to provide fairer evaluation of patients' memory abilities.

347

348 Our experimental design offers new recommendations to healthcare professionals to improve 349 the accuracy of predementia diagnosis. Characterizing the memory tests as age fair or 350 informing older adults about the age-based stereotype threat phenomenon are two simple and 351 easily implementable instructions likely to help older people to perform at an optimal level

- 352 during neuropsychological testing. These recommendations may help medical staff to deliver
- 353 more accurate and potentially less threatening information to the patients and/or their family
- 354 (thereby significantly improving their well-being and quality of life). It should be noted,
- 355 however, that these recommendations may need to be adapted to consider other types of
- 356 threat that could be simultaneously triggered by the clinical context (e.g., the specific threat of
- AD as a dramatic disease²⁹, the threat of being at hospital³⁰, and the threat of the white $coat^{31}$).
- 358

359 **ACKNOWLEDGMENTS**:

- Part of this work was supported by Plan Alzheimer Foundation on a Humanities and Social
 Sciences grant (AAP SHS 2013: "Sociocognitive aspects of Alzheimer disease" to F. Rigalleau and
 M. Mazerolle).
- 363

364 **DISCLOSURES:**

- 365 The authors have nothing to disclose.
- 366

367 **REFERENCES**:

- 368 1. Steele, C. M., Aronson, J. Stereotype threat and the intellectual test performance of African
 369 Americans. *Journal of Personality and Social Psychology*. 69 (5), 797–811 (1995).
- 2. Steele, C. M. A threat in the air: How stereotypes shape intellectual identity and performance. *American Psychologist.* **52**, 613–629 (1997).
- 372 3. Barber, S. J. An examination of age-based stereotype threat about cognitive decline: 373 Implications for stereotype-threat research and theory development. *Perspectives on* 374 *Psychological Science.* **12** (1), 62–90 (2017).
- 4. Chasteen, A. L., Kang, S. K., Remedios, J. D. Aging and stereotype threat: Development, process, and interventions. In *Stereotype threat: Theory, process, and application*. Edited by Inzlicht M., Schmader T., 202–216, Oxford University Press. New York, NY (2012).
- 378 5. Lamont, R. A, Swift, H. J., Abrams, D. A. Review and Meta-Analysis of Age-Based Stereotype
- Threat: Negative Stereotypes, Not Facts, Do the Damage. *Psychology and Aging.* **30** (1), 180–193 (2015).
- 381 6. Haslam, C. et al. "When the age is in, the wit is out": Age related self-categorization and
- deficit expectations reduce performance on clinical tests used in dementia assessment.
 Psychology and Aging. 27, 778–784 (2012).
- 384 7. Desrichard, O., Köpetz, C. A threat in the elder: The impact of task-instructions, self-efficacy 385 and performance expectations on memory performance in the elderly. *European Journal of* 386 Castel Deschalase, **25**, 552 (2005)
- 386 Social Psychology. **35**, 537–552 (2005).
- 387 8. Kang, S. K., Chasteen, A. L. The moderating role of age-group identification and perceived
 388 threat on stereotype threat among older adults. *The International Journal of Aging and Human*
- 389 Development. 69, 201–220 (2009).
- 390 9. Rahhal, T. A., Hasher, L., Colcombe, S. J. Instructional manipulations and age differences in 391 memory: Now you see them, now you don't. *Psychology and Aging*. **16**, 697–706 (2001).
- memory: Now you see them, now you don't. *Psychology and Aging*. 16, 697–706 (2001).
 10. Hess, T. M., Auman, C., Colcombe, S. J., Rahhal, T. A. The Impact of Stereotype Threat on
- 393 Age Differences in Memory Performance. The Journals of Gerontology Series B: Psychological
- 394 *Sciences and Social Sciences*. **58** (1), P3–P11 (2003).

- 11. Hess, T. M., Hinson, J. T. Age-related variation in the influences of aging stereotypes on
 memory in adulthood. *Psychology and Aging*. 21 (3), 621–625 (2006).
- Levy, B. Improving memory in old age through implicit self-stereotyping. *Journal of Personality and Social Psychology*. **71**, 1092–1107 (1996).
- 399 13. Stein, R., Blanchard-Fields, F., Hertzog, C. The effects of age-stereotype priming on the 400 memory performance of older adults. *Experimental Aging Research*. **28** (2), 169–181 (2002).
- 401 14. Sachdev, P. S. et al.; Sydney Memory, Ageing Study Team. Factors predicting reversion from
- 402 mild cognitive impairment to normal cognitive functioning: a population-based study. *PloS One*.
 403 **8**, e59649 (2013).
- 404 15. Régner, I. et al. Aging stereotypes must be taken into account for the diagnosis of 405 prodromal and early Alzheimer's disease. *Alzheimer Disease & Associated Disorders - An* 406 *International Journal.* **30** (1), 77–79 (2016).
- 407 16. Chasteen, A. L., Bhattacharyya, S., Horhota, M., Tam, R., Hasher, L. How feelings of
 408 stereotype threat influence older adults' memory performance. *Experimental Aging Research*.
 409 **31** (3), 235–260 (2005).
- 410 17. Yesavage, J. A. et al. Development and validation of a geriatric depression screening scale: a 411 preliminary report. *Journal of Psychiatric Research*. **17** (1), 37–49 (1982).
- 412 18. Spielberger, C. D. *Test Anxiety Inventory*. The Corsini Encyclopedia of Psychology. John Wiley
 413 & Sons, Inc. Hoboken, NJ (2010).
- 414 19. Folstein, M. F., Folstein, S. E., McHugh, P. R. "Mini-mental state." *Journal of Psychiatric* 415 *Research.* **12** (3), 189–198 (1975).
- 416 20. Crum, R. M., Anthony, J. C., Bassett, S. S., Folstein, M. F. Population-based norms for the
- 417 Mini-Mental State Examination by age and educational level. *The Journal of the American*
- 418 *Medical Association*. **269** (18), 2386–2391 (1993).
- 419 21. Gimmig, D., Huguet, P., Caverni, J-P., Cury, F. Choking under pressure and working-memory
 420 capacity: When performance pressure reduces fluid intelligence. *Psychonomic Bulletin &*421 *Review.* 13, 1005-1010 (2006).
- 422 22. Mazerolle, M., Régner, I., Morisset, P., Rigalleau, F., Huguet, P. Stereotype Threat 423 Strengthens Automatic Recall and Undermines Controlled Processes in the Elderly.
- 424 *Psychological Science*. **23**, 723-727 (2012).
- 425 23. Daneman, M., Carpenter, P. A. Individual differences in working memory and reading.
 426 Journal of Verbal Learning and Verbal Behavior. 19, 450–466 (1980).
- 427 24. Nasreddine, Z. S. et al. The Montreal Cognitive Assessment, MoCA: A brief screening tool for
- 428 mild cognitive impairment. *Journal of the American Geriatrics Society*. **53**, 695–699 (2005).
- 429 25. Mazerolle, M. et al. Negative Aging Stereotypes Impair Performance on Brief Cognitive Tests
- 430 Used to Screen for Predementia. *Journals of Gerontology, Series B: Psychological Sciences and*
- 431 Social Sciences. **72** (6), 932–936 (2017).
- 432 26. Brown, J. The use and misuse of short cognitive tests in the diagnosis of dementia. *Journal*
- 433 of Neurology, Neurosurgery, and Psychiatry. **86**, 680–685 (2015).
- 434 27. Le Couteur, D. G., Doust, J., Creasey, H., Brayne, C. Political drive to screen for pre-
- 435 dementia: not evidence based and ignores the harms of diagnosis. *BMJ*. **347**, f5125 (2013).
- 436 28. Steele, C.M., Davies, P.G. Stereotype Threat and Employment Testing: A Commentary.
- 437 *Human Performance.* **16** (3), 311–326 (2003).

- 438 29. Scholl, J. M., Sabat, S. R. Stereotypes, stereotype threat and ageing: implications for the
- understanding and treatment of people with Alzheimer's disease. *Ageing & Society.* 28 (1),
 103–130 (2008).
- 441 30. Suhr, J. A., Gunstad, J. "Diagnosis threat": The effect of negative expectations on cognitive
- 442 performance in head injury. Journal of Clinical and Experimental Neuropsychology. 24 (4),
- 443 448–457 (2002).
- 444 31. Schlemmer, M., Desrichard, O. Is Medical Environment Detrimental to Memory? A Test of A
- 445 White Coat Effect on Older People's Memory Performance. *Clinical Gerontologist.* **41** (1), 77–81
- 446 (2018).