



# Impact of the Variability in Hydraulic Parameters of the Vadose Zone on Pollutant Plume Development

Léa Pannecoucke, Mathieu Le Coz, Clémence Houzé, Chantal de Fouquet, D C | Washington

## ► To cite this version:

Léa Pannecoucke, Mathieu Le Coz, Clémence Houzé, Chantal de Fouquet, D C | Washington. Impact of the Variability in Hydraulic Parameters of the Vadose Zone on Pollutant Plume Development. AMERICAN GEOPHYSICAL UNION (AGU) fall meeting, AGU, Dec 2018, WASHINGTON, United States. 2018. hal-02460185

HAL Id: hal-02460185

<https://hal.science/hal-02460185>

Submitted on 29 Jan 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

Copyright

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## INTRODUCTION AND CONTEXT

Simulations of hydraulic parameter fields, spatially variable, and radionuclide plumes have been generated to assess:

- the influence of small scale variability in hydraulic parameters on solute transport;
- the parameter that should be the most precisely characterized when studying a polluted site.




### (I) PARAMETER FIELDS

3 parameters of the Mualem-van Genuchten formalism:  $K_s$ ,  $\alpha$  and  $n$ , represented by random fields (lognormal distribution) characterized by  $\sigma^2$  (variance),  $a_x$  and  $a_z$  (ranges).

### (II) PLUME SIMULATIONS

Simulation of flow and solute transport in variably saturated porous media from a localized source of tritiated water, using random fields as inputs for hydraulic parameters.

Examples of random fields and the corresponding simulated tritium plumes.



The + correspond to centers of mass of all plumes simulated from the same  $\{\sigma^2, a_x \text{ and } a_z\}$  triplet.

Léa PANNECOUCKE<sup>1</sup>, Mathieu LE COZ<sup>2</sup>, Clémence Houzé<sup>3</sup>, Chantal de FOUQUET<sup>1</sup>

<sup>1</sup> MINES ParisTech, PSL University, Centre de géosciences, 35 rue St Honoré, 77300 Fontainebleau, France

<sup>2</sup> IRSN, PSE-ENV/SEDRE, 31 avenue de la Division Leclerc, 92260 Fontenay-aux-Roses, France


<sup>3</sup> GEOPS, Univ. Paris-Sud, CNRS, Université Paris-Saclay, 91405 Orsay, France




### (III) SENSITIVITY ANALYSES

For each triplet  $\{\sigma^2, a_x \text{ and } a_z\}$ , sets of 100 random fields are generated and the corresponding plumes are simulated and characterized (in the vadose zone) thanks to:

- their surface (average and standard deviation for each set);
- the position of their center of mass (average distance between the centers of mass of the set and the center of mass of the plume simulated in a homogeneous medium).



- Variances of hydraulic parameters strongly impact the simulated tritium plumes: when  $\sigma^2$  increases, plumes are more expanded and more spread in space;
- **$n$  is the most critical parameter**: mean variances found at field scale (marked by ■, □ and ▨ on the X-axis) result in an increase of 65% of plume size compared to the homogeneous medium for  $n$  and 20 to 25% for  $K_s$  and  $\alpha$ .



- Parameter ranges have a minor impact on simulated plumes (especially  $a_x$ ) at least for the values tested.

This work is supported by the French National Radioactive Waste Management Agency (ANDRA) under the "Investments for the Future" Program.

