



HAL
open science

Decidability of the isomorphism and the factorization between minimal substitution subshifts

Fabien Durand, Julien Leroy

► **To cite this version:**

Fabien Durand, Julien Leroy. Decidability of the isomorphism and the factorization between minimal substitution subshifts. 2020. hal-02461469v1

HAL Id: hal-02461469

<https://hal.science/hal-02461469v1>

Preprint submitted on 30 Jan 2020 (v1), last revised 11 Aug 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Decidability of the isomorphism and the factorization between minimal substitution subshifts

Fabien Durand* and Julien Leroy†

*Laboratoire Amiénois de Mathématiques Fondamentales et Appliquées, CNRS-UMR 7352, Université de Picardie Jules Verne, 33 rue Saint Leu, 80000 Amiens, France.

`fabien.durand@u-picardie.fr`

†Département de Mathématique, Université de Liège, 12 Allée de la découverte, 4000 Liège, Belgique.

`j.leroy@uliege.be`

January 23, 2020

Abstract

Classification is a central problem for dynamical systems, in particular for families that arise in a wide range of topics, like substitution subshifts. It is important to be able to distinguish whether two such subshifts are isomorphic, but the existing invariants are not sufficient for this purpose. We first show that given two minimal substitution subshifts, there exists a computable constant R such that any factor map between these subshifts (if any) is the composition of a factor map with a radius smaller than R and some power of the shift map. Then we prove that it is decidable to check whether a given sliding block code is a factor map between two prescribed minimal substitution subshifts. As a consequence of these two results, we provide an algorithm that, given two minimal substitution subshifts, decides whether one is a factor of the other and, as a straightforward corollary, whether they are isomorphic.

Contents

1	Introduction	3
2	Basic materials	6
2.1	Words and sequences	6
2.2	Subshifts, minimality, factors, local and induced maps	7

2.3	Morphic and substitutive sequences	8
2.4	Examples	9
3	From uniformly recurrent morphic sequences to purely substitutive sequences w.r.t. proper primitive substitutions	10
3.1	Return words and derived sequences	11
3.2	Proof of Theorem 3.1	14
3.3	Examples	16
4	Bounds for primitive substitutive sequences	17
4.1	Recognizability	17
4.2	Block representations of subshifts and substitutions on the blocks of radius n	20
4.3	Computable bounds for linear recurrence of substitutive sequences	22
5	Bounding the radius of a factor map between primitive substitution subshifts	25
5.1	Orbit-preserving maps, cocycles and dill maps	25
5.2	Specific dill maps associated with factor maps	27
5.3	Factorizations with smaller radius using dill maps	30
6	Decidability of the factorization in the case of periodic subshifts	36
7	Decidability of the factorization	37
7.1	Return words to a set of words	38
7.2	Some sufficient conditions for $f : A \rightarrow B$ to be a factor map . . .	43
7.3	The sufficient conditions of Theorem 7.15 are necessary and algorithmically checkable	44
7.4	Proof of Theorem 1.1 and Corollary 1.2	51
8	The constant length case	51
8.1	Factor maps between constant length primitive substitution subshifts	52
8.2	Decidability of the factorization	56
8.3	How to get rid of the injective substitution assumption	57
8.4	Listing of the factors	58
9	Factor maps between linearly recurrent subshifts	58
10	Other decidability problems for morphic subshifts	61
10.1	Factor maps for subshifts generated by automatic sequences . . .	61
10.2	The non minimal case	61
10.3	Morphic languages	61
10.4	Equality problem of morphic sequences	62
10.5	The listing of the factors	62
10.6	Orbit equivalence	63
10.7	Beyond substitution and linearly recurrent subshifts	64

1 Introduction

Classification is a central problem in the study of dynamical systems, in particular for families of systems that arise in a wide range of topics. Let us mention subshifts of finite type that appear, for example, in information theory, hyperbolic dynamics, C^* -algebra, statistical mechanics and thermodynamic formalism [LM95, Bow75]. The most important and longstanding open problem for this family originates in [Wil73] and is stated in [Boy08] as follows : *Classify subshifts of finite type up to topological isomorphism. In particular, give a procedure which decides when two non-negative integer matrices define topologically conjugate subshifts of finite type.*

The existence of an isomorphism between two given subshifts of finite type is known to be equivalent to the *Strong Shift Equivalence* for matrices over \mathbb{Z}^+ [Wil73], which is not known to be decidable [KR99]. In contrast, the *Shift Equivalence* for matrices over \mathbb{Z}^+ is decidable and provides an invariant of isomorphism for subshifts of finite type.

An other well-known family of subshifts, that is also defined through matrices, with a wide range of interests is the family of substitution subshifts. These subshifts are concerned, for example, with automata theory, first order logic, combinatorics on words, quasicrystallography, fractal geometry, group theory and number theory [AB07, AS03, Fog02, Nek18, Rig14]. In this paper we show that not only the existence of isomorphism between such subshifts is decidable but also the factorization. This answers a question asked in [Dur13b].

Theorem 1.1. *Let (X, S) and (Y, S) be two subshifts generated by uniformly recurrent morphic sequences. It is decidable whether or not there is a factor map $f : (X, S) \rightarrow (Y, S)$.*

Using the fact that minimal substitution subshifts are coalescent [Dur00] one deduces the decidability of the isomorphism problem.

Corollary 1.2. *It is decidable whether or not two subshifts generated by uniformly recurrent morphic sequences are isomorphic.*

Our approach gives a precise and detailed description of the isomorphisms and factor maps. It is thus also concerned with the study of automorphism groups of topological dynamical systems and its recent developments. This was a classical topic in the 70's and 80's [Hed69, Cov72, BLR88] and it recently got a renewal of interest subshifts with low complexity. Cyr and Kra [CK16] obtained that for minimal subshifts (X, S) of sub-quadratic complexity, the quotient $\text{Aut}(X, S)/\langle S \rangle$ is periodic, where $\text{Aut}(X, S)/\langle S \rangle$ stands for the automorphism group of (X, S) and $\langle S \rangle$ the group generated by the shift map. Whereas in [?] and [DDMP16] it was shown that automorphism groups of subshifts with sub-affine complexity along a subsequence (which includes minimal substitution

subshifts) are virtually \mathbb{Z} . In other words, there exists a finite set of automorphisms F such that any automorphism is the composition of an element of F with a power of the shift.

When studying factor maps instead of automorphisms we lose the group structure. The main feature used in the articles cited above is that automorphisms can be iterated, which is not the case for factor maps, and that automorphisms preserve some key features of the dynamics. In [DDMP16], the authors used the concept of asymptotic pairs. Unfortunately, images and preimages of asymptotic pairs via factor maps is neither well controlled nor understood.

Nevertheless, for specific families and with different approaches, similar results can be obtained: Theorem 1.3 for minimal substitution subshifts and Theorem 1.5 for linearly recurrent subshifts.

Durand [Dur00] proved that for linearly recurrent subshifts there are finitely many subshift factors up to isomorphism. Salo and Törmä [ST15] gave a more precise result for factor maps between two minimal substitution subshifts of constant length: there exists a bound R such that any factor map is the composition of a sliding block code of radius R with a power of the shift map. A similar result has also been obtained in [CDKL14] (see also [CKL08]). Salo and Törmä use a renormalization process allowing, given a factor map with some radius, to obtain another one with a smaller radius. It has also been used in [CQY16] to study the automorphism groups of constant length substitution subshifts.

We extend the result of Salo and Törmä to the whole class of aperiodic minimal morphic subshifts.

Theorem 1.3. *Let (X, S) and (Y, S) be two subshifts generated by uniformly recurrent morphic sequences. Suppose (Y, S) is aperiodic. Then, there exists a computable constant R such that any factor map from (X, S) to (Y, S) is the composition of a power of the shift map S with a factor map of radius less than R .*

This drastically extends the results obtained in the papers mentioned above that were considering the restrictive case of constant length substitutions. Theorem 1.3 is one of the two main ingredients of the proof of Theorem 1.1. The other one is the following.

Theorem 1.4. *Let (X, S) and (Y, S) be two minimal morphic subshifts and ϕ be a sliding block code. It is decidable to know whether ϕ defines a factor map from (X, S) to (Y, S) .*

Its proof relies on a different method. We consider constructions based on return words to clopen sets and use the fact that minimal substitution subshifts have finitely many induced systems on cylinder sets up to conjugacy [Dur98a, HZ99]. In the context of constant length substitutions (or more generally, for subshifts generated by automatic sequences), this theorem has been obtained 20 years ago by I. Fagnot [Fag97b] using the first order logic framework of Presburger arithmetic, without assuming minimality.

Let us mention an interesting problem tackled in [CDK17] for minimal constant length substitution subshifts that naturally follows the discussion above.

From [Dur00, Dur13b], minimal morphic subshifts have finitely many aperiodic subshift factors. Thus, it is natural to try to produce (to compute) the finite list of all such factors of a given minimal substitutive subshift (X, S) (up to isomorphism). A strategy would be to consider one by one factor maps of increasing radius. For each new factor map, a factor (Y, S) is obtained. One can compare it to the previously obtained subshifts using Corollary 1.2. It is tempting to say in view of Theorem 1.3 that there are finitely many factor maps to test, those of radius less than R . Unfortunately in our proof this constant strongly depends on the factor (Y, S) , in fact on the constant of recognizability of the substitution defining it. We leave it as an open problem. Nevertheless, if we ask both (X, S) and (Y, S) to be constant length substitution subshifts then this strategy is successful and the list of factors can be given (Section 8.4) as shown in [CDK17].

Organization of the paper

In Section 2 are defined the different classical notions and constructions, and some of their properties, that will be used throughout this paper.

In Section 3 we prove (Theorem 3.1) that minimal morphic subshifts are isomorphic to primitive substitution subshifts. In fact this was already known as it is a straightforward consequence of a combination of results from [DHS99] and [Dur13b]. We provide a proof as we need to precise the radius of the isomorphisms involved. This is used in many proofs of this paper as it substantially simplifies the assumptions.

Dealing with decision problems we need along this paper computable bounds of key features of the substitution subshifts. This is done in Section 4 for the recognizability, the linear recurrence, the return words, the growth of the substitutions, the complexity functions and the ergodic measures. We also recall the classical construction of substitutions on the blocks of length n [Que10].

Section 5 is devoted to the proof of Theorem 1.3. We introduce a crucial concept for this paper. This is what Salo and Törmä [ST15] called dill maps. These dill maps are defined by a renormalization process on factor maps. Salo and Törmä showed that, starting with a factor map between two constant length substitution subshifts, this renormalization process yields to another factor map between the same subshifts and with a smaller radius. This is no longer true for non-constant length substitutions. Nevertheless this produces dill maps for which the notion of radius can also be defined. Furthermore, the renormalization of a dill map provides another dill map and the iteration of the process lead to dill maps with radius smaller than some computable constant R . Using the pigeon hole principle, we obtain twice the same dill map in this iteration process and it happens that these two dill maps are in fact factor maps, thus with a small computable radius. It finally turns out that that these factor maps are equal to the original factor map up to a power of the shift map.

When the factors are periodic the decision problems are easier to deal with. This is done in Section 6.

Theorem 1.1, Corollary 1.2 and Theorem 1.4 are shown in Section 7. We

first prove these results for minimal substitution subshifts and then for minimal morphic subshifts using Theorem 3.1. We use Theorem 1.3 and constructions based on return words to sets of words.

In Section 8 we consider as a particular example the family of constant length substitution subshifts. For this context, we precise Theorem 1.3 showing we have a much better control on the radius of the factor maps. This improves one of the main results in [CDK17]. We also give a different proof of Theorem 1.4 using first order logic of the Presburger arithmetic. This provides a new proof of Theorem 1.1 and more efficient algorithms for this framework.

In Section 9 we consider the wider family of linearly recurrent subshifts. We obtain the following analogue of theorem 1.3 but without any control on the radius of the factor maps.

Theorem 1.5. *Let (X, S) and (Y, S) be aperiodic subshifts. Suppose (X, S) is linearly recurrent for the constant K . Then, there exist $N \leq (2(K+1)^4)^{6(K+1)^6}$ and some factor maps $f_i : (X, S) \rightarrow (Y, S)$, $1 \leq i \leq N$ such that if $f : (X, S) \rightarrow (Y, S)$ is a factor map then there exist $k \in \mathbb{Z}$ and $i \in \{1, \dots, N\}$ such that $f = S^k \circ f_i$.*

In Section 10 we discuss some open questions.

Acknowledgements

The authors would like to thank Samuel Petite for many interesting discussions. This research is supported by the ANR project¹ ‘‘Dyna3S’’ as well as by a Hubert Curien Partnership² (PHC) ‘‘Tournesol’’.

2 Basic materials

2.1 Words and sequences

An *alphabet* A is a set of elements called *letters*. Unless explicitly stated, we consider that the alphabets are finite. A *word* over A is an element of the free monoid generated by A , denoted by A^* . Let $u = u_0u_1 \cdots u_{n-1}$ (with $u_i \in A$, $0 \leq i \leq n-1$) be a word, its *length* is n and is denoted by $|u|$. The *empty word* is denoted by ϵ , $|\epsilon| = 0$. The set of non-empty words over A is denoted by A^+ . A *subword* of u is a finite word y such that there exist two words v and w satisfying $u = v y w$. When v (resp. w) is the empty word, we say that y is a *prefix* (resp. *suffix*) of u . If k, l and integers such that $0 \leq k \leq l < |u|$, we let $u_{[k,l]} = u_{[k,l+1[}$ denote the subword $u_k u_{k+1} \cdots u_l$ of u . If $l < k$, then $u_{[k,l]}$ is the empty word. If y is a subword of u , the *occurrences* of y in u are the integers i such that $u_{[i,i+|y|-1]} = y$. If y has an occurrence in u , we also say that y *occurs* in u .

¹Ref. ANR-13-BS02-0003

²Ref. SCO/AD/FR/SOR/2015/2019880

In this article \mathbb{K} will stand for \mathbb{N} or \mathbb{Z} . The elements of $A^{\mathbb{K}}$ are called *sequences*. Sometimes, when we want to emphasize the type of sequences we are dealing with we say *one-sided* sequences for elements of $A^{\mathbb{N}}$ and *two-sided* sequences for elements of $A^{\mathbb{Z}}$. For $x = (x_n)_{n \in \mathbb{Z}} \in A^{\mathbb{Z}}$, we let x^+ and x^- respectively denote the sequences $(x_n)_{n \geq 0}$ and $(x_n)_{n < 0}$. The notion of subword is naturally extended to sequences, as well as the notion of prefixes when $\mathbb{K} = \mathbb{N}$. The set of subwords of length n of x is written $\mathcal{L}_n(x)$ and the set of subwords of x , or the *language* of x , is denoted by $\mathcal{L}(x)$. We let $p_x : \mathbb{N} \rightarrow \mathbb{N}, n \mapsto \#\mathcal{L}_n(x)$, denote the *word complexity (function)* of x .

The sequence $x \in A^{\mathbb{N}}$ is *ultimately periodic* if there exist a word u and a non-empty word v such that $x = uv^\omega$, where $v^\omega = vvv \dots$. It is *periodic* if u is the empty word. A sequence that is not ultimately periodic is called *aperiodic*. A word u is *recurrent* in x if it occurs in x infinitely many times. A sequence x is *recurrent* if every subword u of x is recurrent in x^+ and in x^- (whenever x is two-sided). It is *uniformly recurrent* if it is recurrent and for any subword u of x , the greatest difference of two successive occurrences of u in x is bounded. It is *linearly recurrent* for the constant K if it is recurrent and if this greatest difference is bounded by $K|u|$.

2.2 Subshifts, minimality, factors, local and induced maps

A (*topological*) *dynamical system* is a couple (X, S) where S is a continuous map from X to X and X is a S -invariant compact metric space, **that is $S^{-1}X = X$** . It is *aperiodic* if there does not exist *S -periodic points*, that is points x such that $S^n x = x$ for some $n \neq 0$.

It is a *subshift* when X is a closed S -invariant subset of $A^{\mathbb{Z}}$ (for the usual topology on $A^{\mathbb{Z}}$) and S is the shift map ($S : A^{\mathbb{Z}} \rightarrow A^{\mathbb{Z}}, (x_n)_{n \in \mathbb{Z}} \mapsto (x_{n+1})_{n \in \mathbb{Z}}$). In the sequel even if the alphabet changes, S will always denote the shift map and we usually say that X itself is a subshift. The *language* $\mathcal{L}(X)$ of a subshift (X, S) is the union of the languages of x for $x \in X$. The word complexity function naturally extends to subshifts. We denote it p_X .

We say that a dynamical system (X, S) is *minimal* whenever the only closed S -invariant subsets of X are X and \emptyset .

Let x be a sequence belonging to $A^{\mathbb{Z}}$ or $A^{\mathbb{N}}$. Consider the set $\Omega(x) = \{y \in A^{\mathbb{Z}} \mid \mathcal{L}(y) \subset \mathcal{L}(x)\}$. Then, $(\Omega(x), S)$ is a subshift. We call it the *subshift generated (or defined)* by x .

If x is two-sided or recurrent, the subshift $(\Omega(x), S)$ is minimal if and only if x is uniformly recurrent. In this case all sequences y in $\Omega(x)$ are uniformly recurrent and fulfill $\Omega(y) = \Omega(x)$, $\mathcal{L}(x) = \mathcal{L}(y)$ and $\Omega(x) = \overline{\{S^n y \mid n \in \mathbb{Z}\}}$.

A subshift (X, S) is *linearly recurrent* if it is a subshift generated by a linearly recurrent sequence x . If K is a constant of linear recurrence of x , then all sequences $y \in X$ are linearly recurrent for the constant K .

Let (X, S) be a subshift on the alphabet A , D be some set endowed with the discrete topology and $g : X \rightarrow D$ be a continuous map. Then g being locally constant, there exists a map $\hat{g} : A^{t+s+1} \rightarrow D$, called a *local map* defining g , such that $g(x) = \hat{g}(x_{[-t, s]})$ for all $x \in X$. We say that \hat{g} defines g , and, t is called

the *memory* and s the *anticipation* of \hat{g} (and g). When $s = t$ we call $r = s$ the *radius* of \hat{g} and a radius of g (there exist other local maps defining f). The *radius* of g is the smallest one among the set of radius of local maps defining g .

Let (X, T) and (Y, R) be two dynamical systems. We say that (Y, R) is a *factor* of (X, T) if there is a continuous and onto map $f : X \rightarrow Y$ such that $f \circ T = R \circ f$. We call f a *factor map*. If, moreover, f is one-to-one we say f is an *isomorphism*.

Suppose $f : (X, S) \rightarrow (Y, S)$ is a factor map between subshifts, where $X \subset A^{\mathbb{Z}}$ and $Y \subset B^{\mathbb{Z}}$. Then, considering the map $g : X \rightarrow B$ given by $g(x) = f(x)_0$ there is a local map $\hat{g} : A^{t+s+1} \rightarrow B$ defining g and satisfying $f(x)_n = \hat{g}(x_{n-t} \dots x_n \dots x_{n+s})$ for any $n \in \mathbb{Z}$ and $x \in X$. This is the Curtis-Hedlund-Lyndon Theorem (see [LM95]). The map \hat{g} is called a *sliding block code* defining f and we extend the notions of memory, anticipation and radius of the map g to the map f . The sliding block code \hat{g} naturally extends to the set of words of length at least $t + s + 1$, and we denote this map also by \hat{g} .

2.3 Morphic and substitutive sequences

Let A and B be finite or infinite alphabets. By a *morphism* from A^* to B^* we mean a homomorphism of free monoids. Let σ be a morphism from A^* to B^* . When $\sigma(A) = B$, we say σ is a *coding*. Thus, codings are onto. We set $|\sigma| = \max_{a \in A} |\sigma(a)|$ and $\langle \sigma \rangle = \min_{a \in A} |\sigma(a)|$. The morphism σ is of *constant length* if $\langle \sigma \rangle = |\sigma|$. We say σ is *erasing* if there exists $b \in A$ such that $\sigma(b)$ is the empty word. If σ is non-erasing, it induces by concatenation a map from $A^{\mathbb{N}}$ to $B^{\mathbb{N}}$ and a map from $A^{\mathbb{Z}}$ to $B^{\mathbb{Z}}$. These maps are also denoted by σ . The language of the endomorphism $\sigma : A^* \rightarrow A^*$ is the set $\mathcal{L}(\sigma)$ of words occurring in some $\sigma^n(a)$, $a \in A$, $n \in \mathbb{N}$. When $\mathcal{L}(\sigma)$ is infinite, we let X_σ denote the set $\{y \in A^{\mathbb{Z}} \mid \mathcal{L}(y) \subset \mathcal{L}(\sigma)\}$. It is closed and S -invariant. We say (X_σ, S) is the subshift generated by σ .

In the sequel we use the definition of substitution that is in [Que10] and the notion of substitutive sequence defined in [Dur98a]. Both are restrictive in general but not in the uniformly recurrent case as shown in [Dur13a]. For non restrictive context we will speak of (prolongable) endomorphisms and (purely) morphic sequences.

Let $\sigma : A^* \rightarrow A^*$ be an endomorphism. If there exist a letter $a \in A$ and a non empty word u such that $\sigma(a) = au$ and moreover, if $\lim_{n \rightarrow +\infty} |\sigma^n(a)| = +\infty$, then σ is said to be *right-prolongable on a* . It is a *substitution* whenever it is right-prolongable on some letter $a \in A$ and *growing* (that is, $\lim_n \langle \sigma^n \rangle = +\infty$). Then, (X_σ, S) is called the *substitution subshift* generated by σ . Suppose that σ is right-prolongable on $a \in A$. Since for all $n \in \mathbb{N}$, $\sigma^n(a)$ is a prefix of $\sigma^{n+1}(a)$ and because $|\sigma^n(a)|$ tends to infinity with n , the sequence $(\sigma^n(aaa \dots))_{n \geq 0}$ converges (for the usual product topology on $A^{\mathbb{N}}$) to a sequence denoted by $\sigma^\omega(a)$ which is a fixed point of σ : $\sigma(\sigma^\omega(a)) = \sigma^\omega(a)$. A sequence obtained in this way by iterating a prolongable endomorphism is said to be *purely morphic* (w.r.t. σ) or *purely substitutive* when σ is a substitution. If $x \in A^{\mathbb{N}}$ is purely morphic and $\phi : A^* \rightarrow B^*$ is a morphism such that $\phi(x)$ is a sequence, then

$y = \phi(x)$ is said to be a *morphic sequence* (w.r.t. (σ, ϕ)) and the subshift it generates is called a *morphic subshift*. When ϕ is a coding and σ a substitution, we say y is *substitutive* (w.r.t. (σ, ϕ)) and the subshift it generates is called *substitutive subshift*.

Two-sided fixed points of σ can be similarly defined: When σ is right-prolongable on $a \in A$ and *left-prolongable* on $b \in A$ (that is $\sigma(b) = vb$ for some non-empty word v and $\lim_{n \rightarrow +\infty} |\sigma^n(b)| = +\infty$), the sequence $\sigma^n(\dots bbb \cdot aaa \dots)$ converges to $\sigma^\omega(b \cdot a)$ which is a fixed point of σ . It can happen that $\sigma^\omega(b \cdot a)$ does not belong to X_σ . In fact, $\sigma^\omega(b \cdot a)$ belongs to X_σ if and only if ba belongs to $\mathcal{L}(\sigma)$. In that case, we say that $\sigma^\omega(b \cdot a)$ is an *admissible* fixed point of σ .

With a morphism $\sigma : A^* \rightarrow B^*$ is naturally associated the *incidence matrix* $M_\sigma = (m_{i,j})_{i \in B, j \in A}$ where $m_{i,j}$ is the number of occurrences of i in the word $\sigma(j)$. Whenever $A = B$ and the matrix associated with σ is primitive (*i.e.*, when it has a power with strictly positive coefficients) we say that σ is a *primitive endomorphism*. In this situation we easily check that $\mathcal{L}(\sigma) = \mathcal{L}(\sigma^\omega(a))$ for any letter $a \in A$ on which σ is right-prolongable. In particular, we also have that for all $x \in X_\sigma$, $\Omega(x) = \Omega(\sigma^\omega(a)) = X_\sigma$. If σ is primitive we say (X_σ, S) is a *primitive substitution subshift*. Such a subshift is minimal [Que10]. A sequence x is *primitive substitutive* if it is substitutive w.r.t. a primitive substitution. The subshift $(\Omega(x), S)$ it generates is called *primitive substitutive subshift*. It is clearly minimal. We say σ is *aperiodic* whenever (X_σ, S) is aperiodic.

It is well known that if a matrix M is primitive, its spectral radius $\rho(M) = \max\{|\lambda| \mid \lambda \in \text{Spec}(M)\}$ is an eigenvalue of M which is algebraically simple. Furthermore, any eigenvalue of M different from $\rho(M)$ has modulus less than $\rho(M)$. We call $\rho(M)$ the *dominant eigenvalue* of M . By abuse of language, when M is the incidence matrix of a primitive endomorphism σ , we call $\rho(M)$ the *dominant eigenvalue* of σ .

2.4 Examples

Throughout the paper, we will try to illustrate our results on examples. To keep computations reasonable, we stick to well-known examples. In particular, we will mainly focus on:

1. The *Fibonacci sequence* is the fixed point $\varphi^\omega(a)$, where φ is the *Fibonacci substitution* defined by

$$\varphi : \begin{cases} a \mapsto ab \\ b \mapsto a \end{cases} .$$

We have

$$\varphi^\omega(a) = abaababaabaababaababaababaababaababaababaababaa \dots$$

The Fibonacci substitution is primitive, hence the Fibonacci sequence is uniformly recurrent. The *Fibonacci subshift* is $\Omega(\varphi^\omega(a))$.

2. The *Thue-Morse sequence* is the fixed point $\nu^\omega(a)$, where ν is the *Thue-Morse substitution* defined by

$$\nu : \begin{cases} a \mapsto ab \\ b \mapsto ba \end{cases} .$$

We have

$$\nu^\omega(a) = abbabaabbaababbabaababbaabbabaabbaababbaabbabaababb \dots$$

The Thue-Morse substitution is primitive, hence the Thue-Morse sequence is uniformly recurrent. The *Thue-Morse subshift* is $\Omega(\nu^\omega(a))$.

3. The *Chacon sequence* is the fixed point $\gamma^\omega(a)$, where γ is the *Chacon morphism* defined by

$$\gamma : \begin{cases} a \mapsto aaba \\ b \mapsto b \end{cases} .$$

We have

$$\gamma^\omega(a) = aabaaabababaaabaaababababababaaababababaaabababaaabababaa \dots$$

The Chacon sequence is uniformly recurrent, but γ is clearly not primitive. The *Chacon subshift* is $\Omega(\gamma^\omega(a))$.

3 From uniformly recurrent morphic sequences to purely substitutive sequences w.r.t. proper primitive substitutions

In this section we show that we can restrict our study to subshifts generated by purely substitutive sequences where the underlying substitution is primitive and proper. A substitution $\sigma : A^* \rightarrow A^*$ is *left proper* if there exists $a \in A$ such that $\sigma(A)$ is included in aA^* . It is *right proper* if there exists $b \in A$ such that $\sigma(A)$ is included in A^*b . It is *proper* whenever it is both left and right proper.

We show in Theorem 3.1 below that any subshift generated by an aperiodic uniformly recurrent morphic sequence is isomorphic to a subshift generated by a purely substitutive sequence whose underlying substitution is primitive and proper. We also give computable bounds for the radii of the isomorphism and of its inverse.

Theorem 3.1. *Let $y \in A^{\mathbb{K}}$ be an aperiodic uniformly recurrent morphic sequence and (Y, S) the minimal subshift it generates. There exist a computable constant K , a computable primitive proper substitution σ and an isomorphism ϕ from (X_σ, S) onto (Y, S) whose radius is 0 and whose radius of ϕ^{-1} is less than K .*

Remark 3.2. Theorem 3.1 is the first step to prove Theorem 1.3 and Theorem 1.4 (hence Theorem 1.1) for aperiodic subshifts. Indeed, suppose that we are given two aperiodic subshifts (X, S) and (Y, S) generated by uniformly recurrent morphic sequences. Using Theorem 3.1, we can compute primitive and proper substitutions σ and τ such that (X, S) is isomorphic to (X_σ, S) and (Y, S) is isomorphic to (X_τ, S) , with isomorphisms of computable radii. In the next sections, we show that for factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$, with σ and τ primitive and (X_τ, S) non periodic, the radius of some $f \circ S^n$ is bounded by a computable constant only depending on σ and τ , Theorem 5.10. We also show that it is decidable whether a given sliding block code defines a factor map from (X_σ, S) onto (X_τ, S) (Theorem 7.27). This will complete the proofs of Theorem 1.3 and Theorem 1.4 by composing the factor maps with the isomorphisms.

Remark 3.3. As the sequence y in Theorem 3.1 is uniformly recurrent, we have $Y = \Omega(y^+)$. Therefore, except for the definitions, we essentially consider one-sided sequences in this section.

3.1 Return words and derived sequences

Let $x \in A^{\mathbb{K}}$ be a sequence and u be a word in $\mathcal{L}(x)$. We call *return word to u* any word $w \in A^*$ such that wu is a word in $\mathcal{L}(x)$ that admits u as a prefix and u occurs exactly twice in wu . The set of return words to u in x is denoted by $\mathcal{R}_{x,u}$. It is easily seen that a sequence x is uniformly recurrent if and only if for all $u \in \mathcal{L}(x)$, the set $\mathcal{R}_{x,u}$ is finite.

Remark 3.4. If w is a return word to u in $x \in A^{\mathbb{K}}$, then the word $w^{1+\lfloor |u|/|w| \rfloor}$ occurs in x . Furthermore, if $|w| \leq |u|$, then the word $w^{1+\lfloor |u|/|w| \rfloor}$ occurs in x , where the fractional power $w^{p/|w|}$, $p \in \mathbb{N}$, of a word w is the word $w^n w_{[0,r[}$ such that $p/|w| = n + r/|w|$ with $n, r \in \mathbb{N}$ and $r < |w|$.

When dealing with return words, it is convenient to enumerate the elements of $\mathcal{R}_{x,u}$ in the order of their first occurrence in x^+ . Formally, we consider the set $R_{x,u} = \{1, 2, \dots\}$ with $\#R_{x,u} = \#\mathcal{R}_{x,u}$ and we define

$$\Theta_{x,u} : R_{x,u}^* \rightarrow \mathcal{R}_{x,u}^*$$

as the unique morphism that maps $R_{x,u}$ bijectively onto $\mathcal{R}_{x,u}$ and such that for all $i \in R_{x,u}$, $\Theta_{x,u}(i)u$ is the i th word of $\mathcal{R}_{x,u}u$ occurring in x^+ . When x is uniformly recurrent, for all $y \in X = \Omega(x)$ we have $\mathcal{R}_{y,u} = \mathcal{R}_{x,u}$ so we let $\mathcal{R}_{X,u}$ denote this set.

Example 3.5. For the Fibonacci sequence x , we have the return words $\mathcal{R}_{x,a} = \{a, ab\}$ and $\mathcal{R}_{x,aa} = \{aab, aabab\}$ and the morphisms $\Theta_{x,a}$, $\Theta_{x,aa}$ are given by

$$\Theta_{x,a} : \begin{cases} 1 \mapsto a \\ 2 \mapsto ab \end{cases} \quad \text{and} \quad \Theta_{x,aa} : \begin{cases} 1 \mapsto aabab \\ 2 \mapsto aaba \end{cases} .$$

For the Thue-Morse sequence y , we have the following sets of return words $\mathcal{R}_{y,b} = \{b, ba, baa\}$, $\mathcal{R}_{y,bb} = \{bbaa, bbaaba, bbabaa, bbabaaba\}$ and the morphisms $\Theta_{y,b}$, $\Theta_{y,bb}$ are given by

$$\Theta_{y,b} : \begin{cases} 1 \mapsto b \\ 2 \mapsto ba \\ 3 \mapsto baa \end{cases} \quad \text{and} \quad \Theta_{y,bb} : \begin{cases} 1 \mapsto bbabaa \\ 2 \mapsto bbaaba \\ 3 \mapsto bbaa \\ 4 \mapsto bbabaaba \end{cases} .$$

For the Chacon sequence z , we have the following sets of return words $\mathcal{R}_{z,a} = \{a, ab\}$, $\mathcal{R}_{z,aab} = \{aba, aabab\}$ and the morphisms $\Theta_{z,a}$, $\Theta_{z,aab}$ are given by

$$\Theta_{z,a} : \begin{cases} 1 \mapsto a \\ 2 \mapsto ab \end{cases} \quad \text{and} \quad \Theta_{z,aab} : \begin{cases} 1 \mapsto aaba \\ 2 \mapsto aabab \end{cases} .$$

The next results are classical when dealing with return words. The first three are proved in [Dur98a] for one-sided sequences and when u is a prefix of x , but the proofs are exactly the same in the two-sided case and when u is a factor but not a prefix. The proof of Proposition 3.7 that is in [Dur98a] only concerns x itself (and not $y \in \Omega(x)$) but again, the proof is the same.

Proposition 3.6 ([Dur98a]). *Let $x \in A^{\mathbb{K}}$ be a uniformly recurrent sequence and u be a word in $\mathcal{L}(x)$. The set $\mathcal{R}_{x,u}$ is a code, i.e., the map $\Theta_{x,u} : R_{x,u}^* \rightarrow \mathcal{R}_{x,u}^*$ is one-to-one.*

Proposition 3.7 ([Dur98a]). *Let x be a uniformly recurrent sequence in $A^{\mathbb{K}}$ and let u be a word in $\mathcal{L}(x)$. Let $y \in \Omega(x)$ and $i \in \mathbb{N}$ be the first occurrence of u in y^+ . Then, there exists a unique sequence $z \in R_{x,u}^{\mathbb{K}}$ satisfying $\Theta_{x,u}(z) = S^i(y)$; we call the sequence z a derived sequence of y (w.r.t. u) and we denote it by $\mathcal{D}_u(y)$. Furthermore, for all $y \in \Omega(x)$, we have $\Omega(\mathcal{D}_u(y)) = \Omega(\mathcal{D}_u(x))$.*

Proposition 3.8 ([Dur98a]). *Let x be a uniformly recurrent sequence in $A^{\mathbb{K}}$. Let u and u' be two prefixes of x^+ where u is a prefix of u' .*

1. *There is a unique non-erasing morphism $\lambda_{x,u,u'} : R_{x,u'}^* \rightarrow R_{x,u}^*$ such that*

$$\Theta_{x,u'} = \Theta_{x,u} \circ \lambda_{x,u,u'};$$

2. *If $\mathcal{D}_u(x^+) = \mathcal{D}_{u'}(x^+)$ and if for all $(w, w') \in \mathcal{R}_{x,u} \times \mathcal{R}_{x,u'}$, wu occurs in $w'u'$, then $\lambda_{x,u,u'}$ is primitive, left proper ($\lambda_{x,u,u'}(R_{x,u'}) \subset 1R_{x,u}^+$) and such that $\mathcal{D}_u(x^+) = \lambda_{x,u,u'}^\omega(1)$.*

Example 3.9. For the Fibonacci sequence x , the Thue-Morse sequence y and the Chacon sequence z , the morphisms $\lambda_{x,a,aa} : R_{x,aa}^* \rightarrow R_{x,a}^*$, $\lambda_{y,b,bb} : R_{y,bb}^* \rightarrow R_{y,b}^*$ and $\lambda_{z,a,aab} : R_{z,aab}^* \rightarrow R_{z,a}^*$ given by the previous proposition are respectively

$$\lambda_{x,a,aa} : \begin{cases} 1 \mapsto 122 \\ 2 \mapsto 121 \end{cases}, \quad \lambda_{y,b,bb} : \begin{cases} 1 \mapsto 123 \\ 2 \mapsto 132 \\ 3 \mapsto 13 \\ 4 \mapsto 1232 \end{cases} \quad \text{and} \quad \lambda_{z,a,aab} : \begin{cases} 1 \mapsto 121 \\ 2 \mapsto 122 \end{cases} .$$

Example 3.14. Using the formula given in the previous remark, the Fibonacci sequence is linearly recurrent for the constant $10^{\frac{1+\sqrt{5}}{2}}$. Numerical experiments suggest that the optimal constant is around 2,65. Similarly, the Thue-Morse sequence is linearly recurrent for the constant 16. Numerical experiments suggest that the optimal constant is 9. For the Chacon sequence, we would first need to consider it as a primitive substitutive sequence (the Chacon morphism not being primitive) to obtain an upper bound on its constant of linear recurrence.

Observe that in the next sections, we do not always suppose that the substitutions are proper, several results being true without that hypothesis.

3.2 Proof of Theorem 3.1

Proof. We make use of Proposition 3.7, Proposition 3.8, Theorem 3.10, Theorem 3.11 and Proposition 3.12.

Assume that y is a one-sided sequence, otherwise, take $y = y^+$. As y is uniformly recurrent, we have $\lim_{n \rightarrow +\infty} \min\{|w| \mid u \in \mathcal{L}_n(y), w \in \mathcal{R}_{y,u}\} = +\infty$. Using Theorem 3.10, we can algorithmically find two prefixes u and u' of y such that

- $\mathcal{D}_{u'}(y) = \mathcal{D}_u(y)$;
- $\Theta_{y,u}(1)u$ is a prefix of u' ;
- for all $(w, w') \in \mathcal{R}_{y,u} \times \mathcal{R}_{y,u'}$, wu occurs in $w'u'$.

Set $R = R_{y,u} = R_{y,u'}$. By Proposition 3.8, there is a primitive and left proper substitution $\tau = \lambda_{y,u,u'}$ ($\tau(1) \subset 1R^+$) such that $\mathcal{D}_u(y) = \tau^\omega(1)$. Since $\Theta_{y,u}$ and $\Theta_{y,u'}$ are computable by Theorem 3.10, the substitution τ is computable as well. In particular, we have $X_\tau = \Omega(\mathcal{D}_u(y))$, $\Theta_{y,u}(X_\tau) \subset Y$ and, by Proposition 3.7, for all $y' \in Y$, there exists a unique pair $(x, n) \in X_\tau \times \mathbb{N}$ such that

- u is prefix of $S^{-n}(y')$ and is not prefix of $S^{-m}(y')$ for all integers $0 \leq m < n$ (in particular, we have $0 \leq n < |\Theta_{y,u}|$);
- $y' = S^n \Theta_{y,u}(x)$.

Set $\Theta = \Theta_{y,u}$ and consider the alphabet $D = \{(r, k) \mid r \in R, 0 \leq k < |\Theta(r)|\}$ and the morphism $\psi : R^* \rightarrow D^*$ defined by:

$$\psi(r) = (r, 0) \dots (r, |\Theta(r)| - 1).$$

Since $\langle \tau \rangle \geq 2$, there is an integer $n \leq |\Theta|$ such that $\langle \tau^n \rangle \geq |\Theta|$. Let σ be the endomorphism from D^* to D^* defined by:

$$\begin{aligned} \sigma((r, k)) &= \psi(\tau^n(r)_{[k,k]}), & \text{if } 0 \leq k < |\Theta(r)| - 1; \\ \sigma((r, |\Theta(r)| - 1)) &= \psi(\tau^n(r)_{[|\Theta(r)|-1, |\tau^n(r)|-1]}), & \text{otherwise.} \end{aligned}$$

Observe that τ being primitive, so is σ . Furthermore, since τ is left proper, it is easily seen that σ^2 is left proper. For all r in R , we have

$$\begin{aligned}\sigma(\psi(r)) &= \sigma((r, 0) \cdots (r, |\Theta(r)| - 1)) \\ &= \psi(\tau^n(r)_{[0,0]}) \cdots \psi(\tau^n(r)_{[|\Theta(r)|-1, |\tau^n(r)|-1]}) \\ &= \psi(\tau^n(r)),\end{aligned}$$

hence $\sigma \circ \psi = \psi \circ \tau^n$. Since τ is left proper, it has a fixed point $z \in R^{\mathbb{N}}$ (which is unique) and it satisfies $\sigma(\psi(z)) = \psi(\tau^n(z)) = \psi(z)$. Hence $\psi(z)$ is a fixed point of the substitution σ and $X_\sigma = \Omega(\psi(z))$. In particular, $\psi(X_\tau) \subset X_\sigma$ and for all $x \in X_\sigma$, there exists a unique pair $(z', n) \in X_\tau \times \mathbb{N}$ such that

- $(S^{-n}(x))_0 = (r, 0)$ for some $r \in R$ and $(S^{-m}(x))_0 \notin R \times \{0\}$ for all integers $0 \leq m < n$ (in particular we have $0 \leq n < |\psi| = |\Theta|$);
- $x = S^n \psi(z')$.

Let ϕ be the coding from D^* to A^* defined by $\phi((r, k)) = \Theta(r)_{[k,k]}$ for all (r, k) in D . For all r in R we obtain

$$\phi(\psi(r)) = \phi((r, 0) \cdots (r, |\Theta(r)| - 1)) = \Theta(r),$$

and then $\phi \circ \psi = \Theta$. Let us show that ϕ defines a factor map of radius 0 from (X_σ, S) onto (Y, S) . We have $\phi(X_\sigma) = Y$: if $x \in X_\sigma$, then $x = S^n \psi(z')$ for some integer n and some $z' \in X_\tau$. Therefore, $\phi(x) = S^n(\phi \circ \psi(z')) = S^n(\Theta(z')) \in Y$. Reciprocally, if $y' \in Y$, there exists a unique pair $(x, n) \in X_\tau \times \mathbb{N}$ such that $y' = S^n \Theta_{y,u}(x)$ where u is not prefix of $S^{-m}(y')$ for all integers $0 \leq m < n$. As $\phi \circ \psi = \Theta$, we then have $y' = \phi(S^n(\psi(x)))$, with $\psi(x) \in X_\sigma$.

To show that ϕ is an isomorphism, we build a factor map f from (Y, S) onto (X_σ, S) and we show that $f \circ \phi$ is the identity. Let L be the constant of linear recurrence of $\mathcal{D}_u(y)$. It is computable by Proposition 3.12 (since τ is computable). Since $y = \theta(\mathcal{D}_u(y))$, y is also linearly recurrent and for the constant $L' = |\Theta|L$. Set $K = (L + 1)|u| + 1$. Let $w = w_{-K} \cdots w_K$, $w_i \in A$ for all i , be a word of length $2K + 1$ of $\mathcal{L}(Y)$. By Theorem 3.11, there is a smallest non-negative integer $i < (L + 1)|u|$ such that $w_{[-i, -i+|u|]} = u$. With the same argument, there is a smallest positive integer $j \leq L|u| + 1$ such that $w_{[j, j+|u|]} = u$. This defines a letter $r_w \in R$ such that $\Theta(r_w) = w_{[-i, j]}$. We define $\hat{f} : A^{2K+1} \rightarrow R$ by $\hat{f}(w) = \psi(r_w)_i = (r_w, i)$ and we call $f : A^{\mathbb{Z}} \rightarrow R^{\mathbb{Z}}$ the map defined by \hat{f} , i.e., $(f(x))_n = \hat{f}(x_{[n-K, n+K]})$ for all x, n . Let us show $f \circ \phi$ is the identity.

Indeed, let $x \in X_\sigma$ and $y = \phi(x)$. To show that $f(y) = x$, it is enough to show that $\hat{f}(y_{[-K, K]}) = x_0$. Let $i \geq 0$ and $j > 0$, be the smallest integers such that $x_{-i} = (r, 0)$ and $x_j = (r', 0)$ for some $r, r' \in R$. Then, $x_{[-i, j-1]} = \psi(r)$ and $y_{[i, j+|u|]} = \phi(x_{[i, j+|u|]}) = \phi(\psi(r))u = \Theta(r)u$. Setting $w = y_{[-K, K]}$, we have $r = r_w$ and $\hat{f}(y_{[-K, K]}) = \psi(r_w)_i = x_0$. This shows that ϕ is a isomorphism of radius 0 from (X_σ, S) onto (Y, S) and that its inverse $f = \phi^{-1}$ has radius K .

To conclude the proof, it suffices to show that (X_σ, S) can be obtained as a primitive substitution subshift with a proper substitution. Indeed, up to now, we only know that σ^2 is left proper. Let $a \in D$ be the letter such that $\sigma^2(D) \subset aD^+$. We consider the substitutions σ' and ξ respectively defined by $a\sigma'(d) = \sigma^2(d)a$ for all $d \in D$ and $\xi = \sigma' \circ \sigma^2$. The substitution ξ is proper, primitive and such that $X_\xi = X_\sigma$, which ends the proof. \square

3.3 Examples

Let us apply Theorem 3.1 on two examples. Our aim is to follow the construction given in the proof of the theorem.

The first one is the Chacon subshift generated by the Chacon sequence z . Let us recall what we know from Example 3.5 and Example 3.9:

$$\Theta_{z,a} : \begin{cases} 1 \mapsto a \\ 2 \mapsto ab \end{cases} \quad ; \quad \Theta_{z,aba} : \begin{cases} 1 \mapsto aaba \\ 2 \mapsto aabab \end{cases} \quad ; \quad \tau_z : \begin{cases} 1 \mapsto 121 \\ 2 \mapsto 122 \end{cases} .$$

We consider the notation of the theorem and successively obtain:

- $\tau = \tau_z$;
- $D = \{(1, 0), (2, 0), (2, 1)\}$ and $n = 1$;
- $\psi : \begin{cases} 1 \mapsto (1, 0) \\ 2 \mapsto (2, 0)(2, 1) \end{cases}$ and $\sigma : \begin{cases} (1, 0) \mapsto (1, 0)(2, 0)(2, 1)(1, 0) \\ (2, 0) \mapsto (1, 0) \\ (2, 1) \mapsto (2, 0)(2, 1)(2, 0)(2, 1) \end{cases}$;
- $\phi : \begin{cases} (1, 0) \mapsto a \\ (2, 0) \mapsto a \\ (2, 1) \mapsto b \end{cases}$.

The theorem states that ϕ is an isomorphism from the Chacon subshift onto the subshift (X_ξ, S) , where ξ is the primitive and proper substitution defined by $\xi = \sigma' \circ \sigma^2$ and σ' is defined by $(1, 0)\sigma'(d) = \sigma^2(d)(1, 0)$ for all $d \in D$.

Let us consider a second example with a sequence y which is not purely morphic like the Chacon sequence. We consider the morphisms

$$\sigma : \begin{cases} 0 \mapsto 0123 \\ 1 \mapsto 02 \\ 2 \mapsto 13 \\ 3 \mapsto 3 \end{cases} \quad \text{and} \quad \psi : \begin{cases} 0 \mapsto abb \\ 1 \mapsto ab \\ 2 \mapsto a \\ 3 \mapsto \epsilon \end{cases} .$$

Let $x = \sigma^\omega(0)$. It is clearly non uniformly recurrent and non periodic as the words 3^n , $n \in \mathbb{N}$, appears infinitely many times separated by some words over the alphabet $\{0, 1, 2\}$. Nonetheless, $y = \psi(x)$ is uniformly recurrent, as could be checked by the algorithm given in [Dur13b].

The algorithm of Theorem 3.10 gives $\mathcal{D}_u(y) = \mathcal{D}_{u'}(y)$, with $u = abba$ and $u' = abbabaabb$. Then, we obtain $z = \mathcal{D}_u(y) = \tau^\omega(1)$, with τ defined by $\Theta_{y,u'} = \Theta_{y,u} \circ \tau$ and

$$\Theta_{y,u} : \begin{cases} 1 \mapsto abbaba \\ 2 \mapsto abbaab \\ 3 \mapsto abbabaab \\ 4 \mapsto abba \end{cases}, \Theta_{y,u'} : \begin{cases} 1 \mapsto abbabaabbaab \\ 2 \mapsto abbabaababba \\ 3 \mapsto abbabaabbaababba \\ 4 \mapsto abbabaab \end{cases}, \tau : \begin{cases} 1 \mapsto 12 \\ 2 \mapsto 34 \\ 3 \mapsto 124 \\ 4 \mapsto 3 \end{cases}.$$

We finally obtain a primitive and proper substitution $\xi : \mathcal{A}^* \rightarrow \mathcal{A}^*$ on the 24 letter alphabet $D = \{(i, j) \mid i \in \{1, 2, 3, 4\}, 0 \leq j < |\Theta_{y,u}(i)|\}$ and a coding $\phi : D^* \rightarrow \{a, b\}^*$ that induces an isomorphism from (X_ξ, S) onto $(\Omega(y), S)$. We could moreover show that $(\Omega(y), S)$ is the Thue-Morse subshift, as it could be checked by using the algorithm that we provide later in this paper.

4 Bounds for primitive substitutive sequences

In the previous section, we have shown that to prove Theorem 1.1, Corollary 1.2, Theorem 1.3 and Theorem 1.4, we can restrict ourselves to the case of subshifts generated by primitive and proper substitutions (see Remark 3.2). In this section, we consider primitive substitution subshifts (X_σ, S) and (X_τ, S) and develop some tools that will help us to bound the radius of factor maps from (X_σ, S) onto (X_τ, S) (whenever such maps exist). In particular, we give some computable constants about the substitutions σ and τ (sometimes assuming that there is a factor map from (X_σ, S) onto (X_τ, S)) and we consider a variation of the notion of substitution on the words of length n [Que10].

Observe that in this section, we do not assume that σ and τ are proper, the results being true without this hypothesis.

4.1 Recognizability

Let $\sigma : A^* \rightarrow A^*$ be a primitive substitution. Taking a power of σ if needed, we can suppose σ has an admissible fixed point x in $A^\mathbb{Z}$. Let $E(x, \sigma) = \{e_n \mid n \in \mathbb{Z}\}$ be the subset of integers defined by:

$$e_n = \begin{cases} -|\sigma(x[n, -1])|, & \text{if } n < 0; \\ 0, & \text{if } n = 0; \\ |\sigma(x[0, n-1])|, & \text{if } n > 0. \end{cases}$$

The integers e_n are usually called the *cutting bars* of the substitution. The following result is fundamental for substitutive subshifts.

Theorem 4.1. *Let $\sigma : A^* \rightarrow A^*$ be an aperiodic primitive substitution and $x \in A^\mathbb{Z}$ be an admissible fixed point of σ .*

1. [Mos96] *There exists $M > 0$ such that*

$$(n \in E(x, \sigma), m \in \mathbb{Z}, x[n-M, n+M] = x[m-M, m+M]) \implies m \in E(x, \sigma).$$

2. [Mos92] There exists $L \geq M$ such that

$$(i, j \in \mathbb{Z}, x[e_i - L, e_i + L] = x[e_j - L, e_j + L]) \implies x_i = x_j.$$

In particular, the constants M and L do not depend on the admissible fixed point x .

The smallest L satisfying (2) in the previous theorem is called the *recognizability constant* for the substitution σ and is denoted L_σ .

Observe that if $u \in \mathcal{L}(\sigma)$ has length at least $2L + 1$ and if $i, j \in \mathbb{Z}$, $i < j$, are two occurrences of u in x , then the first part of Mossé's Theorem ensures that

$$[i + L, i + |u| - L] \cap E(x, \sigma) = ([j + L, j + |u| - L] \cap E(x, \sigma)) - (j - i)$$

and the second part states that the central part of u has the same preimage under σ at positions i and j . We call the set $([i + L, i + |u| - L] \cap E(x, \sigma)) - i$ the set of *relative cutting bars* of u .

Example 4.2. For the Fibonacci substitution φ , we have $E(x, \varphi) = \{i \in \mathbb{Z} \mid x_i = a\}$ and so $M = 1$. Then, the words of length 3 with a letter a as central position are aab, baa, bab . It is immediate that if $x_{[e_i, e_i+1]} = ab$, then $x_i = a$ and if $x_{[e_i, e_i+1]} = aa$, then $x_i = b$. As a consequence, the constant of recognizability for the Fibonacci substitution is $L_\varphi = 1$.

For the Thue-Morse substitution ν , the constant M has to be greater than 1, since the word aba both occurs in the language as $a\nu(b)$ and $\nu(a)b$. On the contrary, any word u in $\mathcal{L}_5(\nu)$ contains an occurrence of the word aa or bb and these two words uniquely determine the relative cutting bars in u . Then for all i , we have $x_{e_i} = x_i$, which shows that the constant of recognizability of ν is $L_\nu = 2$.

The two following lemmas are consequences of Mossé's theorem (Theorem 4.1).

Corollary 4.3 ([Que10, Corollary 5.11]). *Let σ be an aperiodic primitive substitution. For all $n \geq 1$,*

1. σ^n is a homeomorphism from X_σ onto the clopen $\sigma^n(X_\sigma)$;
2. for every $x \in X_\sigma$, there is a unique $y \in X_\sigma$ and a unique k , $0 \leq k < |\sigma^n(y_0)|$, such that $x = S^k(\sigma^n(y))$.

Corollary 4.4. *Let σ be an aperiodic primitive substitution and x, y be two elements in X_σ . Then,*

1. If there exists n such that $\sigma(x) = S^n\sigma(y)$ then there exists m such that $x = S^m(y)$. Moreover, if $n \geq 0$ then $0 \leq m \leq n/\langle \sigma \rangle$;
2. If $\sigma^k(x)$ and $\sigma^k(y)$ coincide on the indices $[-m, n]$, with $m, n \geq L_{\sigma^k}$, then x and y coincide on the indices $\left[\left[\frac{-m + L_{\sigma^k}}{|\sigma^k|} \right], \left[\frac{n - L_{\sigma^k}}{|\sigma^k|} \right] \right]$.

3. If σ has constant length and is one-to-one on the letters, and if $\sigma^k(x)$ and $\sigma^k(y)$ coincide on the indices $[-m, n]$, then x and y coincide on the indices $\left[\left\lfloor \frac{-m}{|\sigma^k|} \right\rfloor, \left\lfloor \frac{n}{|\sigma^k|} \right\rfloor \right]$.

Corollary 4.5 ([Que10, Proposition 5.20]). *Let $\sigma : A^* \rightarrow A^*$ be an aperiodic primitive substitution with recognizability constant L_σ . Then, $\{S^i\sigma([a]) \mid a \in A, 0 \leq i < |\sigma(a)|\}$ is a clopen partition of X_σ . Moreover, if $x, y \in X_\sigma$ coincide on the indices $[-n, n]$ with $n = L_\sigma + |\sigma|$, then they belong to the same atom of this partition.*

Later, we will not only need the existence of the recognizability constant but also a bound on its value depending on the data of the primitive substitution.

Theorem 4.6. [DL17] *Let $\sigma : A^* \rightarrow A^*$ be an aperiodic primitive morphism that has an admissible fixed point $x \in A^{\mathbb{Z}}$. Then*

$$L_\sigma \leq 2|\sigma|^{6(\text{Card}A)^2 + 6(\text{Card}A)|\sigma|^{28(\text{Card}A)^2}} + |\sigma|^{\text{Card}A}.$$

Moreover, if σ is one-to-one on letters then,

$$L_\sigma \leq 2|\sigma|^{6(\text{Card}A)^2 + 6|\sigma|^{28(\text{Card}A)^2}} + |\sigma|.$$

In [KM16] better bounds are given for the two letter alphabet but in terms of “synchronization delay” and “circularity”. As the recognizability constant is bounded by the synchronization delay (see comments in [DL17]), this gives the following theorem that greatly improves our bounds.

Theorem 4.7 ([KM16]). *If $\text{Card}A = 2$ and $\sigma : A^* \rightarrow A^*$ is a primitive substitution of constant length k .*

1. $L_\sigma \leq 8$, if $k = 2$;
2. $L_\sigma \leq k^2 + 3k - 4$, if k is an odd prime number;
3. $L_\sigma \leq k^2 \left(\frac{k}{d} - 1\right) + 5k - 4$, otherwise, where d is the least divisor of k .

Later we will need to control the recognizability constant of σ^k with respect to its growth.

Proposition 4.8. [DL17, Proposition 13] *If $\sigma : A^* \rightarrow A^*$ is recognizable on the admissible fixed point $x \in A^{\mathbb{Z}}$. Then, for all $k > 0$,*

$$L_{\sigma^k} \leq L_\sigma \sum_{i=0}^{k-1} |\sigma^i|.$$

4.2 Block representations of subshifts and substitutions on the blocks of radius n

Let (X, S) be a subshift over the alphabet A . For any positive integer n , we let $A^{(n)}$ denote the set $A^{(n)} = \{(w) \mid w \in \mathcal{L}_{2n+1}(\sigma)\}$ considered as an alphabet. The following result is classical [Que10].

Lemma 4.9. *Let $\phi : A^{2n+1} \rightarrow A^{(n)}$ be the sliding block code defined by*

$$\phi(a_1 \cdots a_{2n+1}) = (a_1 \cdots a_{2n+1}).$$

Then ϕ defines an isomorphism f from (X, S) to $(X^{(n)}, S)$, where $X^{(n)} = f(X)$.

We call $(X^{(n)}, S)$ the n -block representation of (X, S) . When the subshift X of the previous result is a substitutive subshift associated with a substitution σ , then $X^{(n)}$ is also substitutive and associated to the *substitution on the words of length $2n+1$* of σ . The definition we give below of the well-known notion of substitution on the words of length n (see [Que10] for more details) is slightly different from the usual one as we need to define it for a “two-sided context”.

Let $\sigma : A^* \rightarrow A^*$ be a substitution and $n \geq 1$. Below any time we take (uv) in $A^{(n)}$, we suppose that u belongs to A^n and v belongs to A^{n+1} . We define the map $\pi^{(n)} : (A^{(n)})^* \rightarrow A^*$ by, for all $(uv) \in A^{(n)}$, $\pi^{(n)}((uv)) = v_0$, where $v = v_0 v_1 \dots v_n$, $v_i \in A$ for all i . We also define the substitution $\sigma^{(n)} : (A^{(n)})^* \rightarrow (A^{(n)})^*$ by, for $(uv) \in A^{(n)}$ with $\sigma(u) = a_1 \cdots a_{k-1}$, $\sigma(v) = a_k \cdots a_{k+l}$ where the a_i 's belong to A and $p = |\sigma(v_0)|$,

$$\sigma^{(n)}((uv)) = (a_{k-n} \cdots a_{k+n})(a_{k-n+1} \cdots a_{k+n+1}) \cdots (a_{k-n+p-1} \cdots a_{k+p+n-1}).$$

In other words, $\sigma^{(n)}((uv))$ consists in the ordered list of the first p factors of length $2n+1$ of $\sigma(uv) = a_1 \cdots a_k a_{k+1} \cdots a_{k+l}$ starting from the index $k-n$. We say that $\sigma^{(n)}$ the *substitution on the blocks of radius n* .

Example 4.10. Let us build the substitution $\gamma^{(5)}$ on the blocks of radius 2 associated with the Chacon morphism γ . The words of length 5 in $\mathcal{L}(\gamma)$ are

$$\mathcal{L}_5(\gamma) = \{aabaa, abaaa, baaab, aaaba, aabab, ababa, babaa, abaab, baaba\}.$$

We obtain

$$\gamma^{(5)} : \begin{cases} (aabaa), (aabab), (babaa) & \mapsto (babaa) \\ (abaaa), (abaab) & \mapsto (abaab)(baaba)(aabab)(ababa) \\ (baaba), (aaaba) & \mapsto (baaab)(aaaba)(aabab)(ababa) \\ (baaab) & \mapsto (baaab)(aaaba)(aabaa)(abaaa) \\ (ababa) & \mapsto (abaab)(baaba)(aabab)(ababa) \end{cases}.$$

Proposition 4.11. *Let σ be a primitive substitution, then for all n , $\sigma^{(n)}$ is computable, primitive and such that $X_\sigma^{(n)} = X_{\sigma^{(n)}}$. In particular, for all i and*

n , if x is a two-sided admissible fixed point of σ and $x^{(n)} = ((x_{[-n+i, n+i]})_{i \in \mathbb{Z}})$, then $x^{(n)}$ is an admissible fixed point of $\sigma^{(n)}$ and

$$\{j \in \mathbb{Z} \mid S^j(x) \in \sigma^i(X_\sigma)\} = \{j \in \mathbb{Z} \mid S^j(x^{(n)}) \in (\sigma^{(n)})^i(X_{\sigma^{(n)}})\}. \quad (1)$$

Moreover, for all n and i , $(\sigma^{(n)})^i = (\sigma^i)^{(n)}$ and $|\sigma^{(n)i}((uv))| = |\sigma^i(v_0)|$ for all $uv \in A^{(n)}$ where $v = v_0 v_1 \cdots v_n$. In particular, the incidence matrices M_σ and $M_{\sigma^{(n)}}$ have the same spectral radius.

Proof. First, let us prove by induction that for every $i \geq 1$,

$$(\sigma^{(n)})^i = (\sigma^i)^{(n)}. \quad (2)$$

The base case is trivial so let us assume that the result holds for some $i \geq 1$. For any $(uv) \in A^{(n)}$, we have

$$\begin{aligned} (\sigma^{(n)})^{i+1}((uv)) &= \sigma^{(n)}(\sigma^{(n)})^i((uv)) \\ &= \sigma^{(n)}(\sigma^i)^{(n)}((uv)), \end{aligned}$$

where we have used the induction hypothesis. Let us write

$$\begin{aligned} \sigma^i(uv) &= a_1 \cdots a_p, & a_j \in A \text{ for all } j \\ \sigma^{i+1}(uv) &= b_1 \cdots b_q, & b_j \in A \text{ for all } j. \end{aligned}$$

By definition, we have, with $k_i = |\sigma^i(u)|+1$, $k_{i+1} = |\sigma^{i+1}(u)|+1$, $\ell_i = |\sigma^i(v_0)|-1$ and $\ell_{i+1} = |\sigma^{i+1}(v_0)|-1$,

$$\begin{aligned} (\sigma^i)^{(n)}((uv)) &= (a_{k_i-n} \cdots a_{k_i+n}) \cdots (a_{k_i-n+\ell_i} \cdots a_{k_i+n+\ell_i}); \\ (\sigma^{i+1})^{(n)}((uv)) &= (b_{k_{i+1}-n} \cdots b_{k_{i+1}+n}) \cdots (b_{k_{i+1}-n+\ell_{i+1}} \cdots b_{k_{i+1}+n+\ell_{i+1}}). \end{aligned}$$

For all $m \in \{0, 1, \dots, \ell_i\}$, if we set

$$\begin{aligned} r_m &= |\sigma(a_1 \cdots a_{k_i-n+m-1})| \\ s_m &= |\sigma(a_{k_i-n+m} \cdots a_{k_i+n+m})| \\ t_m &= |\sigma(a_{k_i-n+m} \cdots a_{k_i+m-1})| + 1 \\ z_m &= |\sigma(a_{k_i+m})| - 1, \end{aligned}$$

we get

$$\sigma(a_{k_i-n+m} \cdots a_{k_i+n+m}) = b_{r_m+1} \cdots b_{r_m+s_m}$$

and thus

$$\begin{aligned} \sigma^{(n)}((a_{k_i-n+m} \cdots a_{k_i+n+m})) &= (b_{r_m+t_m-n} \cdots b_{r_m+t_m+n}) \cdots \\ &\quad (b_{r_m+t_m-n+z_m} \cdots b_{r_m+t_m+n+z_m}). \end{aligned}$$

To finish the proof of (2), it suffices to notice that we have

$$\begin{aligned} r_0 + t_0 &= k_{i+1}; \\ r_m + t_m + z_m + 1 &= r_{m+1} + t_{m+1}, \quad \text{for all } m \in \{0, \dots, \ell_i - 1\}; \\ r_{\ell_i} + t_{\ell_i} + z_{\ell_i} &= k_{i+1} + \ell_{i+1}. \end{aligned}$$

Using (2), for the sake of simplicity we let $\sigma^{(n)i}$ denote the substitution $(\sigma^{(n)})^i$. We deduce that for all i

$$\begin{aligned}\sigma^{(n)i}((x_{[-n,n]})) &= (x_{[-n,n]}) \cdots (x_{[|\sigma^i(x_0)|-1-n, |\sigma^i(x_0)|-1+n]}) \\ \sigma^{(n)i}((x_{[-n-1, n-1]})) &= (x_{[-|\sigma^i(x_{-1})|-n, -|\sigma^i(x_{-1})|+n]}) \cdots (x_{[-n-1, n-1]})\end{aligned}$$

and that $x^{(n)} = ((x_{[-n+i, n+i]}))_{i \in \mathbb{Z}}$ is an admissible two-sided fixed point of $\sigma^{(n)}$. In particular, for all $(uv) \in A^{(n)}$, we have

$$|\sigma^{(n)i}((uv))| = |\sigma^i(v_0)|,$$

from which we deduce (1) and the fact that M_σ and $M_{\sigma^{(n)}}$ have the same spectral radius.

Let us show that $\sigma^{(n)}$ is primitive. If i is large enough, every $w \in \mathcal{L}(\sigma)$ of length $2n+1$ is a factor of $\sigma^i(a)$ for every $a \in A$. Thus, every $(uv) \in A^{(n)}$ occurs in $\sigma^{(n)i}((u'v'))$ for every $(u'v') \in A^{(n)}$.

For the equality $X_\sigma^{(n)} = X_{\sigma^{(n)}}$, as $x^{(n)}$ is uniformly recurrent, we have hence $X_\sigma^{(n)} = \Omega(x^{(n)})$. Furthermore, as $x^{(n)}$ is also an admissible fixed point of the primitive substitution $\sigma^{(n)}$, we have $X_{\sigma^{(n)}} = \Omega(x^{(n)})$. \square

4.3 Computable bounds for linear recurrence of substitutive sequences

We now develop some lemmas that allow to relate the quantities $|\sigma^n|$ and $\langle \sigma^n \rangle$ to the spectral radius of M_σ^n . First recall that if (X, T) is a dynamical system, an *invariant measure* for (X, T) is a probability measure μ on the σ -algebra of Borel sets such that $\mu(T^{-1}B) = \mu(B)$ for all Borel sets B . Such a measure is *ergodic* if every T -invariant Borel set has measure 0 or 1. The system (X, T) is said to be *uniquely ergodic* if there exists a unique invariant measure (which is then ergodic). Let us recall the ergodic theorem for such systems: Let μ be the unique T -invariant ergodic measure of (X, T) and $f : X \rightarrow \mathbb{R}$ be a continuous function, then $((1/n) \sum_{i=0}^{n-1} f \circ T^i)_n$ converges uniformly to $\int_X f d\mu$. We refer the reader to [HK95] for more details on ergodicity.

Proposition 4.12 ([Dur00, Proposition 13]). *If (X, S) is an aperiodic linearly recurrent subshift, then it is uniquely ergodic. Furthermore, if μ denotes the unique invariant measure of (X, S) and if K is its constant of linear recurrence, we have for all $u \in \mathcal{L}(X)$,*

$$1/K \leq |u|\mu([u]) \leq K.$$

Lemma 4.13 ([HJ90, Corollary 8.1.33]). *Let $M = (m_{i,j})$ be a $d \times d$ -matrix with non-negative entries. If M has a positive eigenvector x , then for all $n = 1, 2, \dots$ and all $i = 1, \dots, d$ we have*

$$\left[\frac{\min_{1 \leq k \leq d} x_k}{\max_{1 \leq k \leq d} x_k} \right] \rho(M)^n \leq \sum_{j=1}^d m_{ij}^{(n)} \leq \left[\frac{\max_{1 \leq k \leq d} x_k}{\min_{1 \leq k \leq d} x_k} \right] \rho(M)^n,$$

where $\rho(M)$ is the spectral radius of M and $M^n = (m_{ij}^{(n)})_{i,j}$.

Any substitution subshift (X_σ, S) , where σ is primitive, has a unique ergodic measure μ [Que10]. We denote by $\mu^{(k)}$ the vector whose entries are $\mu([u])$, $u \in \mathcal{L}_k(\sigma)$.

As we already said above, our definition of n -block substitutions is slightly different from the usual one in [Que10], nevertheless the following result remains true for our definition.

Lemma 4.14 ([Que10, Corollary 5.4 and Proposition 5.10]). *Let $\sigma : A^* \rightarrow A^*$ be an aperiodic primitive substitution and let μ be the unique invariant measure of (X_σ, S) . The matrix $M_{\sigma^{(k)}}$ has the same dominant eigenvalue as M_σ . Furthermore, the vector $\mu^{(k)}$ is a positive eigenvector of $M_{\sigma^{(k)}}$ associated with this dominant eigenvalue.*

The next result is a direct consequence of Proposition 4.12, Lemma 4.13 and Lemma 4.14.

Corollary 4.15. *Let σ be an aperiodic primitive substitution. We have for all n ,*

$$(1/K_\sigma^2)\rho(M_\sigma)^n \leq \langle \sigma^n \rangle \leq |\sigma^n| \leq K_\sigma^2 \rho(M_\sigma)^n; \quad (3)$$

$$|\sigma^n| \leq K_\sigma^4 \langle \sigma^n \rangle. \quad (4)$$

Let us show the same result holds for subshift factors of (X_σ, S) keeping the same constant K_σ where K_τ should be used. It is inspired from Theorem 3.11. The following proposition quantitatively expresses how key features of the dynamics of an aperiodic factor (X_τ, S) of a primitive substitution subshift (X_σ, S) are controlled by σ . But not only, observe that Item (1) and (2) claim that the relative growth of τ is controlled by a constant only depending on σ .

Proposition 4.16. *Let (X, S) be an aperiodic linearly recurrent subshift for the constant K and (Y, S) be one of its aperiodic factors given by a factor map of radius r . We have the following.*

1. For all n , $p_Y(n) \leq K(n + 2r)$.
2. For all n , every word of $\mathcal{L}_n(Y)$ occurs in every word of $\mathcal{L}_{(K+1)n+2rK}(Y)$.
3. For all $v \in \mathcal{L}(Y)$ and all $w \in \mathcal{R}_{Y,v}$,

$$\frac{|v| - 2rK}{K} \leq |w| \leq K(|v| + 2r).$$

4. For all $v \in \mathcal{L}(Y)$ with $|v| \geq 2rK + 1$ one has

$$\#(\mathcal{R}_{Y,v}) \leq \frac{(K+1)K^2(|v| + 2r) + 2rK^2}{|v| - 2rK}.$$

5. Let δ be the unique ergodic measure of (Y, S) . For all $v \in \mathcal{L}(Y)$, with $|v| > 2rK$,

$$\frac{1}{K(1 + 2r/|v|)} \leq |v|\delta([v]) \leq \frac{K}{1 - 2rK/|v|}.$$

Proof. Let \hat{f} be a block map with radius r defining a factor map from (X, S) to (Y, S) .

(1) We have $\mathcal{L}_n(Y) = \hat{f}(\mathcal{L}_{n+2r}(X))$ so the result follows from Theorem 3.11.

(2) Let $u \in \mathcal{L}_n(Y)$ and $v \in \mathcal{L}_{(K+1)n+2rK}(Y)$. There exist $u' \in \mathcal{L}_{n+2r}(X)$ and $v' \in \mathcal{L}_{(K+1)(n+2r)}(X)$ such that $u = \hat{f}(u')$ and $v = \hat{f}(v')$. By Theorem 3.11, u' occurs in v' . Thus u occurs in v .

(3) Let $v \in \mathcal{L}(Y)$. For any return word $w \in \mathcal{R}_{Y,v}$, there exist $v_1, v_2 \in \hat{f}^{-1}(\{v\})$ and $w' \in \mathcal{L}(X)$ such that v_1 is a prefix of $w'v_2$, $w'v_2$ contains only two occurrences of words in $\hat{f}^{-1}(\{v\})$ and $\hat{f}(w'v_2) = wv$. In particular w' is a subword of a return word in \mathcal{R}_{X,v_1} and, using Theorem 3.11, we have $|w| = |w'| \leq K(|v| + 2r)$.

Let us now show by contradiction that $\frac{|v|-2rK}{K} \leq |w|$. If $\frac{|v|-2rK}{K} > |w|$, then, by Remark 3.4, the word $u = w^{1+|v|/|w|}$ belongs to $\mathcal{L}(Y)$ and this word has length $|u| = |w| + |v| > (K+1)|w| + 2rK$. By Item (2), u contains an occurrence of all words in $\mathcal{L}_{|w|}(Y)$. As the number of different words of length $|w|$ occurring in u is at most $|w|$, this implies by Morse and Hedlund's theorem [MH40] that Y is periodic, which contradicts the hypothesis.

(4) Let $v \in \mathcal{L}(Y)$ with $|v| \geq 2rK + 1$. From (3), the length of return words to v are less than $K(|v| + 2r)$. Let w be a word of length $(K+1)K(|v| + 2r) + 2rK$. From (2) all return words to v appear in w . Hence, from (3) the number of return words to v is bounded by

$$\frac{(K+1)K(|v| + 2r) + 2rK}{\frac{|v|-2rK}{K}}.$$

(5) Let $v \in \mathcal{L}(Y)$. The ergodic theorem applied to the indicator function $1_{[v]}$ gives that for all $x \in Y$

$$\frac{1}{N} \sum_{n=0}^{N-1} 1_{[v]}(S^n x) \longrightarrow_{N \rightarrow +\infty} \delta([v]).$$

By definition, we have

$$\sum_{n=0}^{N-1} 1_{[v]}(S^n x) = \text{Card}(\{i \in \{0, \dots, N-1\} \mid x_{[i, i+|v|-1]} = v\}).$$

Using item (3), one obtains

$$\frac{N}{K(|v| + 2r)} - 1 \leq \sum_{n=0}^{N-1} 1_{[v]}(S^n x) \leq \frac{NK}{|v| - 2rK} + 1,$$

from which the inequalities follow. \square

Proposition 4.17. *Let σ and τ be two primitive substitutions such that (X_τ, S) is an aperiodic factor of (X_σ, S) given by a factor map of radius r . For all $n \geq 1$, we have the following.*

1. $(1/K_\sigma^2)\rho(M_\tau)^n \leq \langle \tau^n \rangle \leq |\tau^n| \leq K_\sigma^2\rho(M_\tau)^n$.
2. If $\rho(M_\sigma) = \rho(M_\tau)$, then $\max(|\sigma^n|, |\tau^n|) \leq K_\sigma^4 \min(\langle \sigma^n \rangle, \langle \tau^n \rangle)$.

Proof. (1) First notice that from Proposition 4.11 for all k and n one has $\langle \tau^n \rangle = \langle \tau^{(k)n} \rangle \leq |\tau^{(k)n}| = |\tau^n|$. Let μ be the unique ergodic measure of (Y, S) . Using the notation introduced before Lemma 4.14, from lemmas 4.13 and 4.14, one gets

$$\left[\frac{\min_i \delta_i^{(k)}}{\max_i \delta_i^{(k)}} \right] \rho(M_\tau)^n \leq \langle \tau^n \rangle \leq |\tau^n| \leq \left[\frac{\max_i \delta_i^{(k)}}{\min_i \delta_i^{(k)}} \right] \rho(M_\tau)^n.$$

Then, if we fix n whereas k goes to infinity, Item (5) of Proposition 5 allows to conclude.

- (2) It is a direct consequence of Item (1) and Corollary 4.15. □

5 Bounding the radius of a factor map between primitive substitution subshifts

In this section we show that for aperiodic primitive proper substitutions σ and τ , the radius of a factor map from X_σ to X_τ can be chosen smaller than a bound that only depends on σ and τ and that one can algorithmically compute.

5.1 Orbit-preserving maps, cocycles and dill maps

Let (X_1, T_1) and (X_2, T_2) be two topological dynamical systems. A map $f : X_1 \rightarrow X_2$ is said to be *orbit-preserving* if $f(\mathcal{O}_{T_1}(x)) \subset \mathcal{O}_{T_2}(f(x))$ holds for all $x \in X_1$, where $\mathcal{O}_{T_1}(x)$ stands for $\{T_1^n(x) \mid n \in \mathbb{Z}\}$. Observe that, in this case, f satisfies

$$f(T_1(x)) = T_2^{c(x)}(f(x)), \tag{5}$$

for all $x \in X$, for some map $c : X_1 \rightarrow \mathbb{Z}$ we call a *cocycle* for f . If f and c are continuous and if for all x , $c(x) \geq 0$ and there exists some y such that $c(y) > 0$, then we say f is a *dill map*. Observe that if f is a dill map, then the associated cocycle c is bounded. Furthermore, when (X_2, T_2) has no periodic point, f is a factor map if and only if $c(x) = 1$ for all x . Other examples of dill maps are given by morphisms $\Theta : A^* \rightarrow B^*$. Indeed, take a minimal subshift (X, S) generated by the sequence $x \in A^{\mathbb{Z}}$ and let (Y, S) be generated by $y = \Theta(x) \in B^{\mathbb{Z}}$. Then Θ defined a dill map from X to Y .

In the definition of dill maps, the existence of y such that $c(y) > 0$ is only required to avoid degenerated cases such as trivial maps satisfying $f(\mathcal{O}_{T_1}(x)) = \{f(x)\}$ for all x . In the rest of the paper, we will only consider non-trivial dill maps between minimal subshifts. In that situation, the next lemma shows that the existence y is just a consequence of the continuity of f .

Lemma 5.1. *Let (X_1, T_1) and (X_2, T_2) two topological dynamical systems such that (X_1, T_1) is minimal. If $f : X_1 \rightarrow X_2$ is a continuous orbit-preserving map such that $f(X_1)$ is not a singleton and the cocycle $c : X_1 \rightarrow \mathbb{Z}$ associated with f is continuous and non-negative, then f is a dill map.*

Proof. Let assume that $c(x) = 0$ for all $x \in X_1$. Then, $f(T_1(x)) = f(x)$ for all $x \in X_1$. Hence, $f(T_1^n(x)) = f(x)$ for all $n \in \mathbb{Z}$. Since $\mathcal{O}_{T_1}(x)$ is dense in X_1 and f is continuous, we get that $f(X_1)$ is a singleton, which contradicts the hypothesis. \square

The notion of dill map has been introduced by Salo and Törmä [ST15] in a one-sided context and they call a *nice cocycle* a cocycle associated with a non-trivial dill map.

Suppose (X, S) and (Y, S) are subshifts and $f : (X, S) \rightarrow (Y, S)$ is a dill map. A continuous function $\phi : X \rightarrow A^*$ satisfying

$$f(x) = \cdots \phi(S^{-1}(x)) \cdot \phi(x) \phi(S(x)) \phi(S^2(x)) \cdots \quad (6)$$

for all $x \in X$ is called an *implementation of f* . Observe that $\phi(x)$ could be the empty word.

Lemma 5.2. *[ST15, Lemma 8] Let (X, S) and (Y, S) be subshifts and $f : (X, S) \rightarrow (Y, S)$ a dill map with cocycle c . Then, f has an implementation ϕ given by*

$$\phi(x) = f(x)_{[0, c(x)-1]} \text{ for all } x \in X.$$

Notice that $|\phi(x)| = c(x)$ and that $\phi(x)$ is the empty word whenever $c(x) = 0$.

It is straightforward to check (as observed in [ST15]) that if Y does not contain S -periodic points then c and ϕ are unique. As we will deal with infinite minimal subshifts, there will be no S -periodic points in Y , and, thus, both maps will be unique.

Let $f : (X, S) \rightarrow (Y, S)$ be a dill map between subshifts without S -periodic points and with implementation map ϕ . We set

$$\begin{aligned} r &= \min\{R \in \mathbb{N} \mid x_{[-R, R]} = y_{[-R, R]} \implies \phi(x) = \phi(y)\} \text{ and} \\ m &= \max\{|\phi(x)| \mid x \in X\}. \end{aligned}$$

The couple (r, m) is called the *radius pair of f* . It is clear that r is a radius for ϕ and c . If f is a factor map, then its radius pair is $(r, 1)$ and ϕ is a sliding block code defining f .

Let f and g be two continuous maps from (X_1, T_1) to (X_2, T_2) . We say f and g are *orbitally related* (and we write $f \equiv g$) whenever there exist continuous maps $c, d : X_1 \rightarrow \mathbb{Z}$ satisfying, for all $x \in X_1$,

$$T_2^{c(x)} \circ f \circ T_1^{d(x)}(x) = g(x). \quad (7)$$

It is clear that the orbit relation is an equivalence relation.

Lemma 5.3. *Let f and g be two orbitally related factor maps from (X_1, T_1) to (X_2, T_2) . Suppose (X_1, T_1) is minimal and (X_2, T_2) is aperiodic. Then, there exists $e \in \mathbb{Z}$ such that $f = T_2^e \circ g$.*

Proof. Let $c, d : X_1 \rightarrow \mathbb{Z}$ be continuous maps satisfying, for all $x \in X_1$,

$$T_2^{c(x)} \circ f \circ T_1^{d(x)}(x) = g(x).$$

Thus, $T_2^{c(x)+d(x)} \circ f(x) = g(x)$. Therefore

$$T_2^{c(T_1(x))+d(T_1(x))} \circ f(T_1(x)) = g(T_1(x)) = T_2 \circ g(x) = T_2^{c(x)+d(x)} \circ f(T_1(x)).$$

As (X_2, T_2) is aperiodic, one gets $c(T_1(x)) + d(T_1(x)) = c(x) + d(x)$ for all x . From the minimality of (X_1, T_1) and by continuity of c and d , one concludes that $c + d$ is a constant function. \square

5.2 Specific dill maps associated with factor maps

Let $\sigma : A^* \rightarrow A^*$ and $\tau : B^* \rightarrow B^*$ be two aperiodic primitive substitutions. In this section, starting with a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$, we build pairwise orbitally related dill maps $F_n : (X_\sigma, S) \rightarrow (X_\tau, S)$, $n \in \mathbb{N}$. We show that for n large enough, these dill maps have bounded radius pairs and that there exist only finitely many such maps. In Section 5.3, we will use these dill maps to build a new factor map from (X_σ, S) to (X_τ, S) with a radius only depending on σ and τ , not on the radius of f .

Let us thus consider $f : (X_\sigma, S) \rightarrow (X_\tau, S)$ be a factor map with radius r . Let us recall a Cobham's type theorem proven in [Dur98b].

Theorem 5.4. *If (X_τ, S) is aperiodic then the dominant eigenvalues of σ and τ share a non trivial common power.*

Consequently, we can suppose that σ and τ have the same dominant eigenvalues (taking powers of the substitutions does not change the subshifts). As the dominant eigenvalue is greater than 1, we also may assume (by taking powers if needed) that it is at least equal to 2.

For all $n \geq 0$ we define the map $F_n : X_\sigma \rightarrow X_\tau$ by

$$\tau^n \circ F_n = S^{r_n \circ f \circ \sigma^n} \circ f \circ \sigma^n, \quad (8)$$

where $r_n : X_\tau \rightarrow \mathbb{N}$ is the return map to the clopen set $\tau^n(X_\tau)$ (see Corollary 4.3):

$$r_n(x) = \min\{i \geq 0 \mid S^i(x) \in \tau^n(X_\tau)\}.$$

Thus, we have $\tau^n \circ F_n \equiv f \circ \sigma^n$. Observe that, using Mossé's theorem (Theorem 4.1), r_n is continuous and F_n is well-defined.

Now let us show that F_n is a dill map and that, for n large enough, its radius pair is bounded by a computable bound only depending on σ and τ .

Proposition 5.5. *For all n , F_n is a dill map. Furthermore, if n is such that $r/|\tau^n| \leq 1$, the radius pair (s_1, s_2) of F_n satisfies*

$$s_1 \leq K_\sigma^4(2K_\sigma^4 + 2 + L_\tau) \quad \text{and} \quad s_2 \leq 2K_\sigma^4.$$

In particular, we have $\#\{F_n \mid r/|\tau^n| \leq 1\} \leq (\#B)^{3K_\sigma^9(2K_\sigma^4+2+L_\tau)}$.

Proof. Let n be a positive integer. By Corollary 4.3, F_n is a continuous map. We have :

$$\begin{aligned} \tau^n \circ F_n(S(x)) &= S^{r_n \circ f \circ \sigma^n(S(x))} \circ f \circ \sigma^n(S(x)) \\ &= S^{r_n \circ f \circ S^{|\sigma^n(x_0)|} \circ \sigma^n(x)} \circ f \circ S^{|\sigma^n(x_0)|} \circ \sigma^n(x) \\ &= S^{r_n \circ f \circ S^{|\sigma^n(x_0)|} \circ \sigma^n(x) + |\sigma^n(x_0)|} \circ f \circ \sigma^n(x) \\ &= S^{r_n \circ S^{|\sigma^n(x_0)|} \circ f \circ \sigma^n(x) + |\sigma^n(x_0)| - r_n \circ f \circ \sigma^n(x)} \circ \tau^n \circ F_n(x). \end{aligned} \tag{9}$$

The map

$$x \in X_\sigma \mapsto r_n \circ S^{|\sigma^n(x_0)|} \circ f \circ \sigma^n(x) + |\sigma^n(x_0)| - r_n \circ f \circ \sigma^n(x)$$

is continuous and $F_n(X_\sigma)$ is obviously not a singleton as (X_τ, S) is aperiodic. Therefore, from Corollary 4.4 and Lemma 5.1, to prove that F_n is a dill map it suffices to show that

$$r_n \circ S^{|\sigma^n(x_0)|} \circ f \circ \sigma^n(x) + |\sigma^n(x_0)| - r_n \circ f \circ \sigma^n(x)$$

is non-negative for all x . This is clear if $r_n \circ f \circ \sigma^n(x) \leq |\sigma^n(x_0)|$.

Suppose $r_n \circ f \circ \sigma^n(x) > |\sigma^n(x_0)|$. This means that $S^i \circ f \circ \sigma^n(x)$ does not belong to $\tau^n(X_\tau)$ for all $0 < i \leq |\sigma^n(x_0)|$. Hence $r_n \circ S^{|\sigma^n(x_0)|} \circ f \circ \sigma^n(x) + |\sigma^n(x_0)| = r_n \circ f \circ \sigma^n(x)$ and F_n is a dill map.

Now suppose that $n \in \mathbb{N}$ is such that $r/|\tau^n| \leq 1$. We let $c : X_\sigma \rightarrow \mathbb{N}$ denote the continuous map such that $F_n \circ S(x) = S^{c(x)} \circ F_n(x)$ for all x and ϕ be the implementation of F_n defined in Lemma 5.2. Let us also denote (s_1, s_2) the radius pair of F_n . From Equation (9) and Corollary 4.4, we get

$$\begin{aligned} c(x) &\leq \left(r_n \circ S^{|\sigma^n(x_0)|} \circ f \circ \sigma^n(x) + |\sigma^n(x_0)| - r_n \circ f \circ \sigma^n(x) \right) / \langle \tau^n \rangle \\ &\leq (|\tau^n| + |\sigma^n|) / \langle \tau^n \rangle. \end{aligned}$$

Thus from Corollary 4.15, Proposition 4.17 and the fact that $\rho(M_\tau) = \rho(M_\sigma)$ (see the observation below Theorem 5.4) we get $s_2 = \max\{c(x) \mid x \in X_\sigma\} \leq 2K_\sigma^4$.

Let $s = K_\sigma^4(2K_\sigma^4 + 2 + L_\tau)$ and let us show that $\phi(x)$ only depends on $x_{[-s, s]}$. This will imply that s_1 is at most equal to s . Consider $x, y \in X_\sigma$ such that $x_{[-s, s]} = y_{[-s, s]}$. Then $S^{r_n \circ f \circ \sigma^n} \circ f \circ \sigma^n(x)$ and $S^{r_n \circ f \circ \sigma^n} \circ f \circ \sigma^n(y)$ coincide on the indices $[-s \langle \sigma^n \rangle + r - k, s \langle \sigma^n \rangle - r - k]$ for some integer k satisfying $0 \leq k \leq |\tau^n|$. From Corollary 4.4, $F_n(x)$ and $F_n(y)$ coincide on the indices

$$\left[\frac{-s \langle \sigma^n \rangle + r - k + L_{\tau^n}}{|\tau^n|}, \frac{s \langle \sigma^n \rangle - r - k - L_{\tau^n}}{|\tau^n|} \right]$$

and, since $0 \leq k, r \leq |\tau^n|$, on the indices

$$\left[\frac{-s\langle\sigma^n\rangle + L_{\tau^n}}{|\tau^n|} + 1, \frac{s\langle\sigma^n\rangle - L_{\tau^n}}{|\tau^n|} - 2, \right],$$

where L_{τ^n} is a recognizability constant of τ^n . Recalling that $\rho(M_\tau) \geq 2$ and that $|\tau^n| \leq K_\sigma^2 \rho(M_\tau)^n$ by Proposition 4.17, we get from Proposition 4.8 that

$$L_{\tau^n} \leq K_\sigma^2 L_\tau \rho(M_\tau)^n.$$

Hence, $F_n(x)$ and $F_n(y)$ coincide on the indices

$$\left[\frac{-s\langle\sigma^n\rangle}{K_\sigma^2 \rho(M_\tau)^n} + L_\tau + 1, \frac{s\langle\sigma^n\rangle}{K_\sigma^2 \rho(M_\tau)^n} - L_\tau - 2 \right].$$

Then, it suffices to observe that, using Corollary 4.15 and the fact that $\rho(M_\sigma) = \rho(M_\tau)$,

$$\frac{s\langle\sigma^n\rangle}{K_\sigma^2 \rho(M_\tau)^n} - L_\tau - 2 \geq \frac{s}{K_\sigma^4} - 2 - L_\tau \geq s_2.$$

Similarly, we observe that

$$\frac{-s\langle\sigma^n\rangle}{K_\sigma^2 \rho(M_\tau)^n} + L_\tau + 1 \leq \frac{-s}{K_\sigma^4} + L_\tau + 1 \leq 0,$$

which shows that $s_1 \leq s$.

To conclude the proof, the number of possible dill maps F_n is at most equal to the number of maps from $\mathcal{L}_s(\sigma)$ to $\bigcup_{i=0}^{2K_\sigma^4} \mathcal{L}_i(\tau)$. We get, using Theorem 3.11,

$$\#\{F_n \mid r/|\tau^n| \leq 1\} \leq \left(\sum_{i=0}^{2K_\sigma^4} p_{X_\tau}(i) \right)^{K_\sigma s} \leq (\#B)^{3K_\sigma^9(2K_\sigma^4+2+L_\tau)}.$$

□

Corollary 5.6. *The set $\{F_n \mid n \in \mathbb{N}\}$ is finite. In particular, there exists an increasing sequence $(n_i)_{i \in \mathbb{N}}$ of integers such $F_{n_i} = F_{n_0}$ for all i .*

We finally show that all maps F_n are orbitally related.

Lemma 5.7. *If $F_n \equiv F_m$ for some $n \neq m$, then $F_{n-1} \equiv F_{m-1}$.*

Proof. Suppose $F_n \equiv F_m$. As $\tau^{n-1} \circ F_{n-1} \equiv f \circ \sigma^{n-1}$, one gets $\tau^{n-1} \circ F_{n-1} \circ \sigma \equiv f \circ \sigma^n \equiv \tau^n \circ F_n$. Thus, from Corollary 4.4, $F_{n-1} \circ \sigma \equiv \tau \circ F_n$. We also have $F_{m-1} \circ \sigma \equiv \tau \circ F_m$. Hence, $F_{n-1} \circ \sigma \equiv F_{m-1} \circ \sigma$ from which we deduce, using the definitions of F_{n-1} and F_{m-1} , that $F_{n-1} \equiv F_{m-1}$. □

5.3 Factorizations with smaller radius using dill maps

In this section, we show that if σ is proper and if there exists a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$, then there exists another factor map $g : (X_\sigma, S) \rightarrow (X_\tau, S)$ whose radius is bounded by some computable constant depending on σ and τ . Observe that if $\sigma : A^* \rightarrow A^*$ is proper and if $a, b \in A$ are such that $\sigma(A) \subset aAb$, then $x = \sigma^\omega(b \cdot a)$ is the unique admissible fixed point of σ and we have $\{x\} = \bigcap_{n \in \mathbb{N}} \sigma^n(X_\sigma)$.

Let (X, T) be a minimal dynamical system. A continuous function $c : X \rightarrow \mathbb{R}$ is a *coboundary* of (X, T) if there is some continuous function $d : X \rightarrow \mathbb{R}$ such that $c = d \circ T - d$.

Let $c : X \rightarrow \mathbb{R}$ be a continuous function. For $n \in \mathbb{N}$, we write

$$c^{(n)} = \begin{cases} 0 & \text{if } n = 0, \\ \sum_{k=0}^{n-1} c \circ T^k & \text{if } n > 0, \\ \sum_{k=n}^{-1} c \circ T^k & \text{if } n < 0, \end{cases} \quad (10)$$

We remind the following useful identity : $c^{(n+m)}(x) = c^{(n)}(x) + c^{(m)}(T^n(x))$ and recall the following well-known result of Gottschalk and Hedlund [GH55]. A proof can be found in [HK95] (Theorem 2.9.4 p. 102).

Theorem 5.8. *Let (X, T) be a minimal dynamical system and $c : X \rightarrow \mathbb{R}$ be a continuous function. The following are equivalent.*

1. c is a coboundary;
2. The sequence of functions $(c^{(n)})_{n \in \mathbb{N}}$ is uniformly bounded;
3. The sequence $(c^{(n)}(x))_{n \in \mathbb{N}}$ is bounded for some $x \in X$.

Let us show this result in a particular case but obtaining more information. Let (X, S) be a subshift and $c : X \rightarrow \mathbb{Z}$ be a continuous map of radius r defined by \hat{c} : c is constant, and equal to $\hat{c}(uv)$, on any cylinder $[u.v]$ where $|u| = r$ and $|v| = r + 1$. In the sequel we identify \hat{c} with the vector $(\hat{c}(uv))_{uv \in \mathcal{L}_{2r+1}(\sigma)}$.

The proof of the following lemma follows the lines of a proof of Theorem 5.8 given by Host [Hos95] for integer functions instead of real functions.

Lemma 5.9. *Let $\sigma : A^* \rightarrow A^*$ be a primitive, aperiodic and proper substitution, and let $x \in A^\mathbb{Z}$ be its admissible fixed point. Suppose that $c : X_\sigma \rightarrow \mathbb{Z}$ is a continuous map with radius r , defined by \hat{c} , such that $(c^{(n)})_n$ is bounded. We have the following.*

1. There exists $k > 0$ such that $\sum_{i=e_1}^{e_2-1} c(S^i(x)) = 0$ for all $e_1, e_2 \in E(\sigma^k, x)$, where $E(\sigma^k, x) = \{i \in \mathbb{Z} \mid S^i(x) \in \sigma^k(X_\sigma)\}$.
2. \hat{c} belongs to $\text{Ker}((M_{\sigma(r)}^t)^{p_\sigma(2r+1)})$, where M^t denotes the transposed matrix of M and p_σ denotes the factor complexity function of X_σ .
3. In (1), k can be chosen equal to $p_\sigma(2r + 1)$.

4. $c = d \circ S - d$ for some continuous map $d : X \rightarrow \mathbb{Z}$ having a radius bounded by

$$|\sigma^{p_\sigma(2r+1)}| + \max\{L_{\sigma^{p_\sigma(2r+1)}}, r\}$$

and such that

$$\max_{x \in X_\sigma} |d(x)| \leq |\sigma^{p_\sigma(2r+1)}| \max_{x \in X_\sigma} |c(x)|.$$

Proof. Let $U_k = \sigma^k(X_\sigma)$. Since the admissible fixed point x of σ satisfies $\{x\} = \bigcap_{k \in \mathbb{N}} U_k$, for any neighborhood V of x there exists k such that U_k is included in V . We set

$$\Lambda_k = \overline{\{c^{(n)}(x) \mid n \in \mathbb{N}, S^n(x) \in U_k\}} \text{ and } \Lambda = \bigcap_{k \in \mathbb{N}} \Lambda_k.$$

Obviously 0 belongs to Λ . Let us show that $\Lambda = \{0\}$. The map c being continuous and thus bounded, it is sufficient to prove that $2a$ belongs to Λ whenever a belongs to Λ .

Let $a \in \Lambda$ and $k \in \mathbb{N}$. There exists $n \geq 0$ such that $S^n(x)$ belongs to U_k and $c^{(n)}(x) = a$. Because S^n and $c^{(n)}$ are continuous, there exists a clopen set V included in U_k and containing x such that for all $y \in V$,

$$S^n(y) \in U_k \text{ and } c^{(n)}(y) = a.$$

Observe that there exists l such that U_l is included in V . Thus, one can suppose $V = U_l$. As a belongs to Λ_l , there exists $m \geq 0$ with $S^m(x)$ in U_l and $c^{(m)}(x) = a$. We get that $S^{n+m}(x)$ belongs to U_k and $c^{(n+m)}(x) = c^{(n)}(S^m(x)) + c^{(m)}(x) = 2a$. Thus $2a$ belongs to Λ_k . As this is true for all k , we get that $2a$ belongs to Λ .

Observe that, since c takes integer values and $\Lambda_{l+1} \subset \Lambda_l$ for all l , there exists k such that $\Lambda_l = \{0\}$ for all $l \geq k$. Thus (1) is proven.

Consider $\sigma^{(r)} : A^{(r)*} \rightarrow A^{(r)*}$ the substitution on the blocks of radius r defined in Section 4.2 and the notation used therein. Let e_1 and e_2 be two consecutive integers in $E(\sigma^k, x)$ and $uv \in \mathcal{L}(\sigma)$ with $|u| = r$ and $|v| = r + 1$. The number of occurrences of uv in $x_{[e_1-r, e_2-1+r]}$ is equal to the number of occurrences of (uv) in

$$W = (x_{[e_1-r, e_1+r]})(x_{[e_1-r+1, e_1+r+1]}) \cdots (x_{[e_2-1-r, e_2-1+r]}).$$

Then, as $c^{(e_1)}(x) = c^{(e_2)}(x) = 0$, one gets

$$0 = \sum_{i=e_1}^{e_2-1} c(S^i(x)) = \sum_{i=e_1}^{e_2-1} \hat{c}(x_{[i-r, i+r]}) \quad (11)$$

Observe that we can write

$$\sum_{i=e_1}^{e_2-1} \hat{c}(x_{[i-r, i+r]}) = \langle \hat{c}, \vec{W} \rangle, \quad (12)$$

where $\vec{W} = (|W|_w)_{w \in \mathcal{L}_{2r+1}(\sigma)}$, $\hat{c} = (\hat{c}(w))_{w \in \mathcal{L}_{2r+1}(\sigma)}$ and $\langle \cdot, \cdot \rangle$ stands for the usual scalar product.

From Proposition 4.11, we have

$$E(\sigma^k, x) = E(\sigma^{(r)k}, x^{(r)}).$$

Hence, e_1 and e_2 also belong to $E(\sigma^{(r)k}, x^{(r)})$ and $W = \sigma^{(r)k}(\beta)$ for some $\beta \in A^{(r)}$.

Let $\mathbf{1}_\beta$ be the column vector defined by $\mathbf{1}_\beta(\beta) = 1$ and 0 elsewhere. From the conjunction of (11) and (12) we obtain

$$0 = \langle \hat{c}, M_{\sigma^{(r)}}^k \mathbf{1}_\beta \rangle.$$

As this is true for all such e_1 and e_2 , it is true for all $\beta \in A^{(r)}$ and one gets $\hat{c}^t M_{\sigma^{(r)}}^k = 0$. The dimension of M_{σ^r} is $p_\sigma(2r+1) \times p_\sigma(2r+1)$, hence $\hat{c}^t M_{\sigma^{(r)}}^{p_\sigma(2r+1)} = 0$ and (2) is proven. Tracing back the previous arguments shows that for all $e_1, e_2 \in E(\sigma^{p_\sigma(2r+1)}, x)$ one gets

$$0 = \sum_{i=e_1}^{e_2-1} c(S^i(x))$$

and that k could be chosen to be $p_\sigma(2r+1)$. This proves (3) and we set $k = p_\sigma(2r+1)$.

Let us find a continuous map $d : X \rightarrow \mathbb{Z}$ such that $c = d \circ S - d$.

For all $n \in \mathbb{N}$, we set $d(S^n(x)) = c^{(n)}(x)$. On the positive orbit $\mathcal{O}_+(x) = \{S^n(x) \mid n \in \mathbb{N}\}$, we trivially have $c = d \circ S - d$. Let us show that d is uniformly continuous on $\mathcal{O}_+(x)$. By density of $\mathcal{O}_+(x)$ in X , the function will then uniquely extend to a continuous function d on X still satisfying $c = d \circ S - d$.

For all $n \in \mathbb{N}$, there exists $\ell \in \{0, 1, \dots, |\sigma^k|\}$ such that $S^{n+\ell}(x) \in U_k$. Observe that all functions $c^{(\ell)}$ and S^ℓ , $\ell \in \{0, 1, \dots, |\sigma^k|\}$, are uniformly continuous on X . Thus there exists $L > 0$ such that for all $m, n \in \mathbb{N}$ such that $S^m(x)$ and $S^n(x)$ coincide on the indices $[-L, L]$, there exists $\ell \in \{0, 1, \dots, |\sigma^k|\}$ such that $S^{\ell+m}(x)$ and $S^{\ell+n}(x)$ belong to U_k and $c^{(\ell)}(S^m(x)) = c^{(\ell)}(S^n(x))$. Observe that, as $S^{\ell+m}(x)$ and $S^{\ell+n}(x)$ belong to U_k , we have $c^{(m+\ell)}(x) = c^{(n+\ell)}(x) = 0$. We thus have

$$\begin{aligned} |d(S^m(x)) - d(S^n(x))| &= |c^{(m)}(x) - c^{(n)}(x)| \\ &= |c^{(m)}(x) + c^{(\ell)}(S^m(x)) - c^{(n)}(x) - c^{(\ell)}(S^n(x))| \\ &\leq |c^{(m+\ell)}(x)| + |c^{(n+\ell)}(x)|, \end{aligned}$$

which shows that d is uniformly continuous on $\mathcal{O}_+(x)$.

Let us end showing that the radius of d is at most $|\sigma^k| + \max\{L_{\sigma^k}, r\}$. We recall (see Corollary 4.5) that $\mathcal{P} = \{S^m \sigma^k([a]) \mid a \in A, 0 \leq m < |\sigma^k(a)|\}$ is a clopen partition of $X_{\sigma^k} = X_\sigma$.

Let $n \in \mathbb{N}$ with $n \geq \max_{a \in A} |\sigma^k(a)|$. We have $S^n(x) \in S^{m(S^n(x))} \sigma^k([a])$ for some $a \in A$ and some $m(S^n(x)) \in \{0, \dots, |\sigma^k(a)| - 1\}$. In particular, we have $n - m(S^n(x)) \geq 0$ and, since $S^{n-m(S^n(x))}(x) \in U_k$, $c^{(n-m(S^n(x)))}(x) = 0$. We get

$$d(S^n(x)) = c^{(n)}(x) = c^{(m(S^n(x)))}(S^{n-m(S^n(x))}(x)).$$

By continuity of d , for all $y \in X$, we have

$$d(y) = c^{(m(y))}(S^{-m(y)}y), \quad (13)$$

where $0 \leq m(y) < |\sigma^k(a)|$ for some $a \in A$ such that $y \in S^{m(y)} \sigma^k([a])$. By Corollary 4.5, the function $m : X \rightarrow \{0, \dots, |\sigma^k(a)|\}$ is continuous and has radius at most $L_{\sigma^k} + |\sigma^k|$, hence d has radius at most $|\sigma^k| + \max\{L_{\sigma^k}, r\}$. The inequality

$$\max_{x \in X_\sigma} |d(x)| \leq |\sigma^{p_\sigma(2r+1)}| \max_{x \in X_\sigma} |c(x)|$$

directly follows from (13). \square

Below we prove a more detailed version of Theorem 1.3 when the subshift are generated by substitutions and where σ is a proper substitution.

Theorem 5.10. *Let $\sigma : A^* \rightarrow A^*$ and $\tau : B^* \rightarrow B^*$ be two aperiodic and primitive substitutions, σ being proper. Suppose σ and τ have the same dominant eigenvalue. The following are equivalent.*

1. *There exists a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$.*

2. *There exists a dill map $F : (X_\sigma, S) \rightarrow (X_\tau, S)$ such that*

- *its radius (s_1, s_2) satisfies*

$$s_1 \leq K_\sigma^4(2K_\sigma^4 + 2 + L_\tau) \quad \text{and} \quad s_2 \leq 2K_\sigma^4;$$

- *for all $x \in X_\tau$, $F \circ S(x) = S^{c(x)} \circ F(x)$ where $c : X \rightarrow \mathbb{Z}$ is a continuous map satisfying that $(\sum_{k=0}^{n-1} (c \circ S^k - 1))_n$ is bounded.*

3. *There exists a factor map $g : (X_\sigma, S) \rightarrow (X_\tau, S)$ having a radius bounded by, with $R = K_\sigma^4(2K_\sigma^4 + 2 + L_\tau)$,*

$$2K_\sigma^4(K_\sigma + 1)(2R + 1)|\sigma^{p_\sigma(2R+1)}|$$

Moreover, once given f one can find g (as above) such that $f = S^n \circ g$ for some n .

Proof. Of course (3) implies (1). Let us first show that (1) implies (2). Recall that for all n , $F_n : X_\sigma \rightarrow X_\tau$ is the dill map defined by

$$\tau^n \circ F_n = S^{r_n \circ f \circ \sigma^n} \circ f \circ \sigma^n.$$

Let r be a radius of f and L_τ the constant of recognizability of τ . From Proposition 5.5, we have

$$\#\{F_n \mid r/|\tau^n| \leq 1\} \leq (\#B)^{3K_\sigma^9(2K_\sigma^4+2+L_\tau)}.$$

Thus there exist n and m , $n \geq m$, verifying $|\tau^{n-m}| \geq r$ and such that $F_n = F_m$. From Lemma 5.7 we obtain that $f \circ S = F_0 \equiv F_{n-m}$.

We fix $N = n - m$, $F = F_N$ and we let (s_1, s_2) be the radius pair of F . From Proposition 5.5 one has

$$s_1 \leq K_\sigma^4(2K_\sigma^4 + 2 + L_\tau) \quad \text{and} \quad s_2 \leq 2K_\sigma^4.$$

Let $c, d : X_\sigma \rightarrow \mathbb{Z}$ be the unique continuous maps verifying for all $x \in X_\sigma$,

$$F(Sx) = S^{c(x)}F(x) \quad \text{and} \quad F(Sx) = S^{d(x)}f(x).$$

One has $|c(x)| \leq s_2$ for all $x \in X_\sigma$. Furthermore, we have, for all x ,

$$F(Sx) = S^{d(x)+1}f(S^{-1}x) = S^{d(x)+1-d(S^{-1}(x))}F(x),$$

hence $c(x) = d(x) - d(S^{-1}(x)) + 1$. This proves (2).

Let us show that (2) implies (3). Observe that the radius R of c is at most $K_\sigma^4(2K_\sigma^4 + 2 + L_\tau)$. From Lemma 5.9, there exists a continuous map $D : X \rightarrow \mathbb{Z}$ with radius

$$H \leq |\sigma^{p_\sigma(2R+1)}| + \max\{L_{\sigma^{p_\sigma(2R+1)}}, R\}$$

such that $c - 1 = D \circ S - D$ and for all $x \in X_\sigma$,

$$|D(x)| \leq 2K_\sigma^4|\sigma^{p_\sigma(2R+1)}|.$$

Let $G : X \rightarrow Y$ be the continuous map defined by $G(x) = S^{-D(x)} \circ F(x)$. We have:

$$\begin{aligned} G \circ S(x) &= S^{-D(S(x))} \circ F(S(x)) = S^{-D(S(x))+c(x)-1+1} \circ F(x) \\ &= S^{-D(x)+1} \circ F(x) = S \circ G(x). \end{aligned}$$

Thus G is a factor map from (X_σ, S) to (X_τ, S) .

Let us give a bound on the radius of G . Let us set $M = 2K_\sigma^4|\sigma^{p_\sigma(2R+1)}|$. The radius of G is at most equal to N , where N is such that for all $x \in X_\sigma$, $x_{[-N, N]}$ uniquely determine $F(x)_{[-M, M]}$, i.e., for all $x, y \in X_\sigma$,

$$x_{[-N, N]} = y_{[-N, N]} \implies F(x)_{[-M, M]} = F(y)_{[-M, M]}.$$

Let ϕ be the implementation of F given by Lemma 5.2. For all n , we have

$$\begin{aligned} F(x)_{[0, c^{(n)}(x)-1]} &= \phi(x)\phi(Sx) \cdots \phi(S^{n-1}x), & \text{if } n > 0; \\ F(x)_{[c^{(n)}(x), -1]} &= \phi(S^n x)\phi(S^{n+1}x) \cdots \phi(S^{-1}x), & \text{if } n < 0. \end{aligned}$$

Furthermore, there is a local map $\hat{\phi} : A^{2s_1+1} \rightarrow B^*$ defining ϕ so that

$$\begin{aligned} F(x)_{[0, c^{(n)}(x)-1]} &= \prod_{i=0}^{n-1} \hat{\phi}(x_{[-s_1+i, s_1+i]}), & \text{if } n > 0; \\ F(x)_{[c^{(n)}(x), -1]} &= \prod_{i=0}^{n-1} \hat{\phi}(x_{[-s_1-n+i, s_1-n+i]}), & \text{if } n < 0. \end{aligned}$$

As F is a dill map, there is a word $u \in \mathcal{L}(\sigma)$ of length $2s_1+1$ such that $\hat{\phi}(u) \neq \epsilon$. By Theorem 3.11, u occurs in any word $v \in \mathcal{L}(\sigma)$ of length $(K_\sigma + 1)(2s_1 + 1)$. Therefore, u occurs at least M times both in

$$x_{[-s_1, s_1 + M(K_\sigma + 1)(2s_1 + 1)]} \quad \text{and in} \quad x_{[-s_1 - 1 - M(K_\sigma + 1)(2s_1 + 1), s_1 - 1]}.$$

The radius of G is thus at most equal to

$$\begin{aligned} N &\leq M(K_\sigma + 1)(2s_1 + 1) \\ &\leq 2K_\sigma^4(K_\sigma + 1)(2(K_\sigma^4(2K_\sigma^4 + 2 + L_\tau)) + 1)|\sigma^{p_\sigma(2R+1)}|, \end{aligned}$$

which ends the proof. \square

As a corollary we obtain Theorem 1.3. Observe that the hypothesis on the dominant eigenvalues is only needed to prove the implication (1) implies (2).

Let us consider an application of Theorem 1.3. When dealing with morphic sequences it is rather natural to ask whether it is a purely morphic sequence. Indeed, it is much easier to work directly with the endomorphism without to take care to some coding identifying letters. A lot, most, of the papers dealing with morphic sequences only consider the case of purely morphic sequences.

A first answer can be given in the non uniformly recurrent case. A well-known result of Pansiot [Pan84] asserts that the word complexity of purely morphic sequences is in $\Theta(1)$, $\Theta(n)$, $\Theta(n \log \log n)$, $\Theta(n \log n)$ or $\Theta(n^2)$, whereas there are morphic words with word complexity in $\Theta(n^{1+1/k})$ for any $k \in \mathbb{N} \setminus \{0\}$ [Pan85] (see also [Dev15]). Other examples of morphic sequences that are not purely morphic are given in [ACSZ18].

The argument of complexity cannot be used anymore in the uniformly recurrent case as the complexity of such morphic sequences is in $\Theta(n)$. Again in [ACSZ18], examples are given. Nevertheless they do not consider a dynamical framework and some of their examples (Example 23) presents uniformly recurrent morphic sequences that are not purely morphic but that clearly belongs to a subshift generated by a purely morphic sequence. Thus, one may ask whether there are minimal morphic subshifts (X, S) that are not purely morphic subshifts, that is $X \neq X_\sigma$ for any endomorphism σ . Of course, the same question but up to isomorphism has already been answered in Theorem 3.1. The following result provides an answer to this question.

Proposition 5.11. *Let (X, S) be an aperiodic linearly recurrent subshift for the constant K . There exist a factor map $\phi : (X, S) \rightarrow (Y, S)$ such that (Y, S) is not purely morphic.*

Proof. Let A be the alphabet of X . Since (X, S) is minimal, there exists an integer $r \geq 0$ such that for all $u \in \mathcal{L}_{2r+1}(X)$, there exists a return word $w \in \mathcal{R}_{X,u}$ of length bigger than $4K$. Let us build a sliding block code $\phi : A^{2r+1} \rightarrow B$ such that $(\phi(X), S)$ is aperiodic and not generated by a primitive substitution.

For all $u \in \mathcal{L}_{2r+1}(X)$, let $\phi_u : A^{2r+1} \rightarrow B$ be defined by

$$\phi_u(v) = \begin{cases} 1, & \text{if } v = u; \\ 0, & \text{otherwise.} \end{cases}$$

There exists $u \in \mathcal{L}_{2r+1}(X)$ such that $(\phi_u(X), S)$ is aperiodic. Indeed, if for all $u \in \mathcal{L}_{2r+1}(X)$, $(\phi_u(X), S)$ is periodic of period p_u , then (X, S) is periodic of period at most $\prod_{u \in \mathcal{L}_{2r+1}(X)} p_u$.

Consider $Y = \phi_u(X)$ such that (Y, S) is aperiodic. By the assumption on r , the word 0^{4K} belongs to $\mathcal{L}(Y)$. If Y is generated by a primitive substitution τ , then $(\tau^n(0))^{4K}$ belongs to $\mathcal{L}(Y)$ for all $n \in \mathbb{N}$. As τ is primitive, we have $|\tau^N(0)| \geq r$ for some N . Using Proposition 4.16, all words of $\mathcal{L}_{|\tau^N(0)|}(Y)$ occur in $(\tau^N(0))^{4K}$. As there are at most $|\tau^N(0)|$ words of length $|\tau^N(0)|$ in $(\tau^N(0))^{4K}$, Morse and Hedlund's Theorem implies that (Y, S) is periodic, which is a contradiction. \square

Example 5.12. Let (X, S) be the Thue-Morse subshift that is linearly recurrent for the constant 16 (see Example 3.14). Considering the word $u = b\nu^3(a)ba \in \mathcal{L}(X)$, we can check that the word $b\nu^4(aba)\nu^3(b)\nu^2(a)\nu(b)a$ is a return word to u in X and has length 80 and that the sliding block code $\phi_u : \{a, b\}^{19} \rightarrow \{0, 1\}$ defined by

$$\phi_u(v) = 1 \Leftrightarrow v = u$$

is such that $(\phi_u(X), S)$ is aperiodic. Indeed, it could be checked by hand (or using Walnut [Mou16]) that if x is the Thue-Morse sequence, then for any $n \in \mathbb{N}$, there is no positive integer p such that $x_{n+kp} = x_n$ for all $k \in \mathbb{N}$, which implies the aperiodicity of $(\phi_u(X), S)$. Since 0^{79} belongs to $\mathcal{L}(\phi_u(X))$, $(\phi_u(X), S)$ cannot be generated by a primitive substitution.

6 Decidability of the factorization in the case of periodic subshifts

In this section we consider Theorem 1.1, Corollary 1.2, Theorem 1.3 and Theorem 1.4 under the assumptions that (X, S) and, or, (Y, S) are periodic. As we are dealing with decision problems, it is important to recall that it is decidable to know whether a morphic sequence, and thus the subshift it generates, is periodic or ultimately periodic, or not, and that a period can be computed.

Theorem 6.1 ([Dur13a]). *Let x be a uniformly recurrent sequence. It is decidable whether x is periodic. Moreover some u such that $x = uuu \dots$ can be computed.*

Let us give the following notation that will facilitate the description of the different situations. Let (X, T) a minimal topological dynamical system and consider the set

$$P(X, T) = \{p \in \mathbb{N} \mid (X, T) \text{ admits a factor of cardinality } p\}.$$

Proof of Theorem 1.1 and Corollary 1.2 for the periodic case

If both (X, S) and (Y, S) are periodic with respective periods p and q then it suffices to test whether q divides p to decide whether (Y, S) is a factor of (X, S) . Suppose (Y, S) is periodic and (X, S) is not. Then the following result allows to conclude for the proofs of 1.1 and Corollary 1.2 in this periodic context.

Theorem 6.2. *[DG18] Let (X, S) be a uniformly recurrent morphic subshift and p be a positive integer. It is decidable to check whether p belongs to $P(X, S)$ or not.*

Theorem 1.3 is not true for factors that are periodic

In [Dur00] it is shown $P(X, S)$ might be infinite. For the Thue-Morse subshift for example one has $P(X, S) = \{n/2^k \mid k \geq 0\}$. This disproves Theorem 1.3 if we suppose (Y, S) is periodic.

Proof of Theorem 1.4 for the periodic case

As recalled above one can decide whether (Y, S) is periodic or not. Taking some block representation of (X, S) , one can suppose ϕ is a coding, see Section 4.2. This can be done algorithmically.

When (Y, S) is not periodic, we need Theorem 1.3 to prove Theorem 1.4. If we suppose (Y, S) periodic we cannot use Theorem 1.3 anymore as explained above.

Nevertheless Theorem 1.4 remains true in this context. Suppose (Y, S) is periodic. Then one can compute some word v such that Y is generated by the sequence $y = \cdots vvv \cdots$. To decide whether ϕ is a factor map from (X, S) to (Y, S) it suffices to check whether $\phi(x)$ is equal to $S^k(y)$ for some $k \in [1, |v|]$. It is clearly decidable using Theorem 6.1.

7 Decidability of the factorization

In this section we prove that the factorization and isomorphism problems between minimal morphic subshifts are decidable, that is Theorem 1.1 and Corollary 1.2. Let us sketch briefly how we proceed.

From Theorem 3.1, and in particular Remark 3.2, it is enough to consider primitive substitution subshifts. Let σ and τ be two primitive substitutions.

In Section 5, we have shown there is a computable constant R such that if there is a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$, then there is another factor map $F : (X_\sigma, S) \rightarrow (X_\tau, S)$ with radius at most R . In this section, we show that

there is an algorithm that answers whether or not a given sliding block code $f : A^{2R+1} \rightarrow B$ defines a factor map from (X_σ, S) to (X_τ, S) . Using Lemma 4.9 and Proposition 4.11, we can assume that f has radius 0, i.e., $f : A \rightarrow B$ is a coding. As a subshift is uniquely determined by its language, we want to check whether or not $f(\mathcal{L}(\sigma)) = \mathcal{L}(\tau)$. If x is an admissible fixed point of σ , as well as y for τ , all we have to check is that $\mathcal{L}(f(x)) = \mathcal{L}(y)$.

To this end, we will deeply use return words. Roughly speaking, we will show that we have $\mathcal{L}(f(x)) = \mathcal{L}(y)$ if, and only if, there exists a prefix u of x , whose length is bounded by a computable constant, such that the return words to $f(u)$ in y and the return words to $f^{-1}(\{f(u)\})$ in x (as defined in Section 7.1) are identically concatenated in y and in x respectively. A key argument is to find some ‘‘canonical morphisms’’, i.e., morphisms that only depend on the combinatorics of x and y and not on the substitutions σ and τ . One of the main steps for this is to extend the results of Section 3.1 to return words to sets of words instead of words.

7.1 Return words to a set of words

Definition 7.1. Let $x \in A^{\mathbb{N}}$ be a sequence and let $U \subset \mathcal{L}(x)$ be a set of non-empty words of the same length. We call *return word to U* any non-empty word $w_{[i,j-1]} = w_i w_{i+1} \cdots w_{j-1}$ for which there exist $u, v \in U$ such that wv is a word in $\mathcal{L}(x)$ that admits u as a prefix and that contains no other occurrences of a word in U . The set of return words to U in x is denoted by $\mathcal{R}_{x,U}$.

With such a definition, we can define a bijection $\Theta_{x,U}$ as in the case of return words to a single word. We consider the set $R_{x,U} = \{1, 2, \dots\}$ with $\#R_{x,U} = \#\mathcal{R}_{x,U}$ and we define

$$\Theta_{x,U} : R_{x,U}^* \rightarrow \mathcal{R}_{x,U}^*$$

as the unique morphism that maps $R_{x,U}$ bijectively onto $\mathcal{R}_{x,U}$ and such that for all $i \in R_{x,U}$, $\Theta_{x,U}(i)$ is the i th word of $\mathcal{R}_{x,U}$ occurring in x . Then Proposition 3.7 can be generalized as follows.

Example 7.2. Considering the Thue-Morse sequence y and the set of words $U = \{aa, bb\}$, we have $\mathcal{R}_{y,U} = \{aa, bb, aaba, bbab\}$ and

$$\Theta_{y,U} : \begin{cases} 1 \mapsto bbab \\ 2 \mapsto aa \\ 3 \mapsto bb \\ 4 \mapsto aaba \end{cases} .$$

Similarly, for the Fibonacci sequence x and the set $U = \{aa, ab\}$, we have $\mathcal{R}_{x,U} = \{a, ab\}$ and

$$\Theta_{x,U} : \begin{cases} 1 \mapsto ab \\ 2 \mapsto a \end{cases} .$$

Proposition 7.3. *Let x be a sequence in $A^{\mathbb{N}}$ and let U be a set of non-empty words in $\mathcal{L}(x)$ with the same length. Suppose that x has infinitely many occurrences of elements of U and let i be the first occurrence of a word of U in x . Then,*

1. *There exists a unique sequence $(w_n)_n \in \mathcal{R}_{x,U}^{\mathbb{N}}$ such that for all n , $S^i(x)$ has a prefix in $w_0w_1 \cdots w_nU$;*
2. *There exists a unique sequence $z \in \mathcal{R}_{x,U}^{\mathbb{N}}$ satisfying $\Theta_{x,U}(z) = S^i(x)$;*

Proof. We only prove Item (1), Item (2) being a direct consequence of it. The existence of a sequence $(w_n)_n \in \mathcal{R}_{x,U}^{\mathbb{N}}$ satisfying the statement is obvious so let us prove the uniqueness.

Suppose that $(w_n)_n$ and $(w'_n)_n$ are two sequences in $\mathcal{R}_{x,U}^{\mathbb{N}}$ such that $S^i(x)$ has a prefix in $w_0w_1 \cdots w_nU$ for all n . Let $n \in \mathbb{N}$ be the smallest integer such that $w_n \neq w'_n$ and let $u, v, u', v' \in U$ be words such that $S^i(x)$ has for prefixes $w_0 \cdots w_{n-1}u$, $w_0 \cdots w_{n-1}v$, $w'_0 \cdots w'_{n-1}u'$ and $w'_0 \cdots w'_{n-1}v'$. As $w_i = w'_i$ for all $0 \leq i \leq n-1$, $w_n \neq w'_n$ and $w_0 \cdots w_{n-1}w_n$ and $w_0 \cdots w_{n-1}w'_n$ are prefixes of $S^i(x)$, we necessarily have $|w_n| \neq |w'_n|$. Assuming that $|w_n| < |w'_n|$, we reach a contradiction with the definition of return words: the word $w'_n v'$ contains three occurrences of words in U : u' as a prefix v' as a suffix and v . \square

Remark 7.4. The previous proposition shows that $S^i(x)$ can be uniquely decomposed into a sequence $(w_n)_n$ of return words to U . Furthermore, since all words in U have the same length, we have that for all n , there is a unique word u_n such that $w_0 \cdots w_n u_n$ is a prefix of $S^i(x)$. However, at a “local scale” it can happen that for two different integers m and n , we have $w_m = w_n$ but $u_m \neq u_n$. Thus in some sense such a return word w_m represents several return words (because it returns to several words of U). On the opposite way, if w is a return word to U , there may exist $u \in U$ such that wu does not have any prefix in U . This is illustrated on the next example.

Example 7.5. Continuing Example 7.2, we see that, with the Thue-Morse sequence, not every word wu with $w \in \mathcal{R}_{y,U}$ and $u \in U$ belongs to $\mathcal{L}(y)$. For instance, we have $bbbb \notin \mathcal{L}(y)$ although $bb \in \mathcal{R}_{y,U} \cap U$. On the other side with the Fibonacci sequence, the return word ab returns to different words in U since $abaa$ and $abab$ both belong to $\mathcal{L}(x)$.

Let us recall that our aim is to prove a result similar to Proposition 3.8, but for return words to sets of words. In the case where U is a singleton $\{u\}$, a key argument in the proof of Proposition 3.8 is that $\mathcal{R}_{x,u}$ is a code (see Proposition 3.6). This is in particular needed for the morphism $\lambda_{x,u,u'}$ of that proposition to be properly defined. Unfortunately, this property is not true anymore when U is not a singleton. Indeed, in [LR13], the authors exhibit an admissible fixed point x of a primitive substitution for which, if we take U_n to be the set of left special words of length n , we have $\{a, b, ab\} \subset \mathcal{R}_{x,U_n}$ for infinitely many values of n . We thus introduce the following notion (see also [Dur13a]).

Definition 7.6. Let $x \in A^{\mathbb{N}}$ be a sequence and let $U \subset \mathcal{L}(x)$ be a set of non-empty words with the same length. A *return pair* to U is a pair $(w, u) \in \mathcal{R}_{x,U} \times U$ such that wu belongs to $\mathcal{L}(x)$. We let $\tilde{\mathcal{R}}_{x,U}$ denote the set of return pairs to U in x .

Using these return pairs, we again enumerate the elements (w, u) in $\tilde{\mathcal{R}}_{x,U}$ in the order of the first occurrence of wu in x . Formally, we consider the set $\tilde{R}_{x,U} = \{1, 2, \dots\}$ with $\#\tilde{R}_{x,U} = \#\tilde{\mathcal{R}}_{x,U}$ and we define

$$\tilde{\Theta}_{x,U} : \tilde{R}_{x,U}^* \rightarrow \tilde{\mathcal{R}}_{x,U}^*$$

as the unique morphism that maps bijectively $\tilde{R}_{x,U}$ onto $\tilde{\mathcal{R}}_{x,U}$ and for which for all $i \in \tilde{R}_{x,U}$, the element $\tilde{\Theta}_{x,U}(i) = (w, u)$ is such that wu is the i th such word occurring in x . Observe that we should only consider words $(w_1, u_1) \cdots (w_n, u_n) \in \tilde{\mathcal{R}}_{x,U}^*$ that correspond to real concatenations of return words. More precisely, we say that $(w_1, u_1) \cdots (w_n, u_n) \in \tilde{\mathcal{R}}_{x,U}^*$ is *admissible* if

1. $w_1 w_2 \cdots w_n u_n$ is a factor of x ;
2. for all $i \in \{1, 2, \dots, n-1\}$, u_i is a prefix of $w_{i+1} w_{i+2} \cdots w_n u_n$.

By extension, a sequence $(w_1, u_1)(w_2, u_2)(w_3, u_3) \cdots \in \tilde{\mathcal{R}}_{x,U}^{\mathbb{N}}$ is admissible if so is each of its prefixes. By symmetry, we also say that a finite or infinite word w on $\tilde{R}_{x,U}$ is admissible if so is $\tilde{\Theta}_{x,U}(w)$.

Example 7.7. We continue our running examples with the Thue-Morse sequence and the Fibonacci sequence. For the Thue-Morse sequence, we have $\tilde{\mathcal{R}}_{y,U} = \{(aa, bb), (bb, aa), (aaba, bb), (bbab, aa)\}$ and

$$\tilde{\Theta}_{y,U} : \begin{cases} 1 \mapsto (bbab, aa) \\ 2 \mapsto (aa, bb) \\ 3 \mapsto (bb, aa) \\ 4 \mapsto (aaba, bb) \end{cases} .$$

Observe that the word $(aa, bb)(bb, aa)(aaba, bb)$ is admissible (because the word $abbaababb$ occurs in y), but $(aa, bb)(bb, aa)(aa, bb)$ is not.

For the Fibonacci sequence, we have $\tilde{\mathcal{R}}_{x,U} = \{(ab, aa), (a, ab), (ab, ab)\}$ and

$$\tilde{\Theta}_{x,U} : \begin{cases} 1 \mapsto (ab, aa) \\ 2 \mapsto (a, ab) \\ 3 \mapsto (ab, ab) \end{cases} .$$

This notion of return pairs satisfies a generalization of Proposition 3.6.

Proposition 7.8. *Let x be a sequence in $A^{\mathbb{N}}$ and let $U \subset \mathcal{L}(x)$ be a set of non-empty words with the same length.*

1. If $(w_1, u_1) \cdots (w_k, u_k) \in \tilde{\mathcal{R}}_{x,U}^*$ is admissible and if $(r, v) \in \tilde{\mathcal{R}}_{x,U}$ is such that rv occurs in $w_1 w_2 \cdots w_k u_k$, then $(r, v) = (w_i, u_i)$ for some $i \in \{1, \dots, k\}$.
2. If two admissible words $(w_1, u_1) \cdots (w_k, u_k)$ and $(r_1, v_1) \cdots (r_\ell, v_\ell)$ in $\tilde{\mathcal{R}}_{x,U}^*$ are such that $w_1 w_2 \cdots w_k u_k = r_1 r_2 \cdots r_\ell v_\ell$, then $k = \ell$ and for all $i \in \{1, \dots, k\}$, $(w_i, u_i) = (r_i, v_i)$.

Proof. Let us prove (1). Let $i \leq k$ be the largest integer such that rv occurs in $w_i \cdots w_k u_k$. If $(r, v) \neq (w_i, u_i)$, there exists a non-empty word p such that prv is a prefix of $w_i \cdots w_k u_k$. Furthermore, by choice of i , we have $|p| < |w_i|$. Let v' denote the prefix of rv which is in U . If $|pr| \leq |w_i|$, then $w_i u_i$ contains at least three occurrences of words of U : u_{i-1} , v' and u_i , where u_0 is the prefix of $w_1 u_1$ which is in U . This contradicts the fact that (w_i, u_i) is a return pair. If $|pr| > |w_i|$, then we get an analogous contradiction with rv containing occurrences of v' , u_i and v . In all cases, we get a contradiction, so we have $(r, v) = (w_i, u_i)$.

Let us now prove (2). The proof goes by induction on k and uses the same reasoning as for (1). Assume that $k = 1$ and let u' be the prefix of $w_1 u_1$ which is in U . Since $w_1 u_1 = r_1 r_2 \cdots r_\ell v_\ell$, all words u, r_1, \dots, r_ℓ occur in $w_1 u_1$. By definition of a return pair, we must have $\ell = 1$. As $|u_1| = |v_1|$, this shows that $(w_1, u_1) = (r_1, v_1)$.

Assume that the result is true for $k \geq 1$ and let us prove it for $k+1$. With the same reasoning as for the case $k = 1$, we must have $\ell > 1$. Since $|u_{k+1}| = |v_\ell|$, we again have $u_{k+1} = v_\ell$. If $w_{k+1} \neq r_\ell$, then since $w_1 \cdots w_{k+1} = r_1 \cdots r_\ell$, we have $|w_{k+1}| \neq |r_\ell|$. Let us consider the case $|w_{k+1}| < |r_\ell|$, the other one being symmetric. Then $r_\ell v_\ell$ contains at least three occurrences of words of U : $v_{\ell-1}$ has a prefix, v_ℓ as a suffix and u_k . This contradicts the definition of a return pair, so we have $(w_{k+1}, u_{k+1}) = (r_\ell, v_\ell)$, hence $w_1 \cdots w_k u_k = r_1 \cdots r_{\ell-1} v_{\ell-1}$, which ends the proof. \square

Let $\delta_{x,U,1} : \tilde{\mathcal{R}}_{x,U}^* \rightarrow \mathcal{R}_{x,U}^*$ and $\delta_{x,U,2} : \tilde{\mathcal{R}}_{x,U}^* \rightarrow U^*$ be respectively defined

$$\delta_{x,U,1}(w, u) = w \quad \text{and} \quad \delta_{x,U,2}(w, u) = u.$$

Let also $d_{x,U,1} : \tilde{\mathcal{R}}_{x,U}^* \rightarrow \mathcal{R}_{x,U}^*$ and $d_{x,U,2} : \tilde{\mathcal{R}}_{x,U}^* \rightarrow \mathcal{R}_{x,U}^*$ be respectively defined by

$$d_{x,U,1}(i) = \delta_{x,U,1}(\tilde{\Theta}_{x,U}(i)) \quad \text{and} \quad d_{x,U,2}(i) = \delta_{x,U,2}(\tilde{\Theta}_{x,U}(i)).$$

We get the following generalization of Proposition 3.7 whose proof is similar to the proof of Proposition 7.3 and left to the reader.

Proposition 7.9. *Let x be a sequence in $A^{\mathbb{N}}$ and let $U \subset \mathcal{L}(x)$ be a set of non-empty words with the same length. Suppose that x has infinitely many occurrences of elements of U and let $i \in \mathbb{N}$ be the first occurrence of a word of U in x . Then,*

1. *There exists a unique admissible sequence $(w_n, u_n)_n \in \tilde{\mathcal{R}}_{x,U}^{\mathbb{N}}$ such that for all n , the word $w_0 w_1 \cdots w_n u_n$ is a prefix of $S^i(x)$;*

2. There exists a unique admissible sequence $z \in \tilde{R}_{x,U}^{\mathbb{N}}$ satisfying $d_{x,U,1}(z) = S^i(x)$.

Definition 7.10. The sequence $z \in \tilde{R}_{x,U}^{\mathbb{N}}$ of the previous proposition is called a *derived sequence* of x (w.r.t. U). We denote it by $\mathcal{D}_U(x)$.

Remark 7.11. Observe that when U is a singleton $\{u\}$, we have $\tilde{\mathcal{R}}_{x,\{u\}} = \mathcal{R}_{x,u} \times \{u\}$ and $\mathcal{D}_{\{u\}}(x) = \mathcal{D}_u(x)$.

The next result directly follows from Proposition 7.8.

Corollary 7.12. Let $x \in A^{\mathbb{N}}$ be uniformly recurrent and let $U \subset \mathcal{L}(x)$ be a set of non-empty words with the same length. Then the derived sequence $\mathcal{D}_U(x)$ is uniformly recurrent.

We can now state a result analogous to Proposition 3.8. We first independently define the morphism $\lambda_{x,U,V}$ in a more general setting.

Lemma 7.13. Let x be a uniformly recurrent sequence in $A^{\mathbb{N}}$. Let U and V be subsets of $\mathcal{L}(x)$ such that all words of U have the same length $\ell_U \geq 1$ and all words of V have the same length $\ell_V > \ell_U$. Assume that any word of V has a prefix in U . There exists a unique morphism $\lambda_{x,U,V} : \tilde{R}_{x,V}^* \rightarrow \tilde{R}_{x,U}^*$ such that

$$\begin{cases} d_{x,U,1} \circ \lambda_{x,U,V} = d_{x,V,1}; \\ d_{x,U,2}(\lambda_{x,U,V}(i)_{|\lambda_{x,U,V}(i)|-1}) = (d_{x,V,2}(i))_{[0,\ell_U-1]}, \quad \text{for all } i \in \tilde{R}_{x,U}. \end{cases} \quad (14)$$

In particular, we have $\lambda_{x,U,V}(\mathcal{D}_V(x)) = S^k \mathcal{D}_U(x)$, where $k \geq 0$ is the smallest integer such that $d_{x,U,1}(S^k \mathcal{D}_U(x))$ has a prefix in V .

Proof. Let us first show that the morphism $\lambda_{x,U,V}$ is well defined. Let $i \in \tilde{R}_{x,V}$ and let us call $w \in \mathcal{R}_{x,V}$ and $v', v \in V$ the words such that $\tilde{\Theta}_{x,V}(i) = (w, v)$ and v' is a prefix of wv . As any word in V has a prefix in U , there exists an admissible word $(w_1, u_1) \cdots (w_k, u_k) \in \tilde{\mathcal{R}}_{x,U}^*$ such that $wv_{[0,\ell_U-1]} = w_1 w_2 \cdots w_k u_k$. By Proposition 7.8, this admissible word is unique. We thus define $\lambda_{x,U,V}(i)$ as the unique word $i_1 \cdots i_k \in \tilde{R}_{x,U}^*$ such that $\tilde{\Theta}_{x,U}(i_j) = (w_j, u_j)$ for all $1 \leq j \leq k$. Observe that this definition is equivalent to (14).

Now let us prove that $\lambda_{x,U,V}(\mathcal{D}_V(x)) = S^k \mathcal{D}_U(x)$. By Proposition 7.8, $\mathcal{D}_U(x)$ is the unique sequence $z \in \tilde{R}_{x,U}$ satisfying $d_{x,U,1}(z) = S^i(x)$, where i is the first occurrence in x of a word of U . Similarly, the sequence $\mathcal{D}_V(x)$ is such that $d_{x,V,1}(\mathcal{D}_V(x)) = S^j(x)$, where j is the first occurrence in x of a word of V . The result thus directly follows from (14). \square

Proposition 7.14. Let x be a linearly recurrent sequence for the constant K . Let U and V be subsets of $\mathcal{L}(x)$ such that all words of U have the same length $\ell_U \geq 1$ and all words of V have the same length $\ell_V \geq 1$. Assume the following:

1. $\ell_V \geq K(K+1)\ell_U$;
2. any word of V has a prefix in U ;

3. any return word to V has length at least ℓ_V/K ;
4. $\mathcal{D}_U(x) = \mathcal{D}_V(x)$.

Then the morphism $\lambda_{x,U,V}$ is a primitive substitution and $X_{\lambda_{x,U,V}} = \Omega(\mathcal{D}_U(x))$.

Proof. Let us show that $\lambda_{x,U,V}$ is primitive, that is: for all $(i, j) \in \tilde{R}_{x,V} \times \tilde{R}_{x,U}$, j occurs in $\lambda_{x,U,V}(i)$. Let $\tilde{\Theta}_{x,U}(j) = (w, u) \in \tilde{R}_{x,U}$ be a return pair to U . We have $|w| \leq \max\{|r| \mid r \in \mathcal{R}_{x,u}\}$. Hence, by Theorem 3.11, we have $|wu| \leq (K+1)\ell_U$ and wu occurs in any factor of length $(K+1)^2\ell_U$ of x . Since any return pair $(r, v) \in \tilde{R}_{x,V}$ is such that $|rv| \geq \ell_V \frac{K+1}{K} \geq (K+1)^2\ell_U$, we deduce from Theorem 3.11 that j occurs in $\lambda_{x,U,V}(i)$ for all $i \in \tilde{R}_{x,V}$.

With Lemma 7.13 one deduces that $\lambda_{x,U,V}(\mathcal{D}_V(x)) = S^k(\mathcal{D}_V(x))$ for some k , which achieves the proof. \square

7.2 Some sufficient conditions for $f : A \rightarrow B$ to be a factor map

Let σ and τ be two primitive substitutions. In this section, we give some sufficient conditions for $f : A \rightarrow B$ to be a factor map from (X_σ, S) to (X_τ, S) . These conditions are defined on pairs of words $(u, v) \in \mathcal{L}(\tau) \times \mathcal{L}(\tau)$. In the next sections, we will show that they are algorithmically checkable and that if $f : A \rightarrow B$ really defines a factor map, then there must exist such a pair (u, v) where u, v have length bounded by a computable constant.

Theorem 7.15. *Let $\sigma : A^* \rightarrow A^*$ and $\tau : B^* \rightarrow B^*$ be primitive substitutions and let $x \in A^{\mathbb{N}}$ and $y \in B^{\mathbb{N}}$ be fixed points of σ and τ respectively. Let K_σ and K_τ denote the respective constants of linear recurrence of x and y and let $K = \max\{K_\sigma, K_\tau\}$. Let $f : A \rightarrow B$ be a coding. If there exist non-empty words $u, v \in \mathcal{L}(\tau)$ such that*

1. u is a prefix of v such that $|v| \geq K(K+1)|u|$,
2. $u, v \in f(\mathcal{L}(\sigma))$;
3. $\Omega(\mathcal{D}_u(y)) = \Omega(\mathcal{D}_v(y))$,
4. $\Omega(\mathcal{D}_{f^{-1}(\{u\})}(x)) = \Omega(\mathcal{D}_{f^{-1}(\{v\})}(x))$,

then f defines a factor map from (X_σ, S) to (X_τ, S) .

Proof. By minimality, it suffices to show that $f(\mathcal{L}(\sigma)) \subset \mathcal{L}(\tau)$. Let us consider the sets $U = f^{-1}(\{u\})$ and $V = f^{-1}(\{v\})$. Since for any return word w to V in x , $f(w)$ is a return word to v in y and $|w| = |f(w)|$, we have from Theorem 3.11 that $|w| \geq |v|/K$. We are thus in the conditions of Proposition 3.8 and Proposition 7.14 and we can define the primitive morphisms $\lambda_{y,u,v}$ and $\lambda_{x,U,V}$ such that $X_{\lambda_{y,u,v}} = \Omega(\mathcal{D}_u(y))$ and $X_{\lambda_{x,U,V}} = \Omega(\mathcal{D}_U(x))$.

We let $\varphi_u : \tilde{R}_{x,U}^* \rightarrow R_{y,u}^*$ and $\varphi_v : \tilde{R}_{x,V}^* \rightarrow R_{y,v}^*$ denote the unique morphisms satisfying

$$\begin{aligned} f \circ d_{x,U,1} &= \Theta_{y,u} \circ \varphi_u, \\ f \circ d_{x,V,1} &= \Theta_{y,v} \circ \varphi_v. \end{aligned}$$

By construction, the morphisms φ_u and φ_v are codings. Furthermore, using Proposition 3.8 and Equation (14) and the two equalities above, we get

$$\begin{aligned} \Theta_{y,u} \circ \lambda_{y,u,v} \circ \varphi_v &= \Theta_{y,v} \circ \varphi_v \\ &= f \circ d_{x,V,1} \\ &= f \circ d_{x,U,1} \circ \lambda_{x,U,V} \\ &= \Theta_{y,u} \circ \varphi_u \circ \lambda_{x,U,V}. \end{aligned}$$

Using Proposition 3.6, we then obtain

$$\lambda_{y,u,v} \circ \varphi_v = \varphi_u \circ \lambda_{x,U,V}.$$

By definition of return words, we have

$$\begin{aligned} \mathcal{L}(\sigma) &= \mathcal{L}(d_{x,U,1}(\mathcal{D}_U(x))) = \mathcal{L}(d_{x,U,1}(\mathcal{L}(\lambda_{x,U,V}))) \\ \mathcal{L}(\tau) &= \mathcal{L}(\Theta_{y,u}(\mathcal{D}_u(y))) = \mathcal{L}(\Theta_{y,u}(\mathcal{L}(\lambda_{y,u,v}))) \end{aligned}$$

Let w be a word in $\mathcal{L}(\sigma)$. By primitiveness, there is a positive integer n such that w occurs in $d_{x,U,1}(\lambda_{x,U,V}^n(1))$. Thus $f(w)$ occurs in the word

$$\begin{aligned} f(d_{x,U,1}(\lambda_{x,U,V}^n(1))) &= \Theta_{y,u}(\varphi_u(\lambda_{x,U,V}^n(1))) \\ &= \Theta_{y,u}(\lambda_{y,u,v}^n(\varphi_v(1))), \end{aligned}$$

which belongs to the set $\mathcal{L}(\Theta_{y,u}(\mathcal{L}(\lambda_{y,u,v}))) = \mathcal{L}(\tau)$ and this concludes the proof. \square

7.3 The sufficient conditions of Theorem 7.15 are necessary and algorithmically checkable

The aim of this section is to show that for $f : A \rightarrow B$ to be a factor map from (X_σ, S) to (X_τ, S) , there must exist some words u, v of bounded length (with a computable bound) that satisfy the conditions of Theorem 7.15. We also show that the conditions (3) and (4) are algorithmically checkable. This can be achieved through return substitutions.

7.3.1 Return substitutions for a single word

We first recall the classical notion of return substitution that has been introduced in [Dur98a]. Observe that the notion that we develop here are closely related to the discussion made in Section 3.1. We decide to introduce them in this section because we want to extend them to the more general notion of return words to a set of words.

Proposition 7.16 ([Dur98a]). *Let σ be an aperiodic primitive substitution, $x \in A^{\mathbb{N}}$ be a fixed point of σ and u be a non-empty prefix of x . The sequence $\mathcal{D}_u(x)$ is the admissible fixed point starting with 1 of the primitive substitution $\sigma_{x,u} : R_{x,u}^* \rightarrow R_{x,u}^*$ defined by*

$$\Theta_{x,u} \circ \sigma_{x,u} = \sigma \circ \Theta_{x,u}. \quad (15)$$

Furthermore, if K is the constant of linear recurrence of x , then $\mathcal{D}_u(x)$ is linearly recurrent for the constant K^3 .

Definition 7.17. The morphism $\sigma_{x,u}$ of the previous proposition is called a *return substitution* (w.r.t. u).

From Theorem 3.11 and the definition of $\sigma_{x,u}$ we deduce the following corollary.

Corollary 7.18. *Let σ be an aperiodic primitive substitution and $x \in A^{\mathbb{N}}$ be a fixed point of σ . We have*

$$\text{Card}(\{\sigma_{x,x_{[0,n]}} \mid n \in \mathbb{N}\}) \leq Q_\sigma = (1 + (K_\sigma + 1)^3)^{|\sigma|K_\sigma^2(1+(K_\sigma+1)^3)}.$$

The following lemma is clear from [Dur98a]. We briefly present the algorithm for the sake of completeness.

Lemma 7.19. *Let σ be an aperiodic primitive substitution, $x \in A^{\mathbb{N}}$ be a fixed point of σ and u be a non-empty prefix of x . The morphisms $\sigma_{x,u}$ and $\Theta_{x,u}$ are algorithmically computable.*

Sketch of the algorithm. Since x is a fixed point of σ , the word u is a prefix of $\sigma(u)$ and there is an integer n such that u occurs at least twice in $\sigma^n(u)$. We define $\Theta_{x,u}(1)$ as the shortest non-empty word w such that wu is a prefix of $\sigma^n(u)$. We then compute $\sigma(\Theta_{x,u}(1))$. By construction, the word $\sigma(\Theta_{x,u}(1))u$ is a prefix of x and has u as prefix. By Proposition 3.6, there exist some unique words $w_1, w_2, \dots, w_k \in \mathcal{R}_{x,u}$ such that $\sigma(\Theta_{x,u}(1)) = \Theta_{x,u}(1)w_1 \cdots w_k$. Furthermore the word $\Theta_{x,u}(1)w_1 \cdots w_k u$ is a prefix of x . We add some letters $2, 3, \dots, \ell \leq k$ to $R_{x,u}$ and define $\Theta_{x,u}$ on $\{1, \dots, \ell\}$ by $\{\Theta_{x,u}(i) \mid 1 \leq i \leq \ell\} = \{\Theta_{x,u}(1), w_1, \dots, w_k\}$ and in the order of occurrences of the words w_i 's in $\sigma(\Theta_{x,u}(1))$. This also defines the image $\sigma_{x,u}(1)$. We iterate the process on the added letters. The algorithm stops when no new return word appears when we compute the images $\sigma(\Theta_{x,u}(i))$ for $i \in R_{x,u}$. \square

Example 7.20. Let ν be the Thue-Morse substitution and y be the Thue-Morse sequence. Let us follow the algorithm described above to compute the return substitution $\nu_{x,a}$.

1. We have $\nu^2(a) = abba$, so we set $\Theta_{y,a}(1) = abb$.
2. We compute $\nu \circ \Theta_{y,a}(1) = \nu(abb) = abbaba$. The return words ab and a do not belong yet to $\mathcal{R}_{y,a}$ so we add the letters 2 and 3 to $R_{y,a}$ and set $\Theta_{y,a}(2) = ab$, $\Theta_{y,a}(3) = a$ and $\nu_{y,a}(1) = 123$.

3. We compute $\nu \circ \Theta_{y,a}(2) = \nu(ab) = abba$. The return words abb and a already belong to $\mathcal{R}_{y,a}$ so we do not add any letter to $R_{y,a}$. We set $\nu_{y,a}(2) = 13$.
4. We compute $\nu \circ \Theta_{y,a}(3) = \nu(a) = ab$. The return word ab already belongs to $\mathcal{R}_{y,a}$ so we do not add any letter to $R_{y,a}$. We set $\nu_{y,a}(3) = 2$.
5. No more letters have to be checked so the algorithm stops and returns

$$\Theta_{y,a} : \begin{cases} 1 \mapsto abb \\ 2 \mapsto ab \\ 3 \mapsto a \end{cases} \quad \text{and} \quad \nu_{x,a} : \begin{cases} 1 \mapsto 123 \\ 2 \mapsto 13 \\ 3 \mapsto 2 \end{cases} .$$

7.3.2 Return substitutions for a set of words U

When x is an admissible fixed point of a primitive substitution σ , Proposition 7.16 states that the derived sequence w.r.t. any prefix u of x is again an admissible fixed point of a primitive substitution $\sigma_{x,u}$. This result can be generalized to the case where U is not a singleton but we need to be more careful. Indeed, the existence of $\sigma_{x,u}$ uses the fact that $\sigma(\mathcal{R}_{x,u})u \subset \mathcal{R}_{x,u}^*u \cap \mathcal{L}(\sigma)$, which may be not be the case when U is not a singleton.

Example 7.21. Consider the Thue-Morse substitution ν and the set $U = \{aa, bb\}$. We obtain $\mathcal{R}_{y,U} = \{aa, bb, aaba, bbab\}$ and, for instance, $\nu(aa) = abab$ is not prefix-comparable to any word in $\mathcal{R}_{y,U}^*U$.

Proposition 7.22. *Let σ be a primitive substitution and x be an element of X_σ . Let $U \subset \mathcal{L}(X_\sigma)$ be a set of non-empty words with the same length ℓ_U . There is a computable primitive substitution $\sigma_{x,U} : \tilde{\mathcal{R}}_{x,U}^* \rightarrow \tilde{\mathcal{R}}_{x,U}^*$ such that $X_{\sigma_{x,U}} = \Omega(\mathcal{D}_U(x))$. Furthermore, there is a computable constant C , not depending of U , such that*

$$|\sigma_{x,U}| \leq C \frac{\max_{w \in \mathcal{R}_{x,U}} |w|}{\min_{w \in \mathcal{R}_{x,U}} |w|}.$$

Proof. Let us first define the morphism $\sigma_{x,U}$. Let k be the smallest integer such that $\langle \sigma^k \rangle \geq (K_\sigma + 1)^2 \ell_U$. For any return word w to U in x and all $u \in U$, we have, by Theorem 3.11,

$$|w| \leq \max\{|r| \mid r \in \mathcal{R}_{x,u}\} \leq K_\sigma \ell_U,$$

where K_σ is the constant of linear recurrence of x . Still by Theorem 3.11, we thus have

1. for all $(w, u), (w', u') \in \tilde{\mathcal{R}}_{x,U}$, wu occurs in $\sigma^k(w')$;
2. for all $u \in U$, $\sigma^k(u)$ contains at least one occurrence of a word of U .

For any $w \in \mathcal{R}_{x,U} \cup U$ we set

$$\sigma^k(w) = p(w)m(w)s(w)$$

where $m(w)$ belongs to U and $p(w)m(w)$ has a unique occurrence of a word of U . Now take $(w, u) \in \tilde{\mathcal{R}}_{x,U}$. Then

$$\sigma^k(wu) = p(w)m(w)s(w)p(u)m(u)s(u)$$

and, by Proposition 7.8, there exist a unique admissible word $(w_1, u_1) \cdots (w_\ell, u_\ell)$ in $\tilde{\mathcal{R}}_{x,U}^*$ such that

$$w_1 \cdots w_\ell u_\ell = m(w)s(w)p(u)m(u)$$

We set

$$\tilde{\sigma}_{x,U}(w, u) = (w_1, u_1) \cdots (w_\ell, u_\ell).$$

This defines an endomorphism of $\tilde{\mathcal{R}}_{x,U}^*$. Then, we define $\sigma_{x,U}$ as the unique endomorphism of $\tilde{R}_{x,U}^*$ satisfying

$$\tilde{\sigma}_{x,U} \circ \tilde{\Theta}_{x,U} = \tilde{\Theta}_{x,U} \circ \sigma_{x,U}.$$

The morphism $\sigma_{x,U}$ is obviously computable. Furthermore, by the choice of k , $\sigma_{x,U}$ is a primitive morphism. We also have

$$|\sigma_{x,U}| \leq \frac{\max_{(w,u) \in \tilde{\mathcal{R}}_{x,U}} |\sigma^k(wu)|}{\min_{(w,u) \in \tilde{\mathcal{R}}_{x,U}} |wu|} \leq |\sigma^k| \frac{\max_{w \in \mathcal{R}_{x,U}} |w|}{\min_{w \in \mathcal{R}_{x,U}} |w|} \quad (16)$$

and the constant $C = |\sigma^k|$ is clearly computable and does not depend on U . Indeed, with the help of Theorem 3.11, it is sufficient to take k such that $\langle \sigma^k \rangle / K_\sigma \geq (K_\sigma + 1)^2$ and $\langle \sigma^k \rangle \geq (K_\sigma + 1)$. Using Corollary 4.15, we can choose k such that

$$\alpha^k \geq K_\sigma^3 (K_\sigma + 1)^2,$$

where α is the dominating eigenvalue of M_σ .

Now let us show that $X_{\sigma_{x,U}} = \Omega(\mathcal{D}_U(x))$. By Corollary 7.12, the subshift $\Omega(\mathcal{D}_U(x))$ is minimal. It is thus enough to show that $X_{\sigma_{x,U}}$ is included in $\Omega(\mathcal{D}_U(x))$. Again, one needs to show that any long enough word in $\mathcal{L}(\sigma_{x,U})$ belongs to $\mathcal{L}(\mathcal{D}_U(x))$ hence, by primitiveness of $\sigma_{x,U}$, that $\sigma_{x,U}^n(a)$ belongs to $\mathcal{L}(\mathcal{D}_U(x))$ for any letter $a \in \tilde{R}_{x,U}$ and for any $n \in \mathbb{N}$.

We proceed by induction on n . By construction of $\sigma_{x,U}$, the result is true for $n = 1$. Let us suppose it is true for $n - 1$. Let $a \in \tilde{R}_{x,U}$ and let us write $\sigma_{x,U}^{n-1}(a) = a_1 a_2 \cdots a_\ell \in \mathcal{L}(\mathcal{D}_U(x))$, with $a_1, \dots, a_\ell \in \tilde{R}_{x,U}$.

For $i \in \{1, \dots, \ell\}$, let us write $\tilde{\Theta}_{x,U}(a_i) = (w_i, u_i)$. As $a_1 a_2 \cdots a_\ell$ belong to $\mathcal{L}(\mathcal{D}_U(x))$, the word $w_1 w_2 \cdots w_\ell u_\ell$ belongs to $\mathcal{L}(x)$ and for all $1 \leq i \leq \ell$, $w_1 w_2 \cdots w_i u_i$ is a prefix of $w_1 w_2 \cdots w_\ell u_\ell$. We have to show that $\sigma_{x,U}(a_1 a_2 \cdots a_\ell)$ belongs to $\mathcal{L}(\mathcal{D}_U(x))$. By definition of $\sigma_{x,U}$, we have for all $i \in \{1, \dots, \ell\}$, $\sigma_{x,U}(a_i) = b_{i,1} \cdots b_{i,L_i}$, where $b_{i,1}, \dots, b_{i,L_i}$ are the unique letters in $\tilde{R}_{x,U}$ that satisfy both conditions

1. $\sigma^k(w_i u_i) = p(w_i) d_{x,U,1}(b_{i,1} \cdots b_{i,L_i}) m(u_i) s(u_i)$;
2. for all $j \in \{1, \dots, L_i\}$, the word $d_{x,U,1}(b_{i,1} \cdots b_{i,j}) d_{x,U,2}(b_{i,j})$ is a prefix of $d_{x,U,1}(b_{i,1} \cdots b_{i,L_i}) m(u_i)$.

Observe that for all $i \in \{1, \dots, \ell - 1\}$, the word $w_i u_i$ is a prefix of $w_i w_{i+1} u_{i+1}$ and we have

$$\sigma^k(w_i u_i) = p(w_i) m(w_i) s(w_i) p(u_i) m(u_i) s(u_i)$$

and

$$\begin{aligned} \sigma^k(w_i w_{i+1} u_{i+1}) &= p(w_i) m(w_i) s(w_i) \\ &\quad p(w_{i+1}) m(w_{i+1}) s(w_{i+1}) p(u_{i+1}) m(u_{i+1}) s(u_{i+1}). \end{aligned}$$

Consequently, we have $p(u_i) = p(w_{i+1})$ and $m(u_i) = m(w_{i+1})$.

In definitive, we have that $\sigma^k(w_1 w_2 \cdots w_\ell u_\ell)$ belongs to $\mathcal{L}(x)$ and the letters $b_{1,1}, \dots, b_{1,L_1}, \dots, b_{\ell,1}, \dots, b_{\ell,L_\ell}$ are the unique letters in $\tilde{R}_{x,U}$ that satisfy

1. $\sigma^k(w_1 w_2 \cdots w_\ell u_\ell) = p(w_1) d_{x,U,1}(b_{1,1} \cdots b_{1,L_1} \cdots b_{\ell,1} \cdots b_{\ell,L_\ell}) m(u_\ell) s(u_\ell)$;
2. for all $i \in \{1, \dots, \ell\}$, for all $j \in \{1, L_i\}$, the word

$$d_{x,U,1}(b_{1,1} \cdots b_{1,\ell_1} \cdots b_{i,1} \cdots b_{i,j}) d_{x,U,2}(b_{i,j})$$

is a prefix of

$$d_{x,U,1}(b_{1,1} \cdots b_{1,\ell_1} \cdots b_{\ell,1} \cdots b_{\ell,L_\ell}) m(u_\ell).$$

Hence, the word $\sigma_{x,U}(a_1 \cdots a_\ell)$ belongs to $\mathcal{L}(\mathcal{D}_U(x))$, which ends the proof. \square

Definition 7.23. The substitution $\sigma_{x,U}$ of the previous proposition is called a *return substitution* (w.r.t. U).

Example 7.24. Let us consider our running example with the Fibonacci sequence and follow the algorithm given by Proposition 7.22. Recall that we have $U = \{aa, ab\}$, $\tilde{\mathcal{R}}_{x,U} = \{(ab, aa), (a, ab), (ab, ab)\}$ and

$$\tilde{\Theta}_{y,U} : \begin{cases} 1 \mapsto (ab, aa) \\ 2 \mapsto (a, ab) \\ 3 \mapsto (ab, ab) \end{cases} .$$

For all $(w, u), (w', u') \in \tilde{\mathcal{R}}_{x,U} \times U$, we have

1. wu occurs in $\varphi^4(w')$;
2. $\varphi^4(u)$ contains at least one occurrence of a word of U .

Indeed, we have

$$\varphi^4 : \begin{cases} a \mapsto abaababa \\ ab \mapsto abaababaabaab \\ aa \mapsto abaababaabaababa \end{cases} .$$

Corollary 7.26. *Let $\sigma : A^* \rightarrow A^*$ be an aperiodic primitive substitution, x be an element of X_σ and $f : A \rightarrow B$ be a coding. There is a computable constant \tilde{Q}_σ such that*

$$\text{Card}(\{\sigma_{x,f^{-1}(\{v\})} \mid v \in f(\mathcal{L}(\sigma)) \setminus \{\epsilon\}\}) \leq \tilde{Q}_\sigma,$$

where ϵ is the empty word of B^* .

Proof. It suffices to show that there exist computable constants D, E such that for any non-empty word $v \in f(\mathcal{L}(\sigma))$,

$$\text{Card}(\tilde{\mathcal{R}}_{x,f^{-1}(\{v\})}) \leq D \quad \text{and} \quad |\sigma_{x,f^{-1}(\{v\})}| \leq E.$$

In that case, we can take $\tilde{Q}_\sigma = D^{(1+E)D}$. Let K_σ be the constant of linear recurrence of x . Of course, $f(x)$ is also linearly recurrent for the constant K_σ . Let v be a non-empty word in $f(\mathcal{L}(\sigma))$ and $U = f^{-1}(\{v\})$. Let us give a bound on $\text{Card}(\tilde{\mathcal{R}}_{x,U})$. For any $(w, u) \in \tilde{\mathcal{R}}_{x,U}$, we have $f(wu) = w'v$ for some concatenation w' of return words to v in $f(x)$. Since $|w| = |w'|$, we get, by Theorem 3.11, that $|v|/K_\sigma \leq |w| \leq K_\sigma|v|$. Still by Theorem 3.11, any word of $\mathcal{L}(\sigma)$ of length at most $(K_\sigma + 1)|v|$ has an occurrence in any word of $\mathcal{L}(\sigma)$ of length $(K_\sigma + 1)^2|v|$. Let z be a word of length $(K_\sigma + 1)^2|v|$ in $\mathcal{L}(\sigma)$. With the previous reasoning, there exist an admissible word $(w_1, u_1) \cdots (w_n, u_n) \in \tilde{\mathcal{R}}_{x,U}^*$ such that

1. $w_1 \cdots w_n u_n$ is a factor of z ;
2. $\{(w_i, u_i) \mid 1 \leq i \leq n\} = \tilde{\mathcal{R}}_{x,U}$.

We thus have

$$\text{Card}(\tilde{\mathcal{R}}_{x,U}) \leq n \leq \frac{(K_\sigma + 1)^2|v|}{|v|/K_\sigma} = K_\sigma(K_\sigma + 1)^2.$$

From Proposition 7.22, we also deduce that

$$|\sigma_{x,U}| \leq CK_\sigma^2,$$

where C is a computable constant depending on σ but not on U . □

7.3.3 It suffices to compute a finite number of return substitutions

We are now ready to state a necessary and sufficient condition for a coding to be a factor map between two minimal substitution subshifts.

Theorem 7.27. *Let $\sigma : A^* \rightarrow A^*$ and $\tau : B^* \rightarrow B^*$ be aperiodic primitive substitutions and let $x \in A^{\mathbb{N}}$ and $y \in B^{\mathbb{N}}$ be admissible fixed points of σ and τ respectively. Let $f : A \rightarrow B$ be a coding. Let $K = \max\{K_\sigma, K_\tau\}$ and let us consider the computable constants Q_τ and \tilde{Q}_σ of Corollary 7.18 and Corollary 7.26 respectively. Then f defines a factor map from (X_σ, S) to (X_τ, S) if, and only if, there exist two distinct prefixes u, v of y such that*

1. $1 \leq |u| \leq Q_\tau \tilde{Q}_\sigma$ and $K(K+1)|u| \leq |v| \leq K(K+1)|u| + Q_\tau \tilde{Q}_{\sigma,f}$;
2. $u, v \in f(\mathcal{L}(X_\sigma))$;
3. the return substitutions $\tau_{y,u}$ and $\tau_{y,v}$ are equal;
4. for $U = f^{-1}(\{u\})$ and $V = f^{-1}(\{v\})$, the return substitutions $\sigma_{x,U}$ and $\sigma_{x,V}$ are equal.

Proof. Theorem 7.15 shows that the conditions are sufficient. They are also necessary. Indeed, suppose that we can find prefixes u, v of y satisfying $1 \leq |u| \leq Q_\tau \tilde{Q}_\sigma$ and $K(K+1)|u| \leq |v| \leq K(K+1)|u| + Q_\tau \tilde{Q}_\sigma$ and u or v does not belong to $f(\mathcal{L}(\sigma))$. Then f can not be a factor from (X_σ, S) to (X_τ, S) . The result thus directly follows from the number of possible pairs of return substitutions of x and y (using Corollary 7.18 and Corollary 7.26). \square

Since all data in Theorem 7.27 are computable from σ and τ , we obtain Theorem 1.4 as a corollary.

7.4 Proof of Theorem 1.1 and Corollary 1.2

We deduce Theorem 1.1 from Theorem 3.1, Lemma 4.9, Proposition 4.11 and Theorem 7.27.

To prove Corollary 1.2, observe that one can decide whether there exist two factor maps $f : (X_\sigma, S) \rightarrow (X_\tau, S)$ and $g : (X_\tau, S) \rightarrow (X_\sigma, S)$. And thus that $f \circ g : (X_\sigma, S) \rightarrow (X_\sigma, S)$ is an endomorphism. But it is known from [Dur00] (see also [Cov72] for binary alphabets) that endomorphisms of minimal substitution subshifts are isomorphism. This is what is called the *coalescence property*. This shows Corollary 1.2.

8 The constant length case

In this section we look in more details to the factor maps between constant length substitution subshifts and in particular to the maps F_n defined by (8). The main theorem in [CDK17] shows that when there exists a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$ between aperiodic minimal one-to-one constant length substitution subshifts then there exists an other factor map $g : (X_\sigma, S) \rightarrow (X_\tau, S)$ given by a sliding block code of radius 1. In [CQY16] the result is improved for automorphism $f : (X_\sigma, S) \rightarrow (X_\sigma, S)$ with some extra assumptions. In this case g can be chosen with memory 0 and anticipation 1.

In this section we improve these results by removing the assumption of injectiveness and the additional assumptions in [CQY16] and by giving a link between f and g . This is Theorem 8.1.

8.1 Factor maps between constant length primitive substitution subshifts

We say that a substitution $\sigma : A^* \rightarrow A^*$ is *one-to-one on letters* when $\sigma(a) \neq \sigma(b)$ whenever $a \neq b$. When σ has constant length, it is clear that if $\sigma : A^* \rightarrow A^*$ is one-to-one on letters then $\sigma : A^* \rightarrow A^*$ is one-to-one and σ^n is one-to-one for all $n \geq 1$.

Theorem 8.1. *Let σ and τ be two primitive substitutions of constant length. Suppose there exists a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$ of radius r with (X_τ, S) aperiodic. Let $N_n = \inf\{k \geq 0 \mid S^k f(x) \text{ belongs to } \tau^n(X_\tau), x \in \sigma^n(X_\sigma)\}$, and, set $L = 0$ if τ is one-to-one on letters and $L = L_\tau$ otherwise.*

Then, there exists a factor map $F : (X_\sigma, S) \rightarrow (X_\tau, S)$ such that :

1. *If $r \geq N_n$ for all large enough n , then the anticipation of F is bounded by $L + 1$ and its memory by $L + 1$;*
2. *If $r < N_n \leq |\tau^n| - r$ for all large enough n , then the anticipation of F is bounded by L and its memory by $L + 1$;*
3. *If $N_n \geq |\tau^n| - r$ for all large enough n , then the anticipation of F is bounded by L and its memory by $L + 2$;*
4. *$f = F \circ S^n$ for some $n \in \mathbb{Z}$.*

Our proof follows the strategy used in [ST15] to show that the automorphism group of subshifts generated by primitive one-to-one constant length substitutions is virtually \mathbb{Z} .

We then use Theorem 8.1 to show the decidability of the factorization between two minimal constant length substitution subshifts.

Observe that when dealing with one-to-one property it is important to distinguish $\sigma : A^* \rightarrow A^*$ and $\sigma : A^{\mathbb{K}} \rightarrow A^{\mathbb{K}}$ as there exists substitutions σ such that $\sigma : A^* \rightarrow A^*$ is not one-to-one but $\sigma : X_\sigma \rightarrow X_\sigma$ is. For example not all primitive aperiodic substitutions $\sigma : A^* \rightarrow A^*$ are one-to-one but all of them are such that $\sigma : X_\sigma \rightarrow X_\sigma$ is one-to-one (Corollary 4.3).

As taking powers of a substitution does not change the associated subshift, the next result implies that we can restrict ourselves to the case of substitutions of the same constant length.

Proposition 8.2 ([Fag97b, Théorème 15]). *Let σ and τ be two primitive substitutions of constant length. If (X_τ, S) is aperiodic and if there exists a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$, then there exist $k, l \geq 1$ such that $|\sigma^k| = |\tau^l|$.*

Lemma 8.3. *Let σ and τ be two primitive substitutions of constant length p . Suppose there exists a factor map $\phi : (X_\sigma, S) \rightarrow (X_\tau, S)$ with (X_τ, S) aperiodic. Then, there exists a unique i belonging to $[0, \dots, p - 1]$ such that if x belongs to $\sigma(X_\sigma)$ then $\phi(x)$ belongs to $S^{-i}(\tau(X_\tau))$.*

Proof. Let $x \in \sigma(X_\sigma)$. From Corollary 4.5, there exists a unique $i(x)$ belonging to $[0, \dots, p-1]$ such that $\phi(x)$ belongs to $S^{-i(x)}(\tau(X_\tau))$. Let us show $i(x)$ is constant on $\sigma(X_\sigma)$. Observe that $S^j(x)$ belongs to $\sigma(X_\sigma)$ if and only if $j = k|\sigma|$ for some k and, in that case, we have $\phi(S^{k|\sigma|}(x)) = S^{k|\sigma|}\phi(x) \in S^{-i(x)}(\tau(X_\tau))$. Let $y \in \sigma(X_\sigma)$. Using the minimality of (X_σ, S) , there exists a sequence $(S^{k_n|\sigma|}x)_n$ tending to y . As $\phi(S^{k_n|\sigma|}x)$ belongs to $S^{-i(x)}\tau(X_\tau)$ for all n , we get that $\phi(y)$ also belongs to $S^{-i(x)}\tau(X_\tau)$. \square

Proposition 8.4. *Let σ and τ be two primitive substitutions of the same constant length and let $l \geq 0$. Suppose there exists a factor map $f : (X_\sigma, S) \rightarrow (X_\tau, S)$ with radius r and that (X_τ, S) is aperiodic. Then,*

1. *There exist n and a factor map $F : (X_\sigma, S) \rightarrow (X_\tau, S)$ uniquely defined by*

$$\tau^l \circ F = S^n \circ f \circ \sigma^l, \quad 0 \leq n < |\tau|^l. \quad (17)$$

2. *The memory t and the anticipation s of F satisfy*

$$\begin{aligned} t &\leq \max(0, L + \lceil (r-n)/|\tau|^l \rceil), \\ s &\leq L + \lceil (r+n)/|\tau|^l \rceil, \end{aligned}$$

where $L = 0$ if τ is one-to-one on letters and $L = L_\tau$ otherwise.

Proof. (1) By Lemma 8.3, there exists a unique n such that $0 \leq n < |\tau|^l$ and for all $x \in X_\sigma$, $S^n \circ f \circ \sigma^l(x)$ belongs to $\tau^l(X_\tau)$. Thus, using Corollary 4.3, Equation (17) uniquely defines a continuous map F . Using again Corollary 4.3 and noticing that

$$\tau^l \circ F \circ S(x) = S^{n+|\sigma|^l} \circ f \circ \sigma^l(x) = S^{|\sigma|^l} \circ \tau^l \circ F(x) = \tau^l \circ S \circ F(x),$$

one obtains that F is a factor map between (X_σ, S) and (X_τ, S) .

(2) We know from the Curtis-Hedlund-Lyndon theorem that F is a factor map if and only if there exist $s, t \geq 0$ such that for all x , $F(x)_0$ only depends on $x_{[-t, s]}$.

Let $s, t \geq 0$ and $x \in X_\sigma$. Then the word $\sigma^l(x)_{[-t|\sigma|^l, s|\sigma|^l]}$ only depends on $x_{[-t, s]}$. Thus, it is also the case of the words $f \circ \sigma^l(x)_{[-t|\sigma|^l+r, s|\sigma|^l-r]}$ and $\tau^l \circ F(x)_{[-t|\sigma|^l+r-n, s|\sigma|^l-r-n]}$. Let $L_l = 0$ if τ is one-to-one on letters and $L_l = L_{\tau^l}$ otherwise. From Corollary 4.4,

$$F(x) \left[\left\lceil \frac{-t|\sigma|^l+r-n+L_l}{|\tau|^l} \right\rceil, \left\lfloor \frac{s|\sigma|^l-r-n-L_l}{|\tau|^l} \right\rfloor \right],$$

and thus, from Proposition 4.8,

$$F(x) \left[\left\lceil \frac{-t|\sigma|^l+r-n+|\tau|^l L_l}{|\tau|^l} \right\rceil, \left\lfloor \frac{s|\sigma|^l-r-n-|\tau|^l L_l}{|\tau|^l} \right\rfloor \right]$$

only depends on $x_{[-t,s]}$, that is,

$$F(x)_{[[-t+L_1+(r-n)/|\tau|^l, [s-L_1-(r+n)/|\tau|^l]]}$$

only depends on $x_{[-t,s]}$. Hence it is sufficient that s and t satisfy $s - L_1 - (r + n)/|\tau|^l \geq 0$ and $-t + L_1 + (r - n)/|\tau|^l \leq 0$, which ends the proof. \square

Consider $\Delta(\sigma, \tau)$ the set of factor maps from (X_σ, S) to (X_τ, S) . From Proposition 8.4 there is a map $\Phi_{\sigma, \tau}$ from $\Delta(\sigma, \tau)$ to itself defined by

$$\tau \circ \Phi_{\sigma, \tau}(f) = S^{N_{\sigma, \tau}(f)} \circ f \circ \sigma, \quad \forall f \in \Delta(\sigma, \tau), \quad (18)$$

for some uniquely defined integer $N_{\sigma, \tau}(f)$ verifying $0 \leq N_{\sigma, \tau}(f) < |\tau|$. Observe $N_{\sigma, \tau}(f)$ is the unique integer n such that $0 \leq n < |\tau|$ and $S^n \circ f \circ \sigma(X_\sigma) \subset \tau(X_\tau)$.

Lemma 8.5. *Let σ and τ be two primitive substitutions of the same constant length. Suppose (X_τ, S) is non-periodic. Let $g, h \in \Delta(\sigma, \tau)$. We have the following properties.*

1. *If $h = S^n g$ with $N_{\sigma, \tau}(g) - |\tau| < n \leq N_{\sigma, \tau}(g)$ then $N_{\sigma, \tau}(S^n g) = N_{\sigma, \tau}(g) - n$ and $\Phi_{\sigma, \tau}(S^n g) = \Phi_{\sigma, \tau}(g)$.*
2. *If $\Phi_{\sigma, \tau}(g) = S^k \circ \Phi_{\sigma, \tau}(h)$ for some k then $g = S^l \circ h$ for some l .*
3. *For all $m, n > 0$ one has*

$$N_{\sigma^{m+n}, \tau^{m+n}}(g) = |\tau|^m |N_{\sigma^n, \tau^n}(\Phi_{\sigma^m, \tau^m}(g)) + N_{\sigma^m, \tau^m}(g)$$

$$\text{and } \Phi_{\sigma^n, \tau^n} \circ \Phi_{\sigma^m, \tau^m}(g) = \Phi_{\sigma^{n+m}, \tau^{n+m}}(g).$$

4. *For all $n > 0$, $\Phi_{\sigma, \tau}^n = \Phi_{\sigma^n, \tau^n}$.*

Proof. For the first assertion, we observe that for all n such that $N_{\sigma, \tau}(g) - |\tau| < n \leq N_{\sigma, \tau}(g)$, we have $0 \leq N_{\sigma, \tau}(g) - n < |\tau|$ and

$$S^{N_{\sigma, \tau}(g)} \circ g \circ \sigma = S^{N_{\sigma, \tau}(g) - n} \circ S^n \circ g \circ \sigma.$$

We thus have $N_{\sigma, \tau}(S^n g) = N_{\sigma, \tau}(g) - n$, which implies, using Corollary 4.3, that $\Phi_{\sigma, \tau}(S^n g) = \Phi_{\sigma, \tau}(g)$.

Let us now prove the second assertion and consider g and h in $\Delta(\sigma, \tau)$ such that $\Phi_{\sigma, \tau}(g) = S^k \Phi_{\sigma, \tau}(h)$ for some k . Then, one gets

$$\begin{aligned} S^{N_{\sigma, \tau}(g)} \circ g \circ \sigma &= \tau \circ \Phi_{\sigma, \tau}(g) \\ &= \tau \circ S^k \circ \Phi_{\sigma, \tau}(h) \\ &= S^{k|\tau|} \circ \tau \circ \Phi_{\sigma, \tau}(h) \\ &= S^{k|\tau|} \circ S^{N_{\sigma, \tau}(h)} \circ h \circ \sigma. \end{aligned}$$

By minimality of (X_σ, S) we deduce that $g = S^{k|\tau| + N_{\sigma, \tau}(h) - N_{\sigma, \tau}(g)} \circ h$.

We now prove the third assertion. Let m, n be two positive integers. One has the following equalities:

$$\begin{aligned}\tau^m \circ \Phi_{\sigma^m, \tau^m}(g) &= S^{N_{\sigma^m, \tau^m}(g)} \circ g \circ \sigma^m, \\ \tau^n \circ \Phi_{\sigma^n, \tau^n} \circ \Phi_{\sigma^m, \tau^m}(g) &= S^{N_{\sigma^n, \tau^n}(\Phi_{\sigma^m, \tau^m}(g))} \circ \Phi_{\sigma^m, \tau^m}(g) \circ \sigma^n.\end{aligned}$$

Thus, we get

$$\begin{aligned}\tau^{n+m} \circ \Phi_{\sigma^n, \tau^n} \circ \Phi_{\sigma^m, \tau^m}(g) &= \tau^m \circ S^{N_{\sigma^n, \tau^n}(\Phi_{\sigma^m, \tau^m}(g))} \circ \Phi_{\sigma^m, \tau^m}(g) \circ \sigma^n \\ &= S^{|\tau^m| N_{\sigma^n, \tau^n}(\Phi_{\sigma^m, \tau^m}(g))} \circ \tau^m \circ \Phi_{\sigma^m, \tau^m}(g) \circ \sigma^n \\ &= S^{|\tau^m| N_{\sigma^n, \tau^n}(\Phi_{\sigma^m, \tau^m}(g)) + N_{\sigma^m, \tau^m}(g)} \circ g \circ \sigma^{n+m}.\end{aligned}$$

As

$$\begin{aligned}0 \leq |\tau^m| N_{\sigma^n, \tau^n}(\Phi_{\sigma^m, \tau^m}(g)) + N_{\sigma^m, \tau^m}(g) &\leq |\tau^m| (|\sigma^n| - 1) + |\sigma^m| - 1 \\ &\leq |\sigma^{m+n}| - 1,\end{aligned}$$

we obtain, by definition, that

$$N_{\sigma^{m+n}, \tau^{m+n}}(g) = |\tau^m| N_{\sigma^n, \tau^n}(\Phi_{\sigma^m, \tau^m}(g)) + N_{\sigma^m, \tau^m}(g),$$

hence that $\Phi_{\sigma^n, \tau^n} \circ \Phi_{\sigma^m, \tau^m}(g) = \Phi_{\sigma^{n+m}, \tau^{n+m}}(g)$.

For the (4), the equality $\Phi_{\sigma, \tau}^n = \Phi_{\sigma^n, \tau^n}$ is a direct consequence of (3). \square

Proof of Theorem 8.1. Let r be the radius of f . From Proposition 8.2, one can assume, taking powers of the substitutions if needed, that $|\sigma| = |\tau|$. We recall that from Lemma 8.5, $\Phi_{\sigma^n, \tau^n}(f) = \Phi_{\sigma, \tau}^n(f)$ is a factor map from (X_σ, S) to (X_τ, S) . We will find F among these maps.

Observe that the quantity N_n defined in the statement of Theorem 8.1 is exactly $N_{\sigma^n, \tau^n}(f)$.

Let us show that for some $n > 0$ the memory and the anticipation of $\Phi_{\sigma, \tau}^n(f)$ are as required. Indeed, let $n \in \mathbb{N}$. Let t and s be, respectively, the memory and the anticipation of $\Phi_{\sigma, \tau}^n(f)$. From Proposition 8.4 one can take

$$\begin{aligned}t &= \max(0, L + \lceil (r - N_{\sigma^n, \tau^n}(f)) / |\tau^n| \rceil) \\ s &= L + \lceil (r + N_{\sigma^n, \tau^n}(f)) / |\tau^n| \rceil.\end{aligned}$$

This shows the first three statements in the conclusion.

As there are finitely many such factor maps of radius at most 1, there exist p, q , with $p < q$, such that $\Phi_{\sigma, \tau}^p(f) = \Phi_{\sigma, \tau}^q(f)$. From Lemma 8.5, one gets $\Phi_{\sigma, \tau}^{q-p}(f) = S^m \circ f$ for some m . \square

An example in [CKL08] studying the subshifts isomorphic to the Thue-Morse subshift shows that these bounds are sharp. In particular, this implies that, considering the notations of Theorem 8.1, if τ is a one-to-one primitive constant length substitution, then (X_τ, S) can be a factor of (X_σ, S) only if τ is defined

on an alphabet of size at most $\#\mathcal{L}_3(\sigma)$. This result is not true anymore for non-constant length substitutions. Indeed, using a slightly different definition of the substitution ξ of Theorem 3.1, we would build a primitive and non-constant length substitution ξ' that is injective on the letters, defined on a 24-letters alphabet and such that $(X_{\xi'}, S)$ is isomorphic to the Thue-Morse subshift (see also Section 3.3).

8.2 Decidability of the factorization

Assume that we are given two primitive substitutions of constant length σ and τ . By Theorem 8.1, to decide whether there exists a factor map between (X_σ, S) and (X_τ, S) , it suffices to test all sliding block codes of radius $L + 1$. Thus, for each such sliding block code f , we have to decide whether $\mathcal{L}(f(X_\sigma))$ is equal to $\mathcal{L}(X_\tau)$. But it can be observed that both languages are languages of automatic sequences. A result attributed to V. Bruyère in [Fag97b] allows to conclude as it states that the inclusion of automatic languages is decidable. We give some details below.

8.2.1 Background on automatic sequences

For more details concerning this section, we refer to [BHMV94] and [Rig14].

A *k-automatic sequence* is a substitutive sequence w.r.t. some constant substitution σ such that $k = |\sigma|$.

Let $k \geq 2$ be an integer and define the function $V_k : \mathbb{N} \rightarrow \mathbb{N}$ by $V_k(0) = 1$ and

$$V_k(n) = k^p, \quad \text{where } p = \max\{r \mid k^r \text{ divides } n\}, \quad n \neq 0.$$

Let us consider the first order logical structure $\langle \mathbb{N}, +, V_k, = \rangle$. A subset $E \subset \mathbb{N}$ is *k-definable* if there exists a formula in $\langle \mathbb{N}, +, V_k, = \rangle$ describing E . For example, the formula “ $V_2(n) = n$ ” describes the set $\{2^n \mid n \in \mathbb{N}\}$ as the power of 2 are exactly the fixed points of V_2 . Consider the formula $\phi(n)$ defined by $(\exists m)(n = m + m)$, then $\{n \mid \phi(n)\} = 2\mathbb{N}$. As ϕ is a formula expressed in $\langle \mathbb{N}, +, = \rangle$, this means that $2\mathbb{N}$ is *k-definable* for all $k \geq 2$.

A sequence $x \in A^{\mathbb{N}}$ is *k-definable* if for all $a \in A$ the set $\{n \mid x_n = a\}$ is *k-definable*.

Theorem 8.6 ([Buc60, Cob72]). *Let $k \geq 2$ and A be a finite alphabet. A sequence $x \in A^{\mathbb{N}}$ is *k-automatic* if and only if it is *k-definable*.*

Theorem 8.7. [Buc60] *The theory $\langle \mathbb{N}, +, V_k, = \rangle$ is decidable, that is, for each closed formula expressed in the first order logical structure $\langle \mathbb{N}, +, V_k, = \rangle$, there is an algorithm deciding whether it is true or not.*

We refer to [CRS12, Cha18] for expository texts on this decision procedure. In particular, it has been implemented in the software *Walnut* [Mou16] and can be downloaded from www.cs.uwaterloo.ca/shallit/papers.html.

As a consequence of Theorem 8.7 the following theorem is proven in [Fag97b] and attributed to V. Bruyère.

Theorem 8.8. *Let x and y be two k -automatic sequences. It is decidable whether $\mathcal{L}(x)$ is included in $\mathcal{L}(y)$.*

8.2.2 Decidability of the factorization for constant length primitive substitution subshifts

Let us prove that the factorization is decidable for aperiodic subshifts generated by primitive constant length substitutions. Let σ and τ be two primitive substitutions of constant length. Let us describe the algorithm testing whether there exists a factor map F from (X_σ, S) to (X_τ, S) .

Algorithm : Test all sliding block codes of radius $L + 1$ using the algorithm given by Theorem 8.8 where L is defined in Theorem 8.1.

Let us give some explanations.

First we check whether (X_τ, S) is periodic or not as this is decidable [Hon86, Hon08, ARS09]. If (X_τ, S) is periodic then one can decide whether there exists a factor map (Section 6). Suppose (X_τ, S) is non periodic. One has first to test whether σ and τ have some (non trivial) power of their length that are equal. This is clearly decidable. If they do not, then there is no factor map (Proposition 8.2). If they do, then Theorem 8.4 ensures that it suffices to consider sliding block codes of radius $L + 1$. There are finitely many such maps. Thus it suffices to test each of them. Let f be such a sliding block code of radius $L + 1$. To test whether it defines a factor map from (X_σ, S) to (X_τ, S) is equivalent to prove that $\mathcal{L}(f(x)) = \mathcal{L}(y)$ where x and y are respectively fixed points of σ and τ . Due to Theorem 8.8 it suffices to prove that $f(x)$ is an automatic sequence. This is clear using the substitutions on the blocks of radius $L + 1$ (see Section 4.2). Thus we proved the following theorem.

Theorem 8.9. *Let (X, S) and (Y, S) be two minimal constant length substitution subshifts. It is decidable to know whether there exists a factor map between these two systems.*

8.3 How to get rid of the injective substitution assumption

The proof of Theorem 8.9 involves sliding block codes of radius $L + 1$ where $L = 0$ if the substitutions are one-to-one. In this section, we show one can always suppose the substitutions are one-to-one, up to isomorphism. This will reduce the number of sliding block codes to test and the size of the involved alphabets. Hence, this improves the algorithm to decide the factorization.

Let $\sigma : A^* \rightarrow A^*$ be a primitive substitution such that (X_σ, S) is not periodic. Let $B \subset A$ and $\phi : A^* \rightarrow B^*$ be a coding such that:

$$\text{if } \phi(a) = \phi(b) \text{ then } \sigma(a) = \sigma(b). \quad (19)$$

Then, there exists a unique endomorphism $\tau : B^* \rightarrow B^*$ defined by

$$\tau \circ \phi = \phi \circ \sigma.$$

It is clear that τ is primitive. We say τ is an *injectivation* of σ .

Proposition 8.10. [BDM04] *Let τ be an injectivation of the non periodic primitive endomorphism σ . Then, (X_σ, S) is conjugate to (X_τ, S) . Moreover τ can be chosen injective and such a τ can be algorithmically found.*

Proof. The first claim is proven in [BDM04]. The two other claims are straightforward but we provide a proof below.

Suppose there exist a and b such that $\sigma(a) = \sigma(b)$. Let $B = A \setminus \{b\}$. Consider the map $\phi : A^* \rightarrow B^*$ defined by $\phi(c) = c$ if $c \neq b$ and $\phi_1(b) = a$. It satisfies (19) and thus defines an injectivation ζ of σ . Observe that the cardinality of the alphabet decreases. If ζ is not injective, we proceed in the same way to obtain a new substitution τ that will be an injectivation of ζ and of σ and that will be defined on a smaller alphabet. As A is finite, iterating the process, we will obtain an injectivation τ of σ that is injective. This achieves the proof. \square

8.4 Listing of the factors

Let σ be a primitive constant length substitution. We know from [Dur00] that the number of aperiodic subshift factors of (X_σ, S) is finite and all such factors are conjugate to a substitution subshift (Theorem 3.1). We are not able to establish the list of its subshift factors but what precedes allows to find the exhaustive list of its constant length substitution subshift factors. Indeed, periodic factors can be easily found with the main result in [Dek78], see Section 6. For the aperiodic constant length substitution subshift factors, Section 8.3 allows us to suppose the substitutions are one-to-one on letters. Then, Theorem 8.1 indicates it suffices to test finitely many sliding block codes. Theorem 8.8 can decide whether or not they are conjugate one to the other. This gives the exhaustive list of aperiodic constant length substitution subshift factors. However, some subshift factors might be of non-constant length. More precisely, it is not known whether a factor of a constant length substitution subshift is isomorphic to a constant length substitution subshift. As a consequence, it is not clear that the list obtained above corresponds to the list of all subshift factors.

In [CDK17] an other algorithm is given but with more assumptions on the subshifts.

We leave open the question to find the list of all aperiodic substitution subshift factors.

9 Factor maps between linearly recurrent subshifts

In this section we prove Theorem 1.5.

Given a dynamical system (X, T) , $\text{Aut}(X, T)$ will stand for the group of automorphisms of (X, T) , that is the group of continuous and bijective maps defined on X and commuting with T . In [DDMP16] (see also [CK16]) has been proven the following result.

Theorem 9.1. *Let (X, S) be an aperiodic minimal subshift. If*

$$\liminf_{n \in \mathbb{N}} \frac{p_X(n)}{n} < +\infty,$$

then $\text{Aut}(X, S)$ is virtually \mathbb{Z} , i.e., the quotient of $\text{Aut}(X, S)$ by the group generated by the shift map is a finite group.

The previous result means that there exist finitely many elements f_1, \dots, f_n of $\text{Aut}(X, S)$ such that for any $f \in \text{Aut}(X, S)$ there exist $m \in \mathbb{Z}$ and $i \in \{1, \dots, n\}$ satisfying $f = f_i \circ S^m$. In this section we extend this result to factor maps but in the restrictive context of the linearly recurrent subshifts (which are known to have linear complexity). Observe that for factor maps we loose the group structure which is a key argument in both papers [DDMP16] and [CK16].

Theorem 1.5 includes answers to questions stated in [ST15] but in the more general context of factor maps. The proof is inspired by the one of Theorem 1 in [Dur00].

Proof of Theorem 1.5. Let A and B be the alphabets of X and Y respectively. Consider the constant $N = (2(K+1)^3 + 1)^{6K^2(2(K+1)^3 + 1)} + 1$ and suppose that for all $i \in \{1, \dots, N\}$, $f_i : (X, S) \rightarrow (Y, S)$ is a factor map. It suffices to show that there exist $i \neq j$ and $k \in \mathbb{Z}$ such that $f_i = S^k \circ f_j$.

By the Curtis-Hedlund-Lyndon Theorem there exists a positive integer r such that for all $i \in \{1, \dots, N\}$ there is a sliding block code $\phi_i : A^{2r+1} \rightarrow B$ satisfying

$$(f_i(x))_j = \phi_i(x_{[j-r, j+r]}) \quad \forall x \in X, \forall j \in \mathbb{Z}. \quad (20)$$

Let us fix some $x \in X$ and some $y \in Y$. For all $i \in \{1, \dots, N\}$, there exists an integer k_i such that $(S^{k_i} f_i(x))_{[0, 2r-1]} = y_{[0, 2r-1]}$. By Proposition 4.16, we can assume that $0 \leq k_i \leq 4rK$. We then define the factor map $\tilde{f}_i : (X, S) \rightarrow (Y, S)$ by $\tilde{f}_i(x') = S^{k_i} f_i(x')$ for all $x' \in X$. Thus \tilde{f}_i has a radius of $4rK + r$: there exists a sliding block code $\tilde{\phi}_i : A^{2r(4K+1)+1} \rightarrow B$ such that

$$(\tilde{f}_i(x'))_j = \tilde{\phi}_i(x'_{[j-r(4K+1), j+r(4K+1)]}) \quad \forall x' \in X, \forall j \in \mathbb{Z}.$$

Furthermore, by choice of k_i we have for all i

$$\tilde{\phi}_i(x_{[-r(4K+1), r(4K+1)+2r-1]}) = y_{[0, 2r-1]}. \quad (21)$$

We will now show that there exist i and j , $i \neq j$, such that $\tilde{f}_i = \tilde{f}_j$, hence $f_i = S^{k_j - k_i} f_j$.

We fix $r' = r(4K+1)$ and we consider the block representation $X^{(r')}$ of X which is isomorphic to X through the isomorphism h associated with the sliding block code $\phi : A^{2r'+1} \rightarrow A^{(r')}$, $a_1 \cdots a_{2r'+1} \mapsto (a_1 \cdots a_{2r'+1})$ (see Lemma 4.9).

Observe that, from Theorem 3.11, for all $u \in \mathcal{L}(X^{(r')})$ we have:

1. for all words $v \in \mathcal{R}_{X^{(r')}, u}$, $\frac{1}{K}(|u| + 2r') \leq |v| \leq K(|u| + 2r')$, and,
2. $\#(\mathcal{R}_{X^{(r')}, u}) \leq (K+1)^3$.

For all $i \in \{1, \dots, N\}$ let $\psi_i : A^{(r')*} \rightarrow B^*$ be the morphism defined by, for $(u) \in A^{(r')}$, $\psi_i((u)) = \tilde{\phi}_i(u)$. It defines a factor map $g_i : (X^{(r')}, S) \rightarrow (Y, S)$. Since for all i , $f_i = g_i \circ h$, it suffices to prove that there exist i and j , $i \neq j$, such that $g_i = g_j$. By minimality, it is then enough to show that $g_i(x^{(r')}) = g_j(x^{(r')})$ for some $i \neq j$. Indeed, for all $z^{(r')} \in X^{(r')}$, there is an increasing sequence of integers $(n_\ell)_{\ell \in \mathbb{N}}$ such that $z^{(r')} = \lim_{\ell \rightarrow +\infty} S^{n_\ell} x^{(r')}$. If $g_i(x^{(r')}) = g_j(x^{(r')})$, we get by continuity that

$$\begin{aligned} g_i(z^{(r')}) &= \lim_{\ell \rightarrow +\infty} g_i(S^{n_\ell} x^{(r')}) = \lim_{\ell \rightarrow +\infty} S^{n_\ell} g_i(x^{(r')}) \\ &= \lim_{\ell \rightarrow +\infty} S^{n_\ell} g_j(x^{(r')}) = g_j(z^{(r')}). \end{aligned}$$

Take $u = (x^{(r')})_{[0, r'-1]}$ and $v = y_{[0, r'-1]}$. From (21) we have $\psi_i(u) = v$ for all i . Hence for each return word $w \in \mathcal{R}_{x^{(r')}, u}$, the word $\psi_i(w)$ is a concatenation of return words in $\mathcal{R}_{y, v}$. This induces a non-erasing morphism λ_i from $R_{x^{(r')}, u}^*$ to $R_{y, v}^*$ defined by

$$\psi_i \circ \Theta_{x^{(r')}, u} = \Theta_{y, v} \circ \lambda_i.$$

This morphism naturally induces a map from $R_{x^{(r')}, u}^{\mathbb{Z}}$ to $R_{y, v}^{\mathbb{Z}}$ that we also denote by λ_i and that satisfies

$$g_i \circ \Theta_{x^{(r')}, u} = \Theta_{y, v} \circ \lambda_i.$$

As ψ_i is a coding and $|v| = r'$, for all $b \in R_{x^{(r')}, u}$ we have from (1) above and (3) of Proposition 4.16

$$|\lambda_i(b)| \leq \frac{K(|u| + 2r')}{\frac{|v| - 2rK}{K}} = \frac{3K^2 r'}{r' - 2rK} = \frac{3K^2(4K + 1)r}{(2K + 1)r} \leq 6K^2.$$

Since, from (4) of Proposition 4.16, we also have

$$\begin{aligned} \#(\mathcal{R}_{Y, v}) &\leq \frac{(K + 1)K^2(r(4K + 1) + 2r) + 2rK^2}{2rK + 1} \\ &\leq (K + 1)K^2 \frac{4K + 3}{2K} + K \leq 2(K + 1)^3, \end{aligned}$$

we get $\#\{\lambda_i \mid 1 \leq i \leq N\} < N$ which implies that there exist i, j , $i \neq j$, such that $\lambda_i = \lambda_j$. We set $\lambda = \lambda_i$.

Since u is a prefix of $x^{(r')+}$, from Proposition 3.7 there exists a unique sequence $z \in R_{x^{(r')}, u}^{\mathbb{Z}}$ such that $\Theta_{x^{(r')}, u}(z) = x^{(r')}$. We get

$$g_i(x^{(r')}) = g_i \circ \Theta_{x^{(r')}, u}(z) = \Theta_{y, v} \circ \lambda(z) = g_j \circ \Theta_{x^{(r')}, u}(z) = g_j(x^{(r')}),$$

which ends the proof. \square

10 Other decidability problems for morphic subshifts

In this section we mention some open problems of decidability concerning morphic subshifts. The inputs will always be the morphism ϕ and the endomorphism σ defining the morphic sequence or pairs of such couples (ϕ, σ) .

10.1 Factor maps for subshifts generated by automatic sequences

Fixed points of constant length substitutions are a special kind of automatic sequences. Even if Theorem 1.3 applies to minimal automatic subshifts, to have a better radius for factor maps it would be interesting to know whether Theorem 8.1 remains valid for minimal automatic subshifts. This would make more efficient the algorithm to find its automatic subshift factors.

10.2 The non minimal case

We recall that primitive substitutions, as well as uniformly recurrent sequences, generate minimal subshifts. Is it possible to extend the main results stated in the introduction (Theorem 1.1, Corollary 1.2, Theorem 1.3 and Theorem 1.4) to (non uniformly recurrent) morphic sequences?

Let us focus on Theorem 1.4. Extension to the non minimal case, in this case, is equivalent to the decidability problem of the equality problem of morphic languages.

Equality problem of morphic languages. Given two morphic sequences x and y , is it decidable whether $\mathcal{L}(x) = \mathcal{L}(y)$?

Indeed, suppose (X, S) and (Y, S) are two morphic subshifts generated by, respectively, the morphic sequences x and y , and let $\phi : A^{\mathbb{Z}} \rightarrow B^{\mathbb{Z}}$ be a factor map. It is easy to show that ϕ is a factor map from (X, S) to (Y, S) if and only if $\mathcal{L}(\phi(x))$ is equal to $\mathcal{L}(y)$.

If we do not ask the factor map $\phi : (X, S) \rightarrow (Y, S)$ to be onto, then it is equivalent to the inclusion problem of morphic languages.

Inclusion problem of morphic languages. Given two morphic sequences x and y , is it decidable whether $\mathcal{L}(x)$ is included in $\mathcal{L}(y)$?

Of course the Inclusion problem implies the Equality problem, both problems being open.

10.3 Morphic languages

Let us consider the Inclusion and Equality problems.

In [Fag97a] is given positive answers to the Inclusion problem for some families of purely morphic sequences, one for those defined on a 2-letter alphabet and

an other for those generated by primitive substitutions under some mild assumptions. Observe that Theorem 1.4 solves this problem for uniformly recurrent morphic sequences as inclusion of languages of uniformly recurrent sequences implies the equality.

Theorem 10.1. *Inclusion problem for morphic languages is decidable for uniformly recurrent morphic sequences.*

Proof. It is left as an exercise. □

Both problems are open if we do not assume uniform recurrence. Nevertheless, I. Fagnot [Fag97b] solved both decidability problem for automatic sequences, without the assumption of uniform recurrence. She considered two cases : the multiplicative dependence and independence of the dominant eigenvalues p and q (which are integers in these settings) of the underlying substitutions, that is whether $\log p / \log q$ belongs or not to \mathbb{Q} . She managed to treat the multiplicatively independent case using the extension of Cobham's theorem for automatic languages she obtained in the same paper, then testing the ultimate periodicity of the languages to see whether they are equal. For the dependent case she used the logical framework of Presburger arithmetic associated with automatic sequences.

Theorem 10.2. *[Fag97b] Inclusion problem for morphic languages is decidable for automatic sequences.*

10.4 Equality problem of morphic sequences

Given two morphic sequences x and y , is it decidable whether $x = y$?

For purely morphic sequences, the Equality problem of morphic sequences is called the *D0L ω -equivalence problem* and was solved in 1984 by K. Culik II and T. Harju [CH84]. In [Dur12], it was shown to be true for primitive morphic sequences and that more can be said for the D0L ω -equivalence problem in the primitive case: the two (underlying) primitive substitutions have some non trivial powers that should coincide on some cylinders if $x = y$.

The general case of the Equality problem remains open.

10.5 The listing of the factors

As we mentioned in Section 8.4 for constant length substitution subshifts it would be of interest to find an algorithm that, given a minimal morphic subshift, provides the list of all its (finitely many) aperiodic subshift factors. More precisely, to consider the following problem.

Computability of the set of aperiodic factor subshifts of a minimal morphic subshift. Given a minimal morphic subshift (X, S) . Can we compute a finite set of primitive substitutions $\{\sigma_1, \dots, \sigma_N\}$ such that any aperiodic factor subshift of (X, S) is isomorphic to some (X_{σ_n}, S) ?

From [Dur00] and Theorem 3.1 we know that such a set of substitutions exists but we do not have an algorithm to find them.

Nevertheless, this has been partly realized in [CDK17] in the case of minimal constant length substitution subshifts to the detail that they only consider constant length substitution factors. It is not clear that with such a restriction one can obtain all aperiodic subshifts factors. Maybe some factors are not isomorphic to constant length substitution subshifts. What we know from the language version of Cobham's theorem [Dur98b] is that the dominant eigenvalues of the underlying substitutions should have a non trivial common power. In this case, i.e., starting with a primitive substitution of constant length k , the substitutions σ_i above should have a dominant eigenvalue $k^{n(i)}$ for some integer $n(i)$. But this does not ensure that the σ_i can be chosen of constant length k^n for some n . At least we do not know such a result. Hence we address the following question: Does there exist aperiodic subshift factors of constant length substitution subshifts that are not isomorphic to constant length substitution subshifts?

10.6 Orbit equivalence

A factor map $\phi : (X, S) \rightarrow (Y, T)$ between minimal dynamical has the property that orbits are sent to orbits:

$$\phi(\{S^n(x) \mid n \in \mathbb{Z}\}) = \{S^n(\phi(x)) \mid n \in \mathbb{Z}\} \quad (22)$$

for all $x \in X$. When considering isomorphisms, this defines an equivalence relation. One can relax the conditions and consider ϕ is just an homeomorphism satisfying (22). In this case we say (X, S) and (Y, T) are *orbit equivalent*. It is classical to observe that in this situation there exists $N : X \rightarrow \mathbb{Z}$ such that

$$\phi(S(x)) = S^{N(x)}\phi(x). \quad (23)$$

When N has at most one point of discontinuity, we say (X, S) and (Y, T) are *strong orbit equivalent*.

A remarkable result in [GPS95] characterizes strong orbit equivalence of minimal Cantor systems by means of dimension groups. In [BJKR01] it is shown that the isomorphism of stationary simple dimension groups is decidable. This shows, as dimension groups of minimal substitution subshifts are simple and stationary [For97, DHS99] that the strong orbit equivalence is decidable for minimal substitution subshifts. However, this result does not provide a new proof of Corollary 1.2 since for a given minimal substitution subshift (X, S) , there might be other minimal substitution subshifts that are not isomorphic to (X, S) , but that are in the same strong orbit equivalence class of (X, S) [Wer09]. Thus, decidability of the strong orbit equivalence for minimal substitution subshifts is not sufficient to decide the isomorphism between minimal substitution subshifts.

For the orbit equivalence we address the following question: Is orbit equivalence decidable for minimal substitution subshifts?

10.7 Beyond substitution and linearly recurrent subshifts

Considering Theorem 9.1 and Theorem 1.5 we wonder whether the conclusion of the second is true under the assumptions of the first. We leave this as an open question. We know that assuming zero entropy is too weak to have such a conclusion [DDMP16]. We have no counter example for finite topological rank subshifts (see [BDM10] for the definition).

10.8 For tilings

All these questions could also be asked for self-similar tilings or multidimensional substitutions. The more tractable problems would certainly concern “square multidimensional substitutions” (see, for example, [ČG86, Sal87a, Sal87b, Sal89] or, in an interesting logical context, the nice survey [BHMV94]). Moreover, the striking papers [Moz89] and [GS98] should also be mentioned as they show that in higher dimension the situation is not as “simple” as it is in dimension 1.

References

- [AB07] B. Adamczewski and Y. Bugeaud. On the complexity of algebraic numbers. I. Expansions in integer bases. *Ann. of Math. (2)*, 165:547–565, 2007.
- [ACSZ18] J.-P. Allouche, J. Cassaigne, J. Shallit, and L. Q. Zamboni. A taxonomy of morphic sequences. *arXiv:1711.10807*, 2018.
- [ARS09] J.-P. Allouche, N. Rampersad, and J. Shallit. Periodicity, repetitions, and orbits of an automatic sequence. *Theoret. Comput. Sci.*, 410:2795–2803, 2009.
- [AS03] J.-P. Allouche and J. O. Shallit. *Automatic Sequences, Theory, Applications, Generalizations*. Cambridge University Press, 2003.
- [BDM04] F. Blanchard, F. Durand, and A. Maass. Constant-length substitutions and countable scrambled sets. *Nonlinearity*, 17:817–833, 2004.
- [BDM10] X. Bressaud, F. Durand, and A. Maass. On the eigenvalues of finite rank Bratteli–Vershik dynamical systems. *Ergod. Th. & Dynam. Sys.*, 30:639–664, 2010.
- [BHMV94] V. Bruyère, G. Hansel, C. Michaux, and R. Villemaire. Logic and p -recognizable sets of integers. *Bull. Belg. Math. Soc.*, 1:191–238, 1994. Corrigendum, *Bull. Belg. Math. Soc.* **1** (1994), 577.
- [BJKR01] O. Bratteli, P. Jorgensen, K. H. Kim, and F. Roush. Decidability of the isomorphism problem for stationary AF-algebras and the associated ordered simple dimension groups. *Ergod. Th. & Dynam. Sys.*, 21:1625–1655, 2001.

- [BLR88] M. Boyle, D. Lind, and D. Rudolph. The automorphism group of a shift of finite type. *Trans. Amer. Math. Soc.*, 306:71–114, 1988.
- [Bow75] R. Bowen. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*. Lecture Notes in Mathematics, Vol. 470. Springer-Verlag, Berlin-New York, 1975.
- [Boy08] M. Boyle. Open problems in symbolic dynamics. In *Geometric and probabilistic structures in dynamics*, volume 469 of *Contemp. Math.*, pages 69–118. Amer. Math. Soc., Providence, RI, 2008.
- [Buc60] J. R. Buchi. Weak second-order arithmetic and finite automata. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, 6:66–92, 1960. Reprinted in S. Mac Lane and D. Siefkes, eds., *The Collected Works of J. Richard Buchi*, Springer-Verlag, 1990, pp. 398–424.
- [CDK17] E. Coven, M. Dekking, and M. Keane. Topological conjugacy of constant length substitution dynamical systems. *Indag. Math. (N.S.)*, 28:91–107, 2017.
- [CDKL14] E. Coven, A. Dykstra, M. Keane, and M. LeMasurier. Topological conjugacy to given constant length substitution minimal systems. *Indag. Math. (N.S.)*, 25:646–651, 2014.
- [ČG86] A. Černý and J. Gruska. Modular trellises. In G. Rozenberg and A. Salomaa, editors, *The Book of L*, pages 45–61. Springer-Verlag, 1986.
- [CH84] K. Culik and T. Harju. The ω -sequence equivalence problem for D0L systems is decidable. *J. Assoc. Comput. Mach.*, 31:282–298, 1984.
- [Cha18] E. Charlier. *First-Order Logic and Numeration Systems*, pages 89–141. Springer International Publishing, Cham, 2018.
- [CK16] V. Cyr and B. Kra. The automorphism group of a shift of subquadratic growth. *Proc. Amer. Math. Soc.*, 144:613–621, 2016.
- [CKL08] E. Coven, M. Keane, and M. Lemasurier. A characterization of the morse minimal set up to topological conjugacy. *Ergod. Th. & Dynam. Sys.*, 28:1443–1451, 2008.
- [Cob72] A. Cobham. Uniform tag sequences. *Math. Systems Theory*, 6:164–192, 1972.
- [Cov72] E. Coven. Endomorphisms of substitution minimal sets. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 20:129–133, 1971/1972.

- [CQY16] E. Coven, A. Quas, and R. Yassawi. Computing automorphism groups of shifts using atypical equivalence classes. *Discrete Anal.*, pages Paper No. 611, 28, 2016.
- [CRS12] E. Charlier, N. Rampersad, and J. Shallit. Enumeration and decidable properties of automatic sequences. *Internat. J. Found. Comput. Sci.*, 23:1035–1066, 2012.
- [DDMP16] S. Donoso, F. Durand, A. Maass, and S. Petite. On automorphism groups of low complexity minimal subshifts. *Ergod. Th. & Dynam. Sys.*, 36:64–95, 2016.
- [Dek78] M. Dekking. The spectrum of dynamical systems arising from substitutions of constant length. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 41:221–239, 1977/78.
- [Dev15] R. Devyatov. On Subword Complexity of Morphic Sequences. *arXiv:1502.02310*, February 2015.
- [DG18] F. Durand and V. Goyheneche. Decidability, arithmetic subsequences and eigenvalues for substitution subshifts. *preprint*, pages 1–20, 2018.
- [DHS99] F. Durand, B. Host, and C. Skau. Substitutive dynamical systems, Bratteli diagrams and dimension groups. *Ergod. Th. & Dynam. Sys.*, 19:953–993, 1999.
- [DL17] F. Durand and J. Leroy. The constant of recognizability is computable for primitive morphisms. *J. Integer Seq.*, 20:Art. 17.4.5, 15, 2017.
- [Dur98a] F. Durand. A characterization of substitutive sequences using return words. *Discrete Math.*, 179:89–101, 1998.
- [Dur98b] F. Durand. Sur les ensembles d’entiers reconnaissables. *J. Théor. Nombres Bordeaux*, 10:65–84, 1998.
- [Dur00] F. Durand. Linearly recurrent subshifts have a finite number of non-periodic subshift factors. *Ergod. Th. & Dynam. Sys.*, 20:1061–1078, 2000.
- [Dur12] F. Durand. HD0L ω -equivalence and periodicity problems in the primitive case (dedicated to the memory of G. Rauzy). *J. of Uniform Distribution Theory*, 7:199–215, 2012.
- [Dur13a] F. Durand. Decidability of the HD0L ultimate periodicity problem. *RAIRO - Theoretical Informatics and Applications*, 47:201–214, 2013.
- [Dur13b] F. Durand. Decidability of uniform recurrence of morphic sequences. *Internat. J. Found. Comput. Sci.*, 24:123–146, 2013.

- [Fag97a] I. Fagnot. On the subword equivalence problem for morphic words. *Discrete Appl. Math.*, 75:231–253, 1997.
- [Fag97b] I. Fagnot. Sur les facteurs des mots automatiques. *Theoret. Comput. Sci.*, 172:67–89, 1997.
- [Fog02] N. Pytheas Fogg. *Substitutions in dynamics, arithmetics and combinatorics*, volume 1794 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002. Edited by V. Berthé, S. Ferenczi, C. Mauduit and A. Siegel.
- [For97] A. Forrest. K -groups associated with substitution minimal systems. *Israel J. Math.*, 98:101–139, 1997.
- [GH55] W. H. Gottschalk and G. A. Hedlund. *Topological Dynamics*, volume 36 of *AMS Colloquium Publications*. Amer. Math. Soc., 1955.
- [GPS95] T. Giordano, I. Putnam, and C. F. Skau. Topological orbit equivalence and C^* -crossed products. *Internat. J. Math.*, 469:51–111, 1995.
- [GS98] C. Goodman-Strauss. Matching rules and substitution tilings. *Ann. of Math. (2)*, 147:181–223, 1998.
- [Hed69] G. A. Hedlund. Endomorphisms and automorphisms of the shift dynamical system. *Math. Systems Theory*, 3:320–375, 1969.
- [HJ90] R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, 1990. Corrected reprint of the 1985 original.
- [HK95] B. Hasselblatt and A. Katok. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995.
- [Hon86] J. Honkala. A decision method for the recognizability of sets defined by number systems. *RAIRO Inform. Théor. App.*, 20:395–403, 1986.
- [Hon08] J. Honkala. Cancellation and periodicity properties of iterated morphisms. *Theoret. Comput. Sci.*, 391:61–64, 2008.
- [Hos95] B. Host. Dimension groups and substitution dynamical systems. Prétirage IML 95-13, 1995.
- [HZ99] C. Holton and L. Q. Zamboni. Descendants of primitive substitutions. *Theory Comput. Systems*, 32:133–157, 1999.
- [KM16] K. Klouda and K. Medková. Synchronizing delay for binary uniform morphisms. *Theoret. Comput. Sci.*, 615:12–22, 2016.
- [KR99] K. H. Kim and F. W. Roush. The Williams conjecture is false for irreducible subshifts. *Ann. of Math. (2)*, 149:545–558, 1999.

- [LM95] D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, 1995.
- [LR13] J. Leroy and G. Richomme. A combinatorial proof of S -adicity for sequences with linear complexity. *Integers*, 13:Paper No. A5, 19, 2013.
- [MH40] M. Morse and G. A. Hedlund. Symbolic dynamics II. Sturmian trajectories. *Amer. J. Math.*, 62:1–42, 1940.
- [Mos92] B. Mossé. Puissances de mots et reconnaissabilité des points fixes d’une substitution. *Theoret. Comput. Sci.*, 99:327–334, 1992.
- [Mos96] B. Mossé. Reconnaissabilité des substitutions et complexité des suites automatiques. *Bull. Soc. Math. France*, 124:329–346, 1996.
- [Mou16] H. Mousavi. Automatic theorem proving in walnut. *CoRR*, abs/1603.06017, 2016.
- [Moz89] S. Mozes. Tilings, substitution systems and dynamical systems generated by them. *J. Analyse Math.*, 53:139–186, 1989.
- [Nek18] V. Nekrashevych. Palindromic subshifts and simple periodic groups of intermediate growth. *Ann. of Math. (2)*, 187:667–719, 2018.
- [Pan84] J.-J. Pansiot. Complexité des facteurs des mots infinis engendrés par morphismes itérés. In J. Paredaens, editor, *ICALP84*, volume 172 of *Lecture Notes in Computer Science*, pages 380–389. Springer-Verlag, 1984.
- [Pan85] J.-J. Pansiot. Subword complexities and iteration. *Bulletin of the EATCS*, 26:55–62, 1985.
- [Que10] M. Queffélec. *Substitution dynamical systems—spectral analysis. Second Edition*, volume 1294 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2010.
- [Rig14] M. Rigo. *Formal Languages, Automata and Numeration Systems 2, Applications to Recognizability and Decidability*. ISTE Wiley, 2014.
- [Sal87a] O. Salon. Suites automatiques à multi-indices. In *Sém. Théor. Nombres Bordeaux*, volume 4, pages 1–27, 1986-1987.
- [Sal87b] O. Salon. Suites automatiques à multi-indices et algébricité. *C. R. Acad. Sci. Paris*, 305:501–504, 1987.
- [Sal89] O. Salon. Quelles tuiles! (Pavages aperiodiques du plan et automates bidimensionnels). *Sém. Théor. Nombres Bordeaux*, 1:1–26, 1989.

- [ST15] V. Salo and I. Törmä. Block maps between primitive uniform and pisot substitutions. *Ergod. Th. & Dynam. Sys.*, 35:2292–2310, 2015.
- [Wer09] B. M. Werner. An example of Kakutani equivalent and strong orbit equivalent substitution systems that are not conjugate. *Discrete Contin. Dyn. Syst. Ser. S*, 2:239–249, 2009.
- [Wil73] R. F. Williams. Classification of subshifts of finite type. *Ann. of Math. (2)*, 98:120–153; errata, *ibid.* (2) 99 (1974), 380–381, 1973.