



## **Pneumonia in acute ischemic stroke patients requiring invasive ventilation: Impact on short and long-term outcomes**

J. Bourenne, N. Brule, C. Bretonniere, C. Cheval, J. Carvelli, E. Coupez, M. Cour, L. Dopeux, A. S. Dumenil, C. Dupuis, et al.

### **► To cite this version:**

J. Bourenne, N. Brule, C. Bretonniere, C. Cheval, J. Carvelli, et al.. Pneumonia in acute ischemic stroke patients requiring invasive ventilation: Impact on short and long-term outcomes. *Journal of Infection*, 2019, 79 (3), pp.220–227. 10.1016/j.jinf.2019.06.012 . hal-02466193

**HAL Id: hal-02466193**

**<https://hal.science/hal-02466193>**

Submitted on 25 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

# **Pneumonia in acute ischemic stroke patients requiring invasive ventilation: impact on short and long-term outcomes**

## **Running title**

Pneumonia in acute ischemic stroke patients

## **Author names and affiliations**

Etienne de Montmollin, MD<sup>a,b</sup>; Stephane Ruckly, MSc<sup>a</sup>; Carole Schwebel, MD, PhD<sup>c</sup>; Francois Philippart, MD, PhD<sup>d,e</sup>; Christophe Adrie, MD, PhD<sup>f</sup>; Eric Mariotte, MD<sup>g</sup>; Guillaume Marcotte, MD<sup>h</sup>; Yves Cohen, MD, PhD<sup>i</sup>; Benjamin Sztrymf, MD<sup>j</sup>; Daniel da Silva, MD<sup>k</sup>; Fabrice Bruneel, MD<sup>l</sup>; Marc Gainnier, MD, PhD<sup>m</sup>; Maité Garrouste-Orgeas, MD<sup>a,n</sup>; Romain Sonnevile, MD, PhD<sup>b,o</sup>; and Jean-François Timsit, MD, PhD<sup>a,b</sup> for the OUTCOMEREA Study Group

<sup>a</sup> UMR 1137, IAME, Paris Diderot University, Sorbonne Paris Cité, 16 rue Henri Huchard, 75018 Paris, France.

<sup>b</sup> Medical and Infectious Diseases Intensive Care Unit, AP-HP, Bichat-Claude Bernard Hospital, 46 rue Henri Huchard, 75018 Paris, France.

<sup>c</sup> Medical Intensive Care Unit, Grenoble University Hospital, Grenoble 1 University, Avenue Maquis du Grésivaudan, 38700 La Tronche, France.

<sup>d</sup> Intensive Care Unit, Saint Joseph Hospital Network, 185 Rue Raymond Losserand, 75014 Paris, France.

<sup>e</sup> UMR 9891, CNRS-CEA-Paris Saclay university, 1 Avenue De La Terrasse, 91190 Gif-sur-Yvette, France.

<sup>f</sup> Physiology department, AP-HP, Cochin Hospital, 27 Rue du Faubourg Saint-Jacques, 75014 Paris, France.

<sup>g</sup> Medical Intensive Care Unit, AP-HP, Saint-Louis Hospital, 1 Avenue Claude Vellefaux, 75010 Paris, France.

<sup>h</sup> Surgical Intensive Care Unit, Lyon University Hospital, 5 Place d'Arsonval, 69003 Lyon, France.

<sup>i</sup> Intensive Care Unit, AP-HP, Avicenne Hospital, Paris 13 University, 125 Rue de Stalingrad, 93000 Bobigny, France.

<sup>j</sup> Intensive Care Unit, AP-HP, Antoine Bécère Hospital, 157 Rue de la Porte de Trivaux, 92140 Clamart, France.

<sup>k</sup> Intensive Care Unit, Delafontaine Hospital, 2 rue du Dr Delafontaine, 93200 Saint-Denis, France.

<sup>l</sup> Intensive Care Unit, André Mignot Hospital, 177 Rue de Versailles, 78150 Le Chesnay, France.

<sup>m</sup> Intensive Care Unit, AP-HM, La Timone Hospital, Aix-Marseille University, 278 Rue Saint-Pierre, 13005 Marseille, France.

<sup>n</sup> Medical Unit, French British Hospital, 4 rue Kléber, 92300 Levallois-Perret, France.

<sup>o</sup> UMR 1148, LVTS, Paris Diderot University, Sorbonne Paris Cité, 46 rue Henri Huchard, 75018 Paris, France.

### **Corresponding author**

Dr Etienne de Montmollin, Service de Réanimation Médicale et infectieuse

Hôpital Bichat–Claude Bernard

46 rue Henri Huchard, 75018 Paris

Phone: + 33 1 40 25 77 03 / Fax: + 33 1 40 25 88 37

Mail: [etienne.demontmollin@aphp.fr](mailto:etienne.demontmollin@aphp.fr)

## ABSTRACT

**Objectives:** To describe the epidemiology and prognostic impact of pneumonia in acute ischemic stroke patients requiring invasive mechanical ventilation.

**Methods:** Retrospective analysis from a prospective multicenter cohort study of critically ill patients with acute ischemic stroke requiring invasive mechanical ventilation at admission. Impact of pneumonia was investigated using Cox regression for 1-year mortality, and competing risk survival models for ICU mortality censored at 30-days.

**Results:** We included 195 patients. Stroke was supratentorial in 62% and 64% of patients had a Glasgow coma scale score <8 on admission. Mortality at day-30 and 1 year were 56%, and 70% respectively. Post-stroke pneumonia was identified in 78 (40%) patients, of which 46/78 (59%) episodes were present at ICU admission. Post-stroke pneumonia was associated with an increase in 1-year mortality (adjusted HR 1.49, 95%CI [1.01 – 2.20]). Post-stroke pneumonia was not associated with ICU mortality but was associated with a 1.6-fold increase in ICU length of stay (CSHR 0.62 [0.39– 0.99], p=0.06).

**Conclusions:** In ischemic stroke patients requiring invasive ventilation, pneumonia occurred in 40% of cases and was associated with a 49% increase in 1-year mortality. Post-stroke pneumonia did not impact day-30 mortality but increased ICU length of stay.

## **KEYWORDS**

Pneumonia

Stroke

Critical Care

Mechanical Ventilation

Mortality

Epidemiology

## INTRODUCTION

Infection is a frequent medical complication after stroke <sup>1</sup> with an observed pooled frequency from a recent meta-analysis of 21%, most frequently involving the lungs (12%) and urinary tract (8%) <sup>2</sup>. The mechanisms leading to post-stroke infections are complex and include a combination of brain injury-induced immunosuppression, neurological impairment that fosters lung aspiration, immobilization, and use of urinary catheters <sup>3,4</sup>. Post-stroke infections, in general, are associated with lower survival and less favorable neurologic outcomes <sup>5</sup>. When analyzing by infection type, post-stroke pneumonia but not urinary tract infections seem to be the main driver of this negative impact <sup>5</sup>.

Studies focusing on critically ill stroke patients report the highest infection rates <sup>4</sup>, ranging from 9 to 57 %. Single-center studies suggest that occurrence of post-stroke pneumonia in the intensive care unit (ICU) is associated with an increase of ICU length of stay <sup>6-8</sup> and hospital mortality <sup>6,9</sup>, and poorer functional outcomes in survivors <sup>9</sup>. However, studies conducted on post-stroke infection in critically ill patients have many biases, including retrospective design, non-standardized definition of pneumonia, small number of patients, pooled analysis of ischemic/hemorrhagic strokes, and variable rates of invasive mechanical ventilation (ranging from 2 to 100%), indicating variable severity of studied populations. Moreover, most of these studies only focused on short-term outcomes, namely hospital mortality. Finally, decisions of withholding and withdrawal of life-sustaining treatments during ICU stay that may impact survival analyses were never reported or considered in most of studies, despite being a major confounder <sup>10</sup>. Finally, there are very few data regarding the impact of ICU post-stroke pneumonia on long term survival and functional outcomes.

Therefore, we aimed to describe the epidemiology and impact on short and long-term prognosis of post-stroke pneumonia in critically ill ischemic stroke patients requiring invasive mechanical ventilation and taking withholding/withdrawal of life-sustaining treatments into account. Some of the results of this study have been previously reported in the form of an abstract (French Intensive Care Society annual congress, Paris, January 2019).

## **MATERIALS AND METHODS**

### **Patient data source.**

This study was conducted using data from the French prospective multicenter (n=30 ICUs) OUTCOMEREA database, from patients included between 1997 and 2016. The OUTCOMEREA database has been described in previous publications, and has been approved by the French Advisory Committee for Data Processing in Health Research (CCTIRS) and the French Informatics and Liberty Commission (CNIL, registration no. 8999262) <sup>11,12</sup>. The database protocol was submitted to the Institutional Review Board of the Clermont-Ferrand University hospital (Clermont-Ferrand, France), who waived the need for informed consent (IRB no. 5891).

### **Study population and definitions.**

We included adult patients with acute ischemic stroke, admitted to the Intensive Care Unit (ICU) and requiring invasive mechanical ventilation. All ICU stays in the database were screened for a diagnosis of stroke, using the International Classification of Diseases 10th Revision (ICD-10) codes I63 ("Cerebral infarction") and I64 ("Stroke, not specified as hemorrhage or infarction"). ICU stays were considered as related to acute stroke in cases of (1) direct ICU admission following stroke onset, or (2) ICU admission during the initial acute care hospital stay following stroke onset. We excluded patients without hospitalization reports, and if stroke was related to traumatic brain injury.

Severity of illness was graded at ICU admission with use of the Simplified Acute Physiology Score (SAPS II) <sup>13</sup>. Organ failure was assessed with each component of the sepsis-related organ failure assessment (SOFA) score, and a modified SOFA score was calculated, excluding the neurological component <sup>14,15</sup>. Neurologic status was assessed using the



modified Rankin Scale (mRS) <sup>16</sup>. Pneumonia was defined as new or persistent pulmonary infiltrates on chest X-ray combined with purulent tracheal secretions and/or fever or hypothermia (body temperature greater  $\geq 38.5$  or  $\leq 36.5$  °C, respectively) and/or leukocytosis or leukopenia (white blood cells count  $\geq 10 \times 10^9$  or  $\leq 4 \times 10^9$ /L, respectively). **Community-acquired** pneumonia (CAP) was defined as pneumonia occurring after stroke onset and before or within 48 hours of hospital admission. Hospital-acquired pneumonia (HAP) was defined as pneumonia occurring after 48 hours of hospital length of stay in patients without mechanical ventilation or with a duration of mechanical ventilation less than 48 hours. Ventilator-acquired pneumonia was defined as pneumonia occurring after 48 hours of mechanical ventilation <sup>17</sup>. **ICU admission** pneumonia was defined as pneumonia occurring within 1 day of ICU admission (**irrespective of the type of admission (direct admission from home/Emergency department or indirect from a hospitalization ward)**). Early-onset pneumonia (EOP) was defined as pneumonia occurring between day 2 and 7 of ICU admission, and late-onset pneumonia (LOP) as pneumonia occurring from day 8 of ICU admission <sup>18</sup>. Sepsis was defined as infection associated with organ dysfunction (SOFA score  $\geq 2$ ) <sup>19</sup>. Septic shock was defined as sepsis with persisting hypotension requiring vasopressors to maintain MAP  $\geq 65$  mmHg and a serum lactate level  $> 2$  mmol/L despite adequate volume resuscitation.

#### **Data collection.**

Data were prospectively collected at admission (demographics, chronic diseases, admission features, baseline severity indexes, admission diagnosis, and admission type), and on a daily basis throughout the ICU stay (clinical and biological parameters, assessment of organ functions, requirement for MV, fluid challenges, ICU-acquired infections, length of

stay (LOS), decision to withhold or withdraw life-sustaining therapies, and vital status at ICU and hospital discharge), through an anonymized electronic case report form using the Vigirea, Rhea and e-Rhea softwares (OutcomeRea, Aulnay-sous-Bois, France). Long term survival after hospital discharge was collected by each local investigator. For each stay, we retrospectively collected the following data in the medical charts: (1) stroke history, including date of stroke, localization, specific treatment (i.e. thrombolysis, hemicraniectomy, etc.), (2) chronic diseases potentially related to stroke, including arterial hypertension, atrial fibrillation, history of ischemic or hemorrhagic stroke, (3) modified Rankin scale (mRS) at ICU discharge.

### **Statistical analysis.**

Quantitative variables are presented as median, 1<sup>st</sup> and 3<sup>rd</sup> quartiles, and compared between groups with the Wilcoxon test. Qualitative variables are presented as frequency and corresponding percentage and compared with the Chi-square test or Fisher exact test as appropriate. Association between the occurrence of post-stroke pneumonia and 30-day ICU mortality and length of stay was assessed using a cause specific hazard model. It provided a cause-specific hazard ratio (CSHR) which estimates the direct effect of the time-dependent exposure of interest (i.e., post-stroke pneumonia) on various outcomes (i.e., ICU mortality and ICU discharge)<sup>20,21</sup>. We considered that determinants of 1-year survival were not affected by competing risks, and used a multivariable Cox regression model. Competing risk models were adjusted with non-collinear and clinically pertinent factors associated ( $p < 0.05$ ) with the outcome of interest in univariate analysis and no selection procedure. Cox model for 1-year mortality was adjusted for potentially confounding factors (non-collinear and clinically pertinent factors associated ( $p < 0.1$ ) with the outcome of interest in univariate

analysis) with a backward selection procedure. The occurrence of pneumonia and decisions to withhold/withdraw life-sustaining treatments were handled as time-dependent variables. Only the first episode of pneumonia was considered for analysis, and subsequent episodes were disregarded. For missing data, simple imputation with median/most frequent method was used when missing was completely at random with less than 10% missing value per variable <sup>22</sup>. All statistical analyses were carried out with SAS 9.4 (SAS Institute Inc., Cary, NC, USA). A *p*-value of 0.05 and lower was considered statistically significant.

## RESULTS

### Patients.

Among 22106 ICU admissions from 30 ICUs over the study period, we identified 229 stays involving acute ischemic stroke. We excluded 34 patients without invasive ventilation at admission and included 195 patients from 10 ICUs in the analysis (Supplemental digital content, **SDC 1**). Two ICUs represented 150/195 (77%) of included patients (**SDC 2**). Patients admitted during the second half of the study period (2007-2016) accounted for 71% of the cohort. Baseline characteristics of patients are presented in **Table 1**. Patients were predominantly males (68%), aged 69.4 [61.2 – 76.5] years. Ischemic stroke was supratentorial in 121 (62%) cases, and the median time from stroke onset to ICU admission was 2 [1 – 4] days. Patients received specialized referral for acute stroke therapy (i.e. thrombolysis or endovascular treatment) or neurosurgery in 190 (97%) cases, and acute stroke therapy was carried out in 35 (18%) cases. Patients were admitted mainly for coma (56%) and respiratory failure (20%), with a SAPS2 of 56 [45 – 66] and a SOFA score of 7 [5 – 9]. Glasgow Coma Scale score at admission was 6 [3 – 10]. During ICU stay, 92 (47%) patients required vasopressor support, and 17 (9%) renal replacement therapy (**Table 2**). Duration of invasive mechanical ventilation was 5 [3 – 12] days. A decision to withhold/withdraw life-sustaining treatments was made in 65 (33%) cases, with a delay of 5 [2 – 8] days.

### Epidemiology of post-stroke pneumonia.

During ICU stay, 78 (40%) patients presented at least one pneumonia episode, among which 12 presented 2 or more episodes (**Table 3**). Among those 78 episodes, pneumonia was diagnosed at ICU admission in 46/78 (59%) cases. Of the 32/78 pneumonias acquired during ICU stay, 11/32 (34%) were early-onset and 21/32 (66%) were late-onset. The supplemental

digital content 3 (**SDC 3**) shows the distribution of pneumonia (ICU admission pneumonia/EOP/LOP) according to the type of ICU admission (direct or from hospitalization ward). Cases of post-stroke pneumonia were classified as CAP, HAP and VAP in 22 (28%), 28 (36%) and 28 (36%) cases, respectively. Pneumonia was associated with sepsis or septic shock in 62 (79%) cases. Diagnostic incidence of pneumonia during ICU stay was highest at ICU admission, as shown by the cumulative incidence curve presented in the Supplemental digital content (**SDC 4**). Among pneumonia episodes (n=78), 56 had a microbiological identification, and 67 pathogens were identified, 25 (37%) being gram-positive cocci and 42 (63%) gram-negative bacilli (Supplemental digital content, **SDC 5**). When analyzing bacterial identification according to the date of pneumonia onset, we show that cases of pneumonia diagnosed at ICU admission appeared to be more frequently undocumented than EOP and LOP (20/53 (38%) vs 2/24 (8%) vs 0/12 (0%), p=0.14). Conversely, EOP and even more LOP appeared to be more frequently due to *Enterobacteriaceae* than ICU admission pneumonia (11/53 (21%) vs 7/24 (29%) vs 5/12 (42%), p=0.24). The same pattern was found with non-fermenting gram-negative bacilli (2/53 (4%) vs 2/24 (8%) vs 5/12 (42%), p=0.23) (**Figure 1**). Having had pneumonia at day 1 or 2 was not associated in univariate analysis with an increased risk of pneumonia between day 3 and 30 (20% vs 19%, p=0.92). In univariate analysis, intravenous thrombolysis was associated with an increased frequency of post-stroke pneumonia (**Table 1**, 15% vs 4%, p=0.02). Characteristics of patients who received intravenous thrombolysis are presented in the supplemental digital content (**SDC 6**) and show that patients with intravenous thrombolysis were younger and more frequently admitted from the hospitalization ward before ICU admission. ICU admission for respiratory failure was also associated with an increased frequency of post-stroke pneumonia (**Table 1**, 35% vs 11%, p<0.01). Patients admitted for respiratory failure were more frequently

admitted from hospitalization ward (75% vs 48%,  $p<0.01$ ), with a longer delay between hospital and ICU admission (2 [1 – 8] days vs 1 [1 – 3] days,  $p=0.03$ ) (**SDC 7**). Patients admitted for respiratory failure had lower neurological failure (neurologic SOFA 2 [0.5 – 3] vs 3 [2 – 4],  $p=0.03$ ), and higher respiratory (respiratory SOFA 3 [2 – 4] vs 2 [0 – 3],  $p<0.01$ ) and cardiovascular (cardiovascular SOFA 3 [0 – 4] vs 1 [0 – 3],  $p=0.01$ ) failures. Among the 78 patients with a pneumonia episode, only 4 (5%) had had a prior decision to withhold/withdraw life-sustaining treatments. Occurrence of pneumonia was associated with neither an increase in the risk of decisions to withhold or withdraw life-sustaining treatments (36% vs 32%,  $p=0.54$ ), nor the delay of such decision (5 [3 – 9] days vs 4 [2 – 7],  $p=0.33$ ).

#### **Prognostic impact of the occurrence of post-stroke pneumonia.**

ICU, 30-day, hospital and 1-year mortality rates were respectively 53%, 56%, 63%, and 70%. Kaplan-Meier curves of 30-day and 1-year survival are presented as supplemental digital content (**SDC 8**). ICU length of stay of the total population was 7 [4 – 15] days, and 12 [6 – 23] days in ICU survivors only. Good neurologic outcome at ICU discharge, defined as a mRS score  $\leq 3$ , was present in 22/195 (11%) patients. In ICU survivors, good neurologic outcome at ICU discharge was more frequent in patients without post-stroke pneumonia (17/49 (35%) vs 5/43 (12%),  $p<0.01$ ). Median follow-up for the vital status of ICU survivors was 747 [58 – 2112] days. At 1-year, vital status was available for 182 (93%) patients.

Univariate analysis of factors associated with 30-day mortality is presented in the Supplemental digital content (**SDC 9**). Of note, neither age nor stroke characteristics were associated with 30-day mortality. We show that the period of inclusion was associated with both 30-day and 1-year mortality. We explored this association (Supplemental digital

content, **SDC 10**) and showed that compared to the first period (1997-2006), patients admitted during the second period (2007-2016) had higher admission SOFA scores (8 [6 – 10] vs 6 [1 – 8],  $p<0.1$ ). There was no difference between the 2 inclusion periods with regard to age (69.1 [61 – 77.3] vs 70.4 [61.6 – 75.5],  $p=0.71$ ), GCS score (6.5 [3 – 10] vs 5 [3 – 10],  $p=0.13$ ) and pneumonia rate (40% vs 39%,  $p=0.9$ ).

In the multivariable competing risk survival analysis, the occurrence of pneumonia was associated with an adjusted ICU discharge cause specific hazard ratio (CSHR) of 0.62 [0.39–0.99], corresponding to a 1.6-fold increase in the duration of ICU stay (**Table 4**). The occurrence of pneumonia was associated with an adjusted mortality CHSR of 0.96 [0.59–1.57] indicating the absence of effect of post-stroke pneumonia on ICU mortality (**Table 4**). In ICU survivors, pneumonia during ICU stay was significantly associated with a lower rate of good neurological functional outcome at ICU discharge (12% vs 35%,  $p<0.01$ ) (**Table 2**). Long term impact of post-stroke pneumonia was assessed at 1 year. In a Cox proportional hazard model adjusted on decisions to withhold/withdraw life-sustaining treatments, GCS score at admission and period of inclusion, post-stroke pneumonia was significantly associated with 1-year mortality (adjusted HR 1.49, 95%CI [1.01 – 2.20],  $p=0.046$ ) (Supplemental digital content, **SDC 11**).

## DISCUSSION

In this retrospective analysis of a prospective database of critically ill ischemic stroke patients requiring invasive mechanical ventilation, we show that post-stroke pneumonia occurs in 40% of cases, and was associated with an increase in 1-year mortality. On the short term, post stroke-pneumonia was associated with an increased ICU length of stay but not 30-day mortality. Post-stroke pneumonia seemed to be associated with a reduced functional outcome at ICU discharge.

We show that the prognosis of ischemic stroke patients requiring invasive mechanical ventilation is poor, with a hospital mortality of 63%. The 1-year mortality of 70% suggests that the vast majority of death occur during the initial hospital stay. We report a higher hospital mortality rate than the 47% from a recent United States population-based study<sup>23</sup> of mechanically ventilated stroke patients. Hospital mortality rates in previous studies focusing on post-stroke pneumonia and with high mechanical ventilation rates appear to be much lower, ranging from 14 to 35%<sup>6,8,24,25</sup>. This is explained by the fact that these studies excluded patients that died in the first 48 hours. Because our multivariable models were adjusted on decision to withhold/withdraw life-sustaining treatments and because the incidence of post-stroke pneumonia is maximum at ICU admission, we chose to keep in the analysis all patients admitted to the ICU.

Post-stroke pneumonia rates reported in the literature are highly variable, ranging from 9.5 to 75%<sup>4</sup>. This is due to the variability of mechanical ventilation rate in the studied populations and to the non-standardized definition of post-stroke pneumonia. We report here a pneumonia rate of 40% in a population of mechanically ventilated patients from 10 non-neurologic ICUs, using a standardized methodology for pneumonia diagnosis using



prospectively collected data. We show that incidence of pneumonia is the highest at ICU admission. The EOP rate of 26% we report is in accordance with published rates (20-60%) in traumatic brain injury and subarachnoid hemorrhage patients<sup>26,27</sup>. When considering only VAP (**Table 3**), we report a rate of 14% which is similar to VAP rate reported in general ICU populations<sup>28,29</sup>. Our univariate analysis shows that intravenous thrombolysis appears to be associated with an increased frequency of post-stroke pneumonia. This association has not been noted before, and is likely due to the fact that patients who undergo intravenous thrombolysis are a subset of less severe patients (intravenous thrombolysis is contraindicated for older and comatose patients) at hospital admission, and are more frequently admitted to the ICU for respiratory failure.

Contrarily to the majority of studies performed in stroke units and acute hospital wards where post-stroke pneumonia was an independent risk factor for hospital mortality<sup>4</sup>, MICU/NICU studies<sup>6,8,9,24,30,31</sup> have not shown such an association except for one with only 13% of mechanically ventilated patients and an inadequate methodology (univariate analysis only)<sup>9</sup>. In accordance with previously published ICU studies, we show that post-stroke pneumonia has no impact on short-term mortality, but increases ICU length of stay. One of the reasons could be the higher frequency of post-stroke pneumonia in patients admitted for respiratory failure compared to patients admitted for coma. Indeed, we hypothesize that the negative impact of post-stroke pneumonia could be balanced by the better prognosis of patients admitted for respiratory failure compared to patients admitted for coma<sup>32</sup>. It has also been shown previously in general ICU populations that EOP were not associated with ICU or hospital mortality, but increased ICU length of stay<sup>18,33,34</sup>.

As for long-term impact, the majority of MICU/NICU studies have used only in-hospital mortality<sup>8,24,30,31</sup>, which is not the preferred endpoint. We chose to use 1-year mortality as the endpoint, and show that post-stroke pneumonia is associated with an increase in mortality. **This is, to the best of our knowledge, the first report of an impact of ICU post-stroke pneumonia on long term mortality of stroke patients<sup>4</sup>.** It is interesting to note the discrepancy between the fact that a majority of deaths occur before day-30 and the difference in the effect of pneumonia on short-term and long-term mortality. We believe that these early deaths are directly related to the acute neurologic injury of the ischemic stroke, while post-stroke pneumonia impacts survival of patients that have survived these initial critical days. This hypothesis is comforted by the fact that ICU survivors exposed to a pneumonia had a worse mRS score at ICU discharge.

The strengths of our study include a large multicenter population from a high-quality database, a focus on a well-defined population of ischemic stroke patients requiring invasive mechanical ventilation, adjustment for decisions to withhold/withdraw life-sustaining treatments, and use of an adequate statistical methodology. Our study has also limitations. First, the OUTCOMEREA database has not been built specifically for stroke studies, and all data regarding stroke characteristics has been retrospectively recorded from hospitalization records. Therefore, data on potentially useful scores for prognostication in this setting, such as the NIHSS scores, are lacking<sup>35</sup>. Furthermore, only long-term vital status was available, and evaluation of long-term functional outcome with an appropriate tool (i.e. the modified Rankin scale) would have been more relevant. Second, the inclusion period of 20 years is very wide, and management of stroke patients has greatly improved over the past decades (e.g., development of stroke units, thrombolysis, mechanical thrombectomy...). However, 77% of our cohort has been included between 2007 and 2016, and analyses have been

adjusted on the inclusion period when necessary. Third, diagnostic adequacy of pneumonia in ICU is difficult, and despite the use of prospective standard screening criteria for the diagnosis of pneumonia, we cannot exclude that some pneumonia cases were misdiagnosed. In the same line, discrimination between aspiration pneumonia and chemical pneumonitis is complex and it cannot be excluded that pneumonitis cases were wrongfully classified as pneumonia. Fourth, we were not able to adjust our results for left informative censoring (that is, critically ill acute stroke patients not referred to the ICU for reasons such as decisions to withhold/withdraw life-sustaining treatments).

## **CONCLUSIONS**

In this retrospective analysis of a prospective database of critically ill ischemic stroke patients requiring invasive mechanical ventilation, we show that post-stroke pneumonia occurs in 40% of cases, and is associated with an increase in 1-year mortality. Post-stroke pneumonia was not associated with 30-day ICU mortality, but increased ICU length of stay. These results suggest that early deaths are directly related to the acute neurologic injury while the detrimental effect of post-stroke pneumonia impacts survival of patients that have survived these initial critical days. Further studies are needed to properly assess the effect of post-stroke pneumonia on long-term functional outcome in ICU survivors.

## AKNOWLEDGEMENTS

**Members of the OUTCOMEREA Study Group—Scientific Committee:** Jean-François Timsit (Medical and Infectious Diseases ICU, Bichat-Claude Bernard Hospital, Paris, France; UMR 1137 Inserm-Paris Diderot university IAME, F75018, Paris); Elie Azoulay (Medical ICU, Saint Louis Hospital, Paris, France); Maïté Garrouste-Orgeas (ICU, Saint-Joseph Hospital, Paris, France); Jean-Ralph Zahar (Infection Control Unit, Angers Hospital, Angers, France); Christophe Adrie (Physiology, Cochin Hospital, Paris, France); Michael Darmon (Medical ICU, Saint Etienne University Hospital, St Etienne, France); and Christophe Clec'h (ICU, Avicenne Hospital, Bobigny, and UMR 1137 Inserm, Paris Diderot university IAME, F75018, Paris, France).

**Biostatistical and information system expertise:** Jean-Francois Timsit (Medical and Infectious Diseases ICU, Bichat-Claude Bernard Hospital, Paris, France; UMR 1137 Inserm - Paris Diderot university IAME, F75018, Paris); Corinne Alberti (Medical Computer Sciences and Biostatistics Department, Robert Debré Hospital, Paris, France); Adrien Français (Integrated Research Center U823, Grenoble, France); Aurélien Vesin (OUTCOMEREA organization and Integrated Research Center U823, Grenoble, France); Stephane Ruckly (OUTCOMEREA organization and Inserm UMR 1137 IAME, F75018, Paris); Sébastien Bailly (Grenoble university hospital Inserm UMR 1137 IAME, F75018, Paris) and Christophe Clec'h (ICU, Avicenne Hospital, Bobigny, and Inserm UMR 1137 IAME, F75018, Paris, France); Frederik Lecorre (Supelec, France); Didier Nakache (Conservatoire National des Arts et Métiers, Paris, France); and Aurélien Van-nieuwenhuyze (Tourcoing, France).

**Investigators of the OUTCOMEREA database:** Dr Romain HERNU, Christophe Adrie (ICU, CH Melun, and Physiology, Cochin Hospital, Paris, France); Carole Agasse (medical ICU, university hospital Nantes, France); Bernard Allaouchiche (ICU, Pierre benite Hospital, Lyon, France); Olivier Andremont (ICU, Bichat Hospital, Paris, France); Pascal Andreu (CHU Dijon, Dijon, France); Laurent Argaud (Medical ICU, Hospices Civils de Lyon, Lyon, France); Claire Ara-Somohano (Medical ICU, University Hospital, Grenoble, France); Elie Azoulay (Medical ICU, Saint Louis Hospital, Paris, France); François Barbier (medical-surgical ICU, Orleans, France), Déborah Boyer (ICU, CHU Rouen, France), Jean-Pierre Bedos (ICU, Versailles

Hospital, Versailles, France); Thomas Baudry (Medial ICU, Edouard Heriot hospital, Lyon France), Jérôme Bedel (ICU, Versailles Hospital, Versailles, France), Julien Bohé (ICU, Hôpital Pierre Benite, Lyon France), Lila Bouadma (ICU, Bichat Hospital, Paris, France); Jeremy Bourenne (Réanimation des urgencies, Timone-2; APMH, Marseille, France); Noel Brule (medical ICU, university hospital Nantes, France); Cédric Brétonnière (medical ICU, university hospital Nantes, France); Christine Cheval (ICU, Hyeres Hospital, Hyeres, France); Julien Carvelli (Réanimation des urgencies, Timone-2; APMH, Marseille, France); Christophe Clec'h (ICU, Avicenne Hospital, Bobigny, France); Elisabeth Coupez (ICU, G Montpied Hospital, Clermont-Ferrand, France); Martin Cour Medial ICU, Edouard Heriot hospital, Lyon France), Michael Darmon (ICU, Saint Etienne Hospital, Saint Etienne, France); Etienne de Montmollin (Bichat hospital and UMR 1137 Inserm –Paris Diderot university IAME, F75018, Paris, France), Loa Dopeux (ICU, G Montpied Hospital, Clermont-Ferrand, France); Anne-Sylvie Dumenil (Antoine Béclère Hospital, Clamart, France); Claire Dupuis (Bichat hospital and UMR 1137 Inserm –Paris Diderot university IAME, F75018, Paris, France), Jean-Marc Forel (AP HM, Medical ICU, Hôpital Nord Marseille), Marc Gainnier (Réanimation des urgencies, Timone-2; APMH, Marseille, France), Charlotte Garret (medical ICU, university hospital Nantes, France); Steven Grangé (ICU, CHU Rouen, France), Antoine Gros (ICU, Versailles Hospital, Versailles, France), Akim Haouache (Surgical ICU, H Mondor Hospital, Creteil, France); Romain Hernu (Medical ICU, Hospices Civils de Lyon, Lyon, France); Tarik Hissem (ICU, Eaubonne, France), Vivien Hon Tua Ha (ICU, CH Meaux, France); Sébastien Jochmans (ICU, CH Melun); Jean-Baptiste Joffredo (ICU, G Montpied Hospital, Clermont-Ferrand, France); Hatem Kallel (ICU, Cayenne General Hospital, Cayenne, France); Guillaume Lacave (ICU, Versailles Hospital, Versailles, France), Alexandre Lautrette (ICU, G Montpied Hospital, Clermont-Ferrand, France); Virgine Lemiale (Medical ICU, Saint Louis Hospital, Paris, France); Mathilde Lermuzeaux (ICU, Bichat Hospital, Paris, France), Guillaume Marcotte (Surgical ICU, Hospices Civils de Lyon, Lyon, France); Jordane Lebut (ICU, Bichat Hospital, Paris, France); Maxime Lugosi (Medical ICU, University Hospital Grenoble, Grenoble, France); Eric Magalhaes (ICU, Bichat Hospital, Paris, France), Sibylle Merceron (ICU, Versailles Hospital, Versailles, France), Bruno Mourvillier (ICU, Bichat Hospital, Paris, France); Benoît Misset (ICU, Saint-Joseph Hospital, Paris, France and Medical ICU CHU Rouen, France); Bruno Mourvillier (ICU, Bichat Hospital, Paris, France); Mathilde Neuville (ICU, Bichat Hospital, Paris, France), Laurent Nicolet (medical ICU, university hospital Nantes, France); Johanna Oziel (Medico-surgical

ICU, hôpital Avicenne APHP, Bobigny, France), Laurent Papazian (Hopital Nord, Marseille, France), Benjamin Planquette (pulmonology ICU, George Pompidou hospital Hospital, Paris, France); Jean-Pierre Quenot (CHU Dijon, Dijon, France); Aguila Radjou (ICU, Bichat Hospital, Paris, France), Marie Simon (Medial ICU, Edouard Heriot hospital, Lyon France), Romain Sonnevile (ICU, Bichat Hospital, Paris, France), Jean Reignier (medical ICU, university hospital Nantes, France); Bertrand Souweine (ICU, G Montpied Hospital, Clermont-Ferrand, France); Carole Schwebel (ICU, A Michallon Hospital, Grenoble, France); Shidasp Siami (ICU, Eaubonne, France); Roland Smonig (ICU, Bichat Hospital, Paris, France); Gilles Troché (ICU, Antoine Béchère Hospital, Clamart, France); Marie Thuong (ICU, Delafontaine Hospital, Saint Denis, France); Guillaume Thierry (ICU, Saint-Louis Hospital, Paris, France); Dany Toledano (ICU, Gonesse Hospital, Gonesse, France); Guillaume Van Der Meersch, Medical Surgical ICU, university hospital Avicenne), Marion Venot (Medical ICU, Saint Louis Hospital, Paris, France); Olivier Zambon (medical ICU, university hospital Nantes, France).

**Study Monitors:** Julien Fournier, Caroline Tournegros, Stéphanie Bagur, Mireille Adda, Vanessa Vindrieux, Sylvie de la Salle, Pauline Enguerrand, Loic Ferrand, Vincent Gobert, Stéphane Guessens, Helene Merle, Nadira Kaddour, Boris Berthe, Samir Bekkhouche, Kaouttar Mellouk, Mélaine Lebrazic, Carole Ouisse, Diane Maugars, Christelle Aparicio, Igor Theodose, Manal Nouacer, Veronique Deiler, Myriam Moussa, Atika Mouaci, Nassima Viguier, Fariza Lamara and Sophie Letrou.

## REFERENCES

- 1 Chamorro Ángel, Urra Xabier, Planas Anna M. Infection After Acute Ischemic Stroke: A Manifestation of Brain-Induced Immunodepression. *Stroke* 2007;**38**(3):1097–103. Doi: 10.1161/01.STR.0000258346.68966.9d.
- 2 Badve Monica S, Zhou Zien, van de Beek Diederik, Anderson Craig S, Hackett Maree L. Frequency of post-stroke pneumonia: Systematic review and meta-analysis of observational studies. *Int J Stroke* 2019;**14**(2):125–36. Doi: 10.1177/1747493018806196.
- 3 Emsley Hedley CA, Hopkins Stephen J. Acute ischaemic stroke and infection: recent and emerging concepts. *Lancet Neurol* 2008;**7**(4):341–53. Doi: 10.1016/S1474-4422(08)70061-9.
- 4 Hannawi Yousef, Hannawi Bashar, Rao Chethan P. Venkatasubba, Suarez Jose I., Bershad Eric M. Stroke-Associated Pneumonia: Major Advances and Obstacles. *Cerebrovasc Dis* 2013;**35**(5):430–43. Doi: 10.1159/000350199.
- 5 Vermeij Frederique H., Scholte op Reimer Wilma J. M., de Man Peter, van Oostenbrugge Robert J., Franke Cees L., de Jong Gosse, et al. Stroke-associated infection is an independent risk factor for poor outcome after acute ischemic stroke: data from the Netherlands Stroke Survey. *Cerebrovasc Dis* 2009;**27**(5):465–71. Doi: 10.1159/000210093.
- 6 Berger Benjamin, Gumbinger Christoph, Steiner Thorsten, Sykora Marek. Epidemiologic features, risk factors, and outcome of sepsis in stroke patients treated on a neurologic intensive care unit. *J Crit Care* 2014;**29**(2):241–8. Doi: 10.1016/j.jcrc.2013.11.001.
- 7 Yang Chih-Chieh, Shih Nai-Ching, Chang Wen-Chiung, Huang San-Kuei, Chien Ching-Wen. Long-term medical utilization following ventilator-associated pneumonia in acute stroke and traumatic brain injury patients: a case-control study. *BMC Health Serv Res* 2011;**11**(1):289. Doi: 10.1186/1472-6963-11-289.
- 8 Upadya Anupama, Thorevska Natalya, Sena Kanaga N., Manthous Constantine, Amoateng-Adjepong Yaw. Predictors and consequences of pneumonia in critically ill patients



with stroke. *J Crit Care* 2004;**19**(1):16–22.

9 Hilker Ruediger, Poetter Carsten, Findeisen Nahide, Sobesky Jan, Jacobs Andreas, Neveling Michael, et al. Nosocomial Pneumonia After Acute Stroke: Implications for Neurological Intensive Care Medicine. *Stroke* 2003;**34**(4):975–81. Doi: 10.1161/01.STR.0000063373.70993.CD.

10 Kelly A. G., Hoskins K. D., Holloway R. G. Early stroke mortality, patient preferences, and the withdrawal of care bias. *Neurology* 2012;**79**(9):941–4. Doi: 10.1212/WNL.0b013e318266fc40.

11 Lautrette Alexandre, Garrouste-Orgeas Maïté, Bertrand Pierre-Marie, Goldgran-Toledano Dany, Jamali Samir, Laurent Virginie, et al. Respective impact of no escalation of treatment, withholding and withdrawal of life-sustaining treatment on ICU patients' prognosis: a multicenter study of the Outcomerea Research Group. *Intensive Care Med* 2015;**41**(10):1763–72. Doi: 10.1007/s00134-015-3944-5.

12 Truche Anne-Sophie, Darmon Michael, Bailly Sébastien, Clec'h Christophe, Dupuis Claire, Misset Benoit, et al. Continuous renal replacement therapy versus intermittent hemodialysis in intensive care patients: impact on mortality and renal recovery. *Intensive Care Med* 2016;**42**(9):1408–17. Doi: 10.1007/s00134-016-4404-6.

13 Le Gall J. R., Lemeshow S., Saulnier F. A new Simplified Acute Physiology Score (SAPS II) based on a European/North American multicenter study. *JAMA* 1993;**270**(24):2957–63.

14 Timsit Jean-François, Fosse Jean-Philippe, Troché Gilles, De Lassence Arnaud, Alberti Corinne, Garrouste-Orgeas Maïté, et al. Calibration and discrimination by daily Logistic Organ Dysfunction scoring comparatively with daily Sequential Organ Failure Assessment scoring for predicting hospital mortality in critically ill patients. *Crit Care Med* 2002;**30**(9):2003–13. Doi: 10.1097/01.CCM.0000025210.75241.3E.

15 Vincent J. L., Moreno R., Takala J., Willatts S., De Mendonça A., Bruining H., et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996;**22**(7):707–10.

- 16 van Swieten J. C., Koudstaal P. J., Visser M. C., Schouten H. J., van Gijn J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke J Cereb Circ* 1988;**19**(5):604–7.
- 17 Torres Antoni, Niederman Michael S., Chastre Jean, Ewig Santiago, Fernandez-Vandellos Patricia, Hanberger Hakan, et al. International ERS/ESICM/ESCMID/ALAT guidelines for the management of hospital-acquired pneumonia and ventilator-associated pneumonia: Guidelines for the management of hospital-acquired pneumonia (HAP)/ventilator-associated pneumonia (VAP) of the European Respiratory Society (ERS), European Society of Intensive Care Medicine (ESICM), European Society of Clinical Microbiology and Infectious Diseases (ESCMID) and Asociación Latinoamericana del Tórax (ALAT). *Eur Respir J* 2017;**50**(3):1700582. Doi: 10.1183/13993003.00582-2017.
- 18 Bornstain C., Azoulay E., De Lassence A., Cohen Y., Costa M. A., Mourvillier B., et al. Sedation, Sucralfate, and Antibiotic Use Are Potential Means for Protection against Early-Onset Ventilator-Associated Pneumonia. *Clin Infect Dis* 2004;**38**(10):1401–8. Doi: 10.1086/386321.
- 19 Singer Mervyn, Deutschman Clifford S., Seymour Christopher Warren, Shankar-Hari Manu, Annane Djillali, Bauer Michael, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016;**315**(8):801. Doi: 10.1001/jama.2016.0287.
- 20 Lau B., Cole S. R., Gange S. J. Competing Risk Regression Models for Epidemiologic Data. *Am J Epidemiol* 2009;**170**(2):244–56. Doi: 10.1093/aje/kwp107.
- 21 Wolkewitz M., Cooper B. S., Bonten M. J. M., Barnett A. G., Schumacher M. Interpreting and comparing risks in the presence of competing events. *BMJ* 2014;**349**(aug21 5):g5060–g5060. Doi: 10.1136/bmj.g5060.
- 22 Vesin Aurélien, Azoulay Elie, Ruckly Stéphane, Vignoud Lucile, Rusinovà Kateřina, Benoit Dominique, et al. Reporting and handling missing values in clinical studies in intensive care units. *Intensive Care Med* 2013;**39**(8):1396–404. Doi: 10.1007/s00134-013-2949-1.
- 23 Lahiri Shouri, Mayer Stephan A., Fink Matthew E., Lord Aaron S., Rosengart Axel, Mangat Halinder S., et al. Mechanical Ventilation for Acute Stroke: A Multi-state Population-

Based Study. *Neurocrit Care* 2015;**23**(1):28–32. Doi: 10.1007/s12028-014-0082-9.

24 Kasuya Yusuke, Hargett James L., Lenhardt Rainer, Heine Michael F., Doufas Anthony G., Rimmel Kerri S., et al. Ventilator-associated pneumonia in critically ill stroke patients: Frequency, risk factors, and outcomes. *J Crit Care* 2011;**26**(3):273–9. Doi: 10.1016/j.jcrc.2010.09.006.

25 Kostadima E., Kaditis A.G., Alexopoulos E.I., Zakynthinos E., Sfyras D. Early gastrostomy reduces the rate of ventilator-associated pneumonia in stroke or head injury patients. *Eur Respir J* 2005;**26**(1):106–11. Doi: 10.1183/09031936.05.00096104.

26 Bronchard Régis, Albaladejo Pierre, Brezac Gilles, Geffroy Arnaud, Seince Pierre-François, Morris William, et al. Early onset pneumonia: risk factors and consequences in head trauma patients. *Anesthesiology* 2004;**100**(2):234–9.

27 Lepelletier Didier, Roquilly Antoine, Demeure dit latte Dominique, Mahe Pierre Joachim, Loutrel Olivier, Champin Philippe, et al. Retrospective analysis of the risk factors and pathogens associated with early-onset ventilator-associated pneumonia in surgical-ICU head-trauma patients. *J Neurosurg Anesthesiol* 2010;**22**(1):32–7. Doi: 10.1097/ANA.0b013e3181bdf52f.

28 Bekaert Maarten, Timsit Jean-Francois, Vansteelandt Stijn, Depuydt Pieter, Vésin Aurélien, Garrouste-Orgeas Maité, et al. Attributable Mortality of Ventilator-Associated Pneumonia: A Reappraisal Using Causal Analysis. *Am J Respir Crit Care Med* 2011;**184**(10):1133–9. Doi: 10.1164/rccm.201105-0867OC.

29 Dananché Cédric, Vanhems Philippe, Machut Anaïs, Aupée Martine, Bervas Caroline, L'Hériveau François, et al. Trends of Incidence and Risk Factors of Ventilator-Associated Pneumonia in Elderly Patients Admitted to French ICUs Between 2007 and 2014. *Crit Care Med* 2018;**46**(6):869–77. Doi: 10.1097/CCM.0000000000003019.

30 Yeh Shin-Joe, Huang Kuang-Yu, Wang Tyng-Guey, Chen Yee-Chun, Chen Chung-Hwa, Tang Sung-Chun, et al. Dysphagia screening decreases pneumonia in acute stroke patients admitted to the stroke intensive care unit. *J Neurol Sci* 2011;**306**(1–2):38–41. Doi: 10.1016/j.jns.2011.04.001.

- 31 Josephson S. Andrew, Moheet Asma M., Gropper Michael A., Nichols Amy D., Smith Wade S. Ventilator-Associated Pneumonia in a Neurologic Intensive Care Unit Does Not Lead to Increased Mortality. *Neurocrit Care* 2010;**12**(2):155–8. Doi: 10.1007/s12028-009-9285-x.
- 32 Meyfroidt Geert, Bollaert Pierre-Edouard, Marik Paul E. Acute ischemic stroke in the ICU: to admit or not to admit? *Intensive Care Med* 2014;**40**(5):749–51. Doi: 10.1007/s00134-014-3289-5.
- 33 Moine Pierre, Timsit Jean-François, De Lassence Arnaud, Troché Gilles, Fosse Jean-Philippe, Alberti Corrine, et al. Mortality associated with late-onset pneumonia in the intensive care unit: results of a multi-center cohort study. *Intensive Care Med* 2002;**28**(2):154–63. Doi: 10.1007/s00134-001-1172-7.
- 34 Ibn Saied Wafa, Souweine Bertrand, Garrouste-Orgeas Maité, Ruckly Stéphane, Darmon Michael, Bailly Sébastien, et al. Respective impact of implementation of prevention strategies, colonization with multiresistant bacteria and antimicrobial use on the risk of early- and late-onset VAP: An analysis of the OUTCOMEREA network. *PLOS ONE* 2017;**12**(11):e0187791. Doi: 10.1371/journal.pone.0187791.
- 35 Sonnevile Romain, Gimenez Laetitia, Labreuche Julien, Smonig Roland, Magalhaes Eric, Bouadma Lila, et al. What is the prognosis of acute stroke patients requiring ICU admission? *Intensive Care Med* 2017;**43**(2):271–2. Doi: 10.1007/s00134-016-4553-7.


## FIGURE LEGENDS

### **Figure 1.** Microbiological documentation according to date of pneumonia onset

ICU admission pneumonia: pneumonia occurring within 1 day of ICU admission. Early-onset pneumonia: pneumonia occurring between day 2 and 7 of ICU admission. Late-onset pneumonia: pneumonia occurring from day 8 of ICU admission.

## FIGURES

Figure 1.



## TABLES

**Table 1.** Baseline characteristics

Variable N (%) or median [Q1 – Q3]	All N=195	No pneumonia N=117	Pneumonia N=78	p
Age, years	69.4 [61.2 – 76.5]	71.1 [62.1 – 78.3]	68 [60.9 – 75.3]	0.08
Male sex	132 (67.7)	71 (60.7)	61 (78.2)	0.01
<b>History</b>				
Atrial fibrillation/flutter	29 (14.9)	17 (14.5)	12 (15.4)	0.87
Previous ischemic stroke	47 (24.1)	28 (23.9)	19 (24.4)	0.95
Hypercholesterolemia	46 (23.6)	28 (23.9)	18 (23.1)	0.89
Hypertension	123 (63.1)	81 (69.2)	42 (53.8)	0.03
Coronary artery disease	32 (16.4)	19 (16.2)	13 (16.7)	0.94
Peripheral vascular disease	25 (12.8)	19 (16.2)	6 (7.7)	0.08
Obstructive pulmonary disease	16 (8.2)	10 (8.5)	6 (7.7)	0.83
Diabetes mellitus	48 (24.6)	30 (25.6)	18 (23.1)	0.68
BMI $\geq 30$ kg/m <sup>2</sup>	33 (16.9)	21 (17.9)	12 (15.4)	0.64
Pre-admission mRS score $\leq 2$	175 (89.7)	106 (90.6)	69 (88.5)	0.63
<b>Stroke characteristics</b>				
Supratentorial stroke	121 (62.4)	74 (63.2)	47 (61)	0.76
Specialized neurologic referral	190 (97.4)	113 (96.6)	77 (98.7)	0.65
Specific treatment				0.02
Intravenous thrombolysis	17 (8.7)	5 (4.3)	12 (15.4)	.
<i>In situ</i> thrombolysis	10 (5.1)	6 (5.1)	4 (5.1)	.
Thrombus aspiration	8 (4.1)	7 (6)	1 (1.3)	.
Hemorrhagic transformation	18 (9.3)	14 (12.1)	4 (5.2)	0.11
Time from stroke to ICU admission, days	2 [1 – 4]	2 [1 – 3]	2 [1 – 8]	<0.01
Time from hospital to ICU admission, days	1 [1 – 4]	1 [1 – 3]	1 [1 – 7]	0.08
<b>ICU admission</b>				
Type of ICU admission				0.77
Direct (from ED or home)	90 (46.2)	55 (47)	35 (44.9)	.
Transfer from ward	105 (53.8)	62 (53)	43 (55.1)	.
Main symptom at admission				<0.01
Coma	110 (56.4)	74 (63.2)	36 (46.2)	.
Respiratory failure	40 (20.5)	13 (11.1)	27 (34.6)	.
Seizure	19 (9.7)	14 (12)	5 (6.4)	.
Other	26 (13.3)	16 (13.7)	10 (12.8)	.

Glasgow Coma Scale score	6 [3 – 10]	5 [3 – 9]	6.5 [3 – 11]	0.03
SAPS 2	56 [45 – 66]	57 [46 – 67]	54.5 [44 – 63]	0.33
SOFA score	7 [5 – 9]	7 [5 – 9]	7 [5 – 10]	0.60

---

Abbreviations: BMI, Body Mass Index; mRS, modified Rankin Scale; ICU, Intensive Care Unit; ED, Emergency Department; SAPS, Simplified Acute Physiology Score; SOFA, Sequential Organ Failure Assessment.



**Table 2.** ICU management and outcomes

Variable N (%) or median [Q1 – Q3]	All N=195	No pneumonia N=117	Pneumonia N=78	p
<b>Organ support</b>				
Duration of invasive ventilation, days	5 [3 – 12]	4 [3 – 7]	9 [5 – 20]	<0.01
Use of renal replacement therapy	17 (8.7)	6 (5.1)	11 (14.1)	0.03
Use of vasopressors	92 (47.2)	53 (45.3)	39 (50)	0.52
Duration of vasopressor therapy, days	0 [0 – 2]	0 [0 – 2]	0.5 [0 – 4]	0.11
<b>Decisions to withhold and withdraw life-sustaining treatments</b>				
Any decision	65 (33.3)	37 (31.6)	28 (35.9)	0.54
Time to first decision, days	5 [2 – 8]	4 [2 – 7]	5 [3 – 9]	0.33
Withholding	39 (20)	21 (17.9)	18 (23.1)	0.38
Time to withholding decision, days	5 [2 – 8]	4 [2 – 7]	6 [2 – 18]	0.32
Withdrawal	34 (17.4)	21 (17.9)	13 (16.7)	0.82
Time to withdrawal decision, days	5.5 [4 – 9]	5 [3 – 9]	6 [4 – 9]	0.38
First decision before pneumonia onset (n=78)	4 (5.1)	–	–	.
<b>Outcomes</b>				
ICU length of stay, days	7 [4 – 15]	5 [3 – 9]	12 [7 – 23]	<0.01
Hospital length of stay, days	15 [6 – 36]	12 [4.5 – 28]	28 [10 – 62]	<0.01
mRS at ICU discharge				<0.01
≤ 3	22 (11.3)	17 (34.7)	5 (11.6)	.
≥ 4	70 (35.9)	32 (65.3)	38 (88.4)	.
30-day mortality (N=193)	108 (55.9)	72 (62.1)	36 (46.8)	0.03
ICU mortality	103 (52.8)	68 (58.1)	35 (44.9)	0.07
Cause of ICU death				0.11
Brain death / cerebral herniation	44 (22.6)	34 (50)	10 (28.6)	.
Systemic causes	13 (6.7)	7 (10.3)	6 (17.1)	.
Following limitation or withdrawal of care	46 (23.6)	27 (39.7)	19 (54.3)	.
Hospital mortality	123 (63.1)	77 (65.8)	46 (59)	0.33
1-year mortality (N=182)	128 (70.3)	79 (70.5)	49 (70)	0.94

Abbreviations: ICU, Intensive Care Unit; mRS, modified Rankin Scale

**Table 3.** Characteristics of post-stroke pneumonia

Variable	All
N (%) or median [Q1 – Q3]	N=195
At least one pneumonia during ICU stay	78 (40)
Number of pneumonia cases per ICU stay	
1	66 (84.6)
2	10 (12.8)
3 or more	2 (2.6)
Type of pneumonia, according to modality of acquisition (n=78)	
Community acquired	22 (28.2)
Hospital-acquired	28 (35.9)
Ventilator-acquired	28 (35.9)
Type of pneumonia, according to timing <sup>a</sup> (n=78)	
At ICU admission	46 (59)
Early-onset	20 (25.6)
Late-onset	12 (15.4)
Pneumonia severity	
Infection	16 (20.5)
Sepsis	28 (35.9)
Septic shock	34 (43.6)

Abbreviations: ICU, Intensive Care Unit

<sup>a</sup> Early-onset pneumonia: pneumonia occurring between day 2 and 7 of ICU admission. Late-onset pneumonia: pneumonia occurring from day 8 of ICU stay.

**Table 4.** Multivariable competing risk analysis for ICU mortality and ICU discharge at day-30 <sup>a</sup>

Variable	ICU mortality			ICU discharge		
	CSHR	95% CI	p	CSHR	95% CI	p
Pneumonia	0.96	[0.59 – 1.57]	0.88	0.62	[0.39 – 0.99]	0.047
Decision to withhold and withdraw life-sustaining treatments	4.43	[2.80 – 7.02]	<0.01	0.65	[0.28 – 1.53]	0.32
Participating ICUs <sup>b</sup>			0.24			0.04
ICU 1	Ref	.	.	Ref	.	.
ICU 2	1.50	[0.93 – 2.42]	.	1.87	[1.12 – 3.11]	.
Other ICUs	1.24	[0.72 – 2.12]	.	1.05	[0.57 – 1.93]	.
Study period			0.04			0.13
1997-2006	Ref	.	.	Ref	.	.
2007-2016	1.79	[1.02 – 3.15]	.	1.49	[0.89 – 2.51]	.
Chronic obstructive pulmonary disease	1.56	[0.85 – 2.89]	0.15	0.43	[0.13 – 1.39]	0.16
GCS at admission			0.01			0.11
8-15	Ref	.	.	Ref	.	.
3-7	1.90	[1.16 – 3.10]	.	0.69	[0.43 – 1.09]	.
Renal SOFA score at admission	1.03	[0.86 – 1.23]	0.75	0.87	[0.71 – 1.06]	0.12
Respiratory SOFA score at admission	1.02	[0.86 – 1.21]	0.78	0.99	[0.81 – 1.22]	0.96
Hyperglycemia <sup>c</sup> within 24h of ICU admission	0.98	[0.60 – 1.58]	0.92	0.27	[0.14 – 0.51]	<0.01

Abbreviations: CSHR, Cause-Specific Hazard Ratio; CI, Confidence Interval; GCS, Glasgow Coma Scale; SOFA, Sequential Organ Failure Assessment.

Models were adjusted with non-collinear and clinically pertinent factors associated ( $p < 0.05$ ) with the outcome of interest in univariate analysis and no selection procedure

<sup>a</sup> Multivariable competing risk analysis provides the cause-specific hazard ratio, which estimates the direct effect of an event on both ICU discharge and ICU mortality

<sup>b</sup> ICU 1 and 2 represent the 2 ICUs (out of 10) with the highest number of patients included, representing 77% of the cohort

<sup>c</sup> Hyperglycemia was defined as glycemia  $> 11\text{mmol/L}$