

Metabolic endotoxemia: a potential underlying mechanism of the relationship between dietary fat intake and risk for cognitive impairments in humans?

P. Andre, F. Laugerette, C. Feart

To cite this version:

P. Andre, F. Laugerette, C. Feart. Metabolic endotoxemia: a potential underlying mechanism of the relationship between dietary fat intake and risk for cognitive impairments in humans?. Nutrients, 2019, 11 (8), pp.1887. 10.3390/nu11081887. hal-02466234

HAL Id: hal-02466234 <https://hal.science/hal-02466234>

Submitted on 10 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Distributed under a Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

Communication

Metabolic Endotoxemia: A Potential Underlying Mechanism of the Relationship between Dietary Fat Intake and Risk for Cognitive Impairments in Humans?

Perrine André 1,*, Fabienne Laugerette ² and Catherine Féart ¹

- ¹ Team Lifelong Exposure Health and Aging, Inserm, Bordeaux Population Health Research Center, Univ. Bordeaux U1219, F-33000 Bordeaux, France
- ² Univ-Lyon, CarMeN Laboratory, INRA U1397, Inserm U1060, Université Claude Bernard Lyon 1, INSA Lyon, Charles Mérieux Medical School, FR-69600 Oullins, France
- ***** Correspondence: perrine.andre@u-bordeaux.fr; Tel.: +33-5-57-57-12-99

Received: 31 May 2019; Accepted: 9 August 2019; Published: 13 August 2019

Abstract: (1) Background: Nutrition is a major lifestyle factor that can prevent the risk of cognitive impairment and dementia. Diet-induced metabolic endotoxemia has been proposed as a major root cause of inflammation and these pathways emerge as detrimental factors of healthy ageing. The aim of this paper was to update research focusing on the relationship between a fat-rich diet and endotoxemia, and to discuss the potential role of endotoxemia in cognitive performances. (2) Methods: We conducted a non-systematic literature review based on the PubMed database related to fat-rich meals, metabolic endotoxemia and cognitive disorders including dementia in humans. A total of 40 articles out of 942 in the first screening met the inclusion criteria. (3) Results: Evidence suggested that a fat-rich diet, depending on its quality, quantity and concomitant healthy food components, could influence metabolic endotoxemia. Since only heterogeneous cross-sectional studies are available, it remains unclear to what extent endotoxemia could be associated or not with cognitive disorders and dementia. (4) Conclusions: A fat-rich diet has the capability to provide significant increases in circulating endotoxins, which highlights nutritional strategies as a promising area for future research on inflammatory-associated diseases. The role of endotoxemia in cognitive disorders and dementia remains unclear and deserves further investigation.

Keywords: nutrition; dietary fat; high-fat; endotoxemia; lipopolysaccharide; Alzheimer's disease; dementia; humans

1. Introduction

As a consequence of the ageing population, the prevalence of dementia, characterized by a progressive deterioration of cognitive performances in multiple domains (i.e., memory, reasoning, judgement and ability to perform daily activities) that evolves into a pathological diagnosis, is increasing [1]. The aetiology of dementia is multi-factorial and consists of a dynamic interaction between genetic susceptibility, non-modifiable factors (i.e., age and sex), pathological processes and environmental factors, some of them being potentially preventable [2,3].

In this context, nutrition—a major lifelong environmental factor—is of growing interest and offers an interesting strategy for the prevention of cognitive decline and subsequent dementia [4]. Numerous studies have suggested a benefit of higher adherence to healthy dietary patterns, with a beneficial balance in favour of unsaturated fatty acids to the detriment of saturated fatty acids such as the plant-based Mediterranean diet, on age-related cognitive impairments, whether they are due to degenerative or vascular origin [5–8].

Conversely, the Western diet, characterized among other factors by a high intake of saturated and trans fatty acids, has been associated with an increased risk of developing dementia [5,7]. Moreover, adopting this fat-rich dietary pattern can lead to metabolic disorders such as obesity or Type 2 Diabetes Mellitus (T2DM), these latter being associated with a higher risk of developing dementia in later life [9,10].

Nutritional strategies represent a promising area to prevent neurocognitive impairments and subsequent dementia. While the precise mechanisms underlying the relationships between these dietary habits on age-related cognitive disorders are not yet completely understood (for instance, inflammatory pathways, vascular factors, oxidative stress or amyloidogenesis [11–14]), we suggest that the activation of the innate immune system by endotoxins—which are still poorly described—could be considered as one potential cellular mechanism involved in these relationships.

Endotoxins (Lipopolysaccharides, LPS), a major component of the outer membrane of Gram-negative bacteria, are now well as contributors to the inflammation [15]. Although LPS may originate from skin and mucous membranes or local sites of bacterial infection for instance, the gut microbiota is considered the main natural reservoir of pro-inflammatory endotoxins in the body [16]. Endotoxins are released when bacteria die, and then dissociated endotoxins are able to cross the gastro-intestinal barrier to end up in the bloodstream. The presence of LPS in the bloodstream is defined as endotoxemia. Briefly, the endotoxic metabolic pathway includes the binding of circulating LPS to LPS-Binding Protein (LBP) and its transfer to the CD14 receptor, which is present both in a membrane-anchored form (mCD14) and in a soluble circulating form (sCD14) [17]. The complex LPS–LBP–CD14 initiates the secretion of pro-inflammatory cytokines, such as Interleukin-6 (IL-6) or Tumor Necrosis Factor α (TNF α), through a TLR4-dependent mechanism [18].

Although LPS is detectable at low concentrations in the circulation of healthy individuals, there is evidence that LPS levels transiently increase following ingestion of fat-rich meals [19]. Such endotoxemia is defined as "metabolic endotoxemia", in contrast to other sources of endotoxemia such as exogenous bacterial infection or sepsis. For instance, in mice fed with a four-week high-fat diet, plasma levels of LPS were similar to those observed following a four-week subcutaneous infusion of 300 µg/kg/day of LPS [20]. Metabolic endotoxemia has been proposed as a major cause of inflammation, including chronic low-grade inflammation. Indeed, animal and experimental studies have demonstrated that postprandial state may result in an inflammatory response closely associated with the increase in the circulating levels of LPS [21,22]. Interestingly, not all dietary interventions lead to increased metabolic endotoxemia and the associated postprandial inflammatory status. Data from animal studies have suggested that dietary fats act as the main macronutrients responsible for postprandial endotoxemia, and that both the quantity and the quality of the dietary fat differentially influence metabolic endotoxemia [23–25]. Moreover, healthy diets rich in unsaturated fatty acids have been associated with lower postprandial circulating levels of LPS, closely associated with lower pro-inflammatory markers. Conversely, the consumption of high-energy or high saturated-fat diets has been associated with increased postprandial levels of LPS and increased circulating levels of pro-inflammatory markers [23–25].

Currently, the available literature points out that elevated circulating levels of LPS are detrimental for healthy ageing. More specifically, elevated levels of LPS have been associated with a large range of diseases such as obesity, T2DM, coronary artery disease and depression [19]. In the context of dementia, studies have summarized evidence that neuroinflammation is considered not only an epiphenomenon of Alzheimer's disease (AD) but also a triggering factor that contributes to exacerbated AD pathology [26,27]. In addition, systemic exposure to an overload of endotoxins is widely used in animal studies to induce neuroinflammation and neurodegeneration and is associated with increased disruption of the amyloid-β precursor, hyperphosphorylated tau and neuronal cell death, which are hallmarks of AD [15,28–31]. Based on the biological plausibility taken from animal studies, we

hypothesized that lifelong exposure to endotoxins, worsened by a chronic consumption of unhealthy diets rich in saturated fat, might exacerbate the detrimental outcomes of endotoxins in the human brain. The aim of this paper is therefore to discuss the potential role of endotoxins, as an underappreciated mechanism that could be involved, at least in part, in the relationship between dietary fat intake and the pathogenesis of dementia in humans.

This article was thus conducted to emphasize research that focused on two major approaches: (i) to determine the impact of fat-rich meal intake on postprandial and fasting endotoxemia and then (ii) to clarify the relationship between LPS and cognitive performances, with a specific emphasis on dementia or AD wherever possible.

2. Methods

We did not follow the strict methodology described for conducting a systematic literature review; since not all selected studies were reviewed by the three co-authors, there was no standardized quality assessment of selected studies and we restricted our research to a single database (i.e., PubMed). The literature search was conducted and closed in February 2019 to encompass the two main aspects described above.

We used the following nutritional search terms: "high-fat", "fat intake", "dietary fat", "(un)saturated fatty acids", "Western diet", "(un)healthy diet", "Mediterranean diet", "prudent diet" combined with at least one of the search terms relative to LPS, which included "lipopolysaccharide", "LPS", "endotoxin", "endotoxemia". These search terms were applied for the selection of articles related to the approach focusing on the relation between dietary fat intake and metabolic endotoxemia. Regarding the approach focusing on the relation between endotoxemia and cognitive performances/dementia, identical search terms have been applied for LPS, combined with at least one of the following cognition-related terms: "dementia", "Alzheimer's disease", "cognition", "memory", "neurocognitive disorders". A literature search was also applied with the combination of all three domain terms (i.e., nutritional AND LPS AND cognitive terms) to select any article that would concern the whole approach. Other studies were obtained manually from the reference lists of review or original articles.

Through this research strategy, we identified a total of 942 original articles conducted on human samples (299 related to the relation between dietary fat intake and metabolic endotoxemia and 643 related to the relation between endotoxemia and cognitive performances/dementia).

Article selection was based on a two-step procedure with a first step selection based on the title and a second step selection based on the abstract reading. Articles were selected applying the following inclusion criteria: articles published in English, in scientific journals, original research, conducted on human samples. Articles including animal studies, in vitro and ex vivo studies, studies on vaccination with LPS or other pre-treatment with LPS, population aged less than 18 y, systematic review and meta-analyses were excluded. No restriction was performed on study design or date of publication.

Due to the possible sickness behavior induced by endotoxin injection (e.g., fever, malaise, headache and so forth) which may influence the subjective aspect of well-being and cognitive feeling, we chose not to discuss studies about self-reported feelings of the cognitive aspect or mood disorders. Finally, a total of 40 articles were included to establish this "state-of-the-art" article.

Due to the heterogeneity observed in nutritional intervention studies and regarding the cognitive domains assessed, no meta-analyses were performed in this article.

3. Results

3.1. Dietary Fat and Endotoxemia

3.1.1. Role of Dietary Fat Interventions on Postprandial Metabolic Endotoxemia

The role of dietary fats is of growing interest among macronutrients in many studies on the link between nutrition and metabolic endotoxemia. As described below (Table 1), evidence accumulated over the past few decades has demonstrated a transient increase in circulating LPS following the consumption of dietary fats.

Dietary Fats

Deopurkar et al. [32] attempted to determine which macronutrient was responsible for postprandial metabolic endotoxemia by comparing isocaloric meals (300 kcal) from a glucose-drink, orange juice or dairy cream among 42 healthy adults. In this study, only dairy cream consumption increased plasma LPS levels, suggesting that dietary fats act as the main macronutrient responsible for postprandial endotoxemia. Among healthy men, postprandial LPS was increased by 35% to 50% following a single acute high-fat bolus [21,33,34].

Amount of Fats

In a comparative study using different caloric doses of a high-fat diet (500, 1000 or 1500 kcal corresponding to 34 to 102 g of fat), Schwander et al. [35] observed a positive association between fat caloric doses and increased LPS levels. Along this line, another study found higher postprandial levels of LPS following ingestion of 40 g of fat compared to 10 g among obese participants [36].

Quality of Fats

More recently, a study suggested that not only the quantity but also the quality of dietary fats may be of importance. In a randomized crossover study among 20 healthy young adults, Lyte et al. [37] demonstrated that serum levels of LPS were decreased by 50% following the consumption n-3 polyunsaturated fatty acids (PUFA) and increased by 60% following the consumption of saturated fatty acids (SFA). A sustained response (until 3 h after meal) was also observed in high-fat diet compared to low-fat-diet (peak at 1 h after meal).

Structure of Fats

Interestingly, beyond the quantity and quality of fat intake, Vors et al. [38] suggested that the structure of lipids could also be one of the determinants of metabolic endotoxemia and compared the effect of 40 g of milk fat, which only differed by fat structure, either spread or emulsified. Compared with spread fat consumption, emulsified fat consumption induced more pronounced postprandial levels of LPS in obese subjects.

Supplementation with Healthy Food Components

Acute peanut consumption within a high-fat meal was associated with lower LPS concentrations 3 h postprandially compared with a control biscuit which provided similar quantities of macronutrients, fibre and energy [39]. This result suggested that the addition of healthy food components to a high-fat diet could contribute to limiting postprandial metabolic endotoxemia. In this line, Ghanim et al. demonstrated that the increased levels of LPS following the intake of a high-fat high-calorie meal [40] could be prevented by the addition of 30 g of fibre, among healthy adults [41]. In parallel, no increase in postprandial endotoxemia was observed after the consumption of a low-fat meal rich in fibre and fruit [40]. Using the same high-fat high-calorie meal as Ghanim et al. [40], Milan et al. [42] explored the postprandial effect of dietary fat intake among healthy younger (20–25 y) and older adults (60–75 y) and failed to reproduce the postprandial effect observed by Ghanim et al., although a trend for higher LPS levels was observed with the high-fat high caloric diet. Moreover, no difference in LPS levels was observed between younger and older adults following meal intake.

Overall, these studies suggested that the matrix effect of foods (i.e., the quantity, the quality, and structure of the ingested fats and the overall food components of the diet) could be a key component of the diet intake-related metabolic endotoxemia response, although no report has specifically developed this hypothesis.

Metabolic Endotoxemia among Different Disease Phenotypes

The acute effects of dietary fats intake on metabolic endotoxemia among different disease phenotypes including obesity, impaired glucose tolerance (IGT) and T2DM have also been explored. Interestingly, the overall results suggested that metabolic endotoxemia was significantly associated with metabolic health as described below.

In 2012, Harte et al. [43] found that metabolic endotoxemia was exacerbated after dietary fat intake among individuals with IGT and T2DM compared with non-obese participants. Following the ingestion of whipping cream, postprandial endotoxin levels were increased by 20% in obese or IGT subjects and by 125% in T2DM individuals compared with non-obese subjects. Among 40 morbidly obese adults (i.e., BMI > 40 kg/m²), the ingestion of a fat overload increased levels of LPS in participants with the highest postprandial hypertriglyceridemia (>80 mg/dL) [44]. More recently, Al-Disi et al. [45] demonstrated that a high-fat diet (75 g of fat per $m²$ body surface) exacerbated the postprandial endotoxemia in normal-weight and T2DM participants, but with a different impact on cardio-metabolic health: non-diabetic subjects appeared to have better metabolic resistance to the insult of a high-fat diet as seen by a postprandial increase in triglycerides and insulin and a decrease in HDL and LDL cholesterol [45].

Overall, evidence suggests that a single high-fat meal, especially one devoid of healthy food components, has an undoubted capacity to enhance transient postprandial endotoxemia in humans, this latter finding appears to be more pronounced in metabolically impaired individuals. To date, only a few studies have investigated the long-term effect of diet on postprandial and fasting levels of LPS, as described below.

Table 1. Selective evidence on the association between a single high-fat meal and metabolic endotoxemia.

Table 1. *Cont.*

ccharide; LAL Limulus Amebocyte Lysate; SFA Saturated fatty acids; MUFA n meostasis model assessment of insulin resistance; TG Triglycerides; IGT impaired glucose tolerance. ds; MUFA monounsaturated fatty acids; PUFA polyunsaturated fatty acids; T2DM Type ly acius, 12DM Type 2 aride; LAL Limulus Amebocyte Lysate; SFA Saturated fatty acids; MUFA monounsaturated fatty acids; PUFA polyunsaturated fatty acids and the server of the metals and the care and the control of insulin resistance, TG Triglycerides; IGT imparentials and 18 years of the care age 42 years of the control stasis model assessment of insulin resistance; TG Triglycerides; IGT impaired glucose tolerance. $\frac{1}{2}$ compared to baseline to baseline the baseline to baseline the set of $\sum_{i=1}^n$ ride: LAL Limulus Amebocyte Lysate: SFA Saturated fatty acids: MUFA monounsaturated fatty acids: PUFA polyunsaturated fatty acids: T2DM Type 2 aride; LAL Limulus Amebocyte Lysate; SFA Saturated fatty acids; MUFA mon JFA : 1 fatty acids; PUFA polyunsaturated fatty acids; T2DM Type 2 charide; LAL Limulus Amebocyte Lysate; SFA Saturated fatty acids; MUFA monounsaturated fatty acids; PUFA polyunsaturated fatty acids; T2DM Type 2 meostasis model assessment of insulin resistance; TG Triglycerides; IGT imp quantified by d glucose tolerance Diabetes Mellitus; HOMA-IR Homeostasis model assessment of insulin resistance; TG Triglycerides; IGT impaired glucose tolerance. Abbreviations: LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate; SFA Saturated fatty acids; MUFA monounsaturated fatty acids; PUFA polyunsaturated fatty acids; T2DM Type 2
Diabetes Mellitus: HOMA-IR Homeostasis model a

3.1.2. Associations between Long-Term Dietary Interventions and Circulating LPS

In an intervention study, the increase in energy intake (+70 g fat for eight weeks) was associated with an acute rise in endotoxin levels in the postprandial state, but not in the fasting state [46] (Table 2). More recently, three weeks of an intervention based on a low-fat high-carbohydrate diet enriched in n-3 PUFA has been shown to increase the postprandial levels of LPS, but decrease the fasting levels of LPS, compared with a Mediterranean diet enriched in MUFA or a SFA-rich diet, among healthy older subjects [47]. Among 75 metabolically impaired subjects, adherence to a high-fat high-saturated-fatty acid diet for 12 weeks led to an increase in the postprandial levels of LPS, but not the fasting levels [48].

In a crossover study among eight healthy older adults (55 y and over), Pendyala et al. [49] demonstrated that following a Western-type diet for four weeks significantly increased fasting plasma levels of LPS by 71%. In contrast, a 38% decrease in fasting plasma LPS levels was observed after four weeks of adherence to a prudent-type diet with equivalent energy intake to the Western-type diet. Regardless of the source of energy, Breusing et al. [50] observed that the 31% increase in fasting endotoxemia after one week of overfeeding (+50% of the energy requirement) was reversed by three weeks of caloric restriction (−50% of the energy requirement) among 15 healthy adults.

3.1.3. Associations between Lifestyle Dietary Patterns and Circulating LPS (Table 2)

Although of major interest, only very few studies have investigated the relationship between usual diet and circulating levels of LPS, all using a cross-sectional design analysis and exhibiting inconsistent results. In a subsample of 201 healthy men, fat and energy, but not carbohydrate or protein, intake was positively associated with fasting levels of LPS [51]. More recently, Kallio et al. [52] reported an association between energy intake and the levels of LPS observed only among lean subjects. Surprisingly, no association between fat intake and the fasting levels of LPS was observed in this study. On the other hand, two studies did not find an association between nutrient intake and fasting LPS levels among overweight and obese pregnant women [53] or type 1 diabetes patients [54]. In this latter study [54], higher consumption of fish and healthy snacks (including fruits and berries, fresh vegetables, soft drinks, yoghurt and low-fat cheese) and higher adherence to a modern diet (composed of fresh vegetables, pasta and rice, poultry, meat dishes and fried or grilled foods) were all significantly associated with lower fasting LPS levels in serum [54]. Finally, among elderly patients with nonvalvular atrial fibrillation, higher adherence to a Mediterranean-type diet was inversely correlated with fasting circulating LPS levels [55]. Interestingly, among Mediterranean diet food components, higher intake of fruits and legumes showed a major association with lower levels of LPS.

Overall, these results underlined that a transient rise in circulating levels of LPS can be induced by a large variety of high-fat diets, especially those devoid of healthy food components. Moreover, fasting or postprandial assessment of LPS could also partly explain the discrepancies observed in several studies. Considering that exposure to an overload of endotoxins could contribute to the development of adverse health outcomes such as elevated systemic inflammation, neuroinflammation, neurodegeneration and neural death in experimental studies [15,56], we hypothesized that lifelong exposure to endotoxins inherent to adherence to unhealthy diets and to ageing could worsen the detrimental outcomes of endotoxemia. The second part of this state-of-the-art paper describes the few studies focusing on the relationship between endotoxemia and cognitive disorders and dementia in humans.

Table 2. Selective evidence on the association between long-term diet and endotoxemia. ϵ n iong-term diet and endow **Table 1.** Selective evidence on the association between a single high-fat meal and metabolic endotoxemia.

weight and obese

Table 2. *Cont.* **Reference Sample Characteristics Endotoxemia**

s: AD Alzheir et al., J Nutr, 19 normal-weight (mean age 41 y) and 18 obese nebocyte Ly $\frac{1}{\sqrt{2}}$ $cide$ PHEA pol $\frac{1}{\sqrt{2}}$ AD Alzh 'S Lipopolysaccharide, ocyte Lysate; SFA Saturated fatty acids, M<mark>L</mark> surface) control to baseline of the bas atty acids, PUFA polyunsa β . AD Alzheid σ Disease, Er σ Eipoporysacchariae, EAE Emidia, E SFA Saturated fatty acids, MUF achas, i CIT poi Abbreviations: AD Alzheimer's Disease; LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate; SFA Saturated fatty acids, MUFA monounsaturated fatty acids, PUFA polyunsaturated fatty acids.

3.2. LPS, Cognitive Disorders and Dementia

3.2.1. LPS Injection and Short-Term Cognitive Function Assessment in Interventional Studies

In the field of LPS-related inflammation in association with dementia or cognitive performances, few studies have been identified following our research strategy.

First, all studies were interested in the impact of LPS injection on the inflammatory response of the host. Transient increases in pro-inflammatory cytokines (for instance IL-6 and $TNFx$, on average between 1 and 6 h post-infection) followed by the release of anti-inflammatory cytokines (for instance IL1-Ra and IL-10, on average between 3 and 8 h post-infection), were observed after LPS injection in all intervention studies. This result was consistent regardless of the LPS dose injected. Second, several lines of evidence suggested that inflammatory cytokines such as IL-6 or $TNF\alpha$ could be involved in cognitive disturbances [26,27]. However, it remains unknown whether and to what extent cognitive functions could be affected during transient immune activation induced by a single injection of LPS, as provided by the selected intervention studies described below (Table 3).

In a crossover study, Reichenberg et al. [57] reported a significant impairment in declarative memory until 10 h after the injection of LPS. With this intervention, decreased performances in declarative memory have also been observed in a subsample of subjects, as well as improvements in working memory compared to the injection of placebo [58]. These last controversial results are, however, not generalizable to the whole study sample [57], which raises questions. Finally, in both studies [57,58], there were no statistical or clinical differences regarding attention or executive functions following the injection of LPS compared to placebo.

In another study, using an injection of a lower dose of LPS (0.2 ng/kg body weight) a negative correlation was found between the increased IL-6 levels and memory and learning performance after 4.5–6 h [59]. Among all results, the injection of LPS did not alter working or executive functions or attention among young healthy volunteers. With higher doses of LPS (2 ng/kg body weight), the results are also controversial. This treatment did not induce any alteration of working memory, psychomotor speed capacity and information processing ability, fine control motor and attention performances until 10 h post-injection compared with the placebo group [60]. Surprisingly, the authors observed an increase in attention performance in the treated group compared to the placebo group.

In 2010, Grigoleit et al. [61] found that the injection of LPS did not affect the subscales of the Wechsler Memory Scale, analysing performances in verbal, visual or delayed memory, as well as attention and executive control processes. Using a double-blind crossover study, the same authors [62] observed that LPS injection did not affect accuracy in working memory performance, but improved reaction time in the high-dose (0.8 ng/kg body weight) 2 h post-injection compared to placebo; a result that was not observed with the lowest LPS dose (0.4 ng/kg body weight).

Alteration in emotional/social processing was observed following the injection of LPS at 0.8 ng/kg body weight [63], but not with a lower dose of LPS (0.4 ng/kg body weight) [64].

Altogether, these results underlined that the LPS dose, the delay and the targeted samples have different responses on the cognitive performances assessed (themselves being heterogeneous between studies), which deserves further research with emphasis on both experimental conditions and outcomes.

3.2.2. Endotoxemia and HIV-Associated Neurocognitive Disorders in Observational Studies

In the field of LPS and cognition, we identified 2 cross-sectional observational studies on non-healthy individuals, focusing on HIV-infected participants with or without HIV-associated neurocognitive disorders (HAND) [65,66] (Table 3). The median fasting levels of LPS were higher among the HAND group than among the no-HAND participants (116.1 $\rho g/mL$ vs. 98.2 $\rho g/mL$) [65]. Interestingly, circulating levels of LPS were not associated with the severity of HAND. In the other study, plasma LPS levels did not differ according to the score of a neurocognitive test battery designed to assess several domains of cognitive function (i.e., motor skills, speed of information processing, attention,

learning, memory, language fluency and executive function) [66]. More recently, Jespersen et al. [67] reported a lack of association between fasting levels of LPS and markers of axonal damage or monocyte activation in the central nervous system among HIV-infected adults without evidence of impaired cognitive function. In this specific sample, LPS was not detectable in the cerebrospinal fluid.

Although not tested in a human model of endotoxemia, repeated lifelong exposure to endotoxin may lead to a long-term alteration in all cognitive domains. Dementia diagnosis is the result of a long insidious process where cognitive disorders evolve to pathology. However, the relationship between endotoxemia and dementia has been explored in very few studies, which are described below (Table 4).

Table 3. Selective evidence on the association between lipopolysaccharides and cognitive function.

Table 3. Cont.

 $\overline{1a$ h aimear/s. D 2022222
2010 Eli LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate $\sum_{k=1}^{\infty}$ man Immunode levels in Hangler and the distribution of Abbreviations: AD Alzheimer's Disease; LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate; HIV Human Immunodeficiency Viruses; HAND HIV-associated neurocognitive
disorders: CSE Cerebresinal Fluid: CNS Central Newsus Syst disorders; CSF Cerebrospinal Fluid; CNS Central Nervous System; NFL Neurofilament light chain protein.

First, we identified two studies interested in the anatomy of the brain in post-mortem patients. Recent investigations reported higher LPS abundance in grey matter (superior temporal gyrus lobe) and white matter (frontal lobe) in brains from patients with Alzheimer's disease (AD) than in those from participants free of dementia [68]. Of greater interest, LPS colocalized with $\mathbf{A}\beta\mathbf{1}\text{-}40/42$ in amyloid plaques and with Aβ1-40/42 around blood in AD brains [68]. Another study reported an average three-fold higher abundance of in the hippocampus—an anatomical region of the AD brain that develops the earliest and most profound neuropathology—from four AD brains compared to two age-matched control brains, as well as a two-fold higher abundance in neocortical extracts from six AD brains and six age-matched control brains [69]. In some advanced AD patients (criteria not defined in the study), hippocampal brain lysate exhibited up to a 26-fold increase in LPS [69].

In 2008, Ancuta et al. [70] observed that plasma LPS levels were significantly higher among dementia-associated HIV participants than among participants without neurocognitive impairment, independent of plasma viral load and CD4 counts. Surprisingly, LPS levels in participants with minor cognitive and motor disorders, asymptomatic neurocognitive impairments or neuropsychiatric impairments did not differ from those in participants without neurocognitive impairment. Finally, the last study we identified reported that plasma LPS levels were three-fold higher in 18 AD participants (mean 61 ρ g/mL) than in 18 healthy controls (mean 21 ρ g/mL) [71].

Altogether, the scarce available literature in this field suggests that higher LPS levels are observed (i) among the brains of AD patients, in several regions of interest and (ii) among living AD participants, and that these results could be partially explained by higher immunosenescence also among HIV patients. These results allow us to speculate that not only transient immune activation induced by LPS, as described earlier, but also increased fasting levels of LPS (from different sources) could be partly involved in the pathogenesis of dementia. Longitudinal studies are still required to test this hypothesis.

Table 4. Selective evidence on the association between lipopolysaccharides and dementia. Four meals (35% fat provided en iipopolysaccharides and de

Abbreviations: AD Alzheimer's Disease; LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate; HIV Human Immunodeficiency Viruses. tions: AD Alzheimer's Disease; LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate; HIV Human Abbreviations: AD Alzheimer's Disease; LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate; HIV Human Immunodeficiency Viruses. Endocrinol $\frac{1}{2}$ and $\frac{1}{2}$ $\overline{}$ $\overline{}$ and $\overline{}$ or $\overline{}$ to $\overline{}$ the $\overline{}$ Abbreviations: AD Alzheimer's Disease; LPS Lipopolysaccharide; LAL Limulus Amebocyte Lysate; HIV Human Immunodeficiency Viruses. s_{SUSY}

4. Discussion

The hypothesis tested in this "state-of-the-art" paper was that metabolic endotoxemia could be an underlying mechanism of the relationship between nutrition (and mainly fats intake) and age-related cognitive impairments.

First, to the best of our knowledge, there was no study focusing on this whole association (i.e., nutritional habits that could modulate endotoxemia, which itself could be part of the pathological processes leading to dementia) in a single sample setting. Therefore, we conducted two approaches, based on the available literature: we synthesized studies focusing (i) on the role of nutritional habits and interventions on the modulation of metabolic endotoxemia, and (ii) on the association between endotoxemia and cognitive impairments.

To some extent, overall metabolic endotoxemia may be attributed to a balance between the number of LPS-containing Gram-negative bacteria in the gut microbiota and the subsequent translocation of the LPS across the gastro-intestinal barrier into the bloodstream; the fat content of the diet being a putative key actor of this translocation.

Our synthesis provided convincing evidence that fat-rich meals have an undoubted capacity to transiently modulate postprandial metabolic endotoxemia in humans. Indeed, recent studies have demonstrated that fat absorption and digestion is a step where dissociated LPS can be incorporated into chylomicrons (i.e., lipoproteins responsible for the transport of lipids through the gut barrier) thereby enabling their translocation into the bloodstream [21,22]. We highlighted that not only the quantity but also the quality of dietary fats may influence metabolic endotoxemia. Precisely, imbalanced diets in favour of saturated fatty acids have been associated with higher postprandial levels of LPS while a combination of these diets with other healthy nutritional components, such as fibre [41], is able to limit this increase.

On the other hand, long-term consumption of a high-energy-density diet, especially those derived from fat, has been associated with gut microbiota dysbiosis [72]. By shifting the balance in favour of, or at the expense of, LPS-containing Gram-negative bacteria, diet could also contribute to the amount of LPS in the gut and their translocation into the bloodstream. Metabolic diseases such as obesity and T2DM are often associated with gut microbiota dysbiosis [73,74], most likely worsened by lifelong consumption of an unhealthy fat-rich diet. This latter result could explain, at least in part, the results reporting increased fasting levels of LPS and an exacerbated postprandial increase of LPS following a high-fat-saturated meal in metabolically impaired individuals compared to healthy individuals.

Moreover, gut dysbiosis, and the subsequent release of LPS can be managed by several dietary factors, that are not discussed in the present paper, including, for instance, the intake of refined sugars, alcohol or nutritional supplements with pre- and probiotics for details see, [19]. In particular, high consumption of glucose or fructose, which is part of the Western diet, could induce an increase in circulating levels of LPS in mice [75]. Pronounced intestinal permeability and increased plasma levels of LPS were found in patients with chronic alcohol abuse [76]. Pre- and probiotics have demonstrated the ability to manipulate gut microbiota and to influence the circulating levels of LPS [19]. However, we did not intend to review all potential and nutritional factors that might increase levels of LPS and prefer to limit our hypothesis to lipids, which is a well-known risk factor for AD [77].

As a limitation of the selected studies, the discrepancies observed between postprandial and fasting levels of circulating LPS after dietary interventions are questionable. These observations suggest that metabolic endotoxemia has a fluctuating nature in humans, and that fasting levels of LPS may therefore not be an accurate marker of chronic exposure to endotoxins over time. Some authors have also suggested the measurement of LBP, which is considered a longer-term marker of endotoxin-related exposure than LPS [78,79].

The main consequence of chronic exposure to endotoxemia is the onset and maintenance of a low-grade inflammation state, with associated deleterious outcomes in elderly individuals [26,27]. However, it remains unclear to what extent acute exposure to LPS could induce age-related cognitive disorders. Indeed, our literature research showed no convincing evidence that exposure to an

intravenous overload of endotoxins was associated with major cognitive alterations in healthy individuals; a result that is in line with a previous review [80]. As already mentioned, the large heterogeneity of interventional studies reported in this paper (i.e., in terms of the injected doses of LPS, the neurocognitive tests used to assess short-term cognitive performances or the various delays between injection of LPS and cognitive assessment, for instance) limits us in drawing definitive conclusions regarding the association between endotoxemia and cognitive performances. Additionally, speculative, controversial results of LPS injection on some selected cognitive tests also ask for questions on possible compensatory mechanisms, which deserves further research.

Regarding the association between endotoxemia and dementia, the results seem to be more consistent, while they are based on a very small number of studies. Dementia results in a long insidious process accompanied by molecular and physiological changes, including oxidative stress, impairment in neuronal function and the death of neuronal cells, which may be caused or worsened by neuroinflammation [26,81]. We thought that accumulated lifelong exposure to endotoxin may therefore be associated with more severe stages of cognitive disorders and dementia. Due to the lack of prospective studies to support this hypothesis, we cannot exclude a possible reverse causation. For instance, individuals with advanced stages of dementia are more likely to develop bacterial infections (i.e., in part due to a lack of hygiene inherent to the deterioration of cognitive performances and disability status) and therefore to be exposed to higher endotoxemia. In addition, and inherent to ageing, increased permeability in physiological barriers (i.e., intestinal and blood-brain barriers) is observed in AD individuals [82,83], which could also promote the translocation of higher amounts of neurotoxic molecules such as LPS.

Finally, the LPS doses used in most interventional studies may also be discussed, in addition to those of natural exposure pathways which may gradually and intermittently deliver smaller amounts of endotoxin over time.

Among the limitations of this state-of-the-art paper, we acknowledge the lack of adopting a strict systematic literature review methodology. However, only partial studies have been identified to respond to our whole hypothesis which confirmed our innovative approach. As a limitation, we cannot exclude a publication bias; studies that are not statistically significant have been available more often than those with significant results. The strengths of the present analysis are therefore to update, by a holistic approach, in a single article the experimental and observational literature in the field of fat-rich nutrition, endotoxemia and cognition in humans and to identify some gaps to be completed in the near future.

5. Conclusions

Nutrition has been proposed as a promising non-medical strategy to prevent cognitive decline and subsequent dementia. Affected by the quantity and the quality of ingested fats, metabolic endotoxemia, involving a potent pro-inflammatory response of the host, could be one of the underlying mechanisms. As the postprandial state represents a stressful condition in which our current society spends most of its time, the identification of an individual-adapted dietary pattern associated with lower metabolic endotoxemia and subsequent inflammation is a promising area for future research focusing on inflammatory-associated diseases. However, there is an important need for research to understand to what extent transient but also chronic low-exposure to LPS, through repeated measurements of postprandial and fasting levels of LPS over time, could be associated with long-term cognitive changes until the diagnosis of dementia.

Author Contributions: Conceptualization, P.A. and C.F.; Methodology, P.A. and C.F.; Investigation, P.A.; Writing—original draft preparation, P.A.; Writing—review and editing P.A., C.F. and F.L.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- 1. Prince, M.; Bryce, R.; Albanese, E.; Wimo, A.; Ribeiro, W.; Ferri, C.P. The global prevalence of dementia: A systematic review and metaanalysis. *Alzheimers Dement.* **2013**, *9*, 63–75.e2. [\[CrossRef\]](http://dx.doi.org/10.1016/j.jalz.2012.11.007) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/23305823)
- 2. Baumgart, M.; Snyder, H.M.; Carrillo, M.C.; Fazio, S.; Kim, H.; Johns, H. Summary of the evidence on modifiable risk factors for cognitive decline and dementia: A population-based perspective. *Alzheimers Dement.* **2015**, *11*, 718–726. [\[CrossRef\]](http://dx.doi.org/10.1016/j.jalz.2015.05.016) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/26045020)
- 3. Samieri, C.; Perier, M.-C.; Gaye, B.; Proust-Lima, C.; Helmer, C.; Dartigues, J.-F.; Berr, C.; Tzourio, C.; Empana, J.-P. Association of Cardiovascular Health Level in Older Age with Cognitive Decline and Incident Dementia. *JAMA* **2018**, *320*, 657–664. [\[CrossRef\]](http://dx.doi.org/10.1001/jama.2018.11499) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/30140876)
- 4. Canevelli, M.; Lucchini, F.; Quarata, F.; Bruno, G.; Cesari, M. Nutrition and Dementia: Evidence for Preventive Approaches? *Nutrients* **2016**, *8*, 144. [\[CrossRef\]](http://dx.doi.org/10.3390/nu8030144) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/26959055)
- 5. Kalmijn, S.; Launer, L.J.; Ott, A.; Witteman, J.C.; Hofman, A.; Breteler, M.M. Dietary fat intake and the risk of incident dementia in the Rotterdam Study. *Ann. Neurol.* **1997**, *42*, 776–782. [\[CrossRef\]](http://dx.doi.org/10.1002/ana.410420514) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/9392577)
- 6. Lourida, I.; Soni, M.; Thompson-Coon, J.; Purandare, N.; Lang, I.A.; Ukoumunne, O.C.; Llewellyn, D.J. Mediterranean diet, cognitive function, and dementia: A systematic review. *Epidemiology* **2013**, *24*, 479–489. [\[CrossRef\]](http://dx.doi.org/10.1097/EDE.0b013e3182944410) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/23680940)
- 7. Morris, M.C.; Tangney, C.C. Dietary fat composition and dementia risk. *Neurobiol. Aging* **2014**, *35* (Suppl. 2), S59–S64. [\[CrossRef\]](http://dx.doi.org/10.1016/j.neurobiolaging.2014.03.038) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/24970568)
- 8. Melo, H.M.; Santos, L.E.; Ferreira, S.T. Diet-Derived Fatty Acids, Brain Inflammation, and Mental Health. *Front Neurosci* **2019**, *13*, 265. [\[CrossRef\]](http://dx.doi.org/10.3389/fnins.2019.00265) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/30983955)
- 9. Ninomiya, T. Diabetes mellitus and dementia. *Curr. Diab. Rep.* **2014**, *14*, 487. [\[CrossRef\]](http://dx.doi.org/10.1007/s11892-014-0487-z) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/24623199)
- 10. Dye, L.; Boyle, N.B.; Champ, C.; Lawton, C. The relationship between obesity and cognitive health and decline. *Proc. Nutr. Soc.* **2017**, *76*, 443–454. [\[CrossRef\]](http://dx.doi.org/10.1017/S0029665117002014)
- 11. Lee, H.J.; Seo, H.I.; Cha, H.Y.; Yang, Y.J.; Kwon, S.H.; Yang, S.J. Diabetes and Alzheimer's Disease: Mechanisms and Nutritional Aspects. *Clin. Nutr. Res.* **2018**, *7*, 229–240. [\[CrossRef\]](http://dx.doi.org/10.7762/cnr.2018.7.4.229) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/30406052)
- 12. Malcomson, F.C.; Mathers, J.C. Nutrition and Ageing. *Subcell. Biochem.* **2018**, *90*, 373–424. [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/30779016)
- 13. Gardener, S.L.; Rainey-Smith, S.R.; Martins, R.N. Diet and Inflammation in Alzheimer's Disease and Related Chronic Diseases: A Review. *J. Alzheimers Dis.* **2016**, *50*, 301–334. [\[CrossRef\]](http://dx.doi.org/10.3233/JAD-150765) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/26682690)
- 14. Pistollato, F.; Sumalla Cano, S.; Elio, I.; Masias Vergara, M.; Giampieri, F.; Battino, M. Role of gut microbiota and nutrients in amyloid formation and pathogenesis of Alzheimer disease. *Nutr. Rev.* **2016**, *74*, 624–634. [\[CrossRef\]](http://dx.doi.org/10.1093/nutrit/nuw023) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/27634977)
- 15. Catorce, M.N.; Gevorkian, G. LPS-induced Murine Neuroinflammation Model: Main Features and Suitability for Pre-clinical Assessment of Nutraceuticals. *Curr. Neuropharmacol.* **2016**, *14*, 155–164. [\[CrossRef\]](http://dx.doi.org/10.2174/1570159X14666151204122017) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/26639457)
- 16. Savage, D.C. Microbial ecology of the gastrointestinal tract. *Annu. Rev. Microbiol.* **1977**, *31*, 107–133. [\[CrossRef\]](http://dx.doi.org/10.1146/annurev.mi.31.100177.000543) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/334036)
- 17. Hailman, E.; Lichenstein, H.S.; Wurfel, M.M.; Miller, D.S.; Johnson, D.A.; Kelley, M.; Busse, L.A.; Zukowski, M.M.; Wright, S.D. Lipopolysaccharide (LPS)-binding protein accelerates the binding of LPS to CD14. *J. Exp. Med.* **1994**, *179*, 269–277. [\[CrossRef\]](http://dx.doi.org/10.1084/jem.179.1.269) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/7505800)
- 18. Stoll, L.L.; Denning, G.M.; Weintraub, N.L. Potential role of endotoxin as a proinflammatory mediator of atherosclerosis. *Arterioscler. Thromb. Vasc. Biol.* **2004**, *24*, 2227–2236. [\[CrossRef\]](http://dx.doi.org/10.1161/01.ATV.0000147534.69062.dc) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/15472123)
- 19. Brown, B.I. Nutritional Management of Metabolic Endotoxemia: A Clinical Review. *Altern. Ther. Health Med.* **2017**, *23*, 42–54.
- 20. Cani, P.D.; Amar, J.; Iglesias, M.A.; Poggi, M.; Knauf, C.; Bastelica, D.; Neyrinck, A.M.; Fava, F.; Tuohy, K.M.; Chabo, C.; et al. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* **2007**, *56*, 1761–1772. [\[CrossRef\]](http://dx.doi.org/10.2337/db06-1491)
- 21. Laugerette, F.; Vors, C.; Géloën, A.; Chauvin, M.-A.; Soulage, C.; Lambert-Porcheron, S.; Peretti, N.; Alligier, M.; Burcelin, R.; Laville, M.; et al. Emulsified lipids increase endotoxemia: Possible role in early postprandial low-grade inflammation. *J. Nutr. Biochem.* **2011**, *22*, 53–59. [\[CrossRef\]](http://dx.doi.org/10.1016/j.jnutbio.2009.11.011) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/20303729)
- 22. Ghoshal, S.; Witta, J.; Zhong, J.; de Villiers, W.; Eckhardt, E. Chylomicrons promote intestinal absorption of lipopolysaccharides. *J. Lipid Res.* **2009**, *50*, 90–97. [\[CrossRef\]](http://dx.doi.org/10.1194/jlr.M800156-JLR200) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/18815435)
- 23. Netto Candido, T.L.; Bressan, J.; Alfenas R de, C.G. Dysbiosis and metabolic endotoxemia induced by high-fat diet. *Nutr. Hosp.* **2018**, *35*, 1432–1440. [\[CrossRef\]](http://dx.doi.org/10.20960/nh.1792) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/30525859)
- 24. Laugerette, F.; Vors, C.; Peretti, N.; Michalski, M.-C. Complex links between dietary lipids, endogenous endotoxins and metabolic inflammation. *Biochimie* **2011**, *93*, 39–45. [\[CrossRef\]](http://dx.doi.org/10.1016/j.biochi.2010.04.016) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/20433893)
- 25. Huang, H.; Liu, T.; Rose, J.L.; Stevens, R.L.; Hoyt, D.G. Sensitivity of mice to lipopolysaccharide is increased by a high saturated fat and cholesterol diet. *J. Inflamm.* **2007**, *4*, 22. [\[CrossRef\]](http://dx.doi.org/10.1186/1476-9255-4-22) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/17997851)
- 26. Heppner, F.L.; Ransohoff, R.M.; Becher, B. Immune attack: The role of inflammation in Alzheimer disease. *Nat. Rev. Neurosci.* **2015**, *16*, 358–372. [\[CrossRef\]](http://dx.doi.org/10.1038/nrn3880) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/25991443)
- 27. Eikelenboom, P.; van Exel, E.; Hoozemans, J.J.M.; Veerhuis, R.; Rozemuller, A.J.M.; van Gool, W.A. Neuroinflammation—An early event in both the history and pathogenesis of Alzheimer's disease. *Neurodegener. Dis.* **2010**, *7*, 38–41. [\[CrossRef\]](http://dx.doi.org/10.1159/000283480) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/20160456)
- 28. Brugg, B.; Dubreuil, Y.L.; Huber, G.; Wollman, E.E.; Delhaye-Bouchaud, N.; Mariani, J. Inflammatory processes induce beta-amyloid precursor protein changes in mouse brain. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 3032–3035. [\[CrossRef\]](http://dx.doi.org/10.1073/pnas.92.7.3032) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/7708769)
- 29. Medina-Fernández, F.J.; Luque, E.; Aguilar-Luque, M.; Agüera, E.; Feijóo, M.; García-Maceira, F.I.; Escribano, B.M.; Pascual-Leone, Á.; Drucker-Colín, R.; Túnez, I. Transcranial magnetic stimulation modifies astrocytosis, cell density and lipopolysaccharide levels in experimental autoimmune encephalomyelitis. *Life Sci.* **2017**, *169*, 20–26. [\[CrossRef\]](http://dx.doi.org/10.1016/j.lfs.2016.11.011)
- 30. Miklossy, J.; Kis, A.; Radenovic, A.; Miller, L.; Forro, L.; Martins, R.; Reiss, K.; Darbinian, N.; Darekar, P.; Mihaly, L.; et al. Beta-amyloid deposition and Alzheimer's type changes induced by Borrelia spirochetes. *Neurobiol. Aging* **2006**, *27*, 228–236. [\[CrossRef\]](http://dx.doi.org/10.1016/j.neurobiolaging.2005.01.018)
- 31. Lee, J.W.; Lee, Y.K.; Yuk, D.Y.; Choi, D.Y.; Ban, S.B.; Oh, K.W.; Hong, J.T. Neuro-inflammation induced by lipopolysaccharide causes cognitive impairment through enhancement of beta-amyloid generation. *J. Neuroinflammation* **2008**, *5*, 37. [\[CrossRef\]](http://dx.doi.org/10.1186/1742-2094-5-37) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/18759972)
- 32. Deopurkar, R.; Ghanim, H.; Friedman, J.; Abuaysheh, S.; Sia, C.L.; Mohanty, P.; Viswanathan, P.; Chaudhuri, A.; Dandona, P. Differential effects of cream, glucose, and orange juice on inflammation, endotoxin, and the expression of Toll-like receptor-4 and suppressor of cytokine signaling-3. *Diabetes Care* **2010**, *33*, 991–997. [\[CrossRef\]](http://dx.doi.org/10.2337/dc09-1630) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/20067961)
- 33. Erridge, C.; Attina, T.; Spickett, C.M.; Webb, D.J. A high-fat meal induces low-grade endotoxemia: Evidence of a novel mechanism of postprandial inflammation. *Am. J. Clin. Nutr.* **2007**, *86*, 1286–1292. [\[CrossRef\]](http://dx.doi.org/10.1093/ajcn/86.5.1286) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/17991637)
- 34. Schmid, A.; Petry, N.; Walther, B.; Bütikofer, U.; Luginbühl, W.; Gille, D.; Chollet, M.; McTernan, P.G.; Gijs, M.A.M.; Vionnet, N.; et al. Inflammatory and metabolic responses to high-fat meals with and without dairy products in men. *Br. J. Nutr.* **2015**, *113*, 1853–1861. [\[CrossRef\]](http://dx.doi.org/10.1017/S0007114515000677) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/25990454)
- 35. Schwander, F.; Kopf-Bolanz, K.A.; Buri, C.; Portmann, R.; Egger, L.; Chollet, M.; McTernan, P.G.; Piya, M.K.; Gijs, M.A.M.; Vionnet, N.; et al. A dose-response strategy reveals differences between normal-weight and obese men in their metabolic and inflammatory responses to a high-fat meal. *J. Nutr.* **2014**, *144*, 1517–1523. [\[CrossRef\]](http://dx.doi.org/10.3945/jn.114.193565) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/24812072)
- 36. Vors, C.; Pineau, G.; Drai, J.; Meugnier, E.; Pesenti, S.; Laville, M.; Laugerette, F.; Malpuech-Brugère, C.; Vidal, H.; Michalski, M.-C. Postprandial Endotoxemia Linked with Chylomicrons and Lipopolysaccharides Handling in Obese Versus Lean Men: A Lipid Dose-Effect Trial. *J. Clin. Endocrinol. Metab.* **2015**, *100*, 3427–3435. [\[CrossRef\]](http://dx.doi.org/10.1210/jc.2015-2518) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/26151336)
- 37. Lyte, J.M.; Gabler, N.K.; Hollis, J.H. Postprandial serum endotoxin in healthy humans is modulated by dietary fat in a randomized, controlled, cross-over study. *Lipids Health Dis.* **2016**, *15*, 186. [\[CrossRef\]](http://dx.doi.org/10.1186/s12944-016-0357-6)
- 38. Vors, C.; Drai, J.; Pineau, G.; Laville, M.; Vidal, H.; Laugerette, F.; Michalski, M.-C. Emulsifying dietary fat modulates postprandial endotoxemia associated with chylomicronemia in obese men: A pilot randomized crossover study. *Lipids Health Dis.* **2017**, *16*, 97. [\[CrossRef\]](http://dx.doi.org/10.1186/s12944-017-0486-6) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28545546)
- 39. Moreira, A.P.B.; Teixeira, T.F.S.; Alves, R.D.M.; Peluzio, M.C.G.; Costa, N.M.B.; Bressan, J.; Mattes, R.; Alfenas, R.C.G. Effect of a high-fat meal containing conventional or high-oleic peanuts on post-prandial lipopolysaccharide concentrations in overweight/obese men. *J. Hum. Nutr. Diet* **2016**, *29*, 95–104. [\[CrossRef\]](http://dx.doi.org/10.1111/jhn.12284)
- 40. Ghanim, H.; Abuaysheh, S.; Sia, C.L.; Korzeniewski, K.; Chaudhuri, A.; Fernandez-Real, J.M.; Dandona, P. Increase in plasma endotoxin concentrations and the expression of Toll-like receptors and suppressor of cytokine signaling-3 in mononuclear cells after a high-fat, high-carbohydrate meal: Implications for insulin resistance. *Diabetes Care* **2009**, *32*, 2281–2287. [\[CrossRef\]](http://dx.doi.org/10.2337/dc09-0979)
- 41. Ghanim, H.; Batra, M.; Abuaysheh, S.; Green, K.; Makdissi, A.; Kuhadiya, N.D.; Chaudhuri, A.; Dandona, P. Antiinflammatory and ROS Suppressive Effects of the Addition of Fiber to a High-Fat High-Calorie Meal. *J. Clin. Endocrinol. Metab.* **2017**, *102*, 858–869. [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/27906549)
- 42. Milan, A.M.; Pundir, S.; Pileggi, C.A.; Markworth, J.F.; Lewandowski, P.A.; Cameron-Smith, D. Comparisons of the Postprandial Inflammatory and Endotoxaemic Responses to Mixed Meals in Young and Older Individuals: A Randomised Trial. *Nutrients* **2017**, *9*, 354. [\[CrossRef\]](http://dx.doi.org/10.3390/nu9040354) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28368340)
- 43. Harte, A.L.; Varma, M.C.; Tripathi, G.; McGee, K.C.; Al-Daghri, N.M.; Al-Attas, O.S.; Sabico, S.; O'Hare, J.P.; Ceriello, A.; Saravanan, P.; et al. High fat intake leads to acute postprandial exposure to circulating endotoxin in type 2 diabetic subjects. *Diabetes Care* **2012**, *35*, 375–382. [\[CrossRef\]](http://dx.doi.org/10.2337/dc11-1593) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/22210577)
- 44. Clemente-Postigo, M.; Queipo-Ortuño, M.I.; Murri, M.; Boto-Ordoñez, M.; Perez-Martinez, P.; Andres-Lacueva, C.; Cardona, F.; Tinahones, F.J. Endotoxin increase after fat overload is related to postprandial hypertriglyceridemia in morbidly obese patients. *J. Lipid Res.* **2012**, *53*, 973–978. [\[CrossRef\]](http://dx.doi.org/10.1194/jlr.P020909) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/22394503)
- 45. Al-Disi, D.A.; Al-Daghri, N.M.; Khan, N.; Alfadda, A.A.; Sallam, R.M.; Alsaif, M.; Sabico, S.; Tripathi, G.; McTernan, P.G. Postprandial Effect of a High-Fat Meal on Endotoxemia in Arab Women with and without Insulin-Resistance-Related Diseases. *Nutrients* **2015**, *7*, 6375–6389. [\[CrossRef\]](http://dx.doi.org/10.3390/nu7085290) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/26247966)
- 46. Laugerette, F.; Alligier, M.; Bastard, J.-P.; Drai, J.; Chanséaume, E.; Lambert-Porcheron, S.; Laville, M.; Morio, B.; Vidal, H.; Michalski, M.-C. Overfeeding increases postprandial endotoxemia in men: Inflammatory outcome may depend on LPS transporters LBP and sCD14. *Mol. Nutr. Food Res.* **2014**, *58*, 1513–1518. [\[CrossRef\]](http://dx.doi.org/10.1002/mnfr.201400044) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/24687809)
- 47. Lopez-Moreno, J.; Garcia-Carpintero, S.; Gomez-Delgado, F.; Jimenez-Lucena, R.; Vals-Delgado, C.; Alcala-Diaz, J.F.; Roncero-Ramos, I.; Rangel-Zuñiga, O.A.; Yubero-Serrano, E.M.; Malagon, M.M.; et al. Endotoxemia is modulated by quantity and quality of dietary fat in older adults. *Exp. Gerontol.* **2018**, *109*, 119–125. [\[CrossRef\]](http://dx.doi.org/10.1016/j.exger.2017.11.006) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/29129737)
- 48. López-Moreno, J.; García-Carpintero, S.; Jimenez-Lucena, R.; Haro, C.; Rangel-Zúñiga, O.A.; Blanco-Rojo, R.; Yubero-Serrano, E.M.; Tinahones, F.J.; Delgado-Lista, J.; Pérez-Martínez, P.; et al. Effect of Dietary Lipids on Endotoxemia Influences Postprandial Inflammatory Response. *J. Agric. Food Chem.* **2017**, *65*, 7756–7763. [\[CrossRef\]](http://dx.doi.org/10.1021/acs.jafc.7b01909) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28793772)
- 49. Pendyala, S.; Walker, J.M.; Holt, P.R. A high-fat diet is associated with endotoxemia that originates from the gut. *Gastroenterology* **2012**, *142*, 1100–1101.e2. [\[CrossRef\]](http://dx.doi.org/10.1053/j.gastro.2012.01.034) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/22326433)
- 50. Breusing, N.; Lagerpusch, M.; Engstler, A.J.; Bergheim, I.; Mueller, M.J.; Bosy-Westphal, A. Influence of Energy Balance and Glycemic Index on Metabolic Endotoxemia in Healthy Men. *J. Am. Coll. Nutr.* **2017**, *36*, 72–79. [\[CrossRef\]](http://dx.doi.org/10.1080/07315724.2016.1156036)
- 51. Amar, J.; Burcelin, R.; Ruidavets, J.B.; Cani, P.D.; Fauvel, J.; Alessi, M.C.; Chamontin, B.; Ferriéres, J. Energy intake is associated with endotoxemia in apparently healthy men. *Am. J. Clin. Nutr.* **2008**, *87*, 1219–1223. [\[CrossRef\]](http://dx.doi.org/10.1093/ajcn/87.5.1219) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/18469242)
- 52. Kallio, K.A.E.; Hätönen, K.A.; Lehto, M.; Salomaa, V.; Männistö, S.; Pussinen, P.J. Endotoxemia, nutrition, and cardiometabolic disorders. *Acta Diabetol.* **2015**, *52*, 395–404. [\[CrossRef\]](http://dx.doi.org/10.1007/s00592-014-0662-3) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/25326898)
- 53. Röytiö, H.; Mokkala, K.; Vahlberg, T.; Laitinen, K. Dietary intake of fat and fibre according to reference values relates to higher gut microbiota richness in overweight pregnant women. *Br. J. Nutr.* **2017**, *118*, 343–352. [\[CrossRef\]](http://dx.doi.org/10.1017/S0007114517002100) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28901891)
- 54. Ahola, A.J.; Lassenius, M.I.; Forsblom, C.; Harjutsalo, V.; Lehto, M.; Groop, P.-H. Dietary patterns reflecting healthy food choices are associated with lower serum LPS activity. *Sci. Rep.* **2017**, *7*, 6511. [\[CrossRef\]](http://dx.doi.org/10.1038/s41598-017-06885-7) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28747701)
- 55. Pastori, D.; Carnevale, R.; Nocella, C.; Novo, M.; Santulli, M.; Cammisotto, V.; Menichelli, D.; Pignatelli, P.; Violi, F. Gut-Derived Serum Lipopolysaccharide is Associated with Enhanced Risk of Major Adverse Cardiovascular Events in Atrial Fibrillation: Effect of Adherence to Mediterranean Diet. *J. Am. Heart Assoc.* **2017**, *6*, e005784. [\[CrossRef\]](http://dx.doi.org/10.1161/JAHA.117.005784) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28584074)
- 56. Cunningham, C.; Wilcockson, D.C.; Campion, S.; Lunnon, K.; Perry, V.H. Central and systemic endotoxin challenges exacerbate the local inflammatory response and increase neuronal death during chronic neurodegeneration. *J. Neurosci.* **2005**, *25*, 9275–9284. [\[CrossRef\]](http://dx.doi.org/10.1523/JNEUROSCI.2614-05.2005) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/16207887)
- 57. Reichenberg, A.; Yirmiya, R.; Schuld, A.; Kraus, T.; Haack, M.; Morag, A.; Pollmächer, T. Cytokine-associated emotional and cognitive disturbances in humans. *Arch. Gen. Psychiatry* **2001**, *58*, 445–452. [\[CrossRef\]](http://dx.doi.org/10.1001/archpsyc.58.5.445) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/11343523)
- 58. Cohen, O.; Reichenberg, A.; Perry, C.; Ginzberg, D.; Pollmächer, T.; Soreq, H.; Yirmiya, R. Endotoxin-induced changes in human working and declarative memory associate with cleavage of plasma "readthrough" acetylcholinesterase. *J. Mol. Neurosci.* **2003**, *21*, 199–212. [\[CrossRef\]](http://dx.doi.org/10.1385/JMN:21:3:199)
- 59. Krabbe, K.S.; Reichenberg, A.; Yirmiya, R.; Smed, A.; Pedersen, B.K.; Bruunsgaard, H. Low-dose endotoxemia and human neuropsychological functions. *Brain Behav. Immun.* **2005**, *19*, 453–460. [\[CrossRef\]](http://dx.doi.org/10.1016/j.bbi.2005.04.010)
- 60. Van den Boogaard, M.; Ramakers, B.P.; van Alfen, N.; van der Werf, S.P.; Fick, W.F.; Hoedemaekers, C.W.; Verbeek, M.M.; Schoonhoven, L.; van der Hoeven, J.G.; Pickkers, P. Endotoxemia-induced inflammation and the effect on the human brain. *Crit. Care* **2010**, *14*, R81. [\[CrossRef\]](http://dx.doi.org/10.1186/cc9001)
- 61. Grigoleit, J.-S.; Oberbeck, J.R.; Lichte, P.; Kobbe, P.; Wolf, O.T.; Montag, T.; del Rey, A.; Gizewski, E.R.; Engler, H.; Schedlowski, M. Lipopolysaccharide-induced experimental immune activation does not impair memory functions in humans. *Neurobiol. Learn. Mem.* **2010**, *94*, 561–567. [\[CrossRef\]](http://dx.doi.org/10.1016/j.nlm.2010.09.011) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/20875866)
- 62. Grigoleit, J.-S.; Kullmann, J.S.; Wolf, O.T.; Hammes, F.; Wegner, A.; Jablonowski, S.; Engler, H.; Gizewski, E.; Oberbeck, R.; Schedlowski, M. Dose-dependent effects of endotoxin on neurobehavioral functions in humans. *PLoS ONE* **2011**, *6*, e28330. [\[CrossRef\]](http://dx.doi.org/10.1371/journal.pone.0028330) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/22164271)
- 63. Moieni, M.; Irwin, M.R.; Jevtic, I.; Breen, E.C.; Eisenberger, N.I. Inflammation impairs social cognitive processing: A randomized controlled trial of endotoxin. *Brain Behav. Immun.* **2015**, *48*, 132–138. [\[CrossRef\]](http://dx.doi.org/10.1016/j.bbi.2015.03.002) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/25770082)
- 64. Kullmann, J.S.; Grigoleit, J.-S.; Wolf, O.T.; Engler, H.; Oberbeck, R.; Elsenbruch, S.; Forsting, M.; Schedlowski, M.; Gizewski, E.R. Experimental human endotoxemia enhances brain activity during social cognition. *Soc. Cogn. A*ff*ect. Neurosci.* **2014**, *9*, 786–793. [\[CrossRef\]](http://dx.doi.org/10.1093/scan/nst049)
- 65. Vassallo, M.; Dunais, B.; Durant, J.; Carsenti-Dellamonica, H.; Harvey-Langton, A.; Cottalorda, J.; Ticchioni, M.; Laffon, M.; Lebrun-Frenay, C.; Dellamonica, P.; et al. Relevance of lipopolysaccharide levels in HIV-associated neurocognitive impairment: The Neuradapt study. *J. Neurovirol.* **2013**, *19*, 376–382. [\[CrossRef\]](http://dx.doi.org/10.1007/s13365-013-0181-y) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/23846287)
- 66. Lyons, J.L.; Uno, H.; Ancuta, P.; Kamat, A.; Moore, D.J.; Singer, E.J.; Morgello, S.; Gabuzda, D. Plasma sCD14 is a biomarker associated with impaired neurocognitive test performance in attention and learning domains in HIV infection. *J. Acquir. Immune Defic. Syndr.* **2011**, *57*, 371–379. [\[CrossRef\]](http://dx.doi.org/10.1097/QAI.0b013e3182237e54) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/21646912)
- 67. Jespersen, S.; Pedersen, K.K.; Anesten, B.; Zetterberg, H.; Fuchs, D.; Gisslén, M.; Hagberg, L.; Trøseid, M.; Nielsen, S.D. Soluble CD14 in cerebrospinal fluid is associated with markers of inflammation and axonal damage in untreated HIV-infected patients: A retrospective cross-sectional study. *BMC Infect. Dis.* **2016**, *16*, 176. [\[CrossRef\]](http://dx.doi.org/10.1186/s12879-016-1510-6) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/27103116)
- 68. Zhan, X.; Stamova, B.; Jin, L.-W.; DeCarli, C.; Phinney, B.; Sharp, F.R. Gram-negative bacterial molecules associate with Alzheimer disease pathology. *Neurology* **2016**, *87*, 2324–2332. [\[CrossRef\]](http://dx.doi.org/10.1212/WNL.0000000000003391) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/27784770)
- 69. Zhao, Y.; Jaber, V.; Lukiw, W.J. Secretory Products of the Human GI Tract Microbiome and Their Potential Impact on Alzheimer's Disease (AD): Detection of Lipopolysaccharide (LPS) in AD Hippocampus. *Front. Cell. Infect. Microbiol.* **2017**, *7*, 318. [\[CrossRef\]](http://dx.doi.org/10.3389/fcimb.2017.00318) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28744452)
- 70. Ancuta, P.; Kamat, A.; Kunstman, K.J.; Kim, E.-Y.; Autissier, P.; Wurcel, A.; Zaman, T.; Stone, D.; Mefford, M.; Morgello, S.; et al. Microbial translocation is associated with increased monocyte activation and dementia in AIDS patients. *PLoS ONE* **2008**, *3*, e2516. [\[CrossRef\]](http://dx.doi.org/10.1371/journal.pone.0002516)
- 71. Zhang, R.; Miller, R.G.; Gascon, R.; Champion, S.; Katz, J.; Lancero, M.; Narvaez, A.; Honrada, R.; Ruvalcaba, D.; McGrath, M.S. Circulating endotoxin and systemic immune activation in sporadic amyotrophic lateral sclerosis (sALS). *J. Neuroimmunol.* **2009**, *206*, 121–124. [\[CrossRef\]](http://dx.doi.org/10.1016/j.jneuroim.2008.09.017) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/19013651)
- 72. Cândido, F.G.; Valente, F.X.; Grześkowiak, Ł.M.; Moreira, A.P.B.; Rocha, D.M.U.P.; Alfenas R de, C.G. Impact of dietary fat on gut microbiota and low-grade systemic inflammation: Mechanisms and clinical implications on obesity. *Int. J. Food Sci. Nutr.* **2018**, *69*, 125–143. [\[CrossRef\]](http://dx.doi.org/10.1080/09637486.2017.1343286)
- 73. Graham, C.; Mullen, A.; Whelan, K. Obesity and the gastrointestinal microbiota: A review of associations and mechanisms. *Nutr. Rev.* **2015**, *73*, 376–385. [\[CrossRef\]](http://dx.doi.org/10.1093/nutrit/nuv004)
- 74. Arora, T.; Bäckhed, F. The gut microbiota and metabolic disease: Current understanding and future perspectives. *J. Intern. Med.* **2016**, *280*, 339–349. [\[CrossRef\]](http://dx.doi.org/10.1111/joim.12508)
- 75. Do, M.H.; Lee, E.; Oh, M.-J.; Kim, Y.; Park, H.-Y. High-Glucose or -Fructose Diet Cause Changes of the Gut Microbiota and Metabolic Disorders in Mice without Body Weight Change. *Nutrients* **2018**, *10*, 761. [\[CrossRef\]](http://dx.doi.org/10.3390/nu10060761) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/29899272)
- 76. Parlesak, A.; Schäfer, C.; Schütz, T.; Bode, J.C.; Bode, C. Increased intestinal permeability to macromolecules and endotoxemia in patients with chronic alcohol abuse in different stages of alcohol-induced liver disease. *J. Hepatol.* **2000**, *32*, 742–747. [\[CrossRef\]](http://dx.doi.org/10.1016/S0168-8278(00)80242-1)
- 77. La Rosa, F.; Clerici, M.; Ratto, D.; Occhinegro, A.; Licito, A.; Romeo, M.; Iorio, C.D.; Rossi, P. The Gut-Brain Axis in Alzheimer's Disease and Omega-3. A Critical Overview of Clinical Trials. *Nutrients* **2018**, *10*, 1267. [\[CrossRef\]](http://dx.doi.org/10.3390/nu10091267) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/30205543)
- 78. Wan, Y.; Freeswick, P.D.; Khemlani, L.S.; Kispert, P.H.; Wang, S.C.; Su, G.L.; Billiar, T.R. Role of lipopolysaccharide (LPS), interleukin-1, interleukin-6, tumor necrosis factor, and dexamethasone in regulation of LPS-binding protein expression in normal hepatocytes and hepatocytes from LPS-treated rats. *Infect. Immun.* **1995**, *63*, 2435–2442.
- 79. Yao, Z.; Mates, J.M.; Cheplowitz, A.M.; Hammer, L.P.; Maiseyeu, A.; Phillips, G.S.; Wewers, M.D.; Rajaram, M.V.S.; Robinson, J.M.; Anderson, C.L.; et al. Blood-Borne Lipopolysaccharide Is Rapidly Eliminated by Liver Sinusoidal Endothelial Cells via High-Density Lipoprotein. *J. Immunol.* **2016**, *197*, 2390–2399. [\[CrossRef\]](http://dx.doi.org/10.4049/jimmunol.1600702) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/27534554)
- 80. Bollen, J.; Trick, L.; Llewellyn, D.; Dickens, C. The effects of acute inflammation on cognitive functioning and emotional processing in humans: A systematic review of experimental studies. *J. Psychosom. Res.* **2017**, *94*, 47–55. [\[CrossRef\]](http://dx.doi.org/10.1016/j.jpsychores.2017.01.002) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/28183402)
- 81. Anderson, I.; Adinolfi, C.; Doctrow, S.; Huffman, K.; Joy, K.A.; Malfroy, B.; Soden, P.; Rupniak, H.T.; Barnes, J.C. Oxidative signalling and inflammatory pathways in Alzheimer's disease. *Biochem. Soc. Symp.* **2001**, *67*, 141–149. [\[CrossRef\]](http://dx.doi.org/10.1042/bss0670141)
- 82. Russell, R.M. Changes in gastrointestinal function attributed to aging. *Am. J. Clin. Nutr.* **1992**, *55*, 1203S–1207S. [\[CrossRef\]](http://dx.doi.org/10.1093/ajcn/55.6.1203S)
- 83. Osipova, E.D.; Komleva, Y.K.; Morgun, A.V.; Lopatina, O.L.; Panina, Y.A.; Olovyannikova, R.Y.; Vais, E.F.; Salmin, V.V.; Salmina, A.B. Designing in vitro Blood-Brain Barrier Models Reproducing Alterations in Brain Aging. *Front. Aging Neurosci.* **2018**, *10*, 234. [\[CrossRef\]](http://dx.doi.org/10.3389/fnagi.2018.00234) [\[PubMed\]](http://www.ncbi.nlm.nih.gov/pubmed/30127733)

© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://[creativecommons.org](http://creativecommons.org/licenses/by/4.0/.)/licenses/by/4.0/).