



HAL
open science

Laryngeal flow due to longitudinal sweeping motion of the vocal folds and its contribution to auto-oscillation

Henri Boutin, John Smith, Joe Wolfe

► **To cite this version:**

Henri Boutin, John Smith, Joe Wolfe. Laryngeal flow due to longitudinal sweeping motion of the vocal folds and its contribution to auto-oscillation. *Journal of the Acoustical Society of America*, 2015, 10.1121/1.4922470 . hal-02470031

HAL Id: hal-02470031

<https://hal.science/hal-02470031v1>

Submitted on 21 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Laryngeal flow due to longitudinal sweeping motion of the vocal folds and its contribution to auto-oscillation (L)

Henri Boutin,^{a)} John Smith, and Joe Wolfe

School of Physics, The University of New South Wales, Sydney, New South Wales 2052, Australia

(Received 10 November 2014; revised 23 May 2015; accepted 1 June 2015; published online 8 July 2015)

Analysis of published depth-kymography data [George, de Mul, Qiu, Rakhorst, and Schutte (2008). *Phys. Med. Biol.* **53**, 2667–2675] shows that, for the subject studied, the flow due to the longitudinal sweeping motion of the vocal folds contributes several percent of a typical acoustic flow at the larynx. This sweeping flow is a maximum when the glottis is closed. This observation suggests that assumption of zero laryngeal flow during the closed phase as a criterion when determining parameters in inverse filtering should be used with caution. Further, these data suggest that the swinging motion contributes work to overcome mechanical losses and thus to assist auto-oscillation.

© 2015 Acoustical Society of America. [<http://dx.doi.org/10.1121/1.4922470>]

[AH]

Pages: 146–149

I. INTRODUCTION

The acoustic flow entering the vocal tract at the larynx (hereafter the laryngeal flow) plays a fundamental role in the production of speech. However, it cannot usually be measured directly. The waveform of this laryngeal flow (often called the glottal flow) is often estimated using inverse filtering of the radiated sound (e.g., Alku 1992, 2011; Alku *et al.*, 2006a; Alku *et al.*, 2006b; Alku *et al.*, 2009; Airas, 2008). This approach involves separating two unknowns with complex behavior (the laryngeal flow and the transfer function of the tract) using only one measured signal (the radiated sound). This procedure cannot be unequivocal (Drugman, 2012; Chu *et al.*, 2013), and consequently assumptions about the waveform of the laryngeal flow are often made in order to adjust the anti-resonances of a set of inverse filters to remove the filtering effects of the vocal tract and the radiation from the lips. To this end, it is sometimes assumed that the laryngeal flow is very close to zero while the glottis is closed. This assumption is not necessarily true: if the vocal folds, while closed, are moving in the vertical direction, then they displace air in that direction, as observed by Granqvist *et al.* (2003) in the case of “flow phonation,” using inverse filtering. We refer to this hereafter as sweeping flow: the component of laryngeal flow due to the sweeping motion of the folds. [If one used the term glottal flow to describe only the mass injection associated with the flow through the glottis, i.e., through the (vibrating) aperture between the vocal folds, one could therefore write: Laryngeal flow equals glottal flow plus sweeping flow.] The sweeping flow is quite distinct from the flows caused by displacement of the folds in the horizontal plane (Rothenberg, 1973) and by the deformation of the folds during collision (Pelorson *et al.*, 1994). It is therefore of interest to compare the flow rates of total laryngeal flow and sweeping flow, which is one purpose of this short communication. Another purpose is to discuss how the

work done on the folds during their swinging motion can contribute to the auto-oscillation of the vocal folds.

II. SWEEPING FLOW DUE TO THE VOCAL FOLDS' MOTION

For an upright speaker, define the lateral direction (i.e., the horizontal left-right axis) as x , the forwards direction as y , and the (nearly vertical, longitudinal) direction parallel to the supraglottal tract, as z . The z motion of the vocal folds has been quantified by George *et al.* (2008), who used triangulation of laser beams in combination with nasendoscopy. (The cited study measured only one subject, and the motions of this subject's vocal folds were not symmetrical—in the following analysis the data for both vocal folds are combined.) George *et al.* plot $z(x, t)$ at a constant y position roughly midway across the glottis. Its amplitude, about 2 mm, has the same order of magnitude as the estimate given by Granqvist *et al.* (2003), between 1 and 2 mm, and the measurements of Larsson (2009), between 0.3 and 1.5 mm.

To calculate the sweeping flow from these data, $z(x, t)$ is integrated along the x -direction for both vocal folds, differentiated with respect to time and multiplied by an effective length L of the moving vocal folds in the y -direction. L is obviously less than the total length (9.1 mm) measured on Fig. 2 from George *et al.* (2008). If the shape along this length were a sine function from 0 to π , the effective length would be $9.1 \text{ mm} \times 2/\pi = 5.8 \text{ mm}$. If it varied as \sin^2 from 0 to π , the effective length would be $9.1 \text{ mm}/2 = 4.6 \text{ mm}$. The effective length is set at 5 mm for this order-of-magnitude calculation. The differentiation is performed by taking the Fourier transform, multiplying each component at angular frequency ω by $j\omega$ and then taking the inverse transform. Figure 1(top) displays this sweeping flow averaged over four complete cycles before (pale curve) and after (dark curve) the harmonics above the third have been removed. Figure 1(bottom) shows the open area of the glottis (i.e., the glottal aperture) $A(t)$ calculated from the same data set. (Two complete cycles are shown.)

Considering the filtered curve in Fig. 1(top), the peak of the sweeping flow corresponds closely with the beginning of

^{a)}Electronic mail: boutin@lam.jussieu.fr




FIG. 1. Sweeping flow and glottal aperture calculated using the data of George *et al.* (2008), during a sustained phonation by a trained professional.

the closed glottis phase and has a magnitude of 34 mL s^{-1} . Ideally, this sweeping flow would be compared with total laryngeal flow measured simultaneously. However, the laryngeal flow was not measured by George *et al.* (2008), whose subject was “a trained professional able to produce a sustained phonation over a long period of time.” A lower limit on the ratio of sweeping flow to glottal flow can be estimated by considering the extreme case of a phonation in which 5 liters of air is expended in 5 s—a very loud phonation. Approximating the laryngeal flow as a square wave, the peak value would be 2 liters s^{-1} . In this case, the sweeping flow in Fig. 1 would be only 1.7% of the total laryngeal flow. On the other hand, if the laryngeal flow corresponding to the data of Fig. 1 had a more typical amplitude of $0.3 \text{ liters s}^{-1}$ (see examples given by Södersten *et al.*, 1995), then the sweeping flow would be about 11% of the laryngeal flow. Further, its maximum occurs near the beginning of the period of glottal closure. Using a numerical model of vocal fold motion that allows both longitudinal and lateral motion, Flanagan and Ishizaka (1978) calculate the sweeping and laryngeal flows. With the parameters used, they find that the peak-to-peak sweeping flow is five percent of the total peak-to-peak laryngeal flow. Their lower value presumably reflects differences between their model and the vocal folds measured by George *et al.*

A sweeping flow with a relative magnitude of several percent or more might have implications for inverse filtering if the filtering algorithms used (whether automated or manual) assume a zero laryngeal flow during the closed glottis phase, or indeed a ripple-free flow over that phase (e.g., Bäckström *et al.*, 2005; Lehto *et al.*, 2007; Airas, 2008).

It is also interesting that other researchers often find a small flow during the “closed” phase, which is usually attributed to glottal leakage (e.g., Holmberg *et al.*, 1988; Cranen and Schroeter, 1995; Iwarsson *et al.*, 1998). This could be at least partially a consequence of the presence of a sweeping flow.

III. AUTO-OSCILLATION OF THE VOCAL FOLDS

Apart from its possible implications for inverse filtering, the presence of a non-negligible longitudinal motion has another potentially important implication for speech science.

In the following discussion, several phenomena are not mentioned. These include not only the modulation of the flow between the vocal folds, but also the deformation of the folds on collision. This omission from discussion does not, of course, imply that these are negligible.

Consider the highly simplified, one-mass, two-spring model for the vocal folds shown in Fig. 2, which is similar to that of Awrejcewicz (1990), Adachi and Yu (2005), and others. Consider the case where the frequency of vibration is well below those of the resonances of the supraglottal and subglottal tracts. (This is the case of the data analysed here, and is usually the case for both men’s and women’s speech.) At such a frequency, the supraglottal and subglottal impedances are small, so the supraglottal and subglottal pressures are approximately constant. Flow separation at such a constriction is complicated (see, e.g., Hirschberg, 1992). For the very simple model presented here, flow separation is assumed to occur at the narrowest aperture with the consequence that the pressure between the folds and everywhere downstream from them is approximately uniform at P_2 , and that the pressure upstream is uniform at $P_1 > P_2$. What is the work done on the folds by $(P_1 - P_2)$ over one complete cycle?

Assume that (as shown in Fig. 2) the phase of the motion in the z direction is ahead of that of the separation of the vocal folds in the x direction, in agreement with the observations of vocal fold vibration of Baer (1981) and the model of Adachi and Yu (2005). In consequence, the length of the folds in the x direction is greater when the folds are moving upwards [phase (a)–(b) in Fig. 2] than when they are moving downwards [phase (c)–(d) in Fig. 2]. Consequently, the positive component of longitudinal laryngeal flow due to the upwards swinging motion of the longer folds is greater than the negative component due to downwards swinging motion of the shorter folds, see Fig. 2. Subject to the conditions listed above, positive work is then performed by the pressure difference $(P_1 - P_2)$ over one complete cycle (Boutin *et al.*, 2015) solely due to the swinging motion of the vocal folds. This work done can overcome losses, and might provide a significant contribution towards auto-oscillation, particularly in cases where the phases of the load impedances may not be favorable.




FIG. 2. A simple rotating and stretching model of vocal folds composed of two compressible beams, which undergo both shear and compression-dilation due to the pressure upstream P_1 and downstream P_2 of the glottis. The dark and light gray areas represent respectively positive and negative volumes swept by the vocal folds during the intervals (a)-(b) and (c)-(d). In the lower part, the black and gray sine waves are approximations to the oscillations in the z and x directions, and illustrate their phase relation. (This sine approximation requires deformation of the vocal folds during their collision.)

An order-of-magnitude estimate of the rate of positive work done on the vocal folds by $(P_1 - P_2)$ can be calculated using the data of [George et al. \(2008\)](#). Assuming a transglottal pressure difference of 1 kPa, the average power produced by this swinging motion of the vocal folds is 0.5 mW. The radiated acoustic power of the voice is in the range 10^{-3} to 10 mW ([Titze, 1992](#)), but much of this power comes from the modulation by the vibrating vocal folds of the flow between them. The energy stored in the vibratory motion of the folds themselves and the fraction of this that is lost each cycle are not known. If, as seems likely, the rate of loss of vibratory energy is on the order of a milliwatt, the power calculated above may be a significant contribution to auto-oscillation.

IV. CONCLUSION

The measurements of [George et al. \(2008\)](#) were made on just one subject. Consequently much could be learned from similar measurements made on a larger sample, especially if simultaneous measurements could be made of flow at the lips, of the acoustical properties of the tract and/or of glottal behavior using electroglottography. One of the purposes of this paper is to encourage that. However, one immediate conclusion from this analysis is that caution should be exercised in assuming zero laryngeal flow in the closed glottal phase as a criterion for inverse filtering. Another is that the effect of pressure on the swinging motion can contribute a power term of a magnitude that might make a significant contribution to auto-oscillation, particularly in situations where the phases of the acoustic loads were unfavorable.

ACKNOWLEDGMENTS

We thank Harm K. Schutte and Frits F.M. de Mul for sending us the original versions of the plots in [George et al. \(2008\)](#) and Associate Editor A. Hirschberg for helpful comments. The support of the Australian Research Council is gratefully acknowledged.

- Adachi, S., and Yu, J. (2005). "Two-dimensional model of vocal fold vibration for sound synthesis of voice and soprano singing," *J. Acoust. Soc. Am.* **117**(5), 3213–3224.
- Airas, M. (2008). "TKK Aparat: An environment for voice inverse filtering and parameterization," *Logop. Phonatr. Voco.* **33**(1), 49–64.
- Alku, P. (1992). "Glottal wave analysis with pitch synchronous iterative adaptive inverse filtering," *Speech Commun.* **11**, 109–118.
- Alku, P. (2011). "Glottal inverse filtering analysis of human voice production—A review of estimation and parameterization methods of the glottal excitation and their applications," *Sādhanā* **36**, 623–650.
- Alku, P., Horacek, J., Airas, M., Griffond-Boitier, F., and Laukkanen, A.-M. (2006a). "Performance of glottal filtering as tested by aeroelastic modelling of phonation and FE modelling of vocal tract," *Acta Acust. Acust.* **92**, 717–724.
- Alku, P., Magi, C., Yrttiaho, S., Bäckström, T., and Story, B. (2009). "Closed phase covariance analysis based on constrained linear prediction for glottal inverse filtering," *J. Acoust. Soc. Am.* **125**(5), 3289–3305.
- Alku, P., Story, B., and Airas, M. (2006b). "Estimation of the voice source from speech pressure signals: Evaluation of an inverse filtering technique using physical modelling of voice production," *Folia Phoniatr. Logo.* **58**, 102–113.
- Awrejcewicz, J. (1990). "Bifurcation portrait of the human vocal cord oscillations," *J. Sound Vib.* **136**(1), 151–156.
- Bäckström, T., Airas, M., Lehto, L., and Alku, P. (2005). "Objective quality measures for glottal inverse filtering of speech pressure signals," in *Proceedings of IEEE International Conference on Acoustics, Speech, and Signal Processing*, Philadelphia, pp. 897–900.
- Baer, T. (1981). "Observation of vocal fold vibration: Measurement of excised larynges," in *Vocal Fold Physiology*, edited by K. N. Stevens and M. Hirano (University of Tokyo Press, Tokyo, 1981), pp. 119–133.

- Boutin, H., Fletcher, N., Smith, J., and Wolfe, J. (2015). "Relationships between pressure, flow, lip motion, and upstream and downstream impedances for the trombone," *J. Acoust. Soc. Am.* **137**, 1195–1209.
- Chu, D. T. W., Li, K., Epps, J., Smith, J., and Wolfe, J. (2013). "Experimental evaluation of inverse filtering using physical systems with known glottal flow and tract characteristics," *J. Acoust. Soc. Am.* **133**, EL358–EL362.
- Cranen, B., and Schroeter, J. (1995). "Modeling a leaky glottis," *J. Phonetics* **23**(1), 165–177.
- Drugman, T., Bozkurt, B., and Dutoit, T. (2012). "A comparative study of glottal source estimation techniques," *Comput. Speech Lang.* **26**, 20–34.
- Flanagan, J. L., and Ishizaka, K. (1978). "Computer model to characterize the air volume displaced by the vibrating vocal cords," *J. Acoust. Soc. Am.* **63**, 1559–1565.
- George, N. A., de Mul, F. F. M., Qiu, Q., Rakhorst, G., and Schutte, H. K. (2008). "Depth-kymography: high-speed calibrated 3D imaging of human vocal fold vibration dynamics," *Phys. Med. Biol.* **53**, 2667–2675.
- Granqvist, S., Hertegård, S., Larsson, H., and Sundberg, J. (2003). "Simultaneous analysis of vocal fold vibration and transglottal airflow: exploring a new experimental setup," *J. Voice* **17**(3), 319–330.
- Hirschberg, A. (1992). "Some fluid dynamic aspects of speech," *Bull. Commun. Parlée* **2**, 7–30.
- Holmberg, E. B., Hillman, R. E., and Perkell, J. S. (1988). "Glottal airflow and pressure measurements for soft, normal and loud voice by male and female speakers," *J. Acoust. Soc. Am.* **84**, 511–529.
- Iwarsson, J., Thomasson, M., and Sundberg, J. (1998). "Effects of lung volume on the glottal voice source," *J. Voice* **12**(4), 424–433.
- Larsson, H. (2009). "Methods for measurement of vocal fold vibration and viscoelasticity," Ph.D. thesis, Karolinska Institutet, Institutionen för klinisk vetenskap/Department of Clinical Sciences, Stockholm, Sweden.
- Lehto, L., Airas, M., Björkner, E., Sundberg, J., and Alku, P. (2007). "Comparison of two inverse filtering methods in parametrization of the glottal closing phase in different phonation types," *J. Voice* **21**(2), 138–150.
- Pelorson, X., Hirschbert, A., van Hassel, R. R., Wijnands, A. P. J., and Auregan, Y. (1994). "Theoretical and experimental study of quasisteady flow separation within the glottis during phonation. Application to a modified two mass model," *J. Acoust. Soc. Am.* **96**(6), 3416–3431.
- Rothenberg, M. (1973). "A new inverse-filtering technique for deriving the glottal air flow waveform during voicing," *J. Acoust. Soc. Am.* **53**, 1632–1645.
- Södersten, M., Hertegård, S., and Hammarberg, B. (1995). "Glottal closure, transglottal airflow, and voice quality in healthy middle-aged women," *J. Voice* **9**, 182–197.
- Titze, I. R. (1992). "Vocal efficiency," *J. Voice* **6**, 135–138.