



Rare earths elements cycling in reclaimed ion-adsorption mine tailings

Noemie Janot, Hermine Huot, Sophie Leguédois, Geoffroy Séré, Ye-Tao Tang, Jean-Louis Morel, Rong-Liang Qiu, Jan E. Groenenberg

► To cite this version:

Noemie Janot, Hermine Huot, Sophie Leguédois, Geoffroy Séré, Ye-Tao Tang, et al.. Rare earths elements cycling in reclaimed ion-adsorption mine tailings. Goldschmidt, Geochemical Society, Aug 2018, Boston, United States. hal-02473680

HAL Id: hal-02473680

<https://hal.science/hal-02473680>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC0 - Public Domain Dedication 4.0 International License

Rare earths elements cycling in reclaimed ion-adsorption mine tailings

NOÉMIE JANOT^{1,2}, HERMINE HUOT³, SOPHIE LEGUÉDOIS¹,
GEOFFROY SÉRÉ¹, YE-TAO TANG³, JEAN-LOUIS MOREL¹,
RONG-LIANG QIU³, JAN E. GROENENBERG^{2,4}.

¹Laboratoire Sols et Environnement, INRA-Université de
Lorraine, Vandœuvre-lès-Nancy, France

²Laboratoire Interdisciplinaire des Environnements Continentaux,
CNRS-Université de Lorraine, Vandœuvre-lès-Nancy, France

³School of Environmental Science and Engineering, Sun Yat-Sen
University, Guangzhou, China

⁴Department of Soil Quality Wageningen University, Wageningen
University and Research, Wageningen, The Netherlands

Rare Earth Elements (REEs) have become essential components of many high-tech products. Among the various known REEs deposits, the ion-adsorption ones, only found in Southern China, represented around 30% of the worldwide production in 2017. Reclamation of the resulting mine tailings is limited by their poor physical conditions, their low organic matter and nutrient levels as well as their relatively high residual REEs content. Reclamation practices of such sites can include natural attenuation or phytoremediation, using soil amendment or construction, and planting of tolerant species or economic crops. These practices induce drastic changes in the biogeochemical conditions of the soils and thus potentially modify REEs and nutrients availability, impacting the sustainability of the reclamation.

To gain understanding on the impact of reclamation practices on REEs dynamics, we study a mine tailing in Ganzhou city, Jiangxi province, that has been abandoned for 10y. Phytoremediation techniques are currently tested on the tailings material in experimental plots (2x2m) in which plants and soil development are monitored over time. More specifically, the techniques include phytostabilization using perennial grasses or fiber plants, and phytoextraction using REEs hyperaccumulators.

An objective of this study is to develop a geochemical model to be able to evaluate the various practices. A process-based multi-surface model has been parameterized for REEs in soils and sediments in the platform ORCHESTRA. Based on measured contents of REEs, clays and Al/Fe-(hydr)oxides and solution pH and DOC concentration, we predicted dissolved REEs concentrations in pore water and CaCl₂-extract. The model is evaluated by comparing modelled vs. measured REEs solid:solution partitioning. Once validated, such a model could be used to gain more detailed insights on the impact of organic exudate or amendment rates on reclamation sustainability.