



**HAL**  
open science

# Invertebrates facing environmental contamination by endocrine disruptors: Novel evidences and recent insights

Virginie Cuvillier-Hot, Alain Lenoir

► **To cite this version:**

Virginie Cuvillier-Hot, Alain Lenoir. Invertebrates facing environmental contamination by endocrine disruptors: Novel evidences and recent insights. *Molecular and Cellular Endocrinology*, 2020, 504, pp.110712. 10.1016/j.mce.2020.110712 . hal-02480067

**HAL Id: hal-02480067**

**<https://hal.science/hal-02480067v1>**

Submitted on 21 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Invertebrates facing environmental contamination by endocrine disruptors: novel evidences and**  
2 **recent insights**

3  
4 **Corresponding Author: Dr. Virginie Cuvillier-Hot, Ph.D.**

5 **Corresponding Author's Institution: Univ. Lille, CNRS, UMR 8198 - Evo-**

6 **Eco-Paleo**

7 **First Author: Virginie Cuvillier-Hot, Ph.D.**

8 **Authors: Virginie Cuvillier-Hot, Ph.D.; Alain Lenoir, Pr**

9  
10

11 **Abstract**

12 The crisis of biodiversity we currently experience raises the question of the impact of anthropogenic chemicals  
13 on wild life health. Endocrine disruptors are notably incriminated because of their possible effects on  
14 development and reproduction, including at very low doses. As commonly recorded in the field, the burden they  
15 impose on wild species also concerns invertebrates, with possible specificities linked with the specific physiology  
16 of these animals. A better understanding of chemically-mediated endocrine disruption in these species has  
17 clearly gained from knowledge accumulated on vertebrate models. But the molecular pathways specific to  
18 invertebrates also need to be reckoned, which implies dedicated research efforts to decipher their basic  
19 functioning in order to be able to assess its possible disruption. The recent rising of omics technologies opens  
20 the way to an intensification of these efforts on both aspects, even in species almost uninvestigated so far.

21 **Keywords:** endocrine disruption; endocrine disrupting chemicals (EDCs); nuclear receptors; oestrogeno-  
22 mimetics; ecdysteroids ; Juvenile hormone; omics; mollusks; annelids; arthropods

23

24 **1. Introduction**

25 Anthropogenic pollutants are seen as one of the main causes concurring to the ongoing collapse of global  
26 biodiversity. Among them endocrine disrupting chemicals (EDCs) pervade virtually all ecosystems on Earth. They  
27 are largely reported in marine as well as freshwater environments, notably since actual treatment systems fail to  
28 completely remove them from wastewater (*e.g.* Aris et al., 2014; Huang et al., 2019; Salgueiro-González et al.,  
29 2015). They also complex to sediments where they can accumulate and persist, sometimes leading to  
30 permanent release in the milieu following soil erosion. At last, many EDCs (notably phthalates, polychlorinated  
31 biphenyl (PCBs), polycyclic aromatic hydrocarbons (PAHs), brominated flame retardants, dioxins, alkylphenols,  
32 perfluorinated chemicals (PFCs) and also some pesticides) have been shown to be present in the atmosphere at  
33 worrying concentrations (Annamalai and Namasivayam, 2015), and maybe conveyed by the microplastic  
34 particles recently detected in the atmospheric compartment (Dris et al., 2016). Although they belong to various  
35 classes of synthetic and natural compounds, the common feature of EDCs lays in that they can interfere at  
36 different levels of the endocrine system of animals, disrupting physiological, biochemical and /or molecular  
37 processes that control development, growth or reproduction; recent works add neurological and immune  
38 processes to the list (Jones et al., 2017). Toxicological studies have uncovered multiple effects of EDCs, many of  
39 which are hard to predict due to non-monotonic dose-response, frequent cocktail effects and transgenerational  
40 implications (Flint et al., 2012; Skinner, 2014; Vandenberg et al., 2012; Xin et al., 2015; Z. Xu et al., 2017). Beyond  
41 actual concerns as regards human health, the large spread of these molecules, through all ecosystems, from

42 Arctic ices to primitive rainforests, also raises concern about the effects of such contamination on animal health  
43 and, ultimately, on worldwide biodiversity (Lenoir et al., 2016; Lyons, 2006; Sonne, 2010). In this respect,  
44 studying endocrine disruption in invertebrate species is meaningful. First, from an anthropocentric point of view,  
45 invertebrates are tremendous tools to detect and quantify the presence of EDCs in natural substrates, and thus  
46 may serve as useful bioindicators. They also allow us to gain knowledge about EDC toxicological properties at the  
47 scale of the individual's lifetime, which is much more difficult with vertebrate models (Hutchinson, 2007).  
48 Besides, due to their small size which make them easy to handle and stock in the laboratory, and to the large  
49 quantity of material they provide when rearing is possible, they represent a convenient and quick option to set  
50 up multigenerational assays to investigate potential transgenerational effects of EDCs. Second and most  
51 importantly, from an ecological point of view, invertebrates represent a huge part of worldwide biodiversity and  
52 fulfill many crucial ecological roles in every ecosystem (Wilson, 2006); this implies that their present decline is a  
53 clear worrying issue that requires to be handled urgently.

54 Because of their specific physiology (that diverged from that of vertebrates more than 600 MA ago) it appears  
55 necessary to develop and use invertebrate models allowing to detect and predict the whole set of effects of  
56 environmental EDCs on invertebrates wild populations. Contamination routes are likely multiple, with  
57 proportions depending on species' life history, and often different from that of vertebrates. In particular,  
58 because of their small size and thus relatively high surface/volume ratio compared to vertebrates, invertebrates  
59 suffer from higher relative exposure levels, which can imply a higher bioaccumulation, or the need to develop  
60 greater excretion capacities. In the aquatic environment, absorption coefficients may be higher compared to  
61 vertebrate species due to the properties of invertebrates body wall, which is often a functional surface of  
62 exchange, for respiratory gases for instance. As regards terrestrial invertebrates, direct contact and oral route  
63 are so far considered as the most probable ways of contamination. However, the possible role of arthropods  
64 cuticle as a trap for atmospheric pollutants just begins to be studied. It could constitute an effective way of  
65 contamination disregarded so far (Lenoir et al., 2014, 2012). Noteworthy, invertebrates generally lack the blood-  
66 gonad barrier that provide some protection to reproductive organs in mammals. At last, as for fishes and  
67 amphibians, mature gametes are in most cases directly released in the environment, which implies an exposure  
68 of organisms all along the life cycle – from gametogenesis to adulthood – including the highly vulnerable first  
69 stages of development. In short, apprehending the consequences of invertebrates exposure to EDC implies to  
70 take into account many of their peculiarities, and specific researches are required in this field, which is reflected  
71 by the growing body of literature dedicated to the subject.

72 In this review, we will first try to gauge the burden environmental EDCs place on invertebrate populations in  
73 nature, focusing as much as possible on studies that evaluate disruption in wild caught animals, exposed to EDCs  
74 in real situations. Then, we will produce an overview of the most recent molecular insights that may enlighten  
75 our comprehension of endocrine disruption in invertebrates. Finally, we will discuss the contribution of the  
76 recent approaches that rely on omics, to get a wider view of EDCs impact on invertebrate physiology. Note that  
77 the vast majority of our current knowledge relates to protostome species (mainly arthropods, mollusks and  
78 annelids), which explains that the present review will mainly focus on this taxonomic group.

## 79 80 **2. Delineating the burden of endocrine disruption on wild invertebrates**

81 Risk assessment procedures have been using invertebrate models (mainly aquatic species) for long, both to  
82 detect potential EDC activities in mixtures, and to gain a basic understanding of the ecological effects of known  
83 EDCs. Conversely, field studies demonstrating EDCs effects on invertebrate wild populations and communities  
84 are scarcer, in aquatic as well as in terrestrial ecosystems (Amiard and Amiard-Triquet, 2015).

### 85 **2.1 In aquatic ecosystems**

86 In aquatic ecosystems, the most relevant case certainly remains the imposex phenomenon that affects  
87 gastropod mollusks living in areas contaminated by organotin: tributyltin [TBT] and triphenyltin [TPT], at levels  
88 as low as few ng per liter (DeFur, 1999). These organometallic compounds have been used as antifouling agents

89 for several decades until their progressive ban by most countries, in the 80s. However, incomplete observation  
90 of worldwide legislation, as well as organotins desorption from contaminated sediments explain that organotins  
91 contamination is still a major toxicological concern (Laranjeiro et al., 2018). At the individual level, endocrine  
92 disruptive effects in contaminated areas result in the imposition of male sex characteristics on female snails,  
93 with a clear relationship between the extent of the masculinization process and the dose of organotins detected  
94 in the environment (Graceli et al., 2013). At some point, affected females suffer from impaired reproduction  
95 ending up to sterility and even to death (DeFur, 1999). Individual defects due to environmental contamination  
96 thus translate into population-level concerns, with a demographic fall and acute risk of local extinction in the  
97 worst polluted areas (Bryan et al., 1986; Langston et al., 2015; Roach and Wilson, 2009). In the field, TBT was  
98 also shown to affect other invertebrates, disturbing shell calcification and reproduction in bivalve mollusks, and  
99 affecting crustacean growth, carbohydrate and lipid metabolism and sexual maturation (Graceli et al., 2013;  
100 Vogt et al., 2018). More recently, concerns were raised over the toxicity of TBT in terrestrial ecosystems that can  
101 be contaminated notably through soil enrichment with marine sediments or sewage sludge. Silva and colleagues  
102 (2014) assessed TBT toxicity in terrestrial invertebrates and observed negative effects on the food consumption  
103 and assimilation capabilities in isopods (*Porcellionides pruinosus*), and on juvenile production and/or mortality in  
104 collembolans (*Folsomia candida*). But additional studies in field conditions are still required, notably in terms of  
105 TBT doses, especially in soils continuously enriched with sewage sludge.

106 In line with the vertebrate (and especially fish) situation, field exposure cases to estrogenic compounds (e.g.  
107 xeno-estrogens such as 17 $\alpha$ -ethinylestradiol (EE2), bisphenol A (BPA), nonylphenols and octylphenols) provide  
108 another illustration of endocrine disruption in wild aquatic invertebrates (Amiard and Amiard-Triquet, 2015; Jin  
109 et al., 2012), with effects including sex-ratio modification, delayed sexual maturity and intersexuality. Numerous  
110 studies have reported disruptive effects of estrogens in invertebrate models, including at concentrations  
111 compatible with doses measured in the field (ng/l) (Aris et al., 2014; Bovier et al., 2018; Flint et al., 2012; Herrero  
112 et al., 2015; Leonard et al., 2017; Liu et al., 2012; Morales et al., 2018; Oehlmann et al., 2009, 2007; Oetken et  
113 al., 2004; Wright-Walters et al., 2011). Experimental studies have also demonstrated that the reproduction of  
114 the *Potamopyrgus antipodarum* snail (a freshwater mollusk quite abundant in Europe) is impaired by the  
115 exposure to a mixture of environmental estrogens. This response was found to be comparable to that of four  
116 species of freshwater fishes, with a similar sensitivity to these molecules (Jobling et al., 2004), suggesting that  
117 EDC concerns extend to wild invertebrate populations exposed to estrogenic effluents, especially as regards  
118 sessile species. In good agreement with this study, when caged downstream of an effluent discharge, the same  
119 snail species was found to bioaccumulate more alkylphenols, BPA, Estradiol and Testosterone than when  
120 animals were located upstream. These experiments also demonstrated a sharp decrease of reproductive  
121 parameters in these animals (reduction in the number of embryos produced) after 6 weeks of exposure (Gust et  
122 al., 2014). Concomitant measures of egg proteins and mRNA levels allowed authors to demonstrate that  
123 reproductive disruption did not affect oocyte development, but rather targeted early embryonic development,  
124 suggesting the disruption of estrogen signaling pathways (Gust et al., 2014). Recent *in situ* surveys on the marine  
125 mussel *Mytilus trossulus* also evidenced some reproductive impairment in bivalve wild populations exposed to  
126 presumed estrogenic water wastes. Individuals living in the vicinity of a sewage purification plant outlet showed  
127 an increased frequency of gonadal regression and atresia and malformations, which were successfully mimicked  
128 by an experimental exposure to 50 or 500 ng/dl of EE2 (Smolarz et al., 2017). Such experimental exposure in  
129 natural conditions is a powerful way to evidence local contamination by EDCs, but requires a careful choice of  
130 the biomarkers recorded. The level of yolk protein vitellogenin, most often estimated by the Alkali Labile  
131 Phosphate method, has been commonly used as a biomarker of feminization in several aquatic species. Both the  
132 method (Sánchez-Marín et al., 2017) and its results (Boulangé-Lecomte et al., 2017; Short et al., 2014) have  
133 recently been questioned by proteomic and transcriptomic approaches, respectively.

134 In freshwater ecosystems, chironomids are increasingly imposing themselves as a convenient and suited model  
135 of study. Studies carried out in *Chironomus riparius* recently contributed to reveal the endocrine disruptive  
136 effects of complex mixtures of heavy metals in experimental settings in which larvae were reared in sediments  
137 (more or less contaminated) originating from abandoned mines, polluted shores or reference sites. Among other  
138 negative effects recorded as regards biomass and respiration rate, exposed individuals showed disrupted

139 expression of ecdysone-related genes, impaired reproduction and growth, both at the larval and adult stages  
140 (Arambourou et al., 2020, 2019).

## 141 **2.2 In terrestrial ecosystems**

142 In terrestrial ecosystems, the main cases for which endocrine disruption effects are well identified under field  
143 conditions relate to insect growth disruptors (IGD, formerly named insect growth regulators or IGR). Such  
144 substances have been designed by purpose to affect insect growth and development through a specific  
145 disruption of their endocrine regulatory pathways, including juvenile hormone analogues (JHA), and steroidal  
146 ecdysone agonists (Pener and Dhadialla, 2012). For these molecules, the drop of target insect populations is the  
147 hallmark of their efficacy. Unfortunately, although the design of these molecules was intent to produce more  
148 selective pesticides (they are often sold as “reduced-risk insecticides”), non-target species sharing similar  
149 hormone systems can also be affected. Pener and Dhadialla (2012) reviewed several studies demonstrating  
150 lethal or sublethal effects of IGD on non-target species, especially predators or parasitoids of the target species,  
151 proving the diffusion of IGD in food webs. Lawler (2017) recently reviewed the environmental effects of  
152 methoprene use for mosquito control and pinpointed some negative issues at realistic environmental levels,  
153 mainly on aquatic invertebrate species: lobster larvae, mysid shrimp embryos, small diptera and zooplankton-  
154 sized crustacean. Although often disregarded, these species, and especially planktonic species which occupy key  
155 positions in aquatic food webs, for sure deserve further attention on the subject of endocrine disruption in field  
156 context. Regarding IGD’s use for agronomic purposes, the potential threatening of pollinators and other  
157 beneficial insects’ survival is a big subject of concern, due to the huge ecosystemic services they fulfill. This  
158 concerns domestic honeybees but also Lepidoptera, parasitic wasps, Coleoptera and numerous wild bees that  
159 play crucial roles in plant-insect mutualistic networks (Quesada and Sadof, 2019; Steffan-Dewenter et al., 2005).  
160 Beyond direct mortality provoked by many pesticides, some IGD are responsible of sublethal effects that may  
161 contribute to the actual decline of (wild and managed) pollinators and beneficial insects populations (Johnson et  
162 al., 2010; Mommaerts and Smagghe, 2011). Bees illustrate how complex and multiple the routes of exposure can  
163 be: in addition to direct contact associated with spraying as well as the interaction with treated plants, foragers  
164 collect nectar and pollen susceptible to be contaminated. They also bring it back to the nest to feed all the  
165 members of the colony (social bees) or their larvae (solitary bees). For brood as well as for hive bees,  
166 contamination thus probably mainly arise through food intake, which may strongly lead to specific toxic effects  
167 (Sanchez-Bayo and Goka, 2014; Stanley and Preetha, 2016). Recent studies evidenced a clear contamination of  
168 the pollen stored into beehives, by many pesticides and herbicides at sublethal concentrations, among which  
169 ecdysone analogues (tebufenozide, methoxyfenozide) and JHA (fenoxycarb, pyriproxyfen, methoprene) (Böhme  
170 et al., 2018; Calatayud-Vernich et al., 2018; Hakme et al., 2017). Experimental work had previously  
171 demonstrated that exposure of micro-colonies of *Bombus terrestris* to JHA (pyriproxifen or kinoprene), through  
172 pollen contamination (e.g. not through sugar water contamination) at the maximum field recommended  
173 concentration, has no acute toxicity on workers but enhances larval mortality. This effect was hypothesized to  
174 be linked with a lethal blocking of development before metamorphosis (Mommaerts et al., 2006). In the same  
175 study, authors evidenced a stimulatory effect of a very low dose of kinetoprene (0.0650 mg/l) on the  
176 development of the female reproductive system, together with a higher reproductive output, strongly arguing in  
177 favor of an endocrine nature of the disruption. Even though neurotoxic pesticides – and in the first place all  
178 neonicotinoids – are the most incriminated molecules in the complex interactions that lead to bee CCD (colony  
179 collapse disorder; Sanchez-Bayo and Goka, 2014), IGD chronic exposure should not be disregarded, as regards its  
180 potential sublethal effects and possible involvement in synergistic effects. Furthermore, neonicotinoids  
181 themselves were recently shown to exert EDC-like effects in bees, generating sublethal effects suspected to  
182 largely contribute to population extinctions over a long time scale (Woodcock et al., 2016; Baines et al., 2017).  
183 More field studies are now needed to assess and cross-reference several key parameters of pollinators’ potential  
184 poisoning by pesticides. These include (1) the dynamics of pollen contamination in relation with its species of  
185 origin and the nature of the contaminants; (2) the persistency of chemicals under specific conditions inside real  
186 hives and along seasons; (3) the link between external exposure (direct contact and/or chronic ingestion) and  
187 potential risk; (4) the possible interaction between contaminants, which needs to be addressed by testing

188 realistic pesticide mixture (see Böhme et al., 2017); (5) the direct effects on bee survival and activity but also  
189 sublethal effects (fertility, abnormal larval development) that may affect colony productivity; (6) the side-effects  
190 that may arise through the disturbance of other functions such as olfactory performance (see e.g. Chakrabarti et  
191 al., 2015), orientation and learning capacities, which are essential to foraging process and feeding, flight activity  
192 and foraging efficiency (see Prado et al., 2019), pheromonal communication (notably blurring of nestmate  
193 identification) and social interactions (see e.g. Fourrier et al., 2015); and (7) the indirect adverse effects that may  
194 ensue from microbiota modifications.

195 More generally, all herbivorous invertebrates may come across EDCs through the contamination of their food:  
196 some well identified or emerging organic contaminants including BPA, nonylphenol or triclosan are largely  
197 distributed in the water cycle and end up in reclaimed water eventually used for food crop irrigation. This can  
198 lead to their accumulation in leafy vegetables (Dodgen et al., 2013) and to a direct exposure of herbivorous  
199 species. In the insect crop pest *Spodoptera littoralis*, Maria and colleagues (2019) evidenced a higher mortality at  
200 pupal stage and an increase in some larval instar duration associated with modifications of ecdysteroid titers  
201 and nuclear receptor expression, after consumption of food contaminated with BPA concentrations similar to  
202 that found in plants. At last, atmospheric contamination also potentially concerns all epigeic species. Although  
203 this route of exposure has barely been considered so far, we recently evidenced a chronic contamination of  
204 several species of aerial insects (mainly ants but also crickets and honeybees) by phthalates, which are potent  
205 EDCs (Oehlmann et al., 2009; e.g. Mankidy et al., 2013). These phthalates impregnate the cuticle and mix with  
206 the cuticular components (Lenoir et al., 2012). In these studies, all tested individuals were found to be  
207 contaminated, with varying contamination levels depending on the species. Phthalates accounted for 0.11 to  
208 2.66% of the cuticular compounds in the different ant species tested, 0.73% in honey bees and 2.76% in the  
209 wood cricket *Nemobius sylvestris*. This contamination was clearly established to originate from the atmosphere.  
210 It seems that, due to its biochemical properties, insects cuticle can trap the phthalates present in both the  
211 vapor phase and the particulate phase (adsorbed to atmospheric particles; Lenoir et al., 2014; Teil et al., 2006).  
212 The insect cuticle keeps trace of this contamination in the form of a basal level of phthalates mixed to cuticular  
213 compounds, which can be measured by GC-MS. In addition, phthalates were also clearly demonstrated to be  
214 able to cross the epidermal barrier in these insects, as they were detected inside the body of animals, mainly in  
215 fat. Experimental contaminations by environmental doses spiked onto the cuticle induced a reduction of queens'  
216 eggs output and disturbed immune gene expression in workers of the black garden ant (*Lasius niger*), suggesting  
217 possible endocrine disruptive effects in natural populations (Cuvillier-Hot et al., 2014). Considering that these  
218 observations may potentially apply to all invertebrate species possessing a cuticle or a lipophilic integument, it  
219 appears necessary to pursue investigations on other species, in order to examine whether it is necessary to  
220 include this mode of contamination in risk assessment protocols for terrestrial invertebrates.  
221

### 222 3. New insights in invertebrate molecular disruption

223 As an emerging science, EDC ecotoxicology first based on a vertebrate-centered approach, taking advantage of  
224 our better knowledge of vertebrates physiology, to detect and identify disturbances linked with environmental  
225 exposure to contaminants. Then, knowledge acquired from vertebrate models was transposed to invertebrates  
226 and researches have focused on susceptible endocrine pathways shared between invertebrates and vertebrates,  
227 and in a first place on the possible disruption of the estrogen receptor (ER)-associated cascades. But, as more  
228 than 600 Million years have elapsed since the split between protostomes and deuterostomes, the former  
229 acquired endocrine systems of regulation unique to them (e.g. ecdysteroids, Juvenile hormones), which are  
230 potential targets for environmental endocrine disruption and which call for dedicated researches. Even  
231 nowadays, while substantive knowledge has been accumulated for some groups (e.g. insects), a main gap of our  
232 current knowledge over endocrine disruption in invertebrates remains our lack of detailed comprehension of  
233 endocrine signaling pathways in many disregarded groups of invertebrates (Castro and Santos, 2014;  
234 Hutchinson, 2007). This concerns receptors and ligands, associated cascades and regulation, but also the  
235 dynamics of bioaccumulation and elimination of potential disruptors. This basic knowledge is necessary to  
236 ascertain that an adverse effect observed in the field is indeed the result of an endocrine disruption mechanism

237 rather than a more general direct or indirect toxic effect (Barata et al., 2004; Hutchinson, 2007; Lagadic et al.,  
238 2007; Weltje and Schulte-Oehlmann, 2007). In the past years, some attempts have been made to develop  
239 arthropods-specific *in vitro* assays dedicated to screen environmental contaminants for ecdysteroid  
240 agonist/antagonist activities (Dinan et al., 2001; Kontogiannatos et al., 2015; Soin et al., 2009; Swevers et al.,  
241 2004; Yokota et al., 2011; Zotti et al., 2013). However, it appears that ecdysteroid receptors have ligand-binding  
242 properties that can substantially vary between invertebrate species, even among members of a same order  
243 (Graham et al., 2007). This suggests that the results of such molecular assays can hardly be generalized among  
244 species on a large scale and involve the need to develop taxa-specific tests (Santos et al., 2018; Yokota et al.,  
245 2011).

246 EDCs can interfere with the endocrine pathways of animals in many ways. They can affect endogenous hormonal  
247 levels, or those of hormone receptors by disturbing their specific synthesis and/or catabolism; they can also  
248 disturb hormonal receptor functions by spatially mimicking endogenous hormones, thus affecting receptors  
249 availability or leading to unexpected signalization (agonist or antagonist). For decades, physiological and  
250 developmental endpoints have been used to detect and characterize EDCs in invertebrates, all the more that  
251 several model species, easy to breed and possessing a short generation time are available in several taxonomic  
252 groups. Many reviews have reported about lists of suspected EDCs and proposed modes of action based on the  
253 examined studies (e.g. Hutchinson, 2002; Lagadic et al., 2007; LeBlanc, 2007; Oehlmann et al., 2007; Oetken et  
254 al., 2004; Rodríguez et al., 2007; Soin and Smaghe, 2007; Zou, 2005). More recently, molecular approaches  
255 allowed a deeper investigation of the mode of action of these EDCs in invertebrates, which we will focus on in  
256 the next chapter. Figure 1 sums-up the main molecular regulatory pathways hypothesized or demonstrated to  
257 be affected by chemicals in protostomes.

### 258 **3.1 Molecular pathways susceptible to endocrine disruption and shared with vertebrates**

#### 259 3.1.1. Estrogen receptors and relatives

260 In vertebrates, many EDCs exert their action as agonists of estrogen receptors (ER) or/and antagonists of  
261 androgen receptors (AR). The invertebrate orthologs of these receptors are thus potential targets for  
262 environmental endocrine disruption. To date, the only steroid receptor evidenced in protostomes belongs to the  
263 ER family and shares a common ancestor with vertebrate steroid receptors (SR) named AncSR, which would  
264 have arose before protostome and deuterostome cleavage (Castro and Santos, 2014; Jones et al., 2017; Markov  
265 et al., 2009). Absent from the Ecdysozoa, ERs orthologs have been characterized in mollusks (Hultin et al., 2014;  
266 e.g. Keay et al., 2006; Matsumoto et al., 2007; Raingeard et al., 2013; Thornton, 2003; Zhang et al., 2012),  
267 annelids (Keay and Thornton, 2009; Lv et al., 2017) and recently in rotifers (Jones et al., 2017). As in vertebrates,  
268 these receptors have genomic signaling pathways that activate gene transcription through the direct binding of  
269 the nuclear ER complex to ERE (estrogen-response elements) sites (or indirectly through AP-1 or Sp-1 binding  
270 sites) upstream of target genes (Hamilton et al., 2017). However, the modalities of ER commitment clearly differ  
271 among invertebrate groups. Indeed, the Mollusca receptor was shown to have been vestigialized through some  
272 decisive substitutions that froze the ER in an active conformation, resulting in its constitutive activity.  
273 Subsequent mutations then filled the ligand pocket with bulky residues precluding any ligand binding and  
274 rendering unlikely any reversal to the ancestral function (Bridgham et al., 2014). ERs in mollusks are thus ligand-  
275 independent transcriptional activators, which implies that mechanisms of endocrine disruption by estrogenic  
276 environmental contaminants in these species are necessarily independent of any nuclear ER activation (Canesi  
277 and Fabbri, 2015). On the contrary, annelids and rotifers ERs display the classic properties of vertebrates ER,  
278 including apparent sensitivity to estrogens (Castro and Santos, 2014; Jones et al., 2017; Keay and Thornton,  
279 2009). However, the exact nature of their endogenous ligand is still unknown. Steroidogenesis exists in  
280 protostomes but has evolved independently from vertebrate cascades, partly through the recruitment of  
281 cytochrome P450 enzymes formerly involved in xenobiotic detoxification (Markov et al., 2009). Recent work by  
282 Blalock and collaborators (2018) evidenced, for the first time in a protostome (the mussel *Mytilus edulis*), a  
283 partial protein equivalent to vertebrate CYP11A enzyme, responsible for the side-chain cleavage of cholesterol  
284 to pregnenolone. Hence, within the whole steroid biosynthesis pathway, only the last step of aromatization of  
285 testosterone into estradiol would still be missing in lophotrochozoa, the aromatase CYP19A appearing restricted

286 to chordates (Goldstone et al., 2016; Markov et al., 2009). For this reason and also because unbiased methods to  
287 detect steroids in invertebrate tissues so far failed to find estrogens, some authors refute the idea of an  
288 estrogen-like endogenous ligand for protostome ERs (Holzer et al., 2017a; Scott, 2013). But whatever their origin  
289 – endogenous or exogenous – estrogens clearly affect the physiology of many invertebrates, including mollusks,  
290 paving the way to possible endocrine disruption by oestrogeno-mimetics. In vertebrates, ERs can be activated by  
291 several estrogenic endogenous ligands (estradiol,  $\Delta^5$ -androstenediol, 5 $\alpha$ -androstenediol, and 27-  
292 hydroxycholesterol). They also are responsive to several plant-derived compounds with estrogenic activity such  
293 as genistein, coumestrol, and resveratrol (Baker and Lathe, 2018). This promiscuous responsiveness is proposed  
294 to explain their response to synthetic chemicals such as BPA and phthalates (Engel et al., 2017; Flint et al., 2012).  
295 Similar cross-binding may be considered for ligand-sensitive ER described in protostomes. Keay and Thornton  
296 (2009) for example report an agonist effect of genistein and an antagonist effect of BPA on *Platynereis dumerilii*  
297 ER.

298 In vertebrate models, it is well established that ER signaling can be modulated by estrogen-related receptors  
299 (ERRs). These are also nuclear receptors sharing sequence similarity with ERs, but that respond very weakly to  
300 endogenous estrogens while being stimulated by some xenoestrogens. They may thus interfere with ER signaling  
301 upon environmental EDC exposure, along several modalities: (1) by directly interacting with ERs, (2) by  
302 competing with ERs for ERE site binding or (3) by targeting similar genes as ERs via the presence of ERR-response  
303 elements in their promoter region, alongside with ERE (Xu et al., 2017). As a consequence, influence of ERR  
304 activation on ER signaling will clearly depend on the level of expression and tissue distribution of both receptors.  
305 Sequences homologous to ERRs have been described in annelids (the marine worm *Capitella* and the leech  
306 *Helobdella*, Baker, 2008) and in mollusks (the snail *Lottia*, Baker, 2008; the snail *Physa*, Morales et al., 2018; two  
307 *Mytilus* species, Nagasawa et al., 2015). Quite expectedly, mRNA of ER homologues were abundantly detected in  
308 reproductive tissues (ovary but also pedal ganglion) in the mussels *Mytilus edulis* and *M. galloprovincialis*, while  
309 mRNA of ERR homologues, otherwise involved in metabolism regulation and mitochondrial function (Hubbard et  
310 al., 2015), had high levels of expression in the gill and digestive gland ; however, both receptors had basal levels  
311 of expression in all tissues examined and may thus interfere with each other (Nagasawa et al., 2015). In *M.*  
312 *edulis*, ER expression in ovary appeared regulated by *in vitro* exposure to 17 $\beta$ -estradiol, while that of ERR did not  
313 (Nagasawa et al., 2015). In the snail *Marisa cornuarietis*, various vertebrate-like estrogens were tested for their  
314 possible ER and ERR gene transcription modulation, but gave negative results for both genes. Only genistein and  
315 BPA up-regulated gene expression of ER and ERR orthologs respectively (Bannister et al., 2013). In this species,  
316 BPA exposure at environmentally relevant concentrations had clear disruptive effects, leading to a  
317 superfeminization phenomenon characterized by enlargement and malformations of female organs, and higher  
318 egg mass production (Oehlmann et al., 2006). Also in the snail *Physa acuta*, BPA exposure led to a significant  
319 increase in the mRNA levels of both ER and ERR, suggesting that these receptors could be involved in molecular  
320 events that regulate the endocrine disruptor activity of this chemical in Gastropods (Morales et al., 2018).  
321 Interestingly, ERR gene homologues have also been characterized in insects and their expression appeared  
322 modulated by EE2 (Bovier et al., 2018) and ethylparaben (Liu et al., 2014) in *Drosophila melanogaster*, and by  
323 Bisphenol S (BPS), Triclosan, BPA, nonylphenol or diethylhexyl phthalate (DEHP) in *Chironomus riparius* larvae  
324 (Herrero et al., 2018, 2015; Martínez-Paz et al., 2017; Park and Kwak, 2010). In these cases, disruption is  
325 proposed regarding the 20-Hydroxyecdysone pathway (see § 3.2), highlighting the multi-target character of  
326 many EDCs.

327 Alongside and apart from direct interactions of estrogeno-mimetics with nuclear estrogen receptors, many  
328 studies in vertebrates pointed out more rapid effects of these molecules (within seconds to minutes) through  
329 non-genomic signaling pathways (Xu et al., 2017). They may be initiated by estrogen-responsive non-nuclear  
330 receptors and may lead to local effects such as modification of ion fluxes or to activation of cytosolic kinase  
331 cascades that ultimately affect gene transcription. Several studies reviewed by Janer & Porte (2007) suggested  
332 the involvement of such non-genomic processes in endocrine regulation also in invertebrates. Recent  
333 contributions from Omics studies brought further support: individuals of the mudsnail *Potamopyrgus*  
334 *antipodarum* exposed to an effluent discharge, while accumulating alkylphenols, BPA and vertebrate-like sex-  
335 steroid hormones, showed an inhibition of the expression of genes involved in non-genomic signaling pathways,



336 together with an induction of the repressors of the genomic pathway; this led to a drastic decrease in embryo  
337 production (Gust et al., 2014). Similarly, a transcriptomic approach of mussels exposed to EE2 or 4-nonylphenol  
338 (leading notably to female-skewed sex-ratios) revealed similar patterns of gene dysregulation with main targets  
339 belonging to the non-genomic estrogen signaling pathway (Blalock et al., 2018).

340

341

### 3.1.2. Other nuclear receptors

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

At the molecular level as at the individual and population scales, the case of endocrine disruption by organotins was much investigated and led to many relevant trails. In the imposex phenomenon, endocrine disruption is highly suspected at different levels. High levels of testosterone observed in TBT treated prosobranch females could ensue from an inhibition of enzymes involved in steroid metabolism (reviewed in Graceli et al., 2013, Lafont and Mathieu, 2007). TBT may also induce adverse neuromodulation through the production of APGWamide which was shown to induce imposex in mud snails (*Ilyanassa obsoleta*, Oberdörster and McClellan-Green, 2002). Finally Pascoal and colleagues (2013) strengthened the hypothesis of a disruption of the retinoid X receptor (RXR) – peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) pathway by evidencing a strong transcriptional response of the retinoid receptors and of putative members of this signaling pathway after TBT treatment in *Nucella lapillus*. They also showed that activating this pathway with a vertebrate ligand of PPAR $\gamma$  led to imposex in this species. As additional support to this third presumed mechanism, it had been previously established that TBT was able to bind and activate RXR–PPAR vertebrate heterodimers, by mimicking 9cis-retinoic acid interaction with RXR (le Maire et al., 2009). All these clues suggest a strong involvement of this signaling pathway, quite shared between vertebrates and invertebrates, in imposex physiopathology (André et al., 2014; Iguchi and Katsu, 2008).

RXRs are nuclear receptors involved in numerous signaling pathways and biological functions since they can signal as homodimers, but also form heterodimers with many partners, such as PPAR (as exemplified above), retinoic acid receptor (RAR) or thyroid hormone receptor (TR). In Ecdysozoans, RXRs homologues (known as ultraspiracle (USP) in insects) are the obligate heterodimeric partners of ecdysteroid receptors (EcR, see §3.2). RXRs are widely distributed among metazoans and show a great degree of homology between taxa. Furthermore, their sensitivity to 9-cis-retinoic acid (RA) appears as an ancestral character already present in the common ancestor of Cnidaria and Bilateria, widely shared among metazoans and secondarily lost in ecdysozoans (André et al., 2017). Their functional partner in RA sensing, RARs, have also been detected in some annelids and mollusks. In both cases, the essential of the signaling machinery is present so that functional RA signaling pathways is expected, but according to specific processes, given that the molluscan RAR were proved insensitive to retinoids (Gutierrez-Mazariegos et al., 2014; Handberg-Thorsager et al., 2018). Finally, genome data mining recently identified key players possibly involved in retinoid metabolism and storage process in some protostomes, providing new potential targets for disruption. All this lead André et al. (2017, 2014) to suggest that RXR-dependent modulation by organotins might be evolutionary conserved among lophotrochozoans and vertebrates, and foresee that many more metazoan species might be potential targets for EDCs through disruption of the retinoid system. The identification of the physiologic effects of such disruptions clearly deserves future research in all invertebrate groups.

Over the last years, the endocrine disruption of vertebrate biological processes controlled by thyroid hormone receptors (TRs) has been of growing concern, mainly because we progressively uncover the importance of this endocrine system on embryogenesis and early neurogenesis. Several EDCs including PCBs, the pesticide chlorpyrifos, BPA, poly-fluorinated surfactants and polybrominated diethyl ethers (PBDEs), are suspected to disrupt TH regulated pathways (Préau et al., 2015). TRs are also nuclear receptors functioning as ligand-dependent transcription factors; they mediate thyroid hormone (TH) effects by genomic and non-genomic mechanisms. If the genome of *Drosophila melanogaster* and *Caenorhabditis elegans* appeared deprived of TR-like sequences, lophotrochozoans on the contrary clearly possess TR orthologs (Orozco et al., 2017; Sainath et al., 2019). Note that putative TR orthologs have been proposed from *Daphnia pulex* and *D. magna* genomes, but their high sequence divergence poses question about their real origin (Sainath et al., 2019). Activating ligands are clearly different for protostome and deuterostome TRs, in agreement with the poor degree of conservation of their respective ligand-binding domains; and so far, endogenous ligands of protostome TRs remain unknown.

386 Still, endogenous synthesis of TH by the sea hare *Aplysia californica* is supported by its enzymatic equipment  
387 (Heyland et al., 2006), T4 and T3 have been extracted from *Crassostrea gigas* hemolymph (Huang et al., 2015),  
388 and TH were shown to induce settlement and metamorphosis in several mollusk species suggesting that TH  
389 signaling is possible in Lophotrochozoa (Holzer et al., 2017b; Taylor and Heyland, 2017). Although up to now no  
390 clear link has been established between TH and activation of TR in this group, the question remains open about  
391 the possible effect of environmental thyroid disruptors on invertebrates development and physiology, and  
392 urgently requires fundamental scientific work to understand the role of the thyroid endocrine system in  
393 protostomes.

394  
395

### 3.2 Molecular pathways susceptible to endocrine disruption and specific to Protostomes

396 Protostomes have developed specific endocrine regulatory pathways that can also be affected by environmental  
397 EDCs. In arthropods, there are three main classes of hormones: peptide hormones, ecdysteroids (mainly 20-  
398 hydroxyecdysone (20E) in insects and crustaceans, together with ponasterone A in chelicerate species), and  
399 terpenoids (mainly juvenile hormones (JH) in insects, methyl farnesoate (MF) in other arthropods). All three  
400 systems interact to regulate notably growth, development and reproduction. The hormonal action of 20E exerts  
401 via the activation of the ecdysone receptor (EcR), again a member of the superfamily of nuclear hormone  
402 receptors, that regulates gene transcription through heterodimerization with RXR (called USP in insects) and  
403 genomic binding to EcRE (ecdysone responsive element). It is now clear that the developmental events that lead  
404 to prepupal and pupal molt depend on a tightly timed dynamic of hemolymphatic 20E titer. A sharp rise  
405 followed by a clear decrease in the circulating levels of 20E are both important to regulate the expression of the  
406 different waves of transcription factors that coordinate the different molting processes, trigger apolysis, timely  
407 induce the expression of chitin synthase genes, regulate the production of the molting fluid needed to degrade  
408 the old cuticle and provoke its shedding (Guittard et al., 2011; Song et al., 2017). Numerous EDCs, beginning  
409 with IGDs that have been purposely developed to disrupt molting in insect pests (see §2.2), are able to  
410 negatively affect synthesis or secretion of 20E, or to interfere with EcR (Soin and Smagghe, 2007). As a  
411 consequence, it deregulates the fine-tuned signalization by 20E, causing notably molting failure and death, or  
412 impairing gametogenesis or embryogenesis. All the adverse consequences of chemically mediated disruption of  
413 molting in arthropods have recently been reviewed and illustrated by Song and colleagues (2017), highlighting  
414 the multiplicity of potential EDCs targets in the process that can lead to mortality by ecdysis failure. In many of  
415 the investigated cases, EDCs exert 20E agonistic effects, evidenced notably by an up-regulation of EcR or other  
416 genes involved in the ecdysone cascade (e.g. four oestrogeno-mimetics in an amphipod, Gismondi, 2018; BPA in  
417 the Lepidoptera *Sesamia nonagrioides*, Kontogiannatos et al., 2015; the insecticide fipronil in a copepod,  
418 Gaertner et al., 2012; Triclosan, Martínez-Paz et al., 2017; Nonylphenol, Nair and Choi, 2012; several commonly  
419 used UV-filters, Ozáez et al., 2014, 2013; the phthalate BBP, Planelló et al., 2011; BPA, Planelló et al., 2008; and  
420 Bisphenol S, a substitute of BPA, in the Chironome aquatic larvae, Herrero et al., 2018). For most of these  
421 chemicals, gene expression changes came along with affected developmental endpoints such as molting  
422 malformations, increased pupation time and reduced emergence success in *S. nonagrioides* (Kontogiannatos et  
423 al., 2015), emergence failures (Nonylphenol, Lee and Choi, 2006), delay in embryo hatching (Ozáez et al., 2014)  
424 and reduced survival (Planelló et al., 2008) in *Chironomus riparius*. Noteworthy, a cell-based reporter assay in  
425 *Bombyx mori* Bm5 cell line confirmed the agonistic effect of BPA in Lepidoptera (Kontogiannatos et al., 2015).  
426 Less frequently, EDCs act as anti-ecdysteroids (e.g. 5-chloro-1H-benzotriazole, tamoxifen or testosterone in  
427 *Daphnia magna*; resp. Giraud et al., 2017a; Jo et al., 2018; Mu and LeBlanc, 2002). It is clearly the case for the  
428 phthalate DEHP, which was shown to induce a significant drop in EcR transcription both at short exposures to  
429 high doses or at long exposures to low doses (i.e. environmental concentrations and below) in the Chironome  
430 larvae (Herrero et al., 2017; Planelló et al., 2011). Similar antagonist effects have been observed in the  
431 lepidopteran *Spodoptera littoralis*, also manifest through longer larval / pupal stages and global delay in adult  
432 emergence (Aviles et al., 2019). On the other hand, very few is known over ecdysteroid function and possible  
433 disruption outside of the arthropod group. EcR orthologs have been detected in the genome of mollusks,  
434 annelids and nematods and *in silico* analyses predict possible binding to an ecdysone-related steroid (Laguerre

435 and Veenstra, 2010). Interestingly, while the exact role of such molecules in lophotrochozoa physiology is so far  
436 unknown, the expression of EcR in the earthworm *Eisenia fetida* have been shown to be down-regulated in male  
437 reproductive tissues after exposure to low doses of BPA (Novo et al., 2018), and up-regulated in full body  
438 extracts after exposure to sublethal doses of 4-hydroxybenzophenone, a main product of degradation of  
439 benzophenone-3 that is currently used as UV-filter in sunscreens (Novo et al., 2019). The endogenous roles of  
440 such receptors in mollusks and annelids need clarification.

441 The sesquiterpenoid hormones, JH and MF, are also key regulators in arthropods, regulating various  
442 developmental and reproductive processes such as molting, growth, metamorphosis, and gonad maturation.  
443 The elucidation of the signaling pathway of sesquiterpenoid hormones in the last decade evidenced  
444 Methoprene-tolerant (Met; germ-cell expressed, Gce, in *Drosophila*) as their main receptor. The Met/Gce-  
445 hormone complexes are translocated to the nucleus, where they bind, together with the co-activator SRC (Tai in  
446 *Drosophila*), the specific JH response elements (JHRE) to activate transcription of the downstream target genes  
447 (Qu et al., 2018). Easy screening of putative EDCs with JH-like activity is now possible with the OECD TG211  
448 ANNEX 7 assay that use daphnids. Daphnid species are parthenogenetic species that produce only females in  
449 controlled conditions, but may generate male offspring in response to JHs. The validated assay takes advantage  
450 of this specificity to specifically detect JH analogs. Recently, Abe and colleagues (2015b) developed a derivative  
451 short-term screening assay using adult *Daphnia magna* and with a chemical exposure of only 7 days. This test  
452 notably ascertained the JH agonist effect of the IGD Diofenolan. In parallel, a two-hybrid assay evidenced that  
453 diofenolan induced heterodimerization of Met and SRC, suggesting its direct binding to the JH receptor;  
454 microarray analyses confirmed the dysregulation of many gene markers of JH action, evidencing this chemical as  
455 a strong JH agonist (Abe et al., 2015a). The same study evidenced a concomitant disruption of the ecdysteroid  
456 signaling pathway, reinforcing the hypothesis of a cross-talk between JH and ecdysteroid signaling pathways in  
457 arthropod developmental processes, as previously suggested with other JH agonists such as fenoxycarb,  
458 pyriproxyfen or Tris(2-butoxyethyl) phosphate, a flame retardant (Giraud et al., 2017b; Tuberty, 2005).  
459 Sesquiterpenoid hormones had long been thought restricted to arthropods. However, Schenk and colleagues  
460 (2016) recently established in *Platynereis dumerilii* that the hormone called nereidin, produced by head ganglia  
461 and known to regulate growth and sexual maturation in annelids by inhibiting the switch to reproductive state,  
462 corresponds to MF. They also evidenced that eleocytes, the cells that produce vitellogenin then captured by  
463 maturing oocytes, express a Met ortholog, whose transcription is up-regulated both by nereidin and by  
464 exogenous MF. Interestingly, the JHA methoprene and pyriproxyfen mimicked the effect of nereidin at similar  
465 concentration (10nM), reducing vitellogenin expression in cultured eleocytes. Recent insights over EcR and JH  
466 presence in annelids raise clear concerns about this group, notably as crucial members of the soil infauna. Their  
467 endogenous way of life makes them largely exposed to IGD sprayed on fields and cultures. We need to acquire  
468 background knowledge over the role of such putative regulatory pathways in this group so as to assess their  
469 possible disruption in natural conditions.

#### 470 **4. The promises of the high-throughput omics era**

471 Over the last decades, omics techniques became within reach of researchers to decipher how expressed genes  
472 (transcriptomics), proteins (proteomics) and metabolites (metabolomics) are altered by exposure to  
473 environmental pollutants (Colin et al., 2016). This gave rise to a new research field called ecotoxicogenomics  
474 that contributes to an ongoing mutation of our comprehension of the effect of endocrine disruption. Indeed,  
475 these recent techniques now make possible to compare different taxa on a large scale, to consider the  
476 multiplicity of the endogenous targets of individual toxics, and to monitor endocrine disruption in natural  
477 conditions much more precisely (Oliveira et al., 2016). Proteomics and metabolomics for instance enable to  
478 investigate how entire pathways react to toxic exposure, giving more weight to subtle changes that may have  
479 gone unnoticed if only few markers had been considered. Thereby, metabolomics profiling suggested that DEHP  
480 has negative effects on energy production and more particularly on TCA cycle, as in vertebrates, in the crop pest  
481 *Spodoptera littoralis* (Aviles et al., 2019). Tests on *Daphnia magna* metabolome confirmed that dissolved organic  
482 matter specifically affects the bioavailability of some drugs present in waste water (Kovacevic et al., 2019), and  
483 revealed convergent metabolic effects of some psychiatric drugs, despite different modes of action (Garreta-

484 Lara et al., 2018). Lastly, metabolomics approaches highlighted alterations in energy metabolism, amino acids  
485 metabolism and glycerophospholipid metabolism in mussels exposed to a synthetic progestin while no changes  
486 in gonad maturation and in steroids level were recorded (Cappello et al., 2017). For their part, coupled to NGS  
487 technologies, genomics and transcriptomics offer via *de novo* approaches an unprecedented power of  
488 description and analyses of gene sequences whose expression is altered after exposure, even in  
489 environmentally-relevant species for which physiological and genomic knowledge had been historically poor. It  
490 can reveal so far undetected disruption, and identify new biomarkers for the monitoring of given species in  
491 specific environmental conditions. For example, the transcriptomic analysis of the response to mercury exposure  
492 of two copepod species recently revealed an endocrine-disrupting potential for this metal, evidencing a  
493 dysregulation of the estrogen signaling pathway in both species (Wang et al., 2017). Similarly, microarray  
494 analyses of oysters exposed to microplastic particles revealed a molecular signature of endocrine disruption, in  
495 the form of differential expression of hormone receptors or transcripts involved in hormonal pathways  
496 (Sussarellu et al., 2016). Interestingly no EDC could have been detected in the biological samples of this study,  
497 illustrating the pre-eminence of bio-assays over chemical quantification when incriminated substances are active  
498 at very low doses and possibly below detection limits. Even more remarkable, omics approaches can unveil  
499 endocrine disruption even in cases where compensatory mechanisms suppress any noticeable physiological  
500 effects. For instance, in *Daphnia magna*, several studies screened the gene regulation changes after exposition  
501 to different potent EDC (Giraud et al., 2015; Houde et al., 2016). Contrary to other benzotriazoles, 1H-  
502 benzotriazole (BTR) has no effect on molting frequency but RNAseq analyses revealed the up-regulation of  
503 cuticular proteins after BTR exposure, suggesting the latter could have compensated a possible molting  
504 disruption (Giraud et al., 2017a).

505 We seemingly are at the beginning of a new era of research possibilities, with much more tools to apprehend  
506 complex toxic effects and intricate organism responses. This is especially true for invertebrates, so far neglected  
507 with respect to the huge biodiversity they represent. Furthermore, constant progresses in high-throughput  
508 sequencing techniques, both in terms of rapidity and prices, enhance the possibilities to compare multiple  
509 species, populations, life-stages or tissues, expanding the scope of experimental possibilities (e.g. An et al., 2014  
510 for a comparative study of testis and digestive gland response to estrogen exposure in *Chlamys farreri*). Finally,  
511 combining genetic, proteic or metabolic changes detected through omics approaches with the description of  
512 altered apical endpoints (e.g. body condition indexes, lipid content, fertility measures...) will undoubtedly  
513 provide useful insights into the genes, proteins and metabolites possibly implicated in physiological disorders,  
514 unveiling unknown molecular mechanisms of the physiological pathways affected (see Ciocan et al., 2012, and  
515 Grilo and Rosa, 2017 for examples about intersex in invertebrates). A good example is probably that of the dog  
516 whelk (*Nucella lapillus*), for which transcriptomics coupled to *in vitro* testing confirmed the alteration of some  
517 processes already evoked (steroid metabolism, neuroendocrine regulation and retinoid mechanisms) following  
518 TBT exposure, but also revealed the involvement in imposex of a regulatory pathway (peroxisome proliferator-  
519 activated receptor or PPAR), not reported until then in invertebrates (Pascoal et al., 2013). The next challenge is  
520 now to integrate omics in routine environmental monitoring studies notably to help regulatory agencies in their  
521 management of environmental risks (Gouveia et al., 2019; Oliveira et al., 2016; Piña et al., 2018).

522

## 523 5. Conclusion

524 A growing research effort coupled with the use of ever more accurate and efficient methods and approaches  
525 now allow us to better identify EDCs and refine our knowledge of their mode of action, even in non-model  
526 animal species for which basic knowledge is limited. The disruption of hormonal regulatory pathways is of  
527 course the hallmark of these harmful chemicals, but additional toxicological modes of action may also be  
528 involved in sub-lethal effects (Flint et al., 2012). In addition to classical endpoints related to reproduction and  
529 development, more and more studies pinpoint additional disturbed processes and pathways in invertebrates  
530 exposed to EDCs, including immune deficiencies that may hamper the ability of individuals to fight their natural  
531 pathogens and parasites (Canesi et al., 2007; Gagné et al., 2008; Lu et al., 2013) and behavioral changes that

532 may negatively affect food finding, habitat choice or sexual partner selection (Baglan et al., 2018; Clotfelter et  
533 al., 2004; Lam et al., 2010; McCallum et al., 2013), with possible transgenerational effects (Giraud et al., 2017b;  
534 Li et al., 2018; Oliveira-Filho et al., 2009). Epigenetic modifications are notably evoked as possible mechanism  
535 entailed (Novo et al., 2018; Schwindt, 2015) but, as in vertebrates, the identification of the precise mechanisms  
536 involved, as well as their implication in invertebrates require deeper investigations. Among future challenges, it  
537 is necessary to accurately identify all involved mechanisms, in order to better characterize organisms as well as  
538 populations response to multiple environmental stressors.  
539

540

#### 541 **Acknowledgments**

542 The authors thank two anonymous referees for their constructive criticism on the manuscript. The first author  
543 gratefully acknowledges the delegation program of CNRS.  
544

545

546

547

#### 548 **Bibliography**

- 549 Abe, R., Toyota, K., Miyakawa, H., Watanabe, H., Oka, T., Miyagawa, S., Nishide, H., Uchiyama, I., Tollefsen, K.E.,  
550 Iguchi, T., Tatarazako, N., 2015a. Diofenolan induces male offspring production through binding to the  
551 juvenile hormone receptor in *Daphnia magna*. *Aquat. Toxicol.* 159, 44–51.  
552 <https://doi.org/10.1016/j.aquatox.2014.11.015>
- 553 Abe, R., Watanabe, H., Yamamuro, M., Iguchi, T., Tatarazako, N., 2015b. Establishment of a short-term, *in vivo*  
554 screening method for detecting chemicals with juvenile hormone activity using adult *Daphnia magna*: In  
555 vivo screening method for detecting juvenile hormone activity. *J. Appl. Toxicol.* 35, 75–82.  
556 <https://doi.org/10.1002/jat.2989>
- 557 Amiard, J.-C., Amiard-Triquet, C., 2015. Ecotoxicological Risk of Endocrine Disruptors, in: *Aquatic Ecotoxicology*.  
558 Elsevier, pp. 355–382. <https://doi.org/10.1016/B978-0-12-800949-9.00015-2>
- 559 An, L., Zheng, B., Liu, R., Fan, Q., Wang, Q., Luo, Y., 2014. Transcriptomic response to estrogen exposure in the  
560 male Zhikong scallop, *Chlamys farreri*. *Mar. Pollut. Bull.* 89, 59–66.  
561 <https://doi.org/10.1016/j.marpolbul.2014.10.035>
- 562 André, A., Ruivo, R., Capitão, A., Froufe, E., Páscoa, I., Costa Castro, L.F., Santos, M.M., 2017. Cloning and  
563 functional characterization of a retinoid X receptor orthologue in *Platynereis dumerilii* : An evolutionary  
564 and toxicological perspective. *Chemosphere* 182, 753–761.  
565 <https://doi.org/10.1016/j.chemosphere.2017.05.064>
- 566 André, A., Ruivo, R., Gesto, M., Castro, L.F.C., Santos, M.M., 2014. Retinoid metabolism in invertebrates: When  
567 evolution meets endocrine disruption. *Gen. Comp. Endocrinol.* 208, 134–145.  
568 <https://doi.org/10.1016/j.ygcen.2014.08.005>
- 569 Annamalai, J., Namasivayam, V., 2015. Endocrine disrupting chemicals in the atmosphere: Their effects on  
570 humans and wildlife. *Environ. Int.* 76, 78–97. <https://doi.org/10.1016/j.envint.2014.12.006>
- 571 Arambourou, H., Llorente, L., Moreno-Ocio, I., Herrero, Ó., Barata, C., Fuertes, I., Delorme, N., Méndez-  
572 Fernández, L., Planelló, R., 2020. Exposure to heavy metal-contaminated sediments disrupts gene  
573 expression, lipid profile, and life history traits in the midge *Chironomus riparius*. *Water Res.* 168, 115165.  
574 <https://doi.org/10.1016/j.watres.2019.115165>
- 575 Arambourou, H., Planelló, R., Llorente, L., Fuertes, I., Barata, C., Delorme, N., Noury, P., Herrero, Ó., Villeneuve,  
576 A., Bonnineau, C., 2019. *Chironomus riparius* exposure to field-collected contaminated sediments: From  
577 subcellular effect to whole-organism response. *Sci. Total Environ.* 671, 874–882.  
578 <https://doi.org/10.1016/j.scitotenv.2019.03.384>

579 Aris, A.Z., Shamsuddin, A.S., Praveena, S.M., 2014. Occurrence of 17 $\alpha$ -ethynylestradiol (EE2) in the environment  
580 and effect on exposed biota: a review. *Environ. Int.* 69, 104–119.  
581 <https://doi.org/10.1016/j.envint.2014.04.011>

582 Aviles, A., Boulogne, I., Durand, N., Maria, A., Cordeiro, A., Bozzolan, F., Goutte, A., Alliot, F., Dacher, M.,  
583 Renault, D., Maibeche, M., Siaussat, D., 2019. Effects of DEHP on post-embryonic development, nuclear  
584 receptor expression, metabolite and ecdysteroid concentrations of the moth *Spodoptera littoralis*.  
585 *Chemosphere* 215, 725–738. <https://doi.org/10.1016/j.chemosphere.2018.10.102>

586 Baglan, H., Lazzari, C.R., Guerrieri, F.J., 2018. Glyphosate impairs learning in *Aedes aegypti* mosquito larvae at  
587 field-realistic doses. *J. Exp. Biol.* 221, jeb187518. <https://doi.org/10.1242/jeb.187518>

588 Baines, D., Wilton, E., Pawluk, A., de Gorter, M., Chomistek, N., 2017. Neonicotinoids act like endocrine  
589 disrupting chemicals in newly-emerged bees and winter bees. *Sci. Rep.* 7.  
590 <https://doi.org/10.1038/s41598-017-10489-6>

591 Baker, M.E., 2008. Trichoplax, the simplest known animal, contains an estrogen-related receptor but no  
592 estrogen receptor: Implications for estrogen receptor evolution. *Biochem. Biophys. Res. Commun.* 375,  
593 623–627. <https://doi.org/10.1016/j.bbrc.2008.08.047>

594 Baker, M.E., Lathé, R., 2018. The promiscuous estrogen receptor: Evolution of physiological estrogens and  
595 response to phytochemicals and endocrine disruptors. *J. Steroid Biochem. Mol. Biol.* 184, 29–37.  
596 <https://doi.org/10.1016/j.jsbmb.2018.07.001>

597 Bannister, R., Beresford, N., Granger, D.W., Pounds, N.A., Rand-Weaver, M., White, R., Jobling, S., Routledge,  
598 E.J., 2013. No substantial changes in estrogen receptor and estrogen-related receptor orthologue gene  
599 transcription in *Marisa cornuarietis* exposed to estrogenic chemicals. *Aquat. Toxicol.* 140–141, 19–26.  
600 <https://doi.org/10.1016/j.aquatox.2013.05.002>

601 Barata, C., Porte, C., Baird, D.J., 2004. Experimental designs to assess endocrine disrupting effects in  
602 invertebrates. A review. *Ecotoxicol. Lond. Engl.* 13, 511–517.

603 Blalock, B.J., Robinson, W.E., Loguinov, A., Vulpe, C.D., Krick, K.S., Poynton, H.C., 2018. Transcriptomic and  
604 Network Analyses Reveal Mechanistic-Based Biomarkers of Endocrine Disruption in the Marine Mussel,  
605 *Mytilus edulis*. *Environ. Sci. Technol.* 52, 9419–9430. <https://doi.org/10.1021/acs.est.8b01604>

606 Böhme, F., Bischoff, G., Zebitz, C.P., Rosenkranz, P., Wallner, K., 2017. Chronic exposure of honeybees, *Apis*  
607 *mellifera* (Hymenoptera: Apidae), to a pesticide mixture in realistic field exposure rates. *Apidologie* 48,  
608 353–363. <https://doi.org/10.1007/s13592-016-0479-x>

609 Böhme, F., Bischoff, G., Zebitz, C.P.W., Rosenkranz, P., Wallner, K., 2018. Pesticide residue survey of pollen loads  
610 collected by honeybees (*Apis mellifera*) in daily intervals at three agricultural sites in South Germany.  
611 *PLOS ONE* 13, e0199995. <https://doi.org/10.1371/journal.pone.0199995>

612 Boulangé-Lecomte, C., Xuereb, B., Trémolet, G., Duflot, A., Giusti, N., Olivier, S., Legrand, E., Forget-Leray, J.,  
613 2017. Controversial use of vitellogenin as a biomarker of endocrine disruption in crustaceans: New  
614 adverse pieces of evidence in the copepod *Eurytemora affinis*. *Comp. Biochem. Physiol. Part C Toxicol.*  
615 *Pharmacol.* 201, 66–75. <https://doi.org/10.1016/j.cbpc.2017.09.011>

616 Bovier, T.F., Rossi, S., Mita, D.G., Digilio, F.A., 2018. Effects of the synthetic estrogen 17- $\alpha$ -ethynylestradiol on  
617 *Drosophila melanogaster* : Dose and gender dependence. *Ecotoxicol. Environ. Saf.* 162, 625–632.  
618 <https://doi.org/10.1016/j.ecoenv.2018.07.020>

619 Bridgham, J.T., Keay, J., Ortlund, E.A., Thornton, J.W., 2014. Vestigialization of an Allosteric Switch: Genetic and  
620 Structural Mechanisms for the Evolution of Constitutive Activity in a Steroid Hormone Receptor. *PLoS*  
621 *Genet.* 10, e1004058. <https://doi.org/10.1371/journal.pgen.1004058>

622 Bryan, G.W., Gibbs, P.E., Hummerstone, L.G., Burt, G.R., 1986. The Decline of the Gastropod *Nucella Lapillus*  
623 Around South-West England: Evidence for the Effect of Tributyltin from Antifouling Paints. *J. Mar. Biol.*  
624 *Assoc. U. K.* 66, 611. <https://doi.org/10.1017/S0025315400042247>

625 Calatayud-Vernich, P., Calatayud, F., Simó, E., Picó, Y., 2018. Pesticide residues in honey bees, pollen and  
626 beeswax: Assessing beehive exposure. *Environ. Pollut.* 241, 106–114.  
627 <https://doi.org/10.1016/j.envpol.2018.05.062>

628 Canesi, L., Fabbri, E., 2015. Environmental Effects of BPA: Focus on Aquatic Species. *Dose-Response* 13,  
629 155932581559830. <https://doi.org/10.1177/1559325815598304>

630 Canesi, L., Lorusso, L.C., Ciacci, C., Betti, M., Rocchi, M., Pojana, G., Marcomini, A., 2007. Immunomodulation of  
631 Mytilus hemocytes by individual estrogenic chemicals and environmentally relevant mixtures of  
632 estrogens: In vitro and in vivo studies. *Aquat. Toxicol.* 81, 36–44.  
633 <https://doi.org/10.1016/j.aquatox.2006.10.010>

634 Cappello, T., Fernandes, D., Maisano, M., Casano, A., Bonastre, M., Bebianno, M.J., Mauceri, A., Fasulo, S., Porte,  
635 C., 2017. Sex steroids and metabolic responses in mussels *Mytilus galloprovincialis* exposed to  
636 drosiprenone. *Ecotoxicol. Environ. Saf.* 143, 166–172. <https://doi.org/10.1016/j.ecoenv.2017.05.031>

637 Castro, L.F.C., Santos, M.M., 2014. “To Bind or Not To Bind” : The Taxonomic Scope of Nuclear Receptor  
638 Mediated Endocrine Disruption in Invertebrate Phyla. *Environ. Sci. Technol.* 48, 5361–5363.  
639 <https://doi.org/10.1021/es501697b>

640 Chakrabarti, P., Rana, S., Bandopadhyay, S., Naik, D.G., Sarkar, S., Basu, P., 2015. Field populations of native  
641 Indian honey bees from pesticide intensive agricultural landscape show signs of impaired olfaction. *Sci.*  
642 *Rep.* 5. <https://doi.org/10.1038/srep12504>

643 Ciocan, C.M., Cubero-Leon, E., Peck, M.R., Langston, W.J., Pope, N., Minier, C., Rotchell, J.M., 2012. Intersex in  
644 *Scrobicularia plana* : Transcriptomic Analysis Reveals Novel Genes Involved in Endocrine Disruption.  
645 *Environ. Sci. Technol.* 46, 12936–12942. <https://doi.org/10.1021/es302948g>

646 Clotfelter, E.D., Bell, A.M., Levering, K.R., 2004. The role of animal behaviour in the study of endocrine-disrupting  
647 chemicals. *Anim. Behav.* 68, 665–676. <https://doi.org/10.1016/j.anbehav.2004.05.004>

648 Colin, N., Porte, C., Fernandes, D., Barata, C., Padrós, F., Carrassón, M., Monroy, M., Cano-Rocabayera, O., de  
649 Sostoa, A., Piña, B., Maceda-Veiga, A., 2016. Ecological relevance of biomarkers in monitoring studies of  
650 macro-invertebrates and fish in Mediterranean rivers. *Sci. Total Environ.* 540, 307–323.  
651 <https://doi.org/10.1016/j.scitotenv.2015.06.099>

652 Cuvillier-Hot, V., Salin, K., Devers, S., Tasiemski, A., Schaffner, P., Boulay, R., Billiard, S., Lenoir, A., 2014. Impact  
653 of ecological doses of the most widespread phthalate on a terrestrial species, the ant *Lasius niger*.  
654 *Environ. Res.* 131, 104–110. <https://doi.org/10.1016/j.envres.2014.03.016>

655 DeFur, P.L. (Ed.), 1999. Endocrine disruption in invertebrates: endocrinology, testing, and assessment. SETAC  
656 Press, Pensacola, FL.

657 Dinan, L., Bourne, P., Whiting, P., Dhadialla, T.S., Hutchinson, T.H., 2001. Screening of environmental  
658 contaminants for ecdysteroid agonist and antagonist activity using the *Drosophila melanogaster* BII cell  
659 in vitro assay. *Environ. Toxicol. Chem.* 20, 2038–2046.

660 Dodgen, L.K., Li, J., Parker, D., Gan, J.J., 2013. Uptake and accumulation of four PPCP/EDCs in two leafy  
661 vegetables. *Environ. Pollut.* 182, 150–156. <https://doi.org/10.1016/j.envpol.2013.06.038>

662 Dris, R., Gasperi, J., Saad, M., Mirande, C., Tassin, B., 2016. Synthetic fibers in atmospheric fallout: A source of  
663 microplastics in the environment? *Mar. Pollut. Bull.* 104, 290–293.  
664 <https://doi.org/10.1016/j.marpolbul.2016.01.006>

665 Engel, A., Buhrke, T., Imber, F., Jessel, S., Seidel, A., Völkel, W., Lampen, A., 2017. Agonistic and antagonistic  
666 effects of phthalates and their urinary metabolites on the steroid hormone receptors ER $\alpha$ , ER $\beta$ , and AR.  
667 *Toxicol. Lett.* 277, 54–63. <https://doi.org/10.1016/j.toxlet.2017.05.028>

668 Flint, S., Markle, T., Thompson, S., Wallace, E., 2012. Bisphenol A exposure, effects, and policy: A wildlife  
669 perspective. *J. Environ. Manage.* 104, 19–34. <https://doi.org/10.1016/j.jenvman.2012.03.021>

670 Fourier, J., Deschamps, M., Droin, L., Alaux, C., Fortini, D., Beslay, D., Le Conte, Y., Devillers, J., Aupinel, P.,  
671 Decourtye, A., 2015. Larval Exposure to the Juvenile Hormone Analog Pyriproxyfen Disrupts Acceptance  
672 of and Social Behavior Performance in Adult Honeybees. *PLOS ONE* 10, e0132985.  
673 <https://doi.org/10.1371/journal.pone.0132985>

674 Gaertner, K., Chandler, G.T., Quattro, J., Ferguson, P.L., Sabo-Attwood, T., 2012. Identification and expression of  
675 the ecdysone receptor in the harpacticoid copepod, *Amphiascus tenuiremis*, in response to fipronil.  
676 *Ecotoxicol. Environ. Saf.* 76, 39–45. <https://doi.org/10.1016/j.ecoenv.2011.09.008>

677 Gagné, F., André, C., Cejka, P., Hausler, R., Fournier, M., Blaise, C., 2008. Immunotoxic effects on freshwater  
678 mussels of a primary-treated wastewater before and after ozonation: A pilot plant study. *Ecotoxicol.*  
679 *Environ. Saf.* 69, 366–373. <https://doi.org/10.1016/j.ecoenv.2007.10.027>

680 Garreta-Lara, E., Checa, A., Fuchs, D., Tauler, R., Lacorte, S., Wheelock, C.E., Barata, C., 2018. Effect of psychiatric  
681 drugs on *Daphnia magna* oxylipin profiles. *Sci. Total Environ.* 644, 1101–1109.  
682 <https://doi.org/10.1016/j.scitotenv.2018.06.333>

683 Giraudo, M., Douville, M., Cottin, G., Houde, M., 2017a. Transcriptomic, cellular and life-history responses of  
684 *Daphnia magna* chronically exposed to benzotriazoles: Endocrine-disrupting potential and molting  
685 effects. *PLOS ONE* 12, e0171763. <https://doi.org/10.1371/journal.pone.0171763>

686 Giraudo, M., Douville, M., Houde, M., 2015. Chronic toxicity evaluation of the flame retardant tris (2-  
687 butoxyethyl) phosphate (TBOEP) using *Daphnia magna* transcriptomic response. *Chemosphere* 132,  
688 159–165. <https://doi.org/10.1016/j.chemosphere.2015.03.028>

689 Giraudo, M., Dubé, M., Lépine, M., Gagnon, P., Douville, M., Houde, M., 2017b. Multigenerational effects  
690 evaluation of the flame retardant tris(2-butoxyethyl) phosphate (TBOEP) using *Daphnia magna*. *Aquat.*  
691 *Toxicol.* 190, 142–149. <https://doi.org/10.1016/j.aquatox.2017.07.003>

692 Gismondi, E., 2018. Identification of molt-inhibiting hormone and ecdysteroid receptor cDNA sequences in  
693 *Gammarus pulex*, and variations after endocrine disruptor exposures. *Ecotoxicol. Environ. Saf.* 158, 9–  
694 17. <https://doi.org/10.1016/j.ecoenv.2018.04.017>

695 Goldstone, J.V., Sundaramoorthy, M., Zhao, B., Waterman, M.R., Stegeman, J.J., Lamb, D.C., 2016. Genetic and  
696 structural analyses of cytochrome P450 hydroxylases in sex hormone biosynthesis: Sequential origin and  
697 subsequent coevolution. *Mol. Phylogenet. Evol.* 94, 676–687.  
698 <https://doi.org/10.1016/j.ympev.2015.09.012>

699 Gouveia, D., Almunia, C., Cogne, Y., Pible, O., Degli-Esposti, D., Salvador, A., Cristobal, S., Sheehan, D., Chaumot,  
700 A., Geffard, O., Armengaud, J., 2019. Ecotoxicoproteomics: A decade of progress in our understanding of  
701 anthropogenic impact on the environment. *J. Proteomics* 198, 66–77.  
702 <https://doi.org/10.1016/j.jprot.2018.12.001>

703 Graceli, J.B., Sena, G.C., Lopes, P.F.I., Zamprogno, G.C., da Costa, M.B., Godoi, A.F.L., dos Santos, D.M., de  
704 Marchi, M.R.R., dos Santos Fernandez, M.A., 2013. Organotins: A review of their reproductive toxicity,  
705 biochemistry, and environmental fate. *Reprod. Toxicol.* 36, 40–52.  
706 <https://doi.org/10.1016/j.reprotox.2012.11.008>

707 Graham, L.D., Johnson, W.M., Pawlak-Skrzecz, A., Eaton, R.E., Bliese, M., Howell, L., Hannan, G.N., Hill, R.J., 2007.  
708 Ligand binding by recombinant domains from insect ecdysone receptors. *Insect Biochem. Mol. Biol.* 37,  
709 611–626. <https://doi.org/10.1016/j.ibmb.2007.03.006>

710 Grilo, T.F., Rosa, R., 2017. Intersexuality in aquatic invertebrates: Prevalence and causes. *Sci. Total Environ.* 592,  
711 714–728. <https://doi.org/10.1016/j.scitotenv.2017.02.099>

712 Guittard, E., Blais, C., Maria, A., Parvy, J.-P., Pasricha, S., Lumb, C., Lafont, R., Daborn, P.J., Dauphin-Villemant, C.,  
713 2011. CYP18A1, a key enzyme of *Drosophila* steroid hormone inactivation, is essential for  
714 metamorphosis. *Dev. Biol.* 349, 35–45. <https://doi.org/10.1016/j.ydbio.2010.09.023>

715 Gust, M., Gagné, F., Berlioz-Barbier, A., Besse, J.P., Buronfosse, T., Tournier, M., Tutundjian, R., Garric, J., Cren-  
716 Olivé, C., 2014. Caged mudsnail *Potamopyrgus antipodarum* (Gray) as an integrated field biomonitoring  
717 tool: Exposure assessment and reprotoxic effects of water column contamination. *Water Res.* 54, 222–  
718 236. <https://doi.org/10.1016/j.watres.2014.01.057>

719 Gutierrez-Mazariegos, J., Nadendla, E.K., Lima, D., Pierzchalski, K., Jones, J.W., Kane, M., Nishikawa, J.-I.,  
720 Hiromori, Y., Nakanishi, T., Santos, M.M., Castro, L.F.C., Bourguet, W., Schubert, M., Laudet, V., 2014. A  
721 Mollusk Retinoic Acid Receptor (RAR) Ortholog Sheds Light on the Evolution of Ligand Binding.  
722 *Endocrinology* 155, 4275–4286. <https://doi.org/10.1210/en.2014-1181>

723 Hakme, E., Lozano, A., Gómez-Ramos, M.M., Hernando, M.D., Fernández-Alba, A.R., 2017. Non-target evaluation  
724 of contaminants in honey bees and pollen samples by gas chromatography time-of-flight mass  
725 spectrometry. *Chemosphere* 184, 1310–1319. <https://doi.org/10.1016/j.chemosphere.2017.06.089>



726 Hamilton, K.J., Hewitt, S.C., Arao, Y., Korach, K.S., 2017. Estrogen Hormone Biology, in: Current Topics in  
727 Developmental Biology. Elsevier, pp. 109–146. <https://doi.org/10.1016/bs.ctdb.2016.12.005>

728 Handberg-Thorsager, M., Gutierrez-Mazariegos, J., Arold, S.T., Kumar Nadendla, E., Bertucci, P.Y., Germain, P.,  
729 Tomançak, P., Pierzchalski, K., Jones, J.W., Albalat, R., Kane, M.A., Bourguet, W., Laudet, V., Arendt, D.,  
730 Schubert, M., 2018. The ancestral retinoic acid receptor was a low-affinity sensor triggering neuronal  
731 differentiation. *Sci. Adv.* 4, eaao1261. <https://doi.org/10.1126/sciadv.aao1261>

732 Herrero, Ó., Aquilino, M., Sánchez-Argüello, P., Planelló, R., 2018. The BPA-substitute bisphenol S alters the  
733 transcription of genes related to endocrine, stress response and biotransformation pathways in the  
734 aquatic midge *Chironomus riparius* (Diptera, Chironomidae). *PLOS ONE* 13, e0193387.  
735 <https://doi.org/10.1371/journal.pone.0193387>

736 Herrero, Ó., Morcillo, G., Planelló, R., 2017. Transcriptional deregulation of genetic biomarkers in *Chironomus*  
737 *riparius* larvae exposed to ecologically relevant concentrations of di(2-ethylhexyl) phthalate (DEHP).  
738 *PLOS ONE* 12, e0171719. <https://doi.org/10.1371/journal.pone.0171719>

739 Herrero, Ó., Planelló, R., Morcillo, G., 2015. The plasticizer benzyl butyl phthalate (BBP) alters the ecdysone  
740 hormone pathway, the cellular response to stress, the energy metabolism, and several detoxication  
741 mechanisms in *Chironomus riparius* larvae. *Chemosphere* 128, 266–277.  
742 <https://doi.org/10.1016/j.chemosphere.2015.01.059>

743 Heyland, A., Price, D.A., Bodnarova-buganova, M., Moroz, L.L., 2006. Thyroid hormone metabolism and  
744 peroxidase function in two non-chordate animals. *J. Exp. Zool. B Mol. Dev. Evol.* 306B, 551–566.  
745 <https://doi.org/10.1002/jez.b.21113>

746 Holzer, G., Markov, G.V., Laudet, V., 2017a. Evolution of Nuclear Receptors and Ligand Signaling, in: Current  
747 Topics in Developmental Biology. Elsevier, pp. 1–38. <https://doi.org/10.1016/bs.ctdb.2017.02.003>

748 Holzer, G., Roux, N., Laudet, V., 2017b. Evolution of ligands, receptors and metabolizing enzymes of thyroid  
749 signaling. *Mol. Cell. Endocrinol.* 459, 5–13. <https://doi.org/10.1016/j.mce.2017.03.021>

750 Houde, M., Douville, M., Giraudo, M., Jean, K., Lépine, M., Spencer, C., De Silva, A.O., 2016. Endocrine-disruption  
751 potential of perfluoroethylcyclohexane sulfonate (PFECHS) in chronically exposed *Daphnia magna*.  
752 *Environ. Pollut.* 218, 950–956. <https://doi.org/10.1016/j.envpol.2016.08.043>

753 Huang, W., Xu, F., Qu, T., Zhang, R., Li, L., Que, H., Zhang, G., 2015. Identification of Thyroid Hormones and  
754 Functional Characterization of Thyroid Hormone Receptor in the Pacific Oyster *Crassostrea gigas* Provide  
755 Insight into Evolution of the Thyroid Hormone System. *PLOS ONE* 10, e0144991.  
756 <https://doi.org/10.1371/journal.pone.0144991>

757 Huang, Y., Li, W., Qin, L., Xie, X., Gao, B., Sun, J., Li, A., 2019. Distribution of endocrine-disrupting chemicals in  
758 colloidal and soluble phases in municipal secondary effluents and their removal by different advanced  
759 treatment processes. *Chemosphere* 219, 730–739. <https://doi.org/10.1016/j.chemosphere.2018.11.201>

760 Hubbard, W.J., Bland, K.I., Chaudry, I.H., 2015. The ERRor of Our Ways: Estrogen-Related Receptors are About  
761 Energy, Not Hormones, and are Potential New Targets for Trauma and Shock. *Shock* 44, 3–15.  
762 <https://doi.org/10.1097/SHK.0000000000000364>

763 Hultin, C.L., Hallgren, P., Persson, A., Hansson, M.C., 2014. Identification of an estrogen receptor gene in the  
764 natural freshwater snail *Bithynia tentaculata*. *Gene* 540, 26–31.  
765 <https://doi.org/10.1016/j.gene.2014.02.039>

766 Hutchinson, T.H., 2007. Small is useful in endocrine disrupter assessment—four key recommendations for  
767 aquatic invertebrate research. *Ecotoxicology* 16, 231–238. <https://doi.org/10.1007/s10646-006-0107-z>

768 Hutchinson, T.H., 2002. Reproductive and developmental effects of endocrine disrupters in invertebrates: in  
769 vitro and in vivo approaches. *Toxicol. Lett.* 131, 75–81.

770 Iguchi, T., Katsu, Y., 2008. Commonality in Signaling of Endocrine Disruption from Snail to Human. *BioScience* 58,  
771 1061–1067. <https://doi.org/10.1641/B581109>

772 Janer, G., Porte, C., 2007. Sex steroids and potential mechanisms of non-genomic endocrine disruption in  
773 invertebrates. *Ecotoxicology* 16, 145–160. <https://doi.org/10.1007/s10646-006-0110-4>

774 Jin, S., Yang, F., Liao, T., Hui, Y., Wen, S., Xu, Y., 2012. Enhanced effects by mixtures of three estrogenic  
775 compounds at environmentally relevant levels on development of Chinese rare minnow (*Gobiocypris*  
776 *rarus*). *Environ. Toxicol. Pharmacol.* 33, 277–283. <https://doi.org/10.1016/j.etap.2011.12.016>

777 Jo, M., Lee, S., Yoon, S., Kim, W.-K., 2018. Developmental and reproductive effects of tamoxifen on *Daphnia*  
778 *magna*. *Environ. Monit. Assess.* 190. <https://doi.org/10.1007/s10661-018-7002-y>

779 Jobling, S., Casey, D., Rodgers-Gray, T., Oehlmann, J., Schulte-Oehlmann, U., Pawlowski, S., Baunbeck, T., Turner,  
780 A.P., Tyler, C.R., 2004. Comparative responses of molluscs and fish to environmental estrogens and an  
781 estrogenic effluent. *Aquat. Toxicol.* 66, 207–222. <https://doi.org/10.1016/j.aquatox.2004.01.002>

782 Johnson, R.M., Ellis, M.D., Mullin, C.A., Frazier, M., 2010. Pesticides and honey bee toxicity – USA. *Apidologie* 41,  
783 312–331. <https://doi.org/10.1051/apido/2010018>

784 Jones, B.L., Walker, C., Azizi, B., Tolbert, L., Williams, L.D., Snell, T.W., 2017. Conservation of estrogen receptor  
785 function in invertebrate reproduction. *BMC Evol. Biol.* 17. <https://doi.org/10.1186/s12862-017-0909-z>

786 Keay, J., Bridgham, J.T., Thornton, J.W., 2006. The Octopus vulgaris Estrogen Receptor Is a Constitutive  
787 Transcriptional Activator: Evolutionary and Functional Implications. *Endocrinology* 147, 3861–3869.  
788 <https://doi.org/10.1210/en.2006-0363>

789 Keay, J., Thornton, J.W., 2009. Hormone-Activated Estrogen Receptors in Annelid Invertebrates: Implications for  
790 Evolution and Endocrine Disruption. *Endocrinology* 150, 1731–1738. <https://doi.org/10.1210/en.2008-1338>

791

792 Kontogiannatos, D., Swevers, L., Zakasis, G., Kourti, A., 2015. The molecular and physiological impact of  
793 bisphenol A in *Sesamia nonagrioides* (Lepidoptera: Noctuidae). *Ecotoxicology* 24, 356–367.  
794 <https://doi.org/10.1007/s10646-014-1384-6>

795 Kovacevic, V., Simpson, A.J., Simpson, M.J., 2019. The concentration of dissolved organic matter impacts the  
796 metabolic response in *Daphnia magna* exposed to 17 $\alpha$ -ethynylestradiol and perfluorooctane sulfonate.  
797 *Ecotoxicol. Environ. Saf.* 170, 468–478. <https://doi.org/10.1016/j.ecoenv.2018.12.008>

798 Lafont, R., Mathieu, M., 2007. Steroids in aquatic invertebrates. *Ecotoxicology* 16, 109–130.  
799 <https://doi.org/10.1007/s10646-006-0113-1>

800 Lagadic, L., Coutellec, M.-A., Caquet, T., 2007. Endocrine disruption in aquatic pulmonate molluscs: few  
801 evidences, many challenges. *Ecotoxicology* 16, 45–59. <https://doi.org/10.1007/s10646-006-0114-0>

802 Laguerre, M., Veenstra, J.A., 2010. Ecdysone receptor homologs from mollusks, leeches and a polychaete worm.  
803 *FEBS Lett.* 584, 4458–4462. <https://doi.org/10.1016/j.febslet.2010.10.004>

804 Lam, C., Neumann, R., Shin, P.K.S., Au, D.W.T., Qian, P.Y., Wu, R.S.S., 2010. Polybrominated Diphenylethers  
805 (PBDEs) Alter Larval Settlement of Marine Benthic Polychaetes. *Environ. Sci. Technol.* 44, 7130–7137.  
806 <https://doi.org/10.1021/es1012615>

807 Langston, W.J., Pope, N.D., Davey, M., Langston, K.M., O’ Hara, S.C.M., Gibbs, P.E., Pascoe, P.L., 2015. Recovery  
808 from TBT pollution in English Channel environments: A problem solved? *Mar. Pollut. Bull.* 95, 551–564.  
809 <https://doi.org/10.1016/j.marpolbul.2014.12.011>

810 Laranjeiro, F., Sánchez-Marín, P., Oliveira, I.B., Galante-Oliveira, S., Barroso, C., 2018. Fifteen years of imposex  
811 and tributyltin pollution monitoring along the Portuguese coast. *Environ. Pollut.* 232, 411–421.  
812 <https://doi.org/10.1016/j.envpol.2017.09.056>

813 Lawler, S.P., 2017. Environmental safety review of methoprene and bacterially-derived pesticides commonly  
814 used for sustained mosquito control. *Ecotoxicol. Environ. Saf.* 139, 335–343.  
815 <https://doi.org/10.1016/j.ecoenv.2016.12.038>

816 le Maire, A., Grimaldi, M., Roecklin, D., Dagnino, S., Vivat-Hannah, V., Balaguer, P., Bourguet, W., 2009.  
817 Activation of RXR–PPAR heterodimers by organotin environmental endocrine disruptors. *EMBO Rep.* 10,  
818 367–373. <https://doi.org/10.1038/embor.2009.8>

819 LeBlanc, G.A., 2007. Crustacean endocrine toxicology: a review. *Ecotoxicology* 16, 61–81.  
820 <https://doi.org/10.1007/s10646-006-0115-z>

821 Lee, S.-B., Choi, J., 2006. Multilevel evaluation of nonylphenol toxicity in fourth-instar larvae of *Chironomus*  
822 *riparius* (Diptera, Chironomidae). *Environ. Toxicol. Chem.* 25, 3006–3014.

823 Lenoir, A., Boulay, R., Dejean, A., Touchard, A., Cuvillier-Hot, V., 2016. Phthalate pollution in an Amazonian  
824 rainforest. *Environ. Sci. Pollut. Res.* 23, 16865–16872. <https://doi.org/10.1007/s11356-016-7141-z>

825 Lenoir, A., Cuvillier-Hot, V., Devers, S., Christidès, J.-P., Montigny, F., 2012. Ant cuticles: A trap for atmospheric  
826 phthalate contaminants. *Sci. Total Environ.* 441, 209–212.  
827 <https://doi.org/10.1016/j.scitotenv.2012.10.003>

828 Lenoir, A., Touchard, A., Devers, S., Christidès, J.-P., Boulay, R., Cuvillier-Hot, V., 2014. Ant cuticular response to  
829 phthalate pollution. *Environ. Sci. Pollut. Res.* 21, 13446–13451. <https://doi.org/10.1007/s11356-014-3272-2>

830

831 Leonard, J.A., Cope, W.G., Hammer, E.J., Barnhart, M.C., Bringolf, R.B., 2017. Extending the toxicity-testing  
832 paradigm for freshwater mussels: Assessing chronic reproductive effects of the synthetic estrogen 17 $\alpha$ -  
833 ethinylestradiol on the unionid mussel *Elliptio complanata*. *Comp. Biochem. Physiol. Part C Toxicol.*  
834 *Pharmacol.* 191, 14–25. <https://doi.org/10.1016/j.cbpc.2016.09.002>

835 Li, S.-W., How, C.M., Liao, V.H.-C., 2018. Prolonged exposure of di(2-ethylhexyl) phthalate induces  
836 multigenerational toxic effects in *Caenorhabditis elegans*. *Sci. Total Environ.* 634, 260–266.  
837 <https://doi.org/10.1016/j.scitotenv.2018.03.355>

838 Liu, T., Li, Y., Zhao, X., Zhang, M., Gu, W., 2014. Ethylparaben affects lifespan, fecundity, and the expression  
839 levels of ERR, EcR and YPR in *Drosophila melanogaster*. *J. Insect Physiol.* 71, 1–7.  
840 <https://doi.org/10.1016/j.jinsphys.2014.09.008>

841 Liu, Y., Tam, N.F.Y., Guan, Y., Gao, B., 2012. Influence of a Marine Diatom on the Embryonic Toxicity of 17 $\alpha$ -  
842 Ethinylestradiol to the Abalone *Haliotis diversicolor supertexta*. *Water. Air. Soil Pollut.* 223, 4383–4395.  
843 <https://doi.org/10.1007/s11270-012-1202-9>

844 Lu, Y., Zhang, P., Li, C., Su, X., Jin, C., Li, Y., Xu, Y., Li, T., 2013. Characterisation of immune-related gene  
845 expression in clam (*Venerupis philippinarum*) under exposure to di(2-ethylhexyl) phthalate. *Fish Shellfish*  
846 *Immunol.* 34, 142–146. <https://doi.org/10.1016/j.fsi.2012.10.015>

847 Lv, L., Dong, X., Lv, F., Zhao, W., Yu, Y., Yang, W., 2017. Molecular cloning and characterization of an estrogen  
848 receptor gene in the marine polychaete *Perinereis aibuhitensis*. *Comp. Biochem. Physiol. B Biochem.*  
849 *Mol. Biol.* 207, 15–21. <https://doi.org/10.1016/j.cbpb.2017.02.001>

850 Lyons, G., 2006. Viewpoint: Policy Requirements for Protecting Wildlife from Endocrine Disruptors. *Environ.*  
851 *Health Perspect.* 114, 142–146. <https://doi.org/10.1289/ehp.8070>

852 Mankidy, R., Wiseman, S., Ma, H., Giesy, J.P., 2013. Biological impact of phthalates. *Toxicol. Lett.* 217, 50–58.  
853 <https://doi.org/10.1016/j.toxlet.2012.11.025>

854 Maria, A., Malbert-Colas, A., Boulogne, I., Braman, V., Boitard, C., Dacher, M., Chertemps, T., Maibeche, M.,  
855 Blais, C., Siaussat, D., 2019. Effects of bisphenol A on post-embryonic development of the cotton pest  
856 *Spodoptera littoralis*. *Chemosphere* 235, 616–625. <https://doi.org/10.1016/j.chemosphere.2019.06.073>

857 Markov, G.V., Tavares, R., Dauphin-Villemant, C., Demeneix, B.A., Baker, M.E., Laudet, V., 2009. Independent  
858 elaboration of steroid hormone signaling pathways in metazoans. *Proc. Natl. Acad. Sci.* 106, 11913–  
859 11918. <https://doi.org/10.1073/pnas.0812138106>

860 Martínez-Paz, P., Morales, M., Urien, J., Morcillo, G., Martínez-Guitarte, J.L., 2017. Endocrine-related genes are  
861 altered by antibacterial agent triclosan in *Chironomus riparius* aquatic larvae. *Ecotoxicol. Environ. Saf.*  
862 140, 185–190. <https://doi.org/10.1016/j.ecoenv.2017.02.047>

863 Matsumoto, T., Nakamura, A.M., Mori, K., Akiyama, I., Hirose, H., Takahashi, Y., 2007. Oyster estrogen receptor:  
864 cDNA cloning and immunolocalization. *Gen. Comp. Endocrinol.* 151, 195–201.  
865 <https://doi.org/10.1016/j.ygcen.2007.01.016>

866 McCallum, M.L., Matlock, M., Treas, J., Safi, B., Sanson, W., McCallum, J.L., 2013. Endocrine disruption of sexual  
867 selection by an estrogenic herbicide in the mealworm beetle (*Tenebrio molitor*). *Ecotoxicology* 22,  
868 1461–1466. <https://doi.org/10.1007/s10646-013-1132-3>

869 Mommaerts, V., Smaghe, G., 2011. Side-Effects of Pesticides on the Pollinator *Bombus*: An Overview, in:  
870 Stoytcheva, M. (Ed.), *Pesticides in the Modern World - Pests Control and Pesticides Exposure and*  
871 *Toxicity Assessment*. InTech. <https://doi.org/10.5772/25254>

872 Mommaerts, V., Sterk, G., Smagghe, G., 2006. Bumblebees can be used in combination with juvenile hormone  
873 analogues and ecdysone agonists. *Ecotoxicology* 15, 513–521. [https://doi.org/10.1007/s10646-006-](https://doi.org/10.1007/s10646-006-0087-z)  
874 [0087-z](https://doi.org/10.1007/s10646-006-0087-z)

875 Morales, M., Martínez-Paz, P., Sánchez-Argüello, P., Morcillo, G., Martínez-Guitarte, J.L., 2018. Bisphenol A (BPA)  
876 modulates the expression of endocrine and stress response genes in the freshwater snail *Physa acuta*.  
877 *Ecotoxicol. Environ. Saf.* 152, 132–138. <https://doi.org/10.1016/j.ecoenv.2018.01.034>

878 Mu, X., LeBlanc, G.A., 2002. Developmental toxicity of testosterone in the crustacean *Daphnia magna* involves  
879 anti-ecdysteroidal activity. *Gen. Comp. Endocrinol.* 129, 127–133.

880 Nagasawa, K., Treen, N., Kondo, R., Otoki, Y., Itoh, N., Rotchell, J.M., Osada, M., 2015. Molecular characterization  
881 of an estrogen receptor and estrogen-related receptor and their autoregulatory capabilities in two  
882 *Mytilus* species. *Gene* 564, 153–159. <https://doi.org/10.1016/j.gene.2015.03.073>

883 Nair, P.M.G., Choi, J., 2012. Modulation in the mRNA expression of ecdysone receptor gene in aquatic midge,  
884 *Chironomus riparius* upon exposure to nonylphenol and silver nanoparticles. *Environ. Toxicol.*  
885 *Pharmacol.* 33, 98–106. <https://doi.org/10.1016/j.etap.2011.09.006>

886 Novo, M., Muñoz-González, A.B., Trigo, D., Casquero, S., Martínez Guitarte, J.L., 2019. Applying sunscreens on  
887 earthworms: Molecular response of *Eisenia fetida* after direct contact with an organic UV filter. *Sci. Total*  
888 *Environ.* 676, 97–104. <https://doi.org/10.1016/j.scitotenv.2019.04.238>

889 Novo, M., Verdú, I., Trigo, D., Martínez-Guitarte, J.L., 2018. Endocrine disruptors in soil: Effects of bisphenol A on  
890 gene expression of the earthworm *Eisenia fetida*. *Ecotoxicol. Environ. Saf.* 150, 159–167.  
891 <https://doi.org/10.1016/j.ecoenv.2017.12.030>

892 Oberdörster, E., McClellan-Green, P., 2002. Mechanisms of imposex induction in the mud snail, *Ilyanassa*  
893 *obsoleta*: TBT as a neurotoxin and aromatase inhibitor. *Mar. Environ. Res.* 54, 715–718.  
894 [https://doi.org/10.1016/S0141-1136\(02\)00118-6](https://doi.org/10.1016/S0141-1136(02)00118-6)

895 Oehlmann, J., Di Benedetto, P., Tillmann, M., Duft, M., Oetken, M., Schulte-Oehlmann, U., 2007. Endocrine  
896 disruption in prosobranch molluscs: evidence and ecological relevance. *Ecotoxicology* 16, 29–43.  
897 <https://doi.org/10.1007/s10646-006-0109-x>

898 Oehlmann, J., Schulte-Oehlmann, U., Bachmann, J., Oetken, M., Lutz, I., Kloas, W., Ternes, T.A., 2006. Bisphenol  
899 A Induces Superfeminization in the Ramshorn Snail (Gastropoda: Prosobranchia) at Environmentally  
900 Relevant Concentrations. *Environ. Health Perspect.* 114, 127–133. <https://doi.org/10.1289/ehp.8065>

901 Oehlmann, J., Schulte-Oehlmann, U., Kloas, W., Jagnytsh, O., Lutz, I., Kusk, K.O., Wollenberger, L., Santos, E.M.,  
902 Paull, G.C., Van Look, K.J.W., Tyler, C.R., 2009. A critical analysis of the biological impacts of plasticizers  
903 on wildlife. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 2047–2062. <https://doi.org/10.1098/rstb.2008.0242>

904 Oetken, M., Bachmann, J., Schulte-Oehlmann, U., Oehlmann, J., 2004. Evidence for Endocrine Disruption in  
905 Invertebrates, in: *International Review of Cytology*. Elsevier, pp. 1–44. [https://doi.org/10.1016/S0074-](https://doi.org/10.1016/S0074-7696(04)36001-8)  
906 [7696\(04\)36001-8](https://doi.org/10.1016/S0074-7696(04)36001-8)

907 Oliveira, E., Barata, C., Piña, B., 2016. Endocrine Disruption in the Omics Era: New Views, New Hazards, New  
908 Approaches. *Open Biotechnol. J.* 10, 20–35. <https://doi.org/10.2174/1874070701610010020>

909 Oliveira-Filho, E.C., Grisolia, C.K., Paumgarten, F.J.R., 2009. Trans-generation study of the effects of nonylphenol  
910 ethoxylate on the reproduction of the snail *Biomphalaria tenagophila*. *Ecotoxicol. Environ. Saf.* 72, 458–  
911 465. <https://doi.org/10.1016/j.ecoenv.2007.10.008>

912 Orozco, A., Lazcano, I., Hernández-Puga, G., Olvera, A., 2017. Non-mammalian models reveal the role of  
913 alternative ligands for thyroid hormone receptors. *Mol. Cell. Endocrinol.* 459, 59–63.  
914 <https://doi.org/10.1016/j.mce.2017.03.003>

915 Ozáez, I., Martínez-Guitarte, J.L., Morcillo, G., 2014. The UV filter benzophenone 3 (BP-3) activates hormonal  
916 genes mimicking the action of ecdysone and alters embryo development in the insect *Chironomus*  
917 *riparius* (Diptera). *Environ. Pollut.* 192, 19–26. <https://doi.org/10.1016/j.envpol.2014.04.038>

918 Ozáez, I., Martínez-Guitarte, J.L., Morcillo, G., 2013. Effects of in vivo exposure to UV filters (4-MBC, OMC, BP-3,  
919 4-HB, OC, OD-PABA) on endocrine signaling genes in the insect *Chironomus riparius*. *Sci. Total Environ.*  
920 456–457, 120–126. <https://doi.org/10.1016/j.scitotenv.2013.03.081>

921 Park, K., Kwak, I.-S., 2010. Molecular effects of endocrine-disrupting chemicals on the *Chironomus riparius*  
922 estrogen-related receptor gene. *Chemosphere* 79, 934–941.  
923 <https://doi.org/10.1016/j.chemosphere.2010.03.002>

924 Pascoal, S., Carvalho, G., Vasieva, O., Hughes, R., Cossins, A., Fang, Y., Ashelford, K., Olohan, L., Barroso, C.,  
925 Mendo, S., Creer, S., 2013. Transcriptomics and in vivo tests reveal novel mechanisms underlying  
926 endocrine disruption in an ecological sentinel, *Nucella lapillus*. *Mol. Ecol.* 22, 1589–1608.  
927 <https://doi.org/10.1111/mec.12137>

928 Pener, M.Paul., Dhadialla, T.S., 2012. An Overview of Insect Growth Disruptors; Applied Aspects, in: *Advances in*  
929 *Insect Physiology*. Elsevier, pp. 1–162. <https://doi.org/10.1016/B978-0-12-391500-9.00001-2>

930 Piña, B., Raldúa, D., Barata, C., Portugal, J., Navarro-Martín, L., Martínez, R., Fuertes, I., Casado, M., 2018.  
931 Functional Data Analysis: Omics for Environmental Risk Assessment, in: *Comprehensive Analytical*  
932 *Chemistry*. Elsevier, pp. 583–611. <https://doi.org/10.1016/bs.coac.2018.07.007>

933 Planelló, R., Herrero, O., Martínez-Guitarte, J.L., Morcillo, G., 2011. Comparative effects of butyl benzyl phthalate  
934 (BBP) and di(2-ethylhexyl) phthalate (DEHP) on the aquatic larvae of *Chironomus riparius* based on gene  
935 expression assays related to the endocrine system, the stress response and ribosomes. *Aquat. Toxicol.*  
936 105, 62–70. <https://doi.org/10.1016/j.aquatox.2011.05.011>

937 Planelló, R., Martínez-Guitarte, J.L., Morcillo, G., 2008. The endocrine disruptor bisphenol A increases the  
938 expression of HSP70 and ecdysone receptor genes in the aquatic larvae of *Chironomus riparius*.  
939 *Chemosphere* 71, 1870–1876. <https://doi.org/10.1016/j.chemosphere.2008.01.033>

940 Prado, A., Pioz, M., Vidau, C., Requier, F., Jury, M., Crauser, D., Brunet, J.-L., Le Conte, Y., Alaux, C., 2019.  
941 Exposure to pollen-bound pesticide mixtures induces longer-lived but less efficient honey bees. *Sci. Total*  
942 *Environ.* 650, 1250–1260. <https://doi.org/10.1016/j.scitotenv.2018.09.102>

943 Préau, L., Fini, J.B., Morvan-Dubois, G., Demeneix, B., 2015. Thyroid hormone signaling during early neurogenesis  
944 and its significance as a vulnerable window for endocrine disruption. *Biochim. Biophys. Acta BBA - Gene*  
945 *Regul. Mech.* 1849, 112–121. <https://doi.org/10.1016/j.bbagr.2014.06.015>

946 Qu, Z., Bendena, W.G., Tobe, S.S., Hui, J.H.L., 2018. Juvenile hormone and sesquiterpenoids in arthropods:  
947 Biosynthesis, signaling, and role of MicroRNA. *J. Steroid Biochem. Mol. Biol.* 184, 69–76.  
948 <https://doi.org/10.1016/j.jsbmb.2018.01.013>

949 Quesada, C.R., Sadof, C.S., 2019. Field evaluation of insecticides and application timing on natural enemies of  
950 selected armored and soft scales. *Biol. Control* 133, 81–90.  
951 <https://doi.org/10.1016/j.biocontrol.2019.03.013>

952 Raingard, D., Bilbao, E., Cancio, I., Cajaraville, M.P., 2013. Retinoid X receptor (RXR), estrogen receptor (ER) and  
953 other nuclear receptors in tissues of the mussel *Mytilus galloprovincialis*: Cloning and transcription  
954 pattern. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* 165, 178–190.  
955 <https://doi.org/10.1016/j.cbpa.2013.03.001>

956 Roach, A.C., Wilson, S.P., 2009. Ecological impacts of tributyltin on estuarine communities in the Hastings River,  
957 NSW Australia. *Mar. Pollut. Bull.* 58, 1780–1786. <https://doi.org/10.1016/j.marpolbul.2009.08.021>

958 Rodríguez, E.M., Medesani, D.A., Fingerman, M., 2007. Endocrine disruption in crustaceans due to pollutants: A  
959 review. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* 146, 661–671.  
960 <https://doi.org/10.1016/j.cbpa.2006.04.030>

961 Sainath, S.B., André, A., Castro, L.F.C., Santos, M.M., 2019. The evolutionary road to invertebrate thyroid  
962 hormone signaling: Perspectives for endocrine disruption processes. *Comp. Biochem. Physiol. Part C*  
963 *Toxicol. Pharmacol.* 223, 124–138. <https://doi.org/10.1016/j.cbpc.2019.05.014>

964 Salgueiro-González, N., Turnes-Carou, I., Besada, V., Muniategui-Lorenzo, S., López-Mahía, P., Prada-Rodríguez,  
965 D., 2015. Occurrence, distribution and bioaccumulation of endocrine disrupting compounds in water,  
966 sediment and biota samples from a European river basin. *Sci. Total Environ.* 529, 121–130.  
967 <https://doi.org/10.1016/j.scitotenv.2015.05.048>

968 Sanchez-Bayo, F., Goka, K., 2014. Pesticide Residues and Bees – A Risk Assessment. *PLoS ONE* 9, e94482.  
969 <https://doi.org/10.1371/journal.pone.0094482>

- 970 Sánchez-Marín, P., Fernández-González, L.E., Mantilla-Aldana, L., Diz, A.P., Beiras, R., 2017. Shotgun Proteomics  
971 Analysis Discards Alkali Labile Phosphate as a Reliable Method To Assess Vitellogenin Levels in *Mytilus*  
972 *galloprovincialis*. *Environ. Sci. Technol.* 51, 7572–7580. <https://doi.org/10.1021/acs.est.7b01734>
- 973 Santos, M.M., Ruivo, R., Capitão, A., Fonseca, E., Castro, L.F.C., 2018. Identifying the gaps: Resources and  
974 perspectives on the use of nuclear receptor based-assays to improve hazard assessment of emerging  
975 contaminants. *J. Hazard. Mater.* 358, 508–511. <https://doi.org/10.1016/j.jhazmat.2018.04.076>
- 976 Schenk, S., Krauditsch, C., Frühauf, P., Gerner, C., Raible, F., 2016. Discovery of methylfarnesoate as the annelid  
977 brain hormone reveals an ancient role of sesquiterpenoids in reproduction. *eLife* 5.  
978 <https://doi.org/10.7554/eLife.17126>
- 979 Schwindt, A.R., 2015. Parental effects of endocrine disrupting compounds in aquatic wildlife: Is there evidence of  
980 transgenerational inheritance? *Gen. Comp. Endocrinol.* 219, 152–164.  
981 <https://doi.org/10.1016/j.ygcen.2015.01.020>
- 982 Scott, A.P., 2013. Do mollusks use vertebrate sex steroids as reproductive hormones? II. Critical review of the  
983 evidence that steroids have biological effects. *Steroids* 78, 268–281.  
984 <https://doi.org/10.1016/j.steroids.2012.11.006>
- 985 Short, S., Yang, G., Kille, P., Ford, A.T., 2014. Vitellogenin is not an appropriate biomarker of feminisation in a  
986 Crustacean. *Aquat. Toxicol.* 153, 89–97. <https://doi.org/10.1016/j.aquatox.2013.11.014>
- 987 Silva, P.V., Silva, A.R.R., Mendo, S., Loureiro, S., 2014. Toxicity of tributyltin (TBT) to terrestrial organisms and its  
988 species sensitivity distribution. *Sci. Total Environ.* 466–467, 1037–1046.  
989 <https://doi.org/10.1016/j.scitotenv.2013.08.002>
- 990 Skinner, M.K., 2014. Endocrine disruptor induction of epigenetic transgenerational inheritance of disease. *Mol.*  
991 *Cell. Endocrinol.* 398, 4–12. <https://doi.org/10.1016/j.mce.2014.07.019>
- 992 Smolarz, K., Hallmann, A., Zabrzeńska, S., Pietrasik, A., 2017. Elevated gonadal atresia as biomarker of endocrine  
993 disruptors: Field and experimental studies using *Mytilus trossulus* (L.) and 17-alpha ethinylestradiol  
994 (EE2). *Mar. Pollut. Bull.* 120, 58–67. <https://doi.org/10.1016/j.marpolbul.2017.04.007>
- 995 Soin, T., Iga, M., Swevers, L., Rougé, P., Janssen, C.R., Smagghe, G., 2009. Towards Coleoptera-specific high-  
996 throughput screening systems for compounds with ecdysone activity: development of EcR reporter  
997 assays using weevil (*Anthonomus grandis*)-derived cell lines and in silico analysis of ligand binding to A.  
998 *grandis* EcR ligand-binding pocket. *Insect Biochem. Mol. Biol.* 39, 523–534.  
999 <https://doi.org/10.1016/j.ibmb.2009.06.003>
- 1000 Soin, T., Smagghe, G., 2007. Endocrine disruption in aquatic insects: a review. *Ecotoxicology* 16, 83–93.  
1001 <https://doi.org/10.1007/s10646-006-0118-9>
- 1002 Song, Y., Villeneuve, D.L., Toyota, K., Iguchi, T., Tollefsen, K.E., 2017. Ecdysone Receptor Agonism Leading to  
1003 Lethal Molting Disruption in Arthropods: Review and Adverse Outcome Pathway Development. *Environ.*  
1004 *Sci. Technol.* 51, 4142–4157. <https://doi.org/10.1021/acs.est.7b00480>
- 1005 Sonne, C., 2010. Health effects from long-range transported contaminants in Arctic top predators: An integrated  
1006 review based on studies of polar bears and relevant model species. *Environ. Int.* 36, 461–491.  
1007 <https://doi.org/10.1016/j.envint.2010.03.002>
- 1008 Stanley, J., Preetha, G., 2016. Pesticide Toxicity to Pollinators: Exposure, Toxicity and Risk Assessment  
1009 Methodologies, in: *Pesticide Toxicity to Non-Target Organisms*. Springer Netherlands, Dordrecht, pp.  
1010 153–228. [https://doi.org/10.1007/978-94-017-7752-0\\_3](https://doi.org/10.1007/978-94-017-7752-0_3)
- 1011 Steffan-Dewenter, I., Potts, S.G., Packer, L., 2005. Pollinator diversity and crop pollination services are at risk.  
1012 *Trends Ecol. Evol.* 20, 651–652. <https://doi.org/10.1016/j.tree.2005.09.004>
- 1013 Sussarellu, R., Suquet, M., Thomas, Y., Lambert, C., Fabioux, C., Pernet, M.E.J., Le Goïc, N., Quillien, V., Mingant,  
1014 C., Epelboin, Y., Corporeau, C., Guyomarch, J., Robbens, J., Paul-Pont, I., Soudant, P., Huvet, A., 2016.  
1015 Oyster reproduction is affected by exposure to polystyrene microplastics. *Proc. Natl. Acad. Sci.* 113,  
1016 2430–2435. <https://doi.org/10.1073/pnas.1519019113>
- 1017 Swevers, L., Kravariti, L., Ciolfi, S., Xenou-Kokoletsi, M., Ragoussis, N., Smagghe, G., Nakagawa, Y., Mazomenos,  
1018 B., Iatrou, K., 2004. A cell-based high-throughput screening system for detecting ecdysteroid agonists

1019 and antagonists in plant extracts and libraries of synthetic compounds. *FASEB J.* 18, 134–136.  
1020 <https://doi.org/10.1096/fj.03-0627fje>

1021 Taylor, E., Heyland, A., 2017. Evolution of thyroid hormone signaling in animals: Non-genomic and genomic  
1022 modes of action. *Mol. Cell. Endocrinol.* 459, 14–20. <https://doi.org/10.1016/j.mce.2017.05.019>

1023 Teil, M.J., Blanchard, M., Chevreuril, M., 2006. Atmospheric fate of phthalate esters in an urban area (Paris-  
1024 France). *Sci. Total Environ.* 354, 212–223. <https://doi.org/10.1016/j.scitotenv.2004.12.083>

1025 Thornton, J.W., 2003. Resurrecting the Ancestral Steroid Receptor: Ancient Origin of Estrogen Signaling. *Science*  
1026 301, 1714–1717. <https://doi.org/10.1126/science.1086185>

1027 Tuberty, S.R., 2005. Ecdysteroid Responses of Estuarine Crustaceans Exposed Through Complete Larval  
1028 Development to Juvenile Hormone Agonist Insecticides. *Integr. Comp. Biol.* 45, 106–117.  
1029 <https://doi.org/10.1093/icb/45.1.106>

1030 Vandenberg, L.N., Colborn, T., Hayes, T.B., Heindel, J.J., Jacobs, D.R., Lee, D.-H., Shioda, T., Soto, A.M., vom Saal,  
1031 F.S., Welshons, W.V., Zoeller, R.T., Myers, J.P., 2012. Hormones and Endocrine-Disrupting Chemicals:  
1032 Low-Dose Effects and Nonmonotonic Dose Responses. *Endocr. Rev.* 33, 378–455.  
1033 <https://doi.org/10.1210/er.2011-1050>

1034 Vogt, É.L., Model, J.F.A., Vinagre, A.S., 2018. Effects of Organotins on Crustaceans: Update and Perspectives.  
1035 *Front. Endocrinol.* 9. <https://doi.org/10.3389/fendo.2018.00065>

1036 Wang, M., Jeong, C.-B., Li, Y., Lee, J.-S., 2017. Different transcriptomic responses of two marine copepods,  
1037 *Tigriopus japonicus* and *Pseudodiaptomus annandalei*, to a low dose of mercury chloride (HgCl<sub>2</sub>).  
1038 *Aquat. Toxicol.* 187, 124–131. <https://doi.org/10.1016/j.aquatox.2017.03.018>

1039 Weltje, L., Schulte-Oehlmann, U., 2007. The seven year itch—progress in research on endocrine disruption in  
1040 aquatic invertebrates since 1999. *Ecotoxicology* 16, 1–3. <https://doi.org/10.1007/s10646-006-0116-y>

1041 Wilson, E.O., 2006. *Naturalist*, 1st Island Press pbk. ed. ed. Island Press/Shearwater Books, Washington, D.C.

1042 Woodcock, B.A., Isaac, N.J.B., Bullock, J.M., Roy, D.B., Garthwaite, D.G., Crowe, A., Pywell, R.F., 2016. Impacts of  
1043 neonicotinoid use on long-term population changes in wild bees in England. *Nat. Commun.* 7.  
1044 <https://doi.org/10.1038/ncomms12459>

1045 Wright-Walters, M., Volz, C., Talbott, E., Davis, D., 2011. An updated weight of evidence approach to the aquatic  
1046 hazard assessment of Bisphenol A and the derivation a new predicted no effect concentration (Pnec)  
1047 using a non-parametric methodology. *Sci. Total Environ.* 409, 676–685.  
1048 <https://doi.org/10.1016/j.scitotenv.2010.07.092>

1049 Xin, F., Susiarjo, M., Bartolomei, M.S., 2015. Multigenerational and transgenerational effects of endocrine  
1050 disrupting chemicals: A role for altered epigenetic regulation? *Semin. Cell Dev. Biol.* 43, 66–75.  
1051 <https://doi.org/10.1016/j.semcdb.2015.05.008>

1052 Xu, Z., Liu, J., Wu, X., Huang, B., Pan, X., 2017. Nonmonotonic responses to low doses of xenoestrogens: A  
1053 review. *Environ. Res.* 155, 199–207. <https://doi.org/10.1016/j.envres.2017.02.018>

1054 Xu, Z.-X., Liu, J., Gu, L.-P., Huang, B., Pan, X.-J., 2017. Biological effects of xenoestrogens and the functional  
1055 mechanisms via genomic and nongenomic pathways. *Environ. Rev.* 25, 306–322.  
1056 <https://doi.org/10.1139/er-2016-0075>

1057 Yokota, H., Eguchi, S., Nakai, M., 2011. Development of an in vitro binding assay for ecdysone receptor of mysid  
1058 shrimp (*Americamysis bahia*). *Aquat. Toxicol.* 105, 708–716.  
1059 <https://doi.org/10.1016/j.aquatox.2011.09.008>

1060 Zhang, H., Pan, L., Zhang, L., 2012. Molecular cloning and characterization of estrogen receptor gene in the  
1061 Scallop *Chlamys farreri*: Expression profiles in response to endocrine disrupting chemicals. *Comp.*  
1062 *Biochem. Physiol. Part C Toxicol. Pharmacol.* 156, 51–57. <https://doi.org/10.1016/j.cbpc.2012.03.007>

1063 Zotti, M.J., De Geyter, E., Swevers, L., Braz, A.S.K., Scott, L.P.B., Rougé, P., Coll, J., Grutzmacher, A.D., Lenardão,  
1064 E.J., Smagghe, G., 2013. A cell-based reporter assay for screening for EcR agonist/antagonist activity of  
1065 natural ecdysteroids in Lepidoptera (Bm5) and Diptera (S2) cell cultures, followed by modeling of  
1066 ecdysteroid-EcR interactions and normal mode analysis. *Pestic. Biochem. Physiol.* 107, 309–320.  
1067 <https://doi.org/10.1016/j.pestbp.2013.09.003>


1068 Zou, E., 2005. Impacts of Xenobiotics on Crustacean Molting: The Invisible Endocrine Disruption. *Integr. Comp.*  
1069 *Biol.* 45, 33–38. <https://doi.org/10.1093/icb/45.1.33>  
1070  
1071



1072

1073 **Figure 1.** Main molecular regulatory pathways susceptible to endocrine disruption in protosomes;  
1074 some are shared with deuterostomes (dark blue background), while others are specific to invertebrates  
1075 (light blue background). Outer colored rings indicate the groups in which corresponding orthologs have  
1076 been evidenced. Hatched display points out non-functional receptors (insensitivity to ligand), and a  
1077 question mark signifies that an ortholog has been described but its functional activity needs further  
1078 confirmation. In italic font below each molecular pathway, are listed examples of EDCs strongly  
1079 suspected to interfere with the corresponding molecular pathway in protosomes (IGD, insect growth  
1080 disruptors; BPA, bisphenol A; BPS, bisphenol S; NP, nonylphenol; DEHP, di-ethylhexyl phthalate; BBP,  
1081 benzyl butyl phthalate; E2, 17 $\beta$ -estradiol; EE2, 17 $\alpha$ -ethinylestradiol; TBT, tributyltin; TPT, triphenyltin).

1082



1083

1084

1085