



HAL
open science

Liquid biopsies for omics-based analysis in sentinel mussels

France Caza, Philippine Granger Joly de Boissel, Richard Villemur, Stéphane Betoulle, Yves St-Pierre

► **To cite this version:**

France Caza, Philippine Granger Joly de Boissel, Richard Villemur, Stéphane Betoulle, Yves St-Pierre. Liquid biopsies for omics-based analysis in sentinel mussels. PLoS ONE, 2019, 14 (10), pp.e0223525. 10.1371/journal.pone.0223525 . hal-02480320

HAL Id: hal-02480320

<https://hal.science/hal-02480320>

Submitted on 19 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Liquid biopsies for omics-based analysis in sentinel mussels

France Caza¹, Philippine Granger Joly de Boissel¹, Richard Villemur¹, Stéphane Betoulle², Yves St-Pierre^{1*}**1** INRS-Institut Armand-Frappier, Laval, Québec, Canada, **2** Université Reims Champagne-Ardenne, UMR-I 02 SEBIO, Stress environnementaux et Biosurveillance des milieux aquatiques, Reims, France* yves.st-pierre@iaf.inrs.ca

OPEN ACCESS

Citation: Caza F, Joly de Boissel PG, Villemur R, Betoulle S, St-Pierre Y (2019) Liquid biopsies for omics-based analysis in sentinel mussels. PLoS ONE 14(10): e0223525. <https://doi.org/10.1371/journal.pone.0223525>

Editor: Silvana Allodi, Federal University of Rio de Janeiro, BRAZIL

Received: March 5, 2019

Accepted: September 23, 2019

Published: October 3, 2019

Copyright: © 2019 Caza et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The metagenomic data from this publication have been deposited to the NCBI Sequence Read Archive with the assigned the identifier PRJNA564645.

Funding: Supported in part by the Fondation Armand-Frappier (YSP), the French Polar Institute Paul Émile Victor (IPEV, Project N°409 IMMUNOTOXKER) (SB), and the Natural Sciences and Engineering Research Council of Canada (RGPIN-2019-06607). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Liquid biopsy of plasma is a simple and non-invasive technology that holds great promise in biomedical research. It is based on the analysis of nucleic acid-based biomarkers with predictive potential. In the present work, we have combined this concept with the FTA technology for sentinel mussels. We found that hemocytes collected from liquid biopsies can be readily fixed on FTA cards and used for long-term transcriptome analysis. We also showed that liquid biopsy is easily adaptable for metagenomic analysis of bacterial profiles of mussels. We finally provide evidence that liquid biopsies contained circulating cell-free DNA (ccfDNA) which can be used as an easily accessible genomic reservoir. Sampling of FTA-fixed circulating nucleic acids is stable at room temperature and does not necessitate a cold-chain protection. It showed comparable performance to frozen samples and is ideally adapted for sampling in remote areas, most notably in polar regions threatened by anthropogenic activities. From an ethical point of view, this minimally-invasive and non-lethal approach further reduces incidental mortality associated with conventional tissue sampling. This liquid biopsy-based approach should thus facilitate biobanking activities and development of omics-based biomarkers in mussels to assess the quality of aquatic ecosystems.

Introduction

Because of their ability to accumulate xenobiotics in their tissues, their wide distribution, and their ecological and economical importance, blue mussels (*Mytilus* spp.) have long been recognized as good biological indicators for monitoring the effects of pollution and climate change in marine habitats [1]. Not surprisingly, a relatively large number of biomarkers have been developed, often using a multi-tier approach. They include functional biomarkers based on enzymatic activities or cellular functions, or on measuring concentrations of specific stress indicators [2–4]. In most cases, measure of these biomarkers requires on-site analysis or tissue biopsy performed by trained personnel [5, 6]. It also often requires an uninterrupted cold chain to maintain sample quality. For sampling in remote areas, such as polar regions, this implies complex logistical challenges and considerable risk and cost associated with transportation and storage [7]. The use of formalin-fixed tissue biopsy is an alternative, although

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: FTA, Flinders Technology Associates; ccfDNA, circulating cell-free DNA; EF1 α , Elongation factor 1-alpha; EF1 γ , Elongation factor 1-gamma; SNX14, Sorting Nexin 14; Hsp70, heat shock protein 70; SOD, superoxide dismutase.

formalin is known to affect DNA quality. There are, however, novel methods for collection, storage and shipping of biological samples that offers some solutions to these problems. The Flinders Technology Associates (FTA) cards, for example, can be used to fix and preserve nucleic acids for long-term periods at ambient temperature with high stability [8,9]. This approach has been particularly useful for identification/genotyping and long-term surveillance of infectious pathogens in remote areas at reduced shipping costs [10–11]. The use of FTA cards for monitoring marine habitat with mussels as sentinel species, however, will require concomitant development of nucleic acid-based biomarkers. Fortunately, immense progress in the development has been achieved thanks to new methods in gene sequencing and the need for accurate predictive biomarkers in medicine. Nucleic acid-based biomarkers are particularly well-adapted for liquid biopsies. Liquid biopsy is a concept-based approach that is gaining significant attention in the biomedical area. The concept is that a clinician is able to use a small sample of blood collected from a patient to obtain vital genetic information that can be used: 1) to predict the onset of a disease, 2) to assess the severity of a disease (prognostic), 3) to evaluate and measure with high precision the efficacy of a given treatment, and 4) to determine long-term survival and relapse. It is designed to replace the use of logistically complicated, high cost invasive tissue biopsies to assess ongoing tissue damage. It is largely based on the analysis of circulating cell-free DNA (ccfDNA), which is considered as a genomic reservoir that is easily accessible for studying genetic traits, including genomic alterations [12]. Recent studies in humans have further shown that ccfDNA can be used to identify damages in specific tissues [13]. Liquid biopsies are also compatible for metagenomic data analysis. Microbiome information is now considered an important component of the biomarker schematic [14, 15]. With the help of recent advances in next generation sequencing, microbiome-associated biomarkers, just like ccfDNA, can be used in clinical practice for risk prediction, diagnosis and progression of a disease, or to predict and modulate response to treatment [16]. In marine ecology, microbial biomarkers are also being used for measuring the impact of anthropogenic activities on marine ecosystems [17, 18]. In the present work, we have combined the concept of liquid biopsy and the use of FTA cards to develop a liquid biopsy-based sampling method that involves fixation of nucleic acids on FTA cards. This method is simple, does not require maintaining a cold chain and is compatible with transcriptome analyses. We have also identified, for the first time, circulating cell-free ccfDNA in hemolymph of mussels and performed 16SrDNA gene-based bacterial microbiota profiles of mussels using FTA-fixed bacterial DNA collected from liquid biopsies.

Material and methods

Liquid biopsy

Adult specimens (55–70 mm length) of blue mussels *Mytilus* spp. were obtained from a commercial supplier (PEI Mussel King Inc., Prince Edward Islands, CANADA) and placed in a temperature-controlled (4°C) aerated aquarium containing 10 L of 32‰ artificial saline water (Reef Crystal artificial marine salt, Instant Ocean, VA, USA). For *M. desolationis* and *Aulacomya ater* (*A. ater*) specimens, they were collected on the intertidal rocky shore of Port-aux-Français (Kerguelen Islands, France) as previously described [19]. In all cases, hemolymph was collected from single individuals, without pooling and centrifugation. For routine preparation of liquid biopsies on FTA cards, the intervalvar liquid was removed with the tip of a knife and hemolymph was withdrawn from the adductor muscle using a syringe fitted with a 25 gauge needle. This method has been shown to be minimally invasive and non-lethal for freshwater bivalves [20]. Samples (<1ml of hemolymph) were immediately transferred into sterile 1.5 ml Eppendorf tube and centrifuged for 3 minutes at maximum speed (approx. 3000 x g) at room

temperature using a battery-powered mini-centrifuge (TOMY, Japan) (Fig 1). After centrifugation, aliquots (50 μ l) of supernatants were spotted on individual discs (10 mm diameter) on Whatman 903™ FTA cards (Whatman, Z694827, Sigma-Aldrich, Oakville, ON, Canada). The remaining supernatant was discarded and hemocyte pellets gently resuspended and spotted on FTA cards. In some experiments, Whatman FTA Gene Cards (GE Healthcare, Marlborough, MA) were used. The cards were dried for 15 min at room temperature and stored individually in small (3" x 4") clear 2 mil plastic ziplock baggies containing one silica gel desiccant (1gr/bag) moisture absorber. When indicated, hemolymph were pooled, transferred into a sterile 50 mL and centrifuged for 15 min at 3000 x g on a table low speed centrifuge (Beckman Coulter Allegra 6R). Unless otherwise indicated, all FTA cards were kept at room temperature in the dark.

Isolation of ccfDNA from hemolymph

ccfDNA from hemolymph (5 ml) was carried out using the NucleoSpin Plasma XS kit (Macherey Nagel) with optimized manufacturer's protocols. Purified DNA was quantified by the PicoGreen assay according to the manufacturer's recommendations [21].

Isolation of nucleic acid

For RNA isolation and purification, individual 12 mm² disks from FTA cards were carefully placed at the bottom of a sterile 0.6 ml microcentrifuge tube with a hole punctured in its bottom by a 18G needle. The tube was then transferred into a sterile 1.5 ml microcentrifuge tube before addition of TRIzol (800 μ l) to the upper tube. The cards were soaked with TRIzol for 10 min at room temperature and centrifuged for 3 min at high speed (8000 rpm) in a table centrifuge (BioFuge Fresco, Heraeus Instrument, Grimsby, ON, Canada). The TRIzol solution was transferred and topped with 0.2 ml of chloroform. Tubes were incubated 3 min at room temperature and centrifuged at 12000 x g for 15 min at 4°C. The aqueous phase was used to perform RNA extraction following a standard protocol. The RNA was dissolved in RNase/DNase free water and quantified using the Thermo Scientific™ NanoDrop™ spectrophotometer. DNA was isolated using an adapted the prepGEM™ Universal kit for Blood storage cards (Zygem, Charlottesville, VA). Briefly, 3 mm disc were placed on an eppendorf tube in washed with 100 μ l of DNA-free water and DNA recovered with the prepGEM solution following 5 min incubations at 75°C and 95°C. Nucleic acid samples were stored at -20°C until further analysis. Purified DNA was quantified by the PicoGreen assay (Lumiprobe, Hunt Valley, MD) according to the manufacturer's recommendations [21].

Thermal stress

Following acclimatization, groups of ten mussels of similar length were transferred into tank containing oxygenated seawater at 30°C or at 4°C (control). After 30 min, liquid biopsies were collected. Hemocytes pellets and hemolymph were either fixed on FTA cards or kept frozen at -20°C.

RT-PCR analysis

Total cellular RNA was isolated from cell pellets fixed on FTA cards using the TRIzol reagent (Life Technologies) as described above. First-strand cDNA was prepared from 2 μ g of cellular RNA in a total reaction volume of 20 μ l using the High Capacity cDNA Reverse Transcription Kit from Thermo-Fisher. qPCR specificity was carried out with a Rotor-Gene® 3000 (Corbett Research) with primers listed in Table 1. All qPCR reactions were carried out in 20 μ l with the




Fig 1. Diagrammatic summary of the sampling protocol. *Left*, for routine preparation of liquid biopsies on FTA cards, hemolymph was withdrawn from the adductor muscle. Samples (approx. 1ml of hemolymph) were immediately transferred into sterile 1.5 ml and centrifuged at room temperature using a

battery-powered mini-centrifuge (TOMY, Japan). After centrifugation, aliquots (50 μ l) of supernatants were spotted on individual discs (10 mm diameter) on FTA® cards. The remaining supernatant was discarded and hemocyte pellets gently resuspended and spotted on FTA cards. The cards were dried for 15 min at room temperature and stored individually in small plastic Ziploc baggies containing a silica gel desiccant moisture absorber. *Right*, to recover nucleic acids, discs were carefully placed at the bottom of a sterile microcentrifuge tube with a hole punctured in its bottom. The tube was then transferred into a larger sterile microcentrifuge tube before isolation of nucleic acids as described in details in the material and methods section.

<https://doi.org/10.1371/journal.pone.0223525.g001>

QuantiFast SYBR Green PCR Kit (Qiagen). qPCRs were amplified using the following conditions: 95°C for 5 min, followed by 40 cycles at 95°C for 10 s and 60°C for 30 s with a single acquisition. A final melting curve analysis was performed using at 95°C for 5 s, 65°C for 60 s and 97°C with a continuous acquisition. To ensure primer specificity, melting temperature curves were determined for each amplicon. PCR products were run on a 1.5% agarose gel to visualize the bands and ensure a unique band is amplified. The expression level (\log_2 of the fold change) was expressed 1) calculating the delta CT (Δ CT) by subtracting the averaged threshold cycle (CT) value of the three housekeeping genes (*EF1 γ* , *EF1 α* and *SNX14*) from the CT value of the target gene and 2) comparing the fold change of non-exposed mussels versus thermal-exposed mussels [22].

Taxonomic analysis of hemolymph microbiota

Following fixation on FTA cards, DNA was extracted from hemocyte pellets, as described above. The whole 16S bacterial DNA V2-V3 region was targeted by the primers 28F-519R primers and pyrosequenced by the Illumina Mi-Seq Next Generation Sequencer at Research & Testing Laboratory (<http://www.researchandtesting.com/>, Texas, USA). A detailed description of bioinformatic filters that were used for analysis can be found at http://www.rtlgenomics.com/docs/Data_Analysis_Methodology.pdf. Briefly, post-sequencing processing of the raw data for both types of samples (frozen and FTA-fixed cell pellets) was carried out by standard methods to eliminate low quality readings, contaminants and chimeras. The remaining filtered sequences were then grouped into Operational Taxonomy Units (OTUs) at a similarity level \geq 95%. For each sample, we have obtained approximately 10 000–20 000 OTUs. For taxonomic classification, we used the Ribosomal Database Project (RDP; <http://rdp.cme.msu.edu/>). The metagenomic data from this publication have been deposited to the NCBI Sequence Read Archives with the assigned the identifier PRJNA564645. To take into account the problem of contamination, we used, as controls, template-free FTA cards that were processed with the same DNA extraction and PCR amplification kits as the real samples, as previously recommended [23].

Statistics

All datasets were validated for normality (with Shapiro-Wilk test) and homoscedasticity (with Bartlett and F of Fisher test). Student's t-test was used as parametric test to compare data set with two groups and a two-way analysis of variance (ANOVA) was used to compare datasets with more than two groups. For non-parametric data, Kruskal Wallis test was used to

Table 1. List of primers used in this study.

Gene	Sense primers	Antisense primers	Product size
<i>EF1γ</i>	5' -AAA GCT CAA ATT GCA GCG GA-3'	5' -ATG cAA TCT cCT CCA GCT GT-3'	158 bp
<i>SXN14</i>	5' -AGA GGC GAG GAT CAG ATG TG-3'	5' -ATC TTC ACC ACC ACC AGA GG-3'	158 bp
<i>HSP70</i>	5' -ACA GAG AGA CTT GTT GGC GA-3'	5' -TTC CGA TCA GTC TCT TGG CA-3'	88 bp
<i>SOD</i>	5' -ACC ATG TGC GTC TAC AGG AT-3'	5' -TTG ATC AGG TTT CCG ATG GC-3'	95 bp

<https://doi.org/10.1371/journal.pone.0223525.t001>

compare datasets. All data are expressed as the mean \pm S.D. Results were considered statistically significant at $P \leq 0.05$.

Results

Isolation and analysis of RNA from hemocyte pellets fixed on FTA cards

Because hemocytes are implicated in several physiological functions, such as phagocytosis of pathogens, homeostasis, and reproduction, there is increasing interest in studying how their transcriptome is modulated by environmental stress factors [24–28]. We thus developed a general framework procedure for collecting hemolymph samples using FTA cards without the need of a cold chain is shown and detailed in Fig 1A. Other than a battery-operated centrifuge, it does not require specialized equipment and can be easily incorporated into a small, light-weight portable unit (S1 Fig).

To test whether such sampling framework is compatible for multi-omics analysis of hemolymph samples, we first compared RNA yields obtained from frozen hemocytes and hemocytes that were fixed on FTA cards. Isolation of RNA from FTA discs was carried out using a simple TRIzol-based protocol which consisted of soaking FTA discs in TRIzol for 10 min at room temperature (Fig 1B). Nucleic acids were then eluted from the discs by brief centrifugation and RNA purified using standard phase separation with chloroform and subsequent precipitation with isopropyl alcohol [29]. Our results showed that similar amount of RNA was recovered from pellets fixed on FTA discs as compared to pellets frozen at -20°C (Fig 2A). Both methods yielded similar ratio of the absorbance at 260 and 280 nm. RNA recovery from FTA discs could also be achieved using a commonly-used silica-based RNA extraction kit, which resulted in lower yield compared to the TRIzol-based method but provided purer RNA (Fig 2B). We also found that the RNA yield from hemocyte pellets fixed on FTA was not significantly affected by long term storage (up to 6 months) at 4°C or ambient temperatures (Fig 2C). We next compared the performance RT-qPCR carried out using frozen RNA and RNA fixed on FTA cards and found that both conservation methods performed equally when measuring expression of three commonly used housekeeping genes (*EF1 α* , *EF1 γ* , and *SNX14*) (Fig 3A). Both methods were also equally efficient measuring the *de novo* expression of stress-related genes (*Hsp70* and *SOD*) in hemocytes collected from mussels subjected to an acute thermal stress (Fig 3B).

Bacterial microbiome analysis

Bacteria microbiome-derived profiles are emerging as potent biomarkers not only in the biomedical field, but also in monitoring environmental stress in a vast number of species, including marine organisms [30, 31]. For these experiments, we compared the 16SrDNA gene-based bacterial profiles from frozen and FTA-conserved DNA from cellular pellets. Our results showed that both methods generated similar bacterial profiles with regards to phylum (Fig 4A). We found no significant differences in genus-level bacterial groups that were most frequently found in hemolymph (Fig 4B). The most common bacteria found in the hemolymph was from the genus *Colwellia*. This genus is commonly found in marine habitats, including the hemolymph of blue mussels [18]. Taken together, these findings show that FTA-fixed DNA is perfectly compatible with metagenomic analysis of circulating microbiota.

Circulating cell-free DNA

Analysis of ccfDNA from liquid biopsies is another promising avenue for the development of predictive biomarkers. In humans, it serves as a genomic reservoir that is easily accessible with




Fig 2. RNA extraction from FTA cards. (A) Recovery of total RNA from hemocytes stored frozen at -20°C or on FTA cards using a TRIzol-based protocol. (B) Comparative RNA recovery from FTA cards using TRIzol or the RNeasy kit. (C) Stability of RNA fixed on FTA cards and kept at different temperatures. * $p < 0.05$. Values are expressed as the total amount of RNA recovered from a 10 mm diameter sample disc FTA cards.

<https://doi.org/10.1371/journal.pone.0223525.g002>




Fig 3. RT-qPCR analysis using frozen and FTA-fixed RNA. (A) Cycle thresholds (Ct) values obtained for expression analysis of three housekeeping genes in pooled hemocytes pellets that were either kept frozen or fixed on FTA cards. (B) Relative gene expression of *Hsp70* and superoxide dismutase (*SOD*) in hemocytes following an acute thermal stress at 30°C for 30 min. Expression levels are relative to controls (4°C). No significant (N.S.) differences were found using both methods.

<https://doi.org/10.1371/journal.pone.0223525.g003>




Fig 4. Bacterial profiles of hemolymph. Upper histogram, pie chart representation of the relative abundance (in percentage) of 16SrRNA of the most common phyla found in liquid biopsies of *M. edulis*. Lower histogram, bacterial species differentially abundant in frozen and FTA-fixed liquid biopsies.

<https://doi.org/10.1371/journal.pone.0223525.g004>

Table 2. Concentrations of ccfDNA.

Samples	ccfDNA (ng/mL)
Human plasma	8,72 ± 1,43
<i>M. edulis</i>	4,23 ± 0,18

<https://doi.org/10.1371/journal.pone.0223525.t002>

a minimally-invasive procedure. To our knowledge, however, the existence of ccfDNA in hemolymph of invertebrates, and in mussels in particular, has not been reported. Using the same method that is commonly used for isolating ccfDNA (see [material and methods](#) section for details) from human plasma, we have thus investigated the presence of ccfDNA in mussels. Our results revealed that we indeed found ccfDNA in hemolymph of *Mytilus edulis* (**Table 2**) at a range of concentrations (nanograms/ml) that were similar to that found in humans [32]. Similar results were obtained for *M. desolationis* and *Aulacomya ater* (*A. ater*). To validate the integrity of the ccfDNA fragments in liquid biopsies (cell-free hemolymph) fixed on FTA cards, we carried out real-time qPCR analysis using primers that generated small amplicons and that were specific for *EF1γ*, *HSP70* and *SNX14* genes [33] (**Table 3**). Our results showed that we could indeed find evidence of ccfDNA fragments encoding *SNX14* and *HSP70* genes but not *EF1γ*, suggesting that ccfDNA fragments are generated in a gene-specific manner, as observed in humans (**Table 3**). Indeed, ccfDNA fragments have been shown to derive from cell-specific nucleosome footprinting from different cell types [13]. These results reveal for the first time the presence of ccfDNA in mussels and open the door for its use as a biomarker in liquid biopsies collected from mussels.

Discussion

Given the growing concerns over the anthropogenic impact on marine ecosystems, it is increasingly critical to develop biomarkers that can be used as diagnostic and predictive tools, as well as for monitoring the success of the remediation efforts. The use of multi-omics biomarkers is a step in that direction. It obliges us, however, to go back to the drawing board to re-examine our sampling strategy not only for logistical reasons, but for economical reasons as well. This is particularly true for polar regions where the logistic complexity of tissue sampling is a major obstacle. In the present work, we have combined the use of FTA cards and the concept of liquid biopsies for the development of novel omics-based biomarkers in mussels, a widely used sentinel species. More specifically, we have shown that: 1) hemocytes pellets can be readily fixed on FTA cards and used for gene expression analysis; 2) FTA-fixed cell pellets are stable for long term periods when stored at room temperature, facilitating the biobanking activities and shipment to third parties at lower cost when compared to frozen samples; 3) our sampling strategy is easily adaptable for microbiome analysis of bacterial profiles, and 4) ccfDNA is present in mussel hemolymph and can be used as a genomic reservoir for the development of novel biomarkers. Taken together, this study lays the foundation for a logistically-friendly, omics-based multi-biomarker approach using mussels to assess the quality of marine ecosystems.

Until now, the concept of liquid biopsies has been almost exclusively used for guiding clinical practice, especially in oncology, prenatal screening, transplantation and presence of non-human ccfDNA (virus-derived sequences, for example) [12, 34, 35]. One of the main benefits of non-invasive liquid biopsies is that it overcomes many of the drawbacks associated with tissue biopsies. It also provides an accurate snapshot of the genomic landscape, bypassing many problems associated with tissue heterogeneity. In fact, to our knowledge, this is the first report of the presence of ccfDNA in invertebrates. In humans, the normal plasma concentration of

Table 3. qPCR analysis of ccfDNA fixed on FTA cards.

Gene	Template	CT	MT	% of detection
EF1 γ	ccfDNA	25,31 \pm 0,85	78,50 \pm 1,71	0%
	NTC	25,79 \pm 0,67	79,65 \pm 0,21	
SNX14	ccfDNA	24,64 \pm 2,44*	83,84 \pm 0,47	56%
	NTC	31,79 \pm 0.01	N.D.	
HSP70	ccfDNA	21,25 \pm 1,55 *	81,51 \pm 0,14 *	89%
	NTC	32,16 \pm 0	80 \pm 0	

NTC, No template control. CT, threshold cycle, MT, melting temperature, and % of amplicon detection obtained for EF1 γ , SNX14 and HSP70 (n = 9 mussels, with 3 replicates for each sample. ND = no detected.

* $p < 0.05$).

<https://doi.org/10.1371/journal.pone.0223525.t003>

ccfDNA is generally below 10–20 ng/ml, which is relatively similar to what we found in hemolymph of mussels. While it can increase by 5–10 times in patients with malignant disease or stress conditions [36]. Future studies will be needed to determine whether measuring ccfDNA levels in mussels can be used as a rapid stress biomarker as it does in humans [32, 36, 37]. Analysis of ccfDNA is also perfectly adaptable for serial samplings of DNA abnormalities, gene expression analysis, point mutations, chromosome translocations, miRegulome, epigenetic changes, etc. For example, analysis of ccfDNA can theoretically be used for measuring mutations in genes associated with hematopoietic neoplasia (such as *p53*), for rapid identification of mussel species, or for any genomic traits. It could also be used to detect the presence of parasitic and viral DNA to reduce the risk for public human health. From an ethical point of view, it reduces incidental mortality associated with conventional tissue biopsies. To develop the full potential of this approach, we are currently using next-generation sequencing to perform a genomic analysis of the entire ccfDNA found in different mussel populations. The results of these analyses will be reported in a separate manuscript.

Another benefit of combining both FTA-based technology and liquid biopsy is its compatibility with microbiome analysis. In humans, although blood is normally considered a sterile environment, there is increasing studies showing the circulating cells harbor a rich bacterial microbiome that can be used as disease biomarkers [38, 39]. This is also true for bivalves. The microbiome of hemolymph samples collected from the adductor muscle from Pacific oysters, for example, has been shown to undergo significant changes following abiotic and biotic stress factors [40]. A recent study showing that bacterial diversity and richness in mussels were higher in the hemolymph compared to other tissues support the idea that liquid biopsies is a valid approach for microbiome analysis in mussels [18]. Indeed, Vezzulli and colleagues have shown that the alpha diversity of the microbiome of blue mussels was higher in the hemolymph compared to the digestive gland, in both [18]. Our results further showed that the bacterial profiles obtained from DNA of cells fixed on FTA was not significantly different from that obtained from frozen cells (S2 Fig). We found no significant differences in those bacteria that are commonly found in marine ecosystems, such as *Alteromonadales* or *Oceanospirillales* [41, 42]. In fact, our results showed that DNA from cells fixed on FTA cards was more prone to detect those rare bacterial DNA, consistent with the ability of the chemical matrix of FTA cards to stabilize and protect nucleic acids from environmental degradation. Not surprisingly, one of the most abundant bacterial DNA that we found in both FTA cards and frozen samples derived from the *Polaribacter* genus, which species are almost always found in polar or marine environments [43]. The other most frequently found genus were *Colwellia*, *Oleispira*, *Cycloclasticus* and *Neptunomonas*. These bacteria are well known to degrade polycyclic aromatic

hydrocarbons (PAH) or short-chain alkanes. *Cycloclasticus* bacterial DNA has been recently found in *Bathymodiolus heckeriae* mussels where they have established a symbiosis at a 3000m depth in oil-rich habitats near asphalt volcanoes in the Gulf of Mexico [44]. They have also been shown to bloom in marine habitats following the Deep Horizon oil spill [44]. Interestingly, the *Oleispira* genus, another bacterium that has bloomed following the Deep Horizon oil spill, was also among the most frequently found bacterium in *M. edulis*. The third most frequent bacteria found was from the genus *Neptunomonas*, another bacterium implicated in degradation of PAHs and normally found in contaminated sediments [45]. Whether the presence of this microbiota reflects a symbiotic relationship with *M. edulis* is unclear at the present time. Overall, our findings indicate that microbiome analysis in liquid biopsies collected from cellular pellets of mussels is an interesting avenue to study dysbiosis in mussels following exposure to environmental stress.

Our study showed that combining liquid biopsies and the FTA technology could be useful for long-term biobanking and retesting of omics biomarkers collected from sentinel mussels, particularly in remote regions. This approach is also economically viable, especially when using the Whatman 903™ FTA cards. Although we have used for this study 10 mm diameter circular FTA discs, we obtained similar RNA yield using smaller discs (S3 Fig). If necessary, one could use the the Whatman FTA Gene Cards, which have been adapted for capture on nucleic acids. Although more expensive than regular Whatman 903™ cards, we found that they are more performant for DNA recovery (S4 Fig). It is important to note, however, that there are multiple options for isolating nucleic acid from FTA cards. In our hands, we have found that TRIzol was an adequate, low cost, and versatile method for routine RT-qPCR analysis. It can also be combined with commercial kits to obtain purer RNA preparations (S1 Table). TRIzol is also suitable for isolation of small RNAs (<200 nucleotides) as compared to the silica-based methods which do not retain small RNAs [46]. This has to be taken into consideration for future studies aimed at testing the possibility of using non-coding RNAs as putative biomarkers in hemolymph of mussels. In humans, non-coding RNAs have shown great promise as potential cancer biomarkers for liquid biopsies.

Another clear benefit of combining liquid biopsy and the FTA technology is from a logistically-point of view. We can now collect samples at remote site with a minimal amount of equipment (S1 Fig). Moreover, this approach addresses increasing concerns in field work ethics [47]. Liquid biopsies require sampling of a very small volume of hemolymph and does not lead to animal death. We believe that such approach could also be extended to other marine and freshwater organisms and offers many advantages for the development of long-term ecological observatories in polar regions threatened by anthropogenic activities.

Supporting information

S1 Fig. Portable laboratory carrying case used for liquid biopsy sampling. The case contains a battery powered minicentrifuge and a portable solar panel battery charger for long term expeditions. The case also contains all necessary reagents and equipment to collect biopsies on site.

(TIFF)

S2 Fig. Relative abundance of order-level bacteria in frozen and FTA-fixed liquid biopsies. Data are shown as mean \pm SD from duplicates.

(TIFF)

S3 Fig. RNA extraction from FTA cards. (A) Recovery of total RNA from a 10 mm disc (100%) and a quarter of a disc (25%). (B) RNA recovery following soaking for 10 min versus

overnight at room temperature before centrifugation. Results are representative of two independent experiments. RNA recovery was carried out using the TRIzol-based method. (TIFF)

S4 Fig. Comparative analysis of the performance of two different types of FTA cards. DNA recovery from hemocyte pellets that were fixed on the Whatman 903™ cards or the Whatman Gene FTA cards. Values are expressed as the total amount of DNA recovered from a 3 mm diameter sample disc using the PropGEM-based protocol. Results are representative of two independent experiments (n = 12/fgroup). * $p < 0.05$. (TIFF)

S1 Table. RT-qPCR analysis of RNA extracted from frozen or FTA-fixed cell pellets. (DOCX)

Acknowledgments

The authors would like to thank Ms. Marlène Fortier for her technical help, all the personnel from the French Polar Institute Paul Emile Victor (IPEV) and the Terres Australes et Antarctiques Françaises (TAAF) for their help and hospitality during our stay in the Kerguelen archipelago.

Author Contributions

Conceptualization: France Caza, Philippine Granger Joly de Boissel, Stéphane Betoulle, Yves St-Pierre.

Formal analysis: France Caza, Philippine Granger Joly de Boissel, Richard Villemur.

Funding acquisition: Stéphane Betoulle, Yves St-Pierre.

Investigation: France Caza, Philippine Granger Joly de Boissel, Stéphane Betoulle, Yves St-Pierre.

Methodology: France Caza, Philippine Granger Joly de Boissel, Stéphane Betoulle, Yves St-Pierre.

Project administration: Yves St-Pierre.

Resources: Stéphane Betoulle, Yves St-Pierre.

Supervision: Yves St-Pierre.

Validation: France Caza, Philippine Granger Joly de Boissel.

Writing – original draft: Yves St-Pierre.

Writing – review & editing: France Caza, Philippine Granger Joly de Boissel, Richard Villemur, Stéphane Betoulle.

References

1. Beyer J, Green NW, Brooks S, Allan IJ, Ruus A, Gomes T, et al. Blue mussels (*Mytilus edulis* spp.) as sentinel organisms in coastal pollution monitoring: a review. *Marine environmental research*. 2017 Sep 1; 130:338–65. <https://doi.org/10.1016/j.marenvres.2017.07.024> PMID: 28802590
2. Bodin N, Burgeot T, Stanisiere JY, Bocquené G, Menard D, Minier C, et al. Seasonal variations of a battery of biomarkers and physiological indices for the mussel *Mytilus galloprovincialis* transplanted into the northwest Mediterranean Sea. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 2004 Aug 1; 138(4):411–27.

3. Dondero F, Dagnino A, Jonsson H, Capri F, Gastaldi L, Viarengo A. Assessing the occurrence of a stress syndrome in mussels (*Mytilus edulis*) using a combined biomarker/gene expression approach. *Aquatic Toxicology*. 2006 Jun 1; 78:S13–24. <https://doi.org/10.1016/j.aquatox.2006.02.025> PMID: 16644027
4. Farcy E, Burgeot T, Haberkorn H, Auffret M, Lagadic L, Allenou JP, et al. An integrated environmental approach to investigate biomarker fluctuations in the blue mussel *Mytilus edulis* L. in the Vilaine estuary, France. *Environmental Science and Pollution Research*. 2013 Feb 1; 20(2):630–50. <https://doi.org/10.1007/s11356-012-1316-z> PMID: 23247516
5. Berg DJ, Haag WR, Guttman SI, Sickel JB. Mantle biopsy: a technique for nondestructive tissue-sampling of freshwater mussels. *Journal of the North American Benthological Society*. 1995 Dec 1; 14(4):577–81.
6. Mioduchowska M, Kaczmarczyk A, Zając K, Zając T, Sell J. Gender-associated mitochondrial DNA heteroplasmy in somatic tissues of the endangered freshwater mussel *Unio crassus* (Bivalvia: Unionidae): implications for sex identification and phylogeographical studies. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*. 2016 Nov; 325(9):610–25.
7. Lacoursière-Roussel A, Howland K, Normandeau E, Grey EK, Archambault P, Deiner K, et al. eDNA metabarcoding as a new surveillance approach for coastal Arctic biodiversity. *Ecology and evolution*. 2018 Aug; 8(16):7763–77. <https://doi.org/10.1002/ece3.4213> PMID: 30250661
8. da Cunha Santos G. FTA Cards for Preservation of Nucleic Acids for Molecular Assays: A Review on the Use of Cytologic/Tissue Samples. *Archives of pathology & laboratory medicine*. 2018 Mar; 142(3):308–12.
9. Muthukrishnan M, Singanallur NB, Ralla K, Villuppanoor SA. Evaluation of FTA® cards as a laboratory and field sampling device for the detection of foot-and-mouth disease virus and serotyping by RT-PCR and real-time RT-PCR. *Journal of virological methods*. 2008 Aug 1; 151(2):311–6. <https://doi.org/10.1016/j.jviromet.2008.05.020> PMID: 18584888
10. Liang X, Chigerwe M, Hietala SK, Crossley BM. Evaluation of Fast Technology Analysis (FTA) Cards as an improved method for specimen collection and shipment targeting viruses associated with Bovine Respiratory Disease Complex. *Journal of virological methods*. 2014 Jun 15; 202:69–72. <https://doi.org/10.1016/j.jviromet.2014.02.022> PMID: 24657552
11. Tam KI, Esona MD, Williams A, Ndze VN, Boula A, Bowen MD. Evaluation of BBL™ Sensi-Discs™ and FTA® cards as sampling devices for detection of rotavirus in stool samples. *Journal of virological methods*. 2015 Sep 15; 222:41–6. <https://doi.org/10.1016/j.jviromet.2015.05.007> PMID: 26022083
12. Fan HC, Blumenfeld YJ, Chitkara U, Hudgins L, Quake SR. Noninvasive diagnosis of fetal aneuploidy by shotgun sequencing DNA from maternal blood. *Proceedings of the National Academy of Sciences*. 2008 Oct 21; 105(42):16266–71.
13. Snyder MW, Kircher M, Hill AJ, Daza RM, Shendure J. Cell-free DNA comprises an in vivo nucleosome footprint that informs its tissues-of-origin. *Cell*. 2016 Jan 14; 164(1–2):57–68. <https://doi.org/10.1016/j.cell.2015.11.050> PMID: 26771485
14. Gilbert JA, Quinn RA, Debelius J, Xu ZZ, Morton J, Garg N, et al. Microbiome-wide association studies link dynamic microbial consortia to disease. *Nature*. 2016 Jul; 535(7610):94. <https://doi.org/10.1038/nature18850> PMID: 27383984
15. Guglielmi G. How gut microbes are joining the fight against cancer. *Nature*. 2018 May; 557(7706):482. <https://doi.org/10.1038/d41586-018-05208-8> PMID: 29795257
16. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinetz TV, et al. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science*. 2018 Jan 5; 359(6371):97–103. <https://doi.org/10.1126/science.aan4236> PMID: 29097493
17. Apprill A. Marine animal microbiomes: toward understanding host–microbiome interactions in a changing ocean. *Frontiers in Marine Science*. 2017 Jul 18; 4:222.
18. Vezzulli L, Stagnaro L, Grande C, Tassistro G, Canesi L, Pruzzo C. Comparative 16SrDNA gene-based microbiota profiles of the Pacific oyster (*Crassostrea gigas*) and the Mediterranean mussel (*Mytilus galloprovincialis*) from a shellfish farm (Ligurian Sea, Italy). *Microbial ecology*. 2018 Feb 1; 75(2):495–504. <https://doi.org/10.1007/s00248-017-1051-6> PMID: 28803409
19. Caza F, Betoulle S, Auffret M, Brousseau P, Fournier M, St-Pierre Y. Comparative analysis of hemocyte properties from *Mytilus edulis* desolationis and *Aulacomya ater* in the Kerguelen Islands. *Marine environmental research*. 2015 Sep 1; 110:174–82. <https://doi.org/10.1016/j.marenvres.2015.09.003> PMID: 26382607
20. Gustafson LL, Stoskopf MK, Bogan AE, Showers W, Kwak TJ, Hanlon S, et al. Evaluation of a nonlethal technique for hemolymph collection in *Elliptio complanata*, a freshwater bivalve (Mollusca: Unionidae). *Diseases of aquatic organisms*. 2005 Jun 30; 65(2):159–65. <https://doi.org/10.3354/dao065159> PMID: 16060269

21. Chiminqgi M, Moutereau S, Pernet P, Conti M, Barbu V, Lemant J, et al. Specific real-time PCR vs. fluorescent dyes for serum free DNA quantification. *Clinical Chemical Laboratory Medicine*. 2007 Aug 1; 45(8):993–5.
22. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *methods*. 2001 Dec 1; 25(4):402–8. <https://doi.org/10.1006/meth.2001.1262> PMID: 11846609
23. Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, et al. Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC biology*. 2014 Dec; 12(1):87.
24. Tanguy M, McKenna P, Gauthier-Clerc S, Pellerin J, Danger JM, Siah A. Sequence analysis of a normalized cDNA library of *Mytilus edulis* hemocytes exposed to *Vibrio splendidus* LGP32 strain. *Results in immunology*. 2013 Jan 1; 3:40–50. <https://doi.org/10.1016/j.rinim.2013.04.001> PMID: 24600557
25. Moreira R, Pereiro P, Canchaya C, Posada D, Figueras A, Novoa B. RNA-Seq in *Mytilus galloprovincialis*: comparative transcriptomics and expression profiles among different tissues. *BMC genomics*. 2015 Dec; 16(1):728.
26. Detree C, Núñez-Acuña G, Roberts S, Gallardo-Escárate C. Uncovering the complex transcriptome response of *Mytilus chilensis* against saxitoxin: implications of harmful algal blooms on mussel populations. *PloS one*. 2016 Oct 20; 11(10):e0165231. <https://doi.org/10.1371/journal.pone.0165231> PMID: 27764234
27. Dong W, Chen Y, Lu W, Wu B, Qi P. Transcriptome analysis of *Mytilus coruscus* hemocytes in response to *Vibrio alginolyticus* infection. *Fish & shellfish immunology*. 2017 Nov 1; 70:560–7.
28. de Boissel PG, Fournier M, Rodriguez-Lecompte JC, McKenna P, Kibenge F, Siah A. Functional and molecular responses of the blue mussel *Mytilus edulis*' hemocytes exposed to cadmium-An in vitro model and transcriptomic approach. *Fish & shellfish immunology*. 2017 Aug 1; 67:575–85.
29. Chomczynski P, Sacchi N. The single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction: twenty-something years on. *Nature protocols*. 2006 Aug; 1(2):581. <https://doi.org/10.1038/nprot.2006.83> PMID: 17406285
30. Bourlat SJ, Borja A, Gilbert J, Taylor MI, Davies N, Weisberg SB, et al. Genomics in marine monitoring: new opportunities for assessing marine health status. *Marine pollution bulletin*. 2013 Sep 15; 74(1):19–31. <https://doi.org/10.1016/j.marpolbul.2013.05.042> PMID: 23806673
31. Carugati L, Corinaldesi C, Dell'Anno A, Danovaro R. Metagenetic tools for the census of marine meiofaunal biodiversity: an overview. *Marine Genomics*. 2015 Dec 1; 24:11–20. <https://doi.org/10.1016/j.margen.2015.04.010> PMID: 25957694
32. Dawson SJ. Characterizing the Cancer Genome in Blood. *Cold Spring Harbor perspectives in medicine*. 2018 May 29:a026880.
33. Cubero-Leon E, Ciocan CM, Minier C, Rotchell JM. Reference gene selection for qPCR in mussel, *Mytilus edulis*, during gametogenesis and exogenous estrogen exposure. *Environmental Science and Pollution Research*. 2012 Aug 1; 19(7):2728–33.
34. Diaz LA Jr, Bardelli A. Liquid biopsies: genotyping circulating tumor DNA. *Journal of clinical oncology*. 2014 Feb 20; 32(6):579. <https://doi.org/10.1200/JCO.2012.45.2011> PMID: 24449238
35. De Vlaminck I, Martin L, Kertesz M, Patel K, Kowarsky M, Strehl C, et al. Noninvasive monitoring of infection and rejection after lung transplantation. *Proceedings of the National Academy of Sciences*. 2015 Oct 27; 112(43):13336–41.
36. Huttunen R, Kuparinen T, Jylhävä J, Aittoniemi J, Vuento R, Huhtala H, et al. Fatal outcome in bacteremia is characterized by high plasma cell free DNA concentration and apoptotic DNA fragmentation: a prospective cohort study. *PloS one*. 2011 Jul 1; 6(7):e21700. <https://doi.org/10.1371/journal.pone.0021700> PMID: 21747948
37. Forsblom E, Aittoniemi J, Ruotsalainen E, Helmijoki V, Huttunen R, Jylhävä J, et al. High cell-free DNA predicts fatal outcome among *Staphylococcus aureus* bacteraemia patients with intensive care unit treatment. *PloS one*. 2014 Feb 10; 9(2):e87741. <https://doi.org/10.1371/journal.pone.0087741> PMID: 24520336
38. Potgieter M, Bester J, Kell DB, Pretorius E. The dormant blood microbiome in chronic, inflammatory diseases. *FEMS microbiology reviews*. 2015 May 4; 39(4):567–91. <https://doi.org/10.1093/femsre/fuv013> PMID: 25940667
39. Qian Y, Yang X, Xu S, Wu C, Qin N, Chen SD, et al. Detection of Microbial 16S rRNA Gene in the Blood of Patients With Parkinson's Disease. *Frontiers in aging neuroscience*. 2018;10.
40. Lokmer A, Wegner KM. Hemolymph microbiome of Pacific oysters in response to temperature, temperature stress and infection. *The ISME journal*. 2015 Mar; 9(3):670. <https://doi.org/10.1038/ismej.2014.160> PMID: 25180968

41. Evans FF, Egan S, Kjelleberg S. Ecology of type II secretion in marine gammaproteobacteria. *Environmental microbiology*. 2008 May; 10(5):1101–7. <https://doi.org/10.1111/j.1462-2920.2007.01545.x> PMID: [18218035](https://pubmed.ncbi.nlm.nih.gov/18218035/)
42. Jensen S, Duperron S, Birkeland NK, Hovland M. Intracellular Oceanospirillales bacteria inhabit gills of *Acesta bivalves*. *FEMS microbiology ecology*. 2010 Dec 1; 74(3):523–33. <https://doi.org/10.1111/j.1574-6941.2010.00981.x> PMID: [21044098](https://pubmed.ncbi.nlm.nih.gov/21044098/)
43. Gosink JJ, Woese CR, Staley JT. *Polaribacter* gen. nov., with three new species, *P. irgensii* sp. nov., *P. franzmannii* sp. nov. and *P. filamentus* sp. nov., gas vacuolate polar marine bacteria of the Cytophaga-Flavobacterium-Bacteroides group and reclassification of 'Flectobacillus glomeratus' as *Polaribacter glomeratus* comb. nov. *International Journal of Systematic and Evolutionary Microbiology*. 1998 Jan 1; 48(1):223–35.
44. Rubin-Blum M, Antony CP, Borowski C, Sayavedra L, Pape T, Sahling H, et al. Short-chain alkanes fuel mussel and sponge *Cycloclasticus* symbionts from deep-sea gas and oil seeps. *Nature microbiology*. 2017 Aug; 2(8):17093.
45. Hedlund BP, Geiselbrecht AD, Bair TJ, Staley JT. Polycyclic aromatic hydrocarbon degradation by a new marine bacterium, *Neptunomonas naphthovorans* gen. nov., sp. nov. *Appl. Environ. Microbiol.* 1999 Jan 1; 65(1):251–9. PMID: [9872786](https://pubmed.ncbi.nlm.nih.gov/9872786/)
46. Sultan M, Amstislavskiy V, Risch T, Schuette M, Dökel S, Ralser M, et al. Influence of RNA extraction methods and library selection schemes on RNA-seq data. *BMC genomics*. 2014 Dec; 15(1):675.
47. Costello M, Beard KH, Corlett RT, Cumming GS, Devictor V, Loyola R, et al. 2016. Field work ethics in biological research.