



HAL
open science

Acclimation to Water Restriction Implies Different Paces for Behavioral and Physiological Responses in a Lizard Species

David Rozen-Rechels, Andréaz Dupoué, Sandrine Meylan, Kenza Qitout, Beatriz Decencièrre, Simon Agostini, Jean-François Le Galliard

► **To cite this version:**

David Rozen-Rechels, Andréaz Dupoué, Sandrine Meylan, Kenza Qitout, Beatriz Decencièrre, et al.. Acclimation to Water Restriction Implies Different Paces for Behavioral and Physiological Responses in a Lizard Species. *Physiological and Biochemical Zoology*, 2020, 93 (2), pp.160-174. <10.1086/707409>. <hal-02480584>

HAL Id: hal-02480584

<https://hal.science/hal-02480584v1>

Submitted on 9 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Physiological and Biochemical Zoology

Acclimation to water restriction implies different paces for behavioral and physiological responses in a lizard species

--Manuscript Draft--

Manuscript Number:	19062R2
Full Title:	Acclimation to water restriction implies different paces for behavioral and physiological responses in a lizard species
Short Title:	Behavioral and physiological responses to water restriction
Article Type:	Research Paper
Corresponding Author:	David Rozen-Rechels Sorbonne Universite Paris cedex 5, Ile-De-France FRANCE
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Sorbonne Universite
Corresponding Author's Secondary Institution:	
First Author:	David Rozen-Rechels
First Author Secondary Information:	
Order of Authors:	David Rozen-Rechels Andréaz Dupoué Sandrine Meylan Kenza Qitout Beatriz Decencièrè Simon Agostini Jean-François Le Galliard
Order of Authors Secondary Information:	
Abstract:	<p>Chronic changes in climate conditions may select for acclimation responses in terrestrial animals living in fluctuating environments and beneficial acclimation responses may be key to the resilience of these species to global changes. Despite evidence that climate warming induces changes in water availability, acclimation responses to water restriction are understudied relative to thermal acclimation. In addition, acclimation responses may involve different modes, paces and trade-offs between physiological and behavioral traits. Here, we tested the dynamical acclimation responses of a dry-skinned terrestrial ectotherm to a chronic water restriction. Yearling common lizards (<i>Zootoca vivipara</i>) were exposed to a sub-lethal water restriction during 2 months of the summer season in laboratory conditions, then released in outdoor conditions for 10 additional months. Candidate behavioral (exploration, basking, and thermal preferences) and physiological traits (metabolism at rest and standard water loss rate) potentially involved in the acclimation response were measured repeatedly during and after water restriction. We observed a sequential acclimation response in water restricted animals (yearlings spent less time basking during the first weeks of water deprivation) which was followed by delayed sex-specific physiological consequences of the water restriction during the following months (thermal depression in males and lower standard evaporative water loss rates in females). Despite short-term negative effects of water restriction on body growth, annual growth, survival and reproduction were not significantly different between water restricted and control yearlings. This demonstrates that beneficial acclimation responses to water restriction involve both short-term flexible behavioral responses and delayed changes in thermal and water biology traits.</p>

1 **Acclimation to water restriction implies different paces for behavioral and physiological**
2 **responses in a lizard species**

3

4 **Running title:** Behavioral and physiological responses to water restriction

5 **Keywords:** body temperature; water availability; activity; evaporative water loss; exploration;
6 metabolism; reptiles

7 **What is already known:** Recent studies on dry-skinned terrestrial ectotherms have shown that behavioral
8 and physiological responses to chronic changes in water availability (acclimation responses) are crucial
9 determinants of their resilience to global changes. The acclimation response is in general a multi-faceted
10 process that involves a range of traits with different sensitivities and paces of response. Unfortunately,
11 quantification of joint behavioral and physiological acclimation responses to chronic water stress are rare and
12 the relative importance of each response mechanism still remains to be elucidated.

13 **What the study adds:** Our results confirm a sequential acclimation response involving short-term
14 behavioral adjustments followed by delayed sex-specific changes of thermal preferences (thermal
15 depression) and of water loss rates (a reduction of cutaneous water loss). This study highlights the
16 importance of jointly quantifying behavior and physiology in order to understand acclimation responses in
17 wild animals.

18 **Abstract**

19 Chronic changes in climate conditions may select for acclimation responses in terrestrial animals
20 living in fluctuating environments and beneficial acclimation responses may be key to the resilience
21 of these species to global changes. Despite evidence that climate warming induces changes in water
22 availability, acclimation responses to water restriction are understudied relative to thermal
23 acclimation. In addition, acclimation responses may involve different modes, paces and trade-offs
24 between physiological and behavioral traits. Here, we tested the dynamical acclimation responses of
25 a dry-skinned terrestrial ectotherm to a chronic water restriction. Yearling common lizards (*Zootoca*
26 *vivipara*) were exposed to a sub-lethal water restriction during 2 months of the summer season in
27 laboratory conditions, then released in outdoor conditions for 10 additional months. Candidate
28 behavioral (exploration, basking, and thermal preferences) and physiological traits (metabolism at
29 rest and standard water loss rate) potentially involved in the acclimation response were measured
30 repeatedly during and after water restriction. We observed a sequential acclimation response in
31 water restricted animals (yearlings spent less time basking during the first weeks of water
32 deprivation) which was followed by delayed sex-specific physiological consequences of the water
33 restriction during the following months (thermal depression in males and lower standard
34 evaporative water loss rates in females). Despite short-term negative effects of water restriction on
35 body growth, annual growth, survival and reproduction were not significantly different between
36 water restricted and control yearlings. This demonstrates that beneficial acclimation responses to
37 water restriction involve both short-term flexible behavioral responses and delayed changes in
38 thermal and water biology traits.

39 **Introduction**

40 Adaptive plasticity is essential for organisms to cope with spatio-temporal variability of their
41 environment and is often the leading mechanism by which they can face the negative ecological
42 impacts of ongoing and future global changes (Somero 2010; Kelly et al. 2012; Seebacher et al.
43 2015; Wong and Candolin 2015). In particular, thermal plasticity refers to the capacity to adjust
44 phenotypes to changes in thermal conditions and is a major component of species' climate warming
45 resistance (Sinervo et al. 2010; Gunderson and Stillman 2015; Mitchell et al. 2018). Global changes
46 also entail modifications of rainfall regimes or drought events and therefore water availability in the
47 environment (Field et al. 2012). Water is a critical resource for most terrestrial animals, altering, for
48 example, their locomotor performances (Cheuvront and Kenefick 2014; Anderson and Andrade
49 2017) and life history strategies (Lorenzon et al. 2001; Marquis et al. 2008). Terrestrial organisms
50 have therefore evolved numerous plastic strategies to cope with spatial and temporal variation in
51 water availability, hereafter referred to as the hydroregulation tactics (Ostwald et al. 2016; Eto et al.
52 2017; Pirtle et al. 2019). Hydroregulation tactics play a key role in the water balance regulation (i.e.
53 the balance between water inputs and water losses) and the plasticity of hydroregulation tactics will
54 be as critical as thermal plasticity to predict the consequences of global changes on organisms (e.g.,
55 Peterman and Semlitsch 2014; Kearney et al. 2018).

56 Chronic changes in water availability or water losses induced by predictable seasonal fluctuations in
57 rainfall and temperature or by sustained weather events (e.g., warm spells) may lead to acclimation
58 responses in terrestrial animals. These acclimation responses may allow organisms to reduce the
59 costs of performance loss associated with dehydration, and thus be beneficial in their response to
60 environmental changes in water balance regulation (i.e. beneficial acclimation hypothesis, Leroi et
61 al. 1994; Huey et al. 1999; see Anderson and Andrade 2017 for an example). The acclimation
62 response of a given species is usually a multi-faceted process that involves a range of reversible

63 morphological and physiological changes to maintain water balance, including modifications of
64 metabolic rates, adjustments in renal function and osmoregulation, modulations of cutaneous and
65 respiratory water loss rates, or adjustments of body temperature (Peterson 1996; McKechnie 2004;
66 Muir et al. 2007). In terrestrial ectothermic vertebrates, three important water-saving strategies are
67 metabolic depression (e.g., Muir et al. 2007), the reduction of trans-cutaneous evaporative water
68 loss (e.g., Anderson et al. 2017) and thermal depression (Ladyman and Bradshaw 2003). A
69 reduction of resting metabolism following chronic water deprivation is expected to reduce
70 respiratory water loss because metabolism scales linearly with ventilation rate (Woods and Smith
71 2010; Dupoué et al. 2017a). Another water conservation mechanism involves a decreased
72 permeability of the water barrier of the skin (Lillywhite 2006; McCormick and Bradshaw 2006).
73 Ectotherms may also respond to water restriction by thermal depression, that is, the lowering of
74 their preferred set temperature (Ladyman and Bradshaw 2003; Köhler et al. 2011; Anderson and
75 Andrade 2017). This is because higher body temperatures are associated with stronger respiratory
76 and cutaneous water loss rates during activity (e.g., Oufiero and Van Sant 2018; Senzano and
77 Andrade 2018). Previous studies focused mostly on one of these acclimation mechanisms, and their
78 prevalence or pace has been little investigated so far, especially in dry skinned ectotherms.
79 Physiological mechanisms involved in acclimation may be energetically or ecologically costly, and
80 cheaper alternative responses to cope with dehydration may consequently limit the implementation
81 of beneficial acclimation (Marais and Chown 2008; Huey et al. 2012). Relevant examples in
82 terrestrial animals are dispersal responses (i.e., behavioral flight) or changes in behavioral activity
83 and micro-habitat choice within the home range (i.e., behavioral flight responses consisting in
84 behavioral and physiological responses to cope with environmental changes while staying in the
85 same home range, inspired from Hertz et al. 1982), which may concur to increase water intake and
86 reduce water loss. In response to drought, individuals could enhance their exploration rate and

87 locomotor activity to find more suitable hydric environments (e.g., Rozen-Rechels et al. 2018), or
88 on the contrary decrease activity and exposure to drying conditions (e.g., less basking) to reduce
89 water loss rates (e.g., Lorenzon et al. 1999; Davis and DeNardo 2010; Kearney et al. 2018).
90 Individuals can also select cooler and wetter micro-habitats (Guillon et al. 2013; Dupoué et al.
91 2015b; Pintor et al. 2016). Such flight and fight behavioral strategies are not mutually exclusive and
92 might take place relatively quickly, within hours or days after exposure to chronic water stress if
93 environmental conditions are conducive to behavioral plasticity (Huey et al. 2003; Rozen-Rechels
94 et al. 2018). According to the “Bogert effect” (Bogert 1949; Huey et al. 2003; Marais and Chown
95 2008), behavioral flexibility buffers environmental variations to which individuals are exposed and
96 should consequently reduce the benefits of physiological acclimation (i.e. “behavioral inertia”
97 evolutionary scenario, see (Huey et al. 2003; Muñoz and Losos 2018). Unfortunately, quantification
98 of joint behavioral and physiological acclimation responses to chronic water stress are rare and the
99 relative importance of each response mechanism is yet to be elucidated.

100 One intuitive alternative to the Bogert effect is that each trait response follows a distinct pace
101 whereby some physiological and behavioral responses come first, followed by other physiological
102 and behavioral adjustments. For example, physiological models of chronic stress responses in
103 vertebrates, such as the allostatic model (McEwen and Wingfield 2003) or the reactive scope model
104 (Romero et al. 2009), predict non-linear dynamics of behavioral, physiological and life history traits
105 when individuals are exposed to a chronic stressor. In these models, behavioral adjustments, being
106 less energetically costly, are expected on the short-term and facilitate the activation of an
107 emergency state. Examples of the kinetics of behavioral and physiological responses to chronic
108 stressors support this hypothesis (Timmerman and Chapman 2004; Romero and Wingfield 2015).
109 To our knowledge, no study to date has examined the kinetics of plastic responses to a chronic

110 water deprivation in terrestrial animals, including potential long-lasting effects and fitness
111 responses.

112 In this study, we investigated temporal patterns of behavioral and physiological adjustments during
113 and after a chronic water restriction in the European common lizard, *Zootoca vivipara* (Lacertidae,
114 Lichtenstein, 1823). This species is widespread in Eurasia and depends on cold wetlands and
115 permanent access to free standing water for demographic persistence (Lorenzon et al. 2001; Dupoué
116 et al. 2017b). Acute or chronic exposure to sub-lethal water deprivation and reductions in water
117 availability in the habitat are associated with dehydration and physiological stress (Dupoué et al.
118 2017c, 2018a), a reduction in behavioral activity and body growth (Lorenzon et al. 1999), and some
119 behavioral flight responses (Rozen-Rechels et al. 2018). In addition, the potential reversal of
120 population growth to population decline is associated with increased habitat dryness and thermal
121 stress (Lorenzon et al. 2001; Dupoué et al. 2017b). Here, we compared morning activity patterns,
122 standard exploration behavior, thermal preferences, resting metabolic rates and standard water loss
123 rates from yearling lizards exposed to a 2-months water restriction period with those of yearlings
124 that had access to water *ad libitum*. During water restriction in the laboratory and after release in
125 outdoor enclosures, lizards were given opportunities to adjust their behavior. We predicted that a
126 chronic water restriction in this age class, characterized by a limited dispersal behavior and fast
127 growth linked with sexual maturation, will induce immediate water conservation behavioral
128 responses (shifts in activity patterns and less exploratory behaviors) together with delayed
129 physiological responses, including thermal and metabolic depression. We further predicted delayed
130 but potentially long-lasting reduction of standard water loss rates, especially from the skin, as a
131 consequence of water restriction.

132 **Material and Methods**

133 *Study species, sampling and rearing conditions*

134 All common lizards were captured from captive populations maintained in 100 m² outdoor
135 enclosures at the CEREEP-Ecotron IleDeFrance, France (48° 17' N, 2° 41' E). Enclosures include a
136 natural vegetation layer with permanent access to free water and piles of rocks and logs for basking
137 and sheltering. Ground and avian predation are avoided with plastic walls and nets. The density of
138 the populations are similar (23 ± 5 (SD) individuals, approximately 67% of yearlings). In these
139 enclosures, hibernation usually lasts from October to February-March and mating season starts
140 upon emergence of female individuals around March-April and lasts around 2 to 4 weeks. In our
141 study populations, age at maturation ranges from 1 (yearlings) to 2 years old.

142 In May 2016, 100 sexually immature yearlings (57 females and 43 males) were captured by hand in
143 10 enclosures, identified by their unique toe-clip, and measured for snout-vent length (SVL \pm 1
144 mm) and body mass ($M_b \pm$ 1 mg). Lizards were then housed in individual terraria (18 × 11 × 12 cm)
145 with a shelter and sterilized peat soil as substratum. A basking site (around 35°C), created using a
146 25W bulb above the shelter, and light from a UVB 30W neon (Reptisun 10.0, white light) were
147 available for each terrarium 9 hours a day (from 09:00 to 18:00). Terraria were located in a
148 temperature-controlled room (23°C from 09:00 to 18:00, 16°C at night). In this set-up with a
149 thermal gradient and some microhabitat heterogeneity, lizard express a range of typical field
150 behaviors such as locomotor activity, basking and hiding in the shelter or in the soil. We therefore
151 expected that these conditions allowed for behavioral responses such as changes in daily activity
152 profiles, propensity of basking and hiding behaviors, or differences in body temperature.

153 Individuals were fed three times a week with 300 ± 10 mg of live house crickets (*Acheta*
154 *domestica*). Under normal housing conditions, water was available *ad-libitum* in a petri dish and
155 sprayed on one of the walls of the terrarium three times a day (09:00, 13:00 and 17:00). Individuals
156 were released in enclosures on August 1st 2016. Individuals shared the enclosure with adults and
157 newly born juveniles (same density conditions in all enclosures: 41 ± 3 (SD), 72% of juveniles).

158 From September 12th to September 16th 2016, corresponding to the late summer activity season, we
159 recaptured as many individuals as we could (33 males and 40 females). We recaptured again all
160 survivors (recapture rate close to 100%, Le Galliard et al. 2004) during the next reproductive season
161 from May 15th to May 27th 2017 (32 males and 34 females). Non-recaptured individuals are
162 supposed to have not survived.

163 *Water restriction manipulation*

164 After acclimation of all individuals to standard housing conditions in late May 2016, we randomly
165 assigned lizards to two experimental conditions of water availability (Lorenzon et al. 1999; Dupoué
166 et al. 2018a; Rozen-Rechels et al. 2018) with a balanced sample of 29 females and 21 males in the
167 water-restricted treatment and 28 females and 22 males in the control treatment. In the water-
168 restricted treatment, individuals were sprayed only once a day at 09:00, and the Petri dish providing
169 water *ad libitum* was removed. These conditions mimic habitats in which water is only available
170 with the morning dew during summer (Lorenzon et al. 1999, 2001; Dupoué et al. 2017b). This
171 experimental water restriction has already been implemented in past studies during shorter periods
172 from a few days to two weeks. This protocol is sub-lethal but is known to enhance physiological
173 responses, although it is relatively less clear in yearlings compared to adults (Dupoué et al. 2018a).
174 It also enhances trans-generational and delayed effects in reproductive females (Dupoué et al.
175 2018a; Rozen-Rechels et al. 2018).

176 In the control treatment, water conditions remained similar to the normal ones described above.
177 These conditions mimic habitats with permanent access to water such as peat bogs or streams.
178 Water restriction treatment lasted from June 10th 2016 to July 31st 2016, which is equivalent to a
179 chronic early summer drought in natural populations (Dupoué et al. 2017b, 2018b). After that,
180 lizards were released in enclosures corresponding to common garden conditions. This would allow
181 us to test if effects of water restriction under laboratory conditions last in time. In case we did not

182 find any, this set-up would allow us to test if individuals have different life-history trajectories
183 depending on the treatment they experienced in laboratory conditions. Individuals were split
184 randomly in 5 groups of 20 individuals each (10 control individuals and 10 water restricted
185 individuals) so that each group was measured on one day, (5 days total for one measurement
186 session). We defined 5 standard measurement sessions of behavioral and physiological data from
187 June 4th 2016 to June 8th 2016 (week 0, before water restriction), from July 2nd to July 6th (week 4,
188 one month later), from July 25th to July 29th (week 7, two months later and before release in outdoor
189 enclosures), in September 2016 and in May 2017. The chronology of the measurements is
190 summarized in Figure 1.

191 *Body mass and size measurements*

192 Body mass (M_b) provides a good indicator of hydration state in reptiles (Lillywhite et al. 2012;
193 Dupoué et al. 2015a, 2018a). Change in M_b (ΔM_b) was calculated as the difference between initial
194 M_b at the onset of experiment (week 0) and M_b which was focally measured all over the water
195 restriction (weeks 1, 2, 4, 6, 7 and 8). In order to standardize measurements, M_b was always
196 measured two days after a feeding event. To assess growth of structural size, snout-vent length
197 (SVL) was also measured in weeks 0 and 4 as well as in September 2016 and in May 2017.
198 Structural growth rate was then defined as the difference in SVL between two measurements
199 sessions divided by the number of days between those sessions. We calculated early growth during
200 the laboratory experiment (growth between week 0 and week 4; $\Delta SVL_{\text{experiment}}$), summer growth
201 (growth between week 4 and September 2016; $\Delta SVL_{\text{summer}}$), and annual growth (growth between
202 September 2016 and May 2017; $\Delta SVL_{\text{annual}}$).

203 *Behavioral tests*

204 We measured individual behavior at all standard sessions when lizards were in post-absorptive state.

205 Emergence time and basking effort in the home cage

206 We scored lizards' behavior in their home cage during weeks 0, 1, 4 and 7. These measurements
207 were made for all 100 individuals in the same day. From 08:30 to 12:00, an observer recorded the
208 lizards' behavior in their cage every 15 mins (15 record per trial) with a minimum disturbance.
209 Behavioral items included: lizard hidden and inactive, lizard basking under the heat source, and any
210 other active behaviors (e.g., moving, scratching the soil). From this, we calculated emergence time
211 (first time the individual was seen active) and basking effort (proportion of observations seen
212 basking).

213 Exploration behavior in a neutral arena

214 In a temperature-controlled room at 25°C, individuals were tested for their exploration behavior in
215 30-min tests between 09:00 and 11:30 in a neutral arena. Neutral arenas (44.5 × 24 × 26 cm)
216 contained a layer of white sand as substrate and were warmed with a 40W light bulb heated basking
217 point. Two UVB 30W neon tubes were suspended above the arena to homogenize light conditions
218 and avoid phototropism. During the first 10 mins, individuals were allowed to habituate to the arena
219 and recover from handling stress (Rozen-Rechels et al. 2018). We then recorded their behaviors
220 using webcams set at 5 images per sec for 20 mins. Prior to each trial, sand layer was dried and
221 sterilized at 150°C for 2 hours in a stove and arenas were washed between each trial in order to
222 suppress conspecifics' odors from the arena. We extracted 12 behavioral units related to exploration
223 (detailed in Appendix 1) from the videos using the same protocol of image analysis as previously
224 described (Rozen-Rechels et al. 2018). We then used a multivariate analysis to calculate a
225 composite exploration score for each individual within each session (see Appendix 1 and Table A1
226 and (Rozen-Rechels et al. 2018) for the complete analytical procedure).

227 Thermal preferences in a neutral arena

228 We measured thermal preferences in a neutral thermal gradient (80 × 15 × 20 cm) filled with a
229 substratum of dried and sterilized peat soil. We placed a plate of wood on the ground (for basking

230 and hiding) and installed a 40W heat bulb 15 cm above the ground at the warm side of the gradient
231 ($49.1 \pm 6.7^\circ\text{C}$). The cold side of the gradient was maintained at ambient temperature ($25.5 \pm 1.9^\circ\text{C}$).
232 A UVB 30W neon provided white, natural light conditions above each thermal gradient. Heat bulbs
233 were turned on at 12:00 for one hour before individuals were randomly placed in one out of 20
234 thermal gradients for habituation for 30 mins. Every 20 mins until 17:10 (12 measurements), skin
235 surface temperature (T_p) of lizards was measured at the same focal distance (ca. 30cm, i.e.
236 measurement in a 6mm diameter circle, that we were able to assess thanks to three lasers indicating
237 the centre of the circle and the diameter of the spot when measuring the temperature) using an
238 infrared thermometer (Raytek, Raynger MX2). This method allows temperature measurements
239 without handling lizards, which are strongly correlated with core temperatures, and provides an
240 accurate measure of preferred body temperature (Bucklin et al. 2010; Artacho et al. 2013). We
241 analyzed preferred temperature in comparison to initial preferred temperatures at the onset of the
242 experiment. To do so, we calculated the difference between each skin surface temperature
243 measurement and the measurement done on the same individual and the same time of the day at
244 week 0, hereafter referred to as ΔT_p .

245 *Metabolic rate and water losses*

246 Resting metabolic rate (RMR) and total evaporative water loss (TEWL) were measured overnight at
247 25°C when individuals are normally inactive. We measured oxygen consumption (VO_2) using
248 closed-system respirometry methods as previously described (Foucart et al. 2014) and we measured
249 TEWL concomitantly. We weighed the lizards (M_{b1}) and let them acclimate at 25°C for 3h in late
250 afternoon. After collecting a baseline air sample with adapted locked syringes at the onset of trial
251 (two 140 mL syringes, Medtronic Monoject Luer Lock), we closed the jar and placed the lizards for
252 12h in a controlled climatic chamber (Aqualytic TC255). We collected a final air sample from each
253 container and weighted the lizard a second time (M_{b2}) to estimate body mass loss ($M_{b2}-M_{b1}$).

254 Oxygen proportion in air samples was determined using an O₂ analyzer (FOXBOX, Sable Systems,
255 Las Vegas, NV, USA). Using an infusion pump (KDS 200, KD Scientific Inc., Holliston, MA,
256 USA), air was sent at a constant flow (60 mL.min⁻¹), dried in a column of Drierite, and in the O₂
257 analyzer, which was calibrated before each trial using outdoor air. Oxygen consumption (mL.h⁻¹)
258 was calculated as the difference between final O₂ and baseline O₂ multiplied by the exact chamber
259 volume (mL, calculated by measuring the mass of water it can contain) divided by trial duration (h).
260 We also measured CO₂ production (mL.h⁻¹) which was highly correlated to oxygen consumption
261 (R² = 0.98). We calculated the respiratory exchange ration defined as the ratio between CO₂
262 produced and O₂ consumed which was not affected by the water restriction (F_{1,98} = 0.04, p = 0.83;
263 ~0.7 in yearlings, ~0.8 in adults).

264 TEWL (mg.h⁻¹) was calculated from body mass loss divided by the time between M_{b1} and M_{b2}
265 measures, a method previously validated for this species (Dupoué et al. 2017c). RMR and TEWL
266 were measured in weeks 0, 4 and 7 during the laboratory experiment (the night prior to behavioral
267 measurements) as well as in May 2017 at the end of the study (the night after behavioral
268 measurements in order to ensure the same post-prandial digestive state of individuals two days after
269 recapture). For each individual, as we expected physiological traits to change with time and within
270 individuals, we calculated the changes in RMR (Δ RMR) and TEWL (Δ TEWL) as the difference
271 between individual record of the measurement session and the one at week 0.

272 *Statistical analyses*

273 Statistical analyses were performed with R version 3.4.1 (R Core Team 2018). In most analyses,
274 trait variation was analyzed with ANCOVAs using linear models and backward model selection
275 from a full model including the three-way interaction between treatment, sex and time
276 (measurement session). In the cases of Δ M_b, Δ SVL, Δ RMR and Δ TEWL, initial value at the onset
277 of the experiment was included as fixed effect. For Δ RMR and Δ TEWL, we included individual M_b

278 as a fixed effect in the ANCOVA to account for body mass scaling. Metabolism is indeed positively
279 related to body mass (Gillooly et al. 2001) and water loss rates depend on the whole-body area and
280 respiratory exchanges, both scaling with body mass. For $\Delta TEWL$, we also considered the presence
281 of feces as fixed, categorical effect to account for corresponding mass loss due to defecation. We
282 analyzed separately treatment effects during the laboratory experiment and delayed effects after
283 recapture in outdoor enclosures, given the substantial changes in sample sizes and individual
284 contribution between sessions. In the first three measurement sessions, we used linear mixed
285 models (*lme* function from “nlme” package, Pinheiro et al. 2007) and set individual identity as a
286 random factor to account for repeated measurements between sessions. Sessions in September 2016
287 and in May 2017 were investigated separately (one observation per individuals in each). We
288 included enclosure identity as a random factor to account for among-enclosures variability. In all
289 cases, the best model fit was determined using log-likelihood ratio tests (LRTs) and elimination of
290 non-significant factors at 5% critical rate.

291 We analyzed variation in ΔT_p in a different way because body temperature was recorded
292 sequentially during the day and we wanted to account for potential daytime trends in thermal
293 preferences (Artacho et al. 2013). For laboratory sessions, the full mixed effect model thus included
294 a two-way interaction between treatment and sex and a three-way interaction between treatment,
295 measurement session and time of day. September 2016 and May 2017 thermal preferences were
296 analyzed in separate similar models.

297 We estimated whether or not the treatment impacted emergence time using Cox survival model
298 from the package “coxme” (Therneau 2018). By analogy to a survival analysis, at each observation,
299 an individual would emerge with a probability P (“die”). If we did not have any observation of the
300 individual at the end of the experiment, we consider that it never went out (“survived”). The model
301 estimates the mean time of emergence. We analyzed the proportion of time spent basking by testing

302 the influence of all possible two-ways interactions between treatment, session, and sex on the
303 number of basking events using generalized linear mixed models from the package “lme4”,
304 considering that the number of basking events is the result of a binomial probability distribution.
305 We controlled the family-wise error rate due to multiple comparisons by calculating the corrected
306 threshold of significance with the Holm-Bonferroni method based on the number of tests m realized
307 at each step of the study (laboratory experiment: $m = 8$, after one month in common gardens: $m = 3$,
308 after winter in common gardens: $m = 5$). In each case, p-values of significance of the treatment
309 effect were ranked according from the lowest to the highest and the new threshold was calculated as

310
$$\alpha_{Holm} = \frac{0.05}{m+1-rank} \quad (\text{Holm 1979}).$$

311

312 **Results**

313 Initial M_b , SVL, T_b , RMR and TEWL values (week 0, before the beginning of the water restriction)
314 are shown in Table 1.

315 *Effects of water restriction on body mass and growth*

316 Body mass increased on average during the laboratory manipulation ($F_{5,490} = 288.9, p < 0.0001$) but
317 with different temporal trends between treatments (treatment: $F_{1,96} = 28.5, p < 0.0001$; treatment \times
318 time: $F_{5,490} = 10.6, p < 0.0001, \alpha_{Holm} = 0.006$). Water-restricted yearlings had a decreased ΔM_b one
319 week and two weeks after the beginning of the experiment, but this difference vanished in
320 subsequent sessions (Fig. 2A). In addition, males had a lower ΔM_b than females ($F_{1,96} = 4.47, p =$
321 0.037) and ΔM_b was negatively correlated to initial body mass ($F_{1,96} = 8.2, p = 0.005$). During the
322 first month, water-restricted individuals had also a marginally lower $\Delta SVL_{\text{experiment}}$ compared to
323 control individuals ($F_{1,96} = 4.235, p = 0.042, \alpha_{Holm} = 0.08$; see Figure 2B). The treatment did not
324 affect $\Delta SVL_{\text{summer}}$ and $\Delta SVL_{\text{annual}}$ (all $p > 0.59$, all $\alpha_{Holm} < 0.05$; see Figure 2B). Males had

325 consistently lower Δ SVL than females (all $p < 0.007$) and Δ SVL was negatively correlated to initial
326 body size (all $p < 0.002$).

327 *Emergence and basking effort in home cage*

328 Time since the beginning of the experiment significantly explained variation in emergence behavior
329 ($\chi^2 = 107.7$, $df = 3$, $p < 0.0001$), but treatment had no effect ($\chi^2 = 1.01$, $df = 1$, $p = 0.31$, $\alpha_{\text{Holm}} =$
330 0.01 , see Figure 3A). Basking effort was also influenced by time since the beginning of the
331 experiment but in interaction with water restriction treatment (treatment \times time: $\chi^2 = 14.2$, $df = 3$, p
332 $= 0.003$, $\alpha_{\text{Holm}} = 0.007$; Fig. 3B) and in interaction with sex (time \times sex: $\chi^2 = 10.3$, $df = 3$, $p = 0.02$).
333 One week after the beginning of the experiment, control yearlings spent significantly more time
334 basking than water-restricted individuals (Fig. 3B), whereas differences between treatments later
335 disappeared (all $p > 0.25$). In addition, no sex differences existed during the first month (all $p >$
336 0.17) but males basked significantly less than females at the end of the experiment ($z = -3.18$; $p =$
337 0.001).

338 *Exploration score in neutral arenas*

339 Exploration score was lower on average one month and two months after the start of the experiment
340 than at the beginning of the study ($F_{2,196} = 10.84$, $p < 0.0001$) but there was no effect of water
341 restriction on exploration ($F_{1,98} = 0.13$, $p = 0.72$; treatment \times time: $F_{4,194} = 0.91$, $p = 0.40$, $\alpha_{\text{Holm}} =$
342 0.016 ; see Figure 4A). One month after the end of the experiment (September 2016), exploration
343 score was influenced by a two-way interaction between treatment and sex ($F_{1,69} = 5.7$, $p = 0.02$,
344 $\alpha_{\text{Holm}} = 0.017 - 0.025$). Control females and males did not differ in their exploration rate ($t_{69} = 1.57$;
345 $p = 0.12$), nor did control and water-restricted females ($t_{69} = 1.78$; $p = 0.08$). However, water-
346 restricted males explored their environment significantly less than control males ($t_{69} = -2.38$; $p =$
347 0.02). One year later, water restriction treatment had no effect on exploration score ($F_{1,64} = 0.02$, $p =$
348 0.9 , $\alpha_{\text{Holm}} = 0.05$).

349 *Thermal preferences*

350 Preferred body temperature did not change significantly during the laboratory study ($F_{1,2273} = 1.73$,
351 $p = 0.19$) nor between treatments ($F_{1,98} = 0.26$, $p = 0.61$; treatment \times time: $F_{1,2273} = 0.23$, $p = 0.63$,
352 $\alpha_{\text{Holm}} = 0.05$; see Figure 4B). One month after the end of the laboratory study, intra-individual ΔT_p
353 was influenced by a two-way interaction between sex and treatment ($F_{1,69} = 5.74$, $p = 0.02$, $\alpha_{\text{Holm}} =$
354 $0.017 - 0.025$; see Figure 4B) as well as by time of day ($F_{1,802} = 7.94$, $p = 0.005$). At this stage,
355 water restriction did not change thermal preferences of females ($t_{69} = 0.38$, $p = 0.47$) but decreased
356 the thermal preferences of males ($t_{69} = -2.39$, $p = 0.02$). One year later, we found no effect of
357 treatment conditions ($F_{1,64} = 0.26$, $p = 0.61$, $\alpha_{\text{Holm}} = 0.017$).

358 *Standard metabolism and water losses*

359 During the laboratory experiment, ΔRMR was negatively related to initial values ($F_{1,97} = 21.3$, $p <$
360 0.0001) but independent of treatment and session groups (treatment: $F_{1,97} = 2.23$, $p = 0.14$;
361 treatment \times time: $F_{1,97} = 0.25$, $p = 0.62$, $\alpha_{\text{Holm}} = 0.025$) and sex (treatment \times sex: $F_{1,95} = 1.15$, $p =$
362 0.29). ΔRMR measured from metabolic rate at recapture during the next reproductive season was
363 marginally lower in water-restricted yearlings ($F_{1,59} = 3.66$, $p = 0.061$, $\alpha_{\text{Holm}} = 0.0125$; see Figure
364 5A). In addition, changes in evaporative water loss (ΔTEWL) decreased during the laboratory
365 experiment ($F_{1,96} = 28.5$, $p < 0.0001$; see Figure 5B) independently from the treatment (treatment:
366 $F_{1,97} = 1.64$, $p = 0.20$; treatment \times time: $F_{1,96} = 0.75$, $p = 0.39$, $\alpha_{\text{Holm}} = 0.01$). ΔTEWL at recapture
367 during the next reproductive season was significantly explained by a two-way interaction between
368 sex and treatment ($F_{1,48} = 4.35$, $p = 0.04$, $\alpha_{\text{Holm}} = 0.01$; see Figure 5B). Water-restricted females had
369 lower ΔTEWL than control females ($t_{48} = -2.69$, $p = 0.01$), whereas no treatment effect was detected
370 in males ($t_{48} = 0.33$, $p = 0.75$). In all cases, ΔTEWL was negatively correlated to the initial TEWL
371 value (all $p < 0.0001$). ΔTEWL during the laboratory experiment were positively correlated to body
372 mass ($p = 0.0001$).

373 **Discussion**

374 We exposed yearling lizards to a sub-lethal, chronic water restriction episode lasting almost two
375 months during which lizards could substantially buffer dehydration with behavioral shifts. Our data
376 revealed acclimation kinetics of the behavioral and physiological responses to water deprivation.
377 The kinetics of these behavioral and physiological responses were generally in agreement with our
378 initial predictions since flexible behavioral changes came first, followed by delayed sex-specific
379 acclimation responses in thermal physiology (thermal depression) and standard evaporative water
380 loss. However, in contrast to our expectations, we did not find any short-term thermoregulation
381 adjustments in our laboratory setting. Despite significant short-term negative effects of water
382 deprivation on body mass change and marginal negative effects on body size growth during the first
383 weeks of manipulation, which could suggest short-term physiological adjustments that we did not
384 measure, the annual life-history strategy of sub-adult lizards from the two treatments did not differ
385 for total annual size growth, nor for annual survival (see Appendix 2) and future reproduction of
386 females (see Appendix 3). Differences in body mass change might be the consequence of
387 differences in muscle volume (storing water and sources of metabolic water) or differences in
388 hydration state. We thus conclude that phenotypic plasticity in *Z. vivipara* compensates to some
389 degree for the homeostatic load imposed by dehydration risk. This supports the beneficial
390 acclimation hypothesis, at least in our laboratory setting (Huey et al. 1999, Kelly et al. 2012) and
391 the kinetics is consistent with previous studies on acclimation to other stressors. In the sailfin molly
392 *Poecilia latipinna*, fishes exposed to extreme hypoxia display an immediate physiological response
393 (increased ventilation) and an immediate behavioral response (increased aquatic surface respiration)
394 that decreases over time as gradual acclimation processes take place (e.g., changes in hemoglobin
395 concentration, Timmerman and Chapman 2004). Testing whether observed adjustments give

396 significant advantages in the water-restricted environment would however be necessary to strictly
397 conclude that we observe beneficial acclimation.

398 Immediate behavioral responses to water restriction included a sharp reduction of basking activity,
399 which was paralleled by a reduction of body size growth. In ectothermic species, an increase of
400 thermoregulation effort and high body temperatures closed to thermal preferences would increase
401 water loss rates compared to resting conditions. A reduction of basking effort can therefore be
402 interpreted as a water-conservation strategy (Lourdais et al. 2017). This reduction to conserve water
403 can however conflict with energy intake and allocation to structural growth or reserves (Adolph and
404 Porter 1993; Niewiarowski 2001). Similar results were observed in a previous comparative study
405 with wild-caught yearlings from dry and humid natural habitats: lizards were also less active after a
406 one-month long water scarcity and grew slower in the laboratory, and the spatial variation in growth
407 rates observed between wet and dry natural habitats paralleled the results of the laboratory study
408 (Lorenzon et al. 1999, 2001). During the same time period of our laboratory manipulation, we
409 found no plastic changes for thermal preferences, resting metabolism and standard water loss rates,
410 which were expected considering recent studies on other ectotherms (Muir et al. 2007; Anderson
411 and Andrade 2017; Anderson et al. 2017). This may suggest a Bogert effect, as short-term changes
412 in a key behavioral trait, here basking activity, compensated for the environmental changes induced
413 by water restriction and protected individuals from its deleterious effects. We, however, cannot
414 exclude that other physiological responses we did not measure in our study were stimulated by
415 water deprivation (e.g., shift from carbohydrates to protein catabolism to increase metabolic water
416 production, see Bruschi et al. 2016 or hormone-induced changes in renal function and
417 osmoregulation to save water lost in urine and feces, see McCormick and Bradshaw 2006). In
418 addition, some traits may not respond because yearling lizards generally have lower water
419 requirements than adults and are therefore less sensitive to water restriction than older lizards

420 (Dupoué et al. 2018a). In support of this hypothesis, no difference in basking activity was detected
421 after the first month of water deprivation and therefore no Bogert effect was expected. Yet, we
422 found no acclimation response for other behavioral or physiological traits during the second half of
423 the laboratory experiment and water-restricted individuals caught up their growth delay. This
424 suggests that fast and reversible physiological adjustments took place.

425 We predicted delayed responses for some physiological traits but did not expect specifically delayed
426 sex-specific effects of water restriction on exploration behavior and thermal preferences at the end
427 of summer, as observed in male yearlings in this study. Based on a previous study of water-
428 dependent exploration behavior (Rozen-Rechels et al. 2018) and outcomes of thermal acclimation
429 experiments (Clusella-Trullas and Chown 2014), we instead assumed that exploration and, to some
430 extent, thermal preferences would be more flexible and would respond earlier following water
431 deprivation (e.g., (Rozen-Rechels et al. 2018). The delayed responses found in males are concordant
432 with a water conservation strategy (i.e., fight response, aiming at buffering the effects of the
433 environmental constraints on the physiology) since lower exploration scores and thermal depression
434 imply lower rates of water loss from locomotor activities and from trans-cutaneous evaporation
435 (Lourdais et al. 2017). The high sensitivity of male common lizards to weather conditions has
436 already been demonstrated in previous studies (Le Galliard et al. 2006; Dupoué et al. 2018a), and is
437 also consistent with recent results showing lower growth rate in juvenile males exposed to
438 abnormally hot summers (Dupoué et al. 2019). Interestingly, during the time period when yearlings
439 were maintained in enclosures in late summer 2016, the study site experienced an abnormal heat
440 wave characterized by very low precipitation in August and a warmer and drier month of September
441 than average (Meteo France: +0.9°C for August average temperature , +2.4°C for September and
442 rain deficits of 60% in August and >30% in September compared to previous years). Thus, one
443 possibility is that the delayed acclimation responses were caused by “a facilitation process”,

444 whereby a new environmental stressor (here a hot and dry summer) causes a higher than expected
445 stress response in animals that were exposed to past stressful conditions (here the laboratory water
446 deprivation, see (Trompeter and Langkilde 2011) for an example of facilitation response to
447 predation stress in a lizard). To test this hypothesis, future studies independently manipulating the
448 environmental conditions experienced by male yearlings in two successive later stages are needed.
449 Another negative delayed effect of water restriction was found for standard water loss rates in
450 female lizards the year after the laboratory experiment, when females had reached sexual maturity
451 and were all gravid. Variations of TEWL were positive, which can be explained by the higher mass
452 of individuals at the adult stage. Water-restricted females had however lower change of TEWL than
453 control ones. This effect is weak and need further confirmation and should therefore be cautiously
454 interpreted but is also consistent to a water conservation strategy. Since metabolic rates and body
455 size were not different between water restricted and control females during this time period, we
456 assume that these differences might reflect changes in cutaneous water loss rates and therefore
457 probably cutaneous resistance to water loss between water restricted and control females. This
458 explanation is also supported by the robustness of the result of our statistical analyses of TEWL to
459 inclusion of Δ RMR (changes in metabolism, hence ventilation rates) as a covariate, indicating that
460 water restriction changes total water loss rates even after corrections for differences in respiratory
461 activity (see Table A2). This slight change in standard water loss rates supports our initial prediction
462 of a water-conservation strategy. Other studies that have tested for plastic changes in cutaneous
463 water loss rates generally found that lizards or snakes acclimated to drier environments or water
464 restricted environments can adjust cutaneous water loss rates within days or weeks of exposure
465 (Kobayashi et al. 1983; Kattan and Lillywhite 1989; Moen et al. 2005 but see Neilson 2002;
466 Gunderson et al. 2011). Reduction of cutaneous water loss rates is also well documented in anurans
467 exposed to chronic water deprivation (Anderson et al. 2017; Senzano and Andrade 2018). It is

468 generally accepted that the skin resistance to water loss depends mainly on the intracellular lipid
469 content and ultrastructure of the mesos layer of the epidermis (reviewed by Lillywhite 2006).
470 Further investigations of skin properties would be necessary to know what are the mechanisms
471 underlying delayed acclimation patterns in the common lizard. This female-specific acclimation
472 response of skin properties may be explained by the stronger reliance of females on water during
473 gestation and embryonic development in this viviparous species (Dupoué et al. 2015a, 2018a).
474 In conclusion, the results of our multivariate analysis of the physiological and behavioral responses
475 to chronic water deprivation in the common lizard are broadly speaking consistent with general
476 predictions about the kinetics of a long-term acclimation process and stress response (Huey et al.
477 1999; Romero and Wingfield 2015). According to this framework, less costly and more flexible
478 adjustments of daily behavioral routines should occur first followed by delayed responses in
479 thermal preferences, metabolism and eventually cutaneous water loss rates (e.g., Timmerman and
480 Chapman 2004). Such dynamical changes may lead to an “emergency life history stage” (sensu
481 Wingfield et al. 1998) that promotes survival in the face of a challenging environment as seen in
482 this study where water restricted lizards had similar annual growth, survival and reproduction than
483 control lizards. Traits involved in acclimation responses were those most strongly linked to
484 thermoregulation (basking and thermal preferences), but we found no evidence of metabolic
485 depression, although this a widespread response to water restriction and energy intake reduction in
486 endotherms (Ruf and Geiser 2015) and dry-skinned ectotherm (Christian et al. 1996a, 1996b). Our
487 results are partly consistent with biophysical models predicting that hydroregulation responses
488 would rely mostly on evaporative water loss changes but also activity pattern changes, and not
489 metabolic depression (Pirtle et al. 2019). Surprisingly, we found sex-specific acclimation responses,
490 in line with previous findings (e.g., (Dupoué et al. 2018a), which could be explained by ecological
491 and physiological differences between males and females. Interestingly, males seem more

492 susceptible to water restriction than females as previous findings might suggest (Le Galliard et al.
493 2006; Dupoué et al. 2019). This study illustrates the complexity of acclimation responses to water
494 restriction in dry-skinned terrestrial ectotherms.

495 **References**

496 Adolph S.C. and W.P. Porter. 1993. Temperature, activity, and lizard life histories. *Am Nat* 142:273–
497 295.

498 Anderson R.C.O. and D.V. Andrade. 2017. Trading heat and hops for water: Dehydration effects on
499 locomotor performance, thermal limits, and thermoregulatory behavior of a terrestrial toad.
500 *Ecol Evol* 7:9066–9075.

501 Anderson R.C.O., R.P. Bovo, C.E. Eismann, A.A. Menegario, and D.V. Andrade. 2017. Not good,
502 but not all bad: dehydration effects on body fluids, organ masses, and water flux through the
503 skin of *Rhinella schneideri* (Amphibia, Bufonidae). *Physiol Biochem Zool* 90:313–320.

504 Artacho P., I. Jouanneau, and J.-F. Le Galliard. 2013. Interindividual variation in thermal sensitivity
505 of maximal sprint speed, thermal behavior, and resting metabolic rate in a lizard. *Physiol*
506 *Biochem Zool* 86:458–469.

507 Bates D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using
508 lme4. *Journal of Statistical Software* 67:1–48.

509 Bogert C.M. 1949. Thermoregulation in reptiles, a factor in evolution. *Evolution* 3:195–211.

510 Bruschi G.A., E.N. Taylor, and S.M. Whitfield. 2016. Turn up the heat: thermal tolerances of lizards
511 at La Selva, Costa Rica. *Oecologia* 180:325–334.

512 Bucklin S.E., G.W. Ferguson, W.H. Gehrmann, and J.E. Pinder. 2010. Use of remote laser sensing
513 equipment to measure surface temperature and to predict deep body temperatures of small
514 lizards in the field. *Herpetol Rev* 41:309–312.

515 Cheuvront S.N. and R.W. Kenefick. 2014. Dehydration: physiology, assessment, and performance
516 effects. *Compr Physiol* 4:257–285.

517 Christian K., B. Green, G. Bedford, and K. Newgrain. 1996a. Seasonal metabolism of a small,
518 arboreal monitor lizard, *Varanus scalaris*, in tropical Australia. *Journal of Zoology* 240:383–
519 396.

520 Christian K.A., A.D. Griffiths, and G.S. Bedford. 1996b. Physiological ecology of frillneck lizards
521 in a seasonal tropical environment. *Oecologia* 106:49–56.

522 Clusella-Trullas S. and S.L. Chown. 2014. Lizard thermal trait variation at multiple scales: a review.
523 *J Comp Physiol B* 184:5–21.

524 Davis J.R. and D.F. DeNardo. 2010. Seasonal Patterns of Body Condition, Hydration State, and
525 Activity of Gila Monsters (*Heloderma suspectum*) at a Sonoran Desert Site. *J Herpetol*
526 44:83–93.

527 Dray S., A.-B. Dufour, and others. 2007. The ade4 package: implementing the duality diagram for
528 ecologists. *Journal of statistical software* 22:1–20.

529 Dupoué A., F. Brischoux, F. Angelier, D.F. DeNardo, C.D. Wright, and O. Lourdais. 2015a.
530 Intergenerational trade-off for water may induce a mother–offspring conflict in favour of
531 embryos in a viviparous snake. *Funct Ecol* 29:414–422.

532 Dupoué A., F. Brischoux, and O. Lourdais. 2017a. Climate and foraging mode explain interspecific
533 variation in snake metabolic rates. *Proc Roy Soc B-Biol Sci* 284:20172108.

534 Dupoué A., J.-F.L. Galliard, R. Josserand, D.F. DeNardo, B. Decencière, S. Agostini, C. Haussy, et
535 al. 2018a. Water restriction causes an intergenerational trade- off and delayed mother–
536 offspring conflict in a viviparous lizard. *Funct Ecol* 32:676–686.

537 Dupoué A., O. Lourdais, S. Meylan, F. Brischoux, F. Angelier, D. Rozen- Rechels, Y. Marcangeli,
538 et al. 2019. Some like it dry: Water restriction overrides heterogametic sex determination in
539 two reptiles. *Ecology and Evolution* 9:6524–6533.

540 Dupoué A., A. Rutschmann, J.F. Le Galliard, J. Clobert, F. Angelier, C. Marciau, S. Ruault, et al.
541 2017b. Shorter telomeres precede population extinction in wild lizards. *Sci Rep* 7.

542 Dupoué A., A. Rutschmann, J.F. Le Galliard, J. Clobert, P. Blaimont, B. Sinervo, D.B. Miles, et al.
543 2018b. Reduction in baseline corticosterone secretion correlates with climate warming and
544 drying across wild lizard populations. (K. Marske, ed.) *J Anim Ecol* 87:1331–1341.

545 Dupoué A., A. Rutschmann, J.F. Le Galliard, D.B. Miles, J. Clobert, D.F. DeNardo, G.A. Bruschi, et
546 al. 2017c. Water availability and environmental temperature correlate with geographic
547 variation in water balance in common lizards. *Oecologia* 185:561–571.

548 Dupoué A., Z.R. Stahlschmidt, B. Michaud, and O. Lourdais. 2015b. Physiological state influences
549 evaporative water loss and microclimate preference in the snake *Vipera aspis*. *Physiol*
550 *Behav* 144:82–89.

551 Eto E.C., P.C. Withers, and C.E. Cooper. 2017. Can birds do it too? Evidence for convergence in
552 evaporative water loss regulation for birds and mammals. *Proc Roy Soc B-Biol Sci*
553 284:20171478.

554 Field C.B., V. Barros, T.F. Stocker, and Q. Dahe. 2012. Managing the risks of extreme events and
555 disasters to advance climate change adaptation: special report of the intergovernmental
556 panel on climate change. Cambridge University Press.

557 Foucart T., O. Lourdais, D.F. DeNardo, and B. Heulin. 2014. Influence of reproductive mode on
558 metabolic costs of reproduction: insight from the bimodal lizard *Zootoca vivipara*. *J Exp*
559 *Biol* 217:4049–4056.

560 Gillooly J.F., J.H. Brown, G.B. West, V.M. Savage, and E.L. Charnov. 2001. Effects of size and
561 temperature on metabolic rate. *Science* 293:2248–2251.

562 Guillon M., G. Guiller, D.F. DeNardo, and O. Lourdais. 2013. Microclimate preferences correlate
563 with contrasted evaporative water loss in parapatric vipers at their contact zone. *Can J Zool*
564 92:81–86.

565 Gunderson A.R., J. Siegel, and M. Leal. 2011. Tests of the contribution of acclimation to geographic
566 variation in water loss rates of the West Indian lizard *Anolis cristatellus*. *J Comp Physiol B*
567 181:965–972.

568 Gunderson A.R. and J.H. Stillman. 2015. Plasticity in thermal tolerance has limited potential to
569 buffer ectotherms from global warming. *Proc Roy Soc B-Biol Sci* 282:20150401.

570 Hertz P.E., R.B. Huey, and E. Nevo. 1982. Fight versus flight: Body temperature influences
571 defensive responses of lizards. *Animal Behaviour* 30:676–679.

572 Huey R.B., D. Berrigan, G.W. Gilchrist, and J.C. Herron. 1999. Testing the adaptive significance of
573 acclimation: a strong inference approach. *Am Zool* 39:323–336.

574 Huey R.B., P.E. Hertz, and B. Sinervo. 2003. Behavioral drive versus behavioral inertia in
575 evolution: a null model approach. *Am Nat* 161:357–366.

576 Huey R.B., M.R. Kearney, A. Krockenberger, J.A. Holtum, M. Jess, and S.E. Williams. 2012.
577 Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and
578 adaptation. *Phil Trans R Soc Lond B* 367:1665–1679.

579 Kattan G.H. and H.B. Lillywhite. 1989. Humidity acclimation and skin permeability in the lizard
580 *Anolis carolinensis*. *Physiol Zool* 62:593–606.

581 Kearney M.R., S.L. Munns, D. Moore, M. Malishev, and C.M. Bull. 2018. Field tests of a general
582 ectotherm niche model show how water can limit lizard activity and distribution. *Ecol*
583 *Monogr* 88:672–693.

584 Kelly S.A., T.M. Panhuis, and A.M. Stoehr. 2012. Phenotypic plasticity: molecular mechanisms and
585 adaptive significance. *Compr Physiol* 2:1417–1439.

586 Kobayashi D., W.J. Mautz, and K.A. Nagy. 1983. Evaporative water loss: humidity acclimation in
587 *Anolis carolinensis* lizards. *Copeia* 1983:701.

588 Köhler A., J. Sadowska, J. Olszewska, P. Trzeciak, O. Berger-Tal, and C.R. Tracy. 2011. Staying
589 warm or moist? Operative temperature and thermal preferences of common frogs (*Rana*
590 *temporaria*), and effects on locomotion. *Herpetol J* 21:17–26.

591 Ladyman M. and D. Bradshaw. 2003. The influence of dehydration on the thermal preferences of
592 the Western tiger snake, *Notechis scutatus*. *J Comp Physiol B* 173:239–246.

593 Le Galliard J.-F., J. Clobert, and R. Ferriere. 2004. Physical performance and Darwinian fitness in
594 lizards. *Nature* 432:502–505.

595 Le Galliard J.-F., M. Massot, M.M. Landys, S. Meylan, and J. Clobert. 2006. Ontogenic sources of
596 variation in sexual size dimorphism in a viviparous lizard. *J Evol Biol* 19:690–704.

597 Leroi A.M., A.F. Bennett, and R.E. Lenski. 1994. Temperature acclimation and competitive fitness:
598 an experimental test of the beneficial acclimation assumption. *Proc Nat Acad Sci* 91:1917–
599 1921.

600 Lillywhite H.B. 2006. Water relations of tetrapod integument. *J Exp Biol* 209:202–226.

601 Lillywhite H.B., F. Brischoux, C.M. Sheehy, and J.B. Pfaller. 2012. Dehydration and drinking
602 responses in a pelagic sea snake. *Integr Comp Biol* 52:227–234.

603 Lorenzon P., J. Clobert, and M. Massot. 2001. The contribution of phenotypic plasticity to
604 adaptation in *Lacerta vivipara*. *Evolution* 55:392–404.

605 Lorenzon P., J. Clobert, A. Oppliger, and H. John-Alder. 1999. Effect of water constraint on growth
606 rate, activity and body temperature of yearling common lizard (*Lacerta vivipara*). *Oecologia*
607 118:423–430.

608 Lourdais O., A. Dupoué, M. Guillon, G. Guiller, B. Michaud, and D.F. DeNardo. 2017. Hydric
609 “costs” of reproduction: pregnancy increases evaporative water loss in the snake *Vipera*
610 *aspis*. *Physiol Biochem Zool* 90:663–672.

611 Marais E. and S.L. Chown. 2008. Beneficial acclimation and the Bogert effect. *Ecol Lett* 11:1027–
612 1036.

613 Marquis O., M. Massot, and J.F. Le Galliard. 2008. Intergenerational effects of climate generate
614 cohort variation in lizard reproductive performance. *Ecology* 89:2575–2583.

615 McCormick S.D. and D. Bradshaw. 2006. Hormonal control of salt and water balance in
616 vertebrates. *Gen Comp Endocr* 147:3–8.

617 McEwen B.S. and J.C. Wingfield. 2003. The concept of allostasis in biology and biomedicine.
618 *Horm Behav* 43:2–15.

619 McKechnie A.E. 2004. Partitioning of evaporative water loss in white-winged doves: plasticity in
620 response to short-term thermal acclimation. *J Exp Biol* 207:203–210.

621 Mitchell D., E.P. Snelling, R.S. Hetem, S.K. Maloney, W.M. Strauss, and A. Fuller. 2018. Revisiting
622 concepts of thermal physiology: Predicting responses of mammals to climate change. *J*
623 *Anim Ecol* 87:956–973.

624 Moen D.S., C.T. Winne, and R.N. Reed. 2005. Habitat-mediated shifts and plasticity in the
625 evaporative water loss rates of two congeneric pit vipers (Squamata, Viperidae,
626 *Agkistrodon*). *Evol Ecol Res* 7:759–766.

627 Muir T.J., J.P. Costanzo, and R.E. Lee. 2007. Osmotic and metabolic responses to dehydration and
628 urea-loading in a dormant, terrestrially hibernating frog. *J Comp Physiol B* 177:917–926.

629 Muñoz M.M. and J.B. Losos. 2018. Thermoregulatory behavior simultaneously promotes and
630 forestalls evolution in a tropical lizard. *Am Nat* 191:E15–E26.

631 Neilson K.A. 2002. Evaporative water loss as a restriction on habitat use in endangered New
632 Zealand endemic skinks. *J Herpetol* 36:342–348.

633 Niewiarowski P.H. 2001. Energy budgets, growth rates, and thermal constraints: toward an
634 integrative approach to the study of life- history variation. *Am Nat* 157:421–433.

635 Ostwald M.M., M.L. Smith, and T.D. Seeley. 2016. The behavioral regulation of thirst, water
636 collection and water storage in honey bee colonies. *J Exp Biol* 219:2156–2165.

637 Oufiero C.E. and M.J. Van Sant. 2018. Variation and repeatability of cutaneous water loss and skin
638 resistance in relation to temperature and diel variation in the lizard *Sceloporus consobrinus*.
639 *J Comp Physiol B* 188:671–681.

640 Peterman W.E. and R.D. Semlitsch. 2014. Spatial variation in water loss predicts terrestrial
641 salamander distribution and population dynamics. *Oecologia* 176:357–369.

642 Peterson C.C. 1996. Anhomeostasis: seasonal water and solute relations in two populations of the
643 desert tortoise (*Gopherus agassizii*) during chronic drought. *Physiol Zool* 69:1324–1358.

644 Pinheiro J., D. Bates, S. DebRoy, D. Sarkar, and R.C. Team. 2007. Linear and nonlinear mixed
645 effects models. R package version 3:57.

646 Pintor A.F.V., L. Schwarzkopf, and A.K. Krockenberger. 2016. Hydroregulation in a tropical dry-
647 skinned ectotherm. *Oecologia* 182:925–931.

648 Pirtle E.I., C.R. Tracy, and M.R. Kearney. 2019. Hydroregulation. A neglected behavioral response
649 of lizards to climate change? Pp. 343–374 in *Behavior of Lizards: Evolutionary and*
650 *Mechanistic Perspectives*. CRC Press.

651 R Core Team. 2018. R: language and environment for statistical computing. R Foundation for
652 Statistical Computing, 2005; Vienna, Austria.

653 Romero L.M., M.J. Dickens, and N.E. Cyr. 2009. The reactive scope model — A new model
654 integrating homeostasis, allostasis, and stress. *Horm Behav* 55:375–389.

655 Romero L.M. and J.C. Wingfield. 2015. Tempests, poxes, predators, and people: stress in wild
656 animals and how they cope. Oxford University Press.

657 Rozen-Rechels D., A. Dupoué, S. Meylan, B. Decencière, S. Guingand, and J.-F. Le Galliard. 2018.
658 Water restriction in viviparous lizards causes transgenerational effects on behavioral anxiety
659 and immediate effects on exploration behavior. *Behav Ecol Sociobiol* 72:23.

660 Ruf T. and F. Geiser. 2015. Daily torpor and hibernation in birds and mammals. *Biol Rev* 90:891–
661 926.

662 Seebacher F., C.R. White, and C.E. Franklin. 2015. Physiological plasticity increases resilience of
663 ectothermic animals to climate change. *Nat Clim Change* 5:61–66.

664 Senzano L.M. and D.V. Andrade. 2018. Temperature and dehydration effects on metabolism, water
665 uptake and the partitioning between respiratory and cutaneous evaporative water loss in a
666 terrestrial toad. *J Exp Biol* 221:jeb188482.

667 Sinervo B., F. Méndez-de-la-Cruz, D.B. Miles, B. Heulin, E. Bastiaans, M.V.-S. Cruz, R. Lara-
668 Resendiz, et al. 2010. Erosion of lizard diversity by climate change and altered thermal
669 niches. *Science* 328:894–899.

670 Somero G.N. 2010. The physiology of climate change: how potentials for acclimatization and
671 genetic adaptation will determine “winners” and “losers.” *J Exp Biol* 213:912–920.

672 Therneau T.M. 2018. *coxme: Mixed Effects Cox Models*. R package version 2.2-10.

673 Timmerman C.M. and L.J. Chapman. 2004. Behavioral and physiological compensation for chronic
674 hypoxia in the sailfin molly (*Poecilia latipinna*). *Physiol Biochem Zool* 77:601–610.

675 Trompeter W.P. and T. Langkilde. 2011. Invader danger: Lizards faced with novel predators exhibit
676 an altered behavioral response to stress. *Hormones and Behavior* 60:152–158.

- 677 Wingfield J.C., D.L. Maney, C.W. Breuner, J.D. Jacobs, S. Lynn, M. Ramenofsky, and R.D.
678 Richardson. 1998. Ecological bases of hormone—behavior interactions: the “Emergency
679 Life History Stage.” *Am Zool* 38:191–206.
- 680 Wong B.B.M. and U. Candolin. 2015. Behavioral responses to changing environments. *Behav Ecol*
681 26:665–673.
- 682 Woods H.A. and J.N. Smith. 2010. Universal model for water costs of gas exchange by animals and
683 plants. *Proc Nat Acad Sci* 107:8469–8474.

684 **Figures legends**

685 **Figure 1.** Summarized chronology of the experiment across months.

686

687 **Figure 2. Average change in body mass during the laboratory experiment (A, ΔM_b in grams \pm**
688 **SE) and change in body size growth during and after the laboratory experiment (B, ΔSVL in**
689 **mm.d⁻¹ \pm SE). Change was calculated since the beginning of the experiment. Control individuals are**
690 **represented by a circle; water-restricted individuals are represented by a square. Symbols indicate**
691 **the significance of the difference between treatments at each measurement: n.s. = non-significant, †**
692 **$p < 0.10$, * = $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.**

693

694 **Figure 3. Time spent before emergence (in 15 minutes steps) (A, \pm SE) and proportion of time**
695 **spent basking (B, \pm SE) by lizards in their home cages during morning (laboratory**
696 **experiment). Symbols similar to those of Figure 2.**

697

698 **Figure 4. Mean exploration score (A, \pm SE) calculated from behavioral measurements in a**
699 **neutral arena at each session, and intra-individual change in thermal preferences since the**
700 **beginning of the experiment (B, ΔT_p in °C, \pm SE). Black and grey symbols represent females and**
701 **males, respectively. Other symbols similar to those of in Figure 2.**

702

703 **Figure 5. Intra-individual change in basal metabolic rate variation calculated from O₂**
704 **consumption (A, ΔRMR in mL.h⁻¹ \pm SE) and standard evaporative water loss rates (B, $\Delta TEWL$**
705 **in mg.h⁻¹ \pm SE) during and after the laboratory experiment. Change was calculated since the**
706 **beginning of the experiment. In B, black and grey symbols represent females and males,**

707

Figure 1 BLACK AND WHITE









Figure 1 COLOR





Figure



Figure



Figure



Figure





Table 1

	Control individuals		Water restricted individuals	
	Females	Males	Females	Males
M_b (g)	1.68 ± 0.04	1.57 ± 0.04	1.64 ± 0.04	1.57 ± 0.03
SVL (mm)	49.6 ± 0.4	46.5 ± 0.5	50.0 ± 0.5	46.3 ± 0.5
T_p (°C)	33.4 ± 0.1	33.2 ± 0.1	32.9 ± 0.1	33.9 ± 0.1
RMR (mL.h ⁻¹)	0.32 ± 0.01	0.30 ± 0.01	0.31 ± 0.01	0.32 ± 0.01
TEWL (mg.h ⁻¹)	3.6 ± 0.2	4.0 ± 0.2	3.8 ± 0.2	4.2 ± 0.3

Physiological and behavioral parameters measured at week 0 (before the start of the water restriction treatment). Values are average \pm standard error. See text for abbreviation definitions.

Appendix

Appendix 1 and Table A1: Estimation methods of exploration rate following Rozen-Rechels et al. (2018).

Appendix 2: Effects of water restriction on survival from the end of the experiment to the next reproductive season.

Appendix 3: Effects of water restriction on reproductive effort of females during the next reproductive season.

Table A2: Model statistics of model explaining variation of ΔTEWL when adding ΔRMR as a fixed additive variable.

Appendix 1

Behavioral statistics were measured from videos, transformed into image sequences, using the image analysis protocol of Rozen-Rechels et al. (2018). We measured the relative time spent active (T), the total distance walked by the lizard (D), the number of activity bouts (i.e., number of sequences of consecutive images with positive step length, N_{bouts}), the average distance walked during a bout (D_{bouts}), the average distance walked between two images in a bout (M_{bouts}), the mean standard deviation of distances walked between two images in a bout (homogeneity of displacements; SD_{bouts}), the time spent by the lizard in the buffer zone along the walls (T_{walls}), the total distance walked out of the buffer zone ($D_{1\text{walls}}$), the mean distance to the walls ($D_{2\text{walls}}$), and the standard deviation of the distance to the walls (SD_{walls}). We further subdivided the neutral arena into 24 equal squares, to estimate the total area visited by the lizards including the proportion of squares visited at least once by the individual (τ_{explo}) and the standard deviation of the number of locations per square or heterogeneity of the exploration (SD_{explo}).

In order to calculate the exploration rate, we transformed each behavioral trait using a Box-Cox power function to meet gaussian requirements. We used principal component analysis (PCA) (R package “ade4”, Dray and Dufour 2007) to identify correlated behaviors (i.e. behavioral syndromes). The first component of the PCA (PC1) corresponded to the one identified by Rozen-Rechels et al. (2018) as an exploration syndrome (same correlations between variables; see Table A1). PC1 scores, positively correlated to the time spent walking or the distance walked, were then extracted and individuals were ranked according to them.

Table A1

Behavioral measurements	Contribution to PC1	Contribution to PC2
D	0.14	0.002
T	0.12	0.0001
N _{bouts}	0.06	0.003
M _{bouts}	0.13	0.003
SD _{bouts}	0.12	0.0001
D _{bouts}	0.12	0.00009
T _{walls}	0.02	0.24
D _{1walls}	0.14	0.015
D _{2walls}	0.02	0.27
SD _{walls}	0.0001	0.31
τ_{explo}	0.06	0.11
SD _{explo}	0.07	0.05

Contributions of each measured behaviors to the axes of the PCA. Bold contributions are those higher than a random contribution (i.e. $100/12=8.33\%$). Axis 1 explained 55% and axis 2 explained 23% of the variance in the data.

Appendix 2

We estimated survival from recapture data collected in May 2017. It was then possible to estimate lizard survival with a high degree of reliability thanks to a recapture probability close to 1.

Recapture effort in September 2016 was not high enough to ensure sufficient confidence in the survival data. Still, we recaptured individuals from the experiment in May 2018 that have not been captured in 2017 (3 individuals). We then analyzed survival from August 2016 to May 2017 considering they were alive even though not captured. We excluded individuals that have been paralyzed due to decalcification at the end of July.

Survival has been analyzed using a mixed-effect logistic regression model including a logit link and a binomial error term (package *lme4*) with a two-way interaction of the water restriction treatment with the sex of the individual as fixed effect and the enclosure identity as a random effect.

Treatment had not significant effect on survival, whatever the sex (treatment \times sex: $\chi^2 = 2.1$, df = 1, p = 0.15; treatment: $\chi^2 = 0.07$, df = 1, p = 0.79). Males had a significantly higher survival than females ($\chi^2 = 4.77$, df = 1, p = 0.03).

Appendix 3

We estimated the reproductive effort of females recaptured in May 2017 by calculating total litter size and mass (i.e. the number and summed mass of all not-fecundated eggs, aborted embryos, dead juveniles and live juveniles in the litter) and calculating litter size and mass (i.e. only in live juveniles). One female was not pregnant and has been excluded from the analysis.

(Total) litter size was analyzed using a mixed-effect logistic regression model including a logit link and a Poisson error term (package *lme4*, Bates et al. 2015) with the water restriction treatment as fixed effect and the enclosure identity as a random effect. (Total) litter mass was analyzed using a mixed-effect linear model (package *nlme*) with the water restriction treatment as fixed effect and the enclosure identity as a random effect.

Neither (total) litter size nor mass were affected by the water restriction treatment (all $p > 0.19$).

Table A2

Sessions	Variable	Numerator DDL	Denominator DDL	<i>F</i>	<i>P</i>
Weeks 4 and 7	Δ RMR	1	95	3.38	0.06
	Time	1	95	22.3	< 0.0001
	Treatment	1	97	1.11	0.30
	Time \times Treatment	1	95	0.61	0.44
May 2017	Δ RMR	1	47	1.19	0.28
	Sex \times Treatment	1	47	4.51	0.0

Effect of the water restriction treatment on Δ TEWL when adding Δ RMR as a fixed additive effect to the previously selected model. Results are unchanged (see Results section).

Dear Mr. Rozen-Rechels:

An editorial decision has been made regarding your revised manuscript, "Acclimation to water restriction implies different paces for behavioral and physiological responses in a lizard species" (PBZ-19062R1).

I have read your paper and received reports from Reviewers 2 and 3, in addition to input by the Handling Editor for your paper, who is a member of the PBZ Editorial Team. The reviewers returned constructive comments, though their recommendations are mixed, with some additional changes requested. We believe that the manuscript could make a good contribution to the field if appropriate final changes are made.

The reports below present you with comments ranging from minor editorial corrections to more substantive concerns, which will need to be fully addressed before a final editorial decision can be made. We would like to provide you with the opportunity to address these remaining remarks in a final revision. Therefore, we are inviting you to revise and resubmit the paper within 30 days. Please contact us if you will need more time.

The reviewers' comments are below. As you revise your paper, please pay careful attention to this input, responding to each remark in detail, and noting what changes you made to the paper in the response. Your detailed responses to reviewers and list of changes to the paper will be very helpful during the final editorial evaluation of your paper.

If you have additional information that you would like to have published with your paper, please consider submitting it as appendices or online supplements.

When your revision is complete, please submit it online at www.editorialmanager.com/pbz, uploading each figure and table separately, with figures as high-resolution image files (300 or more dpi) and tables as editable documents.

Thank you again for considering PBZ as an outlet for your work. We will be looking forward to reading your revised paper.

Sincerely,

Theodore Garland, Jr.
Editor in Chief
Physiological and Biochemical Zoology

Response: First, we would like to thank the journal, its editorial board and the reviewers who all helped significantly improving our manuscript. We thank Pr. Theodore Garland, Jr. for considering our manuscript for publication. We answered the editor and the reviewer concerns point-by-point below and revised the main text accordingly.

Editorial Comments

Thank you for your diligent revisions, which have improved the paper. In addition to the very helpful input from the referees, there are a few editorial recommendations that we would like you to take into consideration as you make final changes to the manuscript.

Response: We thank the associate editor for supporting our paper for publication. We took the recommendation of the editor and the referees in account and responded to it point-by-point. We also updated our acknowledgements to thank the editor and the referees for significantly improving our manuscript.

Changes are highlighted in color in the new version of the manuscript.

First, please outline for readers, within the body of the paper, the same clarifications that you offer to the referees. Our reviewers are representative of the readers of PBZ, and their concerns will likely be shared.

Response: As detailed in our point-by-point response, we updated the text where Reviewer 2 asked to.

Second, please update Figure 1 by providing a grayscale version for potential print publication. The color image can be retained for potential publication in the online version.

Response: We provided two versions of figure 1, a coloured one and a grayscale one.

Third, please consider citing the attached paper, if relevant:

Kelly, S. A., T. M. Panhuis, and A. M. Stoehr. 2012. Phenotypic plasticity: molecular mechanisms and adaptive significance. *Comprehensive Physiology* 2:1417â€“1439.

Response: We thank the associate editor for suggesting this reference. We have decided to cite this paper when introducing adaptive plasticity (line 42) and discussing beneficial acclimation (line 390). See lines 584-585 in the reference list.

Reviewer 2 Comments

The authors have made a good attempt to address the original review comments, and I thank them for their effort. I have a few small items that are still outstanding.

Response: We thank the reviewer for validating the changes we have made which significantly improved our manuscript.

The reference by Kennett and Christian has been removed from the text but still appears in the reference list at the end.

Response: The reference had been deleted from the list.

My comment about adequately taking body size into account may have not been clear. I was not suggesting that some other metric other than mass should have been used - I absolutely agree that mass is the appropriate metric to use. My comment was related to clarification about how the body size (mass) was taken into account statistically. In the Response to Reviewers document, the authors state: "We used body mass instead of body size in our ANCOVA as it ..." So, if ANCOVA was used, then that satisfies my concern. However, when I refer back to the text, there is still no reference to the term "ANCOVA". The authors mention that mass was used in the model, but that is not sufficient for me to understand how it was incorporated in the model. If appropriate, please include the term "ANCOVA" in the text, and this will satisfy my concern. If, on the other hand, ANCOVA

was not used, then further explanation is required to convince a reader like me that body mass was appropriately included in the statistical comparisons.

Response: An ANCOVA was used and in the case of studying RMR and TEWL variations, body mass scaling was taken into account by adding the body mass as a fixed effect. We changed the text accordingly (line 274: “trait variation was analyzed with ANCOVAs using linear models” and line 278 “For Δ RMR and Δ TEWL, we included individual M_b as a fixed effect in the ANCOVA to account for body mass scaling.”).

Similarly, in the Response to Reviewers document, the authors have adequately addressed my concerns about the way the IR thermometer was used. However, that is not reflected in the text. The text makes reference to the 6 mm diameter spot, but there is no mention of the lasers that the thermometer provides. By mentioning the lasers in the text, you will convince readers like me that you were able to direct the beam onto lizards skin rather than simply pointing it in the general direction of the lizard but measuring some part lizard skin and some part surrounding substrate. So, the authors just need to make the text as convincing as they have made the Response to Reviewers.

Response: As suggested by the reviewer we have also updated the manuscript to answer this concern (line 236-237: “, that we were able to assess thanks to three lasers indicating the centre of the circle and the diameter of the spot when measuring the temperature”).

Reviewer 3 Comments

The authors have addressed all my comments and questions. It is a very well designed study with interesting results.

Response: We thank the reviewer for the positive support.