

# Early Levallois core technology between Marine Isotope Stage 12 and 9 in Western Europe

Marie-Hélène Moncel, Nick Ashton, Marta Arzarello, Federica Fontana, Agnès Lamotte, Beccy Scott, Brunella Muttillo, Gabriele Berruti, Gabriele Nenzioni, Alain Tuffreau, et al.

# ▶ To cite this version:

Marie-Hélène Moncel, Nick Ashton, Marta Arzarello, Federica Fontana, Agnès Lamotte, et al.. Early Levallois core technology between Marine Isotope Stage 12 and 9 in Western Europe. Journal of Human Evolution, 2020, 139, pp.102735. 10.1016/j.jhevol.2019.102735. hal-02489414

HAL Id: hal-02489414

https://hal.science/hal-02489414

Submitted on 12 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Early Levallois core technology between MIS 12 and 9 in Western Europe? Marie-Hélène Moncel<sup>a,\*</sup>, Nick Ashton<sup>b</sup>, Marta Arzarello<sup>c</sup>, Federica Fontana<sup>c</sup>, Agnès Lamotte<sup>d</sup>, Beccy Scott<sup>c</sup>, Brunella Muttillo<sup>b</sup>, Gabriele Berruti<sup>b,e</sup>, Gabriele Nenzioni<sup>f</sup>, Alain Tuffreau<sup>d</sup>, Carlo Peretto<sup>b</sup> <sup>a</sup> UMR 7194 CNRS – Département Hommes et environnement, Muséum National d'Histoire Naturelle, Institut de Paléontologie Humaine, Paris, France <sup>b</sup> Department Britain, Europe & Prehistory, British Museum, Franks House, 56 Orsman Road, London N1 5QJ, UK <sup>c</sup> Sezione di Scienze preistoriche e antropologiche, Dipartimento di Studi Umanistici, Università degli Studi di Ferrara, Corso Ercole I d'Este, 32 – I-44121, Italy d University of Lille, UMR 8164, Bâtiment de géographie, avenue Paul Langevin, Villeneuve d'Asca, France <sup>e</sup> Museo di Archeologia e Paleontologia C. Conti, Borgosesia, Italy <sup>f</sup> Museo della Preistoria "L. Donini", Via Fratelli Canova, Italy \*Corresponding author. E-mail address: marie-helene.moncel@mnhn.fr (M.H. Moncel). 

## Acknowledgements

This study belongs to a research project between the CNRS-National Museum of Natural History, the British Museum (UK), the University of Ferrara (Italy), the University of Lille (France), the Museo di Archeologia e Paleontologia C. Conti (Italy) and the Museo della Preistoria L. Donini (Italy). Results were presented at the University of Haïfa (Israel), November 2017 "The Lower to Middle Paleolithic boundary: A view from the Near east" (Israel Science Fondation and Wenner-Gren Foundation workshop). We would like to thank Mina Weinstein-Evron and Yossi Zaidner for the organization.

Early Levallois core technology between MIS 12 and 9 in Western Europe?

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

27

#### Abstract

Early Levallois core technology is usually dated in Europe to the end of Marine Isotope Stage (MIS) 9, and particularly from the beginning of MIS 8 to MIS 6. This technology is considered as one of the markers of the transition from Lower to Middle Paleolithic or from Mode 2 to Mode 3. Recent discoveries show that some lithic innovations actually appeared earlier in western Europe, from MIS 12 to MIS 9, contemporaneous with changes in subsistence strategies and the first appearance of early Neanderthal anatomical features. Among these discoveries, there is the iconic Levallois core technology. A selection of well-dated assemblages in the United Kingdom, France, and Italy dated from MIS 12 to 9, which include both cores and flakes with Levallois features, has been described and compared with the aim of characterizing this technology. The conclusion supports the interpretation that several technical features may be attributed to a Levallois technology similar to those observed in younger Middle Paleolithic sites, distinct from the main associated core technologies in each level. Some features in the sample of sites suggest a gradual transformation of existing core technologies. The small evidence of Levallois could indicate occasional local innovations from different technological backgrounds and would explain the diversity of Levallois methods that is observed from MIS 12. The technological roots of Levallois technology in the Middle Pleistocene would suggest a multiregional origin and diffusion in Europe, and early evidence of regionalization of local traditions through Europe from MIS 12 to 9. The relationships of Levallois technology with new needs and behaviors are discussed, such as flake preference, functional reasons related to hunting and hafting, an increase in the use of mental templates in European populations and changes in the structure of hominin groups adapting to climatic and environmental changes.

**Keywords:** Neanderthals; Early Levallois; Western Europe; Technology

51

52

#### 1. Introduction

Early Levallois core technology is usually dated in Europe to the end of MIS 9, and particularly from the beginning of MIS 8 to MIS 6. This technology is considered as one marker of the transition from the Lower to Middle Paleolithic or from Mode 2 to Mode 3 (Clark, 1969), resulting in the general adoption of more complex flaking strategies and a higher standardization of products (White and Ashton, 2003; Monnier, 2006; Moncel et al., 2011, 2012; Scott, 2011; White et al., 2011; Fontana et al., 2013; Adler et al., 2014; Wiśniewski, 2014; Villa et al., 2016; Picin, 2017). Recent discoveries show that some lithic innovations actually appeared earlier in Western Europe, from MIS 12 to MIS 9, contemporaneous with changes in subsistence strategies and the first appearance of early Neanderthal anatomical features. Evidence of progressive or gradual developments in behaviour is recorded from c. 400 ka with fire use (Roebroecks and Villa, 2011; Gowlett, 2016) and c. 300 ka through organized hunting strategies (e.g., at Schöningen in Germany; Thieme, 1997; Blasco et al., 2013; Conard et al., 2015; Rodriguez-Hidalgo et al., 2017) and. Likewise, paleontological studies and recent DNA analyses suggest the appearance of the earliest Neanderthal features across Western Europe in Homo heidelbergensis populations between 600 and 450 ka (Krings et al., 1997; Hublin, 1998, 2009; Hublin and Pääbo, 2005; Orlando et al., 2006; Bischoff et al., 2007; Rightmire, 2008, Endicott et al., 2010; Green et al., 2010; Stringer, 2012; Meyer et al., 2014, 2016). Among the lithic innovations, the first evidence of Levallois technology can be reinvestigated, as recent findings seem to attest to an earlier practice. A fresh look at old collections named proto-Levallois, pre-Levallois or Prepared core Technology (PCT) prior to MIS 9/8 has to be undertaken in the context of these new discoveries. Levallois technology was first identified by Boucher de Perthes (1857) with the recognition of three main criteria (preparation of the core surface, role of the convexities and subsequent detachment of one flake). The definition varied over time, changing from the production of one main end-product to various predetermined end-products (De Mortillet G., 1883; Commont, 1909; Bordes, 1950; Boëda, 1986). Experiments clarified the definition and technological requirements, often (but not always) facetted platforms, angles of percussion and management of the volume of the core (Breuil and Kelley, 1954; Boëda, 1995; Lenoir and Turq, 1995).

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

All the definitions recognized that this technology enabled control of the shape and standardization of the end-products, and required general preparation of the core volume and management of core convexities. Recognition of these technological features on cores and flakes allows identification of Levallois core technology.

We have selected well-dated assemblages from MIS 12 to 9 where in the past both cores and flakes were described as Levallois, proto-Levallois or 'prepared cores', or have recently been found by new fieldwork. They are located from the northwest to the south of Europe in the UK (Purfleet and consideration of other occurrences), France (Cagny-la-Garenne I-II, Orgnac 3), and Italy (Guado San Nicola, Cave dall'Ollio; Fig. 1) and described as the earliest evidence for each country of Levallois core technology. These assemblages are frequently discussed with the consideration that not all the classical Levallois characteristics are found together (i.e., Malinsky-Buller, 2016b; Soriano and Villa, 2017). We aim to review the attribution of these cores and flakes in the light of the new data, to characterize this technology (accidental or evidence of technical innovation), which coexisted for an extended period with earlier technologies. These technologies will be discussed by region with ideas on its origin, such as technological roots in the Middle Pleistocene, arrival of populations or diffusion from multiple areas, the relationship with new needs and behaviours, and the evolution of European populations. In the light of the recent findings, the period of MIS 12 to 9 can be considered as a threshold in cultural human evolution and testing of new technological behaviours, raising questions on how we term this important period. Are we dealing with a phase of invention, deliberate or by chance (Renfrew 1978), or innovation, namely the adoption of an invention by a large number of individuals? Determining the timing and mode of the onset of Levallois core technology in Europe is crucial to understanding how these behavioural changes developed at the inception of the Neanderthal (or *Homo heidelbergensis*) way of life.

102

103

104

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

# 2. Materials and methods

#### 2.1. The corpus of sites

From the north to south of western Europe, there are well-dated archaeological sites that show isolated examples of core technologies that have been identified in the past by the originality of the preparation of the flaking surface and the control of the form of the end-products. A selection of these sites, dating from MIS 12 to the end of MIS 9 and from a range of environmental and geological contexts, are reviewed to describe the variation in this technology and to discuss the attribution (or not) to an early form of Levallois core technology. These assemblages are often, but not always, associated with bifaces.

While from MIS 8 the recognition of Levallois is not questioned and the definition of Levallois core technology is largely agreed, the multitude of terms for earlier Levallois indicates that the recognition of this core technology older than MIS 8 is more problematic. The terms used include proto-Levallois, pre-Levallois, prepared cores, or simple prepared cores (Wymer, 1968; Roe, 1981; White and Ashton, 2003). Discovery of some sites in the earlier years of the subject, led to the use of the terms proto- or pre-Levallois due to the unusual nature of the cores, which did not resemble the 'classic' Levallois cores from sites such as Baker's Hole in Britain. In the UK, this led to the adoption of the term 'simple prepared cores' in part to try and avoid the implication of an evolutionary progression that was promoted by the terms proto- or pre-Levallois (White and Ashton, 2003). Despite the adoption of the new term, it was still sometimes used to imply an early date (e.g., Bolton, 2015), even though such cores are found in both pre-MIS 8 and post-MIS 8 contexts (see below).

The background of the selected sites for this review are briefly described below in chronological order. The site of Cagny-la-Garenne is located in fluvial deposits of the Middle Terrace of the Somme Valley (North France). Human occupations took place between the alluvial plain and the limestone slope. The gravels have been attributed to MIS 12 based on the strong regional geological framework of the Somme (i.e. Antoine et al., 2007, 2016). The terrace system of the Somme is particularly well represented in the middle part of the valley, between Amiens and Abbeville, where a set of stepped alluvial formations is preserved by a covering of well developed loess-and palaeosols. In this area, 10 stepped alluvial formations have been recognized between + 5/6 m and + 55 m relative height above

the maximum incision of the present day valley. The summary of the data (sedimentology, bio-indicators and geochronology) shows that each alluvial formation corresponds to the morphosedimentary budget of a single glacial-interglacial cycle. The glacial stages are characterized by a braided river system and mainly sand and gravel deposition while interglacial stages correspond to a meandering river system, with overbank silt deposition and marshy soil formation at the top. Interglacial.

The Electro Spin Resonance (ESR) date of the formation at the site of Cagny-la-Garenne I is of 400  $\pm$  101 ka, while other dates on the same alluvial formation (n°V Garenne Formation + 27-29 m) have given ages of 448  $\pm$  68 ka, 443  $\pm$  53 ka and 403  $\pm$  73 ka (Antoine et al., 2003, 2007, 2016). The dates in combination with the evidence of deposition in a cold environment suggest an MIS 12 age for the formation.

At Cagny-la-Garenne I, the six artifact assemblages (Level CA to CXB) were made from the locally available flint and consist of bifaces, core and flake manufacture with notches and denticulates (Tuffreau, 1987; Lamotte, 1994, 2012). Assemblages CXCA and CA are in primary context close to the Chalk slope, LJ and LG come from fluvial silts, CXB from limestone gravels and at the top CXV comes from coarse, periglacial gravels (Tuffreau and Lamotte 2010). At Cagny-la-Garenne II, 100 m from Cagny-la-Garenne I, four archaeological levels (I, J, K and L) were recovered, while at the top, five archaeological levels (I0–I4 and J) came from gravels (Tuffreau and Lamotte, 2001). Once again, the fluvial sequence is banked up against the Chalk slope. Raw material was available on site in the form of large flint nodules. All stages of core working and biface manufacture are present.

The site of Guado San Nicola is located in south central Italy (Molise Region). It is an open-air site systematically excavated from 2008 to 2015 over an area of 98 m² (Peretto et al., 2016). A 20 m stratigraphic core in the immediate vicinity of the excavation, and a series of stratigraphic sections investigated in the area, have confirmed the sequence of the excavation. From bottom to top, the sequence is composed of eight stratigraphic units (S.U.) The 2 m thick-sequence is of polygenic gravelly silty and clayey deposits and contains interstratified tephra layers. It has been dated on the basis of

morphostratigraphic considerations and radio-isotopic dating of volcanic deposits. The <sup>40</sup>Ar/<sup>39</sup>Ar and ESR/U-series dates clearly place the archaeological occupation at the transition between MIS 11 and 10 (400 and 345 ka). Unit S.U.C., rich in lithic and faunal remains, is dated to 400 +- 9 ka by <sup>40</sup>Ar/<sup>39</sup>Ar (Pereira et al., 2016). The faunal assemblage can be attributed to the typical Galerian and to the Fontana Ranuccio Faunal Unit. The faunal assemblage is mainly composed of the remains of *Cervus elaphus acoronatus, Cervidae, Equus ferus* ssp., followed by *Palaeoloxodon* sp., *Bos primigenius* and *Stephanorhinus kirchbergensis, Ursus* sp. and *Dama* sp. The sedimentary succession consists of four archaeological levels (C, B\*C, B, A\*B) with lithic assemblages composed of reduction sequences of both debitage and shaping. The raw material, (mainly flint of good quality with a high degree of silicification and, more rarely, limestone) was collected from a secondary context in the form of cobbles or slabs. The main flaking methods are an opportunistic exploitation (c.f. alternate platform), followed by discoidal and centripetal debitage. The reduction sequences for bifaces are not complete and lack preparation phases (Muttillo et al., 2014). They were made by direct percussion with a hard hummer and final retouch by soft hammer.

The site of Orgnac 3 in southeast France, first developed as a cave and then became an open doline. The archaeological sequence of 10 levels is dated through biostratigraphy and ESR, U/Th dating from MIS 9, while for levels 2 and 1 at the top of the sequence, dated by volcanic mineralogy to the beginning of MIS 8 (Combier, 1967; Debard and Pastre, 1988; Falguères et al., 1988; Masaoudi, 1995; Moncel et al., 2011, 2012). ESR and Uranium/Thorium (U/Th) give ages of 288 – 45/+ 82 ka, 309 ± 34 ka and 374 – 94/+ 165 ka for the bottom of the archaeological sequence (levels 5b and 6) (Falguères et al., 1988; Laurent, 1989; Masaoudi, 1995), attributed to the MIS 9. Four pure calcite samples of the levels 5b–6–7 (bottom of the sequence) have been dated by U/Th by MC-ICPMS (High-precision Mass Spectrometry and Environment Change Laboratory, HISPEC, Taiwan). Dates vary between 255 and 319 ka (Michel et al., 2011). Level 2 contains volcanic minerals from an eruption of the Mont-Dore volcano, eruption dated to the beginning of the MIS 8 (298 ± 55 ka) (Debard and Pastre, 1988). Direct dating by <sup>40</sup>Ar/<sup>39</sup>Ar on sanidine grains (cineritic material) has been applied to level 2 (Org-C1). The 12 dates are between

276 and 326 ka with an average age of 308.2 ± 6.8 ka. The result is in agreement with the age of 298±55 ka by Fission track dating (FT) on zircons (Khatib, 1994). The biostratigraphy associating large mammals, micromammals and pollen date the bottom of the sequence (levels 7 to 3) to an interglacial of the Middle Pleistocene (Mourer-Chauviré, 1975; Tillier, 1991; Jeannet, 1981; Guerin, 1980; Gauthier, 1992; El Hazzazi, 1998; Aouraghe, 1999; Sam, 2009). Level 1 is indirectly attributed to the MIS 8 by *Hemitragus bonali* and *Ursus deningeri*. Levels 2 and 1 attest of open landscape with the replacement *Equus mosbachensis* by *Equus steinheimensis* (Forsten and Moigne, 1988).

Pre-Neanderthal human remains were discovered in the lowest layers (Lumley de, 1981). The lithic assemblages record a mosaic of changes over time, towards Early Middle Paleolithic strategies (Moncel et al., 2012; Moigne et al., 2016). Debitage activity is dominant and bifaces have variable ratios through the sequence with less than 1% in the top levels. Thin slabs of flint were the main blanks collected locally.

The open air site of Cave dall'Olio is located in an alluvial context along the Northern Apennine edge near Bologna, northern Italy. The lithic assemblage was recovered in the 1970s along a stratigraphic profile brought to light by quarry activities at the top of a fersiallitic palaeosol within the gravels of the River Idice at a depth of about 20 m below the present surface. The fersiallitic palaeosol has been referred to the Molino Unit of the Apennine-Po Plain Quaternary stratigraphic framework dated to MIS 9, indicating a terminus ante quem for the chronology of this assemblage (Farabegoli and Onorevoli, 1996, 2000; Fontana et al., 2010). The dating of the gravels and the soil containing the lithic industry of Cave dall'Olio, as well as their paleoenvironmental interpretation, are based on the integration of data derived from the study of the profile of S. Mamante (Faenza); 22 shallow marine to terrestrial Quaternary units were produced by the long-term activity of a right transcurrent fault with various outcrop segments distributed across a sector of the Emilia-Romagna Apennines edge for a total length of more than 150 km (Farabegoli and Onorevoli, 1992, 1996). Within the reconstructed scheme, the first continental units are dated to the Upper Matuyama chron reverse period and to the Bruhnes direct palaeomagnetic chron. The latter contains the earliest evidence of human occupation in this

area (on a stratigraphic basis Bel Poggio and Romanina Bianca are considered to be of the same age as Ca' Belvedere di Montepoggiolo with mode 1 assemblages). The sediments of the continental units correspond in most cases to the glacial-interglacial transition periods and are intercalated with 8 fersiallitic soils typical of warm interglacial phases. Correlations have been established between the different portions of the outcropping terraced deposits, which are recognizable upstream along the valley flanks. These have allowed the fersiallitic paleosoil identified at Cave dall'Olio to be referred to as the Molino Unit of the Apennine-Po plain Quaternary stratigraphic framework, dated to MIS 9. Initial studies led to classification of this lithic industry, which is dominated by debitage with evidence of manufacturing of bifaces as ancient Clactonian and proto-Levallois (Bisi et alii 1982; Lenzi and Biagioli, 1996) after the original definition by Palma di Cesnola (1967).

In the UK, at Purfleet, Essex, Paleolithic artifacts have been recovered from sediments exposed in four chalk quarries, in the Lower Thames Valley. From east to west, these are the Bluelands, Greenlands, Esso and Botany Pits. The pits reveal terrace deposits occupying an abandoned meander loop of the Thames as part of the Lynch Hill/Corbets Tey terrace (Bridgland, 1994), banked up against the north facing chalk slope of the Purfleet anticline. The sequence comprises gravel (Little Thurrock Member) overlain by interglacial deposits rich in faunal material (Purfleet Member) fining upwards to a silty clay and surmounted by gravel (Botany Member). An assemblage of artifacts excavated and collected by Andrew Snelling from the Botany Gravel at Botany Pit was initially described as 'Proto-Levallois' (Wymer 1968), and 'reduced' Levallois with simplified preparatory stages (Roe, 1981). This gravel reflects a return to cold climate gravel deposition following a fully temperate episode, suggesting an MIS 9/8 date, an attribution which is supported by an OSL date of 324 ka (MIS 9) from an equivalent position at Greenlands Pit (E. Rhodes, quoted in White and Ashton 2003).

#### 2.2. Methods

For this paper we use the terminology of Boëda (1986, 1993, 1995) for the overall concepts of Levallois: a volumetric concept with six technological criteria: (1) core maintenance (lateral and distal

convexities), (2) predetermination of end-products, (3) normalization of end-products, (4) potential for resharpening, (5) ramification, and (6) productivity. The lower surface of the core is devoted to the striking platform and the upper surface to Levallois flake production. To distinguish between cores, such as those from Purfleet, we use the term 'simple prepared core', without implication for an early date. As defined by White and Ashton (2003) they are cores where the striking platform has been selected, minimally prepared and orientated in relation to the pre-existing lateral and distal convexities of one flaking surface. The flakes removed from this surface tend to be larger than any of the preparatory flakes (by which the platform was created), and to flake along the surface at an angle close to 90° to the platform, rather than biting excessively into the core volume. For cores, such as those from Baker's Hole where there is a preferential, single removal, we use the term 'classic Levallois', as for the other sites of our corpus, Cagny-la-Garenne I-II, Orgnac 3, Cave dall'Olio and Guado San Nicola.

#### 3. The technological review of lithic material described as Levallois

3.1. Cagny-la-Garenne I and II (North France)

At Cagny-la-Garenne I, from the base of the sequence (level CA) toward the top (level CXV), the appearance of flakes and cores, described in the past as proto-Levallois, increases, but always in low quantity. In the sandy levels of the middle of the sequence (levels Lj, Lg), this kind of production is rare or absent. At Cagny-la-Garenne II, three levels yielded one or two cores (levels I3, I4, J; Table 1).

For each layer, the main core technology is unipolar and unifacial with few scars and a prepared/cortical platform. On a small quantity of cores, various methods (lineal, unipolar, bipolar and centripetal) are employed for the extraction of the end-products with evidence on the cores of management of the distal and lateral convexities and plain or facetted platforms (Figs. 2–4). The removals extend over at least half of the main length of the core surface and their morphology is due to the organization of the convexities. The preparation of the convexities tends to change from unipolar towards centripetal at the top of the sequence for Cagny-la-Garenne I. Most core sizes vary between 50 and 110 mm in maximum dimension. Among the flakes, we can identify some core edge

flakes (débordant flakes) with many scars, which are probably from the preparation of the core convexities. In assemblage CXB at Cagny-la-Garenne I, there are several biface-cores with invasive removals interpreted as attempts at Levallois.

#### 3.2. Guado San Nicola (south central Italy)

The Levallois assemblage is fresh and the main raw material is aphanitic and microbrecciated flint, or occasionally macrobrecciated flint and silicified limestone. It is of better quality than that used for the bifaces. The supports were ovoid cobbles and quadrangular slabs, or occasionally large flakes. Different stages of the reduction process can be identified and reveal careful preparation, management and maintenance of flaking platforms (angles ranging from 55 to 85°) and convexities (mainly centripetal), indicating the ability to prepare and reprepare cores for predetermined flakes (Fig. 5). Various methods were used in equal quantity (single preferential flake, recurrent centripetal, unipolar and bipolar) and there is evidence of facetted platforms for lineal and recurrent unipolar debitage (Table 4; Fig. 5, 6). The Levallois cores are exhausted and some were made on flakes.

Overshot flakes managed the lateral and distal convexities, except for the lineal debitage with centripetal removals. Levallois flakes (n = 55) mostly result from 'plein debitage' or repreparation of the convexities, produced by unipolar and centripetal recurrent method (Fig. 6). The striking platforms are dihedral or flat, rarely facetted. Levallois points and blades, as well as retouched Levallois flakes, are extremely rare.

#### 3.3. Orgnac 3 (south east France)

In the lowest levels (5b and 5a), less than 10% of the cores and less than 3% of the flakes can be classified as Levallois (Table 2; Figs. 7 and 8). The cores are unipolar/bipolar or centripetal recurrent. The knapping surface indicates the utilization of the core edges, one or two partially prepared striking platforms, and maintenance of lateral and distal convexities. The 'Levallois' cores are very different to the frequent unexhausted unifacial and bifacial centripetal cores on slabs or thick flakes, where there

is no sign of management of the convexities. Centripetal and Levallois cores are associated with some prismatic, polyhedral and orthogonal cores. Some 20% of the rare Levallois flakes are débordant flakes, while 30% of platforms are facetted and 10% dihedral.

In levels 4b and 4a, around 40% of the cores and 2–8 % of flakes can be defined as Levallois (Figs. 8–10). The Levallois cores are associated again with centripetal cores. The cores on flakes again indicate evidence of predetermined flaking, preparation of distal and lateral convexities, use of hard hammer and distinctions between striking platform and flaking surface. The methods applied are again unipolar and bipolar recurrent, but the preferential flake method, not used in the lower levels, becomes the most common. In contrast to the underlying levels, the predetermined removals never cover the flaking surface. As for levels 5b-5a, the quantity of Levallois flakes is very low suggesting export of flakes. Débordant flakes total between 20 and 50% of the assemblage. The removals are mainly centripetal and the ratio of flakes with an invasive scar on the upper face increases. The size of flakes is more variable than the cores.

#### 3.4. Cave dall'Olio and other assemblages of the Northern Apennine margin (northern Italy)

The assemblage totals 494 lithic artifacts, with 71 cores, 403 retouched and unretouched blanks, 5 pebble tools and 15 bifaces (Table 3; Fontana et al., 2013). Most of the assemblage was obtained from a dark colored silicified siltstone that is very abundant locally and available in large-sized nodules and pebbles (10–40 cm). Bifaces were mostly obtained from large flakes and were always worked with a small number of deep removals.

Cores are dominated by unidirectional recurrent schemes with either parallel or convergent removals and represent around 30% of the assemblage. Recurrent crossed methods predominate, while lineal, recurrent bipolar and centripetal flaking methods are rarer (Fig. 11). The flaking surfaces were prepared by either débordants or orthogonal removals and core platforms were variably prepared. Levallois cores are associated with a few prismatic types resulting from the application of a laminar reduction process sensu lato—also reported as 'direct non-Levallois reduction sequence'

and laminar reduction. Other methods include Kombewa, opportunistic and possibly discoid reduction.

Levallois end-products vary in shape and elongation according to the method applied. There are elongated blanks with parallel edges and frequently characterized by a backed edge from unidirectional parallel and bidirectional methods (with some items possibly obtained from a laminar reduction). There are also flakes with convergent and frequently déjétés lateral margins derived from application of the recurrent unidirectional convergent scheme, small-middle sized oval-shaped flakes from the lineal method and small oval and subtriangular flakes extracted with centripetal debitage. Most products measure between 40 and 90 mm in length. Platforms are generally flat while facetted types are rare (7%, including one 'chapeau de gendarme'). Although in several cases the condition of

(Révillon, 1995 )—and with a small group of cores featuring mixed characteristics between Levallois

Several other assemblages recovered during field surveys from the river terraces of the Northern Apennine edge, which are dated to the same age as Cave dall'Olio, are characterized by similar technological features. The general features of such assemblages show that this area was intensively occupied by human groups that were able to apply different predetermined debitage reduction sequences prior to MIS 9/8 (Lenzi and Nenzioni, 1996).

the artifact surfaces was altered by fluvial deposition, around 30 retouched blanks were identified,

#### 3.5. Purfleet (UK) and other UK sites with simple prepared cores

especially scrapers and denticulates, including some on Levallois flakes.

The assemblage from Purfleet consists of over 4000 artifacts, including 30 bifaces, but the vast majority are flakes (White and Ashton, 2003; Scott, 2011; Bolton, 2015). Distinguishing end-products is problematic and only rare examples of potential Levallois flakes can be identified. A sample of more than 300 cores has been examined in more detail (Scott, 2011). They consist of 170 migrating platform cores, 28 discoidal cores, 80 simple prepared cores and 25 cores that are considered as Levallois. The simple prepared cores conform to the description above, while the few Levallois cores have mainly lineal exploitation (84% with one invasive removal) with a few examples showing unipolar, bipolar and

recurrent techniques (with 2 to 3 removals on the flaking surface). The mean length of Levallois cores is 87 mm. The preparation method is mainly centripetal (96%) and there are also cores with proximal, lateral and/or distal preparation of the convexities (Fig. 12). The striking platform is mainly partial. There are also four core débordant flakes and overshot distal flakes. A total of 26 flakes are considered as Levallois. They are large and elongated (mean length of 115 mm) and result from lineal exploitation with a centripetal preparation. None has facetted platforms. As the assemblage was recovered from fluvial gravel, it is not certain whether all the elements of the assemblage are contemporary. The condition of the bifaces is broadly similar to most of the cores, although some of the Levallois cores are fresher in condition. Purfleet remains the best dated and described British site showing application of the principles of Levallois flaking prior to MIS 8.

One of the problems with using the term simple prepared core is that this approach to core working is not rigidly defined or necessarily early, but rather reflects variation in the application of the volumetric principles of Levallois flaking. Apart from Purfleet, there are several UK sites within which simple prepared cores, or cores previously described as proto- or reduced Levallois, have been noted. Some of these are of MIS 7 age, such as Ebbsfleet in the Lower Thames Valley, where simple prepared cores are merely the application of the Levallois flaking system to small pebbles alongside full Levallois applied to larger material (Scott et al., 2010).

There are occasional sites that are pre-MIS 8 in age with individual pieces being reported as simple prepared cores. These include a piece described as coming from the 'upper brickearth' at Rickson's Pit, Swanscombe (Burchell, 1931; Roe, 1981), where the Thames terrace deposits date to MIS 11, and Baker's Farm in the Middle Thames from terrace deposits assigned to MIS 10–8 (Wymer, 1999). Although the cores are illustrated, it is not clear in which museum they are curated and cannot be physically located. They may (as with much of the Levallois material from the Middle Thames) actually come from colluvial sediments overlying and sealing the terrace (Ashton et al., 2003; Scott, 2011). Alternatively, the cores only make up a very small component of the whole assemblage, and may reflect the fortuitous end result of exploiting a nodule which favoured the application of this strategy.

There are other sites which contain simple prepared cores, but which are difficult to date. These include assemblages from the middle terraces of rivers, such as at Biddenham, Cuxton, Dunbridge and Woodston. The age of these sites could range anywhere between MIS 11 to possibly as late as MIS 7. The Cottages Site is one of several dolines at Caddington and is also undated, but contains simple prepared cores alongside classic Levallois material, including refitting sequences (Sampson, 1978; Roe, 1981).

Finally, one intriguing site is Frindsbury in the Medway Valley, but again unfortunately undated. The assemblage originally consisted of thousands of artifacts recovered from within a shallow hollow in the chalk (Cook and Killick, 1924). Only 500 artifacts now survive, but include refitting groups of flakes. White and Ashton (2003) described this material as similar to that from Purfleet, with 14 of the 16 cores from the site as simple prepared. They suggested that a sequence of large refitting flakes might provide insight into the flake production at Purfleet (Fig. 12). More recent analysis of the material suggests that five of the simple prepared cores actually result from unipolar recurrent Levallois flaking, as does the refitting sequence of five Levallois flakes. There are also three classic Levallois flakes in the assemblage. Further dating is required to understand this potentially important site.

# 4. Discussion

4.1. Characterization of these early technologies: Levallois or Levallois-type?

The hypothesis of a controlled but not fully standardized technology has sometimes been suggested for these early lithic assemblages (i.e., White and Ashton, 2003; Malinsky-Buller, 2016; Soriano and Villa, 2017). The production of classic Levallois flakes could have been accidental because these flakes did not share all the characters. Several technical features are however common to most of the assemblages:

(a) Flaking is already organized around a plane of intersection with asymmetrical faces hierarchically related.

- (b) Flaking surfaces on cores show maintenance of peripheral convexities with short or more invasive removals (distal and lateral) respecting the plane of intersection. This phase precedes the removal of the predetermined product(s). One face of the core is devoted to the debitage and the opposite face is for preparing the suitable striking platform, which is made by oblique removals and is often a partial function of the form of the support and the type of management of the flaking surface. The location of the removals for the convexities determines the shape of the predetermined products. There is a hierarchy in the management of the two faces of the core.
- (c) Flake platforms are usually plain, but occasionally dihedral and facetted.

- (d) Selection of raw material is of good quality and with a specific morphology to reduce the shapingphase.
  - (e) Unipolar and bipolar schemes, sometimes crossed, dominate over centripetal and lineal ones.
  - (f) Some cores are on flakes with evidence of a fragmentation of the reduction sequences. Some cores show a final retouch or series of small removals on a short section of the periphery of the flaking surface, perhaps recycling as a tool or for future debitage.
    - (g) Few flakes can be related to this technology in the lithic assemblages due perhaps to a higher mobility than other flakes. Alternatively, they are just more difficult to recognize than classic Levallois flakes, especially when resulting from shaping reduction sequences. *Débordant* flakes and maintenance flakes exist, although in low quantity, in the assemblages.

These features indicate a control of the core flaking surface for some pieces and of the form of the end-products with a recurrent management of the cores and a plane of intersection. If we refer to the definition (Boëda, 1986, 1993, 1995; Boëda et al., 1990), these features may be attributed to a Levallois technology similar to those observed in younger Middle Paleolithic sites.

However, two options exist: (1) these pieces are evidence of a mastery of Levallois technology, with occasional evidence prior to MIS 9 and certainlyMIS 8, (2) they are the result of accidental technological events within the main core technology; due to the low number of pieces we have to consider whether these cores and flakes result by chance without application of a predetermined concept. Our corpus

of sites can be divided in two groups: one dated confidently before MIS 9 (Cagny-la-Garenne and Guado San Nicola) and one with sites dated to MIS 9 (Orgnac 3, Cave dall'Olio and some UK sites), which is usually considered as a period where the Levallois core technology is well mastered in several European regions.

In support of the first hypothesis (Levallois core technology), these cores do not seem to be subgroups of the main core technologies of the assemblages, where there is no sign of management of the convexities. The associated core technologies can be summarized for each site:

- (a) At Cagny-la-Garenne I-II, cores were on flint nodules. The surfaces of these nodules were mainly flaked by a few unipolar and unipolar convergent removals using the natural, flat cortical convexity of the nodules. The cores are abandoned after some removals when the flaking surface becomes too flat.
- (b) At Guado san Nicola, the numerous discoidal cores were managed on slabs with typical features of a discoidal debitage (unifacial or bifacial pyramidal cores). A few cores are centripetal unifacial with a plane surface and there is no evidence of a hierarchy in the debitage and the preparation of the striking platform and the convexities.
- (c) At Orgnac 3, the lower levels mainly used flat flint slabs. Cores are thus asymmetrical with centripetal removals covering one or two surfaces, using the natural shape of the slab and the plane cortical surfaces. There is no hierarchy of the two faces and the debitage stopped when the surface became too flat, as at Cagny-la-Garenne.
- (d) At Cave dall'Ollio, cores with centripetal removals on cortical surfaces existed in the assemblages taking advantages of the natural convexities with no evidence of preparation of convexities.

In support of the second hypothesis (accidental debitage), the cores attributed to Levallois share common technological features with the main production of the assemblage. These cores could be the result of accidents supported by the small number of pieces. So, how do we interpret the innovative behaviour, which are removals and thus the management of convexities controlling the forms of the

end-products all over the flaking and maintaining the plane debitage surface? The presence and the specific location of removals of convexities helped the flaking to become independent of the geometry of the stone and of the natural convexities. The flaking could continue even if the natural convexities did not exist anymore and increased the productivity. In most of the Lower Palaeolithic core technologies, the debitage is mainly related to the stone geometry and when it was overcome, the debitage is above all discoid-type (pyramidal flaking surfaces) or polyhedral (Moncel et al., 2013, 2015). When the surface remains flat (centripetal debitage on flakes for instance), the number of removals is in general low and the debitage is uncontrolled. That raises two questions: (1) why do we observe these removals and the possible management of convexities only on some cores?, (2) are these removals accidental, used by convenience to continue the flaking for longer?

A second feature that characterizes these cores is the location of the striking platform in close relationship to the location of the assumed controlled predeterminated or "Levallois" removals on the flaking surface. The peripherical striking platform (sometimes partial) was made before the management of the flaking surface, a feature often considered as a Levallois criterion by the angle and the location of the striking platform required to maintain the flaking surface.

If we analyse these two hypothesis in regard to our two groups of sites and the general technological features of all the cores, the small number of pieces suggest some innovations among the core technology without doubt for the MIS 11-9 assemblages, but could be accidental in the MIS 12 sites.

#### 4.2. Local innovations from existing technologies?

To explain the early occasional presence of Levallois technology in some assemblages, if non accidental, two hypotheses can be investigated: introduction of external inventions (coming from one place), or local innovations (punctual experimentation of new ideas due to internal or external pressures) possibly with earlier roots, in this case mainly Acheulean-type or at least Lower Paleolithic-type technologies.

Some features in the sample of sites suggest a gradual transformation of existing core technologies with the elaboration of reduction processes, such as the use of flakes for the debitage, and, as already suggested, a fusion of elements of both façonnage (bifaces) or/and debitage (discoidal and centripetal cores; 'hierarchical cores'; Dibble and Bar-Yosef, 1995; White and Ashton, 2003; Malinsky-Buller, 2016). The main core technologies at each site share some common technological features. These common characteristics between the main core technologies and the limited evidence of Levallois or Levallois-like core technology could indicate occasional local innovation from different technological backgrounds, with possible connections between groups, and would explain the diversity of Levallois methods that is observed starting with MIS 12 (Fig. 13). Evidence of removals to manage convexities with a plane of intersection on cores, and Levallois and débordant flakes tend to distinguish the few cores and flakes from the rest of the assemblages. This may be evidence of technological innovation from a wider pool of knowledge. Levallois technology in Europe is sometimes suggested to be a progressive phenomenon preceded by a preparation phase, i.e., a proto-stage. This proto-stage could be observed with use of the hierarchical method, which could be described as intermediate, with a limited preparation of the striking platform and lateral-distal convexities (Picin, 2017). This comes under the umbrella of 'prepared core technology' in the UK. Technological data from the selected sites suggests that this interpretation cannot be applied to all sites (from MIS 12-10) as in some cases, such as Guado San Nicola, Levallois technology seems to have been mastered from an early stage.

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

The emergence of this technology could also have been associated with bifacial artifacts. An *in situ* evolution from handaxe technology has been suggested for Cagny-la-Garenne, where biface convexities were used as core faces (Tuffreau, 1987; Mellars, 1996; DeBono and Goren-Inbar, 2001; Villa, 2009; Adler et al., 2014; White and Pettitt, 2016; Tuffreau et al., 2017). In fact, the recycling of bifaces as Levallois cores is a common feature of Somme Valley sites from MIS 12 to 7 (Tuffreau et al., 2007; Lamotte and Tuffreau, 2016). However, other sites in our sample do not provide any evidence of a technical relationship of bifaces and the emergence of Levallois flaking (Fig. 14). It could be considered as a local innovative circumstance to either reduce the thickness of a biface, or produce an

expedient flake. Similar behaviour is observed in Spanish and Levantine Lower Paleolithic sites with evidence of recycling (Baena et al., 2018).

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

495

496

# 4.3. Early evidence of regionalization of local traditions across Europe?

While Adler et al. (2014) and Akhilesh et al. (2018) suggested an arrival of the technology from the Levant and Africa, the Levallois features of the sample of sites used in the current paper look similar to other early European assemblages with a Levallois component, suggesting a multiregional origin and diffusion of this technology. In addition there are other early occurrences, such as the sites of Atapuerca TD10 (MIS 11-10) and Ambrona (Middle complex, MIS 10?, various Levallois methods) in Spain, or Kesselt-Op-de Schans (MIS 11–8?, Levallois recurrent centripetal) and Petit-Spiennes (MIS 10) in Belgium, the French sites of Aldene (TU IV, MIS 9, Levallois recurrent centripetal), Petit Bost (MIS 9, level 2, various Levallois methods), and Etricourt (layer HUD, MIS 9, some Levallois flakes), and in the Netherlands Maastrich-Belvedere (possibly MIS 9, Site C, subunit IV-B (Roebroeks, 1988); Bourguignon et al., 2008; Brenet et al., 2008; Meijer and Cleveringa, 2009; Fontana et al., 2013; Lamotte and Tuffreau, 2015; Peretto et al., 2015; Di Modica et al., 2016; Di Modica and Pirson, 2016; Hérisson et al., 2016a, b; Ollé et al., 2016; Pereira et al., 2016; Rossoni-Notter et al., 2016; Baena et al., 2017; Van Baelen, 2017; Santonja et al., 2018). Moreover, in Italy the assemblage from Torre in Pietra, layer m, dated between 400 and 200 ka, indicates the application of discoid schemes associated with Levallois reduction (Villa et al., 2016). Therefore from the end of the MIS 9 there is a large diffusion of technological choices sharing common rules, but with diverse methods, rather than isolated attempts to produce standardized end-products (Hérisson et al., 2016b; Malinsky-Buller, 2016; Soriano and Villa, 2017). Levallois technology clearly becomes persistent in Europe between MIS 8 and 6 over a vast territory extending from north-western Europe to the Near East, including central Europe (Tuffreau, 1987; Rigaud, 1988; Lamotte and Tuffreau, 2001; White and Ashton, 2003; Brenet et al., 2008; Wiśniewski, 2014; Sánchez-Romero et al., 2016; Soriano and Villa, 2017). Some sites show Levallois schemes, often

accompanied by a trend towards the production of elongated blanks (Révillon, 1995; Moncel, 2003; Kozlowski, 2014). Levallois was not the only means of standardizing debitage, with for example unipolar and centripetal débitage at Cueva del Bolomor in MIS 10—9 (Blasco and Peris, 2012), the centripetal exploitation strategies seen in layer TD11 at Gran Dolina (Atapuerca) in MIS 10 (Garcia-Medrano et al., 2015), or the laminar method at Cave dall'Olio (Fontana et al., 2013).

Among these sites, technological features show early trends towards regionalization of traditions as early as MIS 8 (Picin et al., 2013; Wiśniewski, 2014; Picin, 2017), supporting the hypothesis of multiregional development and local roots. For instance, in south-east France, Orgnac 3 shows from MIS 9 and 8 an emphasis on centripetal Levallois debitage, while in south-west France uni-bipolar is dominant (Moncel et al., 2011). However, in central-eastern Europe several complexes with Levallois debitage are known from MIS 10 without any evidence of an Acheulean origin, such as Korolevo VI (Koulakovska et al., 2010) and Bechov I (Wisniewski, 2014). Other central European sites are younger, and considered to be evidence of arrivals of new populations during favourable climate with availability of good quality raw materials, such as Rheindahlen, Markkleeberg and Becoc I and IV (Wiśniewski, 2014; Picin, 2017). Similar hypotheses on the arrivals of new populations have been put forward for the Levant (Malinsky-Buller, 2016a; Shimelmitz et al., 2016; Shimelmitz and Khün, 2017). The onset of Levallois technology and all the standardized technologies can probably be explained through multiple modes of origin, dependent on area and latitude.

4.4. Explaining Levallois core technology from MIS 12-9 in Western Europe?

Flake selection and preference: Levallois end-products vs. other end-products. If of local origins, the reasons for the onset of this new core technology remain enigmatic in terms of its selection over other technologies. Levallois core technology is often a minor component of the assemblages, associated with different methods of production, such as discoidal (Bolomor, Ambrona), Kombewa, laminar (Cave dall'Olio), centripetal flaking and expedient (unifacial cores with some removals, orthogonal cores with two flaking surfaces and multidirectional cores), which produce a large variety of flakes (Ashton, 1992;

Peris et al., 2008; Santonja et al., 2018; Vaquero and Romagnoli, 2018). When discoidal and centripetal methods are used, the flaking surfaces are not hierarchically ordered and there is no evidence of management of the convexities. The debitage uses the natural forms of the blank and the previous removals for guiding the production. End-products are often thick and the form badly controlled.

Comparison of the size of Levallois-type cores and flakes to other end-products is not consistent. At Guado San Nicola, despite differences in size between preferential and recurrent flakes, Levalloisproducts are similar in size to other end-products. In contrast, at Orgnac 3, Levallois flakes are among the largest end-products of the assemblages (two groups with lengths of 30–50 mm and 65–70 mm for Levallois flakes; Fig. 15). At Purfleet, Levallois cores are slightly smaller (87 mm) than discoidal and simple prepared cores (93–97 mm). The angles and length of cutting edges on Levallois products do not seem to differ between the assemblages.

The morphology of Levallois end-products also varies between sites and regions with different quantities of flakes, elongated flakes or points. Points dominate some sites in north-west Europe compared to both flakes and points in the south.

Compared to less elaborate core technologies, a better control of the form of Levallois products and a higher productivity of Levallois cores through maintenance of convexities, where all products could be used, seems to be the main focus. This is perhaps linked to an increase in the use of mental templates by populations (Lycett et al., 2016). Villa et al. (2016) suggest that Levallois technology provided thinner products compared to Lower Paleolithic-type methods. These products did not require retouch, the form being predetermined, or the 'one tool, one task' of Douze and Delagnes (2016). The morphological regularity of flakes seems to have led to a reduction in retouched products (Eren and Lycett, 2015). For instance at Orgnac 3, the ratio of flake-tools decreases with the increase in frequency of Levallois core technology (ratio of flake tools of 6% in level 1), while the numerous small flakes (10–15 mm long) produced at the end of the Levallois reduction process and from the cores on flakes, are never retouched (Moncel et al., 2011). At Guado San Nicola, retouched Levallois flakes are extremely rare, with just a few scrapers (Peretto et al., 2015). This decrease in flake

modification could have been a cost-benefit in energy. The selection of good quality raw materials (for instance at Orgnac 3) also suggests that attention was paid to this type of debitage.

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

Functional reasons: For hunting and hafting? In parallel to the increase of hunting in subsistence strategies and some changes in land use patterns (e.g. Moncel et al., 2011), the onset of Levallois is sometimes explained by the appearance of hafting of stone points and the use of points as projectiles (Villa et al., 2009; Hardy et al., 2013; Rots, 2013; Iovita and Katsuhiro, 2016). Stone points are often considered as light penetrative tools (Knecht, 1997), and more effective than wooden spears (see Schöningen, MIS 9; Böhner et al., 2015). The early evidence of hafting at Kathu Pan (South Africa), dated to 500 ka, shows points with modification near the base and damage from hafting (Wilkins et al., 2012). The emergence of the Middle Stone Age tradition in East Africa is related to convergent tool technology (Douze and Delagnes, 2016). Modification on small flakes at Gesher Benot Yakov (GBY) is also argued to be evidence of hafting as early as 900 ka (Alperson-Afil and Goren-Inbar, 2016). However, the development of Levallois technology is only associated with the more dominant production of points in north-west Europe, rather than southern Europe. The lithic assemblages of Cagny-La-Garenne I and II and the other sites during MIS 12 clearly do not indicate an emphasis on triangular flake production, but far more oval and rectangular removals. Moreover, microwear analyses indicate that Levallois products were not systematically single purpose tools and also show that form does not necessarily indicate function. Despite little hafting evidence clearly recorded in the European Levallois (Ben-Dor et al., 2011; Rots et al., 2011; Rots, 2013; Iovita and Katsuhiro, 2016; Villa et al., 2016), some patterns show however that Neanderthals were able to haft stone tools and use glues (Mazza et al. 2006; Kozowik et al., 2017) indicating common capabilities to modern humans characterized by abstraction and planning ability (Villa et Roebroeks, 2014; Soressi, 2016).

The role of Levallois products consequently remains obscure in terms of form and awaits more microwear analyses to clarify the specific uses of these tools. At Guado san Nicola, among the 75% of the studied artifacts, only 2-4.5% show traces (n = 82). Some Levallois flakes carry microtraces with one or two different zones of use with the same activity. All show predominantly animal carcass

processing and occasionally plant use with mainly longitudinal actions from cutting (Berruti, 2017). Flakes from Levallois, other core technologies and bifaces equally show occasional wood-plant use (Peretto et al., 2015). At Orgnac 3, the development of the use of Levallois core technology is related to changes in landscape use, with seasonal and specialized occupations focused on horse hunting, such as in level 1 (Moncel et al. 2012).

# 4.5. Increase in the mental templates over time?

If control of the core knapping surface was a major initial feature, the innovation of Levallois could have been in parallel with the long process of the acquisition of Neanderthal features (accretion model; Hublin, 2009) and could be compared to similar isolated attempts that are observed in some sites older than 400 ka in Africa (Pope et al., 2017). This process could explain why this technology became dominant in many areas through the Middle Paleolithic and why it did not emerge earlier (e.g., Lycett and Eren, 2013).

For instance, the Oldowan assemblage of Nyabusosi (Uganda) dated to 1.5 Ma shows the hierarchical relationship of core surfaces (Texier, 1995). The Early Acheulean site of Peninj (1.6–1.2 Ma, Tanzania) shows some evidence of the preparation of core convexities, as at Wonderwerk (800–500 ka, South Africa), GBY with giant Kombewa and 'Levallois' flakes (900–800 ka, Israel), or la Noira (700 ka, France; Texier and Roche, 1995; de la Torre et al., 2003; Tyron et al., 2006; Moncel et al., 2013; Chazan, 2015; Leader et al., 2018). In the past, the different Victoria West methods have been considered as para-, proto- or pre-Levallois evidence, with large, wide asymmetrical flakes removed through planning from the core edge by radial or centripetal flaking (Bordes, 1950; McNabb, 2001; Mourre, 2006; Sharon et al., 2009). Similarly, the Tabelbala-Tachenghit method or the Kombewa technique used the bulb of the ventral face of a flake and were described as a 'preferential-flake method' (Boëda, 1995). The onset of actual Levallois technology is also observed in East Africa by early modern humans with embedded roots as early as 500 ka with local, gradual changes in the Middle

Stone Age (Douze and Delagnes 2016; Deino et al., 2018; Potts et al., 2018). Meanwhile at Misliya cave (Israël), at around 200 ka, there is full Levallois with modern human remains (Hershkovitz et al., 2018).

4.6. Emergence of Neanderthals or adaptation of hominins to climatic and environmental changes?

If we consider the paleoanthropological record in Europe from MIS 13 to 9, data indicate complexity in the acquisition over time of Neanderthal characteristics among *H. heidelbergensis* and Middle Pleistocene populations (Hublin, 2009; Manzi et al., 2010). In western Europe, the anatomical diversity of fossils suggests pre-Neanderthal regional groups, perhaps persistent forms of *H. heidelbergensis* (Ceprano skull dated to 350 ka), as shown by the genetic variability (Rightmire, 2008; Fabre et al., 2009; Manzi et al., 2010; Walker et al., 2011; Stringer, 2012; Meyer et al., 2016; Rightmire, 2017).

Neanderthal characteristics were evolving in Europe as far back as MIS 11 and possibly earlier (Hublin, 2009; Stringer, 2012). An earlier divergence time (>430 ka) between Neanderthals and Denisovans was inferred from the nuclear DNA sequence from Sima de los Huesos, whereas mtDNA places these populations closer to one another (Meyer et al. 2014). During the time span of MIS 12 to MIS 7 (ca. 460–250 ka), Neanderthal populations may have experienced a period of isolation, but contact with African lineages postdating the divergence from the Denisovans is also suggested (Arsuaga et al., 2014; Meyer et al., 2016; Hublin et al., 2017; Richter et al., 2017; Bermúdez de Castro et al., 2019).

At the moment, correlations between types of hominin and technological innovations are not evident (see Levallois evidence in the Ceprano basin contemporaneous with the skull; Pereira et al., 2018). Technological convergence could exist in many places with a variety of hominins, such as *H. heidelbergensis*, *Homo neanderthalensis* and *Homo* sp. (DeBono and Goren-Inbar, 2001; Tyron, 2006; Adler et al., 2014). Neanderthal anatomical features developed in parallel, as with other hominins at this time, with an increase in brain size, but also changes in life history, such as an extended childhood and an adolescent phase (Kyriacou and Bruner, 2011). This allowed an increase in the capability to transmit more complex technological behaviors through social learning (Nowell and White 2010;

White et al. 2011). Similar developments in East Africa could explain the onset of Levallois technology among modern human populations. If we consider the low number of 'Levallois' pieces or the eventuality of an event by chance in parallel to some innovations (hierarchical organization on some cores), the phenomenon would indicate (1) a progressive development of the use of mental templates and (2) a technological shift in some areas. This progressive development could have found its roots among *H. heidelbergensis* (and Middle Pleistocene populations), not just after the speciation to Neanderthals.

The numerous paleoclimatic archives show a transition from 1.25 Ma up to 0.7 Ma (Mid-Pleistocene Revolution) with a change of the dominant periodicity of climate cycles from 41 ka to 100 ka. Combining different archives over the last 800 ka, some particularly marked interglacials (MIS 19, 15, 11, 9 and 5) and glacial maxima (MIS 16, 12 and 2) have been identified (Jouzel et al., 2007). Some of the earliest Levallois evidence is during MIS 11 (Schreve, 2001; Geyh and Müller, 2005; Nitychoruk et al., 2006; Roe et al., 2009; Rohling et al., 2010; Blain et al., 2015; Limondin-Lozouet et al., 2015; Picin, 2017). This period of time (post MIS 12, MIS 11) is crucial, characterized by a large biodiversity, largescale faunal dispersion associated with the regionalization of mammal communities and hominin morphological variability (Stiner, 2002; Bar-Yosef and Belmaker, 2011; Dennell et al., 2011; Palombo, 2015). Such a long-lasting interglacial (MIS 11) after a harsh glacial (MIS 12) could have favored more sustained vegetational systems and hence more stable hominin occupation and connections between groups in Europe dependent on latitude (Guthrie, 2001; Ashton et al., 2017, 2018). However, climate does not seem to play a great role in the earliest onset of Levallois from MIS 12-9. It appears during both cold and temperate events in various areas. But climate change was certainly important for diffusion and some breaks in occupation in some areas due to climatic constraints, and could explain the introduction of this new technology such as the UK during periods of lowered sea levels.

673

674

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

# 5. Conclusions

Between MIS 12 and 9, elements of Levallois technology, some probably intentional, are found intermittently over a vast area in both northern and southern Europe, and are sometimes accompanied by a trend towards the production of elongated blanks. The lithics seem to be evidence of a technological shift in some areas rather than production by chance. Levallois core technology before the end of MIS 9 to the beginning of MIS 8 remains rare (Fig. 1). From MIS 8, it was diffused all over Europe and appears to have been a phase of diversification rather than the initial stage. A discontinuity between the earliest and youngest phases during MIS 7-6 is open to question, as in East Africa with the isolated early appearance of laminar debitage at 500 ka (Roure Johnson and McBrearty, 2010). Is there a progressive phenomenon preceded by a preparation phase, i.e., a protostage ('prepared core technology' and invasive removals on bifaces)? Due to the small number of sites and their distribution over a large area, we do not know if there is clear evidence of a transition. It is not known whether the isolated evidence of Levallois is due only to innovation (in situ), with multiple convergence or roots in the Acheulean in a variety of environments and geographic situations, or also to invention from outside Europe that by its diffusion may have certainly been enhanced by contacts and exchanges of experiences between different groups (Foley and Mirazón Lahr, 1997; Hublin, 2009; Stringer, 2012).

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

The only certainty is the apparent parallel development with the earliest appearance of Levallois core technology and some behavioral changes affecting subsistence, land use and mobility of populations during and after MIS 12 in Europe as shown by the longer distances for raw material procurement and more specialized hunting. Cultural and technical expertise of these populations allowed integration of strategies for making, using, transporting and discarding tools, and the materials needed for their manufacture and maintenance. However, the development of the Early Middle Paleolithic through Europe is not only related to the innovation of Levallois, which appeared later in some regions, indicating a diversity of trajectories.

To conclude, it is important to note that these isolated onsets of Levallois are associated with an increase of archaeological data and human activity all over Eurasia after the Anglian or Elsterian

glaciation (MIS 12), which is considered as a major turning point. The increase in the quantity of sites raises the question of whether this is due to better preservation or reflects larger populations, while genetic data indicates a decrease of the population size after 500 ka (Meyer et al., 2014). Cultural complexity in the form of Levallois technology does not necessarily reflect demographic expansion (Vaesen and Collard 2016), just as an increase in population does not always lead to diffusion of behavioural changes if populations are poorly connected (Premo and Hublin, 2009;Roger, 2017). Data on the Early Middle Paleolithic (MIS 8-6) indicate both a large diversity of technical expertise among groups and some trends of regionalization before a more pronounced regionalization of traditions during the Late Middle Paleolithic (MIS 5-3; Baena et al., 2017; Carmignani et al., 2017). Middle Paleolithic features emerged as a mosaic as early as MIS 12, including a more complex management of local resources and the use of long-distance lithic raw materials in short-term recurrent occupations. This suggests local innovations. The degree of mobility of human groups and connections between different groups is difficult to estimate, depending on topography and climate, and estimating the type and size of the European population is open to discussion, ranging from a well-connected metapopulation or to more isolated networks of sites with small populations (Bocquet-Appel and Degioanni, 2013; Collard et al., 2013; Derex and Boyd, 2016; Fogarty and Creanza, 2017; Grove, 2016, 2017; Ríos et al., 2019). Finally, we have to keep in mind that from at least 500 ka, two technological worlds existed in Eurasia with western Europe standing in contrast to a large area from central Europe to central Asia. In western Europe, the Acheulean and other Lower Paleolithic behaviors are commonly referred to H. heidelbergensis and early Neanderthals from 700 ka, while in central Europe, the traditions are considered as 'Micro and Pre-Mousterian' without bifacial technology (Kozlowski, 2014; Moncel et al., 2015; Golovanova and Doronichev, 2017). Regional differences persisted in Middle Paleolithic traditions from MIS 6 between these two areas (for instance Micoquian in Central Europe) and perhaps between populations (human colonization from Asia from the Middle Pleistocene?) despite a common technical background in core technologies. This feature is unexplained so far and may be due to at

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

least some late exchanges between populations. The pre-existing Lower Paleolithic certainly had a structuring effect on the different adaptive options selected by hominin groups over Europe. The key question now is why Levallois technology became the dominant technological strategy even if other technologies were also used during the Early Middle Paleolithic.

731

732

727

728

729

730

#### References

- 733 Adler, D.S., Wilkinson, K.N., Blockley, S., Mark, D.F., Pinhasi, R., Schmidt-Magee, B.A., Nahapetyan, S.,
- 734 Mallol, C., Berna, F., Glauberman, P.J., 2014. Early Levallois technology and the Lower to Middle
- Paleolithic transition in the Southern Caucasus. Science 345, 1609–1613.
- 736 Akhilesh, K., Pappu, S., Rajapara, H. M., Gunnell, Y., Shukla, A.D., Singhvi, A.K., 2018. Early Middle
- 737 Palaeolithic culture in India around 385–172 ka reframes Out of Africa models. Nature 554, 97.
- 738 Alperson-Afil, N., Goren-Inbar, N., 2016. Acheulian hafting: Proximal modification of small flint flakes
- at Gesher Benot Ya'aqov, Israel. Quaternary International 411, 34-43.
- 740 Antoine, P., Limondin-Lozouet, N., Auguste, P., Lamotte, A., Bahain, J. J., Falguères, C., Laurent, M.,
- 741 Coudret, P., Locht, J-L., Depaepe, P., Fagnart, J. P., Fontugne, M., Hatté, C., Mercier, N., Frechen,
- M., Moigne, A-M., Munaut, A-V., Ponel, P., Rousseau, D-D., 2003. Paléoenvironnements
- 743 pléistocènes et peuplements paléolithiques dans le bassin de la Somme (nord de la France). Bulletin
- de la Société préhistorique française 100(1), 5-28.
- Antoine, P., Lozouet, N. L., Chaussé, C., Lautridou, J. P., Pastre, J. F., Auguste, P., Bahain, J-J., Falguères,
- 746 C., Galheb, B., 2007. Pleistocene fluvial terraces from northern France (Seine, Yonne, Somme):
- 747 synthesis, and new results from interglacial deposits. Quaternary Science Reviews 26(22-24), 2701-
- 748 2723.
- Antoine P., Moncel M.-H., Limondin-Lozouet N., Locht J.-L., Bahain J.-J., Moreno D., Voinchet P.,
- Auguste P., Stoetzel E., Dabkowski J., Bello S. M., Parfitt S.A., Tombret O., Hardy B., 2016.
- 751 Palaeoenvironment and dating of the Early Acheulean from the type area of the (River) Somme

- basin (Northern France): new discoveries from the high terrace at Abbeville-Carrière Carpentier.
- 753 Quaternary Sciences Reviews 149, 338-371.
- 754 Aouraghe, H., 1999. Reconstitution du paléoenvironnement par les grands mammifères : les faunes
- du Pléistocène moyen d'Orgnac 3 (Ardèche, France), L'Anthropologie 103(1), 177-184.
- 756 Arsuaga, J.L., Martínez, I., Arnold, L.J., Aranburu, A., Gracia-Téllez, A., Sharp, W.D., Quam, R.M.,
- Falguères, C., Pantoja-Pérez, A., Bischoff, J., Poza-Rey, E., Parés, J.M., Carretero, J.M., Demuro, M.,
- 758 Lorenzo, C., Sala, N., Martinón-Torres, M., García, N., Alcázar de Velasco, A., Cuenca-Bescós, G.,
- 759 Gómez-Olivencia, A., Moreno, D., Pablos, A., Shen, C.-C., Rodríguez, L., Ortega, A.I., García, R.,
- 760 Bonmatí, A., Bermúdez de Castro, J.M., Carbonell, E., 2014. Neandertal roots: Cranial and
- 761 chronological evidence from Sima de los Huesos. Science 344, 1358-1363.
- Ashton, N., 1992. The High Lodge flint industries. In: Ashton, N., Cook, J, Lewis S.G., Rose, J., (Eds.)
- High Lodge: excavations by G. de G. Sieveking, 1962-8 and J. Cook, 1988 British Museum Press.
- 764 London, pp. 124-163.
- Ashton, N., 2017. Landscapes of habit and persistent places during MIS 11 in Europe: a return
- journey from Britain. In: Pope, M., McNabb, J., Gamble, C. eds. Crossing the Human Threshold.
- 767 Routledge, New York, pp. 164-186.
- Ashton, N., Harris, C.R., Lewis, S.G. 2018. Frontiers and route-ways from Europe: the Early Middle
- 769 Palaeolithic of Britain. Journal of Quaternary Science 33, 194-211.
- Baena, J., Moncel, M.H., Cuartero, F., Navarro, M.G.C., Rubio, D., 2017. Late Middle Pleistocene
- 771 genesis of Neanderthal technology in Western Europe: The case of Payre site (south-east France).
- 772 Quaternary International 436, 212-238.
- 773 Baena, J., Navas, C. T., Sharon, G., 2018. Life history of a large flake biface. Quaternary Science
- 774 Reviews 190, 123-136.
- 775 Bar-Yosef, O., Belmaker, M., 2011. Early and Middle Pleistocene Faunal and hominins dispersals
- through Southwestern Asia. Quaternary Science Reviews 30, 1318–1337.

- Ben-Dor, M., Gopher, A., Hershkovitz, I., Barkai, R., 2011. Man the fat hunter: the demise of *Homo*erectus and the emergence of a new hominin lineage in the Middle Pleistocene (ca. 400 kyr)
- 779 Levant. PLoS One 6, e28689.
- 780 Bermúdez de Castro, J.M., Martinón-Torres, M., Martínez de Pinillos, M., García-Campos, C.,
- 781 Modesto-Mata, M., Martín-Francés, L., Arsuaga, J.L., 2019. Metric and morphological comparison
- between the Arago (France) and Atapuerca-Sima de los Huesos (Spain) dental samples, and the
- origin of Neanderthals. Quaternary Science Reviews 217, 45-61.
- 784 Berruti, G.L., 2017. Use-wear analysis of discoid-conception lithic industries. PhD. Dissertation,
- 785 Universidade de Tras-os-Montes e Alto Douro.
- 786 Bischoff, J.L., Williams, R.W., Rosenbauer, R.J., Aramburu, A., Arsuaga, J.L., García, N., Cuenca-Bescós,
- G., 2007. High-resolution U-series dates from the Sima de los Huesos hominids yields  $600_{-66}$ + $\sim$
- 788 kyrs: implications for the evolution of the early Neanderthal lineage. Journal of Archaeological
- 789 Science 34, 763–770.
- 790 Bisi, F., Cremaschi, M., Peretto, C., 1982. Le industrie del Paleolitico inferiore del conoide
- 791 pleistocenico del Torrente Idice (Bologna). In: Bisi, F., Cremashi, M., Peretto, C. (Eds), Atti della
- 792 XXIII Riunione Scientifica dell'Istituto Italiano di Preistoria e Protostoria (Bologna). IIPP, Firenze,
- 793 Stampa, pp. 259-271.
- 794 Blain, H.-A., Lozano-Fernández, I., Ollé, A., Rodríguez, J., Santonja, M., Pérez-González, A., 2015. The
- 795 continental record of Marine Isotope Stage 11 (Middle Pleistocene) on the Iberian Peninsula
- 796 characterized by herpetofaunal assemblages. Journal of Quaternary Science 30, 667–678.
- 797 Blasco, R., Peris, J.F., 2012. A uniquely broad spectrum diet during the Middle Pleistocene at Bolomor
- 798 Cave (Valencia, Spain). Quaternary International 252, 16-31.
- 799 Blasco, R., Rosell, J., Domínguez-Rodrigo, M., Lozano, S., Pastó, I., Riba, D., Vaquero, M., Peris, J.F.,
- Arsuaga, J.L., de Castro, J.M.B., Carbonell, E., 2013. Learning by heart: cultural patterns in the
- faunal processing sequence during the Middle Pleistocene. PLoS One 8: e55863.

802 Bocquet-Appel, J.-P., Degioanni, A., 2013. Neanderthal demographic estimates. Current 803 Anthropology 54, S202-S213 Boëda, E., 1986. Approche technologique du concept Levallois et évaluation de son champ 804 805 d'application. Étude de trois gisements saaliens et weichsélien de la France septentrionale. Ph.D. 806 Dissertation, Université de Paris X. 807 Boëda, E., 1993. Le débitage discoïde et le débitage Levallois récurrent centripède. Bulletin de la 808 Société Préhistorique Française 90, 392-404. 809 Boëda, E., 1995. Levallois: A volumetric construction, methods, a technique. In: Dibble, H.L., Bar-810 Yosef, O. (Eds.), The Definition and Interpretation of Levallois Technology. Prehistory Press 811 Madison, New York, pp. 41–69. 812 Boëda, E., Geneste, J-M., Meignen, L., 1990. Identification de chaines operatoires lithiques du 813 Paleolithique ancien et moyen. Paléo 2, 43-79. 814 Böhner, U., Serangeli, J., Richter, P., 2015. The Spear Horizon: First spatial analysis of the Schöningen 815 site 13 II-4. Journal of Human Evolution 89, 202-213. 816 Bolton, L., 2015. Assessing the origins of Levallois through Lower Palaeolithic core variation: A 817 comparative study of Simple Prepared Cores in northwest Europe. Ph.D. Dissertation, University 818 of Southampton. 819 Bordes, F., 1950. Principes d'une méthode d'étude des techniques de débitage et de la typologie du 820 Paléolithique ancien et moyen. Anthropologie 54, 19–34. 821 Boucher de Perthes, J., 1857. Antiquités celtiques et antédiluviennes, Mémoire sur l'industrie primitive et les arts à leur origine, T. 2, Jung-Treuttel, Paris. 822 823 Bourguignon, L., Djema, H., Bertran, P., Lahaye, C., Guibert, P., 2008. Le gisement Saalien de Petit-824 Bost (Neuvic, Dordogne) à l'origine du Moustérien d'Aquitaine ? In: Jaubert, J., Bordes, J-G., Ortega, I. (Eds.), Les Sociétés du Paléolithique dans un Grand Sud-Oest de la France. Société 825 826 Préhistorique Française XLVII, 41-57.

827 Brenet, M., Folgado, M., Lenoble, A., Bertrán, P., Vieillevigne, E., Guibert, P., 2006. Interprétation de 828 la variabilité technologique de deux industries du Paléolithique moyen ancien du Bergeracois. In 829 Les sociétés paléolithiques d'un grand Sud-Ouest: nouveaux gisements, nouvelles méthodes, nouveaux résultats. Journées de la Société Préhistorique Française 47, 57-81. 830 831 Breuil H., Kelley H., 1954. Le Paléolithique ancien. Abbevillien. Clactonien. Acheuléen. Levalloisien. 832 Bulletin de la Société Préhistorique Française, 51(8), 1–18. 833 Bridgland, D.R., 1994. Quaternary of the Thames. Chapman and Hall, London. 834 Burchell, J.P.T., 1931. Early Neoanthropic Man and his relation to the Ice Age. Proceedings of the 835 Prehistoric Society of East Anglia 6, 253-303. Carmignani L., Moncel M-H., Fernandes P., Wilson L., 2017. Technological variability during the Early 836 837 Middle Palaeolithic in Western Europe. Reduction systems and predetermined products at the 838 Bau de l'Aubesier and Payre (South-East France). PLoS One 12, e0178550. 839 Chazan, M., 2015. Technological trends in the Acheulean of Wonderwerk cave, South Africa. African 840 Archaeological Review 32, 701-728. 841 Clark, G., 1969. World Prehistory: A New Outline. Cambridge University Press, London. 842 Collard, M., Buchanan, B., O'Brien, M.J., 2013. Population size as an explanation for patterns in the Paleolithic archaeological record. Current Anthropology 54(S8), S388–S396. 843 844 Combier, J., 1967. Le Paléolithique de l'Ardèche dans son cadre bioclimatique. Publications de 845 l'Institut de Préhistoire, Université de Bordeaux, Bordeaux. 846 Commont, V., 1909. Saint-Acheul et Montières: notes de géologie, de paléontologie et de préhistoire. 847 Mémoire de la Société géologique du Nord 6, 1-68. 848 Conard, N.J., Serangeli, J., Böhner, U., Starkovich, B.M., Miller, C.E., Urban, B., van Kolfschoten, T., 849 2015. Excavations at Schöningen and paradigm shifts in human evolution. Journal of Human 850 Evolution 89, 1–17.

851 Cook, W.H., Killick, J.R., 1924. On the discovery of a flint-working site of Palaeolithic date in the 852 Medway Valley at Rochester, Kent, with notes on the drift stages of the Medway. Proceedings of 853 the Prehistoric Society of East Anglia 4, 133-149. 854 DeBono H., Goren-Inbar, N., 2001. Note on a link between Acheulian handaxes and the Levallois 855 method. Journal of the Israel Prehistoric Society 31, 9-23. 856 De Mortillet G., 1883. *La prehistorique antiquité de l'homme*. Reinwald C. ed., Paris. 857 Debard, E., Pastre, J.F., 1988. Un marqueur chronostratigraphique du Pléistocene moyen à la 858 periphérie du Massif Central: la retombée à clinopyroxène vert du Sancy dans le site acheuléen 859 d'Orgnac III (Bas-Vivarais, SE France). Comptes Rendu de l'Académie des Sciences Paris 306, 860 1515–1520. De la Torre de la, I., Mora, R., Domínguez-Rodrigo, M., de Luque, L., Alcalá, L., 2003. The Oldowan 861 862 industry of Peninj and its bearing on the reconstruction of the technological skills of 863 LowerPleistocene hominids. Journal of Human Evolution 44, 203-224. 864 Dennell, R.W., Martinón-Torres, M., Bermúdez de Castro, J.M., 2011. Hominin variability, climatic 865 instability and population demography in Middle Pleistocene Europe. Quaternary Science Reviews 866 30, 1511-1524. 867 Derex M., Boyd R., 2016. Partial connectivity increases cultural accumulation within groups. 868 Proceedings of the National Academy of Sciences USA 113, 2982-2987. 869 Dibble, H. L., Bar-Yosef, O., 1995. The definition and interpretation of Levallois technology. Prehistory 870 Madison Press, Wisconsin. 871 Di Modica, K., Pirson, S., 2016. The Lower to Middle Palaeolithic transition and the onset of prepared-872 core technologies in Belgium. Quaternary International 411, 95-107. 873 Di Modica, K., Toussaint, M., Abrams, G., Pirson, S., 2016. The Middle Palaeolithic from Belgium: 874 Chronostratigraphy, territorial management and culture on a mosaic of contrasting environments.

875

Quaternary International 411, 77-106.

876 Deino, A. L., Behrensmeyer, A. K., Brooks, A. S., Yellen, J. E., Sharp, W. D., Potts, R., 2018. Chronology 877 of the Acheulean to Middle Stone Age transition in eastern Africa. Science 360, 95-98. Douze, K., Delagnes, A., 2016. The pattern of emergence of a Middle Stone Age tradition at 878 879 Gademotta and Kulkuletti (Ethiopia) through convergent tool and point technologies. Journal of 880 Human Evolution 91, 93-121. 881 El Hazzazi, N., 1998. Paléoenvironnement et chronologie des sites du Pleistocène moyen et 882 supérieur, Orgnac 3, Payre et Abri des Pêcheurs (Ardèche, France) d'après l'étude des rongeurs. 883 Phd Museum National d'Histoire Naturelle, Paris. 884 Endicott, P., Ho, S.Y.W., Stringer, C., 2010. Using genetic evidence to evaluate four 885 palaeoanthropological hypotheses for the timing of Neanderthal and modern human origins. 886 Journal of Human Evolution 59, 87–95. 887 Eren, M. I., Lycett, S. J., 2016. A statistical examination of flake edge angles produced during 888 experimental lineal Levallois reductions and consideration of their functional implications. Journal 889 of Archaeological Method and Theory 23, 379-398. 890 Falguères, C., Shen, G., Yokoyama, Y., 1988. Datation de l'aven d'Orgnac III: comparaison par les 891 méthodes de la resonnance de spin électronique (ESR) et du déséquilibre des familles de 892 l'Uranium. Anthropologie 92, 727–730. 893 Farabegoli, F., Onorevoli, G., 1996. Il margine appenninico emiliano-romagnolo durante il 894 Quaternario: stratigrafia ed eventi. In: Lenzi, F., Nenzioni, G. (Eds.). Lettere di pietra. I depositi 895 pleistocenici : sedimenti , industrie e faune del margine appenninico bolognese. Editrice 896 Compositori, Bologna, pp. XXXIX-LIV. 897 Farabegoli, E., Lenzi, F., Nenzioni, G., Onorevoli, G., Peretto, C., 2000. Lithostratigraphie et evolution 898 des industries du Paleolithique inférieur et moyen a l'est de Bologne (Italie). In: Farabegolie, E., Onorevoli, G., Lenzi, F., Nenzioni, G., Peretto, C. (Eds), Actes du congrès international sur science 899 900 et technologie pour la sauvegarde du patrimoine culturel dans les Pays du bassin méditerranée, 901 vol. II. Elsevier, Paris, pp. 1179-1188

902	Fabre, V, Condemi, S, Degioanni, A., 2009. GeneticeEvidence of geographical groups among
903	Neanderthals. PLoS One 4, e5151.
904	Fogarty, L., Creanza, N., 2017. The niche construction of cultural complexity: interactions between
905	innovations, population size and the environment. Philosophical Transactions of the Royal Society
906	B 372, 20160428.
907	Foley, R., Lahr, M. M., 1997. Mode 3 technologies and the evolution of modern humans. Cambridge
908	Archaeological Journal 7, 3-36.
909	Fontana, F., Nenzioni, G., Peretto, C., 2010, The southern Po plain area (Italy) in the mid-late
910	Pleistocene: human occupation and technical behaviours. Quaternary International 223-224, 465
911	471.
912	Fontana, F., Moncel, MH., Nenzioni, G., Onorevoli, G., Peretto, C., Combier, J., 2013. Widespread
913	diffusion of technical innovations around 300,000 years ago in Europe as a reflection of
914	anthropological and social transformations? New comparative data from the western
915	Mediterranean sites of Orgnac (France) and Cave dall'Olio (Italy). Journal of Anthropological
916	Archaeology 32, 478-498.
917	Forsten, A., Moigne, A-M., 1988. The horse from the Middle Pleistocene of Orgnac 3 (Ardèche-
918	France). Faune et Archéologie, Quaternaire 9(4), 315-323.
919	Gauthier, A., 1992. Paléoenvironnement du Pléistocène moyen dans le sud de la France. Apport et
920	limite de l'analyse pollinique de trois sites préhistoriques : Caune de l'Arago, Orgnac 3, grotte du
921	Lazaret. Phd Muséum National d'Histoire Naturelle, Paris.
922	García-Medrano, P., Ollé, A., Mosquera, M., Cáceres, I., Carbonell, E., 2015. The nature of
923	technological changes: The Middle Pleistocene stone tool assemblages from Galería and Gran
924	Dolina-subunit TD10. 1 (Atapuerca, Spain). Quaternary International 368, 92-111.
925	Geyh, M.A., Müller, H., 2005. Numerical 230Th/U dating and a palynological review of the
926	Holsteinian/Hoxnian Interglacial. Quaternary Science Reviews 24, 1861–1872.

927 Gowlett, J.A.J., 2016. The discovery of fire by humans: a long and convoluted process. Philosophical 928 Transactions of the Royal Society B 371, 20150164-. 929 Green, R.E., Krause, J., Briggs, A.W., Maricic, T., Stenzel, U., Kircher, M., Patterson, N., Heng Li, 930 Weiwei Zhai, Hsi-Yang Fritz, M., Hansen, N.F., Durand, E.Y., Malaspinas, A-S., Jensen, J.D., 931 Marques-Bonet, T., Alkan, C., Prüfer, K., Meyer, M., Burbano, H.A., Good, J.M., Schultz, R., Aximu-932 Petri, A., Butthof, A., Höber, B., Höffner, B., Siegemund, M., Weihmann, A., Nusbaum, C., Lander, 933 E.S., Russ, C., Novod, N., Affourtit J., Egholm, M., Verna, C., Rudan, P., Brajkovic, D., Kucan, Z., 934 Gušic, I., Doronichev, V.B., Golovanova, L.V., Lalueza-Fox, C., de la Rasilla, Fortea, J., Rosas, A., 935 Schmitz, R.W., Johnson, P.L.F., Eichler, E.E., Falush, D., Birney, E., Mullikin, J.C., Slatkin, M., 936 Nielsen, R., Kelso, J., Lachmann, M., Reich, D., Pääbo, S., 2010. A draft sequence of the 937 Neanderthal genome. Science 328, 710–722. 938 Grove, M., 2016. Population density, mobility, and cultural transmission. Journal of Archaeological 939 Science 74, 75-84. 940 Grove, M., 2018. Hunter-gatherers adjust mobility to maintain contact under climatic variation. 941 Journal of Archaeological Science: Reports 19, 588–595. 942 Guerin, C., 1980. Les Rhinocéros (Mammalia, Perissodactyla) du Miocène terminal au Pléistocène 943 supérieur en Europe occidentale. Comparaison avec les espèces actuelles. Documents du 944 laboratoire de géologie de Lyon, 79, Lyon. 945 Guthrie, R. D. 2001. Origin and causes of the mammoth steppe: a story of cloud cover, woolly 946 mammal tooth pits, buckles, and inside-out Beringia. Quaternary Science Reviews 20, 549-574. 947 Hardy, B.L., Moncel, M-H., Daujeard, C., Fernandes, P., Béarez, P., Desclaux, E., Chacon Navarro, M. 948 G., Puaud, S., Gallotti, R., 2013. Impossible Neanderthals? Making string, throwing projectiles and 949 catching small game during Marine Isotope Stage 4 (Abri du Maras, France). Quaternary Science 950 Reviews 82, 23-40.

- 951 Hérisson, D., Coutard, S., Goval, E., Locht, J.-L., Antoine, P., Chantreau, Y., Debenham, N., 2016a. A
- 952 new key-site for the end of Lower Palaeolithic and the onset of Middle Palaeolithic at Etricourt-
- 953 Manancourt (Somme, France). Quaternary International 409, 73–91.
- 954 Hérisson, D., Brenet, M., Cliquet, D., Moncel, M. H., Richter, J., Scott, B., Baelen, A.V., Di Modica,
- 955 K., De Loecker, D., Ashton, N., Bourguignon, L., Delagnes, A., Faivre, J-P., Folgado-Lopez, M., Locht,
- 956 J-L., Pope, M., Raynal, J-P., Roebroeks, W., Santagata, C., Turq, A., Van Peer, P., 2016b. The
- 957 emergence of the Middle Palaeolithic in north-western Europe and its southern fringes.
- 958 Quaternary international 411, 233-283.
- Hershkovitz, I., Weber, G.W., Quam, R., Duval, M., Grün, R., Kinsley, L., Ayalon, A., Bar-Matthews, M.,
- Valladas, H., Mercier, N., Arsuaga, J.L., Martinón-Torres, M., Bermúdez de Castro, J.M., Fornai, C.,
- 961 Martín-Francés, L., Sarig, R., May, H., Krenn, V.A., Slon, V., Rodríguez, L., García, R., Lorenzo, C.,
- 962 Carretero, J.M., Frumkin, A., Shahack-Gross, R., Bar-Yosef Mayer, D.E., yaming Cui, Xinzhi Wu,
- 963 Peled, N., Groman-Yaroslavski, I., Weissbrod, L., Yeshurun, R., Tsatskin, A., Zaidner, Y., Weinstein-
- 964 Evron, M., 2018. The earliest modern humans outside Africa. Science 359, 456-459.
- 965 Hublin, J., 1998. Climatic changes, paleogeography, and the evolution of the Neandertals. In:
- 966 Akazawa, T., Aoki, K., Bar-Yosef, O. (Eds.), Neandertals and Modern Humans in Western Asia.
- 967 New-York Press, Springer, pp. 295–310.
- 968 Hublin, J.-J., 2009. The origin of Neandertals. Proceedings of the National Academy of Sciences USA
- 969 106, 16022–16027.
- 970 Hublin, J.-J., Pääbo, S., 2005. Neandertals. Current Biology 16, 113–114.
- 971 Hublin, J. J., Ben-Ncer, A., Bailey, S.E., Freidline, S.E., Neubauer, S., Skinner, M.M., Bergmann, I., Le
- 972 Cabec, A., Benzzi, S., Harvati, K., Gunz, P., 2017. New fossils from Jebel Irhoud, Morocco and the
- pan-African origin of *Homo sapiens*. Nature 546, 289.
- 974 Iovita, R., Katsuhiro, S. (Eds.), 2016. Multidisciplinary Approaches to the Study of Stone Age
- 975 Weaponry. Springer, New York.

- 976 Jeannet, M., 1981. Les rongeurs du gisement acheuléen d'Orgnac 3 (Ardeche). Bulletin de la Société
- 977 Linnéenne de Lyon 50-2, 49-71.
- Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B.,
- Nouet, J., Barnola, J.M., Chappellaz, J., Fischer, H., Gallet, J.C., Johnsen, S., Leuenberger, M.,
- 980 Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander,
- J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J.P., Stenni, B., Stocker, T.F., Tison,
- 982 J.L., Werner, M., Wolff, E.W., 2007. Orbital and millennial antarctic climate variability over the
- 983 Past 800,000 Years. Science 317, 793–796.
- 984 Khatib, S., 1994. Datation des cendres volcaniques et analyses géochimiques du remplissage
- 985 d'Orgnac 3 (Ardèche, France), Quaternaire 5(1), 13-22.
- 986 Knecht, H., 1997. Projectile points of bone, antler, and stone. In: Knecht, H. (Ed.), Projectile
- 987 Technology. IDCA, Boston, pp. 191-212.
- 988 Koulakovska, L., Usik, V., Haesaerts, P., 2010. Early Paleolithic of Korolevo site (Transcarpathia,
- 989 Ukraine). Quaternary International 223, 116-130.
- 990 Kozłowski, J. K., 2014. Middle Palaeolithic variability in central Europe: Mousterian vs Micoquian.
- 991 Quaternary International 326, 344-363.
- 992 Kozowyk, P. R. B., Soressi, M., Pomstra, D., Langejans, G. H. J., 2017. Experimental methods for the
- Palaeolithic dry distillation of birch bark: implications for the origin and development of
- Neandertal adhesive technology. Scientific Reports 7, 8033.
- 995 Krings, M., Stone, A., Schmitz, R.W., Krainitzki, H., Stoneking, M., Pääbo, S., 1997. Neanderthal DNA
- sequences and the origin of modern humans. Cell 90, 19–30.
- 997 Kyriacou, A., Bruner, E., 2011. Special Issue. Brain evolution, innovation, and endocranial variations in
- 998 fossiles Hominids. Special Issue: Innovation and the Evolution of Human Behavior.
- 999 PaleoAnthropology 130-143.
- 1000 Lamotte, A., 1994. Les industries à bifaces du Pléistocène moyen de l'Europe du Nord-Ouest :
- données nouvelles des gisements du bassin de l'Escaut, de la Somme et de la Baie de Saint-Brieuc
- 1002 Ph.D. Dissertation, Université de Lille.

1003 Lamotte, A., Tuffreau, A., 2001. Les industries lithiques de Cagny-la-Garenne II (Somme, France). In: 1004 Tuffreau, A. (Ed.), L'Acheuléen dans la Vallée de la Somme et Paléolithique Moyen dans le Nord 1005 de la France : Données Récentes. CERP, Université de Lille, Lille, pp. 59–91. 1006 Lamotte, A., Tuffreau, A., 2016. Acheulean of the Somme basin (France): Assessment of lithic changes 1007 during MIS 12 to 9. Quaternary International 409, 54–72. 1008 Lamotte, A., 2012. Le Paléolithique inférieur et moyen des bassins-versants de la Somme et de la 1009 Saône: aspects cognitifs de l'environnement pétrographique et considérations techno-1010 typologiques. Habilitation à diriger des recherches dissertation, Université de Lille. 1011 Leader, G.M., Kuman, K., Gibbon, R.J., Granger, D.E., 2018. Early Acheulean organised core knapping 1012 strategies ca. 1.3 Ma at Rietputs 15, Northern Cape Province, South Africa. Quaternary 1013 International 480, 16-28. 1014 Lenoir, M., Turg, A., 1995. Recurrent centripetal debitage (Levallois and Discoidal): Continuity or 1015 discontinuity? In: Dibble, H.L., Bar-Yosef, O. (Eds.), The Definition and Interpretation of Levallois 1016 Technology. Prehistory Press, Madison, pp. 249–256. 1017 Lenzi, F., Nenzioni, G. (Eds.), 1996. Lettere di pietra. I depositi pleistocenici: sedimenti , industrie e 1018 faune del margine appenninico bolognese. Editrice Compositori, Bologna. 1019 Limondin-Lozouet, N., Antoine, P., Bahain, J.-J., Cliquet, D., Coutard, S., Dabkowski, J., Ghaleb, B., 1020 Locht, J.-L., Nicoud, E., Voinchet, P., 2015. North-West European MIS 11 malacological 1021 successions: a framework for the timing of Acheulean settlements. Journal of Quaternary Science 1022 30, 702-712. 1023 Lumley, M.A. de, 1981. Les restes humains d'Orgnac 3. In: De Lumley H. (Ed.). Les premiers habitants 1024 de l'Europe (1 5000 000 - 1000 000 ans), Paris, Catalogue de l'exposition du Laboratoire de 1025 Préhistoire du Musée de l'Homme, Paris, pp.143-145. Lycett, S. J., Eren, M. I., 2013. Levallois economics: an examination of 'waste' production in 1026 1027 experimentally produced Levallois reduction sequences. Journal of Archaeological Science 40, 1028 2384-2392.

1029 Lycett, S.J., von Cramon-Taubadel, N., Eren, M.I., 2016. Levallois: potential implications for learning 1030 and cultural transmission capacities. Lithic Technology 41, 19-38. 1031 McNabb, J., 2001. The shape of things to come. A speculative essay on the role of the Victoria West 1032 phenomenon at Canteen Koppie during the South African Earlier Stone Age. In: Miliken, S., Cook, 1033 J. (Eds.), A Very Remote Period Indeed: Papers on the Palaeolithic Presented to Derek Roe. 1034 Oxbow books, London. pp. 37-46. 1035 Malinsky-Buller, A., 2016a. The Lower-Middle Palaeolithic Transition from the Levantine perspective. 1036 Journal of World Prehistory 29, 1-78. 1037 Malinsky-Buller, A., 2016b. Lost and found: Technological trajectories within Lower / Middle 1038 Paleolithic transition in Western Europe, North of the Pyrenees. Quaternary International 409, 1039 104-148. 1040 Manzi, G., Magri, D., Milli, S., Palombo, M. R., Margari, V., Celiberti, V., Barbieri, M., Barbieri, M., 1041 Melis, R.T., Rubini, M., Ruffo, M., Saracino, B., Tzedakis, P.C., Zarattini, A. Biddittu, I., 2010. The 1042 new chronology of the Ceprano calvarium (Italy). Journal of Human Evolution 59, 580-585. 1043 Masaoudi, H., 1995. Application des méthodes du déséquilibre des familles de l'Uranium 1044 (230th/234U) et de la résonance de spin électronique à la datation des sites d'Orgnac 3, de Payre 1045 et de l'abri des Pêcheurs (Ardèche). Ph.D. Dissertation, Muséum National d'Histoire Naturelle Paris. 1046 1047 Mazza, P.P.A., Martini, F., Sala, B., Magi, M., Perla Colombini, M, Giachi, G., Landucci, F., Lemorini, C., 1048 Modugno, F., Ribechini, E., 2006. A new Palaeolithic discovery: tar-hafted stone tools in a 1049 European Mid-Pleistocene bone-bearing bed. Journal of Archaeological Science 33, 1310-1318. 1050 Meijer, T., Cleveringa, P., 2009. Aminostratigraphy of Middle and Late Pleistocene deposits in The 1051 Netherlands and the southern part of the North Sea Basin. Global and Planetary Change 68, 326-1052 345. 1053 Mellars, P., 1996. Symbolism, language, and the Neanderthal mind. Modelling the early human mind, 1054 University of Cambridge, Cambridge, pp. 15-32.

1055 Meyer, M., Fu, Q., Aximu-Petri, A., Glocke, I., Nickel, B., Arsuaga, J.L., Martínez, I., Gracia, A., 1056 Bermúdez de Castro, J.M., Carbonell, E., Pääbo, S., 2014. A mitochondrial genome sequence of a 1057 hominin from Sima de los Huesos. Nature 505, 403. 1058 Meyer, M., Arsuaga, J.-L., de Filippo, C., Nagel, S., Aximu-Petri, A., Nickel, B., Martínez, I., Gracia, A., 1059 Bermúdez de Castro, J.M., Carbonell, E., Viola, B., Kelso, J., Prüfer, K., Pääbo, S., 2016. Nuclear 1060 DNA sequences from the Middle Pleistocene Sima de los Huesos hominins. Nature 531, 504–507. 1061 Michel, V., Shen, G., Shen, C.-C., Wu, C.-C., Vérat, C., Gallet, S., Moncel, M.-H. Combier, J., Khatib, S., 1062 Manetti, M., 2013. Application of U/Th and 40Ar/39Ar Dating to Orgnac 3, a Late Acheulean and 1063 Early Middle Palaeolithic Site in Ardèche, France, Plos ONE 8(12), e82394. 1064 Moigne, A.-M., 1999. Large mammal assemblages from Lower Palaeolithic sites in France: La Caune 1065 de l'Arago, Terra Amata, Orgnac 3 and Cagny l'Epinette. In : Stiner, M. (Ed.), The role of early 1066 humans in the accumulation of European Lower and Middle Palaeolithic bone assemblages. Mainz 1067 1999, Monographien des Römisch-Germanischen Zentralmuseums 42, pp. 219-235. 1068 Moigne, A.-M., Valensi, P., Auguste, P., García-Solano, J., Tuffreau, A., Lamotte, A., Barroso, C., 2016. 1069 Bone retouchers from Lower Palaeolithic sites: Terra Amata, Orgnac 3, Cagny-l'Epinette and 1070 Cueva del Angel. Quaternary International 409, 195–212. 1071 Moncel, M.-H., 2003. L'exploitation de l'espace et la mobilité des groupes humains au travers des 1072 assemblages lithiques à la fin du Pléistocène moyen et au début du Pléistocène supérieur. La 1073 moyenne vallée du Rhône entre Drôme et Ardèche (France). BAR International Series 1184, 1074 Oxford. 1075 Moncel, M.-H., Moigne, A.-M., Sam, Y., Combier, J., 2011. The emergence of Neanderthal technical 1076 behavior: New evidence from Orgnac 3 (level 1, MIS 8), southeastern France. Current 1077 Anthropology 52, 37-75 Moncel, M.H., Moigne, A.M., Combier, J., 2012. Towards the Middle Palaeolithic in western Europe: 1078 1079 the case of Orgnac 3 (southeastern France). Journal of Human Evolution 63, 653-666.

1080 Moncel, M.H., Despriée, J., Voinchet, P., Tissoux, H., Moreno, D., Bahain, J.J., Courcimault, G., 1081 Falguères, C., 2013. Early evidence of Acheulean settlement in northwestern Europe-La Noira Site, 1082 a 700 000 year-old occupation in the center of France. PLoS One 8, e75529. 1083 Moncel, M. H., Ashton, N., Lamotte, A., Tuffreau, A., Cliquet, D., Despriée, J., 2015. The early 1084 Acheulian of north-western Europe. Journal of Anthropological Archaeology 40, 302-331. 1085 Moncel, M.-H., Arzarello, M., Peretto, C., 2016. The Holsteinian period in Europe (MIS 11-9). 1086 Quaternary International 409, 1-8. 1087 Monnier, G., 2006. An Evaluation of the Lower/Middle Palaeolithic Periodization in Western Europe. 1088 Current Anthropology 47, 709–745. 1089 Mourer-Chauvire, C., 1975. Les Oiseaux du Pléistocène moyen et supérieur de France. Falguères 1090 Documents Laboratoire Faculté des Sciences de Lyon 64, Lyon. 1091 Mourre, V., 2006. Émergence et évolution de la prédétermination au Paléolithique. Normes 1092 Techniques et pratiques sociales. De la simplicité des outillages pré-et protohistoriques. In : 1093 Astruc, L., Bon, F., Léa, V., Milcent, P-Y., Philibert, S. (Eds.), XXVIe Rencontres Internationales 1094 d'Archéologie et d'Histoire d'Antibes. APDCA ed., Antibes, pp. 61-74. 1095 1096 Muttillo, B., Lembo, G., Peretto, C. (Eds.), 2014. L'insediamento a bifacciali di Guado san Nicola. 1097 Monteroduni, Molise, Italia. Annali dell'Universita degli Studi di Ferrara, Ferrara. 1098 Nitychoruk, J., Bińka, K., Ruppert, H., Schneider, J., 2006. Holsteinian Interglacial=Marine Isotope 1099 Stage 11? Quaternary Science Reviews 25, 2678–2681. 1100 Nowell, A., White, M., 2010. Growing up in the Middle Pleistocene. In: Nowell A. and Davidson I. 1101 (Eds.) Stone Tools and the evolution of Human cognition. University Press of Colorado, Boulder, 1102 pp. 67-82. 1103 Ollé, A., Mosquera, M., Rodríguez-Álvarez, X. P., García-Medrano, P., Barsky, D., de Lombera-1104 Hermida, A., Carbonell, E., 2016. The Acheulean from Atapuerca: Three steps forward, one step

back. Quaternary International 411, 316-328.

1105

1106 Orlando, L., Darlu, P., Toussaint, M., Bonjean, D., Otte, M., Hänni, C., 2006. Revisiting Neandertal 1107 diversity with a 100,000 year old mtDNA sequence. Current Biology 16, R400-R402. 1108 Palma di Cesnola, A., 1967. Il Paleolitico della Puglia (giacimenti, periodi, problemi). Memorie del 1109 Museo Civico di Storia Naturale di Verona 15, 1-84. 1110 Palombo, M.R., 2015. To what extent could functional diversity be a useful tool in inferring 1111 ecosystem responses to past climate changes? Quaternary International 413, 15-31. 1112 Pereira, A., Nomade, S., Shao, Q., Bahain, J.-J., Arzarello, M., Douville, E., Falguères, C., Frank, N., 1113 Garcia, T., Lembo, G., Muttillo, B., Scao, V., Peretto, C., 2016. 40Ar/39Ar and ESR/U-series dates for 1114 Guado San Nicola, Middle Pleistocene key site at the Lower/Middle Palaeolithic transition in Italy. 1115 Quaternary Geochronology 36, 67–75. 1116 Pereira, A., Nomade, S., Moncel, M-H., Voinchet, P., Bahain, J-J., Biddittu, I., Falguères, C., Giaccio, B., 1117 Manzi, G., Parenti, F., Scardia, G., Scao, V., Sottili, G., Vietti, A., 2018. Geochronological evidences 1118 of a MIS 11 to MIS 10 age for several pivotal Acheulian sites from the Frosinone province (Latium, 1119 Italy): Archaeological implications. Quaternary Sciences Reviews 187, 112-129. 1120 Peretto, C., Arzarello, M., Bahain, J.-J., Boulbes, N., Dolo, J.-M., Douville, E., Falguères, C., Frank, N., 1121 Garcia, T., Lembo, G., Moigne, A.-M., Muttillo, B., Nomade, S., Pereira, A., Rufo, M.A., Sala, B., 1122 Shao, Q., Hohenstein, U.T., Tessari, U., Turrini, M.C., Vaccaro, C., 2015. The Middle Pleistocene 1123 site of Guado San Nicola (Monteroduni, Central Italy) on the Lower/Middle Palaeolithic transition. 1124 Quaternary International 411, 301-315. 1125 Peris, J. F., González, V. B., Blasco, R., Monteagudo, F. C., Sañudo, P., 2008. El Paleolítico Medio en el 1126 territorio valenciano y la variabilidad tecno-económica de la Cova del Bolomor. Treballs 1127 d'Arqueologia 14, 141-169. 1128 Picin, A., Peresani, M., Falguères, C., Gruppioni, G., Bahain, J.J., 2013. San Bernardino Cave (Italy) and the appearance of Levallois technology in Europe: results of a radiometric and technological 1129

1130

reassessment. PLoS One 8, e76182.

1131 Picin, A., 2017. Technological adaptation and the emergence of Levallois in Central Europe: new 1132 insight from the Markkleeberg and Zwochau open-air sites in Germany. Journal of Quaternary 1133 Science 33, 300-312. 1134 Pope, M., McNabb, J., Gamble, C. (Eds.), 2017. Crossing the Human Threshold: Dynamic 1135 Transformation and Persistent Places during the Middle Pleistocene. Routledge, New York. 1136 Potts, R., Behrensmeyer, A.K., Faith, J.T., Tryon, C.A., Brooks, A. S., Yellen, J.E., Deino, A.L., Kinyanjui, 1137 R., Clark, J.B., Haradon, C.M., Levin, N.E., Meijer, H.J.M., Veatch, E.G., Awen, R.B., Renaut, R.W., 1138 2018. Environmental dynamics during the onset of the Middle Stone Age in eastern Africa. Science 1139 360, 86-90. Premo, L.S., Hublin, J.J., 2008. Culture, population structure, and low genetic diversity in Pleistocene 1140 1141 hominins. Proceedings of the National Academy of Sciences USA 106, 33-37. 1142 Renfrew, C., 1978. The anatomy of innovation. Social Organization and Settlement. In: Green D., 1143 Haselgrove C., and Springs D.. Contributions from Anthropology, Archaeology and Geography. Bar 1144 International Series, Cambridge, pp. 89-117. 1145 Révillon, S., 1995. Technologie du débitage laminaire au Paléolithique moyen en Europe 1146 septentrionale: état de la question. Bulletin de la Société Préhistorique Française 92, 425-441. 1147 Richter, D., Grün, R., Joannes-Boyau, R., Steele, T.E., Amani, F., Rué, M., Fernandes, P., Raynal, J-P., 1148 Geraads, D., Ben-Ncer, A., Hublin, J-J., McPherron, S., 2017. The age of the hominin fossils from 1149 Jebel Irhoud, Morocco, and the origins of the Middle Stone Age. Nature 546, 293-296. 1150 Rigaud, J.P., Aitken, M., Andrieux, C., Beyries, S., Binford, L., 1988. La grotte Vaufrey à Cenac et Saint-1151 Julien (Dordogne)-Paléoenvironnements, chronologie et activités humaines. Mémoires de la 1152 Société Préhistorique Française 19. Paris. 1153 Rightmire, G.P., 2008. Homo in the middle Pleistocene: Hypodigms, variation, and species 1154 recognition. Evolutionary Anthropology 17, 8–21.

- 1155 Rightmire, G.P., 2017. Middle Pleistocene *Homo* Crania from Broken Hill and Petralona: Morphology,
- 1156 Metric Comparisons, and Evolutionary Relationships. In: Marom A., Hovers E. (Eds.). Human
- Paleontology and Prehistory. Springer, Cham, pp. 145-159.
- 1158 Ríos, L., Kivell, T. L., Lalueza-Fox, C., Estalrrich, A., García-Tabernero, A., Huguet, R., Quintino, Y., de la
- 1159 Rasilla, M., Rosas, A., 2019. Skeletal anomalies in the Neandertal family of el Sidrón (Spain) support
- 1160 A Role of Inbreeding in Neandertal extinction. Scientific reports 9, 1697.
- Roe, D.A., 1981. The Lower and Middle Palaeolithic Periods in Britain. Routledge & Kegan Paul, London.
- 1162 Rodriguez-Hidalgo, A., Saladié, P., Ollé, A., Arsuaga, J.L., Bermúdez de Castro, J.M., Carbonell, E.,
- 1163 2017. Human predatory behavior and the social implications of communal hunting based on
- evidence from the TD10. 2 bison bone bed at Gran Dolina (Atapuerca, Spain). Journal of Human
- 1165 Evolution 105, 89-122.
- 1166 Roe, H.M., Coope, G.R., Devoy, R.J.N., Harrison, C.J.O., Penkman, K.E.H., Preece, R.C., Schreve, D.C.,
- 2009. Differentiation of MIS 9 and MIS 11 in the continental record: vegetational, faunal,
- aminostratigraphic and sea-level evidence from coastal sites in Essex, UK. Quaternary Science
- 1169 Reviews 28, 2342–2373.
- 1170 Roebroecks, W., Villa P., 2011. On the earliest evidence for habitual use of fire in Europe.
- 1171 Proceedings of the National Academy of Sciences USA 108, 5209–5214.
- 1172 Roebroeks, W., 1988. A Study of Middle Palaeolithic Riverside Settlements at Maastricht-Belvédère
- 1173 (The Nederlands). Analecta Praehistorica Leidensia 21.
- 1174 Rogers, A.R., Bohlender, R.J., Huff, C.D., 2017. Early history of Neanderthals and Denisovans.
- 1175 Proceedings of the National Academy of Sciences USA 114, 9859-9863.
- 1176 Rohling, E.J., Braun, K., Grant, K., Kucera, M., Roberts, A.P., Siddall, M., Trommer, G., 2010.
- 1177 Comparison between Holocene and Marine Isotope Stage-11 sea-level histories. Earth and
- 1178 Planetary Science Letters 291, 97-105.
- 1179 Rossoni-Notter, E., Notter, O., Simone, S., Simon, P., 2016. Acheulean technical behaviors in Aldène
- cave (Cesseras, Hérault, France). Quaternary International 409, 149–173.

1181 Rots, V., 2013. Insights into early Middle Palaeolithic tool use and hafting in Western Europe. The 1182 functional analysis of level {IIa} of the early Middle Palaeolithic site of Biache-Saint-Vaast (France). 1183 Journal of Archaeological Science 40, 497–506. 1184 Rots, V., Van Peer, P., Vermeersch, P.M., 2011. Aspects of tool production, use, and hafting in 1185 Palaeolithic assemblages from Northeast Africa. Journal of Human Evolution 60, 637–664. 1186 Roure Johnson, C., McBrearty, S., 2010. 500,000 year old blades from the Kapthurin Formation, 1187 Kenya. Journal of Human Evolution 58, 193-200. 1188 Sam, Y., 2009. Etude paléontologique, taphonomique et archéozoologique des grands mammifères du 1189 site Pléistocène moyen d'Orgnac 3 (Ardèche, France). PhD dissertaion University of Perpignan. Sam, Y., Moigne, A.-M., 2011. Rôles des Hommes et des Carnivores dans l'accumulation osseuse des 1190 1191 niveaux profonds d'Orgnac 3 (Ardèche, France). Exemple des niveaux 7-8. In: Brugal, J-P., 1192 Gardeisen, A., Zucker, A. (Eds.), Prédateurs dans tous leurs états, évolution, biodiversité, 1193 interactions, Mythes, Symboles. XXXI rencontres internationales d'archéologie et d'histoire 1194 d'Antibes. Ed. APDCA, Antibes, pp. 351-367 1195 Sampson, C.G. (Ed.), 1978. Paleoecology and Archaeology of an Acheulian Site at Caddington, England. 1196 Southern Methodist University, Dallas. 1197 Sánchez-Romero, L., Benito-Calvo, A., Pérez-González, A., Santonja, M., 2016. Assessment of 1198 accumulation processes at the Middle Pleistocene site of Ambrona (Soria, Spain). Density and 1199 orientation patterns in spatial datasets derived from excavations conducted from the 1960s to the 1200 present. PLoS One 11, e0167595. 1201 Santonja, M., Rubio-Jara, S., Panera, J., Sánchez-Romero, L., Tarriño, A., Pérez-González, A., 2018. 1202 Ambrona revisited: the acheulean lithic industry in the lower stratigraphic complex. Quaternary 1203 International 480, 95-117. 1204 Sharon, G., 2009. Acheulian giant-core technology: a worldwide perspective. Current Anthropology

1205

50, 335-367.

1206 Shimelmitz, R., Kuhn, S.L., 2017. The toolkit in the core: There is more to Levallois production than 1207 predetermination. Quaternary International 464 81-91. 1208 Schreve, D.C., 2001. Mammalian evidence from Middle Pleistocene fluvial sequences for complex 1209 environmental change at the oxygen isotope substage level. Quaternary International 79, 65–74. 1210 Scott, B., 2011. Becoming Neanderthals. The Earlier British Middle Palaeolithic. Oxbow Book, Oxford. 1211 Scott, B., Ashton, N., Penkman, K., Preece, R., White, M., 2010. The position and context of Middle 1212 Palaeolithic industries from the Ebbsfleet Valley, Kent, UK. Journal of Quaternary Science 25, 931-1213 944. 1214 Soressi, M. A., 2016. Neandertals revised. Proceedings of the National Academy of Sciences USA 113, 1215 6372-6379. 1216 Soriano, S., Villa, P., 2017. Early Levallois and the beginning of the Middle Paleolithic in central Italy. 1217 PLoS One 12, 1-28 1218 Stiner, M.C., 2002. Carnivory, coevolution, and the geographic spread of the genus Homo. Journal of 1219 Anthropological Research 10, 1–63. 1220 Stringer, C., 2012. The status of Homo heidelbergensis (Schoetensack 1908). Evolutionary 1221 Anthropology 21, 101–107. 1222 Texier, J.-P., 1995. The Oldowan assemblage from NY18 site at Nyabusosi (Toro-Uganda). Comptes 1223 Rendu de l'Académie des Siences Paris 320, 647-653. 1224 Texier, P-J., Roche, H., 1995. The impact of predetermination on the development of some 1225 Acheulean chaînes opératoires. In: Bermúdez de Castro, J., Arsuaga, J.L., Carbonell, E. (Eds.), 1226 Evolución humana en Europa y los yacimientos de la Sierra de Atapuerca. Junta de Castilla y Leon, 1227 Valladolid, pp. 403-420.

Thieme, H., 1997. Lower Palaeolithic hunting spears from Schöningen, Germany. Nature 358, 807-

1228

1229

810.

- 1230 Tillier, A.-M., 1991. Les plus anciens fossiles humains européens : le cas de la France. In : Bonifay, E.,
- 1231 Vandermeersch, B. (Eds), Les premiers européens. Actes du 114e congrès national des Sociétés
- savantes. CTHS Eds., Paris, pp. 291-298.
- 1233 Tuffreau, A., 1987. Le Paléolithique inférieur et moyen du nord de la France (Nord, Pas-de-Calais,
- 1234 Picardie) dans son cadre stratigraphique. Ph.D. Dissertation, Université de Lille.
- 1235 Tuffreau, A., Lamotte, A., Goval E., 2008. L'Acheuléen de la France septentrionale. L'Anthropologie
- 1236 112, 104-139.
- 1237 Tuffreau, A., Lamotte, A., 2010. Oldest Acheulean Settlements in Northern France. Quaternary
- 1238 International 223–224, 455.
- 1239 Tuffreau, A., Auguste, P., Balescu, S., Bahain, J.-J., 2017. Gentelles-Le Mont de l'Evangile
- 1240 (département de la Somme, France) : un site de plateau occupé de l'Acheuléen au Micoquien.
- 1241 L'Anthropologie 121, 394-427.
- 1242 Tryon, C., 2006. "Early" Middle Stone Age Lithic Technology of the Kapthurin Formation (Kenya).
- 1243 Current Anthropology 47, 367-375.
- 1244 Vaesen, K., Collard, M., 2016. Population size does not explain past changes in cultural complexity.
- 1245 Proceedings of the National Academy of Sciences USA 113, e2241–e2247.
- 1246 Vaquero, M., Romagnoli, F., 2018. Searching for lazy people: the significance of expedient behavior in
- the interpretation of Paleolithic assemblages. Journal of Archaeological Method and Theory 25,
- 1248 334-367.
- 1249 Van Baelen, A., 2017. The Lower to Middle Palaeolithic transition in Northwestern Europe. Evidence
- from Kesselt-Op de Schans. Leuven University Press, Leuven.
- 1251 Villa, P., Boscato, P., Ranaldo, F., Ronchitelli, A., 2009. Stone tools for the hunt: points with impact
- scars from a Middle Paleolithic site in southern Italy. Journal of Archaeological Science 36, 850-
- 1253 859.
- 1254 Villa, P. 2009. Discussion 3: the lower to middle Paleolithic transition. In: Camps, M., Chauhan, P.
- 1255 (Eds), Sourcebook of Paleolithic Transitions . Springer, New York, pp. 265-270

1250	villa, P., Roebroeks, W., 2014. Neandertal demise: an archaeological analysis of the modern numan
1257	superiority complex. PLoS One 9, e96424.
1258	Villa, P., Soriano, S., Grün, R., Marra, F., Nomade, S., Pereira, A., Boschian, G., Pollarolo, L., Fang, F.,
1259	Bahain, JJ., 2016. The Acheulian and Early Middle Paleolithic in Latium (Italy): Stability and
1260	Innovation. PLoS One 11, e0160516.
1261	Walker, M.J., Ortega, J., Parmova, K., Lopez, M.V., Trinkaus, E., 2011. Morphology, body proportion
1262	and postcranial hypertrophy of a female Neandertal from the Sima de las Palomas, southeastern
1263	Spain. Proceedings of the National Academy of Sciences USA 108, 10087–10091
1264	Wilkins, J., Schoville, B.J., Brown, K.S., Chazan, M., 2012. Evidence for early hafted hunting
1265	technology. Science 338, 942-946.
1266	White, M., Ashton, N., 2003. Lower Palaeolithic core technology and the origins of the Levallois
1267	method in north-western Europe. Current Anthropology 44, 598–609.
1268	White, M., Ashton, N., Scott, B., 2011. The emergence, diversity and significance of Mode 3
1269	(Prepared Core) technologies. In: Ashton N., Lewis S.G., Stringer C. (Eds.), The Ancient Human
1270	Occupation of Britain. Quaternary Science 14, 67–91.
1271	White, M., Pettitt, P., 2016. Technology of early Palaeolithic Western Europe: an heuristic
1272	framework. Lithics—The Journal of the Lithic Studies Society (16), 27.
1273	Wiśniewski, A., 2014. The beginnings and diversity of Levallois methods in the early Middle
1274	Palaeolithic of Central Europe. Quaternary International 326–327, 364–380.
1275	Wymer, J.J., 1968. Lower Palaeolithic Archaeology in Britain as Represented by the Thames Valley.
1276	John Baker, London.
1277	Wymer, J.J., 1999. The Lower Palaeolithic Occupation of Britain. Wessex Archaeology and English
1278	Heritage, Salisbury.
1279	
1280	

1282	
1283	Figure captions
1284	
1285	Figure 1. Location of the sites with some early evidence of Levallois core technology (black rounds).
1286	The gray surface indicates the Levallois extension at the end of the MIS 9. The dark gray surface
1287	indicates the extension from the MIS 8.
1288	
1289	Figure 2. Cagny-la-Garenne I. Level I3 (Middle of the alluvial sequence): preferential Levallois cores.
1290	The numbers indicate the order of the removals. Drawings and pictures: A. Lamotte.
1291	
1292	Figure 3. Cagny-La Garenne I – Level I4 (Middle of the alluvial sequence): preferential Levallois cores
1293	The numbers indicate the order of the removals. Drawings and pictures: A. Lamotte.
1294	
1295	Figure 4. 1) Cagny-La-Garenne II – Level J (beginning of the alluvial sequence). 2) Cagny-la-Garenne
1296	– Level CXV (final alluvial sequence). Both preferential Levallois cores. Drawings and pictures: A.
1297	Lamotte.
1298	
1299	Figure 5. Guado San Nicola: n°1-3 recurrent Levallois core on flint. Arrows indicate the direction of
1300	the removals (thick arrows for the "Levallois" removals and thin arrows for the striking platform).
1301	Drawings B. Muttillo, modified.
1302	
1303	Figure 6. Guado San Nicola: 1) Levallois flake on silicified limestone; 2–10) recurrent Levallois flakes
1304	and points on flint. Drawings B. Muttillo, modified.
1305	
1306	Figure 7. Orgnac 3: flakes described as Levallois (level 5b). n° 1 flake with centripetal removals and
1307	convexity scars, curved cross-section; n° 2 flake with convexity scars; n°3 backed flake with

.308	centripetal removals and convexity scars; if 4 flake with centripetal removals, curved cross-section
309	Drawings: M.H.M., modified from Moncel (1999).
310	
311	Figure 8. Orgnac 3: n° 1, 2, level 4b, n° 3 level 5a; n° 1 core with an invasive removal; n° 2 flake
312	described as Levallois; n° 3 core with bipolar invasive removals. Arrows indicate the direction of the
313	removals (thick arrows for the "Levallois" removals and thin arrows for the convexities scars).
314	Drawings: M.H.M., modified from Moncel (1999).
315	
316	Figure 9. Orgnac 3 - Level 4b: n° 1, 3, 4, 5, 6 flakes described as Levallois; n° 2 backed flake. Arrows
.317	indicate the direction of the removals. Drawings: M.H.M., modified from Moncel (1999).
.318	
319	Figure 10. Orgnac 3: Level 4a: n° 1 core with centripetal removals; n° 2, 3, 4 Flakes described as
.320	Levallois. Arrows indicate the direction of the removals (thick arrows for the "Levallois" removals and
321	thin arrows for the striking platform/convexities scars). Drawings: M.H.M., modified from Moncel
322	(1999).
.323	
324	Figure 10. Cave dall'Olio, Levallois cores and blanks: 1) bidirectional core; 2) unidirectional
.325	convergent core; 3) lineal core; 4–6 Levallois blanks. Arrows indicate the direction of the removals
.326	(thick arrows for the "Levallois" removals and thin arrows for the striking platform/convexities scars)
.327	Drawings and pictures: F. Fontana.
.328	
329	Figure 12. British sites: n°1–4 simple prepared flint cores at Purfleet Botany Pit (Essex); n°5 group of
1330	refitting flakes from Frindsbury (Kent). Pictures: N. Ashton.
331	
.332	Figure 13. Local innovations of an Early Levallois core technology over time in Europe.

Figure 14. Cagny-la-Garenne I - level CXB: invasive removal on a biface. Photo A. Lamotte

1335

Figure 15. Comparison of length of Levallois flakes and the other flakes for the levels 4a to 5b at

1337 Orgnac 3.