



Bis[diphenylphosphino]methane and its bridge-substituted analogues as chemically non-innocent ligands for H₂ activation

Nikolay Kireev, Oleg Filippov, Ekaterina Gulyaeva, Elena V Shubina, Laure Vendier, Yves Canac, Jean-Baptiste Sortais, Noël Lugan, Dmitry A. Valyaev

► To cite this version:

Nikolay Kireev, Oleg Filippov, Ekaterina Gulyaeva, Elena V Shubina, Laure Vendier, et al.. Bis[diphenylphosphino]methane and its bridge-substituted analogues as chemically non-innocent ligands for H₂ activation. Chemical Communications, 2020, 56 (14), pp.2139-2142. 10.1039/c9cc07713h . hal-02489712

HAL Id: hal-02489712

<https://hal.science/hal-02489712>

Submitted on 6 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bis[diphenylphosphino]methane and its bridge-substituted analogues as chemically non-innocent ligands for H₂ activation


Received 00th January 20xx,
Accepted 00th January 20xx

Nikolay V. Kireev,^{a,b} Oleg A. Filippov,^{*a} Ekaterina S. Gulyaeva,^a Elena S. Shubina,^{a,c} Laure Vendier,^d Yves Canac,^d Jean-Baptiste Sortais,^{d,e} Noël Lugan^d and Dmitry A. Valyaev^{*d}

DOI: 10.1039/x0xx00000x

Deprotonation of *fac*-[(κ^2P,P -Ph₂PCH(R)PPh₂)Mn(CO)₃Br] (R = H, Me, Ph) produces the corresponding diphosphinomethanide derivatives *fac*-[(κ^3P,C,P -Ph₂PC(R)PPh₂)Mn(CO)₃], which are prone to activate H₂ to form the hydride complexes *fac*-[(κ^2P,P -Ph₂PCH(R)PPh₂)Mn(CO)₃H]. Substitution of the dpmm bridge improves dramatically the reaction efficiency and this was rationalized by DFT calculations.

Since the beginning of 1960's, diphosphinomethanes, including the ubiquitous bis[diphenylphosphino]methane (dpmm), have been extensively used in transition metal chemistry as ligands capable to exhibit both chelating or bridging coordination modes.¹ While several mononuclear dpmm-type complexes have found remarkable applications in homogenous catalysis,^{2,3} the chelating dpmm ligand(s) are generally considered as spectators with the notable exception of the efficient chirality induction brought by P-chirogenic derivatives.⁴ Since the pioneering work of Shaw,⁵ it is well known, however, that the CH₂ bridge of chelating dpmm can be readily deprotonated to generate anionic bis[diphenylphosphino]methanide (dpmm⁻) scaffold (Scheme 1, (a)). The nucleophilic character of the bridging carbon atom in such species has later been exploited for the preparation of various bridge-functionalized dpmm derivatives^{5,6} (Scheme 1, (b)) or bi- or polynuclear compounds⁷ (Scheme 1, (c)). We report herein the first experimental evidence that the deprotonation of dpmm-type ligands can also lead to neutral cyclometallated κ^3P,C,P -diphosphinomethanide



Scheme 1. General overview of the chemistry of transition metal diphosphinomethanide (dpmm⁻) complexes.

complexes relevant for the activation of dihydrogen *via* metal-ligand cooperation^{8,9} (Scheme 1, (d) + (e)).

We have recently reported that the phosphine-*N*-heterocyclic carbene (NHC) ligand in the Mn(I) complex *fac*-[(κ^2P,C -Ph₂PCH₂NHC)Mn(CO)₃Br] can be readily deprotonated at the methylene bridge to form the NHC-phosphinomethanide derivative *fac*-[(κ^3P,C,C -Ph₂PCHNHC)Mn(CO)₃], which is able to activate H₂ *via* a non-classical mode of metal-ligand cooperation implying the NHC-phosphonium ylide *fac*-[(κ^2P,C -Ph₂P=CHNHC)-Mn(CO)₃].¹⁰ In a continuation of this project, we wonder whether the similar reactivity trend could be extended to Mn(I) complexes bearing other chelating R₂PCH₂L ligands and we chose for this study the simple dpmm and some of its readily accessible bridge-substituted derivatives.

Treatment of Mn(I) complexes *fac*-[(κ^2P,P -Ph₂PCH(R)PPh₂)Mn(CO)₃Br] (*fac*-**1^H**, R = H; *fac*-**1^{Me}**, R = Me; *fac*-**1^{Ph}**, R = Ph)¹¹ with KHMDS in toluene at room temperature cleanly affords new products *fac*-**2^R** (Scheme 2), which show in IR similar three ν_{CO} patterns shifted by 10–25 cm⁻¹ to lower frequencies being consistent with the formation of neutral metal species. The deprotonation of the methylene (*fac*-**1^H**) or methine (*fac*-**1^{Me}**,

^a A. N. Nesmeyanov Institute of Organoelement Compounds (INEOS), Russian Academy of Sciences, 28 Vavilov str., GSP-1, B-334, Moscow, 119991, Russia. E-mail: h-bond@ineos.ac.ru


^b Department of Chemistry, M. V. Lomonosov Moscow State University, Leninskie Gory 1/3, 119234 Moscow, Russia.

^c Peoples Friendship University of Russia (RUDN University), 6 Miklukho Maklaya str., Moscow, 117198, Russia.

^d LCC-CNRS, Université de Toulouse, CNRS, 205 route de Narbonne 31077 Toulouse, Cedex 4, France. E-mail: dmitry.valyaev@lcc-toulouse.fr

^e Institut Universitaire de France 1 rue Descartes, F-75231 Paris Cedex 05, France

[†] Electronic Supplementary Information (ESI) available: Complete experimental, characterization and computational data for all Mn(I) complexes. CCDC 1954917-1954918. See DOI: 10.1039/x0xx00000x



Scheme 2. Synthesis of Mn(I) diphosphinomethanide complexes $fac-2^R$ (R = H, Me, Ph) and their reactivity with dihydrogen. (i) 1.1 equiv. KHMDS, toluene, 25°C; (ii) (R = H): 50 atm. H_2 , 50°C, 16h; (iii) (R = Me) 50 atm. H_2 , 25°C, 1 h; (iv) (R = Ph): 1 atm. H_2 , 25°C, 5 min.

$fac-1^{Ph}$) bridge within the chelating ligands was unambiguously confirmed by 1H and ^{13}C NMR spectroscopy showing an aliphatic CH fragment as a singlet at δ_H 1.65 ppm, and a strongly shielded triplet at δ_C -10.1 ppm ($^1J_{PC}$ = 5.3 Hz) for $fac-2^H$, and characteristic quaternary carbon atoms at δ_C 0.2 ppm, and δ_C 12.8 ppm ($^1J_{PC}$ = 2.4 Hz) for $fac-2^{Me}$ and $fac-2^{Ph}$, respectively. The values of $^1J_{PC}$ constants within the PCP moiety in complexes $fac-2^R$ are anomalously small compared to known Mn(I) κ^2P,P -dppm $^-$ derivatives^{6,7,12} suggesting the formation of strained P-containing three-membered cycles.¹³ Crystallization of the most stable complex $fac-2^{Ph}$ at -20°C (isolated in 60% yield) allowed to establish its structure by X-ray diffraction (Fig. 1).

Diphosphinomethanide ligand in $fac-2^{Ph}$ is coordinated to manganese in a tridentate facial κ^3P,C,P coordination mode with three carbonyl ligands completing the metal coordination sphere in a distorted octahedral environment. The coordination mode of the phosphorus-containing fragment here is strikingly different from the κ^2P,P planar structures typically observed in mononuclear diphosphinomethanide transition metal complexes¹² and has been previously evidenced uniquely in a series of complexes of lanthanides.¹⁴ The PCP unit in $fac-2^{Ph}$ is symmetrically bound to Mn and both Mn-P and P-C bond distances are equal within the experimental error (Fig. 1). While Mn-P bonds in $fac-2^{Ph}$ are significantly shorter than those observed in the most structurally similar diphosphinomethanide complex $[(\kappa^2P,P\text{-dppm}^R)\text{Mn}(\text{CO})_4]^{12d}$ (R = 2-(4-methyl)quinolyl, Mn-P: 2.3398(14), 2.3390(15) Å; P-C: 1.738(5), 1.735(5) Å) P-C bonds within PCP moiety are slightly




Fig. 1. A perspective view of complex $fac-2^{Ph}$ (30% probability ellipsoids, phenyl groups are represented as wireframe). Selected bond lengths (Å) and angles (°): Mn1-P1 2.2255(9), Mn1-P2 2.2222(8), Mn1-C4 2.214(3), P1-C4 1.775(3), P2-C4 1.772(3), P1-C4-P2 103.66(15), P1-Mn1-P2 77.64(3)

longer. Finally, the Mn-C4 bond length is in range of typical Mn(I)-C σ -bonds (CCDC average distance of 2.20 Å for Mn(I) alkyl complexes). Actually, the deprotonation of the methine bridge in $fac-1^{Ph}$ induces an intramolecular nucleophilic substitution of the bromide ligand by the resulting carbanion leading to the observed original structure $fac-2^{Ph}$ exhibiting 5-electron donor L_2X -type PCP ligand.¹⁵

A first assessment of the reactivity of complexes $fac-2^R$ toward dihydrogen was carried out under 1 atm. of H_2 at 25°C. Overnight reaction of complex $fac-2^H$ gave an intricate mixture of products (Fig. S15-S16), among which the H_2 activation product $fac-[(\kappa^2P,P\text{-dppm})\text{Mn}(\text{CO})_3\text{H}]$ ($fac-3^H$, Scheme 2), characterized, in particular, by a hydride signal at δ_H -5.10 ppm (td, $^2J_{PH}$ = 43.9 Hz, $^4J_{HH}$ = 5.8 Hz), was detected in trace amount. The reactions of $fac-2^{Me}$ and $fac-2^{Ph}$ were found to be much faster and selective: hydride complex $fac-3^{Me}$ was observed as a major product under atmospheric H_2 pressure after one hour at 25°C (Fig. S26-S27) and a selective formation of $fac-3^{Ph}$ took place after only few minutes (Scheme 2). The reactions of $fac-2^H$ and $fac-2^{Me}$ under H_2 pressure (50 atm.) strongly improve the yield of the corresponding hydrides $fac-3^H$ and $fac-3^{Me}$ isolated in 50 and 92% yield, respectively (Scheme 2). According to NMR data, both complexes $fac-3^{Me}$ and $fac-3^{Ph}$ exist in solution as a mixture of isomers in a similar ca. 5:1 ratio.¹⁶ 1D NOESY experiments (Fig. S34-S35, S46-47) clearly revealed that in both cases the major species correspond to the $fac,anti-3^R$ isomer having a *trans* arrangement of the hydride and the proton of the CH(R) bridge. The structure of $fac,anti-3^{Ph}$ was further confirmed by a single-crystal X-ray diffraction study. (Fig. 2).

The mechanism of the H_2 activation by complexes $fac-2^R$ leading to the hydride complexes $fac-3^R$ has been investigated by DFT calculations at the BP86/def2-TZVP level (see the ESI† for details). The first step of the reaction between $fac-2^R$ and H_2 involves an heterolytic cleavage of the Mn-C bond *via* the transition state $TS1^R$ accompanied by the flipping of dppm $^-$ ligand from the κ^3P,C,P to a κ^2P,P mode and H_2 coordination to afford the η^2-H_2 intermediates $fac-4^R$ (Scheme 3).¹⁷ The latter undergo a concerted H-H bond activation process *via* $TS2^R$ to form the hydride complexes $fac,syn-3^R$. Interestingly, the activation barriers for these two steps show opposite trends depending on the substituent R on the bridge. While the





Fig. 2. A perspective view of complex $fac,anti-3^{Ph}$ (30% probability ellipsoids, phenyl groups are represented as wireframe). Selected bond lengths (Å) and angles (°): Mn1-P1 2.2549(4), Mn1-P2 2.2830(3), P1-C4 1.8741(12), P2-C4 1.8878(12), Mn1-H1 1.53(2), P1-C4-P2 91.44(5), P1-Mn1-P2 72.808(12).




Scheme 3. DFT mechanism of cooperative H_2 activation with complexes fac-2^{R} (BP86/def2-TZVP, toluene SMD model, ΔG^{\ddagger} are given in kcal.mol $^{-1}$ and referred to the reactants ($\text{fac-2}^{\text{R}} + \text{H}_2$), for the ΔH^{\ddagger} profile see Scheme S1 in the ESI).

calculated values for TS1^{R} significantly decrease in the $\text{H} > \text{Me} > \text{Ph}$ order (Scheme 3), the intramolecular proton transfer in fac-4^{R} is more difficult for the phenyl-substituted complex fac-4^{Ph} (energy barriers of TS2^{R} : 12.2, 13.3, and 16.7 kcal.mol $^{-1}$ for fac-4^{H} , fac-4^{Me} , and fac-4^{Ph} , respectively). The both Gibbs energy span values, namely the differences between the starting complexes fac-2^{R} and the highest transition states TS2^{R} , and the barriers for rate limiting step of H_2 addition (TS1^{R}) are fully consistent with experimentally observed reactivity trend $\text{fac-2}^{\text{Ph}} > \text{fac-2}^{\text{Me}} \gg \text{fac-2}^{\text{H}}$. A closer inspection of the electronic structure of $\text{dppm}^{\text{R-}}$ complexes revealed stronger Mn–C bonding in fac-2^{H} with Wiberg bond index (WBI) 18 0.307 compared to its bridge-substituted analogues fac-2^{Me} and fac-2^{Ph} (WBI 0.282 and 0.292, respectively), which could indeed contribute to higher TS1^{H} energy value. The potential energy profile of fac-2^{R} upon increasing the dihedral angle Mn–P–C–P to 180° (Figure S54) revealed the enhancement of the ring opening energy in $\text{fac-2}^{\text{Ph}} < \text{fac-2}^{\text{Me}} < \text{fac-2}^{\text{H}}$ row, thus roughly correlating with the corresponding TS1^{R} energies. This may suggest that in addition to electronic effects, dppm bridge substitution facilitates H_2 coordination step by increasing the strain of the bicyclic $[\kappa^3\text{P,C,P}]\text{Mn}$ system in fac-2^{R} .

While the present mechanism account for the formation $\text{fac,syn-3}^{\text{R}}$ isomers ($\text{R} \neq \text{H}$), experimentally, the fac,anti isomers

are actually found to be predominant in solution. According to the DFT calculations the $\text{fac,anti-3}^{\text{Me}}$ and $\text{fac,anti-3}^{\text{Ph}}$ hydride complexes are indeed thermodynamically more stable by 3.2, and 2.5 kcal.mol $^{-1}$, respectively, than their fac,syn analogues. Considering the reaction conditions, such an isomerization could occur through a base-catalyzed epimerization of the bridging chiral carbon. Still, being aware of thermal fac-to-mer isomerization known for complexes $\text{fac-}[(\text{PR}_3)_2\text{Mn}(\text{CO})_3\text{X}]$, 19 such a possibility in the present case was further investigated by DFT calculations (Scheme 4). As a matter of fact, complexes $\text{fac,syn-3}^{\text{R}}$ can be readily transformed by a rotation of the entire $[\text{Mn}(\text{CO})_3\text{H}]$ moiety to the meridional isomers mer-3^{R} , which then can undergo similar rotation to afford $\text{fac,anti-3}^{\text{R}}$ with activation barriers readily accessible at room temperature (see the ESI † for animated gif files illustrating these steps).



Scheme 4. DFT mechanism of the isomerization of hydride complexes $\text{fac,syn-3}^{\text{R}}$ into $\text{fac,anti-3}^{\text{R}}$ (BP86/def2-TZVP, toluene SMD model, ΔG^{\ddagger} are given in kcal.mol $^{-1}$ (green for $\text{R} = \text{Me}$, black for $\text{R} = \text{Ph}$) and referred to $\text{fac,syn-3}^{\text{R}}$). See the ESI † for geometries of transition states and animated gif files for each step of the process.

In order to validate experimentally some mechanistic clues obtained by DFT calculations, we have carried out low temperature NMR monitoring of the reaction between fac-2^{Ph} and H_2 (2 atm.) in toluene- d_8 (Fig. 3). Unfortunately, we were unable to detect $\eta^2\text{-H}_2$ intermediate fac-4^{Ph} in -80 – -40 °C temperature range. Around -20 °C the formation of $\text{fac,syn-3}^{\text{Ph}}$ as a sole hydride product was evidenced in ^{31}P NMR spectrum at δ_{P} 58.6 ppm. Upon further warming a consumption of starting fac-2^{Ph} (δ_{P} 17.4 ppm) as well as appearance and gradual increase of the signal at δ_{P} 53.8 ppm belonging to $\text{fac,anti-3}^{\text{Ph}}$ was observed, thus confirming the viability of proposed reaction mechanism.




Fig. 3. $^{31}\text{P}\{^1\text{H}\}$ NMR monitoring of low temperature reaction of complex fac-2^{Ph} with dihydrogen (toluene- d_8 , 2 atm. of H_2 , 242.9 MHz, the time between each temperature point is ca. 25–30 min).

In conclusion, this work unveils the possibility for the simple diphosphinomethane scaffold to act upon deprotonation as a non-innocent ligand for cooperative dihydrogen activation. Additionally, our work provides one more illustrative example how a subtle ligand modification can improve the desired reactivity of transition metal complex. We are currently working on the application of such Mn(I) systems in homogeneous catalysis and expansion of this approach to the cooperative activation of other inert chemical bonds.

We thank the joint CNRS/RFBR grant no. 19-53-15014 (PRC2330) for financial support. The Ministry of Science and Higher Education of the Russian Federation, the RUDN University program 5-100, and the IUF (France) are acknowledged for a partial support of this research. Computational studies were performed using HPC resources from CALMIP (Grant no. P18038). N.V.K. is grateful to French Embassy in Moscow for a PhD international mobility fellowship (Ostrogradski program).

Conflicts of interest

There are no conflicts to declare.

Notes and references

- (a) R. J. Puddephatt, *Chem. Soc. Rev.*, 1983, **12**, 99; (b) B. Chaudret, B. Delavaux and R. Poilblanc, *Coord. Chem. Rev.*, 1988, **86**, 191.
- S. M. Mansell, *Dalton Trans.*, 2017, **46**, 15157.
- Selected recent examples: (a) S. M. Hansen, M. A. O. Volland, F. Rominger, F. Eisenträger and P. Hofmann, *Angew. Chem., Int. Ed.*, 1999, **38**, 1273; (b) S. J. Dossett, A. Gillon, A. G. Orpen, J. S. Fleming, P. G. Pringle, D. F. Wass and M. D. Jones, *Chem. Commun.*, 2001, 699; (c) T. Suzuki, M. Tokunaga and Y. Wakatsuki, *Org. Lett.*, 2001, **3**, 735; (d) G. R. M. Dowson, M. F. Haddow, J. Lee, R. L. Wingard and D. F. Wass, *Angew. Chem., Int. Ed.*, 2013, **52**, 9005; (e) M. Castaing, S. L. Wason, B. Estepa, J. F. Hooper and M. C. Willis, *Angew. Chem., Int. Ed.*, 2013, **52**, 13280; (f) A. Prades, M. Fernández, S. D. Pike, M. C. Willis and A. S. Weller, *Angew. Chem., Int. Ed.*, 2015, **54**, 8520; (g) M. Gao and M. C. Willis, *Org. Lett.*, 2017, **19**, 2734.
- (a) I. D. Gridnev, Y. Yamanoi, N. Higashi, H. Tsuruta, M. Yasutake and T. Imamoto, *Adv. Synth. Catal.*, 2001, **343**, 118; (b) G. Hoge, H.-P. Wu, W. S. Kissel, D. A. Pflum, D. J. Greene and J. Bao, *J. Am. Chem. Soc.*, 2004, **126**, 5966; (c) K. Huang, X. Zhang, T. J. Emge, G. Hou, B. Cao and X. Zhang, *Chem. Commun.*, 2010, **46**, 8555; (d) W. Tang, A. G. Capacci, A. White, S. Ma, S. Rodriguez, B. Qu, J. Savoie, N. D. Patel, X. Wei, N. Haddad, N. Grinberg, N. K. Yee, D. Krishnamurthy and C. H. Senanayake, *Org. Lett.*, 2010, **12**, 1104.
- S. Al-Jibori and B. L. Shaw, *J. Chem. Soc., Chem. Commun.*, 1982, 286.
- (a) J. Ruiz, V. Riera, M. Vivanco, M. Lanfranchi and A. Tiripicchio, *Organometallics*, 1998, **17**, 3835; (b) J. Ruiz, V. Riera, M. Vivanco, S. García-Granda and M. R. Díaz, *Organometallics*, 1998, **17**, 4562; (c) J. Ruiz, M. Ceroni, O. V. Quinzani, V. Riera and O. E. Piro, *Angew. Chem., Int. Ed.*, 2001, **40**, 220; (d) J. Ruiz, R. Quesada, V. Riera, E. Castellano and O. Piro, *Organometallics*, 2004, **23**, 175; (e) J. Ruiz, M. Ceroni, M. Vivanco, M. P. Gonzalo, S. García-Granda and F. van der Maelen, *Chem. Commun.*, 2005, 4860; (f) J. Ruiz, M. J. Anton, M. Vivanco, M. E. G. Mosquera and R. Quesada, *Inorg. Chem.*, 2008, **47**, 5540.
- (a) V. Riera, J. Ruiz, X. Solans and E. Tauler, *J. Chem. Soc., Dalton Trans.*, 1990, 1607; (b) J. Ruiz, V. Riera, M. Vivanco, S. García-Granda and A. García-Fernández, *Organometallics*, 1992, **11**, 4077; (c) L. R. Falvello, J. Forníes, R. Navarro, A. Rueda and E. P. Urriolabeitia, *Organometallics*, 1996, **15**, 1309; (d) J. Ruiz, M. E. G. Mosquera, V. Riera, M. Vivanco, C. Bois, *Organometallics*, 1997, **16**, 3388; (e) M. E. G. Mosquera, J. Ruiz, V. Riera, S. García-Granda and M. A. Salvadó, *Organometallics*, 2000, **19**, 5533.
- (a) J. R. Khusnutdinova and D. Milstein, *Angew. Chem., Int. Ed.*, 2015, **54**, 12236; (b) L. Alig, M. Fritz and S. Schneider, *Chem. Rev.*, 2019, **119**, 2681; (c) T. Higashi, S. Kusumoto and K. Nozaki, *Chem. Rev.*, 2019, **119**, 10393.
- For specific examples of metal-ligand cooperation involving the methandiide/methanide forms of oxidized dppm congeners, see: a) R. P. Kamalesh Babu, R. McDonald and R. G. Cavell, *Organometallics*, 2000, **19**, 3462; b) T. Cantat, M. Demange, N. Mézailles, L. Ricard, Y. Jean and P. Le Floch, *Organometallics*, 2005, **24**, 4838; c) H. Heuclin, X. F. Le Goff and N. Mézailles, *Chem. Eur. J.*, 2012, **18**, 16136.
- R. Buhaibeh, O. A. Filippov, A. Bruneau-Voisine, J. Willot, C. Duhayon, D. A. Valyaev, N. Lugan, Y. Canac and J.-B. Sortais, *Angew. Chem., Int. Ed.*, 2019, **58**, 6727.
- Complexes *fac*-**1^R** were prepared in ca. 90% yield by thermal reaction of Mn(CO)₅Br with dppm^{11a} or its readily available bridge-substituted analogues^{11b-c}. (a) R. Colton and M. J. McCormick, *Aust. J. Chem.*, 1976, **29**, 1657; (b) C.-L. Lee, Y.-P. Yang, S. J. Rettig, B. R. James, D. A. Nelson and M. A. Lilga, *Organometallics*, 1986, **5**, 2220; (c) S. A. Al-Jibori, *Transition Met. Chem.*, 1995, **20**, 120.
- (a) J. Ruiz, R. Arauz, V. Riera, M. Vivanco, S. García-Granda and A. Menéndez-Velázquez, *Organometallics*, 1994, **13**, 4162; (b) J. Ruiz, V. Riera, M. Vivanco, M. Lanfranchi and A. Tiripicchio, *Organometallics*, 1996, **15**, 1082; (c) J. Ruiz, M. Ceroni, O. V. Quinzani, V. Riera, M. Vivanco, S. García-Granda, F. Van der Maelen, M. Lanfranchi and A. Tiripicchio, *Chem. Eur. J.*, 2001, **7**, 4422; (d) M. E. G. Mosquera, J. Ruiz, G. García and F. Marquinez, *Chem. Eur. J.*, 2006, **12**, 7706.
- F. Mathey and M. Regitz in *Phosphorus-Carbon Heterocyclic Chemistry* (Ed. F. Mathey), Elsevier, 2001, pp. 17-55.
- (a) H. H. Karsch, A. Appelt and G. Müller, *Angew. Chem., Int. Ed.*, 1986, **25**, 823; (b) S. Hao, J.-I. Song, H. Aghabozorg and S. Gambarotta, *J. Chem. Soc., Chem. Commun.*, 1994, 157; (c) H. Karsch, G. Feruzin and P. Bissinger, *J. Chem. Soc., Chem. Commun.*, 1994, 505; (d) S. T. Liddle and K. Izod, *J. Organomet. Chem.*, 2006, **691**, 2599.
- For relevant examples of formation of L₂X-type μ -1 κ^2P,C ;2 κ^1P five-electron donor diphosphinomethanide ligands in bimetallic complexes see: (a) G. M. Dawkins, M. Green, J. C. Jeffery and F. G. A. Stone, *J. Chem. Soc., Chem. Commun.*, 1980, 1120; (b) *ibid.*, *J. Chem. Soc. Dalton Trans.* 1983, 499; (c) D. A. Vicic, T. J. Anderson, J. A. Cowan and A. J. Schultz, *J. Am. Chem. Soc.*, 2004, **126**, 8132; (d) K. D. Wells, R. McDonald, M. J. Ferguson and M. Cowie, *Inorg. Chem.*, 2011, **50**, 3523.
- Similar situation was observed in Mn(I) complex bearing bridge-substituted κ^2P,P -dppm ligand: R. Jana, S. Chakraborty, O. Blacque and H. Berke, *Eur. J. Inorg. Chem.* 2013, 4574.
- For similar calculated η^2 -H₂ Mn(I) intermediates see ref. 10 and 17a-c. (a) C. Liu, R. van Putten, P. O. Kulyaev, G. A. Filonenko and E. A. Pidko, *J. Catal.*, 2018, **363**, 136; (b) R. van Putten, E. A. Uslamin, M. Garbe, C. Liu, A. Gonzalez-de-Castro, M. Lutz, K. Junge, E. J. M. Hensen, M. Beller, L. Lefort and E. A. Pidko, *Angew. Chem. Int. Ed.*, 2017, **56**, 7531; (c) E. B. Hulley, M. L. Helm and R. M. Bullock, *Chem. Sci.* **2014**, **5**, 4729.
- K. B. Wiberg, *Tetrahedron*, 1968, **24**, 1083.
- (a) R. J. Angelici, F. Basolo and A. J. Poë, *J. Am. Chem. Soc.*, 1963, **85**, 2215; (b) D. J. A. De Waal, R. H. Reimann and E. Singleton, *J. Organomet. Chem.*, 1975, **34**, 339.