



**HAL**  
open science

## Assignment of 5-Hydroxytryptamine Receptor (HTR4) to human chromosome 5 bands q31→q33 by in situ hybridization

S. Claeyen, Patrick Faye, Michèle Sebben, Stéphanie Barrère-Lemaire, Joel Bockaert, Aline Dumuis, Sylvie Taviaux

### ► To cite this version:

S. Claeyen, Patrick Faye, Michèle Sebben, Stéphanie Barrère-Lemaire, Joel Bockaert, et al.. Assignment of 5-Hydroxytryptamine Receptor (HTR4) to human chromosome 5 bands q31→q33 by in situ hybridization. *Cytogenetic and Genome Research*, 1997, 78 (2), pp.133-134. 10.1159/000134646 . hal-02491196

**HAL Id: hal-02491196**

**<https://hal.science/hal-02491196>**

Submitted on 25 Feb 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Assignment<sup>#</sup> of 5-hydroxytryptamine receptor (HTR4) to human chromosome 5 bands q31-->q33 by in situ hybridization.**

Sylvie Claeysen<sup>1</sup>, Patrick Faye<sup>1</sup>, Michèle Sebben<sup>1</sup>, Stéphanie Lemaire<sup>2</sup>, Joël Bockaert<sup>1</sup>, Aline Dumuis<sup>1</sup> and Sylvie Taviaux<sup>2,\*</sup>

<sup>1</sup>CNRS UPR 9023, Centre CNRS-INSERM de Pharmacologie-Endocrinologie (CCIPE), 141 rue de la Cardonille, 34094 Montpellier cedex 05, France

<sup>2</sup>CRBM-CNRS, 1919 route de Mende, BP 5051, 34033 Montpellier Cedex 1, France

\* Author for correspondence

Phone: (33) 04 67-61-33-45 Fax: (33) 04 67-52-15-59

Email: [taviaux@vega.crbm.cnrs-mop.fr](mailto:taviaux@vega.crbm.cnrs-mop.fr)

<sup>#</sup>To our knowledge this is the first time this gene has been mapped.

This work was supported by grants from the Centre National de la Recherche Scientifique, Laboratoires Fournier-Debat (Daix, France) and the Fondation pour la Recherche Médicale.

## **Rationale and significance**

5-Hydroxytryptamine 4 receptors (5-HT<sub>4</sub>R) were first characterized in 1988 (Dumuis *et al.*, 1988) in mouse colliculi neurons. Since then, it has been shown that 5-HT<sub>4</sub>R mediates widespread effects in central and peripheral systems (Eglen *et al.*, 1995). These observations have stimulated the search and discovery of a very large series of selective and potent agonists and antagonists as well as the cloning of the receptor in rat and mouse brain (Eglen *et al.*, 1996; Gerald *et al.*, 1995; Claeysen *et al.*, 1996). In human atria the concentration of 5-HT<sub>4</sub>R and  $\alpha_1$ - and  $\alpha_2$ -adrenoceptors vary, depending on whether the patients have been treated with  $\alpha$ -adrenergic-blocking agents or not (Kaumann *et al.*, 1994; Ouadid *et al.*, 1992). These observations suggest that a "cross-talk" may exist between the regulation of 5-HT<sub>4</sub>R and  $\alpha$ -adrenoceptors in human myocardium. To answer these questions, we cloned human 5-HT<sub>4</sub>R (gene symbol: HTR4; Claeysen *et al.*, 1997) from heart and mapped the chromosomal location by FISH.

## **Materials and methods**

Human metaphase cells were prepared from phytohemagglutinin-stimulated lymphocytes. Probe was labeled with biotinyl-16-dUTP and FISH was performed as previously described (Taviaux and Demaille, 1993).

Probe name(s): h5-HT<sub>4s</sub>

Probe type: cDNA

Insert size: 1.2kb

Proof of authenticity: sequencing

Gene reference: GDB:6381308; Claeysen *et al.* (1997)

## **Results**

Mapping data

Location: 5q31→q33

Number of cells examined: 52

Number of cells with specific signal: 1 (19), 2 (8), 3 (6), 4 (1) chromatids per cell

Most precise assignment: 5q32

Location of background signals (sites with >2 signals): none

## References


- Claeyssen S, Sebben M, Journot L, Bockaert J, Dumuis A: Cloning expression and pharmacology of the mouse 5-HT<sub>4L</sub> receptor. *FEBS Letters* 398:19-25 (1996).
- Claeyssen S, Faye P, Sebben M, Lemaire S, Bockaert J, Dumuis A: Cloning and expression of human 5-HT<sub>4S</sub> receptors: effect of receptor density on their coupling to adenylyl cyclase. *Neuroreport* 8 issue 15 (1997)
- Dumuis A, Bouhelal R, Sebben M, Cory R, Bockaert J: A non classical 5-hydroxytryptamine receptor positively coupled with adenylate cyclase in the central nervous system. *Mol Pharmacol* 34:880-887 (1988).
- Eglen R, Wong EHF, Dumuis A, Bockaert J: Central 5-HT<sub>4</sub> receptors. *Trend Pharmacol* 16:391-398 (1995).
- Eglen RM, Hegde S: 5-Hydroxytryptamine (5-HT<sub>4</sub>) receptors: physiology, pharmacology and therapeutic potential. *Exp Opin Invest Drugs* 5:373-388 (1996).
- Gerald C, Adham N, Kao HT, Olsen MA, Laz TM, Schechter LE, Bard JA, Vaysse PJ, Hartig PR, Branchek TA, Weinshank RL: The 5-HT<sub>4</sub> receptor: molecular cloning and pharmacological characterization of two splice variants. *EMBO J* 14:2806-2815 (1995).
- Kaumann AJ, Sanders L: 5-Hydroxytryptamine causes rate-dependent arrhythmias through 5-HT<sub>4</sub> receptors in human atrium-facilitation by chronic beta-adrenoceptor blockade. *Naunyn-Schmiedeberg's Arch Pharmacol* 349:331-337 (1994).
- Ouadid H, Seguin J, Dumuis A, Bockaert J, Nargeot J: Serotonin increases Ca<sup>++</sup> current in human atrial myocytes via the newly described 5-hydroxytryptamine<sub>4</sub> receptors. *Mol Pharmacol* 41:346-351 (1992).
- Taviaux AS, Demaille JG: Localization of human cell cycle regulatory genes CDC25C to 5q31 and WEE1 TO 11p15.3→p15.1 by fluorescence in situ hybridization. *Genomics* . 15:194-196 (1993).

## Figure legend


### Figure 1

**A)** Hybridization of a human cDNA 5-HT<sub>4</sub> receptor probe to R-band human metaphases. Fluorescent spots are visualized at 5q31→q33. **B)** Ideogram of chromosome 5 homologue showing the fluorescent spot distribution to the 5q31→q33 band for 30 metaphases.

**A**



**B**



**Figure 1**