

## Unorthodox features in two venerid bivalves with doubly uniparental inheritance of mitochondria

Charlotte Capt, Karim Bouvet, Davide Guerra, Brent Robicheau, Donald

Stewart, Eric Pante, Sophie Breton

### ► To cite this version:

Charlotte Capt, Karim Bouvet, Davide Guerra, Brent Robicheau, Donald Stewart, et al.. Unorthodox features in two venerid bivalves with doubly uniparental inheritance of mitochondria. Scientific Reports, 2020, 10 (1), 10.1038/s41598-020-57975-y. hal-02499562

## HAL Id: hal-02499562 https://hal.science/hal-02499562

Submitted on 8 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## SCIENTIFIC REPORTS

natureresearch

### **OPEN**

# Unorthodox features in two venerid bivalves with doubly uniparental inheritance of mitochondria

Charlotte Capt<sup>1\*</sup>, Karim Bouvet<sup>1</sup>, Davide Guerra<sup>1</sup>, Brent M. Robicheau<sup>2</sup>, Donald T. Stewart<sup>3</sup>, Eric Pante<sup>4,5</sup> & Sophie Breton<sup>1,5\*</sup>

In animals, strictly maternal inheritance (SMI) of mitochondria is the rule, but one exception (doubly uniparental inheritance or DUI), marked by the transmission of sex-specific mitogenomes, has been reported in bivalves. Associated with DUI is a frequent modification of the mitochondrial *cox2* gene, as well as additional sex-specific mitochondrial genes not involved in oxidative phosphorylation. With the exception of freshwater mussels (for 3 families of the order Unionida), these DUI-associated features have only been shown in few species [within Mytilidae (order Mytilida) and Veneridae (order Venerida)] because of the few complete sex-specific mitogenomes published for these orders. Here, we present the complete sex-specific mtDNAs of two recently-discovered DUI species in two families of the order Venerida, *Scrobicularia plana* (Semelidae) and *Limecola balthica* (Tellinidae). These species display the largest differences in genome size between sex-specific mitotypes in DUI species (>10 kb), as well as the highest mtDNA divergences (sometimes reaching >50%). An important in-frame insertion (>3.5 kb) in the male *cox2* gene is partly responsible for the differences in genome size. The *S. plana cox2* gene is the largest reported so far in the Kingdom Animalia. The mitogenomes may be carrying sex-specific genes, indicating that general mitochondrial features are shared among DUI species.

Animal mitochondrial DNA (mtDNA) is typically depicted as a strictly maternally inherited (SMI) circular DNA molecule that is relatively small (~16 kb) and genomically streamlined with almost invariant gene content (13 protein-coding genes and 24 structural RNAs)<sup>1,2</sup>. However, important deviations do occur in the mtD-NAs of bivalve molluscs, which not only display dramatic variation in size (<14.7 kb to >67 kb)<sup>3,4</sup> and gene arrangement<sup>5</sup>, but also the presence of additional protein-coding genes not associated with oxidative phosphorylation<sup>6–9</sup>. An even more extreme departure from the norm in bivalve mitochondrial genomes is their mode of doubly uniparental inheritance (DUI) — both egg and sperm mitochondria are transmitted from generation to generation in several bivalve species, but only male offspring retain paternally-transmitted mitochondria (with male or M mtDNA) in their gametes<sup>10–12</sup>. Adult females of DUI-exhibiting species usually possess only the female-transmitted mtDNA (F mtDNA) in their soma and gametes whereas males possess F mtDNA in their soma and M mtDNAs in their gametes<sup>10–13</sup>. The DNA divergence between F and M mtDNAs usually vary from about 8% to 40% depending on the species<sup>12,14</sup>. Genetic analyses suggested that both F and M mtDNAs in DUI bivalves evolve at a faster rate than typical metazoan mtDNA, and that M mtDNA evolves faster than F mtDNA<sup>15–17</sup>. One factor explaining this observation may be that the M genome is subject to weaker selective pressures than

the F genome due to an unequal "division of labor" in the DUI system<sup>16</sup>. Typical animal mtDNA functions in gonads and somatic tissues of both sexes whereas under DUI, F mtDNA functions in female gonads and the soma of both sexes, while M mtDNAs functions primarily in spermatozoa of male gonads and only partially in the male soma<sup>13,16,18</sup>. As opposed to SMI that promotes homoplasmy, a state in which all mtDNA copies are typically genetically identical in each cell, thus preventing potentially harmful genomic conflicts, DUI is a naturally heter-oplasmic system in which two highly divergent mitochondrial lineages coexist in the same nuclear background, enabling the analysis of the consequences of tissue heteroplamy<sup>13</sup>.

<sup>1</sup>Department of Biological Sciences, Université de Montréal, Montréal, QC, Canada. <sup>2</sup>Department of Biology, Dalhousie University, Halifax, NS, Canada. <sup>3</sup>Department of Biology, Acadia University, Wolfville, NS, B4P 2R6, Canada. <sup>4</sup>Littoral, Environnement et Sociétés (LIENSs), UMR 7266 CNRS–La Rochelle Université, 2 rue Olympe de Gouges, 17000, La Rochelle, France. <sup>5</sup>These authors jointly supervised this work: Sophie Breton and Eric Pante. \*email: capt.charlotte@gmail.com; s.breton@umontreal.ca

In addition to a different mode of mitochondrial transmission and evolution rate of mtDNA, two other remarkable differences have been reported between the F and M mtDNAs in DUI bivalves. First, the COX2 protein encoded by M mtDNA (Mcox2 gene) is longer than the FCOX2 protein, which is about the same size as other animal COX2 proteins, although this pattern is not shared by all DUI species for which sex-specific mtDNAs have been completely sequenced (reviewed in Bettinazzi et al.<sup>19</sup>). For example, male freshwater mussels (order Unionida) have an approximately 550 bp 3'-coding extension to the cox2 gene (Mcox2e), that is absent from other animals mtDNAs<sup>20-22</sup>. This extension proved to be translated and localized in both inner and outer mitochondrial membranes<sup>23,24</sup>, leading to the hypothesis that it could act as a mitochondrial tag implicated in paternal mitochondria survival in male embryos<sup>21</sup>. Such a 3'-coding extension of the Mcox2 gene has also been found in the mytilid mussel Musculista senhousia, but in a duplicated version of the cox2 gene<sup>25</sup>. The extension is apparently absent from M mtDNAs of other mytilid DUI species (e.g., Mytilus spp.)<sup>12,26</sup>. In the family Veneridae, the M mtDNA of Meretrix lamarckii presents an insertion of 100 codons within the cox2 gene<sup>19</sup>, whereas a duplicated version of the cox2 gene, similar to that of M. senhousia (i.e., longer at 3'), has been found in Ruditapes philippinarum, but located in the F genome (unpublished GenBank annotation mentioned in Passamonti et al.<sup>25</sup>). As suggested by Bettinazzi et al.<sup>19</sup>, non-canonical features of the cox2 gene are often coupled with DUI, but it is difficult to propose a general function because each major lineage of bivalves that possesses the DUI system exhibits some novel features. Clearly further analyses involving additional species are required to better understand the relationship between these structural variations in the cox2 gene and DUI, as well as other general features of DUI-exhibiting mitochondrial genomes.

To date, DUI has been found in over one hundred bivalve species representing four taxonomic orders and twelve bivalve families (reviewed in Gusman *et al.*<sup>27</sup>). In these species, complete F and M mtDNAs have been sequenced for ~25 DUI spp. of the families Mytilidae (order Mytilida), Veneridae (order Venerida), Unionidae, Margaritiferidae and Hyriidae (order Unionida)<sup>6,7,22,28-35</sup>. To our knowledge, all these species share one DUI-specific feature: they contain additional sex-specific mitochondrial genes without recognizable homologies to other known genes (hereafter called mitochondrial ORFans or mtORFans). These F- and M-specific mtORFans (which have been shown to be expressed) have been respectively called F-orf and M-orf<sup>6-8,36-38</sup>. This discovery is particularly interesting because these mtORFans could be responsible for the different mode of transmission of the mtDNAs and/or the functioning of DUI in bivalves. In unionid freshwater mussels, their discovery established a strong link between DUI and the maintenance of gonochorism (hermaphroditic species possess SMI and usually a highly modified F-orf gene, called H-orf<sup>7,22,39</sup>. Otherwise, their predicted functions support their direct involvement in the DUI mechanism: F-ORF proteins are suggested to interact with nucleic acids, adhere to membranes, and have roles in signalling, and M-ORFs are suggested to interact with the cytoskeleton and take part in ubiquitination and apoptosis<sup>7,8,40</sup>. However, the precise nature of the link between DUI and sex determination and the functions of the F and M mtORFans remains currently unknown. Again, further analyses involving additional species are needed to shed light on this.

Recently, DUI has been discovered in *Scrobicularia plana*<sup>27</sup>, a species of the family Semelidae (order Venerida)<sup>41</sup>, as well as in *Limecola balthica*<sup>42</sup>, a species of the family Tellinidae (order Venerida)<sup>41</sup>, and except for the complete F mtDNA of *L. balthica*<sup>43</sup>, their M mtDNAs and F mtDNA, for *S. plana*, have not previously been sequenced or reported on until now. In this study, we present the complete M and F mitochondrial genomes of these two DUI species. Our main objective was to highlight both unique features and characteristics shared among DUI species of different bivalve families. Obtaining the complete sex-specific mitogenome sequences of the species *S. plana* was particularly interesting to better understand the hypothesized link between DUI and sex determination since an "intersex" condition, i.e., the appearance of oocytes in male gonads following endocrine disruption, has been reported in this species and was associated with down-regulation of male mitochondrial transcripts in males exhibiting intersex compared to "normal" males (see Gusman *et al.*<sup>27</sup> for details). Overall, besides indicating that the newly sequenced mitogenomes may be carrying sex-specific genes like in other DUI species, our data reveal that the *cox2* gene in the M mitogenome of *S. plana* is the largest reported so far in the Kingdom Animalia. It remains to be demonstrated if such unorthodox features play key roles in DUI and sex determination in bivalves.

#### **Results and Discussion**

**Main genomic features.** The complete F mtDNAs of *Scrobicularia plana* and *Limecola balthica* are 16,170 bp and 17,492 bp in length, respectively, whereas the complete M mtDNAs are respectively 26,270 bp and 24,792 bp long (Fig. 1). To our knowledge, these values represent the biggest differences in genome size between the F and M mtDNAs in species with DUI (Table 1), which usually do not exceed 2 kb. These differences in length are partly explained by the presence of an insertion in the protein-coding gene (PCG) *cox2* in the M mtDNA of both species (*Fcox2* = 855 bp and *Mcox2* = 4,815 bp in *L. balthica*, whereas *Fcox2* = 861 bp and *Mcox2* = 5,679 bp in *S. plana*) (Table 2), which is discussed in more details below. Otherwise, the two sex-specific mtDNAs in each species possess the 13 typical mitochondrial PCGs of metazoans<sup>1</sup>, all present on the same strand, as is the case in most bivalve species<sup>14,19,44</sup>. Gene organisation is similar among the four genomes, except for the region between *cytb* and *trnM* in the M mtDNA of *S. plana*. Reorganisation events, such as inversion or transposition, as well as tandem duplication followed by random loss events, are common in animals mtDNAs<sup>1</sup>.

The initiation and termination codons for the typical 13 PCGs encoded by the four mitogenomes are presented in Table 2. There are differences between sex-specific mtDNAs within species and also between species. For example, in *S. plana* a similar start codon between the F and M mtDNA for a particular gene is observed in 4 cases out of 13 (only two in *L. balthica*) whereas 6 PCGs out of 13 have a similar stop codon (4 in *L. balthica*). Overall, most of the PCGs use the ATD start codon (where D means A, T or G) found in metazoan mtDNAs<sup>45</sup>: ATA occurs the most (15/52), followed by ATG (14/52) and ATT (11/52). TTG is also found in seven cases, and GTG in five, an observation that has been previously reported in molluscan species<sup>46,47</sup>. Most of the PCGs are



**Figure 1.** Gene maps of M and F genomes of *Scrobicularia plana* and *Limecola balthica*. All genes are encoded on the heavy strand, total genome lengths are reported inside their corresponding genome. Gene colors correspond to functional groups (OXPHOS gene families, tRNAs and rRNAs).

terminated with the TAA (27/52) and TAG (20/52) codons. Some incomplete termination codons are also found, which are assumed to be completed by polyadenylation of their mRNAs<sup>48</sup>.

The usual 22 tRNAs are found in the four mitochondrial genomes, ranging from 61 to 71 bp, and most of them can be folded into the typical secondary structures (not shown). There is an exception in the M mtDNA of *L. balthica*, which is lacking *trnT*, but instead possess a second copy of *trnA*, which has been identified by MITOS in the same area, between *trnY* and *trnL1* (Fig. S1). This second copy of *trnA* (*trnA\_1*) does not possess the typical secondary structure and it is not clear if it could be functional. This same mitochondrial genome contains also another copy of *trnP* (*trnP\_1*) located next to the first *trnP* (*trnP\_0*), and both possess usual secondary structures (Fig. S1). Additional tRNA copies are common in bivalve mitochondrial genomes<sup>19,49–51</sup>. Concerning the missing *trnT*, it is possible that the *trnA\_1* could be a mutated *trnT* (with a mutation in its anticodon that changed it from *trnT* to a "secondary" *trnA*). It is also possible that an unusual secondary structure makes it unidentifiable by the tRNA search programs, and this will require further analyses.

Length variations among rRNA genes range from 765 bp to 896 bp and from 1249 bp to 1556 bp for 12S (*rrnS*) and 16S (*rrnL*), respectively (Table 2). Their locations are almost identical in the four genomes; *rrnL* is found between *nad6* and *atp6* whereas *rrnS* is found between *trnG* and *trnM*, except for the M genome of *S. plana*, in which it is located between *trnW* and *trnV* (Fig. 1; Tables 2 and 3).

**Intraspecific divergences.** Genetic distances of individual genes between F and M mtDNAs were analyzed in each species. The results show that the level of conservation is higher for both ribosomal RNA genes (*rrnS* and *rrnL*), and also for the PCG *cox1*, whereas NADH dehydrogenase subunit genes (*nad* series) and *atp8* for *L. balthica* are generally less conserved (Fig. 2a,c). These results are consistent with the general findings in DUI bivalves<sup>32</sup>. For *S. plana* the average nucleotide divergence (measured as *p*-distance) of combined protein coding and ribosomal RNA genes is  $0.43 \pm 0.08$  SD (based on values in Fig. 2a). For *L. balthica* the average *p*-distance of

Species	Mitotype	Genome Size (bp)	Genome size difference (bp)	GenBank Accession Number	Reference	
Freshwater mussels (Unionic	la, Unionidae	e)				
Anodonta anatina	F	15,637		KF030965	28	
Anodonta anatina	М	16,906	1,269	KF030963	29	
Freshwater mussels (Unionic	la, Hyriidae)					
Hyridella menziesi	F	16,031		KU873121	22	
Hyridella menziesi	М	18,140	2,109	KU873122	22	
Freshwater mussels (Unionic	la, Margaritif	feridae)				
Cumberlandia monodonta	F	16,099		KU873123	22	
Cumberlandia monodonta	М	17,575	1,476	KU873124	22	
Marine mussels (Mytilida, M	ytilidae)		-			
Musculista senhousia	F	21,557		GU001953	17	
Musculista senhousia	М	20,621	936	GU001954	17	
Marine clams (Venerida, Ven	eridae)		-			
Meretrix lamarckii	F	20,025		KP244451	19	
Meretrix lamarckii	М	19,688	337	KP244452	19	
Marine clams (Venerida, Tell	inidae)		-			
Limecola balthica	F	17,492		MN528028	Present study	
Limecola balthica	М	24,792	7,300	MN528029	Present study	
Marine clams (Venerida, Sen	nelidae)		-			
Scrobicularia plana	F	16,170		MN528026	Present study	
Scrobicularia plana	М	26,270	10,100	MN528027	Present study	

**Table 1.** Mitochondrial genome sizes in DUI species exhibiting extreme length differences between M and F types.

combined protein coding and ribosomal RNA genes is  $0.41 \pm 0.10$  SD (based on values in Fig. 2c). The number of nonsynonymous substitutions per nonsynonymous sites (Ka) relative to the number of synonymous substitutions per synonymous sites (Ks) were also calculated (shown in Fig. 2b,d). This analysis provides an estimate of the degree of selection (either neutral, positive, or purifying) on each PCG. For both species and for all PCGs, all data points fall below the line representing neutrality (expressed as Ka = Ks; see dashed line in Fig. 2b,d), suggesting that the PCGs in S. plana and L. balthica accumulate more synonymous substitutions over evolutionary time relative to nonsynonymous substitutions. Therefore, all Ka and Ks rates supported an initial hypothesis of purifying selection for mitochondrial genes. To statistically test if the genes were indeed under purifying selection, we conducted a Z-test of Selection<sup>52</sup>. For all PCGs in both species, with the exception of *atp8* in *L. balthica*, there is strong statistical support (at  $\alpha = 0.05$ ) for rejecting a null hypothesis of neutrality (H<sub>n</sub>: Ka = Ks) in support of an alternate hypothesis for purifying selection ( $H_a$ : Ka < Ks). This is because all pairwise Z-test p-values were <0.01 for both hypotheses. For *atp8* in *L. balthica* the Z-test of Selection did not support rejecting neutrality (p-value = 0.277 for  $H_n$ : Ka = Ks), a result that might be affected by the small size of this gene (129 bp), which in turn altered the statistical power of the test. With regards to the large insertion within the male cox2 gene for both species, these data are of even greater interest as they suggest that this insertion likely does not render the gene functionless. Rather, cox2 (the alignable parts) actually remains among one of the more relatively conserved mt genes (Fig. 2). Overall, our results are in line with what has been observed in other bivalves with DUI, i.e. despite the considerable divergence of DNA and amino acid sequences of PCGs there is evidence of strong purifying selection acting on these mitochondrial genes<sup>32</sup>.

For comparative purpose, p-distances of individual genes (PCGs only) between F and M mtDNAs were also calculated for three species known for having the greatest F versus M DNA divergences in the families Mytilidae, Veneridae and Unionidae: i.e., the mytilid Modiolus modiolus<sup>53</sup>, the venerid Ruditapes philippinarum<sup>17</sup>, and the unionid *Quadrula quadrula*<sup>14</sup> (Table 4). Our results show an average *p*-distance (for all 13 PCGs) of 0.45 for nucleotide (nt) and 0.53 for amino acid (aa) in S. plana and 0.44 (nt) and 0.53 (aa) in L. balthica (Table 4). To our knowledge, the nucleotide and amino acid divergences reported in S. plana and L. balthica are the greatest reported among DUI species, surpassing those in freshwater mussels (Table 4), which were previously thought to exhibit the greatest divergences between their sex-specific mtDNAs<sup>14</sup>. On the other hand, the average uncorrected nucleotide divergence observed between the F and M PCGs of the marine mussel Mytilus edulis is about  $0.23^{26}$ . This low level of divergence has been proposed to be a consequence of masculinization events, which are characterized by an invasion of the male route of inheritance by an F mtDNA that becomes transmitted through sperm as a standard M mtDNA<sup>12,54</sup>. These events reset the level of divergence between the F and M mitochondrial genomes to zero<sup>12,54</sup>. Conversely, the high level of divergence observed in freshwater mussels has been hypothesized to be a consequence of a complete absence of masculinization events for over 200 million years in Unionida<sup>14,20</sup>. According to these studies, masculinization would be no longer possible in this taxon because of the existence of the M-specific extension of the cox2 gene, a specialized feature of the unionid M mtDNA that would prevent recombination between the F and M mtDNAs, i.e. a step necessary for masculinization to

Scrobiculari	a plana		Scrobicularia	a plana		Limecola bal	thica		Limecola balthica		
F mt genom	t genome			M mt genome			;	-	M mt genome		
Gene/ Region	Length	Start/Stop	Gene/ Region	Length	Start/Stop	Gene/ Region	Length	Start/Stop	Gene/ Region	Length	Start/Stop
cox1	1554	ATA/TAA	cox1	1551	ATT/TAA	cox1	1554	ATA/TAA	cox1	1536	ATT/TAA
nad4	1323	TTG/TAG	nad4	1335	GTG/TAA	nad4	1341	ATA/TAG	nad4	1341	TTG/TAA
trnH	65		trnH	63		trnH	64		trnH	64	
trnS2	64		trnS2	65		trnS2	65		trnS2	64	
trnE	66		trnE	64		trnE	67		trnE	66	
nad3	348	ATA/TAA	nad3	348	ATA/TAA	nad3	348	ATA/TAG	nad3	348	ATG/TAA
trnI	68		trnI	68		trnI	69		trnL	68	
trnK	64		trnK	66		trnK	65		trnK	65	
nad4L	291	ATT/TAA	nad4L	285	ATT/TAG	nad4L	291	GTG/TAA	nad4L	276	TTG/TAG
trnY	64		trnY	64		trnY	63		trnY	67	
trnT	63		trnT	64		trnT	64		trnA_1	71	
trnL1	66		trnL1	65		trnL1	66		trnL1	65	
trnD	63		trnD	66		trnD	63		trnD	65	
trnL2	66		trnL2	66		trnL2	66		trnL2	65	
nad1	924	GTG/TAG	nad1	927	ATT/TAG	nad1	924	ATG/TAG	nad1	927	ATA/TAG
trnN	66		trnN	65		trnN	65		trnN	65	
nad5	1635	TTG/TAA	nad5	1704	ATG/TAG	nad5	1743	TTG/T**	nad5	1767	ATA/TAG
trnR	67		trnR	65		trnR	65		trnR	65	
cob	1135	ATA/T**	cob	1135	ATG/T**	cob	1140	ATT/TAA	cob	1137	ATA/TAA
cox2	861	ATG/TAA	trnW	64		cox2	855	ATG/TAA	cox2	4815#	ATG/T**
trnV	69		rrnS	765		trnV	66		trnV	65	
trnW	66		trnV	66		trnW	66		trnW	67	
trnG	68		trnG	68		trnG	66		trnG	66	
rrnS	896		cox2	5679	ATG/TAG	rrnS	880		rrnS	886	
trnM	66		trnM	67		trnM	65		trnM	64	
atp8	129	ATT/TAG	atp8	129	ATG/TAA	atp8	129	ATT/TAA	atp8	130	TTG/T**
trnS1	71		trnS1	69		trnS1	69		trnS1	69	
nad6	510	ATA/TAA	nad6	510	ATT/TAG	nad6	510	ATA/TAA	nad6	513	TTG/TAG
rrnL	1269		rrnL	1556		rrnL	1249		rrnL	1379	
atp6	714	ATG/TAG	atp6	705	ATT/TAA	atp6	735	ATG/TAA	atp6	741	GTG/TAG
cox3	882	ATA/TAG	cox3	882	ATA/TAG	cox3	882	ATA/TAA	cox3	885	GTG/TAA
nad2	996	ATG/TAA	nad2	1002	ATT/TAA	nad2	990	ATG/TAA	nad2	987	ATG/TAG
trnP	66		trnP	67		trnP	66		trnP_1	64	
trnQ	67		trnQ	67		trnQ	66		trnP_0	65	
trnC	63		trnC	61		trnC	61		trnQ	66	
trnA	65		trnA	65		trnA	67		trnC	63	
trnF	63		trnF	65		trnF	66		trnA_0	66	
									trnF	65	

**Table 2.** General features of *Scrobicularia plana* and *Limecola balthica* M and F mitochondrial genomes. NOTE – T\*\* stands for incomplete stop codon; <sup>#</sup>The exact size of the M*cox2* gene in *Limecola balthica* is uncertain.

.....

occur<sup>12,14,20</sup>. We thus propose that the high divergences observed between the F and M mtDNAs in *S. plana* and *L. balthica* are also related to an absence of masculinization events in these species because of the insertion in their Mcox2 gene (described below).

**Mitochondrial ORFans.** Unassigned regions in the coding strand were searched for the presence of supernumerary PCGs and mtORFans with a minimal length of 150 bp. Only ORFs encoding proteins with at least one predicted transmembrane domain were retained, because all sex-specific mtORFans characterized to date in DUI species (i.e. those encoding F-ORF in females and M-ORF in males), except for one case, possess at least one transmembrane domain or helix (TMH)<sup>7,8,22,40</sup>. In both *S. plana* and *L. balthica* F mtDNAs, only two unassigned regions were susceptible to contain supernumerary ORFs of >150 bp, i.e. between *rrnS-trnM* and *trnF-cox1* (Table 3). However, the region between *trnF* and *cox1* was discarded because of the presence of a possible 5' extension of the *cox1* gene in all four genomes, an issue that will need to be assessed by looking at expression data. Otherwise, no ORFs corresponding to the expected profile were found in the F mtDNAs of *S. plana* and *L. balthica*. However, it is conceivable that smaller ORFs or ORFs encoding proteins without predicted transmembrane domain could be involved in DUI in these distantly-related species. The smallest F-ORF identified to

Boundary	S. plana mtDNA	F	Boundar	y	S. plana M mtDNA	Boundar	y	<i>L. balthica</i> F mtDNA	Boundary		<i>L. balthica</i> M mtDNA
cox1	nd4	10	cox1	nd4	60	cox1	nd4	0	cox1	nd4	15
nd4	trnH	2	nd4	trnH	4	nd4	trnH	6	nd4	trnH	13
trnH	trnS2	0	trnH	trnS2	55	trnH	trnS2	1	trnH	trnS2	16
trnS2	trnE	0	trnS2	trnE	15	trnS2	trnE	1	trnS2	trnE	2
trnE	nd3	15	trnE	nd3	6	trnE	nd3	15	trnE	nd3	18
nd3	trnI	10	nd3	trnI	-17	nd3	trnI	13	nd3	trnI	-17
trnI	trnK	3	trnI	trnK	1	trnI	trnK	2	trnI	trnK	2
trnK	nd4L	0	trnK	nd4L	2	trnK	nd4L	0	trnK	nd4L	10
nd4L	trnY	1	nd4L	trnY	7	nd4L	trnY	4	nd4L	trnY	1
trnY	trnT	-3	trnY	trnT	-2	trnY	trnT	-2	trnY	trnA_1	-2
trnT	trnL1	0	trnT	trnL1	1	trnT	trnL1	0	trnA_1	trnL1	3
trnL1	trnD	-1	trnL1	trnD	-1	trnL1	trnD	-1	trnL1	trnD	-1
trnD	trnL2	2	trnD	trnL2	2	trnD	trnL2	1	trnD	trnL2	6
trnL2	nd1	1	trnL2	nd1	1	trnL2	nd1	1	trnL2	nd1	-14
nd1	trnN	3	nd1	trnN	-19	nd1	trnN	1	nd1	trnN	5
trnN	nd5	0	trnN	nd5	25	trnN	nd5	0	trnN	nd5	0
nd5	trnR	-40	nd5	trnR	4	nd5	trnR	1	nd5	trnR	-10
trnR	cob	3	trnR	cob	72	trnR	cob	31	trnR	cob	111
cob	cox2	110	cob	trnW	610	cob	cox2	71	cob	cox2	216
cox2	trnV	5	trnW	rrnS	3231	cox2	trnV	3	cox2	trnV	533
trnV	trnW	0	rrnS	trnV	101	trnV	trnW	-1	trnV	trnW	0
trnW	trnG	2	trnV	trnG	18	trnW	trnG	5	trnW	trnG	15
trnG	rrnS	4	trnG	cox2	1272	trnG	rrnS	3	trnG	rrnS	410
rrnS	<i>trnM</i>	732	cox2	trnM	57	rrnS	<i>trnM</i>	1890	rrnS	<i>trnM</i>	3697
<i>trnM</i>	atp8	11	<i>trnM</i>	atp8	13	<i>trnM</i>	atp8	9	<i>trnM</i>	atp8	11
atp8	trnS1	-2	atp8	trnS1	28	atp8	trnS1	3	atp8	trnS1	-1
trnS1	nd6	32	trnS1	nd6	20	trnS1	nd6	34	trnS1	nd6	-2
nd6	rrnL	0	nd6	rrnL	0	nd6	rrnL	35	nd6	rrnL	12
rrnL	atp6	54	rrnL	atp6	180	rrnL	atp6	47	rrnL	atp6	40
atp6	cox3	30	atp6	cox3	8	atp6	cox3	52	atp6	cox3	17
cox3	nd2	34	cox3	nd2	56	cox3	nd2	81	cox3	nd2	105
nd2	trnP	52	nd2	trnP	3	nd2	trnP	0	nd2	trnP_1	78
trnP	trnQ	11	trnP	trnQ	25	trnP	trnQ	13	trnP_1	trnP_0	7
trnQ	trnC	15	trnQ	trnC	0				trnP_0	trnQ	35
trnC	trnA	0	trnC	trnA	4	trnQ	trnC	-1	trnQ	trnC	0
trnA	trnF	-1	trnA	trnF	3	trnC	trnA	1	trnC	trnA_0	13
trnF	cox1	161	trnF	cox1	372	trnA	trnF	-1	trnA_0	trnF	20
						trnF	cox1	163	trnF	cox1	250

**Table 3.** Number of nucleotides at gene boundaries in F and M mitochondrial genomes of *Scrobicularia plana* and *Limecola balthica*. NOTE – Negative values indicate that genes overlap.

date in a DUI species, i.e. in the unionid *Venustaconcha ellipsiformis*, is encoding an 89aa-long protein with one predicted TMH<sup>7,40</sup>, whereas the smallest M-ORF is potentially encoding a 30aa-long protein without TMH in *Mytilus californianus*<sup>8</sup>, although the functionality of this latter ORF remains to be demonstrated. Additional F mt sequences and expression data from *S. plana* and *L. balthica* will be necessary to clearly demonstrate the presence (or absence) of the F-*orf* gene in these species.

In both male mitochondrial DNAs, four unassigned regions contained supernumerary ORFs of >150 bp, i.e. between *cob-trnW*, *trnW-rrnS*, *trnG-cox2* and *rrnL-atp6* for *S. plana* and between *cob-cox2*, *cox2-trnW*, *trnG-rrnS* and *rrnS-trnM* for *L. balthica* (Table 3). Five ORFs corresponding to the expected profile were found in *S. plana*, two between *trnW-rrnS* and three between *trnG-cox2*, whereas four ORFs with the expected profile were found in *L. balthica*, one between *trnG-rrnS* and three between *rrnS-trnM* (Fig. S2). Sequence similarity searches using PSI-BLAST<sup>55</sup> against non-redundant protein sequences and SWISSPROT databases failed to detect significant sequence similarity with known proteins for all these ORFs except one (i.e. mtORFan1 in the unassigned region *trnG-rrnS* of *L. balthica* M mtDNA; Fig. S2). For this sequence, our results revealed moderately significant hits (E-values 2e-07) with microtubule-associated proteins. This result is interesting since previous *in silico* analyses of M-ORF sequences in other DUI species also indicated connections with cytoskeleton proteins involved in microtubule-binding (e.g. ankyrin)<sup>8,40</sup>. These observations led to the hypothesis that M-ORF, with its predicted transmembrane domains, may target sites outside sperm mitochondria and be responsible for





their cellular positioning in developing embryos<sup>8</sup>. Indeed, studies have shown that, after fertilization, only in male DUI embryos sperm mitochondria remain grouped together, and are eventually sequestered in the germ line, whereas they are dispersed and/or destroyed in female embryos (reviewed in Zouros<sup>12</sup>). M-ORFs are thus considered as ideal candidates for M mtDNA-derived masculinizing factors in DUI species<sup>8,40</sup>.

At this moment, however, we cannot confirm nor disprove the presence of a M-*orf* gene in the M mtDNAs of *S. plana* and *L. balthica*. Pairwise alignments between the ORFs found in both species were performed but it was not possible to obtain satisfactory alignments that could provide support for identifying a conserved ORF (data not shown). Again, additional M mt sequences and expression data from *S. plana* and *L. balthica* will be necessary to test for the presence of a M-*orf* gene in these species.

**Insertion in the male cox2 gene.** Annotations with MITOS revealed an insertion of >4.5 kb and >3.5 kb in the *Mcox2* gene of *S. plana* and *L. balthica*, respectively, which is absent in the *Fcox2* genes of both species (Figs. 1 and 3). In *S. plana*, this in-frame insertion, if translated, means that the *cox2* gene would be 5,679bp-long and, to our knowledge, would therefore encode for the longest *COX2* protein in the animal kingdom (i.e. 1,893 amino acids). The sequencing of additional *S. plana* male individuals revealed that this insertion is conserved among different individuals from different populations (Tassé *et al.* unpublished), indicating that it is most probably functional. A multiple sequence alignment of *Homo sapiens COX2* and *S. plana* FCOX2 and *MCOX2* amino acids indicates that the insertion is situated between the "heme-patch" region, containing an important residue Trp that functions as the point of electron entry from Cytochrome C, and the first Cu<sub>a</sub>-binding center (Figs. 3

	cox1	nad4	nad3	nad4L	nad1	nad5	cob	cox2	atp8	nad6	atp6	cox3	nad2	Average
(a)														
Scrobicularia plana	0.335	0.466	0.482	0.524	0.435	0.516	0.336	0.419	0.437	0.536	0.420	0.436	0.483	0.449
Limecola balthica	0.305	0.481	0.426	0.446	0.391	0.467	0.329	0.410	0.540	0.544	0.453	0.421	0.513	0.440
Quadrula quadrula	0.287	0.383	0.374	0.404	0.414	0.390	0.365	0.348	0.358	0.475	0.384	0.361	0.443	0.384
Modiolus modiolus*	0.280	0.375	0.373	—	-	—	—	-	-	-	0.466	0.375	—	0.374
Ruditapes philippinarum	0.222	0.300	0.278	0.468	0.240	0.299	0.281	0.336	-	0.313	0.298	0.274	0.347	0.305
(b)														
Scrobicularia plana	0.288	0.610	0.604	0.500	0.480	0.618	0.329	0.506	0.595	0.701	0.493	0.517	0.614	0.527
Limecola balthica	0.300	0.610	0.541	0.527	0.474	0.621	0.332	0.515	0.619	0.690	0.571	0.485	0.663	0.534
Quadrula quadrula	0.318	0.591	0.492	0.763	0.549	0.572	0.380	0.420	0.500	0.722	0.523	0.440	0.661	0.533
Modiolus modiolus*	0.233	0.445	0.449	-	-	-	—	-	-	-	0.576	0.422	—	0.425
Ruditapes philippinarum	0.123	0.334	0.303	0.779	0.216	0.346	0.273	0.419	-	0.560	0.343	0.314	0.444	0.371

**Table 4.** Intraspecific divergence (uncorrected *p*-distances) of the 13 mitochondrial protein-coding genes between M and F mtDNAs in DUI species known for having the greatest F versus M DNA divergences. (a) nucleotides (b) amino acids. NOTE – \*The mtDNA sequence for *Modiolus modiolus is incomplete*.



**Figure 3.** Fcox2 and Mcox2 structural features in *Scrobicularia plana* and *Limecola balthica*. For the sequence alignments of the "heme-patch" regions and  $Cu_a$  centers, identical amino acids are indicated by an \*.

.....

and S3), and *in silico* analyses suggest that it does not contain any additional transmembrane domain (Tassé *et al.* unpublished). In-frame insertions resulting in enlarged *cox2* genes have also been reported in hymenoptera, ciliates, brown algae and microflagellates<sup>56</sup>. As mentioned above, modifications to the *cox2* gene is a particular feature often found in DUI species. In unionid mussels, the longest *cox2* gene is found in the M mtDNA of *Hyridella menziesii* (1,380 bp)<sup>22</sup> and is the result of a 3' extension, as in other DUI unionids<sup>21</sup>. Previous studies indicated an extra-mitochondrial localization of MCOX2 as well as a possible involvement of the protein in reproduction in freshwater mussels<sup>23,24</sup>, further supporting the link between gender and mtDNA transmission patterns. In the order Venerida, i.e. the order to which *S. plana* and *L. balthica* belong, an in-frame insertion of 300 nt has been reported in the species *Meretrix lamarckii* (family Veneridae)<sup>19</sup>. However, such modifications of the *cox2* gene have not been reported in marine mytilid mussels, i.e. *Mytilus* spp., and in the venerid clam *Ruditapes philippinrum*, a duplicated copy of the *cox2* gene with a 3' extension has been reported, but in the F mtDNA<sup>12,19</sup>.

The situation in *L. balthica* is quite different than in *S. plana* regarding the Mcox2 gene, which is split in two by an insertion. Specifically, this insertion divides Mcox2 into Mcox2a, encoding the two transmembrane helices and the "heme-patch" region followed by a complete stop codon (TAA), and Mcox2b, encoding an enlarged intermembrane space and the Cu<sub>a</sub> centers (Figs. 3 and S4). This situation is confirmed by transcriptomic data (Illumina RNAseq reads from sperm cells, used to polish our nanopore reference mitogenomes), which indicate that both regions are transcribed (i.e. with discrete, non-overlapping Mcox2a and Mcox2b transcripts). Although it remains to be determined whether Mcox2a and Mcox2b are translated, it is worth noting that similar cases have been described in the freshwater mussel Anodonta cygnea<sup>33</sup> and the Hymenoptera Campsomeris spp<sup>56</sup>. For example, a translocation of a portion of the nad5 gene has been reported in *A. cygnea*, and since the translocated portion and the rest of the gene both possessed their own start codon, the authors proposed that they could be transcribed and translated separately<sup>33</sup>. In the genus Campsomeris, cox2 is also split into two genes and this likely occurred through intragenic insertion of a cluster of several ORFs, one of which encodes a putative endonuclease that might have been directly involved in the process of *cox2* fission<sup>56</sup>. However, no ORFs with similarity to an endonuclease were found in the *Mcox2* insertion of *L. balthica* (nor in *S. plana*). According to Szafranski *et al.*<sup>56</sup>, COXIIA and COXIIB polypeptides in *Campsomeris* spp. apparently assemble into a functional COXII heterodimer. Further studies will be needed to determine if this is also the case in *L. balthica*, or to clearly verify if a process of mitochondrial RNA splicing, because a group II intron has already been reported in the *cox1* gene of Annelida<sup>57</sup>, could be involved (i.e. *Mcox2* RNA splicing or *Mcox2a* and *Mcox2b* RNA trans-splicing into a single mRNA).

It also remains to be determined if *L. balthica* possesses an enlarged MCOX2 or a MCOX2 protein of a typical length. For example, the stop codon in Mcox2a could be read as a sense codon by a suppressor tRNA, i.e. a mutated (usually) tRNA that would insert an amino acid instead of initiating termination, as seen in several eukaryotes<sup>58</sup>. The presence of duplicated *trn* genes in the M mtDNA of *L. balthica* might be a clue for such scenario. Modification or duplication of tRNAs have already been proven to allow codon reassignment by changing codons from stop-to-sense or sense-to-sense in yeast, green algae and metazoan mitochondria<sup>59,60</sup>. Alternatively, termination can be avoided by ribosomal frame-shifting, a process that allows for protein merging from two or more overlapped ORFs<sup>61</sup>. This has been reported for the mitochondrial *cox1* gene of the diatom *Phaeodactylum tricornutum*: to avoid a stop codon, translational frameshift skips a nucleotide (called + 1)<sup>62</sup>. Similar strategies (+1 or -1 frameshifts) have also been recorded in metazoan mitochondrial genomes including ants, turtles and humans<sup>63-65</sup>. Further transcriptomic and proteomic analyses, and a closest look in mitochondrial tRNAs, will be required to validate any machinery involved in Mcox2 transcription and translation in both *L. balthica* and *S. plana*. The results presented in this study clearly indicate that the relationship between *cox2* variations and DUI deserves much greater attention.

#### Conclusion

In summary, the description and comprehensive analysis of complete F and M mtDNAs of two newly discovered bivalve species with DUI from two additional families (Semelidae and Tellinidae), have led to new insights into DUI mitogenomics. Our results revealed uncorrected amino acid *p*-distance of ~53% for PCGs between the M and F genomes of both species: this is the highest divergences reported in DUI species thus far. Hence the way (or ways) in which males of bivalves with DUI can tolerate heteroplasmy characterized by such high variability remains an important unanswered question (e.g. see Bettinazzi *et al.*<sup>66</sup>). Our results also highlighted an extremely unusual feature of the M genomes of *S. plana* and *L. balthica* compared to their female-transmitted counterparts. This feature is the presence of an important in-frame insertion (>3.5 to 4.5 kb) in the *Mcox2* gene. This is the longest insertion reported in the Kingdom Animalia, which remains to be functionally characterized. The reported data further indicate that the newly-sequenced M mitogenomes may be carrying lineage-specific genes (mtORFans) possibly involved in the DUI process. Analyses of complete mtDNAs from additional bivalve species and further protein-based studies are needed to elucidate the number, taxonomic distribution, evolution, and function of mtORFans and atypical *cox2* genes in this group, as well as the molecular mechanisms underlying DUI.

#### Methods

**Specimen collection and sample preparation.** Adult specimens of *Limecola balthica* were collected at low tide on the Aytré sandy mudflat (France; 46.1259 N, 1.1284 W) in April 2016 and May 2017 (these are periods during which gonads are known to be near sexual maturation<sup>42</sup>. Individuals were carefully opened and gonad tissue nicked with a micro-scalpel. Gametes were then sampled with a micropipette and a small amount was used to sex the animal with a light microscope; the remainder of the sample was flash-frozen in liquid nitrogen and stored at –80 °C. Adult specimens of *Scrobicularia plana* were collected in May 2013 from Concarneau (France; 47.8728°N, 3.9207°W). Individuals were dissected and gonads inspected under the microscope. Female and male mature gonads were conserved in 95% Ethanol and sent to the Université de Montréal.

For *S. plana*, total genomic DNA was extracted separately from male and female gonad tissues with a Qiagen DNeasy Blood & Tissue Kit (QIAGEN Inc., Valencia, CA) using the kit provided animal tissue protocol. The quality and quantity of DNA were assessed by electrophoresis on a 1% agarose gel and also with a BioDrop µLITE spectrophotometer. One female and one male gonad sample with the highest purity and concentration were chosen for sequencing. For *L. balthica*, total genomic DNA from male gonadal tissues (majority of gametic cells, and some somatic carry-over) was extracted using the phenol-chloroform protocol described in<sup>67</sup>, resuspended in molecular grade water, and quantified using a Nanodrop 2000 and a Qubit fluorometer (dsDNA HS assay kit, Molecular Probes). DNA integrity was also checked by electrophoresis on a 1% agarose gel. One sample characterised by the highest purity and concentration was chosen for sequencing.

**Mitochondrial genome sequencing.** Total DNA extractions from *S. plana* male and female gonads were respectively used to prepare two DNA libraries that were paired-end sequenced  $(2 \times 100 \text{ bp})$  using an Illumina HiSeq platform (Génome Québec Innovation Centre, Montréal, QC, Canada). Paired reads were trimmed with Trimmomatic  $0.32^{68}$  and merged with Pear  $0.9.6^{69}$  on the Galaxy online platform<sup>70</sup>, checking the quality of reads at every step with FastQC  $0.11.5^{71}$ . MITObim  $1.8^{72}$  was used to assemble the complete genome sequence, starting from a *S. plana* partial *cox1* sequence (GenBank accession number KX447421 for the male mtDNA and KX447423 for the female mtDNA) as an initial seed.

For *L. balthica*, the library preparation was done following the SQK-LSK108. Nanopore protocol. A DNA shearing step (using Covaris gTubes) was included to linearize circular mtDNA molecules (1 min at 11k RPM in a 5415 R Eppendorf centrifuge). The resulting library was sequenced on the Minion MIN-101B sequencer using a R9 flowcell. Basecalling was performed using Albacore 2.1.3<sup>73</sup>, followed by quality control using Minion\_QC

1.074. Nanopore adapters were removed and chimeric reads were discarded using Porechop 0.2.3 (Wick R. Porechop, available at: https://github.com/rrwick/Porechop). Reads with quality scores <9 were removed using Nanofilt 2.0.075. Cleaned reads were assembled using Canu 1.676 considering an approximate genome size of 2 Gb and disregarding all reads smaller than 500 bp. Resulting unitigs were blasted locally using blastn<sup>55</sup>, using male and female rrnL sequences (Genbank accession numbers KX831969 and KX831970, respectively)<sup>42</sup>. Based on these results, we identified tig00000009 (length of 26,120 bp, assembly based on 61 nanopore reads) as the male mitogenome. We did not retrieve the female mitochondrial genome as one contig but as two (tig00000888: 16,790 bp, 34 reads; tig00000885: 6,319 bp, 1 read). Basecalling errors were corrected using signal-level data with Nanopolish  $0.9.0^{77}$ . We used these two cleaned contigs as reference genomes to map Illumina  $2 \times 150$  bp HiSeq sequences (data quality control and pre-processing step as in<sup>78</sup>) produced from a second individual using Pilon 1.2179. HiSeq sequencing of cDNA was performed based on mRNAs purified from sperm and foot tissue (for mapping onto the male and female mitogenomes, respectively) using the Macherey-Nagel Nucleospin RNA kit for NucleoZOL (RNA quality control, cDNA synthesis, library preparation, sequencing, data quality control and pre-processing as in<sup>80</sup>). By mapping high-quality Illumina reads on our draft nanopore contigs, we were able to further correct sequencing errors. The two F-type contigs (tig00000885 and tig00000888) were then merged. The quality of the mitogenome assembly was checked at each step of the pipeline (initial Canu assembly, Nanopolish polishing, Pilon polishing) by calculating the percent identity between our F-type unitigs and a published F-type mitogenome from the same sampling locality (KM373201)<sup>43</sup> using MUMmer 3.23<sup>81</sup>.

Mitochondrial genome sequences were deposited in GenBank (accession codes MN528026 and MN528027 for the F and M mtDNAs of *S. plana*, respectively, and accession codes MN528028 and MN528029 for the F and M mtDNAs of *L. balthica*, respectively).

**Characterization of mtDNAs and sequence analyses.** The male and female mitogenome sequences obtained were annotated with MITOS2<sup>82</sup>. Protein-coding genes (PCG) were also predicted using NCBI's Open Reading Frame Finder, choosing the invertebrate mitochondrial genetic code and alternative start codons. Candidate transfer RNA (tRNA) genes were also identified using tRNAScan-SE<sup>83</sup>. The ends of 16S and 12S rRNA genes were assumed to extend to the boundaries of their flanking genes. The complete mitochondrial genomes, with their annotated PCG, tRNAs and rRNAs, were illustrated with GenomeVx online tool<sup>84</sup>.

The degree of genetic conservation among the M-versus-F genes in each species was assessed using two approaches: (i) pairwise *p*-distances, and (ii) the rates of synonymous and non-synonymous substitutions, i.e. the number of synonymous substitutions per synonymous sites [*Ks*] and number of non-synonymous substitutions per non-synonymous sites [*Ka*], which were calculated in MEGA v7<sup>52</sup>. Since there were various indels present between the individual M- and F-type genes within both species, pairwise deletion was used to account for gaps/missing data-points and codon-alignments were used to facilitate accurate calculations of genetic distances, as well as *Ka* and *Ks* values. Amino acid sequences were determined using EMBOSS Transeq and by assuming an invertebrate mitochondrial code<sup>85</sup>. Codon-alignments were generated by aligning M and F amino acid sequences for each gene using MUSCLE<sup>86</sup> in the online portal found at http://www.ebi.ac.uk/Tools/msa/muscle/84. Using these protein alignments as references, nucleotide sequences for each respective gene were then codon-aligned via PAL2NAL v14<sup>87</sup>. The exceptionally large indel within the *Mcos2* gene was manually excluded from the gene alignments. MEGA was also used to conduct a Z-test of Selection [via the Nei-Gojobori method] for each gene<sup>52,88</sup>. All plots relating to gene conservation were generated in R version 3.4.1 (R Core Team 2017) using the package ggplot2 v3.2.0<sup>89</sup>.

For comparative purpose, *p*-distances of individual genes between F and M mtDNAs were also calculated for three species known for having the greatest F versus M DNA divergences in the families Mytilidae, Veneridae and Unionidae: i.e. the mytilid *Modiolus modiolus* (GenBank accession: F, KX821782; M, KX821783)<sup>53</sup>, the venerid *Ruditapes philippinarum* (GenBank accession: F, AB065375; M, AB065374)<sup>17</sup>, and the unionid *Quadrula quad-rula* (GenBank accession: F, FJ809750; M, FJ809751)<sup>14</sup>. These comparisons facilitated a better understanding of the selection pressures acting on the PCGs in these species. Finally, the presence of supernumerary mitochondrial protein-coding genes (and mtORFans) in unassigned regions was examined using ORF Finder (NCBI). We retained only ORFs with a minimal length of 150 bp and localized on the coding strand, as mitochondrial genes are only encoded on one strand in our two species. Transmembrane domains were predicted using the TMHMM server v2.0 (http://www.cbs.dtu.dk/services/TMHMM/). Alignments between predicted ORFs were done with MUSCLE<sup>86</sup> and the percentage of identity was visualized with MView<sup>90</sup>, via the EMBL-EBI online web services<sup>85</sup> keeping default parameters.

Received: 19 October 2019; Accepted: 9 January 2020; Published online: 23 January 2020

#### References

- 1. Boore, J. L. Animal mitochondrial genomes. Nucleic Acids Res. 27, 1767-1780 (1999).
- Gissi, C., Iannelli, F. & Pesole, G. Evolution of the mitochondrial genome of Metazoa as exemplified by comparison of congeneric species. *Heredity* 101, 301–20 (2008).
- 3. Park, H. & Ahn, D. H. Complete mitochondrial genome of the Antarctic soft-shelled clam, *Laternula elliptica* (Bivalvia; Laternulidae). *Mitochondrial DNA* 26, 642-3 (2015).
- McCartney, M. A. et al. The Genome of the Zebra Mussel, Dreissena polymorpha: A Resource for Invasive Species Research. bioRxiv 696732 (2019).
- Plazzi, F., Puccio, G. & Passamonti, M. Comparative large-scale mitogenomics evidences clade-specific evolutionary trends in mitochondrial DNAs of Bivalvia. *Genome Biol. Evol.* 8, 2544–2564 (2016).

- Breton, S. *et al.* Comparative mitochondrial genomics of freshwater mussels (Bivalvia: Unionoida) with Doubly Uniparental Inheritance of mtDNA: gender-specific Open Reading Frames (ORFs) and putative origins of replication. *Genetics* 183, 1575–1589 (2009).
- Breton, S. et al. Novel protein genes in animal mtDNA: A new sex determination system in freshwater mussels (Bivalvia: Unionoida)? Mol. Biol. Evol. 28, 1645–1659 (2011).
- Milani, L., Ghiselli, F., Guerra, D., Breton, S. & Passamonti, M. A comparative analysis of mitochondrial ORFans: New clues on their origin and role in species with Doubly Uniparental Inheritance of mitochondria. *Genome Biol. Evol.* 5, 1408–1434 (2013).
- Breton, S. *et al.* A resourceful genome: Updating the functional repertoire and evolutionary role of animal mitochondrial DNAs. *Trends Genet.* 30, 555–564 (2014).
- Breton, S., Beaupré, H. D., Stewart, D. T., Hoeh, W. R. & Blier, P. U. The unusual system of doubly uniparental inheritance of mtDNA: isn't one enough? *Trends Genet.* 23, 465–474 (2007).
- Passamonti, M. & Ghiselli, F. Doubly Uniparental Inheritance: Two mitochondrial genomes, one precious model for organelle DNA inheritance and evolution. DNA Cell Biol. 28, 79–89 (2009).
- 12. Zouros, E. Biparental inheritance through uniparental transmission: The Doubly Uniparental Inheritance (DUI) of mitochondrial DNA. *Evol. Biol.* **40**, 1–31 (2013).
- Breton, S. et al. The extremely divergent maternally-and paternally-transmitted mitochondrial genomes are co-expressed in somatic tissues of two freshwater mussel species with doubly uniparental inheritance of mtDNA. PLoS ONE 12 (2017).
- Doucet-Beaupré, H. *et al.* Mitochondrial phylogenomics of the Bivalvia (Mollusca): Searching for the origin and mitogenomic correlates of doubly uniparental inheritance of mtDNA. *BMC Evol. Biol.* 10, 1–19 (2010).
- Hoeh, W. R., Stewart, D. T., Sutherland, B. W. & Zouros, E. Cytochrome c oxidase sequence comparisons suggest an unusually high rate of mitochondrial DNA evolution in *Mytilus* (Mollusca: Bivalvia). *Mol. Biol. Evol.* 13, 418–21 (1996).
- Stewart, D. T., Kenchington, E. R., Singh, R. K. & Zouros, E. Degree of selective constraint as an explanation of the different rates of evolution of gender-specific mitochondrial DNA lineages in the mussel *Mytilus. Genetics* 143, 1349–57 (1996).
- Passamonti, M. & Scali, V. Gender-associated mitochondrial DNA heteroplasmy in the venerid clam *Tapes philippinarum* (Mollusca Bivalvia). Curr. Genet. 39, 117–124 (2001).
- Dalziel, A. C. & Stewart, D. T. Tissue-specific expression of male-transmitted mitochondrial DNA and its implications for rates of molecular evolution in *Mytilus* mussels (Bivalvia: Mytilidae). *Genome* 45, 348–355 (2002).
- 19. Bettinazzi, S., Plazzi, F. & Passamonti, M. The complete female- and male-transmitted mitochondrial genome of *Meretrix lamarckii*. *PLoS One* **11** (2016).
- Curole, J. P. & Kocher, T. D. Ancient sex-specific extension of the cytochrome c oxidase II gene in bivalves and the fidelity of doublyuniparental inheritance. *Mol. Biol. Evol.* 19, 1323–8 (2002).
- Chapman, E. G. et al. Extreme primary and secondary protein structure variability in the chimeric male-transmitted cytochrome c oxidase subunit II protein in freshwater mussels: Evidence for an elevated amino acid substitution rate in the face of domain-specific purifying selection. BMC Evol. Biol. 8, 165 (2008).
- 22. Guerra, D. et al. Evolution of sex-dependent mtDNA transmission in freshwater mussels (Bivalvia: Unionida). Sci. Rep. 7 (2017).
- Chakrabarti, R. et al. Presence of a unique male-specific extension of C-terminus to the cytochrome c oxidase subunit II protein coded by the male-transmitted mitochondrial genome of *Venustaconcha ellipsiformis* (Bivalvia: Unionoidea). FEBS Lett. 580, 862-866 (2006).
- 24. Chakrabarti, R. *et al.* Reproductive function for a C-terminus extended, male-transmitted cytochrome c oxidase subunit II protein expressed in both spermatozoa and eggs. *FEBS Lett.* **581**, 5213–5219 (2007).
- Passamonti, M., Ricci, A., Milani, L. & Ghiselli, F. Mitochondrial genomes and Doubly Uniparental Inheritance: New insights from Musculista senhousia sex-linked mitochondrial DNAs (Bivalvia Mytilidae). BMC Genomics 12, 442 (2011).
- Breton, S., Burger, G., Stewart, D. T. & Blier, P. U. Comparative analysis of gender-associated complete mitochondrial genomes in marine mussels (*Mytilus* spp.). *Genetics* 172, 1107–1119 (2006).
- Gusman, A., Lecomte, S., Stewart, D. T., Passamonti, M. & Breton, S. Pursuing the quest for better understanding the taxonomic distribution of the system of doubly uniparental inheritance of mtDNA. *PeerJ* 4, e2760 (2016).
- Soroka, M. & Burzyński, A. Complete female mitochondrial genome of Anodonta anatina (Mollusca: Unionidae): Confirmation of a novel protein-coding gene (F ORF). Mitochondrial DNA 26, 267–269 (2015).
- Soroka, M. & Burzyński, A. Complete male mitochondrial genome of *Anodonta anatina* (Mollusca: Unionidae). *Mitochondrial DNA* 27, 1679–1680 (2016).
- Burzyński, A., Soroka, M., Mioduchowska, M., Kaczmarczyk, A. & Sell, J. The complete maternal and paternal mitochondrial genomes of *Unio crassus*: Mitochondrial molecular clock and the overconfidence of molecular dating. *Mol. Phylogenet. Evol.* 107, 605–08 (2017).
- Burzyński, A. & Soroka, M. Complete paternally inherited mitogenomes of two freshwater mussels Unio pictorum and Sinanodonta woodiana (Bivalvia: Unionidae). PeerJ 6, e5573 (2018).
- Plazzi, F. & Passamonti, M. Footprints of unconventional mitochondrial inheritance in bivalve phylogeny: Signatures of positive selection on clades with doubly uniparental inheritance. J. Zool. Syst. Evol. Res. 57, 258–271 (2019).
- Chase, E. E., Robicheau, B. M., Veinot, S., Breton, S. & Stewart, D. T. The complete mitochondrial genome of the hermaphroditic freshwater mussel Anodonta cygnea (Bivalvia: Unionidae): In silico analyses of sex-specific ORFs across order Unionoida. BMC Genomics 19 (2018).
- Gomes-dos-Santos, A. et al. The male and female complete mitochondrial genomes of the threatened freshwater pearl mussel Margaritifera margaritifera (Linnaeus, 1758) (Bivalvia: Margaritiferidae). Mitochondrial DNA 4, 1417–1420 (2019).
- Soroka, M. & Burzynski, A. Doubly uniparental inheritance and highly divergent mitochondrial genomes of the freshwater mussel Unio tumidus (Bivalvia: Unionidae). Hydrobiologia 810, 239–254 (2018).
- Milani, L., Ghiselli, F., Maurizii, M. G., Nuzhdin, S. V. & Passamonti, M. Paternally transmitted mitochondria express a new gene of potential viral origin. *Genome Biol. Evol.* 6, 391–405 (2014).
- Minoiu, I., Burzyński, A. & Breton, S. Analysis of the coding potential of the ORF in the control region of the female-transmitted Mytilus mtDNA. Gene 576, 586–588 (2016).
- 38. Ouimet, P. et al. The ORF in the control region of the female-transmitted Mytilus mtDNA codes for a protein. Gene (In press).
- 39. Soroka, M. & Burzynski, A. Hermaphroditic freshwater mussel *Anodonta cygnea* does not have supranumerary open reading frames in the mitogenome. *Mitochondrial DNA* **2**, 862–864 (2017).
- Mitchell, A., Guerra, D., Stewart, D. T. & Breton, S. In silico analyses of mitochondrial ORFans in freshwater mussels (Bivalvia: Unionoida) provide a framework for future studies of their origin and function. BMC Genomics 17 (2016).
- Bieler, R., Carter, J. G. & Coan E. Classification of Bivalve Families. Pp. 113-133, in: Bouchet P & Rocroi J-P (2010), Nomenclator of Bivalve Families. *Malacologia* 52, 1–184 (2010).
- 42. Pante, E., Poitrimol, C., Saunier, A., Becquet, V. & Garcia, P. Putative sex-linked heteroplasmy in the tellinid bivalve *Limecola balthica* (Linnaeus, 1758). *J. Molluscan Stud.* 83, 226–228 (2017).
- Saunier, A. et al. Mitochondrial genomes of the Baltic clam Macoma balthica (Bivalvia: Tellinidae): Setting the stage for studying mito-nuclear incompatibilities. BMC Evol. Biol. 14 (2014).
- 44. Jameson, D., Gibson, A. P., Hudelot, C. & Higgs, P. G. OGRe: A relational database for comparative analysis of mitochondrial genomes. *Nucleic Acids Res.* **31**, 202–6 (2003).

- 45. Wolstenholme, D. R. Genetic novelties in mitochondrial genomes of multicellular animals. *Curr. Opin. Genet. Dev.* **2**, 918–925 (1992).
- 46. Grande, C., Templado, J. & Zardoya, R. Evolution of gastropod mitochondrial genome arrangements. BMC Evol. Biol. 8, 61 (2008).
- Yuan, Y., Li, Q., Yu, H. & Kong, L. The complete mitochondrial genomes of six heterodont bivalves (Tellinoidea and Solenoidea): Variable gene arrangements and phylogenetic implications. *PLoS One* 7 (2012).
- Ojala, D., Montoya, J. & Attardi, G. tRNA punctuation model of RNA processing in human mitochondria. *Nature* 290, 470–474 (1981).
- Milbury, C. A. & Gaffney, P. M. Complete mitochondrial DNA sequence of the eastern oyster *Crassostrea virginica*. Mar. Biotechnol. 7, 697–712 (2005).
- Xu, K., Kanno, M., Yu, H., Li, Q. & Kijima, A. Complete mitochondrial DNA sequence and phylogenetic analysis of Zhikong scallop Chlamys farreri (Bivalvia: Pectinidae). Mol. Biol. Rep. 38, 3067–3074 (2011).
- Sun, S., Kong, L., Yu, H. & Li, Q. The complete mitochondrial DNA of *Tegillarca granosa* and comparative mitogenomic analyses of three Arcidae species. *Gene* 557, 61–70 (2015).
- 52. Kumar, S., Stecher, G. & Tamura, K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* 33, 1870–1874 (2016).
- Robicheau, B. M., Breton, S. & Stewart, D. T. Sequence motifs associated with paternal transmission of mitochondrial DNA in the horse mussel, *Modiolus modiolus* (Bivalvia: Mytilidae). *Gene* 605, 32–42 (2017).
- 54. Zouros, E. The exceptional mitochondrial DNA system of the mussel family Mytilidae. Genes Genet. Syst. 75, 313-318 (2000).
- 55. Altschul, S. F. et al. Blast and Psi-Blast: Protein Database Search Programs. Nucleid Acid Res. 25, 2289-4402 (1997).
- Szafranski, P. Evolutionarily recent, insertional fission of mitochondrial cox2 into complementary genes in bilaterian Metazoa. BMC Genomics 18, 269 (2017).
- 57. Vallès, Y., Halanych, K. M. & Boore, J. L. Group II introns break new boundaries: Presence in a bilaterian's genome. PLoS One 3 (2008).
- Beier, H. & Grimm, M. Misreading of termination codons in eukaryotes by natural nonsense suppressor tRNAs. *Nucleic Acids Res.* 29, 4767–82 (2001).
- Ling, J., O'Donoghue, P. & Söll, D. Genetic code flexibility in microorganisms: Novel mechanisms and impact on physiology. *Nat. Rev. Microbiol.* 13, 707–721 (2015).
- Noutahi, E., Calderon, V., Blanchette, M., El-Mabrouk, N. & Lang, B. F. Rapid genetic code evolution in green algal mitochondrial genomes. *Mol. Biol. Evol.* 36, 766–783 (2019).
- Rampersad, S. & Tennant, P. Replication and expression strategies of viruses. In: Viruses: Molecular biology, host interactions, and applications to Biotechnology. Elsevier. p. 55–82 (2018).
- 62. Oudot-Le Secq, M.-P. & Green, B. R. Complex repeat structures and novel features in the mitochondrial genomes of the diatoms *Phaeodactylum tricornutum* and *Thalassiosira pseudonana*. *Gene* **476**, 20–6 (2011).
- Beckenbach, A. T., Robson, S. K. A. & Crozier, R. H. Single nucleotide +1 frameshifts in an apparently functional mitochondrial cytochrome b gene in ants of the genus *Polyrhachis, J. Mol. Evol.* 60, 141–52 (2005).
- Russell, R. D. & Beckenbach, A. T. Recoding of translation in turtle mitochondrial genomes: Programmed frameshift mutations and evidence of a modified genetic code. J. Mol. Evol. 67, 682–95 (2008).
- 65. Temperley, R. *et al.* Hungry codons promote frameshifting in human mitochondrial ribosome. *Science* **327**, 301 (2010).
- Bettinazzi, S., Rodríguez, E., Milani, L., Blier, P. U. & Breton, S. Metabolic remodelling associated with mtDNA: Insights into the adaptive value of doubly uniparental inheritance of mitochondria. Proc. R. Soc. B Biol. Sci. 286 (2019).
- Rosel, P. E. & Block, B. A. Mitochondrial control region variability and global population structure in the swordfish, *Xiphias gladius*. Mar. Biol. 125, 11–22 (1996).
- 68. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120 (2014).
- 69. Zhang, J., Kobert, K., Flouri, T. & Stamatakis, A. PEAR: A fast and accurate Illumina paired-end read merger. *Bioinformatics* **30**, 614–620 (2014).
- Afgan, E. et al. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res. 46, W537–W544 (2018).
- Andrews, S. FastQC: a quality control tool for high throughput sequence data. *Babraham Bioinform*. Available from, http://www. bioinformatics.babraham.ac.uk/projects/fastqc/ (2010)
- Hahn, C., Bachmann, L. & Chevreux, B. Reconstructing mitochondrial genomes directly from genomic next-generation sequencing reads – a baiting and iterative mapping approach. Nucleic Acids Res. 41, e129 (2013).
- Pomerantz, A. et al. Real-time DNA barcoding in a rainforest using nanopore sequencing: Opportunities for rapid biodiversity assessments and local capacity building. Gigascience 7 (2018).
- Lanfear, R., Schalamun, M., Kainer, D., Wang, W. & Schwessinger, B. MinIONQC: Fast and simple quality control for MinION sequencing data. *Bioinformatics* 35, 523–525 (2019).
- De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C. NanoPack: Visualizing and processing long-read sequencing data. *Bioinformatics* 34, 2666–2669 (2018).
- Koren, S. et al. Canu: Scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res. 27, 722–736 (2017).
- Loman, N. J., Quick, J. & Simpson, J. T. A complete bacterial genome assembled de novo using only nanopore sequencing data. *Nat. Methods* 12, 733–735 (2015).
- Viricel, A., Becquet, V., Dubillot, E. & Pante, E. De novo assembly and functional annotation of the transcriptome of *Mimachlamys varia*, a bioindicator marine bivalve. *Mar. Genomics* 41, 42–45 (2018).
- Walker, B. J. et al. Pilon: An integrated tool for comprehensive microbial variant detection and genome assembly improvement. PLoS One 9 (2014).
- Vagner, M. et al. Ocean warming combined with lower omega-3 nutritional availability impairs the cardio-respiratory function of a marine fish. J Exp. Biol. 222 (2019).
- 81. Kurtz, S. et al. Versatile and open software for comparing large genomes. Genome Biol. 5, R12 (2004).
- 82. Bernt, M. et al. MITOS: Improved de novo metazoan mitochondrial genome annotation. Mol. Phylogenet. Evol. 69, 313-319 (2013).
- Lowe, T. M. & Eddy, S. R. TRNAscan-SE: A program for improved detection of transfer rna genes in genomic sequence. *Nucleic Acids Res.* 25, 955–964 (1997).
  C. G. Wilfe, M. L. Communication of a distribution of a distribution of the second sec
- Conant, G. C. & Wolfe, K. H. GenomeVx: simple web-based creation of editable circular chromosome maps. *Bioinformatics* 24, 861–862 (2008).
- 85. Madeira, F. et al. The EMBL-EBI search and sequence analysis tools APIs in 2019. Nucleic Acids Res. 47, W636–W641 (2019).
- Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–7 (2004).
  Suyama, M., Torrents, D. & Bork, P. PAL2NAL: Robust conversion of protein sequence alignments into the corresponding codon alignments. *Nucleic Acids Res.* 34, W609–12 (2006).
- Nei, M. & Gojobori, T. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* 3, 418–26 (1986).
- 89. Wickham, H. Ggplot2 Elegant Graphics for Data Analysis (2009).
- Brown, N. P., Leroy, C. & Sander, C. MView: A web-compatible database search or multiple alignment viewer. *Bioinformatics* 14, 380–381 (1998).

#### Acknowledgements

EP thanks the Molecular Core Facility at the University of La Rochelle and the bioinformatics platform Toulouse Midi-Pyrenees (Bioinfo GenoToul). This work was funded by the French Agence Nationale de la Recherche (HySea project ANR-12-BSV7-0011; DRIVE project ANR-18-CE02-0004-01), La Rochelle Université, the CPER/ FEDER (ECONAT project), and by funding from the Natural Sciences and Engineering Research Council of Canada (grant no. RGPIN/435656-2013 to SB).

#### **Author contributions**

C.C., E.P. and S.B. conceived and designed the experiments. C.C. analyzed data and drafted the manuscript. K.B., D.G., B.M.R. and D.T.S. participated in the experimental and analytical work. All authors contributed feedback to the writing process and approved the manuscript.

#### **Competing interests**

The authors declare no competing interests.

#### Additional information

Supplementary information is available for this paper at https://doi.org/10.1038/s41598-020-57975-y.

Correspondence and requests for materials should be addressed to C.C. or S.B.

Reprints and permissions information is available at www.nature.com/reprints.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2020