



HAL
open science

Reviewing *Fasciola hepatica* transmission in the West Indies and novel perceptions from experimental infections of sympatric vs. allopatric snail/fluke combinations

Antonio Vázquez, Mercedes de Vargas, Annia Alba, Jorge Sanchez, Pilar Alda, Emeline Sabourin, Marion Vittecoq, Pedro M. Alarcón-Elbal, Jean-Pierre Pointier, Sylvie Hurtrez-Boussès

► **To cite this version:**

Antonio Vázquez, Mercedes de Vargas, Annia Alba, Jorge Sanchez, Pilar Alda, et al.. Reviewing *Fasciola hepatica* transmission in the West Indies and novel perceptions from experimental infections of sympatric vs. allopatric snail/fluke combinations. *Veterinary Parasitology*, 2019, 275, pp.108955. 10.1016/j.vetpar.2019.108955 . hal-02504268

HAL Id: hal-02504268

<https://hal.science/hal-02504268>

Submitted on 10 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Reviewing *Fasciola hepatica* transmission in the West Indies and novel perceptions from experimental infections of sympatric vs. allopatric snail/fluke combinations

Antonio A. Vázquez, Mercedes de Vargas, Annia Alba, Jorge Sánchez, Pilar Alda, Emeline Sabourin, Marion Vittecoq, Pedro M. Alarcón -Elbal, Jean-Pierre Pointier, Sylvie Hurtrez-Boussès

How to cite : Vázquez A.A., de Vargas M., Alba A., Sánchez J., Alda P., Sabourin E., Vittecoq M., Alarcón-Elbal P.M., Pointier J.-P., Hurtrez-Boussès S. 2019. Reviewing *Fasciola hepatica* transmission in the West Indies and novel perceptions from experimental infections of sympatric vs. allopatric snail/fluke combinations. *Veterinary Parasitology* 275:108955. doi: 10.1016/j.vetpar.2019.108955

Abstract

Fasciolosis is an important food-borne parasitic disease affecting over two million people worldwide with economic losses related to cattle production of up to US\$ 3 billion annually. Despite the long known presence of *Fasciola hepatica* in the Caribbean islands its transmission is not well known. This study reviews historical and recent data on fasciolosis in the West Indies, revealing for the first time the outcomes of sympatric and allopatric fluke/snail interactions in the area by exploring the susceptibility of four lymnaeid species after exposure to *F. hepatica* isolates from Cuba, the Dominican Republic and France. Overall, *Galba cubensis* showed a mean prevalence of 71.8% and appears to be the most suitable intermediate host species irrespective of the isolate used. Sympatric combinations (snail and parasite from the same country) were generally more compatible (higher susceptibility, parasite intensity and snail survival post-exposure) and only the allopatric interaction of French *G. truncatula*/Cuban *F. hepatica* attained 100% prevalence and mean intensity over 33 rediae/snail. However, certain Dominican populations of *Pseudosuccinea columella* showed high parasite intensities (>30 rediae/snail) when infected with Cuban flukes, highlighting the potential risks of biological introductions. Overall, high compatibility in most sympatric combinations compared to low or moderate compatibility in allopatric ones, suggests the existence of local adaptation from a long sustained interaction that has led to high rates of transmission. Interestingly, attempts to infect *G. schirazensis* with sympatric and allopatric flukes failed and coupled with the lowest survival rates which supposes a low risk of fasciolosis transmission in areas where this is the only snail species. Although there are significant gaps in the actual status of fasciolosis transmission from several islands in the West Indies these results show a permanent risk. We conclude that fasciolosis transmission is high in areas where the local snail, *G. cubensis*, occurs, and will be even higher in the presence of the invasive *P. columella*.



Graphical Abstract

Keywords :Fasciola hepatica, Lymnaeidae, , Caribbean, Parasite transmission, Experimental infections

1. Introduction

Fasciolosis is a food-borne disease caused by the parasites *Fasciola hepatica* and *F. gigantica* and transmitted by freshwater snails of the family Lymnaeidae (Andrews, 1999). Fasciolosis affects nearly 17 million people worldwide, with up to 180 million at risk of infection (Mas-Coma et al., 2018). The public health risk keeps increasing, linked to the growing problem of fasciolosis in livestock farming and an increase in the frequency of infected animals that are mainly used for meat and milk production (Khan et al., 2013). Worldwide, estimated economic losses of animal fasciolosis reach nearly US \$3 billion annually with over 600 million animals infected (Toet et al. 2014).

Liver fluke infection by *F. hepatica* is considered to be one of the most widespread trematodes (Mas-Coma et al., 2009), due in part, to the fact that around 30 lymnaeid snail species worldwide are recognized as intermediate hosts (Vázquez et al., 2018). Its introduction into the Western Hemisphere is related to the European colonization of the Americas (1500-1800) by means of non-native infected cattle brought for settlement (Mas-Coma et al., 2009). With a supposed Eurasian origin (Lotfy et al., 2008), *F. hepatica* has found suitable hosts in the local American lymnaeid species of the genera *Galba* and *Pseudosuccinea* (Vázquez et al., 2018).

Within the Americas, the study of fasciolosis transmission in the West Indies (islands of the Caribbean Sea) is particularly interesting given its marked insularity. Although usually not separated by more than a few hundred kilometres, each island probably has its own transmission dynamics related to the mollusc species involved, the particularities of their freshwater ecosystems and the type of cattle husbandry in use (Fig. 1). Only three lymnaeid species —*G. cubensis*, *G. schirazensis* and *P. columella*— are reported in the West Indies and each has been suggested to play different roles in the parasite transmission (Bargues et al., 2011; Gutiérrez et al., 2011; Vázquez et al., 2014).

Even if transmission is not well documented in most of the West Indies, particularly in the Lesser Antilles, the burden of *F. hepatica* infection in either humans or animals (domestic or wild) is expected to be as high as in neighbouring islands with similar epidemiological characteristics (Table 1). In such an insular model, intermediate hosts may reach very high densities caused by low interspecific competition due to the low numbers of sibling species and, in consequence, the chances of becoming infected are high and simplify the arrival of the parasite to the next host. Therefore, this region is highly suitable for fasciolosis transmission and, given its climate, is ideal for an uninterrupted life-cycle of the parasite (Mas-Coma et al., 2018).

However, within the West Indies, only countries such as Cuba, Haiti, Dominican Republic, Puerto Rico, Jamaica and St. Lucia have reports of infections in definitive hosts (Table 1). Cuba is the only country where significant studies have been carried out, and nine outbreaks have been informed in the human population. According to a 1990's study from the WHO, 2594 human cases were acknowledged worldwide particularly from Europe and Asia but also from Cuba and Puerto Rico (Chen and Mott, 1990). In other countries such as Haiti, there are reports of 60% infected cattle (Gentilini et al., 1964) with a supposed high prevalence in the human population (Rojas et al. 2010). In the case of Lesser Antilles reports are only available for animal fasciolosis in St. Lucia and Martinique (Table 1). In the latter, infected cattle imported from Lorraine (France) were observed by the slaughtering services at Fort-de-France (Grétilat, 1966) but autochthonous transmission has not been assessed.

Regarding the intermediate host, freshwater snails of the family Lymnaeidae occur in the vast majority of the Caribbean (Hubendick, 1951). *Galba cubensis* and *P. columella* have been reported in most of the West Indies (Table 1), but so far, *G. schirazensis* has only been reported in the Dominican Republic (Bargues et al., 2011). However, parasites on the snail hosts *Galba cubensis* and *P. columella*, have only been reported in Cuba (Table 1).

Due to the relatively short distances separating each island and the existence of natural biological corridors (e.g. wading birds' migration pathways), the particular transmission scenario of one island may affect transmission on others since snails can easily be introduced (Wesselingh et al., 1999; Cowie and Holland, 2006). Given that snails and/or parasites may migrate (by means of natural or human mediated introduction), more susceptible/infective genotypes could be introduced in the future shaping transmission patterns. Invasive genotypes of the lymnaeid host

P. columella introduced worldwide are supposed to have facilitated the spread of *F. hepatica* in many regions due to a lack of genetic diversity that may counteract infection (Lounnas et al., 2017). In Cuba, for example, the same effect has been observed and the only infected natural populations of *P. columella* carried a widespread genotype particularly present in monomorphic populations (Alba et al. 2019).

The study of compatibility between different parasite and snail populations may serve to better understand and predict transmission. To this end, experimental infections are key to explore traits in host-parasite interactions. It is particularly interesting to include *G. schirazensis*, as to date there is no consensus of its actual role in transmission (Bargues et al., 2011; Dreyfuss et al., 2015; Caron et al., 2017).

This study aims to test the compatibility of several lymnaeid snail populations with sympatric and allopatric fluke isolates. To this end, several *F. hepatica* isolates and snail species from Cuba (two lymnaeid species coexist) and from the Dominican Republic (three lymnaeid species coexist) served as biological models. Also, given the reports of infected cattle in Martinique brought from France (Grétilat, 1967), an isolate of *F. hepatica* and a fourth lymnaeid snail (*G. truncatula*) from France were used to test compatibility over larger allopatric scales. Regarding *G. truncatula*, this species has not been reported in the West Indies but is present in South America (Jabbour-Zahab et al., 1997) as well as in many other world regions (Kock et al., 2003; Rondelaud et al., 2006; Jones et al., 2015) and is considered the main intermediate host of *F. hepatica* in Europe (Correa et al., 2017). These results will give insights into the actual players in the host-parasite interaction in the West Indies by exploring snail-parasite combinations from its largest islands.

2. Material and Methods

2.1. Sampling of snail populations and liver fluke isolates

We sampled several field populations (Fig. 2) of *G. cubensis* (5), *G. schirazensis* (2), *G. truncatula* (1) and *P. columella* (6) from Cuba, the Dominican Republic and France (Table 2). The snails were directly removed from the mud or aquatic vegetation using soft forceps and sieves (up to 100 individuals per locality when possible). The snails were brought to the laboratory and kept isolated in Petri dishes under controlled conditions (i.e. temperature ranging 25–26°C, fed ad libitum with algae; Sánchez et al., 1995) until oviposition to obtain lab-reared colonies for experimental infection trials with *F. hepatica* isolates.

The isolates of *F. hepatica* were retrieved directly from the liver of recently slaughtered cattle in Cuba (Pinar del Río province), the Dominican Republic (Dajabón province) and France (Tour du Valat, Camargue) (Fig. 2). All adult *F. hepatica* samples were kindly donated by the veterinary authorities of each slaughterhouse after routine activities. Adult flukes were placed in a saline solution (0.85%) with glucose (9%) to promote egg laying. Eggs were later stored in tubes containing saline solution (0.85%) at 4 °C in the dark until use (Vázquez et al., 2014).

2.2. Morphological and molecular identification of snail species

Snails were identified using morphological and molecular markers. *Pseudosuccinea columella* can be distinguished without ambiguity from *Galba* species due to its shell morphology and internal anatomy (i.e. shell with a short spire; periostracum ornamented with spiral sculptures; ureter with two distant flexures; Pointier, 2008). In the case of *G. cubensis*, *G. schirazensis*, and *G. truncatula*, these species are very similar in shell morphology and anatomical variation which render them undistinguishable, and can be considered as cryptic species (Correa et al., 2011).

In order to identify these three *Galba* species, up to 30 individuals from each *Galba* field population were tested using a molecular tool designed to this end by Alda et al. (2018). Previously obtained egg clutches of field-collected snails were used for founding colonies after confirmation of the species using multiplex PCR. DNA was extracted using a Chelex protocol following Estoup and Martin (1996) adapted to 96-well plates. The multiplex PCR (Alda et al. 2018) was based on species-specific primers amplifying microsatellite loci targeting cryptic species and producing band sizes that are specific to the three targeted species (179–200 pb in *G. cubensis*, 227–232 pb in *G. schirazensis*, and 111–129 pb in *G. truncatula*).

2.3. Infection trials

Two-week old snails obtained after two successive lab-reared generations were infected with hatched *F. hepatica* miracidia. Eggs of *F. hepatica* were placed in spring water and incubated for a fortnight, at 27 °C, in darkness. Direct light exposure served to induce miracidial hatching. A dose of five miracidia/snail was used to expose 30 individuals from each population following the methodology described in Vázquez et al. (2014). Briefly, lymnaeid snails were placed in a 96 well plate containing the established dose of free-swimming miracidia. Snails were forced to remain submerged by filling the wells with spring water and placing glass slides on top. After three hours of exposure, each snail was returned to their rearing Petri dishes previous confirmation that no miracidia remained in the well. After 25 days post-exposure all snails were dissected to verify their infection status. Dissection was carried out under a stereoscope where each snail was carefully crushed and separated from its shell using forceps (Caron et al. 2008). Only living *F. hepatica* rediae (irrespective of their size or developmental stage) were used to measure parasite intensity. We kept 30 unexposed individuals (experimental controls) of each species/population subjected to the same conditions and procedures. Mortality was recorded daily and the infection status (infected/non-infected) was noted in all dead individuals that died after the first week post-exposure by observing *F. hepatica* rediae. Individuals dying during the first week post-exposure usually have no traceable clues of infection because early developmental stages are difficult to observe. Therefore, in order to prevent introducing uncontrollable bias they were considered non-infected in our experiment. We explored different infection outcomes in relation to sympatric (same country) and allopatric (different country) combinations within the region.

2.4. Data analysis

All gathered data was used to assess compatibility in each snail/fluke combination. Prevalence (proportion of infected individuals within the host sample) and mean parasite intensity (average of intensity values –rediae/snail– calculated for a sample) with 95% confidence intervals (CI) was recorded for each trial according to Reiczigel et al. (2019). Spearman rank correlations were performed to test associations between parasite prevalence and intensities. Parasite intensities were compared among trials by means of one-way ANOVA after normality and homogeneity of variance were verified by the Shapiro-Wilk and Levene tests. A post hoc Tukey test was performed to obtain significant differences for each particular combination. Statistical differences of survivorship in each trial was assessed through a log-rank test of Kaplan-Meier curves built from day 0 to day 25 post-exposure. All statistical tests were performed in Statistica v.12 (StatSoft. Inc., Tulsa, OK, USA 2014) and all differences were considered statistically significant at values of $P < 0.05$.

3. Results

With the exception of *G. schirazensis* that was not-infected (no observation of any larval stages) with any of the *F. hepatica* isolates tested (data not shown), the other three snail species showed a variable degree of susceptibility with prevalence ranging from 0 to 100% (Fig. 3). Overall, *G. cubensis* showed a higher prevalence (mean 72%, range 27–93%) than *P. columella* (48%, 0–70%) irrespective of the isolate used. Other than *G. schirazensis*, only *P. columella* from El Chico (Cuba) failed to develop infection with the Dominican isolate (allopatric combination, data not shown) but was always infected with its sympatric Cuban flukes (Fig. 3). The French population of *G. truncatula* showed a 100% infection prevalence with the Cuban *F. hepatica* (larger allopatric combination) with a mean intensity of 33.5 (95% CI; 27.5–39.6) rediae per snail.

Parasite intensities were also variable and significant differences were found depending on the snail/fluke combination (Fig. 3). Overall, higher parasite intensity was found in sympatric combinations in *G. cubensis*, but some allopatric *P. columella*/*F. hepatica* combinations were also high (e.g. DR.05/Cuba, mean 33.1, 95% CI; 25–41.1). Parasite intensities of 15–20 rediae/snail were observed during infection with the French isolate of *F. hepatica* but were lower than sympatric combinations. In fact, allopatric combinations such as those of *P. columella* (DR.06/Cuba, mean 7.5, 95% CI; 4.8–10.7 and CU.04/DomRep., mean 8.2, 95% CI; 5.2–11.9) and *G. cubensis* (DR.01/Cuba, mean 14.7, 95%CI; 10.8– 19.3 and CU.02/Dom. Rep., mean 14.4, 95% CI; 11–18) showed lower parasite burden. We found no correlation between parasite prevalence and mean intensity in *G. cubensis* ($r = 0.49$, $P = 0.176$) but a positive significant correlation for *P. columella* ($r = 0.86$, $P = 0.003$).

Survival of snails after 25 days post-exposure varied in relation to the lymnaeid species and the isolate of *F. hepatica*, but the most frequent peaks of mortality were observed within the first seven days post-exposure (Fig. 4). Overall, *G. truncatula* (survival 100%) and *G. cubensis* (survival ranged from 73–100%) showed the highest survival rate following infection. Notably, *G. schirazensis* showed a very low survival rate during the 25 days post-exposure (survival ranged from 3.3–13.3%) particularly when exposed to the Cuban isolate (allopatric combination). In this case, this species showed a sustained mortality throughout the experiment even when no infection was observed in dead snails. Overall, lower survival rates were attained with the allopatric French isolate in *G. cubensis* and *P. columella*. Control groups (unexposed individuals, $n = 30$) of any species/population showed zero mortality during the time of the experiment (data not shown). Also, no correlation between the effects of snail survival and parasite prevalence ($r = -0.12$, $P = 0.3$) or intensity ($r = 0.2$, $P = 0.18$) was observed.

4. Discussion

The results of testing different snail/fluke combinations revealed a range of competence for the different snail species, from high to very low suitability as hosts in terms of prevalence, parasite burden and snail survival. Overall, *G. cubensis* and *G. truncatula* were more compatible than *P. columella*, while *G. schirazensis* did not show any sign of compatibility (at least the populations tested). In particular, these results endorse *G. cubensis* as playing the major role of parasite transmission in the region. This is the case in Cuba where *G. cubensis* shows high compatibility with *F. hepatica* both in the laboratory (Vázquez et al., 2014) and in the field (Vázquez et al., 2015; Alba et al., 2016) compared to *P. columella* (Gutiérrez et al. 2011). According to Hubendick (1951), *G. cubensis* has acircum-Caribbean distribution, occurring in most of the West Indies. Moreover, the present study indicates the potential risks of transmission via *G. truncatula* if this species is introduced into the West Indies. This risk is not negligible as this lymnaeid not only showed high compatibility with a local *F. hepatica* isolate (Cuba) but is known to be invasive elsewhere (Meunier et al., 2001).

The understanding of transmission patterns in these countries could be vital to comprehend the overall scenario in the West Indies. According to Mas-Coma et al. (2018), this area is characterized by the ‘Caribbean insular pattern’ characterized by small and frequent outbreaks in human hypoendemic areas. This is consistent with the scenario observed in Cuba where fasciolosis is not yet considered a severe public health problem but outbreaks occur (Rojas et al., 2010). On the contrary, the country is characterized by a hyper endemic situation at the veterinary level with prevalence ranging from 70-100% in cattle (Vázquez et al. 2016). The assessment of the different roles of each intermediate host snail species in the region is important to characterize the epidemiological context and to anticipate it under varying conditions. For this, experimental infections are a valuable approach to investigate differences in compatibility at a phenotypical level since they allow all variables to be managed and controlled (e.g. the combination of snails and parasites tested, the infective parasite dose, etc.), all of which could significantly influence the infection outcome (Sorensen and Minchella, 2001).

Little is known about the immunobiology and the particular mechanism involved in the *F. hepatica* – snail interaction, but variation in infection outcomes, depending on the parasite isolate/snail population combination, are reported here and elsewhere (Rondelaud et al., 2013; Vázquez et al., 2014; Alba et al., 2018). In the widely-studied model of *Biomphalaria glabrata* – *Schistosoma mansoni*, an infectivity mosaic that follows the model of ‘matching alleles’ is also observed (Théron and Coustau, 2005) and is identified as a polymorphism of compatibility. Overall, the *F. hepatica*-lymnaeid snail interplay is apparently characterized by different compatibility that will ultimately depend on the particular host-parasite (genotype) combination (Vázquez et al., 2014; Alba et al., 2018). From the results observed here, the latter seems to be associated with local adaptation as there was high compatibility in most sympatric and low-to-moderate compatibility in most allopatric combinations for *G. cubensis* and *P. columella*. In this sense, it can be hypothesized that local adaptation has rendered sympatric combinations more compatible. The concept of local adaptation is highlighted when we tested the French *F. hepatica* with West Indian snails. This isolate appeared as ‘highly virulent’ and drastically decreased the survival of snails. However, certain allopatric combinations in *P. columella* also showed higher values of prevalence, intensity and survival compared to sympatric combinations. From these findings one could speculate that a random encounter of highly compatible snail/fluke genotypes in this species may trigger the probability of infection if other conditions are met (e.g. ecological, epidemiological, anthropological, etc.).

The case of *G. schirazensis* is particularly interesting given the current results. This species failed to develop the parasite regardless of the *F. hepatica* isolate used. Similarly, a previous study of a one-time series of experimental infections using laboratory reared *F. hepatica* (from which genetic diversity is expected to be reduced) also reported non-infection results for this species when populations derived from Egypt and Spain were challenged (Bargues et al., 2011). Based on this result, Bargues et al. (2011) labelled *G. schirazensis* as a species with no role in the transmission of fasciolosis (refractory to *F. hepatica*). However, while one may agree that the overall infection risk of this snail is considerably lower in relation to other lymnaeid species (*G. cubensis* or *P. columella*), this species should be proposed as highly incompatible to *F. hepatica* (at least the combination tested). The results from the survival particularly if the parasite strain is highly virulent for the host, could explain the overall impairment of survival observed in the allopatric combination.

To understand this problem, a comparison with a different infection related phenotype of resistant snails in this system should be carried out. Some Cuban populations of *P. columella*, that are considered to be resistant to *F. hepatica* infection (Gutiérrez et al., 2003; Alba et al., 2018), show an unusually higher survival following exposure (>70%, Gutiérrez et al., 2002; >80%, Alba et al., 2018) in relation to susceptible individuals. Also, the long-term maintenance in laboratory conditions has no effect on the stability of the phenotype. In these populations, resistance is associated to an effective immune encapsulation of the parasite (Gutiérrez et al., 2003) and it is not restricted to a host genotype – parasite genotype interaction (Gutiérrez et al., 2003; Alba et al., 2018). In fact, no reversion of the phenotype has occurred after different experimental challenges, even with geographically distant parasites (Vázquez et al., 2014; Alba et al., 2018).

In contrast, *G. schirazensis* has been found infected in the field (6% prevalence) in Ecuador (Caron et al., 2017) and Bargues et al. (2011) found parasitic development in 2 out of 10 exposed snails when challenged with a Polish *F. hepatica* isolate. Also, populations of *G. schirazensis* were able to be infected and develop *F. hepatica* rediae and free cercariae after several laboratory reared generations and experimental exposures, pointing to an eventual increase of its compatibility in relation to the selective pressure of particular parasite genotypes (Dreyfuss et al., 2015). Thus, the role of this species as a potential intermediate host of *F. hepatica* should not be totally disregarded, particularly if local adaptation occurs. For instance, *G. schirazensis* has been the only lymnaeid species found after a series of samplings in La Trampa (Venezuela) where *F. hepatica* is prevalent in the bovine population maintained under anthelmintic treatment every six months (Unpublished results).

Further studies are needed to better describe what occurs in *G. schirazensis* after parasitic exposure and which particular conditions/factors (snail and parasite diversity, densities, varying compatibility according to matching genotypes, etc.) might facilitate transmission in the field. Moreover, it should also be considered that the presence of *G. schirazensis* in curves observed here (<20%, this study) and elsewhere (<35%, Dreyfuss et al., 2015) provided some insight on this issue as higher survival should be expected after parasitic challenge in a refractory host (Ibrahim and Trpis, 1987; Abdel-Hamid et al., 2006). The high mortality observed in exposed *G. schirazensis* might be related to a “hypersensitivity” phenomenon occurring in the host in association with the parasitic infection that results in an unsuitable environment for *F. hepatica* development. A highly significant hyper pathological/physiological reaction, the Americas has been largely overlooked and is only relatively recently acknowledged (Correa et al., 2010; Bargues et al., 2011). This species may co-occur with other Galba species (in the present study we found *G. schirazensis* co-occurring with *G. cubensis* in Auyamas, Dominican Republic) and since they cannot be accurately differentiated at the morphological level, assuming an intermediate host role by either of the cryptic Galba, and particularly by *G. schirazensis* could be difficult without molecular discrimination (Alba et al., 2018).

Finally, while in the West Indies *F. hepatica* is probably highly prevalent, more research is needed to give insights into the particular scenarios of islands with unavailable data. Concerning the intermediate hosts, we can predict that the risk of transmission is higher in areas where the local snail *G. cubensis* occurs, and even higher if the invasive *P. columella* co-occurs. Low to very low probability is expected in areas with only *G. schirazensis* unless compatibility increases through local adaptation with the circulating parasite strain.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Competing Interests None

Acknowledgments

We would like to thank Iannis Hurtrez, Mathieu Baudoin and Thomas Gandou for facilitating the transport of samples for experimental infection and molecular identification. We also thank all the students of INTEC in the Dominican Republic, which helped us in the field. We are in debt to veterinary authorities and technical personnel from abattoirs in Cuba (Candelaria), Dominican Republic (Dajabón) and France (Tarascon) for their help in obtaining fluke samples. We are most thankful to the anonymous reviewers for their valuable suggestions and the editor for all the comments that greatly improved the final version of the manuscript. We also thank Dr. Suzanne Mills for smoothing the English writing. These studies have been partially supported by the Institute of Tropical Medicine “Pedro Kourí” (IPK-MINSAP, Cuba), FONDOCYT-MESCyT (Dominican Republic), Ecos-Sud program A16B02, Labex CeMeb, Muse (project REPAS: grant n°170544IA, n°ANR-IDEX-0006), Tour du Valat, IRD, CNRS and University of Montpellier.

References

- Abdel-Hamid, Z., Rawi, S., Arafa, A., 2006. Identification of a genetic marker associated with the resistance to *Schistosoma mansoni* infection using random amplified polymorphic DNA analysis. *Mém. Inst. Oswaldo Cruz* 101, 863–868.
- Agnamey, P., Fortes-Lopes, E., Raccurt, C., Boncy, J., Tote, A., 2011. Cross-sectional serological survey of human fascioliasis in Haiti. *J. Parasitol. Res.* 2012, 751951, doi:10.1155/2012/751951.
- Aguayo, C., 1938. Los moluscos fluviátiles Cubanos. Parte II. – Sistemática. *Mem. Soc. Cub. Hist. Nat.* 12, 253–276.
- Alba, A., Vázquez, A., Sánchez, J., Duval, D., Hernández, H., Sabourin, E., Vittecoq, M., Hurtrez-Boussès, S., Gourbal, B., 2018. *Fasciola hepatica*-*Pseudosuccinea columella* interaction: effect of increasing parasite doses, successive exposures and geographical origin on the infection outcome of susceptible and naturally resistant snails from Cuba. *Parasite & Vectors* 11, 559, doi:10.1186/s13071-018-3155-3.
- Alba, A., Vázquez, A., Sánchez, J., Fraga, J., Hernández, H., Martínez, E., Marcet, R., Figueredo, M., Sarracent, J., 2016. Assessment of the FasciMol-ELISA in the detection of the trematode *Fasciola hepatica* in field-collected *Galba cubensis*: a novel tool for the malacological survey of fasciolosis transmission. *Parasites & Vectors* 9, 22, doi:10.1186/s13071-016-1303-1.
- Alba, A., Vázquez, A., Sánchez, J., Lounnas, M., Pointier, J.-P., Hurtrez-Boussès, S., Gourbal, B., 2019. Patterns of distribution, population genetics and ecological requirements of field-occurring resistant and susceptible *Pseudosuccinea columella* snails to *Fasciola hepatica* in Cuba. *Sci. Rep.* doi: 10.1038/s41598-019-50894-7.
- Alda, P., Lounnas, M., Vázquez, A., Ayaqui, R., Calvopiña, M., Celi-Erazo, M., Dillon, R., Jarne, P., Loker, E., Muñiz, F., Muzzio-Aroca, J., Narváez, A., Noya, O., Martini, L., Rodríguez-Hidalgo, R., Uribe, N., David, P., Pointier, J.-P., Hurtrez-Boussès, S., 2018. A new multiplex PCR assay to distinguish among three cryptic *Galba* species, intermediate hosts of *Fasciola hepatica*. *Vet. Parasitol.* 251, 101–105.
- Andrews, S., 1999. The life cycle of *Fasciola hepatica*, in: Dalton, J. (Ed.), *Fasciolosis*. CABI Publishing, Wallingford, pp. 1–29.
- Bargues, M., Artigas, P., Khoubbane, M., Flores, R., Glöer, P., Rojas-García, R., Ashrafi, K., Falkner, G., Mas-Coma, S., 2011. *Lymnaea schirazensis*, an overlooked snail distorting fascioliasis data: genotype, phenotype, ecology, worldwide spread, susceptibility, applicability. *PLoS ONE* 6, e24567, doi:10.1371/journal.pone.0024567.
- Barnish, G., Prentice, M., Harris, S., 1980. *Fasciola hepatica* in St Lucia, West Indies. *Br. Vet. J.* 136, 299–300.

- Bendezu, P., Frame, A., Frame, E., Bonilla, C., 1988. Watercress cultivation sites and their relationships to fascioliasis in Puerto Rico. *Journal of Agriculture of the University of Puerto Rico* 72, 405–411.
- Bendezu, P., Frame, A., Hillyer, G., 1982. Human fascioliasis in Corozal, Puerto Rico. *J. Parasitol.* 68, 297–299.
- Bundy, D., Arambulo III, P., Grey, C., 1984. La Fasciolosis en Jamaica: aspectos epidemiológicos y económicos de una zoonosis parasitaria transmitida por caracoles. *Boletín de la Organización Panamericana de la Salud* 96, 1–20.
- Caron, Y., Rondelaud, D., Losson, B., 2008. The detection and quantification of a digenean infection in the snail host with special emphasis on *Fasciola* sp. *Parasitol. Res.* 103, 735–744.
- Caron, Y., Celi-Erazo, M., Hurtrez-Boussès, S., Lounnas, M., Pointier, J.P., Saegerman, C., Losson, B., Benítez-Ortíz, W., 2017. Is *Galba schirazensis* (Mollusca, Gastropoda) an intermediate host of *Fasciola hepatica* (Trematoda, Digenea) in Ecuador? *Parasite* 24, 1–9.
- Chen, M., Mott, K., 1990. Progress in assessment of morbidity due to *Fasciola hepatica* infection: a review of recent literature. *Tropical Disease Bulletin* 87, R1–R38.
- Clench, W., 1963. Land and freshwater mollusks of the Crooked Island group, Bahamas. *Bulletin of the Museum of Comparative Zoology at Harvard College* 128, 393–414.
- Correa, A., De Meeûs, T., Dreyfuss, G., Rondelaud, D., Hurtrez-Boussès, S., 2017. *Galba truncatula* and *Fasciola hepatica*: Genetic costructures and interactions with intermediate host dispersal. *Infect. Genet. Evol.* 55, 186–194.
- Correa, A., Escobar, J., Durand, P., David, P., Jarne, J.P., Pointier, J., Hurtrez-Boussès, S., 2010. Bridging gaps in the molecular phylogeny of the Lymnaeidae (Gastropoda: Pulmonata), vectors of fascioliasis. *BMC Evol. Biol.* 10, 381, doi:10.1186/1471-2148-10-381.
- Correa, A., Escobar, J., Noya, O., Velázquez, L., González-Ramírez, C., Hurtrez-Boussès, S., Pointier, J., 2011. Morphological and molecular characterization of Neotropical Lymnaeidae (Gastropoda: Lymnaeidae), vectors of fascioliasis. *Infect. Genet. Evol.* 11, 1978–1988.
- Cowie, R., Holland, B., 2006. Dispersal is fundamental to biogeography and the evolution of biodiversity on oceanic islands. *J Biogeogr.* 33, 193–198.
- Dobsinsky, O., 1969. Helminthoses of cattle under tropical breeding conditions. *Helminthologia* 2, 167–174.
- Dreyfuss, G., Correa, A., Djuikwo-Teukeng, F., Novobilsky, A., Höglund, J., Pankrác, J., Kasny, M., Vignoles, P., Hurtrez-Boussès, S., Pointier, J.P., Rondelaud, D., 2015. Differences in the compatibility of infection between the liver flukes *Fascioloides magna* and *Fasciola hepatica* in a Colombian population of the snail *Galba* sp. *J. Helminthol.* 89, 720–726.
- Estoup, A., Martin, O., 1996. Marqueurs microsatellites: isolement à l'aide de sondes non-radioactives, caractérisation et mise au point, URL: www2.agroparistech.fr/svs/genere/microsat/microsat.htm.
- Frame, A., Bendezu, P., 1978. Bovine fascioliasis in Puerto Rico. *J. Parasitol.* 68, 136.
- Gentilini, M., Laroche, V., Degremont, A., 1964. Aspects de la pathologie tropicale, parasitaire et infectieuse en République d'Haïti. *Bull. Soc. Pathol. Exot.* 57, 299–306.
- Gomez, J., Vargas, M., Malek, E., 1986. Freshwater mollusks of the Dominican Republic. *The Nautilus* 100, 130–134.
- Grétilat, S., 1966. *Lymnaea cubensis* Pfeiffer 1911, mollusque hôte intermédiaire de *Fasciola hepatica* L., 1858 à la Martinique. *Comptes rendus hebdomadaires des seances de l'Academie des sciences. Serie D: Sciences naturelles* 262, 2736–2739.

- Grétilat, S., 1967. Prospections malacologiques aux Antilles françaises: observations sur l'écologie et l'élevage au laboratoire de *Lymnaea cubensis* Pfeiffer. *Rev. Elev. Méd. Vét. Pays Trop.* 20, 279–289.
- Gutiérrez, A., Pointier, J.P., Fraga, J., Jobet, E., Modat, S., Pérez, R., Yong, M., Sánchez, J., Loker, E., Théron, A., 2003. *Fasciola hepatica*: identification of molecular markers for resistant and susceptible *Pseudosuccinea columella* snail hosts. *Exp. Parasitol.* 105, 211–218.
- Gutiérrez, A., Vázquez, A., Hevia, Y., Sánchez, J., Correa, A., Hurtrez-Boussès, S., Pointier, J.P., Théron, A., 2011. First report of larval stages of *Fasciola hepatica* in a wild population of *Pseudosuccinea columella* from Cuba and the Caribbean. *J. Helminthol.* 85, 109–111.
- Gutiérrez, A., Yong, M., Perera, G., Sánchez, J., Théron, A., 2002. *Fasciola hepatica* (Trematoda: Digenea): its effect on the life history traits of *Pseudosuccinea columella* (Gasteropoda: Lymnaeidae), an uncommon interaction. *Parasitol. Res.* 88, 535–539.
- Harry, H., 1964. The foreign freshwater snails now established in Puerto Rico. *American Malacological Union Annual Report*, 4–5.
- Hubendick, B., 1951. Recent Lymnaeidae, their variation, morphology, taxonomy, nomenclature, and distribution. *Kungliga Svenska Vetenskapsakademiens Handlingar* 4, 1–223.
- Ibrahim, M., Trpis, M., 1987. The effect of *Brugia pahangi* infection on survival of susceptible and refractory species of the *Aedes scutellaris* complex. *Med. Vet. Entomol.* 1, 329–337.
- Jabbour-Zahab, R., Pointier, J.P., Jourdane, J., Jarne, P., Oviedo, J., Bargues, M., Mas-Coma, S., Anglés, R., Perera, G., Balzan, C., Khallaayoune, K., Renaud, F., 1997. Phylogeography and genetic divergence of some lymnaeid snails, intermediate hosts of human and animal fascioliasis with special reference to lymnaeids from the Bolivian Altiplano. *Acta Trop.* 64, 191–203.
- Jones, R., Williams, H., Dalesman, S., Brophy, P., 2015. Confirmation of *Galba truncatula* as an intermediate host snail for *Calicophoron daubneyi* in Great Britain, with evidence of alternative snail species hosting *Fasciola hepatica*. *Parasites & Vectors* 8, 656, doi:10.1186/s13071-015-1271-x.
- Khan, M., Sajid, M., Riaz, H., Ahmed, N., He, L., Shahzad, M., Hussain, A., Khan, M., Iqbal, Z., Zhao, J., 2013. The global burden of fasciolosis in domestic animals with an outlook on the contribution of new approaches for diagnosis and control. *Parasitol Res.* 112, 2421–2430.
- Kock, K., Wolmarans, C., Bornman, M., 2003. Distribution and habitats of the snail *Lymnaea truncatula*, intermediate host of the liver fluke *Fasciola hepatica*, in South Africa. *J. South Afr. Vet. Assoc.* 74, 117–122.
- Lotfy, W., Brant, S., DeJong, R., Le, T., Demiaszkiewicz, A., Rajapakse, R., Perera, V., Laursen, J., Loker, E., 2008. Evolutionary origins, diversification, and biogeography of liver flukes (Digenea, Fasciolidae). *Am. J. Trop. Med. Hyg.* 79, 248–255.
- Lounnas, M., Correa, A., Vázquez, A., Dia, A., Escobar, J., Nicot, A., Arenas, J., Ayaqui, R., Dubois, M., Gimenez, T., Gutiérrez, A., González-Ramírez, C., Noya, O., Prepelitchi, L., Uribe, N., Wisnivesky-Colli, C., Yong, M., David, P., Loker, E., Jarne, P., Pointier, J.P., Hurtrez-Boussès, S., 2017. Self-fertilization, long-distance flash invasion and biogeography shape the population structure of *Pseudosuccinea columella* at the worldwide scale. *Mol. Ecol.* 26, 887–903.
- Malek, E., 1980. *Snail transmitted parasitic diseases*. CRC Press, Florida, USA, 348 pp.
- Mas-Coma, S., Bargues, M., Valero, M., 2018. Human fascioliasis infection sources, their diversity, incidence factors, analytical methods and prevention measures. *Parasitology* 145, 1665–1699.
- Mas-Coma, S., Valero, M.A., Bargues, M.D., 2009. *Fasciola*, lymnaeids and human fascioliasis, with a global overview on disease transmission, epidemiology, evolutionary genetics, molecular epidemiology and control. *Adv. Parasitol.* 69, 41–146.

- Meunier, C., Tirard, C., Hurtrez-Boussès, S., Durand, P., Bargues, M., Mas-Coma, S., Pointier, J., Jourdane, J., Renaud, F., 2001. Lack of molluscan host diversity and the transmission of an emerging parasitic disease in Bolivia. *Mol. Ecol.* 10, 1333–1340.
- Pointier, J.P., 1974. Faune malacologique dulçaquicole de l'île de la Guadeloupe (Antilles françaises). *Bulletin du Muséum National d'Histoire Naturelle* 3, 905–933.
- Pointier, J.P., 2008. *Guide to the Freshwater Molluscs of the Lesser Antilles*. ConchBooks, Hackenheim, Germany, 146 pp.
- Reiczigel, J., Marozzi, M., Fâbiân, I., Rózsa, L., 2019. Biostatistics for parasitologists – a primer to quantitative parasitology. *Trends Parasitol.* 35, 277–281.
- Robart, G., Mandahl-Barth, G., Ripert, C., 1977. Inventaire, répartition géographique et écologie des mollusques dulçaquicoles d'Haiti (Caraïbes). *Revue d'Agriculture, Damien Haiti* 2, 20–64.
- Rojas, L., Vázquez, A., Domenech, I., Robertson, L., 2010. Fascioliasis: can Cuba conquer this emerging parasitosis? *Trends Parasitol.* 26, 26–34.
- Rondelaud, D., Sanabria, R., Vignoles, P., Dreyfuss, G., Romero, J., 2013. *Fasciola hepatica*: variations in redial development and cercarial production in relation to the geographic origin of the parasite. *Parasite* 20, 33, doi:10.1051/parasite/2013034.
- Rondelaud, D., Vignoles, P., Dreyfuss, G., Mage, C., 2006. The control of *Galba truncatula* (Gastropoda: Lymnaeidae) by the terrestrial snail *Zonitoides nitidus* on acid soils. *Biological Control* 39, 290–299.
- Sánchez, R., Perera, G., Sánchez, J., 1995. Cultivo de *Fossaria cubensis* (Pfeiffer) (Pulmonata: Lymnaeidae) hospedero intermediario de *Fasciola hepatica* (Linnaeus) en Cuba. *Rev. Cub. Med. Trop.* 47, 71–73.
- Sorensen, R., Minchella, D., 2001. Snail-trematode life history interactions: past trends and future directions. *Parasitology* 123, 3–18.
- Théron, A., Coustau, C., 2005. Are *Biomphalaria* snails resistant to *Schistosoma mansoni*? *J. Helminthol.* 79, 187–191.
- Vázquez, A., Alda, P., Lounnas, M., Sabourin, E., Alba, A., Pointier, J.P., Hurtrez-Boussès, S., 2018. Lymnaeid snails hosts of *Fasciola hepatica* and *Fasciola gigantica* (Trematoda: Digenea): a worldwide review. *CAB Reviews* 13, 62, doi:10.1079/PAVSNNR201813062.
- Toet, H., Piedrafita, D.M., Spithill, T.W., 2014. Liver fluke vaccines in ruminants: strategies, progress and future opportunities. *Int. J. Parasitol.* 44, 915–927.
- Vázquez, A., Lounnas, M., Sánchez, J., Alba, A., Milesi, A., Hurtrez-Boussès, S., 2016. Genetic and infective diversity of the liver fluke *Fasciola hepatica* (Trematoda: Digenea) from Cuba. *J. Helminthol.* 90, 719–725.
- Vázquez, A., Sánchez, J., Alba, A., Pointier, J.P., Hurtrez-Boussès, S., 2015. Natural prevalence in Cuban populations of the lymnaeid snail *Galba cubensis* infected with the liver fluke *Fasciola hepatica*: small values do matter. *Parasitol. Res.* 114, 4205–4210.
- Vázquez, A., Sánchez, J., Pointier, J.P., Théron, A., Hurtrez-Boussès, S., 2014. *Fasciola hepatica* in Cuba: compatibility of different isolates with two intermediate snail hosts, *Galba cubensis* and *Pseudosuccinea columella*. *J. Helminthol.* 88, 434–440.
- Wesselingh, F., Cadée, G., Renema, W., 1999. Flying high: on the airborne dispersal of aquatic organisms as illustrated by the distribution histories of the gastropod genera *Tryonia* and *Planorbarius*. *Geologie en Mijnbouw* 78, 165–174.

Tables

Species	Country	Locality	Code	Coordinates	Habitat
Galba cubensis	Cuba	Arroyo Dolores	CU.01	22°26'27.6"N 79°25'51.6W	brook
		Aurora	CU.02	23°4'44.4"N 81°55'4.8"W	pond
		Playa Jibacoa	CU.03	23°8'38.4"N 81°28'15.6"W	brook
	Dominican Republic	Auyamas	DR.01	18°53'34.8"N 70°44'45.599"W	ditch
		La Victoria	DR.02	18° 34' 22.8"N 69°50'52.799"W	pond
Galba schirazensis	Dominican Republic	Auyamas	DR.03	18°53'31.2"N 70°44'45.599"W	ditch
		Rosa Linda	DR.04	18°58'44.4"N N 70°38'20.399"W	ditch
Galba truncatula	France	Saint Guilhem	FR.01	43°44'2.4"N 3°32'56.399"E	ditch
Pseudosuccinea columella	Cuba	Aurora	CU.04	23°4'44.4"N 81°55'4.8"W	pond
		El Chico	CU.05	22°57'28.8"N 82°27'54"W	channel
	Dominican Republic	Jarabacoa	DR.05	19°6'21.6"N 70°37'44.4"W	watercress beds
		Río Isbe	DR.06	18°30'57.6"N 69°57'7.199"W	Rive
	Pseudosuccinea columella				
St. Lucia	Galba cubensis	NK		Malek (1980)	
		NK	10-23%	Barnish et al. (1980)	

Table 1. Past and recent reports of *Fasciola hepatica* transmission and snail host occurrence in the West Indies. NK: not known

Table 2. Sampling details of parental lymnaeid snails used to establish laboratory populations for infection trials (each code refers to country.locality).

Species	Country	Locality	Code	Coordinates	Habitat
Galba cubensis	Cuba	Arroyo Dolore	CU.01	22°26'27.6"N 79°25'51.6W	brook
		Aurora	CU.02	23°4'44.4"N 81°55'4.8"W	pond
		Playa Jibacoa	CU.03	23°8'38.4"N 81°49'15.6"W	brook
	Dominican Republic	Auyamas	DR.01	18°53'34.8"N 70°44'45.599"W	ditch
		La Victoria	DR.02	18° 34' 22.8"N 69°50'52.799"W	pond
Galba schirazensis	Dominican Republic	Auyamas	DR.03	18°18'53'31.2"N 70°44'45.599 W	ditch
		Rosa Linda	DR.04	18°58'44.4" N 70°38'20.399"W	ditch
Galba truncatula	France	Saint Guilhem	FR.01	43°44'2.4"N 3°32'56.399"E	ditch
Pseudosuccinea columella	Cuba	Aurora	CU.04	23°4'44.4"N81°55'4.8"W	pond
		El Chico	CU.05 DR.05	22°57'28.8"N82°27'54"W	channel
	Dominican Republic	Jarabacoa		19°6'21.6"N 70°37'44.4"W	watercress beds
		Río Isabela	DR.06	18°30'57.6"N 69°57'7.199"W	river

Figure captions




Figure 1. Map of the West Indies indicating the Greater and Lesser Antilles with data on the lymnaeid species reported in each island




Figure 2. Populations of lymnaeid snails and liver fluke isolates used in the experimental infection trials.




Figure 3. Prevalence and mean intensities (with 95% confidence intervals vertical bars) of *Fasciola hepatica* in relation to the combination with a particular population of lymnaeid snail originating from Cuba, the Dominican Republic and France. Intensities are compared by means of a one-way ANOVA, different letters mean significant differences between the means according to the post hoc Tukey test.




Figure 4. Survival curves of each lymnaeid population when exposed to a particular *Fasciola hepatica* isolate (A: *Galba cubensis*, B: *Galba schirazensis*, C: *Galba truncatula*, D: *Pseudosuccinea columella*; population codes are in