



HAL
open science

Aut(F₂) puzzles

Sylvain Barré, Mikael Pichot

► **To cite this version:**

Sylvain Barré, Mikael Pichot. **Aut(F₂) puzzles**. *Geometriae Dedicata*, 2019, 199 (1), pp.225-246.
10.1007/s10711-018-0346-1 . hal-02508106

HAL Id: hal-02508106

<https://hal.science/hal-02508106v1>

Submitted on 5 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Aut(F_2) PUZZLES

SYLVAIN BARRÉ AND MIKAEL PICHOT

ABSTRACT. This paper studies $\text{Aut}(F_2)$ and groups closely related to it from a geometric perspective.

The discussion is centered around geometric structures called “ring puzzles” and “ring complexes”, and groups which can be seen as “randomizations” of the group $\text{Aut}(F_2)$, in the sense of [4], and can be investigated using these structures.

In a recent paper [4] we discussed a class of “random groups of higher rank”. These random groups come in various models, including in particular a “density model” (as in the classical construction of Gromov), and they are associated with “deterministic data” that are used to seed the construction.

The deterministic data in [4] was associated with nonarchimedean lattices, which were chosen in particular for their relations to rank interpolation, and the question of whether

$$\mathbb{R}^2 \hookrightarrow X \xrightarrow{?} \mathbb{Z}^2 \hookrightarrow \Gamma$$

whenever $\Gamma \curvearrowright X$ properly, X is nonpositively curved and X/Γ compact. The so-called “flat closing conjecture” asserts that this has a positive answer.

The present paper uses deterministic data associated with $\text{Aut}(F_2)$, which produces random groups which “resemble $\text{Aut}(F_2)$ ” in a geometrical sense.

We prove that:

Theorem 1. *The random group in the density models associated with $\text{Aut}(F_2)$ satisfies the flat closing conjecture.*

More precisely, the statement holds with overwhelming probability, at arbitrary density $\delta \in (0, 1)$, and with respect to arbitrary deterministic data associated with $\text{Aut}(F_2)$.

The corresponding result for nonarchimedean lattices was left open in [4], and the techniques used in the present paper do not appear to shed light on the problem.

CONTENTS

1. Organization of the proof	2
2. Ring puzzles	3
3. Ring complexes and flat plane problems	19
4. Randomizing $\text{Aut}(F_2)$	23
5. The explicit construction	27
References	27

1. ORGANIZATION OF THE PROOF

In what sense does the random group resemble $\text{Aut}(F_2)$?

The proof of Theorem 1 relies on giving an answer to this question which is precise enough to keep track of the (dis)appearance of (periodic) flat planes in the random groups.

The text is centered around geometric structures called “ring puzzles” and “ring complexes”, which are defined precisely in §2 and §3 respectively.

Ring puzzles are tessellations of the Euclidean plane with constraints on the set of tiles and vertex neighbourhoods (see §2). Associated with $\text{Aut}(F_2)$ are ring puzzles, called $\text{Aut}(F_2)$ puzzles, with two tiles and three types of neighbourhoods. The definition is elementary involving simple local construction rules that are not directly reminiscent of $\text{Aut}(F_2)$. Incidentally, ring puzzles give informative “snapshots” of the geometry of $\text{Aut}(F_2)$ itself.

It is known that $\text{Aut}(F_2)$ acts properly with compact quotient on a 2-dimensional $\text{CAT}(0)$ space called the Brady complex, which is essentially uniquely determined, by results of Crisp and Paoluzzi [5]. The space of $\text{Aut}(F_2)$ puzzles is related to the space of flats in the Brady complex or in any ring complex of type $\text{Aut}(F_2)$. In the case of the Brady complex itself, the space of flat surjects onto the space of puzzles, which is an indication that $\text{Aut}(F_2)$ plays the role of “space of maximal rank” in rank interpolation [1].

The main objective in §2, and the first step in the proof of Theorem 1, is to show that:

Theorem. *There is an explicit classification of all $\text{Aut}(F_2)$ puzzles.*

The classification finds:

- 4 infinite families of puzzles
- 9 exceptional puzzles not belonging to the families

See Theorem 21 for a detailed statement.

(Large portions of a puzzle do not determine the puzzle in general, puzzles are “bifurcating spaces” with an interesting (pointed) Gromov–Hausdorff topology; see Remark 22.)

Ring complexes are defined in §3 as metric simplicial complexes subjects to “ring conditions” on the links. Compared to classical curvature conditions (the girth condition in dimension 2), the ring conditions serve to control the local flatness and shed light on the “intermediate rank” structure of $\text{Aut}(F_2)$ rather than the curvature. Local rules identical to that of $\text{Aut}(F_2)$ puzzles give rise to ring complexes of type $\text{Aut}(F_2)$. The random group in models associated with $\text{Aut}(F_2)$ acts on a ring complex of type $\text{Aut}(F_2)$.

Theorem 1 is proved in §4 using the classification of puzzles. It relies on a more general result in §3 (Theorem 25) establishing the flat closing conjecture under a technical assumption which is satisfied for randomizations of $\text{Aut}(F_2)$.

We do not pursue the study of general ring puzzles in this paper. The definition is very much in the spirit of introducing new rank interpolation tools (compare [1])

or [3]). The ring conditions seem to give rise to an interesting set of objects to study in their own right.

We will further discuss two points related to the Theorem 1 and its proof:

- a) There are several significant differences between randomizations of $\text{Aut}(F_2)$ and randomizations of nonarchimedean lattices as considered in [4]. For example, we show in §3 that:

Theorem. *For some choice of the deterministic data, the random group contains \mathbb{Z}^2 with overwhelming probability at arbitrary density $\delta \in (0, 1)$.*

Here the deterministic data has to be chosen suitably, it is associated the symmetric groups S_n . The property underlying the statement of the theorem, which is satisfied for deterministic data associated with S_n , is the *small tori property* defined in §3, see Definition 33.

- b) The above raises the question of using ring puzzles as “blueprints” to construct groups which resemble $\text{Aut}(F_2)$ in a geometrical sense. While we can produce a wide range of random examples, finding explicit groups does not appear to be an easy task. We give an explicit construction of such a group in §5.

2. RING PUZZLES

A *ring puzzle* is a tessellation of the Euclidean plane \mathbb{R}^2 using planar polygons with coloured angles. The *ring* at a vertex of the puzzle is the coloured circle (of length 2π) of small radius around that vertex.

By *ring puzzle problem* we mean the problem of classifying all puzzles whose coloured polygons and rings belong to prescribed sets.

We consider here the following instance of the problem, which is related to $\text{Aut}(F_2)$.

The two prescribed sets are:



(an equilateral triangle and a lozenge) and




The yellow arcs have length $2\pi/3$, the others $\pi/3$. (For details on the relation with $\text{Aut}(F_2)$, see §3.)

We assume that we have an infinite supply of coloured polygons for each shape type, called the pieces of the puzzle. The position and orientation of the pieces in the plane is free.

In this section we discuss the ring puzzle problem on these two sets. Every solution to the problem is called an $\text{Aut}(F_2)$ *puzzle*.

For example, here is a permitted (bi-infinite) portion of an $\text{Aut}(F_2)$ puzzle which we shall call the *2-strip*.



The main result (Theorem 21) classifies $\text{Aut}(F_2)$ puzzles. This can be achieved after a series of preliminary results on the local-global geometry of these puzzles.

Let us first isolate an important property of the two sets given above, which reflects the fact that $\text{Aut}(F_2)$ puzzles are “nonpositively curved”:

Definition 2. A ring puzzle problem has the θ_0 -*extension property* if the prescribed sets satisfies the following condition: whenever a coloured segment $[0, \theta]$ of length $\theta \in [0, 2\pi]$ can be isometrically embedded in distinct coloured rings respecting the coloration, then $\theta \leq \theta_0$.

For $\text{Aut}(F_2)$ puzzles we have that $\theta_0 = \pi$, which corresponds to nonpositive curvature.

The following consequence is straightforward:

Lemma 3. *Assuming the θ_0 -extension property, every ring puzzle is such that if consecutive pieces meet at a vertex contributing to an angle $> \theta_0$, then the remaining pieces at this vertex are uniquely determined (up to colour preserving isometry).*

Let P be an $\text{Aut}(F_2)$ puzzle.

Definition 4. A \diamond -gallery in P is a sequence of consecutive (= intersecting along an edge) lozenges. Two lozenges are connected if they belong to a \diamond -gallery. The connected components of lozenges in P are called the \diamond -components of P .

Lemma 5. *Let C be a \diamond -component. The inner angle at every vertex of ∂C is equal to either $\pi/3$, $2\pi/3$ or π .*


Proof. The inner angle is a multiple of $\pi/3$. Assume that it is $> \pi$; since we are on the boundary, it can only be $4\pi/3$ or $5\pi/3$. The value $4\pi/3$ would single out the first ring, however, the remaining colour (namely yellow) corresponds to a lozenge, contradicting the fact that we are on the boundary. The value $5\pi/3$ is prohibited by the fact that there is no yellow–green simplicial path of length $5\pi/3$ inside in the prescribed set of rings (by simplicial we mean both of whose end points are vertices of the ring). \square

Let us consider the finite \diamond -components first.

Lemma 6. *Every finite \diamond -component of P is a parallelogram of size $m \times n$ such that $\min(m, n) \leq 2$.*

Proof. The fact that finite \diamond -components are parallelograms follow from Lemma 5. If $m \times n$ denotes the size of such a component with $m \leq n$, then we claim that $m \leq 2$. This is a combinatorial problem which can be solved in the 3-neighbourhood of the component. Note that the 3rd ring has to be used at every point in the boundary, except at the two obtuse angles, where it is the 2nd ring that has to be used.


The solution of the problem, assuming $m \geq 3$ towards a contradiction, is summarized in the following drawing:



The rightmost part does not extend in P given the prescribed sets of types. The drawing explicit one of several similar cases. \square

In general, the prescribed sets in a ring puzzle problem will determine basic blocks (“tiles”) that fit together to build larger portions of the puzzle using the extension property.

An interesting block for Aut(F_2) puzzles is the w -block:




The w -block appears in the proof of Lemma 6.

Observe that w -blocks are “forward analytic” in the following sense:

Lemma 7. *Every w -block in P extends to a unique w -strip in P .*


By w -strip we mean the strip (infinite at least on the right hand side):



Lemma 7 is a consequence of Lemma 3.

Note that two w -strips cannot intersect transversally in the forward direction.

Lemma 8. *The \diamond -components of type 2×2 in P extend uniquely into a bi-infinite strip of height 4 of the form*




Proof. This is a consequence of Lemma 7. □


We call this strip the double w -strip.

Lemma 9. *Finite \diamond -component of type $2 \times n$ with $n \geq 3$ are contained in a unique puzzle.*

Proof. A \diamond -component C of size $2 \times n$ with $n \geq 3$



admits a unique extension of the form




by Lemma 7.

Furthermore using the extension property we have a unique upward extension (and by symmetry a unique downward extension) of the strip. This in turn exhibits a w -strip in the north-east direction (and by symmetry in the south-west direction) that edges an acute sector along with the horizontal w -strip, implying that the horizontal strip belongs to at most one puzzle. A construction by induction then shows that the puzzle exists.


We will call this puzzle the *puzzle of type $2 \times n$* . □

Here is a portion of the puzzle of type 2×4 :



Remark 10. Lemma 8 can be understood as the limiting case “ $n \rightarrow 2$ ” of Lemma 9.

Lemma 11. Assume that P contains a finite \diamond -component C of type $1 \times n$ where $n \geq 3$. Then C is included in a strip S of height n in P (infinite both to the left and to the right) of the form:




Furthermore, the following two conditions are satisfied:


- the \diamond -components of S parallel to C are all of type $1 \times n$, except for at most one component of type $2 \times n$. If there is such a component, then P is the puzzle of type $2 \times n$, otherwise S is as indicated in the figure above.
- every \diamond -component of S remains a \diamond -component in P (otherwise said, it is adjacent to triangles in $P \setminus S$ on both sides, as indicated in the figure).

The strip described in the figure above is called the *bi-infinite strip* of type $1 \times n$ (all \diamond -component of the strip are of type $1 \times n$).

Proof. Starting with a \diamond -component in P of size $1 \times n$



there is a unique extension of the form




which by induction form a half strip $S_{\uparrow} \supset C$ in the upward direction, together with the right 1-neighbourhood of it. The same argument applies in downward direction by symmetry, which delivers the indicated strip $S = S_{\uparrow} \cup S_{\downarrow} \supset C$ together with a 1-neighbourhood of it on both sides.

Inspection of the neighbourhood shows that the \diamond -components parallel to C are \diamond -component in P , which therefore are of the form $m \times n$ for $m = 1, 2$.

We have to prove that a component of type $2 \times n$ can only appear once. This can be done directly noticing that the puzzle of type $2 \times n$ actually contains only one \diamond -component of type $2 \times n$, or with the following direct proof. Suppose we have two components, without loss of generality we may assume that they are separated

only by components of type $1 \times n$.



Note that the eastern neighbourhood of any \diamond -component of type $2 \times n$ contains two w -blocks whose *forward* direction intersect transversally.


By Lemma 7, w -blocks extend uniquely in the forward direction. This contradicts the fact that w -strips do not intersect transversally, and proves the result.

The figure above illustrates the argument. We have indicated the completion in the case of two \diamond -components of type $2 \times n$ separated by precisely one \diamond -component of type $1 \times n$. In this case a direct local contradiction appears in the 1-neighbourhood of the upmost component of type $2 \times n$. \square

Lemma 12. *Assume that every component is finite and of type $1 \times n$ with $n = 1, 2$. Then every finite component is of type 1×2 , and corresponds to a unique corresponding Aut(F_2) puzzle.*

Proof. It is easy to check that there is no puzzle in which every \diamond -component is of type 1×1 . The proof of the lemma, starting from a component of type 1×2 , is then in the same spirit as that of Lemma 11, given the fact that every component

has type $1 \times n$ with $n \leq 2$. The resulting puzzle of type 1×2 is



□

We now turn to infinite \diamond -components.

Lemma 13. *An infinite \diamond -component is either*

- a) *the tessellation of the plane by lozenges (called the \diamond -plane) or half of it (called the half \diamond -plane)*
- b) *a bi-infinite strip of lozenges (\diamond -strip) or half of it (semi-infinite \diamond -strip)*
- c) *a \diamond -sector (with either an acute or an obtuse angle)*

Proof. This follows from Lemma 5. □

Every (semi-infinite) \diamond -strip of lozenges is bounded by a (semi-infinite) 2-strip. In particular both half \diamond -planes and bi-infinite \diamond -strip give rise to an alternating 2-strip/ \diamond -strip $\text{Aut}(F_2)$ puzzles.


Note that:

Lemma 14. *If P contains a semi-infinite \diamond -strip S of height n , then $n = 1$ or $n = 2$.*

The proof is similar to that of Lemma 6.


Lemma 15. *The semi-infinite \diamond -strips of type $2 \times \infty$ can be extended in precisely two ways: a puzzle called the star puzzle of type $2 \times \infty$, or a half-puzzle whose*

boundary is the following strip of height 3:




We refer to the above strip as “the 3-strip”, and the half-puzzle containing it will be called the half puzzle of type $2 \times \infty$.

Proof. Take a semi-infinite (horizontal) \diamond -strip S of height 2. The proof of Lemma 11 shows that there exists a unique sector extending S “in the upper direction”. This sector contains a double w -strip as follows:




There are two extensions of it, corresponding to the two ways a lozenge can be oriented with respect to adjacent the south-east triangles in the figure.

We have already encountered the first extension. It can be derived from the proof of Lemma 9 or Lemma 11 and appears as a “half of” the limit (in the Gromov-Hausdorff sense, say) of the puzzle of type $2 \times n$ as $n \rightarrow \infty$:



The lowermost 3-strip is the 3-strip.

The second extension, which leads to a unique puzzle, is




That it can be extended into at most one puzzle follows by Lemma 7 (and Lemma 3), and it is easy to show that this puzzle (the star puzzle of type $2 \times \infty$) exists by induction. \square

In particular, it appears that the semi-infinite \diamond -strip of type $2 \times \infty$ embeds twice (at most) in an $\text{Aut}(F_2)$ puzzle.


Lemma 16. *If P contains a semi-infinite \diamond -strip of type $1 \times \infty$ then one of the following holds:*

- P is the puzzle of type $2 \times n$ where $n \geq 3$
- P contains the half-puzzle of type $2 \times \infty$
- P is the star puzzle of type $2 \times \infty$
- P contains the double w -strip (see Lemma 8)
- P contains the half-puzzle of type $1 \times n$ (defined below) whose boundary strip of type $1 \times n$ for some $n \geq 3$
- P is the V -puzzle:



Proof. Take a semi-infinite (horizontal) \diamond -strip S of height 1. Again as in the proof of Lemma 11 one can construct a quarter puzzle P_{\uparrow} extending S “in the upper direction” which contains parallel (semi-infinite) 2-strips and (by assumption) semi-infinite \diamond -strip S of height 1.

The lower part of P_{\uparrow} admits a unique extension as follows:




(In the picture the semi-infinite strips are horizontal.)


If the piece at X is a triangle, then we have a \diamond -component of type $2 \times n$ below it heading south-east. There are 4 cases:

- If $n = 1$ then P_{\uparrow} extends uniquely east using a 3-strip. The resulting half-puzzle $P_{\uparrow, \rightarrow}$ piles up horizontal 3-strips.
- If $n = 2$ then by Lemma 8 the quarter puzzle P_{\uparrow} extends uniquely east using the 4-strip (compare Lemma 20).
- If $3 \leq n < \infty$ then P is the puzzle of type $2 \times n$ (see Lemma 9)
- Otherwise, $n = \infty$, and Lemma 15 shows that either P is the star of type $2 \times \infty$ or contains the half-puzzle of type $2 \times \infty$.

We now assume that the piece at X is a square and extend the portion of the puzzle uniquely as follows:



If the piece at Y is a triangle, then by Lemma 6 the puzzle contains a \diamond -component of type $m \times 3$ with $m = 1, 2$, and we can apply Lemma 11 to find either that P is the puzzle of type 2×3 , or if $n = 1$ that P contains the half-puzzle of type 1×3 .




The puzzle of type $1 \times n$ (for $n = 4$)

The conclusion is identical for every $n = 3, 4, \dots$. If $n = \infty$ we obtain a sector in P which is part of the V -puzzle (and corresponds to the limit $n \rightarrow \infty$). Since the horizontal strips alternate 2-strips and \diamond -strips of height 1, this sector in P extends uniquely to the V -puzzle. \square

Let us now turn to the \diamond -sectors, both of which are limiting cases of half \diamond -strips as the height increases. Recall however that \diamond -strips in flat can only have height 1 or 2.

Lemma 17. *If the puzzle contains a sector with acute angle, then it contains a component of type $2 \times n$ for some $n = 1, \dots, \infty$ (if $n = \infty$ this is the semi-infinite \diamond -strip of type 2).*

Proof. Acute sectors extend uniquely (up to symmetry along the bisector) to the following finite puzzle:




This puzzle contains (at the bottom-left) a \diamond -component of type $2 \times n$ with $n = 1, \dots, \infty$. If $n = 1$ then the sector extends uniquely to a half plane containing the 3-strip. If $n = 2$ then it extends uniquely to a half-plane containing the 4-strip. Otherwise

By Lemma 12, we have $n \geq 2$ and by Lemma 8 we have $n \geq 3$. Applying Lemma 11 gives the conclusion. \square

Lemma 18. *There exists a unique puzzle containing a sector with obtuse angle.*

Proof. Obtuse sectors extend to:



which creates a w -strip. Existence and uniqueness of the puzzle is proved by induction using Lemma 3 as earlier. \square


Let us then classify extensions of the 3-strips and the double w -strips.

Lemma 19. *If P contains the 3-strip S , then P contains only parallels 3-strips and at most two copies of the half puzzle of type $2 \times \infty$.*

The 3 -strip puzzle is the puzzle made only of parallels 3-strips. Every other puzzle is called the 3-strip puzzle of height h where $h \geq 1$ is the number of parallels 3-strips it contains. We call *half 3-strip puzzle* the puzzle containing exactly one half-puzzle of type $2 \times \infty$. Every 3-strip puzzle of height $h < \infty$ contains two such puzzles.


Lemma 20. *There are precisely two puzzles containing the double w -strip.*

Proof. Note that, using the symmetry along the vertical axis, the double w -strip extends uniquely as follows:



By Lemma 7 this can be extended north in a unique way. \square

The two puzzles in Lemma 20 will be called the opposite acute sector puzzle and the adjacent acute sector puzzle. The corresponding half puzzle is:



Theorem 21. *The Aut(F_2) puzzles come in 4 infinite series and 9 exceptional puzzles not belonging to the series.*

The 9 exceptional puzzles are:

- a) *the \diamond -puzzle*
- b) *the 2×1 -puzzle (see Lemma 12)*
- c) *the star puzzle of type $2 \times \infty$*
- d) *the 3-strip puzzle and the half 3-strip puzzle*
- e) *the opposite acute sector puzzle and the adjacent acute sector puzzle (see Lemma 8 and Lemma 20).*
- f) *the obtuse sector puzzle (see Lemma 18)*
- g) *the V-puzzle*

The 4 infinite series are:

- A) *the alternating 2-strip/ \diamond -strip series. This contains three subseries:*
 - *the bounded 2-strip/ \diamond -strip series. They are parametrized by biinfinite sequences $(\dots, n_{-1}, n_0, n_1, \dots)$ recording the heights of the \diamond -strips in the given order, where $n_i \geq 1$ is arbitrary. There are uncountably many such puzzles.*
 - *The sided 2-strip/ \diamond -strip series. The 1-sided series is parametrized by right infinite sequences (n_0, n_1, \dots) of heights of \diamond -strips, where $n_i \geq 1$ is arbitrary, giving uncountably many puzzles. The 2-sided series is parametrized by finite sequences (n_0, n_1, n_r) where r is finite but arbitrary. This gives countably many puzzles.*

- B) the $1 \times n$ series. They are constructed using the strip of type $1 \times n$ (Lemma 11) and can be encoded by the biinfinite sequence $(\dots, n_{-1}, n_0, n_1, \dots)$ of heights here $n_i \geq 1$ is arbitrary, and at least one $n_i \geq 3$. There are uncountably many such puzzles. This series also splits into 3 subseries, the bounded $1 \times n$ series and the sided $1 \times \infty$ series, which can either be one-sided or two-sided.
- C) the $2 \times n$ series, where $n = 3, 4, \dots (n < \infty)$. There is a unique puzzle for every $n \times 2$ (Lemma 9).
- D) the 3-strip series of height $h \geq 1$ (Lemma 19).

Several of the exceptional puzzles appear as limits of elements of the infinite series. For example, a), c), and d) can be seen as limiting cases of A), C), and D) respectively, and b) can be seen as a limit of B) when “ $n \rightarrow 2$ ”. The V -puzzle g) can also be seen as a limiting case for B), and D) can be seen as a limiting case for C) when $n \rightarrow \infty$. The obtuse sector puzzle, on the other hand, does not appear to be a limit — and similarly for e).

Proof. Let P be an $\text{Aut}(F_2)$ puzzle. We assume that P is not one of the exceptional puzzles and prove that it belongs to one of the series.

Note that P does not contain an obtuse sector. If it contains an acute sector then it contains a strip of type $2 \times n$. If it contains a semi-infinite strip of type $2 \times \infty$ and since it is not the star puzzle of type $2 \times \infty$ it contains the 3-strip. Not being in d) it belongs to D).

If P contains a semi-infinite strip of type $1 \times \infty$ but no semi-infinite strip of type $2 \times \infty$, then it belongs to B) by Lemma 16.

If P contains a bi-infinite \diamond -strip then it belongs to A).

By Lemma 13 we may now assume that every \diamond -component of P is finite. If it has a component of type $2 \times n$ then we are in case C). Otherwise every component is finite of type $1 \times n$. Then this is case B). \square

Remark 22. One can organize puzzles into a “space of puzzles” as follows. A marked puzzle is a parametrization $f: \mathbb{R}^2 \rightarrow P$ where f is isometric. Define a valuation v on the set $\mathbf{Puzzles}'$ of all marked puzzles (with prescribed shapes and rings) by

$$v(f, f') = \sup\{r, f(B(0, r)) \stackrel{\text{simplicial isometric}}{\simeq} f'(B(0, r))\}$$

with corresponding metric

$$d(f, f') = e^{-v(f, f')}.$$

The space of marked puzzles is

$$\mathbf{Puzzles} := \mathbf{Puzzles}' / \{d = 0\}.$$

Some of the statements regarding convergence of puzzles can be interpreted in the space $\mathbf{Puzzles}$. For example the obtuse sector puzzle “is not a limit” because it corresponds to an isolated point in $\mathbf{Puzzles}$. Note that the space of puzzles is compact if the sets of shapes and rings are finite. It is endowed with an action of \mathbb{R}^2 given by

$$t \cdot f := f \circ t^{-1}$$

for t a translation of \mathbb{R}^2 turning $\mathbf{Puzzles}$ into a lamination whose leaves correspond to puzzles with the given set of shapes and rings.

3. RING COMPLEXES AND FLAT PLANE PROBLEMS

Ring puzzles, or portions thereof, can be “glued together” to form “nonplanar ring puzzles”, which are more complicated objects to understand.

Let us start with a 2-complex X and say that X is a *ring complex* if its faces are flat polygons in the Euclidean plane with coloured angles.

The *rings* at a vertex of X is set of all coloured circles of length 2π of small radius around that vertex.

The *type* of a ring complex is the prescribed set of coloured faces and rings (considered up to coloured isometry).

We study ring complexes of type $\text{Aut}(F_2)$ as considered earlier:

$$T := \left\{ \begin{array}{c} \text{triangle with colored angles} \quad \text{parallelogram with colored angles} \quad \text{three colored circles} \end{array} \right\}$$

Lemma 23. *The group $\text{Aut}(F_2)$ acts properly on a ring complex of type $\text{Aut}(F_2)$ with compact quotient.*

Proof. The Brady complex has the required properties. □

Note that ring complexes are naturally endowed with a global metric, namely the length metric. They are not necessarily nonpositively curved.

A flat plane in X is an isometric embedding $\mathbb{R}^2 \hookrightarrow X$. Every flat plane in a ring complex is a ring puzzle of the same type.

The \mathbb{Z}^2 *embedding problem* for X is the question of whether

$$\mathbb{R}^2 \hookrightarrow X \Rightarrow \mathbb{Z}^2 \hookrightarrow \Gamma$$

whenever $\Gamma \curvearrowright X$ properly with X/Γ compact. Gromov asks for example if this is true for X simplicial and nonpositively curved, say, of dimension 2 (see [7]). When X is nonpositively curved we have

$$\mathbb{Z}^2 \hookrightarrow \Gamma \Rightarrow \mathbb{R}^2 \hookrightarrow X$$

by the flat torus theorem and the “flat closing conjecture” asserts the embedding problem has a positive answer. In §4 and §5 the groups are nonpositively curved.

Definition 24. Consider a simplicial strip $S \simeq \mathbb{R} \times [a, b]$ and a simplicial subset $L \subset S$. The strip S is said to be uniquely L -embeddable in X if for any two simplicial embeddings $f, f': S \hookrightarrow X$ and any translation $g: S \rightarrow S$

$$f' = f \circ g \text{ on } L \Rightarrow f' = f \circ g \text{ on } S.$$

(Note that g is then necessarily a simplicial map.)

Consider the following two 1-strips in a ring complex of type $\text{Aut}(F_2)$:

- the \diamond -strip:



- the Δ -strip:



These strips are said to be *uniquely \diamond -embeddable* if they are uniquely L -embeddable with respect to a lozenge.

We say that X is acylindrical if there is no embedding

$$\mathbb{S}^1 \times \mathbb{R} \not\hookrightarrow X$$

of a bi-infinite cylinder into X .

Theorem 25. *If in an acylindrical ring complex of type $\text{Aut}(F_2)$ every \diamond -strip and every Δ -strip is uniquely \diamond -embeddable, then the \mathbb{Z}^2 embedding problem has a positive solution.*

The proof of Theorem 25 relies on two lemmas. The first lemma uses the classification.

Lemma 26. *Suppose that X is a ring complex X of type $\text{Aut}(F_2)$ and fix $\Gamma \curvearrowright X$ with X/Γ compact. If X contains a flat, then it contains a flat that contains infinitely many parallel \diamond -strips, or a flat that contains infinitely many parallel Δ -strips.*

Proof. We claim that if X contains a flat, then it contains a flat which is isomorphic to either a \diamond -puzzle, a puzzle of uniformly bounded type $1 \times n$ (including $n = 2$), or a puzzle of uniformly bounded alternating 2-strip/ \diamond -strip type. The lemma follows easily from this.

The proof of the claim is a standard sort of compactness argument which relies on the classification. Let Π be a flat and K be a relatively compact fundamental domain for $\Gamma \curvearrowright X$.

Suppose for example that Π contains a \diamond -sector S . Choose vertices x_n in S moving apart from the boundary of S , and $s_n \in \Gamma$ such that $s_n x_n \in K$. For every $r > 0$ we can find infinitely many n for which the flats $s_n \Pi$ coincide on r -neighbourhood of K . A diagonal argument delivers a flat

$$\Pi_\infty := \lim_{n \text{ in a subsequence}} s_n \Pi$$

with $x \in \Pi_\infty$ such that for every r and every n large enough (depending on r) the ball of radius r and center x in Π coincide with the ball of radius r and center $s_n^{-1}x$ in Π_n . This shows that Π_∞ is the \diamond -flat.

This covers cases a), d), e), f) and C) and D) of the classification. The same argument works if Π contains \diamond -strips of arbitrary large height, showing that case B) of the classification leads to either a \diamond -flat, or Π a uniformly bounded flat of type $1 \times n$ (including $n = 2$). This covers cases b) and B).

In cases c) and g) it is easy to choose x_n so that Π_∞ is the alternating 2-strip/ \diamond -strip flat where the \diamond -strip has height 1. Finally in case A), either the height is uniformly bounded, or we can choose x_n so that Π_∞ is the \diamond -flat. \square

Lemma 27. *Suppose that X is a ring complex X of type $\text{Aut}(F_2)$ and fix a proper action $\Gamma \curvearrowright X$ with X/Γ compact. Assume furthermore that*

- X is acylindrical and uniformly locally finite
- X contains a flat that contains infinitely many parallel copies of a simplicial strip S
- S is uniquely L -embeddable in X for some non empty compact simplicial set L
- there are infinitely many distinct simplicial translates of L in S .

Then $\mathbb{Z}^2 \hookrightarrow \Gamma$.

Proof. Let S denote the strip given with simplicial set $L \subset S$.

Let Π be a flat in X that contains infinitely many parallel copies of S , say

$$f_k: S \hookrightarrow X \text{ with } \text{ran } f_k \parallel \text{ran } f_l, \quad f_k \neq f_l \quad (k \neq l \geq 1)$$

Note that $f_k \neq f_l$ on L if $k \neq l$.

Since X/Γ is compact there exist elements $s_k \in \Gamma$ such that

$$s_k \circ f_k = s_1 \circ f_1 \text{ on } L$$

for infinitely many $k \geq 1$.

Since S is uniquely L -extendable we have that

$$s_k \circ f_k = s_1 \circ f_1 \text{ on } S$$

for infinitely many $k \geq 1$.

Choose k large enough so that $\text{ran } f_k \cap \text{ran } f_1 = \emptyset$ and write $f = f_1$, $f' = f_k$ and $s = s_k^{-1} \circ s_1$ so that $f' = s \circ f$.

By assumption we can find infinitely many distinct translates of L in S . Let $g_n: S \rightarrow S$ denote the corresponding translations ($n \geq 1$). Since X/Γ is compact, there exist an index $n_0 \geq 1$ and elements $t_n \in \Gamma$ such that

$$t_n \circ f \circ g_n = f \circ g_{n_0} \text{ on } L$$

for infinitely many $n \geq 1$.

Since S is uniquely L -extendable we have that

$$t_n \circ f \circ g_n = f \circ g_{n_0} \text{ on } S$$

for infinitely many $n \geq 1$.

Choose n large enough so that $f \circ g_n(L) \cap f \circ g_1(L) = \emptyset$ and write $g = g_{n_0} \circ g_n^{-1}$ and $t = t_n$ so that we have the intertwining relation

$$t \circ f = f \circ g$$

on S . Note that g is a simplicial translation. Furthermore $t^m \circ f = f \circ g^m$ for every $m \in \mathbb{Z}$.

Denote by $f \vee f': [a, b] \times \mathbb{R} \rightarrow \Pi \subset X$ the simplicial strip lying “between f and f' ” in Π (whose range is the convex closure of both strips in Π). Thus there is $a < a' < b' < b$ such that S is simplicially isomorphic to both $[a, a'] \times \mathbb{R}$ and $[b', b] \times \mathbb{R}$, and the restriction of $f \vee f'$ to these strips is f and f' , respectively.

Since X is locally finite, there are only finitely many strips in X isometric to $\text{ran } f \vee f'$ containing $\text{ran } f$. Therefore there exists m and a translation $h: [a, b] \times \mathbb{R} \rightarrow [a, b] \times \mathbb{R}$ such that

$$t^m \circ (f \vee f') = (f \vee f') \circ h.$$

Note that h coincide with g^m on $[a, a'] \times \mathbb{R}$, so we will replace t with t^m to obtain the relations

$$t \circ f = f \circ h \text{ and } t \circ (f \vee f') = (f \vee f') \circ h.$$

on $S \simeq [a, a'] \times \mathbb{R}$ and $[a, b] \times \mathbb{R}$. This implies that

$$t^q \circ f = f \circ h^q \text{ and } t^q \circ (f \vee f') = (f \vee f') \circ h^q.$$

on $S \simeq [a, a'] \times \mathbb{R}$ and $[a, b] \times \mathbb{R}$, respectively, for every $q \in \mathbb{Z}$, and in particular that

$$t^q \circ f' = f' \circ h^q$$

on $S \simeq [b', b] \times \mathbb{R}$, for every $q \in \mathbb{Z}$.

Then

$$t^q \circ s \circ f = t^q \circ f' = f' \circ h^q = s \circ f \circ h^q = s \circ t^q \circ f.$$

Choose a point $x \in \text{ran } f$. The relation shows that for every $q \in \mathbb{Z}$ the commutator

$$[t^q, s] \in \text{Stab}_\Gamma(x),$$

which is a finite group by assumption. Therefore we can find $q \neq q'$ such that

$$[t^q, s] = [t^{q'}, s]$$

so

$$[t^{q-q'}, s] = e.$$

Since

$$t^q \circ f = f \circ h^q$$

for every q , it follows that t has infinite order in Γ .

Assume towards a contradiction that s has finite order in Γ . Then the subset of X

$$\bigcup_{k=0}^{\text{order of } s} \text{ran } s^k \circ (f \vee f')$$

is a simplicial cylinder in X , whose existence is precluded by assumption.

Thus, $\langle t^{q-q'}, s \rangle \simeq \mathbb{Z}^2$ in Γ . □

Remark 28. The space of marked flats $\mathbf{Flats}(X)$ is the set

$$\mathbf{Flats}(X) = \{\mathbb{R}^2 \hookrightarrow X\}$$

of all isometric embeddings $\mathbb{R}^2 \hookrightarrow X$. This is a classical object associated with X ; it has been studied in particular by Pansu and is useful when studying the rigidity of symmetric spaces (cf. [8, §14]).

The space $\mathbf{Flats}(X)$ has two commuting actions of \mathbb{R}^2 and Γ

$$t \cdot f := f \circ t^{-1} \text{ and } s \cdot f := s \circ f$$

respectively at the source and the range, for t a translation of \mathbb{R}^2 , $s \in \Gamma$, and $f: \mathbb{R}^2 \hookrightarrow X$ an embedding.

If X is locally finite and X/Γ is compact then $\mathbf{Flats}(X)/\Gamma$ is a compact space. It is not empty if X is not hyperbolic, and it is endowed with a structure of a lamination whose leaves are associated with flat planes in X and given by the action of \mathbb{R}^2 .

If X is a ring complex of type $\text{Aut}(F_2)$ we have an obvious map

$$\mathbf{Flats}(X) \rightarrow \mathbf{Puzzles}(\text{Aut}(F_2))$$

which is continuous and open.

Note that in the case of ring complexes of type $\text{Aut}(F_2)$, the subset $\mathbf{Flats}_x(X)$ corresponding to flats in X of type $x = \text{a), b), \dots$ is closed whenever flats of type x are geometrically periodic (i.e. the simplicial isometry group acts with compact quotient). This holds for flats of type a) or b), for example.

A ring complex X is said to be *full* if the map $\mathbf{Flats}(X) \rightarrow \mathbf{Puzzles}(T)$ is surjective with respect to $T = T(X)$, where $T(X)$ denotes the type of X (minimal sets of shapes and rings in X).

This notion of “fullness” can be thought of as a weakening of the idea of “space of maximal rank” from rank interpolation.

For example the Brady complex is full, and we think of it as a space of maximal rank among ring complexes of type $\text{Aut}(F_2)$. It plays the role of Euclidean buildings

in [4], which are the prototypical “spaces of maximal rank” (in their respective classes, namely among spaces of type \tilde{A}_2 , of type \tilde{B}_2 or of type \tilde{G}_2).

Observe also that the classification of Euclidean buildings into types is primarily a classification of their apartments (flats) and associated Coxeter groups, which in spirit is not very far from the definition of ring complexes being “of type $\text{Aut}(F_2)$ ” described in the present paper.

4. RANDOMIZING $\text{Aut}(F_2)$

Let $\Gamma_p \leq \Gamma_0 := \text{Aut}(F_2)$ be finite index subgroups and consider the deterministic data $(\Gamma_0, \{\Gamma_p\}, X_0)$ in the sense of [4], where X_0 is the Brady space.

We only discuss the density model. Results can also be established in the bounded model (for example).

Recall that if \mathbb{P}_p denotes a process for selecting finite random subsets of elements in the set

$$\mathcal{C}_p := \{\Gamma_p\text{-orbits of faces in } X_0\}$$

then the density model can be defined in the following way, given a parameter $\delta \in (0, 1)$.

Definition 29. The *density model* over $(\Gamma_0, \{\Gamma_p\}, X_0)$ is associated with

$\mathbb{P}_p :=$ “choose $|\mathcal{C}_p|^\delta$ chambers in \mathcal{C}_p , uniformly and independently at random”.

We have the following phase transition at $\delta = \frac{2}{3}$.

Proposition 30 (See [4, §5]). *If Γ denotes the random group in the density model over $(\Gamma_0, \{\Gamma_p\}, X_0)$ then*

- (1) *if $\delta < 2/3$ then Γ acts properly on a CAT(0) space X without boundary such that X/Γ compact*
- (2) *if $\delta > 2/3$ then Γ splits off a free factor isomorphic to a free group F_n on n generators.*

Lemma 31. *The random group in the density model over $(\Gamma_0, \{\Gamma_p\}, X_0)$ acts on a ring complex of type T .*

Proof. By construction the random group Γ at arbitrary density acts on a space X which is locally isometric to a subspace of X_0 and therefore the links of X embed L . Note that if $1/2 < \delta < 2/3$ the contraction of free faces removes free paths in the link, and if $\delta > 2/3$ the links are not necessarily connected. The shape set of X is also a subset of the shape set of X_0 , so X is of type T . This is true more generally of the α -model [4, §4]. \square

The following result is an application of Theorem 25.

Theorem 32. *For arbitrary deterministic data $(\Gamma_0, \{\Gamma_p\}, X_0)$ with $\Gamma_0 = \text{Aut}(F_2)$, Γ_p are finite index subgroups, and X_0 denotes the brady complex, the random group in the density model over $(\Gamma_0, \{\Gamma_p\}, X_0)$ satisfies the flat closing conjecture.*

This may be compared to the randomized lattices in nonarchimedean Lie groups [4], for which the corresponding statement is open.

Proof. Let $\Gamma \curvearrowright X$ denote the action of the random group on its associated random classifying space. Since X is nonpositively curved [4] it is acylindrical. In order to apply Theorem 25 we have to check the unique extension property for X . This

follows from a generalization of [4, Lemma 11] to strips, which are simply connected (see also [1, Lemma 13]). Namely if $A \subset \mathcal{C}_p$ is an arbitrary subset and, in the notation of [4, Lemma 11], if

$$S \subset X_A := X_0 - \bigcup_{C \in A} \overset{\circ}{C}$$

is a strip, then the map $j : S \rightarrow \Pi \subset X_A$ lifts to unique map $\tilde{j} : S \rightarrow \tilde{\Pi} \subset \tilde{X}_A$ through any point $\tilde{j}(s) = \tilde{x} \in \tilde{X}_A$, $s \in S$ such that $\pi_A(\tilde{x}) = j(s) =: x$ and we have a commutative diagram

$$\begin{array}{ccc} & & \tilde{X}_A \\ & \nearrow \tilde{j} & \downarrow \pi_A \\ S & \xrightarrow{j} & X_A \end{array}$$

We prove that the strip S is uniquely L -embeddable in X if it is so in the Brady complex X_0 . Since both the \diamond -strip and the Δ -strip are uniquely embeddable in X_0 (which is an easy exercise), the result will follow.

Consider two simplicial embeddings $f, f' : S \hookrightarrow \tilde{X}_A$ and a translation $g : S \rightarrow S$

$$f' = f \circ g \text{ on } L.$$

We have to show that $f' = f \circ g$ on S .

Let $j := \pi_A \circ f \circ g$ and $j' := \pi_A \circ f'$. These are two embeddings $S \rightarrow X_A \subset X_0$ which coincide on L . By the the unique strip extension property for S in X_0 , we have that $j = j'$ on S .

Let $s \in L$ and $\tilde{x} = f'(s) = f \circ g(s) \in f'(L) = f \circ g(L)$. There is a unique lift \tilde{j} of j such that $\tilde{j}(s) = \tilde{x}$ and a unique lift \tilde{j}' of j' such that $\tilde{j}'(s) = \tilde{x}$ so $\tilde{j} = \tilde{j}'$ on S (since $j = j'$ on S). Applying uniqueness again $f \circ g = \tilde{j}$ on S and $f' = \tilde{j}'$ on S . So $f' = f \circ g$ on S .

This establishes the unique extension property for \tilde{X}_A with respect to an arbitrary — in particular for a random — subset $A \subset \mathcal{C}_p$. \square

The $\text{Aut}(F_2)$ models differ from the nonarchimedean lattices models of [4] in many respects. What we prove below is that for suitable choices of the deterministic data, the presence of periodic flats persists at *arbitrary* densities $\delta \in (0, 1)$. This relies on the fact that the following “small tori property” can be achieved for $\text{Aut}(F_2)$ models, for example for deterministic data associated with the symmetric group, using superexponential growth.

Definition 33. We say that the deterministic data $(\Gamma_0, \{\Gamma_p\}, X_0)$ has the *small tori property* (in the density model) if for any $p \geq 1$ there exists an embedding $\mathbb{Z}^2 \hookrightarrow \Gamma_0$ and a simplicial embedding $\mathbb{R}^2 \hookrightarrow X_0$ with $\mathbb{Z}^2 \curvearrowright \mathbb{R}^2$ (freely and cocompactly) such that

$$\forall \delta > 0, |(\mathbb{R}^2)^{(2)} / (\mathbb{Z}^2 \cap \Gamma_p)| \leq O(|X^{(2)} / \Gamma_p|^\delta).$$

where $^{(2)}$ indicates 2-faces.

This is easily seen to be an algebraic condition on the growth of the finite index subgroups relative to the embedding of \mathbb{Z}^2 , at least if X is a CAT(0) space (see the proof of Theorem 35).

The persistence of periodic flat planes at arbitrary density will follow from:

Lemma 34. *If $(\Gamma_0, \{\Gamma_p\}, X_0)$ has the small tori property, where X_0 is a CAT(0) space, then \mathbb{Z}^2 embeds in the random group at arbitrary density $\delta \in (0, 1)$ with overwhelming probability.*

The proof is extracted from [4, §6].

Proof. Let $\delta \in (0, 1)$ be given. Let \mathcal{C}_p be the set of Γ_p -orbits of 2-cells in X_0 (which can be triangles or lozenges) and let F_p be the set of $\mathbb{Z}^2 \cap \Gamma_p$ orbits of 2-cells in \mathbb{R}^2 viewed as a flat plane in X_0 .

Let $\alpha < 1 - \delta$. By the small tori property we have

$$|F_p| \leq O(|X^{(2)}|/\Gamma_p)^\alpha.$$

Write F'_p for the image of F_p in \mathcal{C}_p so

$$|F'_p| \leq C|\mathcal{C}_p|^\alpha$$

for some constant $C > 0$.

Consider a sequence Y_1, Y_2, Y_3, \dots of i.i.d. random variables with values in \mathcal{C}_p . The probability $\mathbb{P}(E_p)$ of the event:

$$E_p = \{Y_i \notin F'_p, \forall i \leq |\mathcal{C}_p|^\delta\}$$

satisfies

$$\mathbb{P}(E_p) \geq \left(1 - \frac{|F'_p|}{|\mathcal{C}_p|}\right)^{|\mathcal{C}_p|^\delta} \geq e^{-2|\mathcal{C}_p|^{\delta-1}|F'_p|} \geq e^{-2C|\mathcal{C}_p|^{\delta-1+\alpha}}$$

so $\mathbb{P}(E_p) \rightarrow 1$. Applying Lemma 11 in [4] we have that $\mathbb{Z}^2 \hookrightarrow \Gamma$ with overwhelming probability at density δ . \square

This can be applied to the case of $\text{Aut}(F_2)$ in the following way. Write $F_2 = \langle x, y \rangle$ and define automorphisms $\alpha, \beta \in \text{Aut}(F_2)$

$$\alpha: x \rightarrow x, y \rightarrow x^2y, \quad \beta: x \rightarrow xy^2, y \rightarrow y$$

so

$$\bar{\alpha} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \text{ and } \bar{\beta} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$$

where $\bar{\alpha}, \bar{\beta}$ are the reduction relative to the short exact sequence

$$\text{IA}_2 \hookrightarrow \text{Aut}(F_2) \xrightarrow{\bar{\cdot}} \text{GL}_2(\mathbb{Z}).$$

$\text{IA}_2 = \text{Inn}(F_2) = \langle \alpha_0, \beta_0 \rangle \simeq F_2$ is the subgroup of inner automorphisms:

$$\alpha_0: x \rightarrow x, y \rightarrow xyx^{-1}, \quad \beta_0: x \rightarrow yxy^{-1}, y \rightarrow y.$$

It is well-known that $\langle \bar{\alpha}, \bar{\beta} \rangle \simeq F_2$ so we have a split exact sequence

$$\text{IA}_2 \hookrightarrow \text{IA}_2 \rtimes \langle \alpha, \beta \rangle \twoheadrightarrow \langle \bar{\alpha}, \bar{\beta} \rangle.$$

Theorem 35 (Periodic flat planes at arbitrary density). *Let $A_p \triangleleft F_2$ be a characteristic subgroup of F_2 and let $B_p \triangleleft F_2$ be a normal subgroup such that F_2/B_p is the group S_p of permutations of the set with p elements. Suppose that $[F_2 : A_p] \leq e^{O(p)}$. If the deterministic data $(\Gamma_0, \{\Gamma_p\}, X_0)$ satisfy:*

- $\Gamma_0 = \text{Aut}(F_2)$
- X_0 is the Brady complex
- $\Gamma_1 = \text{IA}_2 \rtimes \langle \alpha, \beta \rangle$ (a subgroup of index 24 in Γ_0)

- for $p \geq 2$, we have $\Gamma_p := A_p \times B_p$ where we view A_p as a subgroup of IA_2 and B_p as a subgroup of $\langle \bar{\alpha}, \bar{\beta} \rangle$.

Then with overwhelming probability $\mathbb{Z}^2 \hookrightarrow \Gamma :=$ the random group at density $\delta \in (0, 1)$.

Proof. Since X_0 is a CAT(0) space the triple $(\Gamma_0, \{\Gamma_p\}, X_0)$ has the small tori property if and only if there exists an embedding $\mathbb{Z}^2 \hookrightarrow \Gamma$ such that

$$\forall \delta > 0, |\mathbb{Z}^2 / (\mathbb{Z}^2 \cap \Gamma_p)| \leq O(|\Gamma / \Gamma_p|^\delta).$$

We have

$$|\langle \alpha_0 \rangle / (\langle \alpha_0 \rangle \cap A_p)| \leq e^{O(p)}, \text{ and similarly } |\langle \beta_0 \rangle / (\langle \beta_0 \rangle \cap A_p)| \leq e^{O(p)}$$

by assumption and

$$|\langle \alpha \rangle / (\langle \alpha \rangle \cap B_p)| \leq g(p), \text{ and similarly } |\langle \beta \rangle / (\langle \beta \rangle \cap B_p)| \leq g(p)$$

where

$$g(p) := \text{maximal order of an element in } S_p$$

is the Landau function (see Wikipedia). Note that the value $g(p)$ can be indeed achieved in our situation [6, §2.63].

Landau has proved that

$$\lim_{p \rightarrow \infty} \frac{\ln g(p)}{\sqrt{p \ln p}} = 1$$

therefore

$$|\langle \alpha \rangle / (\langle \alpha \rangle \cap B_p)| \leq e^{O(p)}, \text{ and similarly } |\langle \beta \rangle / (\langle \beta \rangle \cap B_p)| \leq e^{O(p)}.$$

If $\Lambda \simeq \mathbb{Z}^2$ denotes the subgroup of $\text{Aut}(F_2)$ generated either by α_0, α or by β_0, β then we have

$$|\Lambda / (\Lambda \cap \Gamma_p)| \leq e^{O(p)}.$$

On the other hand by Stirling's approximation

$$|\Gamma / \Gamma_p| \geq p! \geq \left(\frac{p}{e}\right)^p.$$

This shows that

$$\forall \delta > 0, |\Lambda / (\Lambda \cap \Gamma_p)| \leq O(|\Gamma / \Gamma_p|^\delta).$$

Then Lemma 34 applies. \square

Remark 36. 1) $\text{Aut}(F_2)$ has many interesting finite quotients leading to interesting phase transitions in the corresponding density model. We hope to come back to this elsewhere.


2) The variation of the small tori property which is implicit in [4] is the assertion that there is $\alpha \geq 0$ such that for any $p \geq 1$ there are $\mathbb{Z}^2 \hookrightarrow \Gamma_0$ and $\mathbb{R}^2 \hookrightarrow X_0$ with $\mathbb{Z}^2 \curvearrowright \mathbb{R}^2$ freely and cocompactly such that

$$\forall \delta > \alpha, |(\mathbb{R}^2)^{(2)} / (\mathbb{Z}^2 \cap \Gamma_p)| \leq O(|X^{(2)} / \Gamma_p|^\delta).$$

5. THE EXPLICIT CONSTRUCTION

The construction provides a non positively curved simplicial complex X and a group Γ acting freely on X with 3 orbits of vertices.

The link at every vertex is a “double” of the link of the Brady complex as explicited on the following drawing (in red and black). See [5, Fig. 6] for the link of the Brady complex itself.



The figure shows X/Γ . The 1-skeleton (in green) of X/Γ supports the following set of faces:

- triangles: $\bullet_1 \bullet_2 \bullet_3$ where $\bullet = a, b, c, d, e, f, x, y$
- lozenges: $\bullet_1 \bullet'_2 \bullet'_3 \bullet_2$, where $\bullet = a, b, c, d, e, f$ and $x_1 d_3 y'_1 b'_1, x_1 c'_3 y'_1 a_1, x_2 f_3 y'_2 c'_1, x_2 e'_3 y'_2 d_1, x_3 a_3 y'_3 e'_1, x_3 b'_3 y'_3 f_1$.

It is easy to see that the complex X thus constructed is a ring complex of type $\text{Aut}(F_2)$. Further inspection shows that X fails to be hyperbolic and has the isolated flats property. Every link contains 8 cycles of length 2π and the corresponding puzzles are of type A.

This group appears to be an intriguing sort of “double” of $\text{Aut}(F_2)$.

REFERENCES

- [1] Barré S., Pichot M., Intermediate rank and property RD, arXiv:0710.1514.
- [2] Barré S., Pichot M., The 4-string braid group has property RD and exponential mesoscopic rank, arXiv:0809.0645.
- [3] Barré S., Pichot M., Removing chambers in Bruhat-Tits buildings, arxiv 1003.4614.
- [4] Barré S., Pichot M., Random groups and nonarchimedean lattices, arXiv:1308.2315

- [5] Crisp, J., Paoluzzi, L. On the classification of $CAT(0)$ structures for the 4-string braid group. *Michigan Math. J.*, 53 (1), 133–163 (2005).
- [6] Dixon J., *Problems in group theory*, Dover publications, 1973.
- [7] Gromov, M., *Geometric group theory. Vol. 2. Proceedings of the symposium held at Sussex University, Sussex, July 1991.* Edited by Graham A. Niblo and Martin A. Roller. London Mathematical Society Lecture Note Series, 182. Cambridge University Press, Cambridge, 1993.
- [8] Mostow, G. D. Strong rigidity of locally symmetric spaces. *Annals of Mathematics Studies*, No. 78. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1973.