



HAL
open science

Reward-related brain activity and behavior are associated with peripheral ghrelin levels in obesity

Volodymyr Bogdanov, Olena Bogdanova, Sandra Dexpert, Ines Delgado, Helen Beyer, Agnès Aubert, Bixente Dilharreguy, Cédric Beau, Damien Forestier, Patrick Ledaguenel, et al.

► To cite this version:

Volodymyr Bogdanov, Olena Bogdanova, Sandra Dexpert, Ines Delgado, Helen Beyer, et al.. Reward-related brain activity and behavior are associated with peripheral ghrelin levels in obesity. *Psychoneuroendocrinology*, 2020, 112, pp.1-8. 10.1016/j.psyneuen.2019.104520 . hal-02517346

HAL Id: hal-02517346

<https://hal.science/hal-02517346>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Reward-related brain activity and behavior are associated with peripheral ghrelin levels in obesity**

2

3 Volodymyr B. Bogdanov^{1,2#}, Olena V. Bogdanova³, Sandra Dexpert¹, Ines Delgado¹, Helen

4 Beyer¹, Agnès Aubert¹, Bixente Dilharreguy⁴, Cédric Beau⁵, Damien Forestier⁵, Patrick

5 Ledaguenel⁵, Eric Magne⁵, Bruno Aouizerate¹, Sophie Layé¹, Guillaume Ferreira¹, Jennifer Felger⁶,

6 Giuseppe Pagnoni^{7,8}, Lucile Capuron^{1#}

7

8 ¹ Univ. Bordeaux, INRA, Bordeaux INP, NutriNeuro, UMR 1286, F-33000, Bordeaux, France;

9 ² Univ. Lyon, Ecole Nationale des Travaux Publics de l'Etat, Laboratoire Génie Civil et Bâtiment, F-

10 69518 Vaulx-en-Velin, France ;

11 ³ INSERM U1028 - CNRS UMR5292, 16 avenue Doyen Lépine, F-69676, Bron, France;

12 ⁴ Univ. Bordeaux, INCIA, CNRS, UMR 5287, F-33076 Bordeaux, France;

13 ⁵ Digestive and Parietal Surgery, Clinique Tivoli, F-33000, Bordeaux, and Clinique Jean Villar, F-33520,

14 Bruges France;

15 ⁶ Dpt of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA 30322,

16 USA;

17 ⁷ Dept of Neural, Biomedical, and Metabolic Sciences, University of Modena and Reggio Emilia, I-

18 41125, Modena, Italy;

19 ⁸ Center for Neuroscience and Neurotechnology, University of Modena and Reggio Emilia, I-41125,

20 Modena, Italy.

21

22

23 **# Corresponding authors:**

24 - Volodymyr B. Bogdanov, PhD, Univ. Lyon, Ecole Nationale des Travaux Publics de l'Etat, Laboratoire

25 Génie Civil et Bâtiment, 3 rue Maurice Audin, F-69518 Vaulx-en-Velin, France; E-mail:

26 vlabogd@yahoo.com

27 - Lucile Capuron, PhD, INRA 1286, NutriNeuro, University of Bordeaux, 146 rue Leo Saignat, F-33076

28 Bordeaux, France; Tel. +33 (0)557571233, Fax. +33 (0) 557571227; E-mail: lucile.capuron@inra.fr

29

30 **Running Title.** Ghrelin-related reward activity in obesity

31

32 **Conflict of Interest.** The authors declare no conflict of interest in relation to the work described.

33

34 **Funding/Support:** This work was supported by the French National Research Agency, ANR (ANR-11-
35 JSV1-0006, Dr Capuron) and funds from the French National Institute for Agricultural Research (INRA).

36

37

38

39 **ABSTRACT**

40 **Background/Objectives:** While excessive food consumption represents a key factor in the development
41 of obesity, the underlying mechanisms are still unclear. Ghrelin, a gut-brain hormone involved in the
42 regulation of appetite, is impaired in obesity. In addition to its role in eating behavior, this hormone was
43 shown to affect brain regions controlling reward, including the striatum and prefrontal cortex, and there is
44 strong evidence of impaired reward processing in obesity. The present study investigated the possibility
45 that disrupted reward-related brain activity in obesity relates to ghrelin deficiency.

46 **Subjects/Methods:** Fifteen severely obese subjects (BMI > 35 kg/m²) and fifteen healthy non-obese
47 control subjects (BMI < 30 kg/m²) were recruited. A guessing-task paradigm, previously shown to
48 activate the ventral striatum, was used to assess reward-related brain neural activity by functional
49 magnetic resonance imaging (fMRI). Fasting blood samples were collected for the measurement of
50 circulating ghrelin.

51 **Results:** Significant activations in the ventral striatum, ventromedial prefrontal cortex and extrastriate
52 visual cortex were elicited by the fMRI task in both obese and control subjects. In addition, greater
53 reward-related activations were present in the dorsolateral prefrontal cortex, and precuneus/posterior
54 cingulate of obese subjects compared to controls. Obese subjects exhibited longer choice times after
55 repeated reward and lower circulating ghrelin levels than lean controls. Reduced ghrelin levels
56 significantly predicted slower post-reward choices and reward-related hyperactivity in dorsolateral
57 prefrontal cortices in obese subjects.

58 **Conclusion:** This study provides evidence of association between circulating ghrelin and reward-related
59 brain activity in obesity and encourages further exploration of the role of ghrelin system in altered eating
60 behavior in obesity.

61

62 *Key words: Obesity, Reward, Behavior, fMRI, Ghrelin*

63 1. INTRODUCTION

64

65 Abnormal feeding behavior resulting in weight gain and obesity has been linked to alterations in
66 reward processing (Burger and Berner, 2014). In support of this, impaired reward-related brain responses
67 have been repeatedly documented in overweight/obese subjects (Carnell et al., 2012; Pursey et al., 2014;
68 Puzziferri et al., 2016). These alterations were found not only in food-related paradigms but also in more
69 generalized experimental conditions, including monetary incentive tasks (Balodis et al., 2013). Monetary
70 delay discounting, which is a bias towards smaller but earlier rewards, is greater in obese subjects
71 pointing to an abnormal reward expectation and decision making in obesity (Price et al., 2016). These
72 effects may be related to a deregulation of hormones that control reward-directed and food-consuming
73 behaviors. In this perspective, a promising target for obesity research is the ghrelin-regulating system,
74 which is involved not only in appetitive behavior but also in higher cognitive functions (Anderberg et al.,
75 2016; Ralevski et al., 2018).

76 Ghrelin is a peptide that is mainly produced by neuroendocrine cells in the stomach mucosa.
77 Secretion of ghrelin has a fluctuating character depending on circadian rhythms and food anticipation
78 (Natalucci et al., 2005). Immediately after food consumption, ghrelin levels fall (English et al., 2002) and
79 then gradually increase to reach a peak before the next meal is initiated. Once secreted, ghrelin is able to
80 act on numerous peripheral tissues and the brain (Muller et al., 2015). Peripheral ghrelin can affect brain
81 function via different pathways (Howick et al., 2017), including vagal or brain stem stimulation (Date,
82 2012), blood stream (Angelidis et al., 2010), passage across the blood-brain barrier via saturated transport
83 system (Solomou and Korbonits, 2014), and transport *via* the cerebrospinal fluid (Uriarte et al., 2018).
84 Ghrelin receptors are widely expressed in the brain, notably in the orexigenic neurons of hypothalamic
85 nuclei (Muller et al., 2015) and in the hippocampus, pituitary, dentate gyrus, substantia nigra,
86 dorsal/median raphe nuclei and ventral tegmental area (Andrews, 2011). Altogether, they contribute to
87 the regulation of appetite, energy balance, olfaction, but also memory/learning, stress response, mood,
88 and reward control (Abizaid, 2019; Goldstone et al., 2014; Howick et al., 2017; Malik et al., 2008; Perello
89 and Dickson, 2015).

90 Studies in animals have shown that ghrelin activates the reward circuitry comprising of the
91 ventral tegmental area, lateral hypothalamus and nucleus accumbens (Revitsky and Klein, 2013). At the
92 clinical level, reductions in dorsolateral prefrontal cortex (dlPFC) activity and decreased self-control,
93 supposed to be related to uncontrolled overeating (Holsen et al., 2012), have been reported in overweight
94 patients with Prader-Willi syndrome (PWS) who also exhibit increased ghrelin levels. In healthy subjects,
95 increased ghrelin levels have been related to increased sensitivity to reward and impulsivity/lower self-
96 control (Ralevski et al., 2018). Deregulation of the ghrelin system may also contribute to the development
97 of obesity (Abizaid, 2019). In obese subjects, not only fasting ghrelin levels are reduced proportionally to
98 body mass index (BMI) (Tschop et al., 2001; Nonogaki, 2008), but also alterations in the circadian
99 rhythm and postprandial inhibition of ghrelin secretion are reported (Yildiz BO et al., 2004). The patterns
100 of phasic and tonic ghrelin regulations are altered (Foster et al., 2007) and ghrelin resistance is proposed
101 as one of the major mechanisms of ghrelin system deficiency in obesity (Zigman et al., 2016). Central
102 action of ghrelin in obesity may be also impaired due to different facets of ghrelin resistance (Zigman et
103 al., 2016). Interestingly, normalization of blood levels of ghrelin after the weight loss is associated with
104 improved brain sensitivity to ghrelin (Briggs et al., 2013).

105 The association of the ghrelin system with both reward response and appetite control/metabolic
106 regulation suggests the potential role of decreased ghrelin levels in the development of brain reward
107 system alterations and disrupted eating behaviors likely to promote the development of obesity. While
108 altered reward-related neural activation in obesity has been frequently associated with changes in other
109 energy balance regulating hormones, including leptin and insulin (Baicy et al., 2007; Farooqi et al., 2007;
110 Kullmann et al., 2012), no studies have investigated the potential link between obesity-related ghrelin
111 deficiency and disruptions in the brain reward system. The present study addressed this issue in a sample
112 of severely obese subjects compared to healthy non-obese controls, using functional magnetic resonance
113 imaging (fMRI) during a non-food related task known to induce reward-related cerebral activity.
114 Behavioral responses and brain processing of reward outcome were investigated and their relationship
115 with individual levels of ghrelin was assessed.

116

117 2. METHODS

118 2.1. Subjects

119 Fifteen severely obese subjects (BMI > 35 kg/m²; 3 males and 12 females) and fifteen healthy
120 non-obese control subjects (BMI < 30 kg/m²; 2 males and 13 females) were included. All subjects were
121 right-handed adults, free of any treatment with psychotropic effects, and had not eaten for the last two
122 hours before the fMRI session. Additional exclusion criteria were age > 65 years old, history of brain
123 damage or neurological disease, acute or chronic inflammatory disorders (other than obesity and obesity-
124 related conditions), severe or uncontrolled medical conditions, psychiatric disorders, metallic implants
125 and eating disorders (in non-obese control subjects). All subjects provided written informed consent and
126 received compensation for their participation in the study. The study was approved by the Committee for
127 the Protection of Persons (CPP) of Bordeaux.

128

129 2.2. Guessing task

130 We used a guessing-task paradigm that was previously shown to activate the ventral striatum,
131 involved in reward processing (Capuron et al., 2012; Reuter et al., 2005). Subjects were presented with
132 two hidden cards on the left and right sides of the screen and were asked to guess which of the two cards
133 was “red” by pressing on a keypad with the index (left key) or middle finger (right key) (**Figure 1A, B**).
134 The time window for the choice was fixed at 2s, after which the selected card was turned over and the
135 outcome was displayed on the screen for another 2s. If the card was “red”, the subject gained one Euro, if
136 “black” one Euro was lost. The sequence of gains and losses was similar for all subjects. Unbeknownst to
137 participants, the outcomes followed a noisy sinusoid with a slight upward trend (**Figure 1A**), as described
138 elsewhere (Reuter et al., 2005). The net score, which was initially equal to 15 Euros, was always shown
139 on the screen and updated each time the chosen card was revealed. The choice time was recorded for each
140 trial. In total, there were 100 trials, and the total duration of the task was 6 min 40 sec. In contrast to
141 previous studies using this paradigm, participants were informed that the task should be considered as a
142 game and the amount of their compensation (related to their participation in the study) would not depend

143 of their gain or lost; furthermore, the number of trials was reduced by a factor of three (Reuter et al.,
144 2005). These methodological adaptations did not impede significant ventral striatum reward response, as
145 shown in **Figure 2**. All stimuli presentations and recordings of reaction time and response keys were
146 performed with E-Prime 2.0 software (Psychology Software Tools, Sharpsburg, PA, USA).

147 Before the acquisition started, a practice session was performed outside the scanner room on a
148 laptop computer under the supervision of the experimenter. An additional practice session was performed
149 in the scanner using the MRI-compatible keypad (4-key response box). After practice, participants
150 performed the task during the scanning session. The task was presented on a screen placed in front of the
151 scanner bed and was monitored through a computer located outside the scanner room. Participants were
152 able to see the screen through a mirror attached to the head coil. Choice time (CT), i.e. the time elapsed
153 from the presentation of the cards on the screen to the volunteer's button press signaling the choice of one
154 of the two hidden decks (left or right), was recorded as a behavioral parameter. We were particularly
155 interested in decision-making processes related to further reward acquisition *after* a reward had already
156 been received, as this is an experimental situation that could provide a model for the prolonged
157 consumption of food in the obese. Accordingly, only CTs after winning trials were analyzed.
158 Furthermore, CTs were grouped into two subsets according to the event-sequence, as this is known to
159 affect cerebral responses in reward tasks (Akitsuki et al., 2003). When there was just one win before the
160 current trial, response latency was classified as CT after short reward (CT1). When there were two or
161 more wins in a row before the current trial, CT was considered as being within the context of long or
162 sustained reward (CT2).

163

164 **2.3. MRI acquisition**

165 Scanning was performed on a 3T scanner GE Discovery MR750 with a 32 channels head coil.
166 First, a high-resolution T1-weighted anatomical scan was acquired via a FSPGR sequence with the
167 following parameters: TR=7.896 ms, TE=3.216 ms, TI=400 ms, FA=10°, FoV=288 mm², 200 sagittal
168 slices with a 288x288 acquisition matrix, 1 mm³ isotropic resolution, total acquisition time: 3 min 43 s.
169 Then, two functional runs of T2*-weighted echo-planar imaging for measurement of the blood oxygen

170 level-dependent (BOLD) contrast were acquired, with the following parameters: TR=2000 ms, TE=30
171 ms, FA=80°, FOV=19.20 cm, 35 64×64 slices with a 3.5 mm thickness and a 0.5 mm between-slice gap
172 (effective voxel size = 3x3x4 mm), with asset factor 2. Slices were prescribed with a negative pitch angle
173 of 30 degrees relative to AC-PC horizontal plane, in order to minimize the ventro-medial BOLD signal
174 dropout, as advised in (Weiskopf et al., 2007). During the first functional run (5 dummy scans + 264
175 dynamic scans; total duration 8 min 58 s), volunteers performed an attention task, whose results will be
176 discussed and reported elsewhere. During the second functional run (5 dummy scans + 204 dynamic
177 scans; total duration 6 min 58 s), subjects performed the guessing task described above.

178

179 **2.4. Imaging preprocessing**

180 MRI images were analyzed using the SPM software package (*SPM8 v4290; Wellcome*
181 *Department of Cognitive Neurology, Institute of Neurology, London, UK;*
182 *<http://www.fil.ion.ucl.ac.uk/spm>) implemented in MATLAB (Mathworks Inc., Sherborn, MA).*

183 For each subject, motion correction to the first functional scan was performed with a six-parameter rigid-
184 body transformation. No excessive motion was found using the criterion of >1 mm instantaneous shift.
185 The structural image was co-registered to the average of the motion-corrected functional images, and then
186 segmented into grey matter, white matter, and cerebrospinal fluid probability maps. The warping
187 parameters obtained from the segmentation algorithm were used to spatially normalize the structural
188 image to the Montreal Neurological Institute (MNI) standard brain space (output voxel size: 1x1x1 mm³).
189 These parameters were also applied to the functional images to bring them into the same MNI space
190 (output voxel size: 3x3x3 mm³), after which they were spatially smoothed with a Gaussian kernel with a
191 FWHM of 6 mm.

192

193 **2.5. Measurement of ghrelin circulating levels**

194 In all participants, overnight fasting blood samples were collected between 8 am and 10 am for
195 the measurement of circulating ghrelin (total). For logistic reasons, blood samples could not be collected

196 on the same day as the fMRI assessment in 60% of study participants. On average, these samples were
197 collected within the 4 weeks preceding the fMRI scan in the obese group, and within the week preceding
198 the scan in the group of healthy controls. After clotting, samples were centrifuged (1000g, 10 min, 4°C)
199 and sera were stored at -80°C until the assay. Serum total ghrelin concentration was determined by
200 enzyme-linked immunosorbent assay (ELISA), according to the manufacturer specifications (EZGRT-
201 89K, Millipore, Billerica, Massachusetts). Sensitivity was 50 pg/mL and intra- and inter-assay variability
202 were $\pm 1.26\%$ and $\pm 7.81\%$, respectively.

203

204 **2.6. Statistical analysis**

205 Participant characteristics, including age, gender, BMI, education level, and circulating levels of
206 ghrelin were compared between the two groups using two-sample *t* tests or Chi-square tests using
207 MATLAB (Mathworks Inc., Sherborn, MA). Two subjects from the control group and 1 subject from the
208 obese group did not properly follow instructions for the performance of the guessing task during the MRI
209 acquisition. Therefore, only data related to ghrelin levels and anatomical brain images were used in those
210 subjects.

211 Cerebral BOLD responses during the fMRI task were assessed using a general linear model
212 approach similar to the one described by Reuter and collaborators (Reuter et al., 2005). Briefly, for each
213 subject, we set up a linear model with two regressors of interest corresponding respectively to the
214 differential "*gain vs. loss*" response to the outcome (each outcome period modeled as 2 second-blocks),
215 and "*net score*" (modeled with 4 second-blocks, with a height tracking the current net gain), both time-
216 locked to the moment when the outcome was revealed (i.e., when the cards were turned up). The two
217 regressors were convolved with the SPM canonical hemodynamic response function. The motion
218 parameters estimated during the realignment phase were added as potential confounds to the model,
219 which also included an autoregressive component of order 1 plus white noise, to account for the presence
220 of autocorrelation in the residuals, and a basis set of discrete cosine functions modeling a high-pass filter
221 with a cut-off period of 240 sec. A random-effects group analysis was then performed using two-sample
222 F-test for identification of any significant "gain vs. loss" statistical effect in any of the two groups (obese

223 and/or control subjects). A similar analysis, but using a one-sample t-test, was conducted to assess the
224 effect of interest separately within the obese and the control groups. Finally, a two-sample t-test was
225 performed to compare the effect between the groups. An *undirected search* (whole-brain analysis) was
226 performed looking for significant voxels with a $p < 0.05$ family wise error rate (FWER), corrected for the
227 whole brain volume. For the between-group and single group cluster level inferences, cluster-forming
228 thresholds for the single-voxel significance were set at $p < 0.001$, while the cluster-level threshold was
229 0.05 FWER with a cluster size threshold of $k_E > 214$ voxels. We wanted to explore the link between
230 individual amplitudes of reward-related cerebral activity, ghrelin levels and choice times. Since the dIPFC
231 turned out to be more activated in the obese group compared to controls, we focused on this region of
232 interest (ROI), extracting the eigenvariates of parameter estimates of the “gain vs. loss” contrast using a
233 bilateral spherical ROI of 6 mm radius around the peak of the group effect in dIPFC. The resulting values
234 were then used to test the main and interaction effects between variables of interest: reward-related brain
235 activation, levels of ghrelin and behavior in groups of healthy volunteers and obese subjects with
236 ANCOVA analysis in MATLAB. We calculated the effect size and reported it as Cohen’s d for between-
237 group comparisons and the partial Eta Squared value, η^2_p , for ANCOVA analysis.

238

239

240 **3. RESULTS**

241 **3.1. Characteristics of study participants**

242 As shown in **Table 2**, there was no significant difference between obese subjects and non-obese
243 participants in terms of age, gender and education level. The average weight and BMI were respectively
244 114.7 kg and 41.2 kg/m² in obese subjects *versus* 60.5 kg and 21.8 kg/m² in non-obese controls (all $p <$
245 0.0001). Circulating ghrelin levels were significantly lower in obese subjects compared to non-obese
246 controls (375.2±47.6 pg/mL vs. 811.6±87.8 pg/mL; $df = 28$; $p < 0.001$, for both parametric t-test and non-
247 parametric Mann-Whitney test, Cohen’s $d = 1.6$).

248

249

250 **3.2. Neuroimaging results**

251 As expected, the fMRI task elicited significant activations in the ventral striatum (left and right
252 sides) (**Figure 2A**), the extrastriate visual cortex and the ventromedial prefrontal cortex (**Table 2**) in both
253 groups. While the peak voxels in the left and right ventral striatum showed a greater response in obese
254 subjects compared to controls (see bar plot in **Figure 2A**), the group difference was not significant when
255 controlling for the FWER across the whole-brain, or when using a small-volume-correction for the two
256 ventral striatum clusters. When looking at the reward-related activity separately for each group, the obese
257 subjects showed a larger extent of activation compared to controls (**Table 2 and Figure 2B**). A direct
258 statistical comparison of the reward-related effect in the two experimental groups showed a greater
259 activation of the left and right dlPFC and the precuneus/posterior cingulate in obese subjects compared to
260 controls (**Table 2 and Figure 3**).

261

262 **3.3. Behavioral performance during the fMRI task**

263 Choice times after one subsequent win (successful guess), CT1, and choice times after two or
264 more subsequent successful guesses, CT2, were compared between the two groups. No significant
265 difference was found in CT1 ($0.96s \pm 0.05s$ in obese subjects *vs.* $0.92s \pm 0.07s$ in controls). In contrast,
266 choice times after two or more successive winning trials, i.e. CT2, were significantly longer in obese
267 subjects compared to controls ($1.17s \pm 0.07s$ *vs.* $0.94s \pm 0.05s$, $df = 25$, $p < 0.05$, Cohen's $d = 0.8$) (**Figure**
268 **4A**).

269

270 **3.4. Relationships among reward-related brain activity, behavioral performance and ghrelin levels**

271 ANCOVAs were performed to test for main and interaction effects among the variables of
272 interest; i.e., reward-related brain activation, levels of ghrelin and behavior (choice time, CT), in obese
273 patients and controls. The first ANCOVA, performed on the individual values for the reward-related
274 dlPFC contrasts of parameter estimates (Win-Lose) with group as factor and ghrelin levels as covariate,
275 revealed a strong effect of group, which was not surprising given that the dlPFC ROI had been identified

276 from the Obese vs. Controls contrast of the fMRI group analysis. Our comparison of interest, however,
277 revealed a significant interaction between groups and ghrelin levels (**Figure 4B**; $F(1, 23)=5.17$, $p=0.033$),
278 indicating that reward-related dlPFC activation was more strongly related to lower ghrelin levels in the
279 group of obese subjects than in control subjects. A similar pattern was identified with the ANCOVA
280 performed on choice times, with group as factor and ghrelin levels as covariate. Here also the slope of
281 decreasing choice times with increasing ghrelin levels was steeper in obese subjects compared to controls,
282 indicating that slower choice behavior was more strongly related to reduced ghrelin levels in the group of
283 obese subjects (**Figure 4C**; Group X Ghrelin interaction: $F(1,23)=5.26$, $p=0.031$). Finally, the ANCOVA
284 performed on the choice times, with group as factor and the values of reward-related dlPFC contrast of
285 parameter estimates as covariate, showed a robust slowing of choice times with increasing reward-related
286 dlPFC activation (**Figure 4D**; covariate effect: $F(1,23)=9.39$, $p=0.005$), with no significant group
287 differences ($F(1,23)=1.01$, $p=0.33$), or interaction effects ($F(1,23)=0.18$, $p=0.68$).

288

289 **4. DISCUSSION**

290

291 This study assessed cerebral activity and behavioral response during a guessing task to identify
292 abnormalities in reward outcome-related processing in severely obese subjects. Reward-related
293 activations in the ventral striatum and ventromedial prefrontal cortex were observed in both obese and
294 non-obese subjects. In obese subjects, group-specific reward-related activations were found in the
295 bilateral dlPFC and precuneus/posterior cingulate. In both obese and non-obese subjects, reward-related
296 activation in the dlPFC correlated with slower choice times for repeated reward. However, significant
297 differences between groups were found with respect to the association of ghrelin with dlPFC activation
298 and reward-related behavior.

299 The comparable ventromedial prefrontal cortex activation during winning trials in obese and
300 control subjects, and the slight obesity-related trend for hyperactivation in the ventral striatum were
301 unexpected in view of the well-documented alterations in reward system function in obesity (Volkow et
302 al., 2011). This result may be due to the nature of the paradigm/incentive stimuli (money-labeled gaming

303 reinforcement task) used in the present study. In support of this possibility, alterations in reward-related
304 brain activity in obesity have been shown to be task-dependent (Volkow et al., 2011) and, while activity
305 in the ventral striatum can be triggered by both monetary and food reward, BMI was found to correlate
306 only with the anticipatory food-reward processing (Simon et al., 2014).

307 The guessing task paradigm used in the present study produced a significant hyperactivity in the
308 precuneus/posterior cingulate and dlPFC in obese subjects, suggesting the recruitment of a more extended
309 brain network for the processing of reward-related information in obesity. This hypothesis is in line with
310 previous reports indicating greater connectivity between reward network brain regions and regions of
311 executive control, emotional arousal, and somatosensory networks in obese individuals (Gupta and
312 Bhatia, 2008). The increased activation of the precuneus, a region of the default mode, found in the group
313 of obese subjects recruited in the present study is consistent with previous reports documenting increased
314 bilateral functional connectivity in the precuneus (Kullmann et al., 2012), together with greater precuneus
315 activity at baseline and in response to food and non-food reward-related cues and during response
316 inhibition and error processing (Bohon, 2017; Hsu et al., 2017; Opel et al., 2015; Tregellas et al., 2011;
317 Zhang et al., 2015) in obese samples. Similarly, greater dlPFC activity and deficits in executive function
318 have been well documented in obesity (Gautier et al., 2000; Holsen et al., 2012; Gentier et al., 2013).
319 Interestingly, in the present study, increased dlPFC activity correlated with slower choice times after
320 repeated (but not single) preceding gains/reward. In view of the greater dlPFC activation in obese
321 subjects, this finding may indicate a greater engagement of effortful, self-control executive processes in
322 rewarded decision-making context (Danner et al., 2012; Gentier et al., 2013), consistent with the known
323 link between greater dlPFC activity and greater task difficulty (Manoach et al., 1997) or high-risk choice
324 (Bembich et al., 2014). In that perspective, the observed overactivation of brain regions related to self-
325 focusing, i.e., precuneus, (Cavanna and Trimble, 2006) and self-control, i.e., dlPFC, (Belfort-DeAguiar et
326 al., 2018) could reflect an increased demand for self-control in rewarding circumstances in obese subjects.
327 A potential alternative explanation is that the occurrence of a string of reward episodes captures more
328 attentional resources in obese subjects, and thus slows down subsequent choices (Kube et al., 2017).

329 Reward-related over-activation in the dlPFC was associated with reduced circulating levels of
330 ghrelin, and this association was stronger in obese subjects. This finding may be interpreted as somewhat
331 surprising in view of previous results indicating increased neural responses to food pictures in brain areas
332 involved in reward processing after ghrelin administration in healthy volunteers (Malik et al., 2008).
333 Nevertheless, it may rely on brain processes differentially involved in reward anticipation and receipt
334 (Oldham et al., 2018). In support of this, prefrontal involvement has been reported specifically at reward
335 receipt, and not during reward anticipation. In line with this, obese subjects were found to exhibit dlPFC
336 hyperactivity in response to food images only after meal (Holsen et al., 2012), which is temporally better
337 aligned with reward receipt than anticipation and consistent with results from the present study where
338 dlPFC hyperactivity was measured during reward receipt (gain vs loss trials). This result is also in line
339 with data indicating reduced dlPFC activation in patients with Prader-Willi syndrome, characterized by
340 hyperphagia and hyperghrelinemia, and increased dlPFC activity in obese subjects with moderate
341 overeating and low ghrelin (Belfort-DeAguiar et al., 2018; Holsen et al., 2012).

342 Similar to the relationship found with the dlPFC, decreased ghrelin levels were found to
343 significantly correlate with slower choice times, particularly in the obese group. This finding is consistent
344 with the role of the ghrelin system in reward and decision-making processes. Given the recent
345 experimental evidence for a role of ghrelin in impulsive behavior/choice (Anderberg et al., 2016), it is
346 possible that this association relies on the effects of the ghrelin system on the central/brain regulation of
347 processes related to self-control. In line with this, higher fasting ghrelin levels were recently found to be
348 associated with greater reward sensitivity and reduced self-control/increased impulsivity (Ralevski et al.,
349 2018). Alternatively, this association may also reflect increased stress-related processes or greater worry
350 of unsuccessful outcome in obese subjects given the associations previously documented between ghrelin
351 and increased susceptibility/altered response to stress (Sarker et al., 2013; Spencer et al., 2012).

352 Albeit they strongly support the link between the ghrelin system and reward-related and decision-
353 making processes in obesity, results from the present study do not provide any information regarding the
354 temporality and causality of events nor their role in the development of overweight/obesity. While
355 previous studies addressing reward processing alterations have shown that hedonically driven food intake,

356 along with stress-induced comfort eating, may contribute to weight gain and obesity maintenance (Burger
357 and Berner, 2014; Lee and Dixon, 2017), further large-scale longitudinal studies are needed to
358 specifically address this issue.

359 This study presents certain limitations that future investigations should also overcome. Due to
360 logistic constraints inherent to clinical research on patients (ghrelin measurement was part of the routine
361 preoperative evaluation of obese patients), ghrelin levels were not measured on the same day of the fMRI
362 acquisition in 60% of the participants. While this limitation should not critically impact results from the
363 present study, based on data from prospective studies showing stable fasting ghrelin levels with time in
364 obese subjects (Hanusch-Enserer et al., 2004) and healthy volunteers (Leidy et al., 2004), future
365 investigations with concomitant measurement of ghrelin and brain activation are needed to comfort the
366 present findings. The exclusive measurement of total ghrelin, and not the active acylated form of the
367 hormone, represents another limitation of this study. Because of its high instability at room temperature,
368 the exact measurement of the active form of ghrelin is highly challenging and requires specific
369 measures/procedures for blood collection and processing (Hosoda and Kangawa, 2004) that were not
370 planned in the present study. While complementary measurements of the active form of ghrelin are highly
371 encouraged in future studies, we believe that assessment of total ghrelin remains highly relevant in the
372 context of the present study since it was shown to be highly sensitive and reliable in multiple studies of
373 obesity. In particular, total ghrelin, was found to correlate with BMI and disease severity in most cases of
374 obesity (except Prader-Willi Syndrome) and to be reduced in hyperphagia and obesity (Nonogaki, 2008;
375 Monti et al., 2006). These associations were not found with acylated ghrelin that seems to be more
376 relevant with respect to metabolic abnormalities, notably in normal subjects (Pacifico et al., 2009).
377 Another aspect that would merit further investigation is the relationship between ghrelin and insulin
378 and/or other neuroendocrine factors, given the known interactions between these different factors and
379 their possible impact on reward signaling (Baicy et al., 2007; Farooqi et al., 2007; Kullmann et al., 2012).
380 In support of this, strong relationships exist between the ghrelin system and the insulin system [e.g.
381 ghrelin and its receptors are expressed within the pancreas and β -cells and ghrelin alters insulin-
382 dependent glucose metabolism in a direct and non-direct manner (Churm et al., 2017)], which has been
383 shown to be involved in the regulation of reward processing in preclinical and clinical studies (Eisenstein

384 et al., 2015; Kleinridders and Pothos, 2019). Finally, another limitation of the study comes from the use
385 of an ROI (dlPFC) identified from the group Win-Lose contrast for subsequent correlational analyses of
386 brain activation with ghrelin and behavior. In this situation, a complete statistical independence may not
387 be assured nor the possibility of additional associations with other brain structures be excluded. While
388 these issues merit further investigation in future studies, they should not substantially influence the
389 findings given that the ROI defined by a between-group test was used for assessing within-group
390 relationships.

391 In conclusion, the present findings indicate an increased activation of brain areas related to self-
392 control and self-referencing during processing of non-food reward in obese subjects. Moreover, fasting
393 levels of circulating ghrelin, the hormone regulating energy balance and food consuming behavior, were
394 associated with generalized reward processing, both at the behavioral and neurobiological levels. The
395 associations of ghrelin with decision times and dlPFC activity suggest the potential involvement of this
396 hormone in the obesity-related behavioral alterations.

397

398

399 **Acknowledgments:** This study was achieved within the context of the Laboratory of Excellence TRAIL
400 ANR-10-LABX-57. The authors thank Professor V. Doucet and his team for their help and assistance
401 with the fMRI assessments.

REFERENCES

- 402
- 403 Abizaid, A., 2019. Stress and obesity: The ghrelin connection. *Journal of neuroendocrinology*, e12693.
- 404 Akitsuki, Y., Sugiura, M., Watanabe, J., Yamashita, K., Sassa, Y., Awata, S., Matsuoka, H., Maeda, Y.,
405 Matsue, Y., Fukuda, H., Kawashima, R., 2003. Context-dependent cortical activation in response to
406 financial reward and penalty: an event-related fMRI study. *Neuroimage*. 19, 1674-1685.
- 407 Anderberg, R.H., Hansson, C., Fenander, M., Richard, J.E., Dickson, S.L., Nissbrandt, H., Bergquist, F.,
408 Skibicka, K.P., 2016. The Stomach-Derived Hormone Ghrelin Increases Impulsive
409 Behavior. *Neuropsychopharmacology* 41, 1199-1209.
- 410 Andrews, Z.B., 2011. The extra-hypothalamic actions of ghrelin on neuronal function. *Trends Neurosci*
411 34, 31-40.
- 412 Angelidis, G., Valotassiou, V., Georgoulas, P., 2010. Current and potential roles of ghrelin in clinical
413 practice. *Journal Endocrinol Invest* 33, 823-838.
- 414 Baicy, K., London, E.D., Monterosso, J., Wong, M.L., Delibasi, T., Sharma, A., Licinio, J., 2007. Leptin
415 replacement alters brain response to food cues in genetically leptin-deficient adults. *Proc Natl Acad*
416 *Sci U S A* 104, 18276-18279.
- 417 Balodis, I.M., Kober, H., Worhunsky, P.D., White, M.A., Stevens, M.C., Pearlson, G.D., Sinha, R., Grilo,
418 C.M., Potenza, M.N., 2013. Monetary reward processing in obese individuals with and without binge
419 eating disorder. *Biol Psychiatry* 73, 877-886.
- 420 Belfort-DeAguiar, R., Seo, D., Lacadie, C., Naik, S., Schmidt, C., Lam, W., Hwang, J., Constable, T.,
421 Sinha, R., Sherwin, R.S., 2018. Humans with obesity have disordered brain responses to food images
422 during physiological hyperglycemia. *Am J Physiol Endocrinol Metab* 314, E522-E529.
- 423 Bembich, S., Clarici, A., Vecchiet, C., Baldassi, G., Cont, G., Demarini, S., 2014. Differences in time
424 course activation of dorsolateral prefrontal cortex associated with low or high risk choices in a
425 gambling task. *Front Hum Neurosci* 8, 464.
- 426 Bohon, C., 2017. Brain response to taste in overweight children: A pilot feasibility study. *PLoS One* 12,
427 e0172604.

428 Briggs, D.I., Lockie, S.H., Wu, Q., Lemus, M.B., Stark, R., Andrews, Z.B., 2013. Calorie-restricted
429 weight loss reverses high-fat diet-induced ghrelin resistance, which contributes to rebound weight
430 gain in a ghrelin-dependent manner. *Endocrinology* 154, 709-717.

431 Burger, K.S., Berner, L.A., 2014. A functional neuroimaging review of obesity, appetitive hormones and
432 ingestive behavior. *Physiol Behav* 136, 121-127.

433 Capuron, L., Pagnoni, G., Drake, D.F., Woolwine, B.J., Spivey, J.R., Crowe, R.J., Votaw, J.R., Goodman,
434 M.M., Miller, A.H., 2012. Dopaminergic mechanisms of reduced basal ganglia responses to hedonic
435 reward during interferon alfa administration. *Arch Gen Psychiatry* 69, 1044-1053.

436 Carnell, S., Gibson, C., Benson, L., Ochner, C.N., Geliebter, A., 2012. Neuroimaging and obesity: current
437 knowledge and future directions. *Obes Rev* 13, 43-56.

438 Cavanna, A.E., Trimble, M.R., 2006. The precuneus: a review of its functional anatomy and behavioural
439 correlates. *Brain*. 129, 564-583.

440 Churm, R., Davies, J.S., Stephens, J.W., Prior, S.L., 2017. Ghrelin function in human obesity and type 2
441 diabetes: a concise review. *Obes Rev* 18, 140-148.

442 Danner, U.N., Ouwehand, C., van Haastert, N.L., Hornsveld, H., de Ridder, D.T., 2012. Decision-making
443 impairments in women with binge eating disorder in comparison with obese and normal weight
444 women. *Eur Eat Disord Rev* 20, e56-62.

445 Date, Y., 2012. Ghrelin and the vagus nerve. *Methods Enzymol* 514, 261-269.

446 Eisenstein, S.A., Gredysa, D.M., Antenor-Dorsey, J.A., Green, L., Arbeláez, A.M., Koller, J.M., KJ, B.,
447 JS, P., SM, M., Hershey, T., 2015. Insulin, Central Dopamine D2 Receptors, and Monetary Reward
448 Discounting in Obesity. *PloS one* 10(7), e0133621.

449 English, P.J., Ghatei, M.A., Malik, I.A., Bloom, S.R., Wilding, J.P., 2002. Food fails to suppress ghrelin
450 levels in obese humans. *J Clin Endocrinol Metab* 87, 2984.

451 Farooqi, I.S., Bullmore, E., Keogh, J., Gillard, J., O'Rahilly, S., Fletcher, P.C., 2007. Leptin regulates
452 striatal regions and human eating behavior. *Science* 317, 1355.

453 Foster, C.M., Barkan, A., Kasa-Vubu, J.Z., Jaffe, C., 2007. Ghrelin concentrations reflect body mass
454 index rather than feeding status in obese girls. *Pediatr Res* 62, 731-734.

455 Gautier, J.F., Chen, K., Salbe, A.D., Bandy, D., Pratley, R.E., Heiman, M., Ravussin, E., Reiman, E.M.,
456 Tataranni, P.A., 2000. Differential brain responses to satiation in obese and lean men. *Diabetes* 49,
457 838-846.

458 Gentier, I., Augustijn, M., Deforche, B., Tanghe, A., De Bourdeaudhuij, I., Lenoir, M., D'Hondt, E.,
459 2013. A comparative study of performance in simple and choice reaction time tasks between obese
460 and healthy-weight children. *Res Dev Disabil* 34, 2635-2641.

461 Goldstone, A.P., Prechtel, C.G., Scholtz, S., Miras, A.D., Chhina, N., Durighel, G., Deliran, S.S.,
462 Beckmann, C., Ghatei, M.A., Ashby, D.R., Waldman, A.D., Gaylinn, B.D., Thorner, M.O., Frost,
463 G.S., Bloom, S.R., Bell, J.D., 2014. Ghrelin mimics fasting to enhance human hedonic, orbitofrontal
464 cortex, and hippocampal responses to food. *Am J Clin Nutr* 99, 1319-1330.

465 Gupta, R., Bhatia, M.S., 2008. Short-lasting unilateral neuralgiform headache with conjunctival injection
466 and tearing: response to antiepileptic dual therapy. *Neurol India* 56, 71-73.

467 Hanusch-Enserer, U., Cauza, E., Brabant, G., Dunky, A., Rosen, H., Pacini, G., Tuchler, H., Prager, R.,
468 Roden, M., 2004. Plasma ghrelin in obesity before and after weight loss after laparoscopic
469 adjustable gastric banding. *J Clin Endocrinol Metab* 89, 3352-3358.

470 Holsen, L.M., Savage, C.R., Martin, L.E., Bruce, A.S., Lepping, R.J., Ko, E., Brooks, W.M., Butler,
471 M.G., Zarcone, J.R., Goldstein, J.M., 2012. Importance of reward and prefrontal circuitry in hunger
472 and satiety: Prader-Willi syndrome vs simple obesity. *Int J Obes (Lond)*. 36, 638-647.

473 Hosoda, H., Kangawa, K. 2004. Ghrelin Measurement: Present and Perspectives. In: Ghigo E., Benso A.,
474 Broglio F. (eds) Ghrelin. *Endocrine Updates*, vol 23. Springer, Boston, MA.

475 Howick, K., Griffin, B.T., Cryan, J.F., Schellekens, H., 2017. From Belly to Brain: Targeting the Ghrelin
476 Receptor in Appetite and Food Intake Regulation. *Int J Mol Sci* 18.

477 Hsu, J.S., Wang, P.W., Ko, C.H., Hsieh, T.J., Chen, C.Y., Yen, J.Y., 2017. Altered brain correlates of
478 response inhibition and error processing in females with obesity and sweet food addiction: A
479 functional magnetic imaging study. *Obes Res Clin Pract* 11, 677-686.

480 Kleinridders, A. Pothos, E.N., 2019. Impact of brain insulin signaling on dopamine function, food intake,
481 reward, and emotional behavior. *Curr Nutr Rep* 8, 83-91.

482 Kube, J., Mathar, D., Horstmann, A., Kotz, S.A., Villringer, A., Neumann, J., 2017. Altered monetary
483 loss processing and reinforcement-based learning in individuals with obesity. *Brain Imaging Behav*
484 12, 1431-1449.

485 Kullmann, S., Heni, M., Veit, R., Ketterer, C., Schick, F., Haring, H.U., Fritsche, A., Preissl, H., 2012.
486 The obese brain: association of body mass index and insulin sensitivity with resting state network
487 functional connectivity. *Hum Brain Mapp* 33, 1052-1061.

488 Lee, P.C., Dixon, J.B., 2017. Food for Thought: Reward Mechanisms and Hedonic Overeating in Obesity.
489 *Curr Obes Rep* 6, 353-361.

490 Leidy, H.J., Gardner, J.K., Frye, B.R., Snook, M.L., Schuchert, M.K., Richard, E.L., Williams, N.I.,
491 2004. Circulating ghrelin is sensitive to changes in body weight during a diet and exercise program in
492 normal-weight young women. *J Clin Endocrinol Metab* 89, 2659-64.

493 Malik, S., McGlone, F., Bedrossian, D., Dagher, A., 2008. Ghrelin modulates brain activity in areas that
494 control appetitive behavior. *Cell Metab* 7, 400-409.

495 Manoach, D.S., Schlaug, G., Siewert, B., Darby, D., Bly, B., Benfield, A., Edelman, R., Warach, S.,
496 1997. Prefrontal cortex fMRI signal changes are correlated with working memory load. *Neuroreport*.
497 8, 545-549.

498 Monti, V., Carlson, J.J., Hunt, S.C., Adams, T.D., 2006. Relationship of ghrelin and leptin hormones with
499 body mass index and waist circumference in a random sample of adults. *J Am Diet Assoc* 106, 822-
500 828.

501 Muller, T.D., Nogueiras, R., Andermann, M.L., Andrews, Z.B., Anker, S.D., Argente, J., Batterham,
502 R.L., Benoit, S.C., Bowers, C.Y., Broglio, F., Casanueva, F.F., D'Alessio, D., Depoortere, I.,
503 Geliebter, A., Ghigo, E., Cole, P.A., Cowley, M., Cummings, D.E., Dagher, A., Diano, S., Dickson,
504 S.L., Dieguez, C., Granata, R., Grill, H.J., Grove, K., Habegger, K.M., Heppner, K., Heiman, M.L.,
505 Holsen, L., Holst, B., Inui, A., Jansson, J.O., Kirchner, H., Korbonits, M., Laferrere, B., LeRoux,
506 C.W., Lopez, M., Morin, S., Nakazato, M., Nass, R., Perez-Tilve, D., Pfluger, P.T., Schwartz, T.W.,
507 Seeley, R.J., Sleeman, M., Sun, Y., Sussel, L., Tong, J., Thorner, M.O., van der Lely, A.J., van der
508 Ploeg, L.H., Zigman, J.M., Kojima, M., Kangawa, K., Smith, R.G., Horvath, T., Tschop, M.H., 2015.
509 Ghrelin. *Mol Metab* 21, 437-60.

510 Natalucci, G., Riedl, S., Gleiss, A., Zidek, T., Frisch, H., 2005. Spontaneous 24-h ghrelin secretion
511 pattern in fasting subjects: maintenance of a meal-related pattern. *Eur J Endocrinol.* 152, 845-850.

512 Nonogaki, K., 2008. Ghrelin and feedback systems. *Vit Horms* 77, 149-170.

513 Oldham, S., Murawski, C., Fornito, A., Youssef, G., Yucel, M., Lorenzetti, V., 2018. The anticipation and
514 outcome phases of reward and loss processing: A neuroimaging meta-analysis of the monetary
515 incentive delay task. *Hum Brain Mapp* 39, 3398-3418.

516 Opel, N., Redlich, R., Grotegerd, D., Dohm, K., Hauptenthal, C., Heindel, W., Kugel, H., Arolt, V.,
517 Dannlowski, U., 2015. Enhanced neural responsiveness to reward associated with obesity in the
518 absence of food-related stimuli. *Hum Brain Mapp* 36, 2330-2337.

519 Pacifico, L., Poggiogalle, E., Costantino, F., Anania, C., Ferraro, F., Chiarelli, F., Chiesa, C., 2009.
520 Acylated and nonacylated ghrelin levels and their associations with insulin resistance in obese and
521 normal weight children with metabolic syndrome. *Eur J Endocrinol* 161, 861-870.

522 Perello, M., Dickson, S.L., 2015. Ghrelin signalling on food reward: a salient link between the gut and the
523 mesolimbic system. *J Neuroendocrinol* 27, 424-434.

524 Price, M., Higgs, S., Maw, J., Lee, M., 2016. A dual-process approach to exploring the role of delay
525 discounting in obesity. *Physiol Behav* 162, 46-51.

526 Pursey, K.M., Stanwell, P., Callister, R.J., Brain, K., Collins, C.E., Burrows, T.L., 2014. Neural
527 Responses to Visual Food Cues According to Weight Status: A Systematic Review of Functional
528 Magnetic Resonance Imaging Studies. *Front Nutr* 1, 7.

529 Puzziferri, N., Zigman, J.M., Thomas, B.P., Mihalakos, P., Gallagher, R., Lutter, M., Carmody, T., Lu,
530 H., Tamminga, C.A., 2016. Brain imaging demonstrates a reduced neural impact of eating in obesity.
531 *Obesity (Silver Spring, Md.)* 24, 829-836.

532 Ralevski, E., Shanabrough, M., Newcomb, J., Gandelman, E., Hayden, R., Horvath, T.L., Petrakis, I.,
533 2018. Ghrelin is Related to Personality Differences in Reward Sensitivity and Impulsivity. *Alcohol*
534 *Alcohol.* 53, 52-56.

535 Reuter, J., Raedler, T., Rose, M., Hand, I., Glascher, J., Buchel, C., 2005. Pathological gambling is linked
536 to reduced activation of the mesolimbic reward system. *Nat Neurosci.* 8, 147-148.

537 Revitsky, A.R, Klein, L.C., 2013. Role of ghrelin in drug abuse and reward-relevant behaviors: a
538 burgeoning field and gaps in the literature. *Curr Drug Abuse Rev* 6, 213-44.

539 Sarker, M.R., Franks, S., Caffrey, J., 2013. Direction of post-prandial ghrelin response associated with
540 cortisol response, perceived stress and anxiety, and self-reported coping and hunger in obese women.
541 *Behav Brain Res* 257, 197-200.

542 Simon, J.J., Skunde, M., Hamze Sinno, M., Brockmeyer, T., Herpertz, S.C., Bendszus, M., Herzog, W.,
543 Friederich, H.C., 2014. Impaired Cross-Talk between Mesolimbic Food Reward Processing and
544 Metabolic Signaling Predicts Body Mass Index. *Front Behav Neurosci* 8, 359.

545 Solomou, S., Korbonts, M., 2014. The role of ghrelin in weight-regulation disorders: implications in
546 clinical practice. *Hormones (Athens)* 13, 458-475.

547 Spencer, S.J., Xu, L., Clarke, M.A., Lemus, M., Reichenbach, A., Geenen, B., Kozicz, T., Andrews, Z.B.,
548 2012. Ghrelin regulates the hypothalamic-pituitary-adrenal axis and restricts anxiety after acute
549 stress. *Biol Psychiatry* 72, 457-465.

550 Tregellas, J.R., Wylie, K.P., Rojas, D.C., Tanabe, J., Martin, J., Kronberg, E., Cordes, D., Cornier, M.A.,
551 2011. Altered default network activity in obesity. *Obesity (Silver Spring, Md.)* 19, 2316-2321.

552 Tschöp, M., Weyer, C., Tataranni, P.A., Devanarayan, V., Ravussin, E., Heiman, M.L., 2001. Circulating
553 ghrelin levels are decreased in human obesity. *Diabetes* 50, 707-9.

554 Uriarte, M., De Francesco, P.N., Fernandez, G., Cabral, A., Castrogiovanni, D., Lalonde, T., Luyt, L.G.,
555 Trejo, S., Perello, M., 2018. Evidence Supporting a Role for the Blood-Cerebrospinal Fluid Barrier
556 Transporting Circulating Ghrelin into the Brain. *Mol Neurobiol* 56, 4120-4134.

557 Volkow, N.D., Wang, G.J., Baler, R.D., 2011. Reward, dopamine and the control of food intake:
558 implications for obesity. *Trends Cogn Sci.* 15, 37-46.

559 Weiskopf, N., Hutton, C., Josephs, O., Turner, R., Deichmann, R., 2007. Optimized EPI for fMRI studies
560 of the orbitofrontal cortex: compensation of susceptibility-induced gradients in the readout direction.
561 *Magma.* 20, 39-49.

562 Yildiz BO, Suchard MA, Wong M-L, McCann SM, J., L., 2004. Alterations in the dynamics of
563 circulating ghrelin, adiponectin, and leptin in human obesity. *Proc Natl Acad Sci USA* 101, 10434-
564 10439.

565 Zhang, B., Tian, D., Yu, C., Li, M., Zang, Y., Liu, Y., Walter, M., 2015. Altered baseline brain activity
566 differentiates regional mechanisms subserving biological and psychological alterations in obese men.
567 Sci Rep 5, 11563.

568 Zigman, J.M., Bouret, S.G., Andrews, Z.B., 2016. Obesity Impairs the Action of the Neuroendocrine
569 Ghrelin System. Trends Endocrinol Metab 27, 54-63.

570

571

572 **TABLES**

573

574 **Table 1. Characteristics of study participants**

	Obese (N=15)	Normal-weight Controls (N=15)	<i>Difference P value</i>
Age, years (SD)	38.7 (9.4)	37 (7.1)	0.6
Gender, men/women (%)	3/12 (20, 80)	2/13 (13.3, 86.7)	0.6
Body Mass Index (BMI), kg/m ²	41.2 (2.1)	21.8 (2.6)	< 0.0001
Weight, kg (SD)	114.7 (9.1)	60.5 (10)	< 0.0001
Diploma ¹	2.9 (0.96)	3.4 (1.40)	0.3
Ghrelin levels, pg/mL (SD)	375.2 (47.6)	811.6 (87.8)	< 0.001

575

576 ¹ Diploma was measured on a scale from 0 (no diploma) to 7 (PhD or MD).

577 Data are shown as mean (SD) or prevalence (%). The statistical significance of group differences

578 (*p*-value) was assessed by Student t-tests (Chi-square test, for Gender).

579

580 **Table 2. Brain regions activated by the reward condition (winning trials > losing trials) in**
581 **obese and non-obese controls**

Brain Region	k _E	T-score	Peak coordinates		
			X	Y	Z
<i>Non-obese controls (cluster-forming threshold p<0.001)</i>					
Ventral striatum	317	9.65*	9	17	-5
Orbitofrontal cortex	107	7.40	21	29	-14
Extrastriate visual cortex (L)	466	9.78*	-36	-88	-11
Extrastriate visual cortex (L)	40	6.73	-39	-46	-23
Extrastriate visual cortex (R)	500	9.42*	42	-64	-20
Posterior cingulate cortex	45	7.68	6	-31	28
Intraparietal sulcus	76	6.94	21	-61	34
<i>Obese subjects (cluster-forming threshold p<0.001)</i>					
Ventral striatum + ventromedial prefrontal cortex + orbitofrontal cortex	681	9.30*	18	47	-11
Extrastriate visual cortex (R)	720	8.78*	30	-88	4
Extrastriate visual cortex (L)	263	6.93	-27	-82	-11
S1(R) + precuneus + posterior cingulate cortex	1516	7.84	39	-22	52
Intraparietal sulcus	159	4.08	-27	-70	52
S1(R)	79	6.17	57	-13	34
Frontal operculum/premotor cortex	93	4.01	-42	2	13
<i>Obese subjects > non-obese controls (cluster-forming threshold p<0.001)</i>					
Dorsolateral prefrontal cortex (R)	81	6.28*	24	17	37
Dorsolateral prefrontal cortex (L)	56	4.92	-30	23	40
Precuneus / posterior cingulate	126	5.32	-6	-43	43

582

583 *Clusters where the T-score of the peak voxel is significant (FWER p<0.05, at the voxel level).

584 The other clusters are significant at the cluster level, but not voxel level (FWER p<0.05, at the

585 cluster level), k_E is the cluster size in voxels. Obese subject, N=14; non-obese controls, N=13.

586 **FIGURE LEGENDS**

587

588 **Figure 1. Reward-task paradigm**

589 **Panel A:** Full timeline of the stimulus presentation design, which was identical across subjects.

590 The net score is indicated by the noisy sinusoid, along with gain and loss trials (white and grey
591 bars)

592 **Panel B:** Trial structure. Examples of trials 2-4 are shown. Red cards indicate gain, black cards
593 indicate loss. Subjects could choose either left or right card, but the outcome in terms of win or
594 loss was predefined (see A).

595

596 **Figure 2. Reward-related responses in obese subjects and non-obese controls**

597 **Panel A:** Reward-related response in the ventral striatum across both experimental groups, as the
598 result of the two-sample (obese, N=14 or controls, N=13) “gain vs. loss” F-test.

599 **Panel B:** Group-wise reward-related activation. Note that the activation in obese subjects
600 extends to precuneus/posterior cingulate and dorsolateral prefrontal cortex (dlPFC), unlike in
601 non-obese controls (within-group activation).

602

603 **Figure 3. Reward-related activity is greater in obese subjects**

604 The statistical contrast Obese (N=14) > Controls (N=13), thresholded for illustration purposes at
605 cluster forming threshold $p < 0.005$ is mapped onto the average of the individual subjects’

606 normalized anatomical images (n=30). Significant clusters and respective peak contrast estimates
607 are shown: (A) left dorsolateral prefrontal cortex, (B) right dorsolateral prefrontal cortex and (C)
608 precuneus/posterior cingulate.

609

610 **Figure 4. Choice times and relationship with dorsolateral prefrontal cortex (dlPFC)**

611 **activity and ghrelin levels in obese subjects and non-obese controls**

612 **Panel A.** Choice time (CT) group effect. A significant difference between groups was observed
613 for choice time during repeated reward (CT2) but not during single reward (CT1); * $p < 0.05$.

614 **Panel B.** Relationship between ghrelin levels and reward-related activity in the dlPFC. The
615 parameter estimates of the “Gain vs. Loss” contrast were extracted from the bilateral dlPFC (i.e.,
616 the first eigenvariates of the individual contrast images within two spherical ROI of 6 mm radius
617 centered on the peaks’ coordinates of [24 17 37] and [-30 23 40] were extracted; as shown in a
618 right upper corner). Main effects and interactions: Group: $F = 14.6$, $p < 0.001$, $\eta^2 p = 0.39$; Ghrelin:
619 $F = 9.4$, $p < 0.01$, $\eta^2 p = 0.29$; Group x Ghrelin: $F = 5.2$, $p < 0.05$, $\eta^2 p = 0.18$.



620 **Panel C.** Relationship between choice time during repeated reward (CT2) and ghrelin levels.
621 Corresponding CIs displayed as gray shadows. Main effects and interactions: Group: $F = 5.9$,
622 $p < 0.05$, $\eta^2 p = 0.20$; Ghrelin: $F = 4.9$, $p < 0.05$, $\eta^2 p = 0.18$; Group x Ghrelin: $F = 5.3$, $p < 0.05$,
623 $\eta^2 p = 0.19$.

624 **Panel D.** Relationship between choice time during repeated reward (CT2) and reward-related
625 activity in the dlPFC. The parameter estimates of the “Gain vs. Loss” contrast were extracted
626 from the bilateral dlPFC (i.e., the first eigenvariates of the individual contrast images within two
627 spherical ROI of 6 mm radius centered on the peaks’ coordinates of [24 17 37] and [-30 23 40]
628 were extracted; as shown in a right upper corner). Main effect: dlPFC: $F = 9.4$, $p < 0.01$, $\eta^2 p = 0.29$.

629 ○ control subjects, N=13 ● obese subjects, N=14

630

Trials





Common activation

Contrast estimates and 90% C.I.

vStr R
[12, 8, -5]




vStr L
[-9, 11, -5]



Within-group activation


Group Control (n=13)

Group Obese (n=14)



Reward F-score
FWE $p < 0.05$


Reward T-score
uncor. $p < 0.005$




A.

B.

dIPFC L [-30, 23, 40]





Contrast estimates and 90% C.I.





A.

dIPFC R [24, 17, 37]




B.

Precuneus/PCC [-6, -43, 43]




C.


A.



B.



C.



D.

