

Reward-related brain activity and behavior are associated with peripheral ghrelin levels in obesity

Volodymyr Bogdanov, Olena Bogdanova, Sandra Dexpert, Ines Delgado, Helen Beyer, Agnès Aubert, Bixente Dilharreguy, Cédric Beau, Damien Forestier, Patrick Ledaguenel, et al.

▶ To cite this version:

Volodymyr Bogdanov, Olena Bogdanova, Sandra Dexpert, Ines Delgado, Helen Beyer, et al.: Reward-related brain activity and behavior are associated with peripheral ghrelin levels in obesity. Psychoneuroendocrinology, 2020, 112, pp.1-8. 10.1016/j.psyneuen.2019.104520 . hal-02517346

HAL Id: hal-02517346 https://hal.science/hal-02517346

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0306453019312612 Manuscript_64e38af6ae801a9edeaae41298d3a2b4

1	Reward-related brain activity and behavior are associated with peripheral ghrelin levels in obesity				
2					
3	Volodymyr B. Bogdanov ^{1,2#} , Olena V. Bogdanova ³ , Sandra Dexpert ¹ , Ines Delgado ¹ , Helen				
4	Beyer ¹ , Agnès Aubert ¹ , Bixente Dilharreguy ⁴ , Cédric Beau ⁵ , Damien Forestier ⁵ , Patrick				
5	Ledaguenel ⁵ , Eric Magne ⁵ , Bruno Aouizerate ¹ , Sophie Layé ¹ , Guillaume Ferreira ¹ , Jennifer Felger ⁶ ,				
6	Giuseppe Pagnoni ^{7,8} , Lucile Capuron ^{1#}				
7					
8	¹ Univ. Bordeaux, INRA, Bordeaux INP, NutriNeuro, UMR 1286, F-33000, Bordeaux, France;				
9	² Univ. Lyon, Ecole Nationale des Travaux Publics de l'Etat, Laboratoire Génie Civil et Bâtiment, F-				
10	69518 Vaulx-en-Velin, France;				
11	³ INSERM U1028 - CNRS UMR5292, 16 avenue Doyen Lépine, F-69676, Bron, France;				
12	⁴ Univ. Bordeaux, INCIA, CNRS, UMR 5287, F-33076 Bordeaux, France;				
13	⁵ Digestive and Parietal Surgery, Clinique Tivoli, F-33000, Bordeaux, and Clinique Jean Villar, F-33520,				
14	Bruges France;				
15	⁶ Dpt of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA 30322,				
16	USA;				
17	⁷ Dept of Neural, Biomedical, and Metabolic Sciences, University of Modena and Reggio Emilia, I-				
18	41125, Modena, Italy;				
19	⁸ Center for Neuroscience and Neurotechnology, University of Modena and Reggio Emilia, I-41125,				
20	Modena, Italy.				
21					
22					
23	# Corresponding authors:				
24	- Volodymyr B. Bogdanov, PhD, Univ. Lyon, Ecole Nationale des Travaux Publics de l'Etat, Laboratoire				
25	Génie Civil et Bâtiment, 3 rue Maurice Audin, F-69518 Vaulx-en-Velin, France; E-mail:				
26	vlabogd@yahoo.com				
27	- Lucile Capuron, PhD, INRA 1286, NutriNeuro, University of Bordeaux, 146 rue Leo Saignat, F-33076				
28	Bordeaux, France; Tel. +33 (0)557571233, Fax. +33 (0) 557571227; E-mail: lucile.capuron@inra.fr				

- 30 Running Title. Ghrelin-related reward activity in obesity

- **Conflict of Interest.** The authors declare no conflict of interest in relation to the work described.
- 34 Funding/Support: This work was supported by the French National Research Agency, ANR (ANR-11-
- 35 JSV1-0006, Dr Capuron) and funds from the French National Institute for Agricultural Research (INRA).

39 ABSTRACT

Background/Objectives: While excessive food consumption represents a key factor in the development of obesity, the underlying mechanisms are still unclear. Ghrelin, a gut-brain hormone involved in the regulation of appetite, is impaired in obesity. In addition to its role in eating behavior, this hormone was shown to affect brain regions controlling reward, including the striatum and prefrontal cortex, and there is strong evidence of impaired reward processing in obesity. The present study investigated the possibility that disrupted reward-related brain activity in obesity relates to ghrelin deficiency.

46 **Subjects/Methods:** Fifteen severely obese subjects (BMI > 35 kg/m^2) and fifteen healthy non-obese 47 control subjects (BMI < 30 kg/m^2) were recruited. A guessing-task paradigm, previously shown to 48 activate the ventral striatum, was used to assess reward-related brain neural activity by functional 49 magnetic resonance imaging (fMRI). Fasting blood samples were collected for the measurement of 50 circulating ghrelin.

Results: Significant activations in the ventral striatum, ventromedial prefrontal cortex and extrastriate visual cortex were elicited by the fMRI task in both obese and control subjects. In addition, greater reward-related activations were present in the dorsolateral prefrontal cortex, and precuneus/posterior cingulate of obese subjects compared to controls. Obese subjects exhibited longer choice times after repeated reward and lower circulating ghrelin levels than lean controls. Reduced ghrelin levels significantly predicted slower post-reward choices and reward-related hyperactivity in dorsolateral prefrontal cortices in obese subjects.

58 Conclusion: This study provides evidence of association between circulating ghrelin and reward-related 59 brain activity in obesity and encourages further exploration of the role of ghrelin system in altered eating 60 behavior in obesity.

61

62 Key words: Obesity, Reward, Behavior, fMRI, Ghrelin

63 1. INTRODUCTION

64

Abnormal feeding behavior resulting in weight gain and obesity has been linked to alterations in 65 reward processing (Burger and Berner, 2014). In support of this, impaired reward-related brain responses 66 have been repeatedly documented in overweight/obese subjects (Carnell et al., 2012; Pursey et al., 2014; 67 Puzziferri et al., 2016). These alterations were found not only in food-related paradigms but also in more 68 generalized experimental conditions, including monetary incentive tasks (Balodis et al., 2013). Monetary 69 70 delay discounting, which is a bias towards smaller but earlier rewards, is greater in obese subjects 71 pointing to an abnormal reward expectation and decision making in obesity (Price et al., 2016). These effects may be related to a deregulation of hormones that control reward-directed and food-consuming 72 73 behaviors. In this perspective, a promising target for obesity research is the ghrelin-regulating system, 74 which is involved not only in appetitive behavior but also in higher cognitive functions (Anderberg et al., 75 2016; Ralevski et al., 2018).

76 Ghrelin is a peptide that is mainly produced by neuroendocrine cells in the stomach mucosa. Secretion of ghrelin has a fluctuating character depending on circadian rhythms and food anticipation 77 (Natalucci et al., 2005). Immediately after food consumption, ghrelin levels fall (English et al., 2002) and 78 79 then gradually increase to reach a peak before the next meal is initiated. Once secreted, ghrelin is able to 80 act on numerous peripheral tissues and the brain (Muller et al., 2015). Peripheral ghrelin can affect brain function via different pathways (Howick et al., 2017), including vagal or brain stem stimulation (Date, 81 2012), blood stream (Angelidis et al., 2010), passage across the blood-brain barrier via saturated transport 82 system (Solomou and Korbonits, 2014), and transport via the cerebrospinal fluid (Uriarte et al., 2018). 83 84 Ghrelin receptors are widely expressed in the brain, notably in the orexigenic neurons of hypothalamic nuclei (Muller et al., 2015) and in the hippocampus, pituitary, dentate gyrus, substantia nigra, 85 dorsal/median raphe nuclei and ventral tegmental area (Andrews, 2011). Altogether, they contribute to 86 the regulation of appetite, energy balance, olfaction, but also memory/learning, stress response, mood, 87 88 and reward control (Abizaid, 2019; Goldstone et al., 2014; Howick et al., 2017; Malik et al., 2008; Perello

89 and Dickson, 2015).

90 Studies in animals have shown that ghrelin activates the reward circuitry comprising of the ventral tegmental area, lateral hypothalamus and nucleus accumbens (Revitsky and Klein, 2013). At the 91 92 clinical level, reductions in dorsolateral prefrontal cortex (dIPFC) activity and decreased self-control, 93 supposed to be related to uncontrolled overeating (Holsen et al., 2012), have been reported in overweight patients with Prader-Willi syndrome (PWS) who also exhibit increased ghrelin levels. In healthy subjects, 94 increased ghrelin levels have been related to increased sensitivity to reward and impulsivity/lower self-95 96 control (Ralevski et al., 2018). Deregulation of the ghrelin system may also contribute to the development 97 of obesity (Abizaid, 2019). In obese subjects, not only fasting ghrelin levels are reduced proportionally to body mass index (BMI) (Tschop et al., 2001; Nonogaki, 2008), but also alterations in the circadian 98 99 rhythm and postprandial inhibition of ghrelin secretion are reported (Yildiz BO et al., 2004). The patterns of phasic and tonic ghrelin regulations are altered (Foster et al., 2007) and ghrelin resistance is proposed 100 101 as one of the major mechanisms of ghrelin system deficiency in obesity (Zigman et al., 2016). Central action of ghrelin in obesity may be also impaired due to different facets of ghrelin resistance (Zigman et 102 al., 2016). Interestingly, normalization of blood levels of ghrelin after the weight loss is associated with 103 104 improved brain sensitivity to ghrelin (Briggs et al., 2013).

The association of the ghrelin system with both reward response and appetite control/metabolic 105 regulation suggests the potential role of decreased ghrelin levels in the development of brain reward 106 107 system alterations and disrupted eating behaviors likely to promote the development of obesity. While 108 altered reward-related neural activation in obesity has been frequently associated with changes in other 109 energy balance regulating hormones, including leptin and insulin (Baicy et al., 2007; Farooqi et al., 2007; Kullmann et al., 2012), no studies have investigated the potential link between obesity-related ghrelin 110 deficiency and disruptions in the brain reward system. The present study addressed this issue in a sample 111 112 of severely obese subjects compared to healthy non-obese controls, using functional magnetic resonance 113 imaging (fMRI) during a non-food related task known to induce reward-related cerebral activity. 114 Behavioral responses and brain processing of reward outcome were investigated and their relationship with individual levels of ghrelin was assessed. 115

117 2. METHODS

118 2.1. Subjects

Fifteen severely obese subjects (BMI > 35 kg/m²; 3 males and 12 females) and fifteen healthy 119 non-obese control subjects (BMI $\leq 30 \text{ kg/m}^2$; 2 males and 13 females) were included. All subjects were 120 121 right-handed adults, free of any treatment with psychotropic effects, and had not eaten for the last two hours before the fMRI session. Additional exclusion criteria were age > 65 years old, history of brain 122 damage or neurological disease, acute or chronic inflammatory disorders (other than obesity and obesity-123 related conditions), severe or uncontrolled medical conditions, psychiatric disorders, metallic implants 124 and eating disorders (in non-obese control subjects). All subjects provided written informed consent and 125 received compensation for their participation in the study. The study was approved by the Committee for 126 the Protection of Persons (CPP) of Bordeaux. 127

128

129 2.2. Guessing task

130 We used a guessing-task paradigm that was previously shown to activate the ventral striatum, 131 involved in reward processing (Capuron et al., 2012; Reuter et al., 2005). Subjects were presented with 132 two hidden cards on the left and right sides of the screen and were asked to guess which of the two cards was "red" by pressing on a keypad with the index (left key) or middle finger (right key) (Figure 1A, B). 133 The time window for the choice was fixed at 2s, after which the selected card was turned over and the 134 outcome was displayed on the screen for another 2s. If the card was "red", the subject gained one Euro, if 135 "black" one Euro was lost. The sequence of gains and losses was similar for all subjects. Unbeknownst to 136 participants, the outcomes followed a noisy sinusoid with a slight upward trend (Figure 1A), as described 137 138 elsewhere (Reuter et al., 2005). The net score, which was initially equal to 15 Euros, was always shown 139 on the screen and updated each time the chosen card was revealed. The choice time was recorded for each 140 trial. In total, there were 100 trials, and the total duration of the task was 6 min 40 sec. In contrast to previous studies using this paradigm, participants were informed that the task should be considered as a 141 game and the amount of their compensation (related to their participation in the study) would not depend 142

of their gain or lost; furthermore, the number of trials was reduced by a factor of three (Reuter et al.,
2005). These methodological adaptations did not impede significant ventral striatum reward response, as
shown in **Figure 2**. All stimuli presentations and recordings of reaction time and response keys were
performed with E-Prime 2.0 software (Psychology Software Tools, Sharpsburg, PA, USA).

147 Before the acquisition started, a practice session was performed outside the scanner room on a laptop computer under the supervision of the experimenter. An additional practice session was performed 148 in the scanner using the MRI-compatible keypad (4-key response box). After practice, participants 149 150 performed the task during the scanning session. The task was presented on a screen placed in front of the 151 scanner bed and was monitored through a computer located outside the scanner room. Participants were 152 able to see the screen through a mirror attached to the head coil. Choice time (CT), i.e. the time elapsed from the presentation of the cards on the screen to the volunteer's button press signaling the choice of one 153 of the two hidden decks (left or right), was recorded as a behavioral parameter. We were particularly 154 155 interested in decision-making processes related to further reward acquisition *after* a reward had already 156 been received, as this is an experimental situation that could provide a model for the prolonged consumption of food in the obese. Accordingly, only CTs after winning trials were analyzed. 157 Furthermore, CTs were grouped into two subsets according to the event-sequence, as this is known to 158 159 affect cerebral responses in reward tasks (Akitsuki et al., 2003). When there was just one win before the 160 current trial, response latency was classified as CT after short reward (CT1). When there were two or more wins in a row before the current trial, CT was considered as being within the context of long or 161 162 sustained reward (CT2).

163

164 2.3. MRI acquisition

Scanning was performed on a 3T scanner GE Discovery MR750 with a 32 channels head coil.
First, a high-resolution T1-weighted anatomical scan was acquired via a FSPGR sequence with the
following parameters: TR=7.896 ms, TE=3.216 ms, TI=400 ms, FA=10°, FoV=288 mm², 200 sagittal
slices with a 288x288 acquisition matrix, 1 mm³ isotropic resolution, total acquisition time: 3 min 43 s.
Then, two functional runs of T2*-weighted echo-planar imaging for measurement of the blood oxygen

level-dependent (BOLD) contrast were acquired, with the following parameters: TR=2000 ms, TE=30 170 ms, FA=80°, FOV=19.20 cm, 35 64×64 slices with a 3.5 mm thickness and a 0.5 mm between-slice gap 171 172 (effective voxel size = 3x3x4 mm), with asset factor 2. Slices were prescribed with a negative pitch angle of 30 degrees relative to AC-PC horizontal plane, in order to minimize the ventro-medial BOLD signal 173 dropout, as advised in (Weiskopf et al., 2007). During the first functional run (5 dummy scans + 264 174 dynamic scans; total duration 8 min 58 s), volunteers performed an attention task, whose results will be 175 discussed and reported elsewhere. During the second functional run (5 dummy scans + 204 dynamic 176 177 scans; total duration 6 min 58 s), subjects performed the guessing task described above.

178

179 2.4. Imaging preprocessing

180 MRI images were analyzed using the SPM software package (SPM8 v4290; Wellcome

181 Department of Cognitive Neurology, Institute of Neurology, London, UK;

182 *http://www.fil.ion.uce.ac.uk/spm*) implemented in MATLAB (Mathworks Inc., Sherborn, MA).

183 For each subject, motion correction to the first functional scan was performed with a six-parameter rigid-

184 body transformation. No excessive motion was found using the criterion of >1 mm instantaneous shift.

185 The structural image was co-registered to the average of the motion-corrected functional images, and then

186 segmented into grey matter, white matter, and cerebrospinal fluid probability maps. The warping

187 parameters obtained from the segmentation algorithm were used to spatially normalize the structural

image to the Montreal Neurological Institute (MNI) standard brain space (output voxel size: 1x1x1 mm³).

189 These parameters were also applied to the functional images to bring them into the same MNI space

190 (output voxel size: 3x3x3 mm³), after which they were spatially smoothed with a Gaussian kernel with a

191 FWHM of 6 mm.

192

193 **2.5. Measurement of ghrelin circulating levels**

In all participants, overnight fasting blood samples were collected between 8 am and 10 am for
the measurement of circulating ghrelin (total). For logistic reasons, blood samples could not be collected

on the same day as the fMRI assessment in 60% of study participants. On average, these samples were collected within the 4 weeks preceding the fMRI scan in the obese group, and within the week preceding the scan in the group of healthy controls. After clotting, samples were centrifuged (1000g, 10 min, 4°C) and sera were stored at -80°C until the assay. Serum total ghrelin concentration was determined by enzyme-linked immunosorbent assay (ELISA), according to the manufacturer specifications (EZGRT-89K, Millipore, Billerica, Massachusetts). Sensitivity was 50 pg/mL and intra- and inter-assay variability were $\pm 1.26\%$ and $\pm 7.81\%$, respectively.

203

204 2.6. Statistical analysis

Participant characteristics, including age, gender, BMI, education level, and circulating levels of ghrelin were compared between the two groups using two-sample *t* tests or Chi-square tests using MATLAB (Mathworks Inc., Sherborn, MA). Two subjects from the control group and 1 subject from the obese group did not properly follow instructions for the performance of the guessing task during the MRI acquisition. Therefore, only data related to ghrelin levels and anatomical brain images were used in those subjects.

Cerebral BOLD responses during the fMRI task were assessed using a general linear model 211 approach similar to the one described by Reuter and collaborators (Reuter et al., 2005). Briefly, for each 212 subject, we set up a linear model with two regressors of interest corresponding respectively to the 213 214 differential "gain vs. loss" response to the outcome (each outcome period modeled as 2 second-blocks), 215 and "net score" (modeled with 4 second-blocks, with a height tracking the current net gain), both time-216 locked to the moment when the outcome was revealed (i.e., when the cards were turned up). The two 217 regressors were convolved with the SPM canonical hemodynamic response function. The motion 218 parameters estimated during the realignment phase were added as potential confounds to the model, 219 which also included an autoregressive component of order 1 plus white noise, to account for the presence 220 of autocorrelation in the residuals, and a basis set of discrete cosine functions modeling a high-pass filter with a cut-off period of 240 sec. A random-effects group analysis was then performed using two-sample 221 222 F-test for identification of any significant "gain vs. loss" statistical effect in any of the two groups (obese

223 and/or control subjects). A similar analysis, but using a one-sample t-test, was conducted to assess the effect of interest separately within the obese and the control groups. Finally, a two-sample t-test was 224 225 performed to compare the effect between the groups. An undirected search (whole-brain analysis) was performed looking for significant voxels with a p<0.05 family wise error rate (FWER), corrected for the 226 227 whole brain volume. For the between-group and single group cluster level inferences, cluster-forming thresholds for the single-voxel significance were set at p<0.001, while the cluster-level threshold was 228 0.05 FWER with a cluster size threshold of $_{k}E>214$ voxels. We wanted to explore the link between 229 230 individual amplitudes of reward-related cerebral activity, ghrelin levels and choice times. Since the dIPFC 231 turned out to be more activated in the obese group compared to controls, we focused on this region of interest (ROI), extracting the eigenvariates of parameter estimates of the "gain vs. loss" contrast using a 232 bilateral spherical ROI of 6 mm radius around the peak of the group effect in dlPFC. The resulting values 233 were then used to test the main and interaction effects between variables of interest: reward-related brain 234 activation, levels of ghrelin and behavior in groups of healthy volunteers and obese subjects with 235 ANCOVA analysis in MATLAB. We calculated the effect size and reported it as Cohen's d for between-236 237 group comparisons and the partial Eta Squared value, $\eta 2p$, for ANCOVA analysis.

238

239

240 3. RESULTS

241 **3.1. Characteristics of study participants**

As shown in **Table 2**, there was no significant difference between obese subjects and non-obese participants in terms of age, gender and education level. The average weight and BMI were respectively 114.7 kg and 41.2 kg/m² in obese subjects *versus* 60.5 kg and 21.8 kg/m² in non-obese controls (all p < 0.0001). Circulating ghrelin levels were significantly lower in obese subjects compared to non-obese controls (375.2±47.6 pg/mL vs. 811.6±87.8 pg/mL; df = 28; p < 0.001, for both parametric t-test and nonparametric Mann-Whitney test, Cohen's *d* = 1.6).

248

250 **3.2. Neuroimaging results**

251 As expected, the fMRI task elicited significant activations in the ventral striatum (left and right sides) (Figure 2A), the extrastriate visual cortex and the ventromedial prefrontal cortex (Table 2) in both 252 groups. While the peak voxels in the left and right ventral striatum showed a greater response in obese 253 254 subjects compared to controls (see bar plot in Figure 2A), the group difference was not significant when 255 controlling for the FWER across the whole-brain, or when using a small-volume-correction for the two 256 ventral striatum clusters. When looking at the reward-related activity separately for each group, the obese 257 subjects showed a larger extent of activation compared to controls (Table 2 and Figure 2B). A direct 258 statistical comparison of the reward-related effect in the two experimental groups showed a greater 259 activation of the left and right dlPFC and the precuneus/posterior cingulate in obese subjects compared to controls (Table 2 and Figure 3). 260

261

262 **3.3. Behavioral performance during the fMRI task**

Choice times after one subsequent win (successful guess), CT1, and choice times after two or more subsequent successful guesses, CT2, were compared between the two groups. No significant difference was found in CT1 (0.96s \pm 0.05s in obese subjects *vs.* 0.92s \pm 0.07s in controls). In contrast, choice times after two or more successive winning trials, i.e. CT2, were significantly longer in obese subjects compared to controls (1.17s \pm 0.07s *vs.* 0.94s \pm 0.05s, df = 25, p<0.05, Cohen's *d* = 0.8) (**Figure 4A**).

269

270 **3.4.** Relationships among reward-related brain activity, behavioral performance and ghrelin levels

ANCOVAs were performed to test for main and interaction effects among the variables of interest; i.e., reward-related brain activation, levels of ghrelin and behavior (choice time, CT), in obese patients and controls. The first ANCOVA, performed on the individual values for the reward-related dlPFC contrasts of parameter estimates (Win-Lose) with group as factor and ghrelin levels as covariate, revealed a strong effect of group, which was not surprising given that the dlPFC ROI had been identified

from the Obese vs. Controls contrast of the fMRI group analysis. Our comparison of interest, however, 276 revealed a significant interaction between groups and ghrelin levels (Figure 4B; F(1, 23)=5.17, p=0.033), 277 278 indicating that reward-related dIPFC activation was more strongly related to lower ghrelin levels in the 279 group of obese subjects than in control subjects. A similar pattern was identified with the ANCOVA performed on choice times, with group as factor and ghrelin levels as covariate. Here also the slope of 280 decreasing choice times with increasing ghrelin levels was steeper in obese subjects compared to controls, 281 indicating that slower choice behavior was more strongly related to reduced ghrelin levels in the group of 282 283 obese subjects (Figure 4C; Group X Ghrelin interaction: F(1,23)=5.26, p=0.031). Finally, the ANCOVA performed on the choice times, with group as factor and the values of reward-related dlPFC contrast of 284 parameter estimates as covariate, showed a robust slowing of choice times with increasing reward-related 285 dlPFC activation (Figure 4D; covariate effect: F(1,23)=9.39, p=0.005), with no significant group 286 differences (F(1,23)=1.01, p=0.33), or interaction effects (F(1,23)=0.18, p=0.68). 287

288

289 4. DISCUSSION

290

291 This study assessed cerebral activity and behavioral response during a guessing task to identify abnormalities in reward outcome-related processing in severely obese subjects. Reward-related 292 293 activations in the ventral striatum and ventromedial prefrontal cortex were observed in both obese and 294 non-obese subjects. In obese subjects, group-specific reward-related activations were found in the 295 bilateral dIPFC and precuneus/posterior cingulate. In both obese and non-obese subjects, reward-related activation in the dIPFC correlated with slower choice times for repeated reward. However, significant 296 297 differences between groups were found with respect to the association of ghrelin with dIPFC activation 298 and reward-related behavior.

The comparable ventromedial prefrontal cortex activation during winning trials in obese and control subjects, and the slight obesity-related trend for hyperactivation in the ventral striatum were unexpected in view of the well-documented alterations in reward system function in obesity (Volkow et al., 2011). This result may be due to the nature of the paradigm/incentive stimuli (money-labeled gaming reinforcement task) used in the present study. In support of this possibility, alterations in reward-related brain activity in obesity have been shown to be task-dependent (Volkow et al., 2011) and, while activity in the ventral striatum can be triggered by both monetary and food reward, BMI was found to correlate only with the anticipatory food-reward processing (Simon et al., 2014).

307 The guessing task paradigm used in the present study produced a significant hyperactivity in the 308 precuneus/posterior cingulate and dIPFC in obese subjects, suggesting the recruitment of a more extended 309 brain network for the processing of reward-related information in obesity. This hypothesis is in line with 310 previous reports indicating greater connectivity between reward network brain regions and regions of 311 executive control, emotional arousal, and somatosensory networks in obese individuals (Gupta and 312 Bhatia, 2008). The increased activation of the precuneus, a region of the default mode, found in the group of obese subjects recruited in the present study is consistent with previous reports documenting increased 313 bilateral functional connectivity in the precuneus (Kullmann et al., 2012), together with greater precuneus 314 315 activity at baseline and in response to food and non-food reward-related cues and during response 316 inhibition and error processing (Bohon, 2017; Hsu et al., 2017; Opel et al., 2015; Tregellas et al., 2011; Zhang et al., 2015) in obese samples. Similarly, greater dIPFC activity and deficits in executive function 317 have been well documented in obesity (Gautier et al., 2000; Holsen et al., 2012; Gentier et al., 2013). 318 Interestingly, in the present study, increased dIPFC activity correlated with slower choice times after 319 320 repeated (but not single) preceding gains/reward. In view of the greater dlPFC activation in obese subjects, this finding may indicate a greater engagement of effortful, self-control executive processes in 321 rewarded decision-making context (Danner et al., 2012; Gentier et al., 2013), consistent with the known 322 link between greater dlPFC activity and greater task difficulty (Manoach et al., 1997) or high-risk choice 323 324 (Bembich et al., 2014). In that perspective, the observed overactivation of brain regions related to selffocusing, i.e., precuneus, (Cavanna and Trimble, 2006) and self-control, i.e., dlPFC, (Belfort-DeAguiar et 325 326 al., 2018) could reflect an increased demand for self-control in rewarding circumstances in obese subjects. 327 A potential alternative explanation is that the occurrence of a string of reward episodes captures more 328 attentional resources in obese subjects, and thus slows down subsequent choices (Kube et al., 2017).

Reward-related over-activation in the dIPFC was associated with reduced circulating levels of 329 ghrelin, and this association was stronger in obese subjects. This finding may be interpreted as somewhat 330 331 surprising in view of previous results indicating increased neural responses to food pictures in brain areas involved in reward processing after ghrelin administration in healthy volunteers (Malik et al., 2008). 332 Nevertheless, it may rely on brain processes differentially involved in reward anticipation and receipt 333 (Oldham et al., 2018). In support of this, prefrontal involvement has been reported specifically at reward 334 335 receipt, and not during reward anticipation. In line with this, obese subjects were found to exhibit dlPFC 336 hyperactivity in response to food images only after meal (Holsen et al., 2012), which is temporally better 337 aligned with reward receipt than anticipation and consistent with results from the present study where dlPFC hyperactivity was measured during reward receipt (gain vs loss trials). This result is also in line 338 with data indicating reduced dIPFC activation in patients with Prader-Willi syndrome, characterized by 339 hyperphagia and hyperghrelinemia, and increased dlPFC activity in obese subjects with moderate 340 overeating and low ghrelin (Belfort-DeAguiar et al., 2018; Holsen et al., 2012). 341

342 Similar to the relationship found with the dlPFC, decreased ghrelin levels were found to 343 significantly correlate with slower choice times, particularly in the obese group. This finding is consistent with the role of the ghrelin system in reward and decision-making processes. Given the recent 344 345 experimental evidence for a role of ghrelin in impulsive behavior/choice (Anderberg et al., 2016), it is 346 possible that this association relies on the effects of the ghrelin system on the central/brain regulation of processes related to self-control. In line with this, higher fasting ghrelin levels were recently found to be 347 associated with greater reward sensitivity and reduced self-control/increased impulsivity (Ralevski et al., 348 2018). Alternatively, this association may also reflect increased stress-related processes or greater worry 349 350 of unsuccessful outcome in obese subjects given the associations previously documented between ghrelin and increased susceptibility/altered response to stress (Sarker et al., 2013; Spencer et al., 2012). 351

Albeit they strongly support the link between the ghrelin system and reward-related and decisionmaking processes in obesity, results from the present study do not provide any information regarding the temporality and causality of events nor their role in the development of overweight/obesity. While previous studies addressing reward processing alterations have shown that hedonically driven food intake, along with stress-induced comfort eating, may contribute to weight gain and obesity maintenance (Burger
and Berner, 2014; Lee and Dixon, 2017), further large-scale longitudinal studies are needed to
specifically address this issue.

359 This study presents certain limitations that future investigations should also overcome. Due to logistic constraints inherent to clinical research on patients (ghrelin measurement was part of the routine 360 preoperative evaluation of obese patients), ghrelin levels were not measured on the same day of the fMRI 361 acquisition in 60% of the participants. While this limitation should not critically impact results from the 362 363 present study, based on data from prospective studies showing stable fasting ghrelin levels with time in 364 obese subjects (Hanusch-Enserer et al., 2004) and healthy volunteers (Leidy et al., 2004), future investigations with concomitant measurement of ghrelin and brain activation are needed to comfort the 365 present findings. The exclusive measurement of total ghrelin, and not the active acylated form of the 366 hormone, represents another limitation of this study. Because of its high instability at room temperature, 367 the exact measurement of the active form of ghrelin is highly challenging and requires specific 368 measures/procedures for blood collection and processing (Hosoda and Kangawa, 2004) that were not 369 370 planned in the present study. While complementary measurements of the active form of ghrelin are highly 371 encouraged in future studies, we believe that assessment of total ghrelin remains highly relevant in the 372 context of the present study since it was shown to be highly sensitive and reliable in multiple studies of 373 obesity. In particular, total ghrelin, was found to correlate with BMI and disease severity in most cases of 374 obesity (except Prader-Willi Syndrome) and to be reduced in hyperphagia and obesity (Nonogaki, 2008; 375 Monti et al., 2006). These associations were not found with acylated ghrelin that seems to be more 376 relevant with respect to metabolic abnormalities, notably in normal subjects (Pacifico et al., 2009). Another aspect that would merit further investigation is the relationship between ghrelin and insulin 377 378 and/or other neuroendocrine factors, given the known interactions between these different factors and 379 their possible impact on reward signaling (Baicy et al., 2007; Farooqi et al., 2007; Kullmann et al., 2012). 380 In support of this, strong relationships exist between the ghrelin system and the insulin system [e.g. ghrelin and its receptors are expressed within the pancreas and β -cells and ghrelin alters insulin-381 dependent glucose metabolism in a direct and non-direct manner (Churm et al., 2017)], which has been 382 shown to be involved in the regulation of reward processing in preclinical and clinical studies (Eisenstein 383

et al., 2015; Kleinridders and Pothos, 2019). Finally, another limitation of the study comes from the use of an ROI (dlPFC) identified from the group Win-Lose contrast for subsequent correlational analyses of brain activation with ghrelin and behavior. In this situation, a complete statistical independence may not be assured nor the possibility of additional associations with other brain structures be excluded. While these issues merit further investigation in future studies, they should not substantially influence the findings given that the ROI defined by a between-group test was used for assessing within-group relationships.

In conclusion, the present findings indicate an increased activation of brain areas related to selfcontrol and self-referencing during processing of non-food reward in obese subjects. Moreover, fasting levels of circulating ghrelin, the hormone regulating energy balance and food consuming behavior, were associated with generalized reward processing, both at the behavioral and neurobiological levels. The associations of ghrelin with decision times and dIPFC activity suggest the potential involvement of this hormone in the obesity-related behavioral alterations.

397

398

Acknowledgments: This study was achieved within the context of the Laboratory of Excellence TRAIL
ANR-10-LABX-57. The authors thank Professor V. Doucet and his team for their help and assistance
with the fMRI assessments.

402 **REFERENCES**

- 403 Abizaid, A., 2019. Stress and obesity: The ghrelin connection. Journal of neuroendocrinology, e12693.
- 404 Akitsuki, Y., Sugiura, M., Watanabe, J., Yamashita, K., Sassa, Y., Awata, S., Matsuoka, H., Maeda, Y.,
- 405 Matsue, Y., Fukuda, H., Kawashima, R., 2003. Context-dependent cortical activation in response to
- 406 financial reward and penalty: an event-related fMRI study. Neuroimage. 19, 1674-1685.
- 407 Anderberg, R.H., Hansson, C., Fenander, M., Richard, J.E., Dickson, S.L., Nissbrandt, H., Bergquist, F.,
- 408 Skibicka, K.P., 2016. The Stomach-Derived Hormone Ghrelin Increases Impulsive
- 409 Behavior.Neuropsychopharmacology 41, 1199-1209.
- Andrews, Z.B., 2011. The extra-hypothalamic actions of ghrelin on neuronal function. Trends Neurosci
 34, 31-40.
- Angelidis, G., Valotassiou, V., Georgoulias, P., 2010. Current and potential roles of ghrelin in clinical
 practice. Journal Endocrinol Invest 33, 823-838.
- 414 Baicy, K., London, E.D., Monterosso, J., Wong, M.L., Delibasi, T., Sharma, A., Licinio, J., 2007. Leptin
- 415 replacement alters brain response to food cues in genetically leptin-deficient adults. Proc Natl Acad
 416 Sci U S A 104, 18276-18279.
- 417 Balodis, I.M., Kober, H., Worhunsky, P.D., White, M.A., Stevens, M.C., Pearlson, G.D., Sinha, R., Grilo,
- 418 C.M., Potenza, M.N., 2013. Monetary reward processing in obese individuals with and without binge
- 419 eating disorder. Biol Psychiatry 73, 877-886.
- 420 Belfort-DeAguiar, R., Seo, D., Lacadie, C., Naik, S., Schmidt, C., Lam, W., Hwang, J., Constable, T.,
- Sinha, R., Sherwin, R.S., 2018. Humans with obesity have disordered brain responses to food images
 during physiological hyperglycemia. Am J Physiol Endocrinol Metab 314, E522-E529.
- 423 Bembich, S., Clarici, A., Vecchiet, C., Baldassi, G., Cont, G., Demarini, S., 2014. Differences in time
- 424 course activation of dorsolateral prefrontal cortex associated with low or high risk choices in a
- 425 gambling task. Front Hum Neurosci 8, 464.
- Bohon, C., 2017. Brain response to taste in overweight children: A pilot feasibility study. PLoS One 12,
- 427 e0172604.

- 428 Briggs, D.I., Lockie, S.H., Wu, Q., Lemus, M.B., Stark, R., Andrews, Z.B., 2013. Calorie-restricted
- 429 weight loss reverses high-fat diet-induced ghrelin resistance, which contributes to rebound weight
- 430 gain in a ghrelin-dependent manner. Endocrinology 154, 709-717.
- Burger, K.S., Berner, L.A., 2014. A functional neuroimaging review of obesity, appetitive hormones and
 ingestive behavior. Physiol Behav 136, 121-127.
- 433 Capuron, L., Pagnoni, G., Drake, D.F., Woolwine, B.J., Spivey, J.R., Crowe, R.J., Votaw, J.R., Goodman,
- 434 M.M., Miller, A.H., 2012. Dopaminergic mechanisms of reduced basal ganglia responses to hedonic
- reward during interferon alfa administration. Arch Gen Psychiatry 69, 1044-1053.
- 436 Carnell, S., Gibson, C., Benson, L., Ochner, C.N., Geliebter, A., 2012. Neuroimaging and obesity: current
- 437 knowledge and future directions. Obes Rev 13, 43-56.
- Cavanna, A.E., Trimble, M.R., 2006. The precuneus: a review of its functional anatomy and behavioural
 correlates. Brain. 129, 564-583.
- Churm, R., Davies, J.S., Stephens, J.W., Prior, S.L., 2017. Ghrelin function in human obesity and type 2
 diabetes: a concise review. Obes Rev 18, 140-148.
- 442 Danner, U.N., Ouwehand, C., van Haastert, N.L., Hornsveld, H., de Ridder, D.T., 2012. Decision-making
- impairments in women with binge eating disorder in comparison with obese and normal weight
- 444 women. Eur Eat Disord Rev 20, e56-62.
- 445 Date, Y., 2012. Ghrelin and the vagus nerve. Methods Enzymol 514, 261-269.
- 446 Eisenstein, S.A., Gredysa, D.M., Antenor-Dorsey, J.A., Green, L., Arbeláez, A.M., Koller, J.M., KJ, B.,
- JS, P., SM, M., Hershey, T., 2015. Insulin, Central Dopamine D2 Receptors, and Monetary Reward
 Discounting in Obesity. PloS one 10(7), e0133621.
- 449 English, P.J., Ghatei, M.A., Malik, I.A., Bloom, S.R., Wilding, J.P., 2002. Food fails to suppress ghrelin
- 450 levels in obese humans. J Clin Endocrinol Metab 87, 2984.
- 451 Farooqi, I.S., Bullmore, E., Keogh, J., Gillard, J., O'Rahilly, S., Fletcher, P.C., 2007. Leptin regulates
- 452 striatal regions and human eating behavior. Science 317, 1355.
- 453 Foster, C.M., Barkan, A., Kasa-Vubu, J.Z., Jaffe, C., 2007. Ghrelin concentrations reflect body mass
- index rather than feeding status in obese girls. Pediatr Res 62, 731-734.

- 455 Gautier, J.F., Chen, K., Salbe, A.D., Bandy, D., Pratley, R.E., Heiman, M., Ravussin, E., Reiman, E.M.,
- Tataranni, P.A., 2000. Differential brain responses to satiation in obese and lean men. Diabetes 49,
 838-846.
- 458 Gentier, I., Augustijn, M., Deforche, B., Tanghe, A., De Bourdeaudhuij, I., Lenoir, M., D'Hondt, E.,
- 459 2013. A comparative study of performance in simple and choice reaction time tasks between obese460 and healthy-weight children. Res Dev Disabil 34, 2635-2641.
- 461 Goldstone, A.P., Prechtl, C.G., Scholtz, S., Miras, A.D., Chhina, N., Durighel, G., Deliran, S.S.,
- 462 Beckmann, C., Ghatei, M.A., Ashby, D.R., Waldman, A.D., Gaylinn, B.D., Thorner, M.O., Frost,
- 463 G.S., Bloom, S.R., Bell, J.D., 2014. Ghrelin mimics fasting to enhance human hedonic, orbitofrontal
- 464 cortex, and hippocampal responses to food. Am J Clin Nutr 99, 1319-1330.
- 465 Gupta, R., Bhatia, M.S., 2008. Short-lasting unilateral neuralgiform headache with conjunctival injection
- 466 and tearing: response to antiepileptic dual therapy. Neurol India 56, 71-73.
- 467 Hanusch-Enserer, U., Cauza, E., Brabant, G., Dunky, A., Rosen, H., Pacini, G., Tuchler, H., Prager, R.,
- 468 Roden, M., 2004. Plasma ghrelin in obesity before and after weight loss after laparoscopical
- 469 adjustable gastric banding. J Clin Endocrinol Metab 89, 3352-3358.
- 470 Holsen, L.M., Savage, C.R., Martin, L.E., Bruce, A.S., Lepping, R.J., Ko, E., Brooks, W.M., Butler,
- 471 M.G., Zarcone, J.R., Goldstein, J.M., 2012. Importance of reward and prefrontal circuitry in hunger
- 472 and satiety: Prader-Willi syndrome vs simple obesity. Int J Obes (Lond). 36, 638-647.
- 473 Hosoda, H., Kangawa, K. 2004. Ghrelin Measurement: Present and Perspectives. In: Ghigo E., Benso A.,
- 474 Broglio F. (eds) Ghrelin. Endocrine Updates, vol 23. Springer, Boston, MA.
- 475 Howick, K., Griffin, B.T., Cryan, J.F., Schellekens, H., 2017. From Belly to Brain: Targeting the Ghrelin
- 476 Receptor in Appetite and Food Intake Regulation. Int J Mol Sci 18.
- 477 Hsu, J.S., Wang, P.W., Ko, C.H., Hsieh, T.J., Chen, C.Y., Yen, J.Y., 2017. Altered brain correlates of
- 478 response inhibition and error processing in females with obesity and sweet food addiction: A
- 479 functional magnetic imaging study. Obes Res Clin Pract 11, 677-686.
- 480 Kleinridders, A. Pothos, E.N., 2019. Impact of brain insulin signaling on dopamine function, food intake,
- reward, and emotional behavior. Curr Nutr Rep 8, 83-91.

- Kube, J., Mathar, D., Horstmann, A., Kotz, S.A., Villringer, A., Neumann, J., 2017. Altered monetary
 loss processing and reinforcement-based learning in individuals with obesity. Brain Imaging Behav
 12, 1431-1449.
- 12, 1451-1449.
- 485 Kullmann, S., Heni, M., Veit, R., Ketterer, C., Schick, F., Haring, H.U., Fritsche, A., Preissl, H., 2012.
- 486 The obese brain: association of body mass index and insulin sensitivity with resting state network
- 487 functional connectivity. Hum Brain Mapp 33, 1052-1061.
- Lee, P.C., Dixon, J.B., 2017. Food for Thought: Reward Mechanisms and Hedonic Overeating in Obesity.
 Curr Obes Rep 6, 353-361.
- 490 Leidy, H.J., Gardner, J.K., Frye, B.R., Snook, M.L., Schuchert, M.K., Richard, E.L., Williams, N.I.,
- 2004. Circulating ghrelin is sensitive to changes in body weight during a diet and exercise program in
 normal-weight young women. J Clin Endocrinol Metab 89, 2659-64.
- Malik, S., McGlone, F., Bedrossian, D., Dagher, A., 2008. Ghrelin modulates brain activity in areas that
 control appetitive behavior. Cell Metab 7, 400-409.
- 495 Manoach, D.S., Schlaug, G., Siewert, B., Darby, D., Bly, B., Benfield, A., Edelman, R., Warach, S.,
- 496 1997. Prefrontal cortex fMRI signal changes are correlated with working memory load. Neuroreport.
 497 8, 545-549.
- 498 Monti, V., Carlson, J.J., Hunt, S.C., Adams, T.D., 2006. Relationship of ghrelin and leptin hormones with
- body mass index and waist circumference in a random sample of adults. J Am Diet Assoc 106, 822-828.
- 501 Muller, T.D., Nogueiras, R., Andermann, M.L., Andrews, Z.B., Anker, S.D., Argente, J., Batterham,
- 502 R.L., Benoit, S.C., Bowers, C.Y., Broglio, F., Casanueva, F.F., D'Alessio, D., Depoortere, I.,
- 503 Geliebter, A., Ghigo, E., Cole, P.A., Cowley, M., Cummings, D.E., Dagher, A., Diano, S., Dickson,
- 504 S.L., Dieguez, C., Granata, R., Grill, H.J., Grove, K., Habegger, K.M., Heppner, K., Heiman, M.L.,
- 505 Holsen, L., Holst, B., Inui, A., Jansson, J.O., Kirchner, H., Korbonits, M., Laferrere, B., LeRoux,
- 506 C.W., Lopez, M., Morin, S., Nakazato, M., Nass, R., Perez-Tilve, D., Pfluger, P.T., Schwartz, T.W.,
- 507 Seeley, R.J., Sleeman, M., Sun, Y., Sussel, L., Tong, J., Thorner, M.O., van der Lely, A.J., van der
- 508 Ploeg, L.H., Zigman, J.M., Kojima, M., Kangawa, K., Smith, R.G., Horvath, T., Tschop, M.H., 2015.
- 509 Ghrelin. Mol Metab 21, 437-60.

- 510 Natalucci, G., Riedl, S., Gleiss, A., Zidek, T., Frisch, H., 2005. Spontaneous 24-h ghrelin secretion
- 511 pattern in fasting subjects: maintenance of a meal-related pattern. Eur J Endocrinol. 152, 845-850.
- 512 Nonogaki, K., 2008. Ghrelin and feedback systems. Vit Horms 77, 149-170.
- 513 Oldham, S., Murawski, C., Fornito, A., Youssef, G., Yucel, M., Lorenzetti, V., 2018. The anticipation and
- 514 outcome phases of reward and loss processing: A neuroimaging meta-analysis of the monetary
- 515 incentive delay task. Hum Brain Mapp 39, 3398-3418.
- 516 Opel, N., Redlich, R., Grotegerd, D., Dohm, K., Haupenthal, C., Heindel, W., Kugel, H., Arolt, V.,
- 517 Dannlowski, U., 2015. Enhanced neural responsiveness to reward associated with obesity in the
- bisence of food-related stimuli. Hum Brain Mapp 36, 2330-2337.
- 519 Pacifico, L., Poggiogalle, E., Costantino, F., Anania, C., Ferraro, F., Chiarelli, F., Chiesa, C., 2009.
- 520 Acylated and nonacylated ghrelin levels and their associations with insulin resistance in obese and
- 521 normal weight children with metabolic syndrome. Eur J Endocrinol 161, 861-870.
- Perello, M., Dickson, S.L., 2015. Ghrelin signalling on food reward: a salient link between the gut and the
 mesolimbic system. J Neuroendocrinol 27, 424-434.
- Price, M., Higgs, S., Maw, J., Lee, M., 2016. A dual-process approach to exploring the role of delay
 discounting in obesity. Physiol Behav 162, 46-51.
- 526 Pursey, K.M., Stanwell, P., Callister, R.J., Brain, K., Collins, C.E., Burrows, T.L., 2014. Neural
- 527 Responses to Visual Food Cues According to Weight Status: A Systematic Review of Functional
- 528 Magnetic Resonance Imaging Studies. Front Nutr 1, 7.
- 529 Puzziferri, N., Zigman, J.M., Thomas, B.P., Mihalakos, P., Gallagher, R., Lutter, M., Carmody, T., Lu,
- 530 H., Tamminga, C.A., 2016. Brain imaging demonstrates a reduced neural impact of eating in obesity.
- 531 Obesity (Silver Spring, Md.) 24, 829-836.
- 532 Ralevski, E., Shanabrough, M., Newcomb, J., Gandelman, E., Hayden, R., Horvath, T.L., Petrakis, I.,
- 533 2018. Ghrelin is Related to Personality Differences in Reward Sensitivity and Impulsivity. Alcohol
- 534 Alcohol. 53, 52-56.
- 535 Reuter, J., Raedler, T., Rose, M., Hand, I., Glascher, J., Buchel, C., 2005. Pathological gambling is linked
- to reduced activation of the mesolimbic reward system. Nat Neurosci. 8, 147-148.

- 537 Revitsky, A.R, Klein, L.C., 2013. Role of ghrelin in drug abuse and reward-relevant behaviors: a
- 538 burgeoning field and gaps in the literature. Curr Drug Abuse Rev 6, 213-44.
- 539 Sarker, M.R., Franks, S., Caffrey, J., 2013. Direction of post-prandial ghrelin response associated with
- 540 cortisol response, perceived stress and anxiety, and self-reported coping and hunger in obese women.
- 541 Behav Brain Res 257, 197-200.
- 542 Simon, J.J., Skunde, M., Hamze Sinno, M., Brockmeyer, T., Herpertz, S.C., Bendszus, M., Herzog, W.,
- 543 Friederich, H.C., 2014. Impaired Cross-Talk between Mesolimbic Food Reward Processing and
- 544 Metabolic Signaling Predicts Body Mass Index. Front Behav Neurosci 8, 359.
- 545 Solomou, S., Korbonits, M., 2014. The role of ghrelin in weight-regulation disorders: implications in
- 546 clinical practice. Hormones (Athens) 13, 458-475.
- 547 Spencer, S.J., Xu, L., Clarke, M.A., Lemus, M., Reichenbach, A., Geenen, B., Kozicz, T., Andrews, Z.B.,
- 548 2012. Ghrelin regulates the hypothalamic-pituitary-adrenal axis and restricts anxiety after acute
- 549 stress. Biol Psychiatry 72, 457-465.
- 550 Tregellas, J.R., Wylie, K.P., Rojas, D.C., Tanabe, J., Martin, J., Kronberg, E., Cordes, D., Cornier, M.A.,
- 551 2011. Altered default network activity in obesity. Obesity (Silver Spring, Md.) 19, 2316-2321.
- 552 Tschöp, M., Weyer, C., Tataranni, P.A., Devanarayan, V., Ravussin, E., Heiman, M.L., 2001.Circulating
- 553 ghrelin levels are decreased in human obesity. Diabetes 50, 707-9.
- 554 Uriarte, M., De Francesco, P.N., Fernandez, G., Cabral, A., Castrogiovanni, D., Lalonde, T., Luyt, L.G.,
- 555 Trejo, S., Perello, M., 2018. Evidence Supporting a Role for the Blood-Cerebrospinal Fluid Barrier
- 556 Transporting Circulating Ghrelin into the Brain. Mol Neurobiol 56, 4120-4134.
- 557 Volkow, N.D., Wang, G.J., Baler, R.D., 2011. Reward, dopamine and the control of food intake:
- implications for obesity. Trends Cogn Sci. 15, 37-46.
- 559 Weiskopf, N., Hutton, C., Josephs, O., Turner, R., Deichmann, R., 2007. Optimized EPI for fMRI studies
- of the orbitofrontal cortex: compensation of susceptibility-induced gradients in the readout direction.
- 561 Magma. 20, 39-49.
- 562 Yildiz BO, Suchard MA, Wong M-L, McCann SM, J., L., 2004. Alterations in the dynamics of
- circulating ghrelin, adiponectin, and leptin in human obesity. Proc Natl Acad Sci USA 101, 10434-
- 564 10439.

- 565 Zhang, B., Tian, D., Yu, C., Li, M., Zang, Y., Liu, Y., Walter, M., 2015. Altered baseline brain activity
- differentiates regional mechanisms subserving biological and psychological alterations in obese men.
 Sci Rep 5, 11563.
- 568 Zigman, J.M., Bouret, S.G., Andrews, Z.B., 2016. Obesity Impairs the Action of the Neuroendocrine
- 569 Ghrelin System. Trends Endocrinol Metab 27, 54-63.
- 570

572 **TABLES**

573

574 Table 1. Characteristics of study participants

	Obese	Normal-weight			
	(N=15)	Controls	Difference		
		(N=15)	P value		
Age, years (SD)	38.7 (9.4)	37 (7.1)	0.6		
Gender, men/women (%)	3/12 (20, 80)	2/13 (13.3, 86.7)	0.6		
Body Mass Index (BMI), kg/m ²	41.2 (2.1)	21.8 (2.6)	< 0.0001		
Weight, kg (SD)	114.7 (9.1)	60.5 (10)	< 0.0001		
Diploma ¹	2.9 (0.96)	3.4 (1.40)	0.3		
Ghrelin levels, pg/mL (SD)	375.2 (47.6)	811.6 (87.8)	< 0.001		

575

 1 Diploma was measured on a scale from 0 (no diploma) to 7 (PhD or MD).

577 Data are shown as mean (SD) or prevalence (%). The statistical significance of group differences

578 (*p*-value) was assessed by Student t-tests (Chi-square test, for Gender).

580 Table 2. Brain regions activated by the reward condition (winning trials > losing trials) in

581 **obese and non-obese controls**

			Peak coordinates		
Brain Region	k _E	T-	X	Y	Z
		score			
Non-obese controls (cluster-forming threshold p<0.001)					
Ventral striatum	317	9.65*	9	17	-5
Orbitofrontal cortex	107	7.40	21	29	-14
Extrastriate visual cortex (L)	466	9.78*	-36	-88	-11
Extrastriate visual cortex (L)	40	6.73	-39	-46	-23
Extrastriate visual cortex (R)	500	9.42*	42	-64	-20
Posterior cingulate cortex	45	7.68	6	-31	28
Intraparietal sulcus	76	6.94	21	-61	34
<i>Obese subjects (cluster-forming threshold p<0.001)</i>					
Ventral striatum + ventromedial prefrontal cortex +	681	9.30*	18	47	-11
orbitofrontal cortex					
Extrastriate visual cortex (R)	720	8.78*	30	-88	4
Extrastriate visual cortex (L)	263	6.93	-27	-82	-11
S1(R) + precuneus + posterior cingulate cortex	1516	7.84	39	-22	52
Intraparietal sulcus	159	4.08	-27	-70	52
S1(R)	79	6.17	57	-13	34
Frontal operculum/premotor cortex	93	4.01	-42	2	13
Obese subjects > non-obese controls					
(cluster-forming threshold p<0.001)					
Dorsolateral prefrontal cortex (R)	81	6.28*	24	17	37
Dorsolateral prefrontal cortex (L)	56	4.92	-30	23	40
Precuneus / posterior cingulate	126	5.32	-6	-43	43

582

584 The other clusters are significant at the cluster level, but not voxel level (FWER p<0.05, at the

cluster level), k_E is the cluster size in voxels. Obese subject, N=14; non-obese controls, N=13.

⁵⁸³ *Clusters where the T-score of the peak voxel is significant (FWER p<0.05, at the voxel level).

586 FIGURE LEGENDS

587

588 Figure 1. Reward-task paradigm

Panel A: Full timeline of the stimulus presentation design, which was identical across subjects.
The net score is indicated by the noisy sinusoid, along with gain and loss trials (white and grey
bars)

592 Panel B: Trial structure. Examples of trials 2-4 are shown. Red cards indicate gain, black cards
593 indicate loss. Subjects could choose either left or right card, but the outcome in terms of win or
594 loss was predefined (see A).

595

596 Figure 2. Reward-related responses in obese subjects and non-obese controls

597 Panel A: Reward-related response in the ventral striatum across both experimental groups, as the

result of the two-sample (obese, N=14 or controls, N=13) "gain vs. loss" F-test.

599 Panel B: Group-wise reward-related activation. Note that the activation in obese subjects

600 extends to precuneus/posterior cingulate and dorsolateral prefrontal cortex (dlPFC), unlike in

601 non-obese controls (within-group activation).

602

603 Figure 3. Reward-related activity is greater in obese subjects

604 The statistical contrast Obese (N=14) > Controls (N=13), thresholded for illustration purposes at

cluster forming threshold p<0.005 is mapped onto the average of the individual subjects'

normalized anatomical images (n=30). Significant clusters and respective peak contrast estimates

are shown: (A) left dorsolateral prefrontal cortex, (B) right dorsolateral prefrontal cortex and (C)

608 precuneus/posterior cingulate.

609

610 Figure 4. Choice times and relationship with dorsolateral prefrontal cortex (dlPFC)

611 activity and ghrelin levels in obese subjects and non-obese controls

Panel A. Choice time (CT) group effect. A significant difference between groups was observed
for choice time during repeated reward (CT2) but not during single reward (CT1); * p<0.05.

614 **Panel B**. Relationship between ghrelin levels and reward-related activity in the dlPFC. The

615 parameter estimates of the "Gain vs. Loss" contrast were extracted from the bilateral dlPFC (i.e.,

the first eigenvariates of the individual contrast images within two spherical ROI of 6 mm radius

centered on the peaks' coordinates of [24 17 37] and [-30 23 40] were extracted; as shown in a

right upper corner). Main effects and interactions: Group: F=14.6, p<0.001, η 2p=0.39; Ghrelin:

619 F=9.4, p<0.01, η 2p=0.29; Group x Ghrelin: F=5.2, p<0.05, η 2p=0.18.

620 **Panel C**. Relationship between choice time during repeated reward (CT2) and ghrelin levels.

621 Corresponding CIs displayed as gray shadows. Main effects and interactions: Group: F=5.9,

622 p<0.05, η2p=0.20; Ghrelin: F=4.9, p<0.05, η2p=0.18; Group x Ghrelin: F=5.3, p<0.05,

623 η2p=0.19.

624 **Panel D.** Relationship between choice time during repeated reward (CT2) and reward-related

activity in the dlPFC. The parameter estimates of the "Gain vs. Loss" contrast were extracted

626 from the bilateral dlPFC (i.e., the first eigenvariates of the individual contrast images within two

spherical ROI of 6 mm radius centered on the peaks' coordinates of [24 17 37] and [-30 23 40]

were extracted; as shown in a right upper corner). Main effect: dlPFC: F=9.4, p<0.01, η 2p=0.29.

629 ○control subjects, N=13 ●obese subjects, N=14





vStr L

Α.

Reward F-score

FWE p<0.05













