

A fragmentation phenomenon for a non-energetic optimal control problem: optimisation of the total population size in logistic diffusive models

Idriss Mazari, Domènec Ruiz-Balet

▶ To cite this version:

Idriss Mazari, Domènec Ruiz-Balet. A fragmentation phenomenon for a non-energetic optimal control problem: optimisation of the total population size in logistic diffusive models. 2020. hal-02523753v1

HAL Id: hal-02523753 https://hal.science/hal-02523753v1

Preprint submitted on 29 Mar 2020 (v1), last revised 21 Oct 2020 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A fragmentation phenomenon for a non-energetic optimal control problem:

optimisation of the total population size in logistic diffusive models

Idriss Mazari^{*}, Domènec Ruiz-Balet^{†‡}

March 29, 2020

Abstract

Following the recent works [9, 17, 30, 31, 37], we investigate the problem of optimising the total population size for logistic diffusive models with respect to resources distributions. Using the spatially heterogeneous Fisher-KPP equation, we obtain a surprising fragmentation phenomenon: depending on the scale of diffusivity (i.e the dispersal rate), it is better to either concentrate or fragment resources. Our main result is that, the smaller the dispersal rate of the species in the domain, the more optimal resources distributions tend to oscillate. This is in sharp contrast with other criteria in population dynamics, such as the classical problem of optimising the survival ability of a species, where concentrating resources is always favourable, regardless of the diffusivity. Our study is completed by numerous numerical simulations that confirm our results.

Keywords: diffusive logistic equation, optimal control, shape optimization.

AMS classification: 35Q92,49J99,34B15.

Contents

Introduction			
1.1	Scope of this article: fragmentation and concentration for spatial ecology		
1.2	Setting and bibliographical references		
1.3	Main result		
Proof of Theorem 1 2.1 The influence of periodisation			
2.2	Technical preliminaries		

*Sorbonne Universités, UPMC Univ Paris 06, UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France (idriss.mazari@sorbonne-universite.fr)

[†]Chair of Computational Mathematics, Fundación Deusto, Av. de las Universidades, 24, 48007 Bilbao, Basque Country, Spain

[‡]Departamento de Mateméticas, Universidad Autónoma de Madrid, 28049 Madrid, Spain

3	Numerical simulations			
	3.1	Simulations in the one-dimensional case		
		3.1.1 $\kappa = 1, m_0 = 0.3$		
		3.1.2 $\kappa = 1, m_0 = 0.6 \dots \dots$		
	3.2	Simulations in the two-dimensional case		
		3.2.1 $\kappa = 1, m_0 = 0.3$		
		3.2.2 $\kappa = 1, m_0 = 0.6 \dots \dots$		

Acknowledgment. This work was started during a research stay of the authors at the the Friedrich-Alexander-Universität at the invitation of Enrique Zuazua and the authors wish to thank E. Zuazua and the FAU University for their hospitality.

I. Mazari was supported by the French ANR Project ANR-18-CE40-0013 - SHAPO on Shape Optimization.

D. Ruiz-Balet was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 694126-DyCon).

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No.765579-ConFlex, grant MTM2017-92996 of MINECO (Spain), ICON of the French ANR and "Nonlocal PDEs: Analysis, Control and Beyond", AFOSR Grant FA9550-18-1-0242 and the Alexander von Humboldt-Professorship program.

1 Introduction

1.1 Scope of this article: fragmentation and concentration for spatial ecology

In this article, we study a problem of great relevance in the field of spatial ecology. Namely, considering a species dispersing in a domain where some resources are available:

How should we spread resources so as to maximise the total population size at equilibrium?

Here, we focus on a fine qualitative analysis of this question and emphasise the crucial role of the characteristic diffusion rate of the population (or, equivalently, of the size of the domain).

Regarding mathematical biology, spatially heterogeneous models are of paramount importance, as acknowledged, for instance, in [11]. Natural questions arise when considering such models: one may for instance think of spatially heterogeneous systems of reaction-diffusion equations, in which case a relevant question is that of existence and stability of (non-trivial) equilibria (see for instance [12, 14, 13, 28]).

Here, we focus on single-species models, in which case two problems have drawn a lot of attention from both the mathematical and the mathematical biology communities: the problem of optimal survival ability, and the problem of optimising the total population size. We expand on bibliographical references in Subsection 1.2 of this Introduction, but let us stress the following fact: while the optimisation of the survival ability with respect to resources distribution is fairly well-understood (in terms of qualitative analysis, see for instance [3, 18, 22]), the problem of the total population size, which has been the subjects of several recent articles (we refer for instance to [2, 9, 17, 30, 31, 37]) is still elusive when considered from a qualitative point of view. For instance, for the optimal survival ability, the following paradigm has been established:

Concentration of resources favors survival ability.

This was first observed in [32], and given a proper mathematical analysis in [3], in terms of rearrangements. One of the other conclusions of [3] is that heterogeneity is favorable to survival ability: under natural assumptions (made precise in Section 1.2 throught the definition of the admissible class, Equation (2)), in order to maximise the survival ability of a population, one should work with patch-models. Here, this means the following: provided the population evolves in Ω and the resources distributions $m : \Omega \to \mathbb{R}$ satisfy pointwise $(0 \leq m \leq 1)$ and integral $(\int_{\Omega} m \leq C)$ bounds, the optimal resources distribution for survival ability m^* satisfies $\Omega = \{m = 0\} \sqcup \{m = 1\}$. Such results generally do not depend on the dispersal rate: regardless of this characteristic speed, resources distributions should be concentrated if we want to optimise the survival ability.

The problem of optimising the total population size, on the other hand, is much more complicated to tackle at the mathematical level. One of the main questions that have been investigated is the influence of diffusion on the population size criterion (which in some models favours the total population size, see [25]), and we refer to [37], as well as the recent survey [9] for a biological perspective on this question. In these two last references, the following question is also presented: can the total population size exceed the total amount of resources? This question, for the model we are going to consider, has been solved in dimension 1 in [2] and, in dimension $n \ge 2$, in the recent preprint [17]. In all of these papers, the dispersal rate plays a crucial role in the analysis.

Regarding qualitative properties, as will be explained further in Section 1.2, very few things are known. The relevance of patch-models for this optimisation problem has been investigated in [30] and [31], but, so far, the only qualitative results can be found in [30]: for large dispersal rates, concentrating resources favours the total population size while, for small diffusivities, fragmentation (i.e. scattering resources across the domain) may be better.

The goals of this article is to give a complete treatment of the case of small dispersal rates for the logistic-diffusive Fisher-KPP equation, and our main result, Theorem 1, may be interpreted as follows:

To maximise the total population size, the smaller the diffusivity, the more one should fragment resources.

From a calculus of variations (or optimal control) perspective, our article's main innovation is that it gives a qualitative analysis of a non-energetic optimisation problems. Such problems are notoriously hard to analyse, given that their structure prohibits using classical tools (e.g rearrangements, symmetrisation) and that the analysis of optimality conditions is very tricky. Here, we propose an approach relying on strong non-monotonicity properties of the functional that is to be optimised.

Finally, we provide several numerical experiments that validate our results. This article is organised as follows:

- In Section 1.2, we present the model under consideration as well as the variational problem under consideration and recall the several qualitative properties available in the literature.
- In Section 1.3, we state our main result.
- In Section 2, we prove our main result.
- In Section 3, we give several numerical simulations to illustrate Theorem 1.

1.2 Setting and bibliographical references

We are working here with the spatially heterogeneous Fisher-KPP equation (which originated in the seminal [10, 20]). Let us consider, in *n*-dimensions, the box

$$\Omega = \prod_{i=1}^{n} [0;1],$$

which will serve as our domain. We could consider more general boxes $\Omega = \prod_{i=1}^{n} (0; a_i)$, but the results and proofs would be the same. We consider a positive parameter $\mu > 0$, which will be referred to as dispersal rate, or diffusivity. To model the spatial heterogeneity, we use resources distributions, i.e functions $m : \Omega \to \mathbb{R}$. Finally, we take into account an intra-specific, non-linear reaction term from the classical Malthusian equation. This gives the following equation: assuming the population density $\theta_{m,\mu}$ has reached an equilibrium, it solves

$$\begin{cases} \mu \Delta \theta_{m,\mu} + \theta_{m,\mu} \left(m - \theta_{m,\mu} \right) = 0 \text{ in } \Omega, \\\\ \frac{\partial \theta_{m,\mu}}{\partial \nu} = 0 \text{ on } \partial \Omega, \\\\ \theta_{m,\mu} > 0 \text{ in } \Omega. \end{cases}$$
(1) {LDE}

For Equation (1) to have a solution, one must restrict the class of resources distributions. If we assume

$$m \in L^{\infty}(\Omega), f_{\Omega} m > 0,$$

then [3, 5, 6] guarantee the existence, uniqueness and stability of a solution to (1).

r

We consider the functional

$$F:(m,\mu)\mapsto \int_{\Omega}\theta_{m,\mu}$$

and the optimisation problem

$$\sup_{n, f_{\Omega} m > 0} F(m, \mu).$$

This problem is ill-posed without further constraints on m. Two natural constraints can be set, a pointwise (L^{∞}) constraint, and a L^1 constraint, which leads to introducing the admissible class:

$$\mathcal{M}(\Omega) := \left\{ m \in L^{\infty}(\Omega) , 0 \leqslant m \leqslant \kappa , \int m = m_0 \right\}.$$
(2) {Eq:Ad}

This admissible class was proposed in [26] and used, for instance, in [30, 31].

The optimisation problem under consideration is

$$\max_{m \in \mathcal{M}(\Omega)} \oint_{\Omega} \theta_{m,\mu}.$$
 (P_µ) {PV}

The direct method of the calculus of variations yields in a straightforward way the existence of a solution $m^*_{\mu} \in \mathcal{M}(\Omega)$.

{Se:Bibl}

A remark on the constraints We would like to stress the importance of the pointwise constraint $0 \leq m \leq \kappa$. As mentioned in the first part of this Introduction, a natural question was that of knowing whether the total population size could exceed the total amount of resources, see [9, 37]. In other words, what can be said about the ratio

$$E(m) := \sup_{\mu > 0} \frac{f_{\Omega} \theta_{m,\mu}}{f_{\Omega} m},$$

where m satisfies $m \ge 0$, $m \ne 0$? It follows from [25] that

$$E(m) \ge 1$$

In the one-dimensional case, Bai, He and Li proved, in [2] that

 $E(m) \leqslant 3$

and that this bound is not attained for any m.

In the *n*-dimensional case, $n \ge 2$, Inoue and Kuto proved, in the preprint [17], that

$$\sup_{m \ge 0, m \ne 0} E(m) = +\infty.$$

Qualitative properties for (P_{μ}) One of the main features of problems such as (P_{μ}) is the bang-bang property: denoting by m_{μ}^* a maximiser for (P_{μ}) , is it true that there exists a set E_{μ}^* such that $m_{\mu}^* = \kappa \mathbb{1}_{E_{\mu}^*}$? Such characteristic functions are called bang-bang functions. This property is of paramount importance in optimisation and, from a mathematical biology point of view, corroborates the relevance of patch-models, see [7].

Let us briefly sum up the main conclusions of [30, 31], which contain the most up to date qualitative informations of that sort about (P_{μ}) :

- 1. A bang-bang property is proved in [31]: if the set $\{0 < m < \kappa\}$ contains an open ball, then m is not a solution of (P_{μ}) . Here, a regularity assumption is thus needed.
- 2. In [30], it is proved that the bang-bang property holds for all large enough diffusivities.
- 3. It is furthermore proved, also in [30] that:
 - (a) In the one-dimensional case $\Omega = (0; 1)$, there exists $\mu_1 > 0$ such that, for every $\mu \ge \mu_1$, the unique solutions of (P_{μ}) are

$$m^* := \kappa \mathbb{1}_{(0;\ell)}$$
 and $m_* := \kappa \mathbb{1}_{(1-\ell;\ell)}$,

with $\kappa \ell = m_0$. We note that these are also optimal configurations for survival ability (see [3, 22]).

- (b) In the 2-dimensional case $\Omega = (0; 1) \times (0; 1)$, concentration holds for large diffusivities in the following sense: any sequence $\{m_{\mu}^*\}_{\mu\to\infty}$ of solutions of (P_{μ}) converges, up to a subsequence, to a bang-bang function $m_{\infty}^* = \kappa \mathbb{1}_{E_{\infty}^*}$ which is decreasing in every direction: $x \mapsto m(x, y)$ (resp. $y \mapsto m(x, y)$) is non-increasing for almost every y (resp. non-increasing for almost every x)
- 4. Finally, in [30] it is proved that, for small enough diffusivities, fragmentation may be better in the following sense: two crenels are better than one crenel.

As already mentioned, our goal is to prove a strong fragmentation phenomenon for small diffusivities. A way to formalise this fragmentation would be to restrict ourselves to looking for bang-bang solutions of (P_{μ}) , i.e solutions of the form $m_{\mu}^* = \kappa \mathbb{1}_{E_{\mu}^*}$ and to prove that

$$Per(E^*_{\mu}) \xrightarrow[\mu \to 0]{} +\infty$$

for some notion of perimeter.

However, the problem

$$\sup_{m \in \mathcal{M}(\Omega), m \text{ bang-bang}} F(m, \mu)$$

does not necessarily have a solution, as remarked above.

Mathematical formulation of fragmentation To quantify the perimeter or the regularity of the optimal resources distribution, we introduce, for a fixed M > 0, the class

$$\mathcal{M}_{M}(\Omega) := \left\{ m \in \mathcal{M}(\Omega), \|m\|_{BV(\Omega)} \leqslant M \right\}.$$
(3) {Eq: AdM}

Here, the $BV(\Omega)$ -norm refers to the bounded-variations norm. We note that, for instance, a set E has a finite perimeter (in the sense of Caccioppoli) if and only if $\mathbb{1}_E$ is a function of bounded variations, so that it gives us a natural extension of the notion of perimeter to the set of admissible resources distributions.

For a general introduction to functions of bounded variations and their link to perimeter, we refer to [1].

1.3 Main result

The main result of this article is the following fragmentation property. {Th:Frag}

Theorem 1. Let, for any $\mu > 0$, m_{μ}^* be a solution of (P_{μ}) . There holds

$$\|m_{\mu}^{*}\|_{BV(\Omega)} \xrightarrow[\mu \to 0^{+}]{} +\infty.$$
 (4) {fragmentation}

 $\{TH\}$

More precisely, we prove:

Remark 1. Since $||m_{\mu}^*||_{L^1(\Omega)} = m_0$, the above statement actually says that the $TV(\Omega)$ -seminorm of m_{μ}^* blows up as $\mu \to 0$.

Remark 2. Two remarks should be made:

• We could actually prove, using our method, that

$$\overline{\lim_{\mu \to 0^+}} \left(\sup_{m \in \mathcal{M}(\Omega)} F(m, \mu) \right) = ||F||_{L^{\infty}(\mathcal{M}(\Omega) \times \mathbb{R}_+)}$$

as will be explained later, see Remark 3. Proving this actually gives a (weaker) fragmentation result (i.e one could find a subsequence of maximisers such that the corresponding sequences of $BV(\Omega)$ -norms diverges to $+\infty$). This seems to indicate that finding the limit problem is very challenging. Finally, we were only able to prove that

$$\lim_{\mu \to 0^+} \left(\sup_{m \in \mathcal{M}(\Omega)} F(m,\mu) \right) > m_0 = \inf_{m \in \mathcal{M}(\Omega), \mu \in \mathbb{R}_+} F(m,\mu).$$

• Theorem 1 can be recast in terms of perimeters. In this case, one may consider the set of admissible subsets

$$\mathcal{O}(\Omega) := \left\{ E \subset \Omega, |E| = \frac{m_0}{\kappa} \right\}$$

and the auxiliary subsets

$$\mathcal{O}_M(E) := \{ E \in \mathcal{O}(\Omega), Per(E) \leq M \}.$$

Here, the perimeter is to be understood in the sense of Caccioppoli. Note that, as already pointed out, the existence of a solution to

$$\sup_{E\in\mathcal{O}(\Omega)}F(\mathbb{1}_E,\mu)$$

is not known for general μ , see [30]. However, we can prove, in the same fashion that, for every M > 0, there exists $\mu_M > 0$ such that, for any $\mu \leq \mu_M$, there holds

$$\sup_{E \in \mathcal{O}_M(\Omega)} F(\mathbb{1}_E, \mu) < \sup_{E \in \mathcal{O}(\Omega)} F(\mathbb{1}_E, \mu).$$

2 Proof of Theorem 1

2.1 The influence of periodisation

The main idea is to exploit the non-monotonicity of the function

$$\mu \mapsto F(m,\mu),$$

for a fixed m.

We recall (see [25]) that

- 1. Setting $F(m, 0) = F(m, +\infty) = m_0$ extends F to a continuous function on $[0; +\infty]$.
- 2. m_0 is a strict, global minimiser of $F(m, \cdot)$ if and only if $m \neq m_0$. If $m \equiv m_0$, then $F(m, \cdot) \equiv m_0$.
- 3. $F(m, \cdot)$ may have several local maxima (see [24], where construct m such that $F(m, \cdot)$ has at least two local maxima).

Our method of proof consists in exploiting this non-monotonicity, as well as Neumann boundary conditions and the fact that we are working in an orthotope.

Indeed, let $k \in \mathbb{N}$. We can extend m and $\theta_{m,\mu}$ to $[-1;1]^n$ by reflecting them across each of the axis segments $\{x_i = 0, 0 \leq x_i \leq 1\}, i = 1, ..., n$ and, then, we can extend them to 2-periodic (in each direction) functions on \mathbb{R}^n . It then makes sense to define, for a given $m \in \mathcal{M}(\Omega)$, the functions

$$m_k(x) := m\left(2^k x\right), \theta_k(x) := \theta_{m,\mu}(2^k x).$$

Straightforward computations show that (m_k, θ_k) solves

$$\begin{cases} \frac{\mu}{2^{2k}} \Delta \theta_k + \theta_k (m_k - \theta_k) = 0 \text{ in } \Omega, \\ \frac{\partial \theta_k}{\partial \nu} = 0 \text{ on } \partial \Omega. \end{cases}$$
(6)

{Proof}

Furthermore,

$$\begin{split} & \oint_{[0;1]^n} \theta_k(x) dx = \frac{1}{(2^k)^n} \int_{[0;2^k]^n} \theta_{m,\mu}(y) dy \\ & = \oint_{[0;1]^n} \theta_{m,\mu}. \end{split}$$

As a consequence of these identities, we have

$$F\left(m_k, \frac{\mu}{2^{2k}}\right) = F(m, \mu). \tag{7} \quad \{\texttt{Eq:Period}\}$$

Visually, if we represent, for instance, $F(m, \cdot)$ as



then $f_k := F(m_k, \cdot)$ can be visualised as



Using (7), we are going to that there exists $\eta > 0$ and $\mu_{\eta} > 0$ such that

$$\inf_{0 \le \mu \le \mu_{\eta}} \left(\sup_{m \in \mathcal{M}(\Omega)} F(m, \mu) \right) \ge m_0 + \eta.$$

Then, we will show that, for any M > 0, for any $\varepsilon > 0$, there exists $\mu_{M,\varepsilon} > 0$ such that

$$\inf_{0 \leqslant \mu \leqslant \mu_{M,\varepsilon}} \sup_{m \in \mathcal{M}_M(\Omega)} F(m,\mu) \leqslant m_0 + \varepsilon.$$

The conclusion of Theorem 1 follows immediately from these two steps.

2.2 Technical preliminaries

Technical background We briefly recall some well-known facts about Equation 1. From the method of sub- and super-solutions (we refer for instance to [3]) we have

$$\forall \mu \in (0; +\infty), \forall m \in \mathcal{M}(\Omega), 0 \leq \theta_{m,\mu} \leq \kappa.$$

Lou, in [25], proves the three following results: first,

$$\forall \mu \in (0; +\infty), \mu \oint_{\Omega} \frac{|\nabla \theta_{m,\mu}|^2}{\theta_{m,\mu}^2} = \oint_{\Omega} \theta_{m,\mu} - m_0.$$
(8)

Then,

$$\forall m \in \mathcal{M}(\Omega), \forall p \in [1; +\infty), \|\theta_{m,\mu} - m\|_{L^p(\Omega)} \xrightarrow[\mu \to 0]{} 0.$$
(9) {Eq:CVM}

Finally, he obtains the following estimate in [25, Claim, Equation 2.4]: there exists a constant C independent of m and $\mu > 0$ such that

$$\forall (m,m') \in \mathcal{M}(\Omega)^2, \forall \mu > 0, \quad \|\theta_{m,\mu} - \theta_{m',\mu}\|_{L^1(\Omega)} \leqslant C \|m - m'\|_{L^1(\Omega)}^{\frac{1}{3}}.$$
 (10) {Eq:Lou}

Although Lou, in [25], assumes that m' is regular, it is readily checked that his proof does not depend on the smoothness of m' and can be extended to all elements of $\mathcal{M}(\Omega)$ in a straightforward way.

Uniform convergence in $\mathcal{M}_M(\Omega)$ (as $\mu \to 0^+$) Our goal is to make the convergence result (9) uniform in $m \in \mathcal{M}_M(\Omega)$. This is the content of the following Lemma:

Lemma 1. For any M > 0, the convergence result (9) is uniform in $\mathcal{M}_M(\Omega)$ in the following sense: let M > 0 be fixed. Then

$$\forall \varepsilon_0 > 0, \exists \mu_{M,\varepsilon_0}, \forall m \in \mathcal{M}(\Omega), \forall 0 \leqslant \mu \leqslant \mu_{M,\varepsilon_0}, \|\theta_{m,\mu} - m\|_{L^1(\Omega)} \leqslant \varepsilon_0.$$
(11) {Eq:Le1}

Proof of Lemma 1. We argue by contradiction. If we assume that (11) does not hold then there exists $\varepsilon_0 > 0$ and a sequence $\{m_k, \mu_k\} \in (\mathcal{M}_M(\Omega) \times (0; +\infty))^{\mathbb{N}}$ such that:

- 1. $\{\mu_k\}_{k \in \mathbb{N}}$ is decreasing and converging to 0,
- 2. There holds:

$$\forall k \in \mathbb{N}, \|\theta_{m_k,\mu_k} - m_k\|_{L^1(\Omega)} \ge \varepsilon_0 > 0.$$
(12)

The embedding $BV(\Omega) \hookrightarrow L^1(\Omega)$ is compact. Hence, there exists $m_\infty \in \mathcal{M}_M(\Omega)$ such that

$$m_k \xrightarrow[k \to \infty]{} m_\infty \text{ strongly in } L^1(\Omega).$$
 (13) {Eq:Int}

Thus, we can write, for any $k \in \mathbb{N}$,

$$\begin{aligned} \|\theta_{m_{\infty},\mu_{k}} - m_{\infty}\|_{L^{1}(\Omega)} &\ge \|\theta_{m_{k},\mu_{k}} - m_{k}\|_{L^{1}(\Omega)} - \|m_{k} - m_{\infty}\|_{L^{1}(\Omega)} - \|\theta_{m_{k},\mu_{k}} - \theta_{m_{\infty},\mu_{k}}\|_{L^{1}(\Omega)} \\ &\ge \varepsilon_{0} + \mathop{o}_{k\to\infty}(1) + \mathop{o}_{k\to\infty}(1) \text{ by, successively, (11), (13) and (10)-(13).} \end{aligned}$$

This is in contradiction with (9). Lemma 1 is proved.

Estimating $\lim_{\mu\to 0} \left(\sup_{\mathcal{M}(\Omega)} F(\cdot,\mu) \right)$ The goal of this paragraph is the following Lemma:

Lemma 2. There exists $\eta > 0$ and $\mu_{\eta} > 0$ such that

$$\inf_{0 \le \mu \le \mu_{\eta}} \left(\sup_{m \in \mathcal{M}(\Omega)} F(m, \mu) \right) \ge m_0 + \eta.$$
(14) {Eq:L2}

Proof of Lemma 2. Let $m \in \mathcal{M}(\Omega)$ be any non-constant admissible resources distribution. We know that m_0 is a strict local minimum of $F(m, \cdot)$ on $[0; +\infty)$, and that $\mu \mapsto F(m, \mu)$ is continuous on $[0; +\infty)$. Let $\mu > 0$ be a real number and consider the interval

$$I_0 := |\mu, 4\mu|$$
.

{Le:Unif}

{Le:2}

Since m_0 is only reached at $\mu = 0$ and $\mu = \infty$, it follows that

$$\inf_{\mu \in I_0} F(m,\mu) > m_0$$

Thus, let $\eta > 0$ be such that

$$\inf_{\mu\in I_0} F(m,\mu) \ge m_0 + \eta.$$

We then consider, for any $k \in \mathbb{N}$, the interval

$$I_k := \left[\frac{\underline{\mu}}{2^{2k}}, \frac{4\underline{\mu}}{2^{2k}}\right].$$

We first remark that, once again setting $m_k(\cdot) = m(k \cdot)$ and, thanks to (7), we have

$$F(m_k, \cdot)(I_k) = F(m, \cdot)(I_0)$$

so that

$$\forall k \in \mathbb{N}, \inf_{\mu \in I_k} F(m_k, \mu) = \inf_{\mu \in I_0} F(m, \cdot) \ge m_0 + \eta.$$

Hence,

$$\inf_{\mu \in I_k} \left(\sup_{m \in \mathcal{M}(\Omega)} F(m, \mu) \right) \ge m_0 + \eta.$$
(15) {eq3}

$$\sup(I_{k+1}) = \inf(I_k)$$

Hence, setting

$$I_{\infty} := \bigcup_{k=1}^{\infty} I_k$$

we can write

$$I_{\infty} = \left(0, 4\underline{\mu}\right)$$

and, as a consequence of (15),

$$\inf_{\mu \in I_{\infty}} \left(\sup_{m \in \mathcal{M}(\Omega)} F(m, \mu) \right) \ge m_0 + \eta.$$

This concludes the proof.

Remark 3. We can use the same method to prove that

$$\overline{\lim_{\mu \to 0^+}} \left(\sup_{m \in \mathcal{M}(\Omega)} F(m, \mu) \right) = ||F||_{L^{\infty}(\mathcal{M}(\Omega) \times \mathbb{R}_+)}.$$

Indeed, consider the problem

$$\sup_{\mu>0,m\in\mathcal{M}(\Omega)}F(m,\mu)=F(m^*,\mu^*).$$

Uniqueness does not hold for this problem, because of the periodisation process we used above.

One can actually see that we can choose $\mu^* > 0$. In this case, considering the sequence $\left(m_k^*, \frac{\mu^*}{2^{2k}}\right)_{k \in \mathbb{N}}$ immediately gives the result.

□ {Rem}

2.3 The proof

Proof of Theorem 1. Let M > 0 be fixed. We are going to prove that there exists $\mu_M > 0$ such that, for any $\mu \leq \mu_M$,

$$\sup_{m \in \mathcal{M}_M(\Omega)} F(m,\mu) < \sup_{m \in \mathcal{M}(\Omega)} F(m,\mu).$$
(16) {Eq:goal}

Theorem 1 follows immediately from (16).

Let $\eta > 0$ and $\mu_{\eta} > 0$ (given by Lemma 2) be fixed throughout the rest of this demonstration:

$$\forall \mu \leqslant \mu_{\eta}, \sup_{m \in \mathcal{M}(\Omega)} F(m, \mu) \geqslant m_0 + \eta.$$
(17) {Eq:4}

From Lemma 1, there exists $\mu_{M,\frac{\eta}{2}} > 0$ such that, for any $\mu \leq \mu_{M,\frac{\eta}{2}}$, we have

$$\sup_{m \in \mathcal{M}_M(\Omega)} \|\theta_{m,\mu} - m\|_{L^1(\Omega)} \leqslant \frac{\eta}{2}.$$

Thus

$$\forall \mu \leqslant \mu_{M,\frac{\eta}{2}}, \forall m \in \mathcal{M}_M(\Omega), \int_{\Omega} \theta_{m,\mu} \leqslant m_0 + \frac{\eta}{2}.$$

Plugging this in (17) then proves that, for $\mu \leq \min(\mu_{\eta}, \mu_{M, \frac{\eta}{2}})$, no solution m_{μ}^* of (P_{μ}) can belong to $\mathcal{M}_M(\Omega)$. This concludes the proof.

Remark 4. We quickly comment on the following, expected, remark: not only does the $BV(\Omega)$ norm blow up, but also, every $X(\Omega)$ -norm, where $X(\Omega)$ is compactly embedded in $L^1(\Omega)$. Indeed, the only part where BV is used is in the proof of Lemma 1, and it is used to get strong $L^1(\Omega)$ convergence.

3 Numerical simulations

We present several numerical simulations in order to emphasise the results of Theorem 1. All of these simulations were obtained using Ipopt [36]. As is expected, the smaller the diffusivity μ , the higher the number of connected components of the set $\{m_{\mu}^* = 1\}$, m_{μ}^* being the optimal resources distribution.

Let h > 0 be the discretization parameter. We work with a uniform space discretisation of size h. Since, numerically, such optimisation problems can be very complicated, we run our optimisation program with different initial guesses $\boldsymbol{m}_{h}^{(1)}, \ldots, \boldsymbol{m}_{h}^{(k)}, \ldots$ to obtain, for each initial guess, a potential candidate to be the optimiser. We then select, among these candidates, the optimal one by comparing the value of the criteria and, to check our results, we apply a gradient descent as a final step.

The simulations are done in the following way (we only present it in the one-dimensional case):

• Generating random initial guesses $(\boldsymbol{\theta}_h^{(k)}, \boldsymbol{m}_h^{(k)})$. We generate a random sample of initial guessest $\boldsymbol{m}_h^{(k)}$ by randomising their first five Fourier coefficients on each discretisation interval. In other words, we define

$$\boldsymbol{m}_{h}^{(k)} = m_{h,i}^{(k)} \text{ on } I_i := [hi; h(i+1)]$$

{Num}

where each of the $m_{h,i}^{(k)}$ is a a random function generated as follows in the one-dimensional case:

$$m_{h,i}^{(k)} = a_0 + \sum_{j=1}^{5} a_j \sin(j\pi ih) + b_j \cos(j\pi ih)$$
(18)

where a_j and b_j are uniform random variables with values in [-0.5, 0.5]. To ensure that the resulting function $\boldsymbol{m}_h^{(k)}$ satisfies the constraint $\boldsymbol{m}_h^{(k)} \in \mathcal{M}(0; 1)$, we apply an affine transformation

$$\begin{cases} T\left(\boldsymbol{m}_{h}^{(k)}\right) = a\boldsymbol{m}_{h}^{(k)} + b \\ a = \min\left(\left|\frac{\kappa - m_{0}}{\max_{i}(m_{h,i}^{(k)} - \sum_{i} m_{h,i}^{(k)})}\right|, \left|\frac{-m_{0}}{\min_{i}(m_{h,i}^{(k)} - \sum_{i} m_{h,i}^{(k)})}\right|\right) \\ b = m_{0} - a\sum_{i} m_{h,i}^{(k)}. \end{cases}$$

The resulting function satisfies $\boldsymbol{m}_{h}^{(k)} \in \mathcal{M}(\Omega)$.

Now, to each of these random initial guess we need to associate an initial guess for the solution of the partial differential equation. We choose an energetic approach: we minimise with Ipopt the discretised energy functional associated with Equation (1) to obtain $\boldsymbol{\theta}_{h}^{(k)}$, which is a piecewise constant function: $\boldsymbol{\theta}_{h}^{(k)} = \boldsymbol{\theta}_{h,i}^{(k)}$ on [ih; (i+1)h]; in other words, $\boldsymbol{\theta}_{h}^{(k)}$ is the minimiser of

$$J^{h}(\boldsymbol{\theta}_{h}) = \frac{\mu}{2} \boldsymbol{\theta}_{h} \left(-\Delta_{h} \right) \boldsymbol{\theta}_{h} - \sum_{i=1}^{N} \left(\frac{1}{2} \theta_{h,i}^{2} m_{h,i}^{(k)} - \frac{1}{3} \theta_{h,i}^{3} \right).$$
(19) {numfun}

where $riangle_h$ is the discrete Laplacian with Neumann boundary conditions, in 1D:

$$\Delta_h := \frac{1}{h^2} \begin{pmatrix} -2 & 2 & 0 & 0 & 0 & \cdots & 0 \\ 1 & -2 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & -2 & 1 & 0 & \cdots & 0 \\ & & & \ddots & \ddots & & \\ & & & \ddots & \ddots & \ddots & \\ 0 & & & \cdots & 0 & 1 & -2 & 1 \\ 0 & & & \cdots & 0 & 2 & -2 \end{pmatrix}$$
(20)

In then end, we get an initial random guess for an optimiser, which we denote $(\boldsymbol{\theta}_h, \boldsymbol{m}_h)$.

• Optimisation under a finite difference scheme constraint We use Ipopt to maximise the total population $\sum \theta_{h,i}^{(k)}$ for every k with respect to \boldsymbol{m}_h . We implement the partial differential equation (1) as a constraint in the scheme:

$$\mu(-\triangle_h \boldsymbol{\theta}_h)_i = \theta_{h,i} m_{h,i} - \theta_{h,i}^2,$$

and, obviously, the constraint $m \in \mathcal{M}(0; 1)$. Among all random initialisations, we choose the optimal solution.

• Gradient descent We recall that, in this context, the adjoint state for the variational problem (P_{μ}) is the function $p\theta_{m,\mu}$ where p solves

$$-\mu\Delta p - p(m - 2\theta_{m,\mu}) = 1$$

with Neumann boundary conditions; in other words, for an admissible perturbation ξ at an admissible resources distribution m (i.e., for every t small enough, $m + t\xi \in \mathcal{M}(0; 1)$) the derivative of the criterion at m in the direction ξ is

$$\int_{\Omega} p\theta_{m,\mu} \xi$$

We refer to [30].

first compute, with the same space discretisation, the discretised adjoint state p_h :

$$-\mu \Delta_h \boldsymbol{p}_h - diag(\boldsymbol{p}_h)(\boldsymbol{m}_h - 2\boldsymbol{\theta}_h) = \boldsymbol{1}$$
(21)

and we find the admissible perturbation $\boldsymbol{\xi}_h$ that gives the highest rise of the total population via maximizing with Ipopt the following quantity:

$$\max_{0 \leqslant m_{h,i} + \xi_{h,i} \leqslant \kappa, \sum_{i} \xi_{h,i} = 0} \sum_{i} p_{h,i} \theta_{h,i} \xi_{h,i}, \tag{22}$$

which corresponds to the highest directional derivative with respect to m, and we apply the gradient descent with Armijo rule. We use a classical stopping criterion, and display the results.

3.1 Simulations in the one-dimensional case

For one-dimensional simulations, we work in

$$\Omega = [0; 1],$$

with $\kappa = 1$ and $N_x = 1000$ discretization points. For each value of the parameter μ , we represent, on the same picture the optimal resources distribution m^*_{μ} (the blue zones correspond to m = 1), which we observe, in each of our case, to be a bang-bang function, and the corresponding solution $\theta_{m^*_{\mu},\mu}$ of (1).

In order to emphasise the influence of the parameter m_0 on the qualitative properties of optimal resources distributions, we present two different values of m_0 .

3.1.1 $\kappa = 1, m_0 = 0.3$







Figure 2



Figure 3

3.1.2 $\kappa = 1, m_0 = 0.6$



Figure 4



Figure 5



Figure 6

3.2 Simulations in the two-dimensional case

For two-dimensional simulations, we work in

$$\Omega = [0; 1]^2,$$

with $\kappa = 1$. For each value of the parameter μ , we represent, on the left picture, the optimal resources distribution m_{μ}^* , which we observe, in each of our case, to be a bang-bang function. On the right, we represent the corresponding solution $\theta_{m_{\mu}^*,\mu}$ of (1).

In order to emphasise the influence of the parameter m_0 on the qualitative properties of optimal resources distributions, we present, as in the one-dimensional case, two different cases. We once again highlight the fact that these simulations prohibit, at a theoretical level, the use of rearrangements to derive qualitative properties but we do notice, in this two dimensional case, the presence of many symmetries. It is a very challenging and interesting project to obtain symmetry properties for this kind of problems. The number of discretisation points in the x and y variable are $N_x = N_y = 60$; the method is otherwise similar to that in the one-dimensional case.









Figure 8

3.2.1 $\kappa = 1, m_0 = 0.3$



Figure 9





Figure 10





Figure 11

3.2.2 $\kappa = 1, m_0 = 0.6$





Figure 12



Figure 13



Figure 14





0.8

0.6

0.4

0.2

0

1

Figure 15





References

[1] L. Ambrosio, N. Fusco, and D. Pallara. Functions of Bounded Variation and Free Discontinuity Problems (Oxford Mathematical Monographs). Oxford University Press, 2000.

- [2] X. Bai, X. He, and F. Li. An optimization problem and its application in population dynamics. Proceedings of the American Mathematical Society, 144(5):2161–2170, Oct. 2015.
- [3] H. Berestycki, F. Hamel, and L. Roques. Analysis of the periodically fragmented environment model: I – species persistence. *Journal of Mathematical Biology*, 51(1):75–113, 2005.
- [4] H. Berestycki and T. Lachand-Robert. Some properties of monotone rearrangement with applications to elliptic equations in cylinders. *Mathematische Nachrichten*, 266(1):3–19, mar 2004.
- [5] R. S. Cantrell and C. Cosner. Diffusive logistic equations with indefinite weights: Population models in disrupted environments II. SIAM Journal on Mathematical Analysis, 22(4):1043– 1064, jul 1991.
- [6] R. S. Cantrell and C. Cosner. The effects of spatial heterogeneity in population dynamics. J. Math. Biol., 29(4):315–338, 1991.
- [7] R. S. Cantrell and C. Cosner. Spatial Ecology via Reaction-Diffusion Equations. John Wiley & Sons, 2003.
- [8] R. S. Cantrell, C. Cosner, and V. Hutson. Permanence in ecological systems with spatial heterogeneity. Proceedings of the Royal Society of Edinburgh Section A: Mathematics, 123(3):533– 559, 1993.
- [9] D. DeAngelis, B. Zhang, W.-M. Ni, and Y. Wang. Carrying capacity of a population diffusing in a heterogeneous environment. *Mathematics*, 8(1):49, Jan. 2020.
- [10] R. A. Fisher. The wave of advances of advantageous genes. Annals of Eugenics, 7(4):355–369, 1937.
- [11] J. D. Goss-Custard, R. A. Stillman, R. W. G. Caldow, A. D. West, and M. Guillemain. Carrying capacity in overwintering birds: when are spatial models needed? *Journal of Applied Ecology*, 40(1):176–187, Feb. 2003.
- [12] W. He and W.-M. Ni. Global dynamics of the Lotka–Volterra competition–diffusion system with equal amount of total resources, III. *Calc. Var. Partial Differential Equations*, 56(5):56:132, 2017.
- [13] X. He and W.-M. Ni. Global dynamics of the lotka-volterra competition-diffusion system: Diffusion and spatial heterogeneity is. *Communications on Pure and Applied Mathematics*, 69(5):981–1014, 2015.
- [14] X. He and W.-M. Ni. Global dynamics of the lotka–volterra competition–diffusion system with equal amount of total resources, ii. *Calculus of Variations and Partial Differential Equations*, 55, 04 2016.
- [15] A. Henrot. Extremum problems for eigenvalues of elliptic operators. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2006.
- [16] A. Henrot and M. Pierre. Variation et optimisation de formes: une analyse géométrique, volume 48. Springer Science & Business Media, 2006.
- [17] J. Inoue and K. Kuto. On the unboundedness of the ratio of species and resources for the diffusive logistic equation, 2020.

- [18] C.-Y. Kao, Y. Lou, and E. Yanagida. Principal eigenvalue for an elliptic problem with indefinite weight on cylindrical domains. *Math. Biosci. Eng.*, 5(2):315–335, 2008.
- [19] B. Kawohl. Rearrangements and Convexity of Level Sets in PDE. Springer Berlin Heidelberg, 1985.
- [20] A. Kolmogorov, I. Pretrovski, and N. Piskounov. étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique. *Moscow University Bulletin of Mathematics*, 1:1–25, 1937.
- [21] K.-Y. Lam and Y. Lou. Persistence, competition, and evolution. In *The Dynamics of Biological Systems*, pages 205–238. Springer International Publishing, 2019.
- [22] J. Lamboley, A. Laurain, G. Nadin, and Y. Privat. Properties of optimizers of the principal eigenvalue with indefinite weight and Robin conditions. *Calculus of Variations and Partial Differential Equations*, 55(6), Dec. 2016.
- [23] J. Langford. Neumann comparison theorems in elliptic pdes. Potential Analysis, 2015.
- [24] S. Liang and Y. Lou. On the dependence of population size upon random dispersal rate. Discrete and Continuous Dynamical Systems - Series B, 17(8):2771–2788, July 2012.
- [25] Y. Lou. On the effects of migration and spatial heterogeneity on single and multiple species. Journal of Differential Equations, 223(2):400–426, Apr. 2006.
- [26] Y. Lou. Some challenging mathematical problems in evolution of dispersal and population dynamics. In *Lecture Notes in Mathematics*, pages 171–205. Springer Berlin Heidelberg, 2008.
- [27] Y. Lou and E. Yanagida. Minimization of the principal eigenvalue for an elliptic boundary value problem with indefinite weight, and applications to population dynamics. Japan J. Indust. Appl. Math., 23(3):275–292, 10 2006.
- [28] I. Mazari. Trait selection and rare mutations: The case of large diffusivities. Discrete & Continuous Dynamical Systems - B, 2019.
- [29] I. Mazari, G. Nadin, and Y. Privat. Optimization of a two-phase, weighted eigenvalue with dirichlet boundary conditions. Preprint, 2019.
- [30] I. Mazari, G. Nadin, and Y. Privat. Optimal location of resources maximizing the total population size in logistic models. *Journal de Mathématiques Pures et Appliquées*, 2020.
- [31] K. Nagahara and E. Yanagida. Maximization of the total population in a reaction-diffusion model with logistic growth. *Calculus of Variations and Partial Differential Equations*, 57(3):80, Apr 2018.
- [32] N. Shigesada and K. Kawasaki. Biological Invasions: Theory and Practice. Oxford University Press, 1997.
- [33] J. G. Skellam. Random dispersal in theoretical populations. *Biometrika*, 38(1-2):196–218, 06 1951.
- [34] K. Taira. Diffusive logistic equations in population dynamics. Adv. Differential Equations, 7(2):237-256, 2002.
- [35] K. Taira. Logistic dirichlet problems with discontinuous coefficients. Journal de Mathématiques Pures et Appliquées, 82(9):1137–1190, Sept. 2003.

- [36] A. Wächter and L. Biegler. On the implementation of an interior-point filter line-search algorithm for large-scale nonlinear programming. *Mathematical Programming*, 106(1):25–57, Mar 2006.
- [37] B. Zhang, A. Kula, K. M. L. Mack, L. Zhai, A. L. Ryce, W.-M. Ni, D. L. DeAngelis, and J. D. V. Dyken. Carrying capacity in a heterogeneous environment with habitat connectivity. *Ecology Letters*, 20(9):1118–1128, July 2017.