



HAL
open science

Current evidence for a role of epigenetic mechanisms in response to ionizing radiation in an ecotoxicological context

N. Horemans, D.J. Spurgeon, C. Lecomte-Pradines, E. Saenen, C. Bradshaw, D. Oughton, I. Rasnaca, J.H. Kamstra, C. Adam-Guillermin

► To cite this version:

N. Horemans, D.J. Spurgeon, C. Lecomte-Pradines, E. Saenen, C. Bradshaw, et al.. Current evidence for a role of epigenetic mechanisms in response to ionizing radiation in an ecotoxicological context. *Environmental Pollution*, 2019, 251, pp.469-483. 10.1016/j.envpol.2019.04.125 . hal-02524800

HAL Id: hal-02524800

<https://hal.science/hal-02524800v1>

Submitted on 7 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Manuscript Details

Manuscript number	ENVPOL_2019_714_R1
Title	Current evidence for a role of epigenetic mechanisms in response to ionizing radiation in an ecotoxicological context
Article type	Review Article

Abstract

The issue of potential long-term or hereditary effects for both humans and wildlife exposed to low doses (or dose rates) of ionising radiation is a major concern. Chronic exposure to ionising radiation, defined as an exposure over a large fraction of the organism's lifespan or even over several generations, can possibly have consequences in the progeny. Recent work has begun to show that epigenetics plays an important role in adaptation of organisms challenged to environmental stimulæ. Changes to so-called epigenetic marks such as histone modifications, DNA methylation and non-coding RNAs result in altered transcriptomes and proteomes, without directly changing the DNA sequence. Moreover, some of these environmentally-induced epigenetic changes tend to persist over generations, and thus, epigenetic modifications are regarded as the conduits for environmental influence on the genome. Here, we review the current knowledge of possible involvement of epigenetics in the cascade of responses resulting from environmental exposure to ionising radiation. In addition, from a comparison of lab and field obtained data, we investigate evidence on radiation-induced changes in the epigenome and in particular the total or locus specific levels of DNA methylation. The challenges for future research and possible use of changes as an early warning (biomarker) of radiosensitivity and individual exposure is discussed. Such a biomarker could be used to detect and better understand the mechanisms of toxic action and inter/intra-species susceptibility to radiation within an environmental risk assessment and management context.

Keywords	Epigenetic marks; Radioecology; gamma radiation; Chronic exposure; multi-transgenerational; Nuclear accidents
Corresponding Author	Nele Horemans
Order of Authors	Nele Horemans, David Spurgeon, Catherine Lecomte-Pradines, Eline Saenen, Clare Bradshaw, Deborah Oughton, Ilze Rasnaca, Jorke Kamstra, christelle Adam-Guillermin
Suggested reviewers	Etiene Bucher, David Copplestone, Almudena Real, Carmel Mothersill, Peter Kille

Submission Files Included in this PDF

File Name [File Type]

letter to the editor.docx [Cover Letter]

rebutal.docx [Response to Reviewers]

2019 bullet points.docx [Highlights]

2019 graphical abstract.pptx [Graphical Abstract]

2019 horemans et al revised.docx [Manuscript File]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Ref: ENVPOL_2019_714

Title: Current evidence for a role of epigenetic mechanisms in response to ionizing radiation in an ecotoxicological context

Journal: Environmental Pollution

Dear editor,

Thank you and your reviewers for giving us the opportunity to revise our manuscript entitled *Current evidence for a role of epigenetic mechanisms in response to ionizing radiation in an ecotoxicological context* (ENVPOL_2019_714). We have tried to address all remarks made by the reviewers as well as journal recommendations. Below you can find our rebuttal specifying the changes made to the text and tables. We hope that our manuscript will now meet the standards of Environmental Pollution and gets approval of the reviewers.

Looking forward to hearing from you,

Kind regards

Nele Horemans

Prof. dr. Nele Horemans

Head of research unit Biosphere Impact Studies (BIS)

Guest lecturer at University of Hasselt (CMK)

+32 14 33 21 15

Expert group of Interdisciplinary Biosciences (BIO)

Institute of Environmental Health and Safety

SCK•CEN | Belgian Nuclear Research Centre

Boeretang 200 - BE-2400 Mol

www.sckcen.be

Rebutal (ENVPOL_2019_714)

Comments from the editors and reviewers:

-Reviewer 1

The authors reviewed the current knowledge of possible involvement of epigenetics in the cascade of responses resulting from environmental exposure to ionizing radiation. They also summarized the evidence on whether changes in the epigenome and in particular the total or locus specific levels of DNA methylation may serve as an early warning (biomarker) of radio-sensitivity and individual exposure, from a comparison of lab and field obtained data. This review paper could help understand the mechanisms of toxic action and inter/intra-species susceptibility to radiation within an environmental risk assessment and management context.

This review paper is certainly interesting and presents valuable information. However it may be suitable for publication in a more subject focused journal, such as Radiation Physics and Chemistry, Radiation Research, Radiation Protection Dosimetry, and the like.

We would like to thank the reviewer for his comments and suggestion to submit our current paper to another journal which is more focused on radiation protection. However, we deliberately chose for Environmental Pollution as this journal, in contrast to the once suggested by the reviewer, is focused on the consequences of pollutants in our natural environment. This review paper specifically deals with radiation as an environmental polluter and the possible long-term consequences to the environment of enhanced exposures. We therefore truly believe it falls within the scope of the journal.

-Reviewer 2

The authors in this review, they bring together and present in a relatively nice way the current knowledge of possible involvement of epigenetics in the responses resulting from environmental exposure to ionising radiation (IR). The need for reliable radiation exposure biomarkers in order to evaluate or even predict the stochastic effects of IR in this case for example mutations, genomic instability 'transmitted' to next generations is certainly important. Now if the epigenetic markers for wildlife or even humans are the best that remains to be seen. The authors need to revise and correct several issues as described below.

The major issues are that they need to embrace the knowledge of radiation bystander effects, the role of IR-stress manifested to the complexity of DNA damage (and probably lipids, proteins etc.). These issues starting from the early work by Munira Kadhim (2004: Interrelationships amongst radiation-induced genomic instability, bystander effects, and the adaptive response. Mutation research, 568, 21-32) to more recent analytical reviews explaining all these (Pouget, J.P., et al. 2018) Targeted and Off-Target (Bystander and Abscopal) Effects of Radiation Therapy: Redox Mechanisms and Risk-Benefit Analysis. Antioxid Redox Signal). Because the DNA damage response is the main regulator of all further effects so the complexity of damage is probably the most prominent parameter for this reviewer. Of course

cases where direct damage to methylating protein genes maybe involved but how possible is this directly at least from radiation only to these genes?

We have added some lines on the possible links between non-targeted effects and epigenetics. We share the view of the recent review by Schofield, that the understanding of the mechanisms linking these two processes is still poorly understood.

"A number of papers have proposed a link between epigenetic effects and non-targeted effects (NTE) such genomic instability and bystander effects (Schofield and Kondratowicz, 2018). However, while the existence of non-targeted effects is well established (Morgan, 2002, Kadhim et al 2004, Pouget et al 2018), and studies have shown an association between the two effects (e.g., Kaup et al., 2006; Wang et al 2018), evidence of a causal relationship is more elusive, since NTE could be either a mechanism or a consequence of epigenetic changes (Schofield and Kondratowicz, 2018)"

Another point and based on several discussion on the radiation ecology communities etc. is for example how one can differentiate between IR-effects and for example airborne particles from wild fires or chemical pollutants? How sure are the authors that their markers are exclusive to IR-exposure? This the reason that probably they need to invest in the unique feature of IR as discussed above and below as a type of environmental stress.

We agree with the author that it in natural environments it will remain questionable whether a specific marker for IR-exposure or any other stressor can be found. In this review we look at evidence for the appearances of changes in methylation or other epigenetic marks after exposure to radiation in either field or lab conditions. It is the first step in the quest whether or not these can serve as biomarkers. Hence we only shortly touch upon the possibility of biomarkers in this review and this mainly in paragraph 9. Within this section it was already indicated at two places that it will be challenging if not impossible to exclude the influence of confounding factors:

- line 671 to 673: However, it is also recognised that significant challenges related to the effects of genetic background and the influence of confounding factors also exist.*
- line 680 to 683: Specific challenges relate to working with some autochtonous species for which genome resources may be lacking and, the influence of confounding factors which may mask the causal response between ionising radiation exposure and epigenetic changes.*

We have added the review of Pernot E, et al. (Mutation Research-Reviews in Mutation Research. 2012;751(2):258-286) as the authors describe the challenges on the specificity of biomarkers for IR.

We do believe that it will be necessary to combine different observations that lead to an adverse outcome in order to identify a set of changes that will be specific for IR as is indicated in the text: epigenetic approaches may be more powerful indicators of effects when linked to known biomarkers using, for example, transcriptional analysis. When used in conjunction with other mechanistic measurements, epigenetic analysis

has the potential to enhance the ecological relevance of molecular biomarkers, as described in the Adverse Outcome Pathway concept (Groh et al. 2015) Line 683-687. We also suggest a thorough comparison between field and lab studies line 696-697. We also added in paragraph 9 the paper of Lourenco J, et al. (J Hazard Mater. 2016;317:503-542) as this is a comprehensive review on the use of biomarkers or bioindicator species for humans and non-human species for IR.

Other specific (not minor) comments:

1. In the abstract, bullets, whole manuscript, the authors need to be very specific. For example when they say radiation they must at least explain initially, what type of ionizing radiation they refer to, what type of exposure acute or chronic as assumed.

To address this remark we have added some more information in table 2 and 3 we have included in the text that exact amounts can be found in the tables (line 333). Throughout the text we also tried to be more precise on this matter.

2. Abstract and elsewhere: "... have consequences in the progeny, which may not be attributable to increased mutation rates alone". The question here is really crucial and based on the long discussion of the systemic effects of ionizing radiation. When they refer to progeny, is the progeny exposed to radiation, or for example the parents have left the irradiation area and the offsprings have been born to a background radiation level area?

We agree with the reviewer that it is important to indicate whether or not the offspring itself is exposed. We have defined this (line 697 to 701) as mutligenerational exposure (both parents and offspring are exposed, transgenerational: only parental line is exposed. It is also true that care must be taken that sometimes for example in pregnant female organisms it is the F3 generation that is the first true transgenerational exposed generation, this concern is also included in the text (Line 313-317)

3. The Figure is way too large.

We agree with the reviewer that the figure is too large, but just provided this figure in the intention to rescale it the size desired for the graphical abstract.

4. Low doses and chronic exposures must be defined. Lines 59-61: One question that arises, is how it is possible radiation accidents like Chernobyl and Fukushima to be considered as low doses exposure examples? For time periods and areas around the accidents, doses where high to very high. Examples and clarifications must be given.

We agree that for Chernobyl and Fukushima the term low dose is not correct for the first and second period after the accident. However dose rates of the last 30 years are estimated to be maximally 0.5 mGy/h. Examples of the dose rates are now included in the text. For the current situation the highest contaminated areas like the Red Forest for CEZ has about 0.5 mGy/h (Beresford personal communication) to 0.1 mGy/h ambient dose rates (Horemans N, et al. Journal of Environmental

Radioactivity. 2018;192:405-416). As it is impossible for these field conditions to exclude the memory effect of previous higher exposures even if it is more than 30 years ago and to avoid confusion we have omitted the use of the term low dose in the text. This review deals with long-term consequences of chronic exposures: we have as suggested included a definition for chronic exposure conditions both in the abstract as in the first paragraph of the introduction.

5. Lines 86-87: "...Long-term exposures to low levels of environmental stressors have been linked to lasting responses in organisms within, but also over multiple exposed generations". Again here wild uncertainties, long term, low levels, stressors of what type? Type of stressors for DNA have been discussed analytically and differentiated from endogenous and a discussion must be done since the main difference between environmental IR stress and endogenous oxidative stress in all living organisms is the complexity of damage (DNA mainly: Nikitaki, Z. et al. 2015. Stress-induced DNA damage biomarkers: applications and limitations. *Frontiers in chemistry*, 3, 35.).

For this part of the manuscript we have not followed the suggestion by the reviewer to include a discussion on the type of stressor as this sentence in the introduction refers to different stressors (not including IR) as evidenced by the references: (Mirbahai and Chipman 2014:review on different chemical stressors; Schultz et al. 2016 on Ag ions, nanoparticles and sulphadised nanoparticles; Jimenez-Chillaron et al. 2015 (different xenobiotics); Marczylo et al. 2016 and Hanson and Skinner 2016 ((again reveals on different contaminants but not IR). We included this sentence to show that the fact that long-term exposure to an environmental stressor can have consequences potentially over generations and that this has been noted before for different pollutants. We do have removed low dose and added sub-lethal concentrations in order to avoid confusion with IR that is expressed in dose.

6. The Tables are nice but the type of assumed or measured or estimated radiation type and levels must be incorporated.

We have tried to follow this request and added where appropriate radiation dose rate (with indication whether it is ambient or total dose rate including internal exposure) and metal concentrations in the tables 2 and 3

7. How one can be sure that animals and living species have not migrated from irradiated areas and through some type of systemic effects alter the epigenome to animals in background levels areas?

We agree with the reviewer that for field studies it is impossible to exactly estimate the exposed dose and one can never be certain whether an organism has not migrated. For the field studies in which the co-authors were involved care was taken to include control sites that are further away from the contaminated sites than the normal action radius of the organism itself. Dosimetry was carefully performed taken into account as much as possible the habitat and frequency of an organism to reside in that habitat. This is described in the references cited in this review. As however you can never exclude changes in exposure we recommend to compare lab and field conditions as much as possible.

8. Earlier studies suggest in many cases no real differences between high- and low-radiation areas (see for example : https://inis.iaea.org/collection/NCLCollectionStore/_Public/23/039/23039160.pdf)

We have tried when describing the field studies to include whether or not the response followed the radiation gradient, for some cases a correlation between the measured endpoint and the gradient is found indicating differential responses to be radiation level dependent (see e.g. for plants Kovalchuk et al. 2003,2004) other plant studies did not find a correlation (Horemans et al 2018) or showed a transient response (Volkova et al 2018). Hence, it seemed study-dependent and the underlying mechanisms have not been identified yet. Hence, in the manuscript we did not make conclusions on radiation gradient dependencies but included it as a challenge for further research to find the underlying mechanism explaining this if you want to use epigenetic marks as a biomarker.

From the suggested IAEA report we picked out the following definition and included it in the manuscript (line first paragraph of the introduction):

An acute exposure is one which is delivered in a time period which is short compared with the time over which any obvious biological response develops. A chronic exposure is one which could continue over a large fraction of the natural life of the organism. These descriptors as a function of time lead naturally to qualifiers of total dose: A high exposure is one which leads to an acute response, usually a severe (and obvious) pathological reaction, with the primary one being mortality. A low exposure would have only marginal and late effects on the normal mortality/time relationship of the organism, but it may produce detectable effects in the normal biological processes of the organism without necessarily producing any obvious harm to the individual.

-Reviewer 3

There are few minor suggestions i have:

1. "Chernobyl" is generally used spelling given that this nuclear catastrophe had happen in Soviet Union, where official language was Russian. It is, however, a Ukrainian city and proper spelling for it is "Chornobyl". This is, however, my recommendation and i would leave it up to the authors to take a decision on it.

We agree with the author that Chornobyl is the correct current name for this city. We are willing to change the names accordingly. We are, however, a bit reluctant as we are afraid Chornobyl is not widespread as a search term such reducing the visibility of our manuscript once accepted and therefore left it as Chernobyl for the time being.

2. Page 6, line 106: should be "as" instead of "a" in front of "cladocerans".

Changed in the text

3. Page 7, lines 135-136: nucleosome positioning is currently considered as one of the major and driving epigenetic mechanisms, along with the other three - DNA methylation, histone modifications and non-coding RNAs.

We have modified the sentence on this in the text as follows:

The epigenetic landscape is shaped by three epigenetic marks; DNA methylation, histones and its post translation modifications and small RNA interactions. Together they shape the structure of the DNA called chromatin (C. David Allis and Thomas Jenuwein, Nature, 2016, The molecular hallmarks of epigenetic control).

4. Page 9, lines 182 - 185. It is now generally accepted that DNMT1, besides its role in maintenance of DNA methylation patterns, can be also involved in de novo methylation; similarly, DNMT3A and 3B have been shown to partake in maintenance as well.


We agree with the reviewer that this might be added to the text and therefore altered the section as follows:

“In vertebrates, maintenance methylation by DNMT1 occurs during the S-phase of mitosis, where the newly synthesized DNA strand is methylated using the original strand as template. De novo DNA methylation is undertaken DNMT3 family members, although recent insights have shown redundancy between two DNMT family members Lyko F. Nature Reviews Genetics. 2018;19(2):81-92.

- Review on long term effects of exposure of wildlife to chronic low dose radiation
- Inter-, multi- and transgenerational studies for both lab and field exposures
- Changes found in epigenetic marks induced by chronic exposure to ionising radiation
- DNA methylation possibly transfers the response from one generation to the next



Arabidopsis
Daphnia
Earthworms
Pine tree
Salmon
Shepherd's purse
Treefrog
Zebrafish...



Δ Epigenetic marks:

DNA methylation, histone modifications, miRNA expression

1 **Current evidence for a role of epigenetic mechanisms in response to ionizing radiation in an**
2 **ecotoxicological context**

3

4 Horemans, Nele ^{a,b †*}, Spurgeon, David J ^{c *}, Lecomte-Pradines Catherine ^d, Saenen Eline ^a,
5 Bradshaw Clare^e, Oughton Deborah^f, Rasnaca, I. ^c, Kamstra H. Jorke^g, Adam-Guillermin,
6 Christelle^h.

7

8 ^a Belgian Nuclear Research Centre, Boeretang 200, B-2400 Mol, Belgium

9 ^b Centre for Environmental Research, University of Hasselt, Agoralaan, 3590, Diepenbeek,
10 Belgium

11 ^c Centre for Ecology and Hydrology, MacLean Building, Benson Lane, Wallingford, Oxon,
12 OX10 8BB, UK.

13 ^d Institut de Radioprotection et de Sûreté Nucléaire, PSE-ENV/SRTE/LECO, Cadarache, Saint
14 Paul Lez Durance, France

15 ^e Department of Ecology, Environment and Plant Sciences, Stockholm University, 106 91
16 Stockholm, Sweden

17 ^f Centre for Environmental Radioactivity (CERAD), Norwegian University of Life Sciences, 1430
18 Aas, Norway

19 ^g Faculty of Veterinary Medicine, Institute for Risk Assessment Sciences, Utrecht University,
20 Utrecht, The Netherlands.

21 ^h Institut de Radioprotection et de Sûreté Nucléaire, PSE-SANTE, Cadarache, Saint Paul Lez
22 Durance, France

23

24 * These authors contributed equally to this manuscript.

25 † Author to whom correspondence should be addressed Nele Horemans

26 nele.horemans@sckcen.be.

27 **Declaration of interest:** none

28 **Keywords:** Epigenetic marks; Radioecology; DNA methylation; gamma radiation; Chronic

29 exposure; multi-transgenerational; Wildlife; Chernobyl; Chernobyl; Fukushima; Nuclear

30 accidents

31 **Abstract**

32 The issue of potential long-term or hereditary effects for both humans and wildlife exposed
33 to low doses (or dose rates) of ionising radiation is a major concern. Chronic exposure to
34 ionising radiation, defined as an exposure over a large fraction of the organism's lifespan or
35 even over several generations, can possibly have consequences in the progeny. Recent work
36 has begun to show that epigenetics plays an important role in adaptation of organisms
37 challenged to environmental stimulæ. Changes to so-called epigenetic marks such as histone
38 modifications, DNA methylation and non-coding RNAs result in altered transcriptomes and
39 proteomes, without directly changing the DNA sequence. Moreover, some of these
40 environmentally-induced epigenetic changes tend to persist over generations, and thus,
41 epigenetic modifications are regarded as the conduits for environmental influence on the
42 genome.

43 Here, we review the current knowledge of possible involvement of epigenetics in the cascade
44 of responses resulting from environmental exposure to ionising radiation. In addition, from a
45 comparison of lab and field obtained data, we investigate evidence on radiation-induced
46 changes in the epigenome and in particular the total or locus specific levels of DNA
47 methylation. The challenges for future research and possible use of changes as an early
48 warning (biomarker) of radiosensitivity and individual exposure is discussed. Such a biomarker
49 could be used to detect and better understand the mechanisms of toxic action and inter/intra-
50 species susceptibility to radiation within an environmental risk assessment and management
51 context.

52 **Capsule:**

53 Review: possible changes in epigenetic marks in wildlife exposed to ionising radiation suggests
54 DNA methylation changes as a key to transfer the response from one generation to the next.

55 **1 Introduction**

56 Activities like ore mining and milling, nuclear accidents and production and testing of nuclear
57 weapons have resulted in enhanced concentrations of radionuclide pollutants in the
58 environment. This can lead to long-term or chronic exposures of organisms defined as an
59 exposure over a considerable fraction of the lifespan of the organism (IAEA 1992). The issue
60 of biological effects induced by chronic sub-lethal doses of ionising radiation along with the
61 question on the potential hereditary effects for both humans and wildlife is a topic of
62 considerable debate and concern. This has been reinforced after the Chernobyl and
63 Fukushima accidents, especially with respect to the quantification (and reduction if possible)
64 of the magnitude of risk to ecosystems when exposed chronically for multiple generations.
65 This concerns both short-term and chronic exposure over several generations and heritable
66 effects on unexposed progeny. To improve the scientific basis for risk assessment for both
67 human and environment in chronic exposure scenarios as observed e.g. in Chernobyl and
68 Fukushima exclusion zones (CEZ and FEZ), an enhanced understanding of the mechanisms that
69 underpin these responses is needed. This will lead to a better understanding of the complex
70 interplay between exposure, organism physiology and phenotypic response over extended
71 timescales (e.g., Marczylo et al. 2016). Comprehensive reviews of the observed phenotypic
72 effects observed in wildlife in CEZ and FEZ have been published e.g. by Hinton et al. (2007),
73 Geras'kin et al. (2008), Lourenco et al. (2016) Steinhauser et al. (2014), Strand et al. (2014),
74 Batlle (2016) and Beresford et al. (2016). The amounts of radionuclides released into the
75 environment after the Chernobyl accident (5300 PBq, excluding noble gases) were about
76 tenfold of those of the accident in Japan (520 PBq) (Steinhauser et al. 2014). Despite this
77 difference both exclusion zones have common features such as (i) for both areas the exposure
78 can be divided in 3 time-periods depending on the exposure rates as described in paragraph

79 6, (ii) the degree to which spatial and temporal heterogeneity is present in the distribution of
80 the radionuclides (including the presence of hot particles); (iii) the presence of other
81 additional pollutants (e.g. from historical land use); (iv) the challenge of finding comparable
82 control conditions and (v) the difficulty to estimate the exact exposure dose rates. Additionally
83 and of importance for interpreting observations made in these contaminated regions, both
84 exclusion zones have undergone changes induced by the removal of human presence and
85 occupancy leading to specific ecological changes that are hard to distinguish from the possible
86 radiological impact (Beresford and Copplestone 2011). The unique nature of these study areas
87 means that the interpretation of field data from these sites needs careful contextual
88 consideration and have led to contrasting and sometimes conflicting reports on effects
89 observed in the CEZ and FEZ (Beresford and Copplestone 2011; Garnier-Laplace et al. 2013).
90 Long-term exposures to environmental stressors have been linked to lasting responses in
91 organisms within, but also over multiple exposed generations (Mirbahai and Chipman 2014;
92 Schultz et al. 2016; Jimenez-Chillaron et al. 2015; Marczylo et al. 2016; Hanson and Skinner
93 2016). Yet, the outcome of a long term-exposure to pollutants is not always predictable. For
94 example, chronic exposure to pollutants or adverse conditions has been shown to lead to
95 changed phenotypes (Singer et al. 2016; Gonzalez et al. 2016; Potters et al. 2007) resulting in
96 adaptation within a population (Costa et al. 2012; Coors et al. 2009; Bible and Sanford 2016).
97 In contrast, there is also evidence suggesting that long term exposures to environmental
98 stressors can lead to an increased population sensitivity (Parisot et al. 2015) that may result
99 in population declines (Vasseur and Cossu-Leguille 2006). This makes predicting the long-term
100 and/or transgenerational consequences of exposure to a stressor a particular challenge for
101 estimating risks to populations (Groh et al. 2015).

102 Selection has been recognised as a major mechanism through which adverse environmental
103 conditions can impact the phenotypes of successive generations. Selection of alleles
104 associated with tolerance can lead to changes in the phenotypic characteristics within a
105 population and, hence, is known to be a key driver of changes in population level sensitivity
106 to pollutant effects (Van Straalen and Roelofs 2007). Detailed studies of populations inhabiting
107 polluted sites have identified numerous cases of modified phenotypes and also of specific
108 genetic selection at loci that lead to biochemical changes that underpin adaptation. Examples
109 cover exposure to radionuclides, trace metals and persistent organic pollutants and taxa such
110 as cladocerans (Hochmuth et al. 2015; Jansen et al. 2015), collembola (Costa et al. 2012; Nota
111 et al. 2013), chironomids (Groenendijk et al. 1999; Loayza-Muro et al. 2014), terrestrial and
112 freshwater annelids (Kille et al. 2013; Langdon et al. 2003; Levinton et al. 2003), fish (Wirgin
113 et al. 2011; Shaw et al. 2014; Reid et al. 2016; Theodorakis and Shugart 1997), plants, birds
114 (Ellegren et al. 1997) and small mammals (Theodorakis et al. 2001). Although selection for
115 enhanced tolerance is a commonly observed phenomenon, some data have shown that rapid
116 adaptation towards heavy-metals or radionuclides in organisms cannot be explained only by
117 increased mutation rates, but could also be due to non-genetic changes in the activity of
118 functional genes and these might be heritable over generations (Geras'kin et al. 2013;
119 Kovalchuk et al. 2003; Mirbahai and Chipman 2014; Kille et al. 2013; Wang et al. 2017). This
120 has revealed further levels of complexity probably provided by relevant epigenetic
121 mechanisms relating to structure and regulation of gene expression and splicing that have the
122 potential to transfer information over generations.

123 In this paper an overview is given of epigenetic changes induced after long-term (within and
124 over generations) exposure to ionising radiation. Although different epigenetic mechanisms

125 will be discussed the main focus of the current review will be on comparing the evidence from
126 both lab and field studies on changes in DNA methylation.

127

128 **2. Overview of epigenetic mechanisms**

129 The first definition of epigenetics, as ‘the causal interactions between genes and their
130 products, which brings the phenotype into being’, was provided by Waddington (1939) long
131 before any mechanistic understanding of the relevant processes had developed. This
132 definition has since been refined. For example, Wu and Morris (2001) defined epigenetics as
133 ‘Nuclear inheritance which is not based on changes in DNA sequence’ or Bird (2007) as ‘the
134 structural adaptation of chromosomal regions so as to register, signal or perpetuate altered
135 activity states’. This reflects that epigenetics is now widely seen as ‘the study of the landscape
136 of mitotically and/or meiotically heritable changes in gene activity and transcript architecture,
137 including splicing variation, that cannot be explained solely by changes in DNA sequence
138 (Vandegheuchte and Janssen 2011; Allis et al. 2007; Berger et al. 2009).

139 The epigenetic landscape is shapen by three epigenetic marks; DNA methylation, histones and
140 it's post translation modifications and small RNA interactions. Together they shape the
141 structure of the DNA called chromatin (Allis and Jenuwein 2016). These major epigenetic
142 players are engaged in a network of interconnected ‘cross-talk’ (Irato et al. 2003; Iorio et al.
143 2010) and orchestrate gene expression that “...underpins the differences between species,
144 ecotypes and individuals” (Mattick et al. 2009; Brautigam et al. 2013). Well established as a
145 key mechanism involved in the aetiology of human disease (Huang et al. 2003), it is only
146 relatively recently that the significance of epigenetic mechanisms in toxicology (Szyf 2007),
147 ecology (Bossdorf et al. 2008) and evolutionary biology (Rapp and Wendel 2005), has begun
148 to emerge. Within ecology, it has been suggested that epigenetics could define “... where the

149 environment interfaces with genomics ... (and could provide a) rapid mechanism by which an
150 organism can respond to its environment without having to change its hardware” (Pray 2004).
151 Studies on plants have indicated that epigenetic systems provide functional links between the
152 detection of environmental change and regulation of gene expression (Bossdorf et al. 2008;
153 Grativol et al. 2012; Whittle et al. 2009; Rasmann et al. 2012; Verhoeven et al. 2016; Sahu et
154 al. 2013; He and Li 2018). Similarly in animals, the role of specific components or changes of
155 the epigenome in species responses to environmental stress has been demonstrated
156 (Vandegheuchte and Janssen 2014; Schott et al. 2014; Marsh and Pasqualone 2014; Mirbahai
157 and Chipman 2014; Wang et al. 2017; Marczylo et al. 2016). Thus epigenetic mechanisms
158 appear to play an important role in determining the physiological responses of species to long-
159 term multigenerational exposure, including to persistent stressors such as radionuclides.

160 To integrate emerging understanding of epigenetic mechanisms with existing mechanistic
161 knowledge in radioecology, a clear understanding of long-term effects induced by ionizing
162 radiation exposure of non-human species and their potential (epigenetic) mechanistic basis is
163 needed. To provide this, we here give a brief overview of the evidence of trans- and
164 multigenerational effects in organisms exposed to ionising radiation. The potential role and
165 value of epigenetic analyses in site-specific studies in radioecology will be discussed, including
166 their relevance for future radiological risk assessment. As the most widely studied mechanism
167 and its potential to be transferred to the next generation, special attention will be given to
168 changes in DNA methylation (locus-specific or total) as a possible marker for exposure to
169 ionising radiation, including under field conditions.

170

171 **3. The biology of epigenetic mechanisms**

172 DNA methylation, histone modifications, and small non-protein coding RNA molecules are the
173 major known epigenetic mechanisms. DNA methylation is the addition of a methyl group to
174 the one of the DNA bases (cytosine or adenine). Most prevalent DNA methylation is on the
175 fifth position of the cytosine ring (5-methyldeoxycytidine, mC). In vertebrates this usually but
176 not exclusively located at in CpG sites. For example, in *Drosophila* methylation is mostly found
177 in the context of CpT dinucleotides (Feil and Fraga 2012), in honey bees there appears to be a
178 clear distinction of CpG sites in exons and non-CpG sites in introns (Cingolani et al. 2013) and
179 in plants and embryonic stem cells also at CHG and CHH sites (H=A,T or C) in addition to CpG
180 (Feil and Fraga 2012; Cingolani et al. 2013).

181 In vertebrates, around 60% of genes are associated with CpG islands that occur at or near the
182 transcription start site of, particularly, housekeeping genes (Gardiner-Garden and Frommer
183 1987). The hypermethylation in CpG rich promoters can be associated with the repression of
184 gene expression (Bock 2012). In invertebrates, methylation is targeted more towards gene
185 body, potentially playing a role in alternative splicing and gene function diversification (Flores
186 et al. 2012; Asselman et al. 2016). Cytosines can be methylated via maintenance and *de novo*
187 methyltransferase enzymes (Law and Jacobsen 2010). In vertebrates, maintenance
188 methylation by DNMT1 occurs during the S-phase of mitosis, where the newly synthesized
189 DNA strand is methylated using the original strand as template. De novo DNA methylation is
190 undertaken DNMT3 family members, although recent insights have shown redundancy
191 between to two DNMT family members (Lyko 2018). *De novo* DNA methylation is undertaken
192 DNMT3 family members. In plants the homologues of DNMT3, DOMAINS REARRANGED
193 METHYLTRANSFERASE 1/2 (DRM1/DRM2) are responsible for the de novo methylation
194 whereas maintenance of CG methylation is conducted by DNA METHYLTRANSFERASE 1
195 (MET1) which is a homolog for DNMT1 (Law and Jacobsen 2010; Chan et al. 2005). In addition

196 the plant specific CHROMOMETHYLASE 3 (CMT3) is responsible for maintaining methylation
197 in a context of CHG and together with DRM1/DRM2 for methylation in a CHH context (Chan
198 et al. 2005). Although the methyltransferase enzymes are the core proteins involved in
199 methylation, they are recruited and guided to their specific interaction targets by proteins,
200 such as UBIQUITIN-LIKE, CONTAINING PHD AND RING FINGER DOMAINS 1 (URHF1) and
201 PROLIFERATING CELL NUCLEAR ANTIGEN (PCNA) (Baubec et al. 2015). A further insight that
202 has recently emerged is that DNA methylation represents only one part of the DNA
203 methylation cycle. Recently, Tet methylcytosine dioxygenases (previously named ten-eleven
204 translocation (TET) proteins) have been identified as crucial proteins in putative
205 demethylation pathways (Coulter et al. 2013; Scourzic et al. 2015). Indeed, the dynamics
206 between methylation and hydroxymethylation exemplifies the balance of DNA methylation at
207 specific regions as well as globally during early developmental reprogramming (Wu and Zhang
208 2014).

209 Histone modifications occur as post-translational modifications predominantly to the N and C
210 terminal tails of histone proteins. Histone proteins are organised in octamer structures
211 forming nucleosomes as the fundamental units of chromatin (Berr et al. 2011). Initially
212 histones were thought of as primarily structural proteins. However, it is now recognised that
213 they play a pivotal role in regulating gene expression via structural changes of chromatin (Jung
214 and Kim 2012; Margueron et al. 2005). Major histone modifications include acetylation,
215 methylation, phosphorylation and ubiquitination (Bannister and Kouzarides 2011). A key role
216 played by histone isoforms and post-translational modifications that is highly relevant to
217 ionising radiation exposure, is their involvement in DNA damage repair (Hunt et al. 2013;
218 Mondal et al. 2016). DNA repair requires multiple steps, including the initial signalling of the
219 break, the opening of the compact chromatin to facilitate access for repair factors, and

220 afterwards the restoration of the chromatin state (Hunt et al. 2013; for details see Huertas et
221 al. 2009). An authoritative overview of the post-translational modifications in histones
222 triggered in response to DNA damage is given by Méndez-Acuña et al. (2010). Changes of
223 histone modifications have also been linked to exposure to different pollutants in both
224 mammalian and non-mammalian species (Kim et al. 2012b; Mendez-Acuña et al. 2010; Santos
225 et al. 2011; Wang et al. 2017). Observations of heterochromatin state maintenance over
226 multiple successive generations following exposure to heat or osmotic stress in *D.*
227 *melanogaster* suggests a mechanism by which the effects of stress are inherited epigenetically
228 via the regulation of chromatin structure (Seong et al. 2011).

229 Short interfering RNAs and microRNAs are functional non-coding RNA molecules. They are not
230 translated into proteins and are involved in gene repression via RNA deactivation and
231 degradation (Castel and Martienssen 2013). Single microRNAs may on average interact with
232 ~400 different protein coding genes. Hence, changes in microRNA expression are proposed to
233 be a key component of organism response to stressor exposure (see e.g. for plant responses
234 Huang et al. 2016). Reduced expression of microRNA has been found in response to insecticide
235 and fungicide exposure (Qi et al. 2014; An et al. 2013). MicroRNAs have been shown to be
236 intimately involved in cellular response to metals such as cadmium and arsenic (Liu et al. 2016;
237 Meng et al. 2011; Gielen et al. 2012). Important roles of non-coding RNAs in the epigenetic
238 inheritance of DNA methylation through cell division and guiding de novo methylation after
239 meiosis indicate key interactions between epigenetic pathways (Calarco et al. 2012; Larriba
240 and del Mazo 2016). In plants e.g. DNA and histone methylation by DRM2 activity and
241 subsequent gene silencing can also be mediated by siRNAs ARGONAUTE (AGO4) and
242 polymerase V (POLV) (Holoch and Moazed 2015; Neeb and Nowacki 2018). Hence dynamic

243 interactions of different epigenetic mechanisms would be expected in response to
244 environmental challenge.

245 The relative role of the different epigenetic mechanisms can vary between species. The
246 majority of eukaryotic phyla possess cytosine methylation ranging from <<1% in some taxa
247 (e.g. many arthropods) to >10% for annelids, molluscs and vertebrates, with species such as
248 *C. elegans* even proposed to lack cytosine methylation completely (Regev et al. 1998) or to be
249 very low (~0.0033%) (Hu et al. 2015). Because of those variations in DNA methylation levels,
250 it was initially uncertain how important cytosine methylation may be among those phyla.
251 However, evidence of the importance of DNA methylation in heritable responses in
252 invertebrates following stressor exposure has begun to emerge, as well as for other epigenetic
253 mechanisms (Seong et al. 2011; Schultz et al. 2016; Stern et al. 2014; Klosin et al. 2017). For
254 some species, and particularly in *C. elegans*, a second DNA modification based on methylation
255 of the N-6 position on adenine may also act as an alternative form of DNA methylation (Greer
256 et al. 2015). In addition, the balance between DNA methylation, post-translational
257 modifications and types of microRNA molecules (both of which are species specific and highly
258 dynamic), presents a challenge to tease apart the roles that different epigenetic mechanism
259 play in gene expression dynamics and ultimately phenotypic responses to stress including
260 those in species exposed to radionuclides and other pollutants over extended timescales (Lim
261 and Brunet 2013).

262

263 **4. Main methods used to detect DNA methylation changes**

264 This review will mainly focus on the evidence for DNA methylation changes induced by
265 radiation in different animals and plants and this in both lab and field conditions. The
266 measurement of total DNA methylation levels is now routine using molecular genetic and

267 biochemical protocols. These analyses provide a useful picture of overall methylation states.
268 The methods have the advantages of reasonable cost per sample, established protocols,
269 sensitivity to overall methylation pattern change and rapid sample processing (Table 1). Two
270 global methylation methods that are commonly used are methylation sensitive amplified
271 fragment length polymorphisms (meAFLP) and measuring the % of methylated cytosine by
272 HPLC-MS/MS. The meAFLP technique is based on the use of two restriction enzymes, HpaII
273 and MspI. Both HpaII and MspI recognize a CCGG sequence. MspI is able to cut both
274 methylated recognition sites as well as unmethylated ones. In contrast, HpaII is unable to cut
275 at such locations when methylated (i.e. only unmethylated recognition sites are cut).
276 Methylation of these restriction sites can be assessed by electrophoretic recording bands cut
277 by MspI but not HpaII on a fragment analyser (e.g. capillary sequencer). The method has been
278 shown to demonstrate limited variability and has the benefit of an internal control (EcoRI) to
279 account for variability in the amount of DNA input. The detection of methyl groups by HPLC-
280 MS/MS allows highly sensitive quantification of methylated and hydroxymethyl cytosines (5
281 mC and 5-hmC) present in a hydrolysed DNA sample. The specific ability to detect and measure
282 5hmC is a specific advantage of this technique, given its recently demonstrated roles in
283 development (Pastor et al. 2011; Song et al. 2011; Xu et al. 2011).

284 Although useful, application of global methylation analysis methods do not allow analysis of
285 the specific methylation states needed to assess functional links between changes in site
286 specific methylation, gene expression changes and phenotypic changes to be made. The use
287 of methylation mapping techniques can provide improved resolution to identify and assess
288 specific genes/regulatory regions of interest that are differentially methylated under specific
289 treatment or exposure conditions. The number of options to study DNA methylation have
290 become more diverse and methods such as reduced representation or whole genome bisulfite

291 sequencing, are now considered close to routine. The value of these genome wide
292 methylation mapping techniques is that they go beyond the level of an overall change to
293 identify the gene associated sites of differential methylation. These methods are of course
294 limited when an organism reference genome is either not available or is poorly assembled or
295 annotated. Hence, significant effort needs to be given to genome resource development
296 before these methods can be used to study autochthonous species.

297

298 ***5. Laboratory evidence for multigenerational and transgenerational effects including those***
299 ***induced by ionising radiation***

300 The interest in understanding the effects of persistent pollutants, including radionuclides, on
301 population exposed for more than a single generation is ongoing. Therefore studies of
302 multigenerational and transgenerational stressor effects on apical phenotypes have become
303 more common. For multigenerational studies, exposure to the stressor in question is
304 maintained in a continuously cultured and exposed population for successive generations
305 (e.g., continuously exposed F0, F1, F2 etc.) to allow the consequences of multigenerational
306 exposure to be assessed. Phenotypes are observed in those generations directly exposed. For
307 these multigenerational cases, the simplest expectation is that the observed toxicity in the
308 offspring is not greater than that in parents exposed over their full life-span (i.e. embryo until
309 death), at least over initial generations, with possible development of tolerance over longer
310 time-scales. Transgenerational experiments, on the other hand, consider not just effects on
311 the exposed generation, but also effects on subsequent unexposed generation(s) reared after
312 hatching in stressor free conditions (Skinner and Guerrero-Bosagna 2009; Skinner 2016; Groot
313 et al. 2016). In such studies, stressor effects may be expected as a result of exposure of the F0
314 mothers in F1 embryo and F2 germline, but not in later offspring. The simplest expectation

315 from transgenerational experiments is thus of physiological effects no greater than those
316 observed in F0s, only in F1s (and possibly F2s), with no further such effects on the later (F3
317 etc.) generations.

318 There are cases where the simplest expectations of multigenerational and transgenerational
319 exposure are met, including examples for plants (Iglesias and Cerdan 2016; Groot et al. 2016;
320 Molinier et al. 2006), earthworms (Hertel-Aas et al. 2011), zebrafish (Baker et al. (2014)
321 (Schwindt et al. (2014) and mice (Ziv-Gal et al. (2015). However, critical analysis of reported
322 multigenerational exposures covering a range of stressor types including radionuclides,
323 metals, nanomaterials, organic chemical and antibiotics, suggests that, at least over the
324 durations used in the laboratory (usually < 10 generations) the simplest expectation of similar
325 sensitivity to F0 in later generations are not always be met. In a number of published cases,
326 an increasing sensitivity in later generations has been observed (see Table 2 and examples
327 below). While this prevalence may partly result from publication bias and from the clonal
328 organisms used, the high frequency of such responses does suggest that increased sensitivity,
329 at least over the initial generations of a multigenerational exposure, may be a common
330 phenomenon (see Table 2).

331 For exposure to radiation and radionuclides there are a number of multigenerational lab-
332 studies that have reported patterns of increased generational sensitivity for continuously
333 exposed populations (see Table 2 for exposure details). For daphnids it has been reported that
334 the progeny of organisms continuously exposed to gamma radiation, Am⁴²¹ (and depleted
335 uranium) show higher sensitivity in the F1 and F2 generations than that for parents depending
336 on the endpoint measured (Pane et al. 2004; Biron et al. 2012; Alonzo et al. 2008b; Parisot et
337 al. 2015). Similarly, Zaka et al. (2004) exposed 5-day old *Pisum sativum* plants over three
338 generations to different acute doses of gamma radiation. Results indicated that doses

339 apparently harmless for the parental plants adversely affected the F2 generation. *Arabidopsis*
340 *thaliana* plants exposed to different dose rates of gamma radiation during the vegetative
341 growth stage for one or two generations also showed greater response in the later generation.
342 In this case, increased responses of antioxidative enzyme activity were measured in
343 multigenerationally exposed plants (van de Walle et al. 2016). This response was accompanied
344 by phenotypic changes, such as accelerated flowering after multigenerational exposure
345 (Horemans et al., pers. comm).

346 Transgenerational studies with radionuclides or after radiation-exposure have shown
347 responses not just in continuously exposed generations, but also in later unexposed
348 generations. A study of reproductive effects of gamma radiation in the nematode *C. elegans*
349 exposed from F0 to F2, either continuously or only at F0 generation also found
350 transgenerational effects in F2 organisms greater than in the initially exposed nematodes
351 (Buisset-Goussen et al. 2014). Daughter cells of chronically gamma-radiation-exposed *Lemna*
352 *minor* plants died off notwithstanding only a limited growth reduction in the exposed mother
353 colonies (10-30%) indicating that the effects were, thus, greater in the recovering non-
354 exposed plants than in the exposed F0s (Van Hoeck et al. 2017). These examples of
355 transgenerational effects leading to increased sensitivity of progeny match similar results
356 found for other stressors, suggesting a possible common mechanism (Schultz et al. 2016;
357 Moon et al. 2017; Annacondia et al. 2018; Groot et al. 2016).

358 The current multigenerational and transgenerational toxicity literature is dominated by lab-
359 studies with relatively high exposure dose rates (7-420 mGy/h, see table 2) and for
360 ecotoxicological relevant species like *C. elegans*, *D. magna* and zebrafish (Table 2). For *C.*
361 *elegans* and *D. magna*, the experimental populations that have been used in most
362 laboratories, multigenerational and transgenerational exposure studies are clonal. Hence, the

363 potential for selection of alleles that may lead to evolution of tolerance in later generations in
364 a multigenerational exposure experiment is limited. This is true especially because the
365 majority of such experiments are conducted over only a relative limited number of
366 generations (<10 and usually ≤ 3). Indeed, when nematodes were continuously exposed for
367 22 generations to U, adaptation was shown to occur (Dutilleul et al. 2014). Although many
368 studies have shown generationally increased sensitivity and its transfer, the clonal nature of
369 species may be accentuated, because the limited genetic variation of the inbred strains. In the
370 study of Dutilleul et al. (2014) for nematodes discussed above, the population used that
371 showed adaptation composed of wild isolates with increased genetic diversity above the
372 clonal *C. elegans* strains used for previous multigenerational studies. Hurem et al (2018b)
373 showed effects on the transcriptome in offspring from irradiated zebrafish that were even
374 accentuated in offspring produced from the same parents does, however, indicated the
375 potential to identify epigenetic responses in a genetically diverse population.

376 Multigenerational exposure experiments by their nature involve continuous incubation of
377 populations with a toxicant or stressor, with generational phenotyping to allow detection of
378 changes in sensitivity. In such studies, increased sensitivity in the progeny could theoretically
379 arise if any toxicant induces "damage" that can be transferred to subsequent exposed
380 generations. Indeed Parisot et al. (2015) highlighted a possible role of DNA damage in
381 multigenerational effects by finding a correlation between increased sensitivity and the
382 transmission of DNA damage in daphnids exposed to gamma radiation. This possible role of
383 DNA damage and genome instability in multigenerational and transgenerational effects may
384 lead to hypotheses about the type of stressors that may cause such phenomena.

385 The role of both paternal and maternal effects has received much research attention in
386 ecology and toxicology (Frost et al. 2010; Wigle et al. 2007). Within these studies there is

387 strong evidence that indicate how the direct exposure of the developing embryo and germline
388 can be adversely affected as a result of exposures to environmental pollutants. However, in
389 addition to these more direct effects, there is evidence of a potential role of the epigenome
390 in the transfer of aberrant phenotypes to F1 offspring and indeed to generations beyond
391 (Bowman and Choudhury 2016; Chen and Baram 2016; Wang et al. 2017). For example,
392 exposing *C. elegans* to nanoparticles resulted in aberrant phenotypes, that were persistent in
393 future unexposed populations for 3 or more generations (Greer et al. 2011; Katz et al. 2009;
394 Rechavi et al. 2014; Schultz et al. 2016). When transgenerational effects occur over these
395 generation scales, germline exposures alone cannot be solely responsible, with the potential
396 that epigenetic mechanisms may be intimately involved.

397

398 ***6. Evidence for long-term effects induced by radiation on the environment coming from field*** 399 ***studies***

400 The nuclear accidents of Chernobyl and Fukushima have made it possible to investigate
401 possible effects of radiation on a whole range of organisms exposed to radionuclides under
402 field conditions over extended timescales. The temporal changes that occurred in radiation
403 exposure in the CEZ and the FEZ, have resulted in a specific time course of responses among
404 non-human biota in the regions (IAEA 2006; Beresford et al. 2016; Beresford and Coplestone
405 2011; IAEA 2015). The most pronounced biological effects were seen in the first and second
406 phases after the accident. In these early stages, the high doses experienced shortly after the
407 accident by the forest located to the west of the Chernobyl reactor, later designated as the
408 Red-forest. In this Red-forest massive death of pine trees was observed, while deciduous
409 species survived despite an early loss of leaves and damage to woody tissues (Arkhipov et al.
410 1994; Kryshev et al. 2005). Similar morphological differences such as loss of apical dominance

411 were recently also reported in Japanese red pine in the FEZ (Yoschenko et al. 2016). In the first
412 phase after the nuclear accidents, direct effects such as a decrease in numbers of small
413 mammals as well as reduced development or survival of embryos was also seen (Geras'kin et
414 al. 2008) and the loss of specific groups of soil biota were also recorded in the most
415 contaminated areas (Krivolutsky 1996; IAEA 2006). These effect could also be linked to the
416 high levels of initial exposure that were experienced following both nuclear accidents. Initial
417 dose rates in the most contaminated areas of CEZ were as high as 5mGy/h (IAEA 2006)

418 The second phase characterised by a decrease in dose rates due to disappearance of short-lived
419 radioisotopes and wash-out and run-off (IAEA 2006). This phase started from two months
420 after the accidents, was associated with reductions (up to a factor of 30) in the density of
421 invertebrates living in the forest litter experiencing greatest contamination. These decreases
422 were linked to radionuclide exposure effects on reproduction and recruitment (Krivolutsky
423 and Pokarzhevskii, 1992; Krivolutsky et al., 1992).

424 In the third exposure phase resulting from the Chernobyl accident, most strongly affected
425 populations of species of pine trees and soil invertebrates were shown to slowly start to
426 recover (Arkhipov et al. 1994; Zelena et al. 2005). Recovery from the initial negative effects
427 was also found in birch pollen, embryonic cells of herbaceous plants like evening primrose
428 embryonic cells (Boubriak et al. 2008) and *Arabidopsis thaliana* (Kovalchuk et al. 2004) and in
429 exposed birds (Galvan et al. 2014). In this phase Cs-137 and Sr-90 are the main contributors
430 to the dose with some additional Am-241 and Pu-isotopes for CEZ and Cs-137/134 for FEZ
431 (Horemans et al. 2018; Saenen et al. 2017). Ambient dose rates now measured are maximally
432 0.5 mGy/h and these can be found in the forest western from the nuclear power plant
433 designated as the Red Forest (Beresford, personal communication).

434 In addition to changes observed at individual or population levels, the radiological impacts
435 within both the CEZ and the FEZ, have also been reported at the sub-organismal level.
436 Aberrant cell frequencies were found in the root meristem of plant seedlings (Geras'kin et al.
437 2011). Increased mutation rate (Kuchma et al. 2011) and gene deregulation (Zelena et al.
438 2005), have been seen in pine trees. Increased mitochondrial DNA haplotype and nucleotide
439 diversity have been reported in bank voles (Matson et al. 2000; Baker et al. 2001),
440 chromosomal aberrations in mice (Kubota et al. 2015) and in soil invertebrates, increased DNA
441 damage in earthworms (Fujita et al. 2014). Most of these studies so far have, however, failed
442 to find a link between these observed sub-organismal effects and impacts at higher level of
443 biological complexity such as radiation-induced phenotypical changes and long-term effects
444 on population dynamics (Meeks et al. 2009; Meeks et al. 2007).

445 The adaptive responses that have been indicated during the extended third phase of exposure
446 following the two accidents at Chernobyl and Fukushima are at least in part due to the
447 reduction over time in dose rates and, hence, exposure. Although a memory-effect of the early
448 high exposures cannot be excluded, the decreased exposure in the third phase might allow
449 both increased *in-situ* recruitment and survival leading to positive population growth, as well
450 as the survival of inwardly migrating individuals (Jackson et al. 2004; Boubriak et al. 2008;
451 Boubriak et al. 2016). Additionally it is also possible that increased tolerance, through
452 selection and as a result of favourable mutations may make a contribution (Kovalchuk et al.
453 2003). However, in *Arabidopsis* no additional mutations compared to plants collected in
454 control sites were found in the CEZ (Abramov et al. 1992). Ostensibly the probability of
455 favourable mutations may be seen as unlikely. Assuming a germline mutation rate in plants of
456 about 10^{-5} to 10^{-6} per gamete, one would expect only one mutation in 500,000 plants
457 (Kovalchuk et al. 2003). Consequently it has been proposed that rapid adaptation may be more

458 strongly linked to epigenetic processes in the development of locally adapted phenotypes at
459 polluted sites (Kovalchuk et al. 2003).

460

461 ***7. Evidence for a role of epigenetics in long-term or transgenerational responses to*** 462 ***radiation-induced stress***

463 Studies on the effects of stressors on the epigenome of organisms under environmentally
464 relevant exposure conditions have covered examples for ionising radiation exposure and for
465 a range of chemical and non-chemical stressors in different species. Within these studies, a
466 range of epigenetic mechanisms and endpoints have been considered (for review see e.g.
467 Aluru 2017; Bruce et al. 2007; Kim et al. 2012b; Mirbahai and Chipman 2014). Initial adaptive
468 changes resulting from exposure to these different stressors have been found for key
469 components of the epigenome, such as DNA methylation (Vandegheuchte and Janssen 2011;
470 Marczylo et al. 2016), non-coding RNAs (Kure et al. 2013; Wang et al. 2013; Song et al. 2012)
471 and histone modifications (Raut and Sainis 2012; Mondal et al. 2016). Changes in microRNA
472 expression have further been shown to be involved in metabolism following starvation and
473 the transfer of longevity (Greer et al. 2011; Katz et al. 2009; Rechavi et al. 2014). In plants,
474 small RNAs play an important role in chromatin remodelling and DNA methylation through
475 RNA-directed DNA methylation also in different abiotic stresses in plants (Hirayama and
476 Shinozaki 2010).

477 Although long a controversial issue and still not fully elucidated, recent evidence has
478 suggested that in plants, vertebrates and invertebrates, epigenetic marks induced by adverse
479 conditions encountered by the parents can be partly stable across generations (Uller et al.
480 2015; Klosin et al. 2017; Whittle et al. 2009; Saze 2012; Pecinka and Mittelsten Scheid 2012;
481 Sudan et al. 2018; Stassen et al. 2018; Norouzitallab et al. 2019). Such retention can potentially

482 lead to transgenerational heritable changes in offspring (Verhoeven et al. 2010; McCarrey
483 2012; Guerrero-Bosagna and Jensen 2015; Guerrero-Bosagna et al. 2012). Evidence has been
484 accumulated for the transfer of DNA methylation patterns in the germline (Verhoeven et al.
485 2010; Verhoeven et al. 2016). As an example of the link between epigenetic mechanisms and
486 transgenerationally altered phenotypes a study of transgenerational response to temperature
487 in *C. elegans* has identified altered trimethylation of histone H3 lysine 9 as a mechanism for
488 transgenerational inheritance (Klosin et al. 2017). On the other hand, in *Arabidopsis*, nickel
489 chloride caused a change in DNA methylation patterns and some of this was inherited by the
490 following generation (Li et al. 2015). In the offspring of mechanically wounded *Mimulus*
491 *guttatus* plants changes in methylation could be associated with transgenerational plasticity
492 (Colicchio et al. 2018). Depending on the methylation context, CG or non-CG methylation,
493 these changes were found to be in gene coding regions or transposable elements, respectively
494 (Colicchio et al. 2018). Dandelions (*Taraxacum officinale*) also showed altered DNA
495 methylation that was largely inherited by the next generation of the asexually reproducing
496 plants when exposed to a number of different stressors (Verhoeven and van Gurp 2012;
497 Verhoeven et al. 2016).

498 A growing number of papers also indicate that exposure to ionising radiation will lead to
499 changes in epigenetic markers (Table 3). For example, scots pine trees present in the most
500 contaminated areas around the Chernobyl nuclear reactor have been found to have
501 hypermethylated DNA, with this hypermethylation directly (Kovalchuk et al. 2003) or
502 transiently associated with the radiation dose received (Volkova et al. 2018). Further work
503 established that the genomes of young trees planted on contaminated soil showed higher
504 levels of cytosine methylation than trees in uncontaminated soil. However, levels of cytosine
505 methylations in plants grown in clean soil from seeds taken from previously exposed plants

506 were not found to differ significantly from controls Kovalchuk et al., (2003). Hence these
507 results are suggestive of a within generation genome methylation effect, rather than of any
508 multigenerational or transgenerational mechanism, as a result of exposure during the somatic
509 development. However, since only overall levels of DNA methylation inheritance was
510 addressed, the potential for loci specific cannot be discounted.

511 In a study of the progeny of *Arabidopsis sp.* sampled in three consecutive years from areas
512 with different levels of contamination within the CEZ, higher resistance to mutagens in
513 progeny of plants from the most contaminated sites compared to unexposed plants was
514 identified (Kovalchuk et al. 2004). This difference in sensitivity could be attributed to higher
515 expression of free radical scavenging enzymes and DNA-repair enzymes and was associated
516 with global genome hypermethylation in the contaminated site plants. It was hypothesised
517 from these data that epigenetic regulation of gene expression and genome stabilization may
518 play a key role in the underlying processes that stabilise *Arabidopsis* genome architecture
519 under exposure to ionizing radiation exposure (Kovalchuk et al. 2004). A number of papers
520 have proposed a link between epigenetic effects and non-targeted effects (NTE) such genomic
521 instability and bystander effects (Schofield and Kondratowicz 2018). However, while the
522 existence of non-targeted effects is well established (Morgan 2002; Kadhim et al. 2004; Pouget
523 et al. 2018; Burdak-Rothkamm and Rothkamm 2018), and studies have shown an association
524 between the two effects (e.g., Kaup et al. 2006; Xu et al. 2015), evidence of a causal
525 relationship is more elusive, since NTE could be either a mechanism or a consequence of
526 epigenetic changes (Schofield and Kondratowicz 2018). Changes in the level of DNA
527 methylation may be intimately linked with transcription remodelling in response to radiation
528 exposures, including changes to the pathways involved in antioxidant defence and DNA repair.
529 Confirmation of such effects would require the use of combined genome wide DNA

530 methylation mapping and transcriptomic approaches to allow loci specific methylation to be
531 associated with gene expression phenotypes in exposed plants.

532 A study of the pale blue grass butterfly *Zizeeria maha* within the FEZ has provided a further
533 indication of the potential for heritable epigenetic changes in a population exposed to ionising
534 radiation (Hiyama et al. 2012; Hiyama et al. 2013). Mild morphological abnormalities were
535 observed on some individuals of adult butterflies collected one month after the accident, but
536 an increase of the severity of these abnormalities occurred in the F1 generation that were
537 further inherited by F2 progeny. These abnormalities and their transgenerational transfer
538 were proposed to be attributable either to random mutation on important genes or through
539 epigenetic mechanisms. As the underlying mechanisms of these effects were not studied by
540 the authors, leaving the mechanistic basis of the observed effects and their inheritance remain
541 an open question.

542 Recently a number of European research groups have combined research efforts to study
543 possible epigenetic changes in organisms exposed to ionizing radiation, in the laboratory or *in*
544 *situ* (Chernobyl or Fukushima), in a range of species (plants, earthworms, fish, frogs) (Table 3).
545 The focus of the combined efforts was to better understand the possible role of these
546 mechanisms in the induction of long-term/transgenerational effects and their relevance as
547 possible biomarkers of ionising radiation (Adam-Guillermin et al. 2013). The organisms chosen
548 were all reproductive non-clonal organisms. Hence the work addresses multigenerational and
549 transgenerational effects in genetically diverse populations. For example, in offspring of
550 zebrafish that were exposed to ionising radiation during gametogenesis, a large number of
551 differentially methylated regions were observed, with five specific loci showing a persistent
552 effect up to the third generation (Kamstra et al. 2018). These methylation changes could be
553 linked to changes in gene pathways and adverse effects found in progeny (Hurem et al. 2017;

554 Hurem et al. 2018b). In the same exposure study, miRNA expression was measured in first
555 filial offspring and histone marks H3K4me3, H3K9me4 and H3K27me3 at 3 specific
556 loci(Lindeman et al. 2019). There were 23 differentially expressed miRNAs indicating a
557 multifaceted response to ionising radiation exposure (Martin et al. 2019, in preparation).
558 Differentially enriched histone marks were observed as well at the three measures loci in F1
559 offspring, but interestingly these effects were diminished in F2 offspring (Lindeman et al. 2019,
560 submitted). Although only exposed embryo's were analysed similar changes in histone markes
561 were found for Atlantic salmon (*Salmo salar*) at higher dose rates (Lindeman et al. 2019).
562 A dose-rate dependent induction of total methylation levels was observed in *A. thaliana* plants
563 exposed in the lab to different levels of gamma radiation for up to three generations (Saenen
564 et al. 2017)). Moreover triple methyltransferase mutants (*drm1drm2cmt3*) of *A. thaliana*
565 showed increased sensitivity to irradiation including an increased induction of oxidative stress
566 (Saenen et al. 2017).
567 In the clonal cladoceran *Daphnia magna*, transgenerational inheritance of DNA methylation
568 changes were studied using bisulphite sequencing, after irradiation of generation F0 to 6.5
569 $\mu\text{Gy/h}$ or 41.3 mGy/h (Trijau et al. 2018). Significant methylation changes at specific CpG
570 positions in every generation were found, independent of dose rate and with a majority of
571 hypomethylation. The total number of common differentially methylated regions was greatest
572 between generations F2 and F3, with three specific persistent loci associated to genes known
573 to play a role during exposure to ionising radiation. The results above suggest a role of
574 enhanced methylation induced by chronic exposure to radiation in lab-conditions and indicate
575 the multi- and transgenerational natures of these responses.
576 For earthworms, studies of DNA methylation in the laboratory and CEZ have shown effects of
577 ionising radiation exposure on DNA methylation pattern as measured by methylated AFLP

578 analysis (Saenen et al. 2017). There are, however, specific challenges in the interpretation of
579 the role of radionuclide exposure in these responses. Large differences in genetic diversity
580 that may occur between morphological similar earthworm “species” may, for example, make
581 it difficult to identify DNA methylation changes unless clades are assessed separately. Indeed
582 clades of the earthworm *Lumbricus rubellus* were found to differ in the nature of their genetic
583 and DNA methylation responses to soil contamination by copper and arsenic (Kille et al. 2013).
584 A similar response was found within an analysed laboratory experiment, where both between
585 and within species allelic differences precluded the identification of a clear DNA methylation
586 profile response to exposure. In CEZ collected earthworm from two species *Aporrectodea*
587 *caliginosa* and *Octolasion lacteum*, a clear site specific change in DNA methylation status was
588 found (Saenen et al. 2017) in *Aporrectodea caliginosa*, while only limited separation was found
589 for *Octolasion lacteum*. While these site specific changes in DNA methylation patterning may
590 indicate a response to radionuclide exposure, a caveat is that the earthworms were collected
591 from sites that differ in the prevailing ecosystem characteristics (wetland and garden sites).
592 An *in situ* study of DNA methylation in frogs collected from a range of differently polluted sites
593 within the Fukushima impacted area indicated that DNA methylation measured as methylated
594 cytosines increased with total absorbed dose rate, up to 7 $\mu\text{Gy/h}$. This increase was
595 concomitant with increased levels of DNA damages (Saenen et al. 2017). As in the study for *A.*
596 *thaliana* in the CEZ (Kovalchuk et al. 2004), this finding of higher DNA methylation associated
597 with increased DNA damage and repair activity supports a functional role of the epigenome in
598 maintaining DNA integrity. These results are in agreement with previous work done on
599 zebrafish exposed to depleted uranium, where changes in DNA methylation patterns both at
600 specific restriction sites and across the whole genome, were observed in F_0 adults and F_1 at
601 the same time as DNA damages (Gombeau et al. 2016; Gombeau et al. 2017). A transient

602 increased methylation with the dose rate was also observed in needles of *Pinus sylvestris*
603 plants collected in radioactively contaminated areas of Belarus (Volkova et al. 2018). In
604 contrast no dose dependent changes in total methylation levels were observed for *C. bursa*
605 *pastoris* plants sampled in spring 2016 in contaminated areas of FEZ. For *A. thaliana* plants
606 collected in CEZ a decrease in global DNA methylation was found in the highest contaminated
607 fields (Horemans et al. 2018).

608 Overall the range of studies of the epigenetic response of species to radionuclide exposure in
609 the laboratory point to a role of the epigenome in adaptive responses. The field studies with
610 plants (pine trees and *Arabidopsis*) showed the potential for ionising radiation to induce
611 changes in DNA methylation levels under field conditions (Georgieva et al. 2017; Kovalchuk et
612 al. 2003; Kovalchuk et al. 2004). For invertebrates, the laboratory and studies in the CEZ and
613 FEZ have partially supported a role of increased methylation in response to radiation among
614 the majority of species studied to date. The challenge from these field studies remains to
615 unequivocally link the observed effects on the epigenome to radiation exposure, rather than
616 to other aspects of environmental variation across the CEZ and FEZ. Studies that specifically
617 investigate changes in mutant lines with reduced DNA methyltransferase activity, as outlined
618 above for *Arabidopsis*, provide initial causal evidence on the validity of such as link.

619

620 **8. Knowledge gaps on epigenetic changes induced by ionising radiation**

621 Although all three different epigenetic layers have been implicated as key mechanisms
622 involved in determining the long-term and transgenerational responses of species to
623 pollutant, including ionising radiation exposure, a majority of studies have to date focussed
624 on the role of DNA methylation (Norouzitallab et al. 2019; Sun et al. 2018; Meehan et al. 2018;
625 Burgio et al. 2018). In cases where difference in DNA methylation response following exposure

626 to ionising radiation are observed, a number of aspects that need further consideration in
627 future work can be drawn.

628 (i) Global methylation alone may be too coarse a measure of epigenetic change to be able to
629 see all biologically relevant differences induced by exposure to low dose rates. As such,
630 differences in methylation might be located in specific sequences of the genome but cannot
631 be detected by global measurements. Therefore, it is important to also include other
632 techniques (e.g. whole genome or reduced representation sequencing) in order to identify
633 specific epigenetic changes and to link these observations to effects on gene expression and
634 physiological change (Paun et al. 2019).

635 (ii) Different DNA methylation response in function of cell type, tissues (as seen in the depleted
636 uranium exposure in zebrafish by Gombeau et al., 2015), or age (as seen in frogs exposed at
637 Fukushima (Saenen et al. 2017), could induce a mosaic of DNA methylation response at the
638 whole organism level, limiting the capability to identify a clear change in methylation pattern.
639 This argues for the analysis of more homogenous tissues or cell types.

640 (iii) Initial changes of DNA methylation resulting from an initial radiation exposure may be lost
641 in individuals exposed over generations of chronic exposure as found for pine trees by
642 Kovalchuk et al. (2003) and in the second generation of lab-exposed *A. thaliana* in a laboratory
643 exposure to gamma radiation. Such results suggest that DNA methylation may be a transient
644 acting potential as an intermediate state preceding later genetic selection and adaptation.

645 (iv) Genetic diversity of species between isolated local populations within the CEZ and FEZ may
646 mean that populations exposed to different levels of radiation may show markedly different
647 epigenetic responses, precluding the identification of a clear exposure response relationship.
648 The presence of natural and man-made barriers to dispersal, which may result in population
649 isolation, across these two zones, may accentuate such differences (Meeks et al. 2007).

650 (v) Although less commonly studied than DNA methylation, the work done to date on the
651 responses of other epigenetic mechanisms like microRNAs or histone modifications to ionising
652 radiation exposure, suggest that these complimentary epigenetic mechanisms may play roles
653 in the response to radiation that may even dominate over DNA methylation changes (Putiri
654 and Robertson 2011; Brautigam et al. 2013);

655 (vii) Long time exposure to radiation might result in selection of alleles linked to tolerance,
656 potentiated potentially by increased mutation (as is seen for frogs in FEZ) that may lead to
657 genetic adaptation that might negate differences in DNA methylation. An interplay between
658 epigenetic changes, notably DNA methylation, and the targeting of mutation has been
659 proposed mechanisms (Putiri and Robertson, 2011 ; Braütigam et al., 2013).

660 (vii) Confounding factors (habitat, soil type, water chemistry; climate etc.) may increase the
661 variability between the samples that may result in changes in DNA methylation that overlie
662 and obscure effects due to ionising radiation making it difficult to link epigenetic change to
663 exposure (see discussion, Garnier-Laplace et al. 2013).

664

665 ***9. Differential DNA methylated regions as possible biomarkers for exposure or effect of a***
666 ***pollutant and its use in risk assessment***

667 There is a strong interest in finding possible biomarkers for exposure and effects of radiation
668 and additionally those that can be markers for long-term effects. Loci specific changes of DNA
669 methylation have been proposed as possible biomarkers for different environmental cues
670 (Meehan et al. 2018) and could possibly be used as molecular fingerprints for e.g. genotoxicity
671 induced when exposed to ionising radiation. However, it is also recognised that significant
672 challenges related to the effects of genetic background and the influence of confounding
673 factors also exist (Pernot et al. 2012). Further studies at environmental realistic doses are

674 needed to assess the prevalence of such responses, including under field conditions. In
675 particular, the use of more targeted methods are needed that identify loci specific changes in
676 DNA methylation, histone modification and the expression of relevant miRNAs.

677 A clear conclusion that emerges from past and ongoing studies concerning the role of the
678 epigenome in response to chronic radiation exposure, lies in the interpretation of changes in
679 methylation patterns from field collected samples in respect to attribution of the principal
680 driver of effects. Specific challenges relate to working with some autochthonous species for
681 which genome resources may be lacking and, the influence of confounding factors which may
682 mask the causal response between ionising radiation exposure and epigenetic changes. In
683 efforts to attribute changes to specific stressor effects, epigenetic approaches may be more
684 powerful indicators of effects when linked to known biomarkers using, for example,
685 transcriptional analysis. When used in conjunction with other mechanistic measurements,
686 epigenetic analysis has the potential to enhance the ecological relevance of molecular
687 biomarkers, as described in the Adverse Outcome Pathway concept (Groh et al. 2015). Given
688 the critical need to establish the nature of effect of prolonged low level exposures, this
689 integrated approach seems a promising way forward, building as it does on existing
690 mechanistic knowledge.

691 The risk assessment process for radiation and radionuclides is largely based on using results
692 from short-term bioassays to predict the effects of exposures in the field. The validity of this
693 laboratory to field extrapolation is one of the key uncertainties in risk assessment (Lourenco
694 et al. 2016). A comparison of field vs laboratory studies has indeed shown that species
695 sampled in the field were 8 times more sensitive than those studied under laboratory
696 controlled conditions (Garnier-Laplace et al., 2013) indicating the need for further thorough lab
697 to field studies. One of the largest differences between laboratory bioassays and field

698 exposures is exposure duration. This is true within a single generation (intergenerational
699 exposure), but even more so when subsequent generations are exposed to the same stressful
700 environment (multigenerational exposure) or when exposure of the parent generation has a
701 subsequent effect on the non-exposed offspring (transgenerational exposure). When
702 multigenerational exposures occur, these may result in effects in later generations that match,
703 and can even exceed those found in exposed F0s (see Table 2). The biological response of
704 species mediated through the genome and epigenome appear to play a role in the
705 development of such effects. Such findings may require a more refined understanding to
706 support and reduce the uncertainty in risk assessment for chronic low dose exposures. Hence,
707 the mechanisms that underlie differential responses within and over generations to previous
708 (sub-lethal) radiation-exposure require further studies to provide a baseline for the
709 development of new approaches such as Adverse Outcome Pathways on low dose radiation
710 exposure, to the risk assessment for both wildlife and human.

711

712 ***10. Conclusions and Recommendations for further development and application.***

713 Work reported to date in both lab and field have indicated changes in DNA methylation
714 resulting from chronic exposure to low dose of ionising radiation. A common conclusion from
715 this work is that both laboratory and field studies have demonstrated changes in overall
716 methylation in organisms exposed chronically to ionising radiation. Generally a chronic
717 enhanced ionising radiation level induced hypermethylation or methylation pattern change
718 which could be taken as a response to induce DNA stability. The main advantage of laboratory
719 studies is the ability to set up controlled multi/transgenerational studies, and avoid
720 confounding factors like local difference in soil characteristics, microclimate. Together with
721 the use of homogeneous populations, this allows for greater insight into the underling

722 mechanisms and processes. Field studies can provide the increased environmental realism of
723 the responses studied. Although data suggest that methylation changes can be observed in
724 different organisms a lower dose rates than those seen in laboratory experiments. The
725 challenge remains to unequivocally link such observations to a specific cause. Furthermore,
726 processes linked to the potential for population adaptation and interactions with other
727 environmental stressors can add a further level of complexity as compared to laboratory
728 studies. Improvements could be made by increasing site coverage and further targeted work
729 on molecular mechanisms, as well as data on the background levels and variations in
730 methylation changes.

731 From the studies presented here, it can be concluded that DNA methylation might be the key
732 to transferring the response to ionising radiation from one generation to the next. Whereas
733 measuring total DNA methylation can be performed without any prior information on genetic
734 background of the species, the rapid technical evolution and the decreasing cost of sequencing
735 analyses will offer a wider comparison of radiologically induced DNA methylation in different
736 biological models and provide greater insight into the underlying mechanisms. An important
737 step will be to compare the sensitivity, reliance and above all specificity of DNA methylation
738 as a possible biomarker of ionising radiation exposure at environmentally relevant levels, with
739 other epigenetic mechanisms such as histone modifications and microRNAs linked to
740 responses at higher level biological complexity e.g. changes in growth and reproduction.

Table 1. Pros and cons of DNA methylation methods. 5-mC (methylcytosine), 5-hmC (hydroxymethylcytosine), AFLP-MS (methylation specific amplification fragment length polymorphism), HPLC-MS/MS (high performance liquid chromatography coupled with tandem mass spectrometry), ELISA assay (enzyme-linked immunosorbent assay), MeDIP seq (methylated DNA immunoprecipitation coupled with next-generation sequencing), WGBS (whole genome bisulfite sequencing), RRBS (reduced representation bisulfite sequencing)

Method	Principle	Methylated base detected	Pros	Cons
AFLP-MS	Cut DNA with restriction enzymes and analyse on a fragment analyser	5-mC	Low cost per sample No need for sequenced genome Low DNA amount (250-500 ng) Low processing time	Detection of global methylation Specific equipment needed
HPLC-MS/MS	Detection of methyl groups on hydrolysed DNA sample	5-mC & 5-hmC	Medium cost per sample No need for sequenced genome Low processing time	Detection of global methylation High DNA amount (50-1000 ng) Specific equipment needed
5 mC ELISA assay	Use of monoclonal antibodies sensitive and specific for 5-mC	5-mC	Low cost per sample No need for sequenced genome No specific equipment needed Low processing time	Detection of global methylation High DNA amount (100-2000 ng)
MeDIP seq	Immunoprecipitation sequencing	5-mC	Detection of site specific methylation Low DNA amount (300 ng)	High cost per sample Need for sequenced genome Specific equipment needed High processing time
WGBS	Bisulfite conversion and DNA sequencing	5-mC & 5-hmC (oxBS-seq)	Detection of site specific methylation Low DNA amount (30 ng)	High cost per sample Need for sequenced genome Specific equipment needed High processing time
RRBS	Bisulfite conversion and DNA sequencing	5-mC & 5-hmC (oxBS-seq)	Detection of site specific methylation	High cost per sample Need for sequenced genome High DNA amount (1000 ng) Specific equipment needed High processing time

Table 2: Overview of lab-based studies in which ecotoxicological relevant model organisms were exposed to radiation, radioisotopes or other toxins for multiple generations; F0=Parental organism, F...= offspring with the number indicating the generation

Species	Chemical	Generations	Observed phenotype	Ref
<i>C. elegans</i>	Gamma radiation 7-42 mGy/h	F0-F2	Greater reproduction effects in multigenerationally and transgenerationally exposed F2s than F0 generation	Buisset-Goussen et al. (2014)
<i>D. magna</i>	Gamma radiation 0.007-35 mGy/h	F0-F2	Toxicity on multiple traits increased from F0 to F2	Parisot et al. (2015)
<i>D. Rerio</i>	Gamma radiation 9-53 mGy/h	F0-F1	Effect on DNA damage, transcription, lipid peroxidation and demographic endpoints in F1	Hurem et al. (2017), Hurem et al. (2018b), (2018a)
<i>D. Rerio</i>	Uranium 20-250 µg/L	F0-F1	Effect on DNA damage, transcription, DNA methylation and demographic endpoints in F1	Bourrachot et al. (2014), Gombeau et al. (2017)
<i>D. magna</i>	Americium 0.3-15 mGy/h	F0-F2	Threshold for effects on reproduction reduced from 1.5 mGyh ⁻¹ in F0 generation to 0.3 mGyh ⁻¹ in F2 and F3	Alonzo et al. (2008)
<i>D. magna</i>	Uranium 2-50 µg/L	F0-F1	Greater reduction in fecundity in F1 than F0 at 50 µg/L	Plaire et al. (2013)
<i>D. magna</i>	Nickel 42-85 µg/L	F0-F1	Greater reduction of ATP levels in F1 compared to F0	Pane et al. (2004)
<i>C. elegans</i>	Ag nanoparticles EC30-value	F0-F10	Greater (10 fold) sensitivity in F2, F5, F8 and F10 generations compared to P generation	Schultz et al. (2016)
<i>D. magna</i>	Ag nanoparticles EC10-EC50	F0-F10	Population growth rate at 10 µg/L reduced by 80% in F2s compared to 21% in F0 generation	Volker et al. (2013)
<i>D. magna</i>	Penta-chlorophenol 0.0002-2 µmol/L	F0-F3	Population growth rate reduction increases from 28.2% to 34.9% to 46.3% in F0, F1, F2 generations	Chen et al. (2014)
<i>D. magna</i>	Tetracycline 0.1-5 mg/L	F0-F1	NOEC decreased from 5 mg/L to 0.1 mg/L from F0 to F3	Kim et al. (2012)
<i>D. magna</i>	Enrofloxacin 13 mg/L	F0-F1	Reproduction NOEC decreased from 30 mg/L to 3.1 mg/L from F0 to F1 generation	Bona et al. (2015)
<i>C. elegans</i>	Uranium 4-50 µg/L	F0-F16	Greater maximal length but increased sensitivity to uranium across the generations	Goussen et al. (2015)
<i>C. elegans</i>	Uranium 4.6 µg/L	F0-F22	Increase of sensitivity from F0 to F6 and subsequent adaptation until F22	Dutilleul et al. (2014)

Tabel 3: Overview of studies in which changes in epigenetic mechanisms (DNA methylation, histone modifications or miRNA's) are measured in organisms exposed to radiation in a long-term set-up (within or over generations) either in laboratory or field conditions. F0=Parental organism, F...= offspring with the number indicating the generation, CEZ: Chernobyl Exclusion Zone, FEZ: Fukushima Exclusion Zone

		Organism	Experimental conditions	Epigenetic changes	Additional endpoints	Reference
Laboratory exposed	Plants	<i>A. thaliana</i>	F1, F2, multigenerational (F0 from CEZ, 1.8-4.4µGy/h) methyl methane sulfonate (140 µM) or Rose Bengal (10 µM)	DNA methylation: hypermethylation in both F1 and F2	Higher resistance to mutagens, increased expression of ROS scavenging enzymes and DNA repair enzymes	Kovalchuk et al. (2004)
		<i>P. sylvestris</i>	F0, trans- and multigenerational set up, on contaminated soil both acute (~10Gy) and chronic (~80Gy) (F0 from CEZ, (absorbed dose 1986: >60Gy, 10-60Gy, 1-10, 0.1-10Gy),	DNA methylation: hypermethylation in exposed	-	Kovalchuk et al. (2003)
		<i>A. thaliana</i>	F1, F2 transgenerational, Progeny of plants collected at CEZ 1.8-4.4µGy/h	DNA methylation: hypermethylation	-	Kovalchuk et al. (2004)
		<i>A. thaliana</i>	F0, F1, F2, mutligenerational, 14 day exposure during vegetative state, 22, 38, 86, 457 mGy/h	DNA methylation: dose-dependent hypermethylation, strongest in F2	Changes in ROS-scavenging enzymes, DNA repair and developmental traits, mutants in methyltransferases showed increased sensitivity to radiation	van de Walle et al. (2016), Saenen et al. (2017)
	Invertebrates	<i>D. magna</i>	F0, F1, F2 and F3 transgenerational, F0 exposed for 25 days, 6,5 µGy/h or 41.3 mGy/h	DNA methylation: hypomethylation but dose-rate independent	Reduction in fecundity in F0, no adverse effects in F1, F2, F3	Trijau et al. (2018)
	Vertebrates	<i>D. rerio</i>	F0, F1, F2, F3, transgenerational, exposure during gametogenesis, 8.7 mGy/h, 28 days	DNA methylation: Genome-wide in F1,	Linked to gene pathways changes and adverse effects in progeny	Hurem et al. (2018b), Kamstra et al. (2018),

				locus-specific regions up to F3		Hurem et al. (2017)
		<i>D. rerio</i>	F0, F1, multigenerational, exposure during gametogenesis, 8.7 mGy/h, 28 days	miRNA expression in F1 embryos	-	Martin et al., in prep
		<i>D. rerio</i>	F0, F1, F2 transgenerational, gametogenesis, 8.7 mGy/h, 28 days	Histone modifications (hypermethylation) at specific loci in F0 and F1 but no longer in F2	-	Lindeman et al. (2019)
		<i>S. salar</i>	F0-embryo's, exposure from one-cell fertilized eggs till early gastrula stage, 1, 10, 20 or 30 mGy/h	Histone modification (hypermethylation) at specific loci at highest dose rate	-	Lindeman et al. (2019)
Field collected	Plants	<i>P. sylvestris</i>	F0, (Belarus, Chernobyl affected area), annual absorbed dose: 10-158 mGy or 1-14 µGy/h	DNA methylation: transient with dose, hypermethylation	-	Volkova et al. (2018)
		<i>C. bursa pastoris</i>	F0, FEZ : total dose rates: 0.13-38 µGy/h	DNA methylation : no change	-	Horemans et al. (2018)
		<i>A. thaliana</i>	F0, CEZ : total dose rates : 0.1-160 µGy/h	DNA methylation : Hypomethylation at highest dose rates	-	Horemans et al. (2018)
		<i>G. max</i>	F0, after 7 generations CEZ, total accumulated dose : 1-132 mGy	DNA methylation: slight increase (10%) in radio-	Increased levels of single and double DNA strand breaks	Georgieva et al. (2017)

				contaminated samples		
Invertebrates	Earthworms (<i>A. caliginosa</i> , <i>O. lacteum</i>)	F0, CEZ, total dose rates 0.12-41 $\mu\text{Gy/h}$	DNA methylation: site-specific differences <i>A. caliginosa</i> . for no or limited changes found for <i>O. lacteum</i>	-		Saenen et al. (2017)
Vertebrates	<i>H. arborea</i>	F0, FEZ, total dose rate 0.38-41,7 $\mu\text{Gy/h}$	DNA-methylation: hypermethylation, dose-dependent	Concomitant with increased DNA damage		Saenen et al. (2017)

Acknowledgements

This work was supported by the EC's COMET (Coordination and implementation of a pan-European instrument for radioecology) program, Fission-2012-3.4.1-604794.

JK was supported by the Research Council of Norway through its Centers of Excellence funding scheme, project number 223268/F50.

References

- Abramov VI, Fedorenko OM, Shevchenko VA (1992) GENETIC CONSEQUENCES OF RADIOACTIVE CONTAMINATION FOR POPULATIONS OF ARABIDOPSIS. *Sci Total Environ* 112 (1):19-28
- Adam-Guillermin C, Horemans N, Oughton D, Spurgeon D, Stark K, Gaschak S, Yoschenko V, Hertel-Aas T, Pereira S, Vandenhove H (2013) Deliverable D4.1, Initial Research Activity, State-of-the art on epigenetics and general approach. COMET program, Fission-2012-3.4.1-604794.
- Allis CD, Jenuwein T (2016) The molecular hallmarks of epigenetic control. *Nature Reviews Genetics* 17 (8):487-500. doi:10.1038/nrg.2016.59
- Allis CD, Jenuwein T, Reinberg D (2007) *Epigenetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA
- Alonzo F, Gilbin R, Zeman FA, Garnier-Laplace J (2008a) Increased effects of internal alpha irradiation in *Daphnia magna* after chronic exposure over three successive generations. *Aquat Toxicol* 87 (3):146-156. doi:10.1016/j.aquatox.2008.01.015
- Alonzo F, Hertel-Aas T, Gilek M, Gilbin R, Oughton DH, Garnier-Laplace J (2008b) Modelling the propagation of effects of chronic exposure to ionising radiation from individuals to populations. *Journal of Environmental Radioactivity* 99 (9):1464-1473. doi:10.1016/j.jenvrad.2007.11.021
- Aluru N (2017) Epigenetic effects of environmental chemicals: insights from zebrafish. *Curr Opin Toxicol* 6:26-33. doi:10.1016/j.cotox.2017.07.004
- An YR, Kim SJ, Oh MJ, Kim HM, Shim IS, Kim PJ, Choi K, Hwang SY (2013) Analysis of microRNA and gene expression profiling in triazole fungicide-treated HepG2 cell line. *Toxicology* 303 (1):94-98. doi:10.1016/j.tox.2012.11.004
- Annacondia ML, Mageroy MH, Martinez G (2018) Stress response regulation by epigenetic mechanisms: changing of the guards. *Physiol Plant* 162 (2):239-250. doi:10.1111/ppl.12662
- Arkhipov NP, Kuchma ND, Askbrant S, Pasternak PS, Musica VV (1994) Acute and long-term effects of irradiation on Pine (*Pinus silvestris*) stands post-Chernobyl. *Sci Total Environ* 157 (1-3):383-386
- Asselman J, De Coninck DIM, Pfrender ME, De Schampelaere KAC (2016) Gene body methylation patterns in *Daphnia* are associated with gene family size. *Genome Biology and Evolution* 8 (4):1185-1196. doi:10.1093/gbe/evw069
- Baker RJ, Bickham AM, Bondarkov M, Gaschak SP, Matson CW, Rodgers BE, Wickliffe JK, Chesser RK (2001) Consequences of polluted environments on population structure: The bank vole (*Clethrionomys glareolus*) at Chernobyl. *Ecotoxicology* 10 (4):211-216. doi:10.1023/a:1016665226716

- Baker TR, Peterson RE, Heideman W (2014) Using Zebrafish as a Model System for Studying the Transgenerational Effects of Dioxin. *Toxicol Sci* 138 (2):403-411. doi:10.1093/toxsci/kfu006
- Bannister AJ, Kouzarides T (2011) Regulation of chromatin by histone modifications. *Cell Res* 21 (3):381-395. doi:10.1038/cr.2011.22
- Battle JVI (2016) Impact of the Fukushima Accident on Marine Biota, Five Years Later. *Integrated Environmental Assessment and Management* 12 (4):654-658. doi:10.1002/ieam.1825
- Baubec T, Colombo DF, Wirbelauer C, Schmidt J, Burger L, Krebs AR, Akalin A, Schubeler D (2015) Genomic profiling of DNA methyltransferases reveals a role for DNMT3B in genic methylation. *Nature* 520 (7546):243-U278. doi:10.1038/nature14176
- Beresford NA, Copplestone D (2011) Effects of ionizing radiation on wildlife: what knowledge have we gained between the Chernobyl and Fukushima accidents? *Integr Environ Assess Manag* 7 (3):371-373. doi:10.1002/ieam.238
- Beresford NA, Fesenko S, Konoplev A, Skuterud L, Smith JT, Voigt G (2016) Thirty years after the Chernobyl accident: What lessons have we learnt? *Journal of Environmental Radioactivity* 157:77-89. doi:10.1016/j.jenvrad.2016.02.003
- Berger SL, Kouzarides T, Shiekhatar R, Shilatifard A (2009) An operational definition of epigenetics. *Genes & Development* 23 (7):781-783. doi:10.1101/gad.1787609
- Berr A, Shafiq S, Shen WH (2011) Histone modifications in transcriptional activation during plant development. *Biochimica Et Biophysica Acta-Genes Regulatory Mechanisms* 1809 (10):567-576. doi:10.1016/j.bbagr.2011.07.001
- Bible JM, Sanford E (2016) Local adaptation in an estuarine foundation species: Implications for restoration. *Biol Conserv* 193:95-102. doi:10.1016/j.biocon.2015.11.015
- Bird A (2007) Perceptions of epigenetics. *Nature* 447 (7143):396-398. doi:10.1038/nature05913
- Biron PA, Massarin S, Alonzo F, Garcia-Sanchez L, Charles S, Billoir E (2012) Population-Level Modeling to Account for Multigenerational Effects of Uranium in *Daphnia magna*. *Environ Sci Technol* 46 (2):1136-1143. doi:10.1021/es202658b
- Bock C (2012) Analysing and interpreting DNA methylation data. *Nature Reviews Genetics* 13 (10):705-719. doi:10.1038/nrg3273
- Bona MD, Zounkova R, Merlanti R, Blaha L, De Liguoro M (2015) Effects of enrofloxacin, ciprofloxacin, and trimethoprim on two generations of *Daphnia magna*. *Ecotox Environ Safe* 113:152-158. doi:10.1016/j.ecoenv.2014.11.018
- Bossdorf O, Richards CL, Pigliucci M (2008) Epigenetics for ecologists. *Ecology Letters* 11 (2):106-115. doi:10.1111/j.1461-0248.2007.01130.x
- Boubriak II, Grodzinsky DM, Polischuk VP, Naumenko VD, Gushcha NP, Micheev AN, McCready SJ, Osborne DJ (2008) Adaptation and impairment of DNA repair function in pollen of *Betula verrucosa* and seeds of *Oenothera biennis* from differently radionuclide-contaminated sites of Chernobyl. *Ann Bot* 101 (2):267-276. doi:10.1093/aob/mcm276
- Boubriak I, Akimkina T, Polischuk V, Dmitriev A, McCready S, Grodzinsky D (2016) Long Term Effects of Chernobyl Contamination on DNA Repair Function and Plant Resistance to Different Biotic and Abiotic Stress Factors. *Cytol Genet* 50 (6):381-399. doi:10.3103/s0095452716060049
- Bourrachot S, Brion F, Pereira S, Floriani M, Camilleri V, Cavalie I, Palluel O, Adam-Guillermin C (2014) Effects of depleted uranium on the reproductive success and F1 generation survival of zebrafish (*Danio rerio*). *Aquat Toxicol* 154:1-11. doi:10.1016/j.aquatox.2014.04.018
- Bowman JD, Choudhury M (2016) Phthalates in neonatal health: friend or foe? *Journal of Developmental Origins of Health and Disease* 7 (6):652-664. doi:10.1017/s2040174416000349
- Brautigam K, Vining KJ, Lafon-Placette C, Fossdal CG, Mirouze M, Marcos JG, Fluch S, Fraga MF, Guevara MA, Abarca D, Johnsen O, Maury S, Strauss SH, Campbell MM, Rohde A,

- Diaz-Sala C, Cervera MT (2013) Epigenetic regulation of adaptive responses of forest tree species to the environment. *Ecol Evol* 3 (2):399-415. doi:10.1002/ece3.461
- Bruce TJA, Matthes MC, Napier JA, Pickett JA (2007) Stressful "memories" of plants: Evidence and possible mechanisms. *Plant Sci* 173 (6):603-608. doi:10.1016/j.plantsci.2007.09.002
- Buisset-Goussen A, Goussen B, Della-Vedova C, Galas S, Adam-Guillermin C, Lecomte-Pradines C (2014) Effects of chronic gamma irradiation: a multigenerational study using *Caenorhabditis elegans*. *Journal of Environmental Radioactivity* 137:190-197. doi:10.1016/j.jenvrad.2014.07.014
- Burdak-Rothkamm S, Rothkamm K (2018) Radiation-induced bystander and systemic effects serve as a unifying model system for genotoxic stress responses. *Mutation Research-Reviews in Mutation Research* 778:13-22. doi:10.1016/j.mrrev.2018.08.001
- Burgio E, Piscitelli P, Migliore L (2018) Ionizing Radiation and Human Health: Reviewing Models of Exposure and Mechanisms of Cellular Damage. An Epigenetic Perspective. *Int J Environ Res Public Health* 15 (9):13. doi:10.3390/ijerph15091971
- Calarco JP, Borges F, Donoghue MTA, Van Ex F, Jullien PE, Lopes T, Gardner R, Berger F, Feijo JA, Becker JD, Martienssen RA (2012) Reprogramming of DNA Methylation in Pollen Guides Epigenetic Inheritance via Small RNA. *Cell* 151 (1):194-205. doi:10.1016/j.cell.2012.09.001
- Castel SE, Martienssen RA (2013) RNA interference in the nucleus: roles for small RNAs in transcription, epigenetics and beyond. *Nature Reviews Genetics* 14 (2):100-112. doi:10.1038/nrg3355
- Chan SWL, Henderson IR, Jacobsen SE (2005) Gardening the genome: DNA methylation in *Arabidopsis thaliana*. *Nature Reviews Genetics* 6 (5):351-360. doi:10.1038/nrg1601
- Chen Y, Huang J, Xing LQ, Liu HL, Giesy J, Yu HX, Zhang XW (2014) Effects of multigenerational exposures of *D. magna* to environmentally relevant concentrations of pentachlorophenol. *Environ Sci Pollut Res* 21 (1):234-243. doi:10.1007/s11356-013-1692-z
- Chen YC, Baram TZ (2016) Toward Understanding How Early-Life Stress Reprograms Cognitive and Emotional Brain Networks. *Neuropsychopharmacology* 41 (1):197-206. doi:10.1038/npp.2015.181
- Cingolani P, Cao XY, Khetani RS, Chen CC, Coon M, Sammak A, Bollig-Fischer A, Land S, Huang Y, Hudson ME, Garfinkel MD, Zhong S, Robinson GE, Ruden DM (2013) Intronic Non-CG DNA hydroxymethylation and alternative mRNA splicing in honey bees. *Bmc Genomics* 14. doi:10.1186/1471-2164-14-666
- Colicchio JM, Kelly JK, Hileman LC (2018) Parental experience modifies the *Mimulus* methylome. *BMC Genomics* 19:15. doi:10.1186/s12864-018-5087-x
- Coors A, Vanoverbeke J, De Bie T, De Meester L (2009) Land use, genetic diversity and toxicant tolerance in natural populations of *Daphnia magna*. *Aquatic Toxicology* 95 (1):71-79. doi:10.1016/j.aquatox.2009.08.004
- Costa D, Marien J, Janssens TKS, van Gestel CAM, Driessen G, Sousa JP, van Straalen NM, Roelofs D (2012) Influence of adaptive evolution of cadmium tolerance on neutral and functional genetic variation in *Orchesella cincta*. *Ecotoxicology* 21 (7):2078-2087. doi:10.1007/s10646-012-0961-9
- Coulter JB, O'Driscoll CM, Bressler JP (2013) Hydroquinone increases 5-Hydroxymethylcytosine formation through ten eleven translocation 1 (TET1) 5-methylcytosine dioxygenase. *Journal of Biological Chemistry* 288 (40):28792-28800. doi:10.1074/jbc.M113.491365
- Dutilleul M, Bonzom JM, Lecomte C, Goussen B, Daian F, Galas S, Reale D (2014) Rapid evolutionary responses of life history traits to different experimentally-induced pollutions in *Caenorhabditis elegans*. *BMC Evol Biol* 14:14. doi:10.1186/s12862-014-0252-6
- Ellegren H, Lindgren G, Primmer CR, Moller AP (1997) Fitness loss and germline mutations in barn swallows breeding in Chernobyl. *Nature* 389 (6651):593-596. doi:10.1038/39303

- Feil R, Fraga MF (2012) Epigenetics and the environment: emerging patterns and implications. *Nature Reviews Genetics* 13 (2):97-109. doi:10.1038/nrg3142
- Flores K, Wolschin F, Corneveaux JJ, Allen AN, Huentelman MJ, Amdam GV (2012) Genome-wide association between DNA methylation and alternative splicing in an invertebrate. *BMC Genomics* 13:9. doi:10.1186/1471-2164-13-480
- Frost PC, Ebert D, Larson JH, Marcus MA, Wagner ND, Zalewski A (2010) Transgenerational effects of poor elemental food quality on *Daphnia magna*. *Oecologia* 162 (4):865-872. doi:10.1007/s00442-009-1517-4
- Fujita Y, Yoshihara Y, Sato I, Sato S (2014) Environmental radioactivity damages the DNA of earthworms of Fukushima Prefecture, Japan. *Eur J Wildl Res* 60 (1):145-148. doi:10.1007/s10344-013-0767-y
- Galvan I, Bonisoli-Alquati A, Jenkinson S, Ghanem G, Wakamatsu K, Mousseau TA, Moller AP (2014) Chronic exposure to low-dose radiation at Chernobyl favours adaptation to oxidative stress in birds. *Functional Ecology* 28 (6):1387-1403. doi:10.1111/1365-2435.12283
- Gardiner-Garden M, Frommer M (1987) CPG islands in vertebrate genomes. *J Mol Biol* 196 (2):261-282. doi:10.1016/0022-2836(87)90689-9
- Garnier-Laplace J, Geras'kin S, Della-Vedova C, Beaugelin-Seiller K, Hinton TG, Real A, Oudalova A (2013) Are radiosensitivity data derived from natural field conditions consistent with data from controlled exposures? A case study of Chernobyl wildlife chronically exposed to low dose rates. *J Environ Radioact* 121:12-21. doi:10.1016/j.jenvrad.2012.01.013
- Georgieva M, Rashydov NM, Hajdich M (2017) DNA damage, repair monitoring and epigenetic DNA methylation changes in seedlings of Chernobyl soybeans. *DNA Repair* 50:14-21. doi:<http://dx.doi.org/10.1016/j.dnarep.2016.12.002>
- Geras'kin S, Evseeva T, Oudalova A (2013) Effects of long-term chronic exposure to radionuclides in plant populations. *Journal of Environmental Radioactivity* 121:22-32. doi:10.1016/j.jenvrad.2012.03.007
- Geras'kin S, Oudalova A, Dikareva N, Spiridonov S, Hinton T, Chernonog E, Garnier-Laplace J (2011) Effects of radioactive contamination on Scots pines in the remote period after the Chernobyl accident. *Ecotoxicology* 20 (6):1195-1208. doi:10.1007/s10646-011-0664-7
- Geras'kin SA, Fesenko SV, Alexakhin RM (2008) Effects of non-human species irradiation after the Chernobyl NPP accident. *Environ Int* 34 (6):880-897
- Gielen H, Remans T, Vangronsveld J, Cuypers A (2012) MicroRNAs in Metal Stress: Specific Roles or Secondary Responses? *Int J Mol Sci* 13 (12):15826-15847. doi:10.3390/ijms131215826
- Gombeau K, Bourdineaud J-P, Ravanat J-L, Armant O, Camilleri V, Cavalie I, Floriani M, Adam-Guillermin C (2017) Epigenetic, histopathological and transcriptomic effects following exposure to depleted uranium in adult zebrafish and their progeny. *Aquat Toxicol* 184:14-25. doi:<http://dx.doi.org/10.1016/j.aquatox.2016.12.004>
- Gombeau K, Pereira S, Ravanat JL, Camilleri V, Cavalie I, Bourdineaud JP, Adam-Guillermin C (2016) Depleted uranium induces sex- and tissue-specific methylation patterns in adult zebrafish. *Journal of Environmental Radioactivity* 154:25-33. doi:10.1016/j.jenvrad.2016.01.004
- Gonzalez APR, Chrtek J, Dobrev PI, Dumalasova V, Fehrer J, Mraz P, Latzel V (2016) Stress-induced memory alters growth of clonal off spring of white clover (*Trifolium repens*). *Am J Bot* 103 (9):1567-1574. doi:10.3732/ajb.1500526
- Goussen B, Pery ARR, Bonzom JM, Beaudouin R (2015) Transgenerational Adaptation to Pollution Changes Energy Allocation in Populations of Nematodes. *Environ Sci Technol* 49 (20):12500-12508. doi:10.1021/acs.est.5b03405

- Grativol C, Hemerly AS, Gomes Ferreira PC (2012) Genetic and epigenetic regulation of stress responses in natural plant populations. *Biochimica Et Biophysica Acta-Gene Regulatory Mechanisms* 1819 (2):176-185. doi:10.1016/j.bbagr.2011.08.010
- Greer EL, Blanco MA, Gu L, Sendinc E, Liu J, Aristizabal-Corrales D, Hsu C-H, Aravind L, He C, Shi Y (2015) *DNA Methylation on N-6-Adenine in C. elegans*. *Cell* 161 (4):868-878. doi:10.1016/j.cell.2015.04.005
- Greer EL, Maures TJ, Ucar D, Hauswirth AG, Mancini E, Lim JP, Benayoun BA, Shi Y, Brunet A (2011) Transgenerational epigenetic inheritance of longevity in *Caenorhabditis elegans*. *Nature* 479 (7373):365-371. doi:<http://www.nature.com/nature/journal/v479/n7373/abs/nature10572.html#supplementary-information>
- Groenendijk D, Kraak MHS, Admiraal W (1999) Efficient shedding of accumulated metals during metamorphosis in metal-adapted populations of the midge *Chironomus riparius*. *Environmental Toxicology and Chemistry* 18 (6):1225-1231. doi:10.1897/1551-5028(1999)018<1225:esoamd>2.3.co;2
- Groh KJ, Carvalho RN, Chipman JK, Denslow ND, Halder M, Murphy CA, Roelofs D, Rolaki A, Schirmer K, Watanabe KH (2015) Development and application of the adverse outcome pathway framework for understanding and predicting chronic toxicity: I. Challenges and research needs in ecotoxicology. *Chemosphere* 120:764-777. doi:10.1016/j.chemosphere.2014.09.068
- Groot MP, Kooke R, Knobens N, Vergeer P, Keurentjes JJB, Ouborg NJ, Verhoeven KJF (2016) Effects of Multi-Generational Stress Exposure and Offspring Environment on the Expression and Persistence of Transgenerational Effects in *Arabidopsis thaliana*. *Plos One* 11 (3):16. doi:10.1371/journal.pone.0151566
- Guerrero-Bosagna C, Covert TR, Haque MM, Settles M, Nilsson EE, Anway MD, Skinner MK (2012) Epigenetic transgenerational inheritance of vinclozolin induced mouse adult onset disease and associated sperm epigenome biomarkers. *Reprod Toxicol* 34 (4):694-707. doi:10.1016/j.reprotox.2012.09.005
- Guerrero-Bosagna C, Jensen P (2015) Globalization, climate change, and transgenerational epigenetic inheritance: will our descendants be at risk? *Clin Epigenetics* 7:3. doi:10.1186/s13148-014-0043-3
- Hanson MA, Skinner MK (2016) Developmental origins of epigenetic transgenerational inheritance. *Environmental Epigenetics* 2 (1):dvw002-dvw002. doi:10.1093/eep/dvw002
- He YH, Li ZC (2018) Epigenetic Environmental Memories in Plants: Establishment, Maintenance, and Reprogramming. *Trends Genet* 34 (11):856-866. doi:10.1016/j.tig.2018.07.006
- Hertel-Aas T, Brunborg G, Jaworska A, Salbu B, Oughton DH (2011) Effects of different gamma exposure regimes on reproduction in the earthworm *Eisenia fetida* (Oligochaeta). *Sci Total Environ* 412:138-147. doi:10.1016/j.scitotenv.2011.09.037
- Hinton TG, Alexakhin R, Balonov M, Gentner N, Hendry J, Prister B, Strand P, Woodhead D (2007) Radiation-induced effects on plants and animals: Findings of the united nations Chernobyl forum. *Health Phys* 93 (5):427-440
- Hirayama T, Shinozaki K (2010) Research on plant abiotic stress responses in the post-genome era: past, present and future. *Plant J* 61 (6):1041-1052. doi:10.1111/j.1365-313X.2010.04124.x
- Hiyama A, Nohara C, Kinjo S, Taira W, Gima S, Tanahara A, Otaki JM (2012) The biological impacts of the Fukushima nuclear accident on the pale grass blue butterfly. *Sci Rep* 2:570. doi:10.1038/srep00570
- Hiyama A, Nohara C, Taira W, Kinjo S, Iwata M, Otaki JM (2013) The Fukushima nuclear accident and the pale grass blue butterfly: evaluating biological effects of long-term low-dose exposures. *BMC Evol Biol* 13:25. doi:10.1186/1471-2148-13-168

- Hochmuth JD, De Meester L, Pereira CMS, Janssen CR, De Schampelaere KAC (2015) Rapid adaptation of a *Daphnia magna* population to metal stress is associated with heterozygote excess. *Environmental Science & Technology* 49 (15):9298-9307. doi:10.1021/acs.est.5b00724
- Holoch D, Moazed D (2015) RNA-mediated epigenetic regulation of gene expression. *Nat Rev Genet* 16 (2):71-84. doi:10.1038/nrg3863
- Horemans N, Nauts R, Vives i Batlle J, Van Hees M, Jacobs G, Voorspoels S, Gaschak S, Nanba K, Saenen E (2018) Genome-wide DNA methylation changes in two Brassicaceae species sampled alongside a radiation gradient in Chernobyl and Fukushima. *Journal of Environmental Radioactivity* 192:405-416. doi:<https://doi.org/10.1016/j.jenvrad.2018.07.012>
- Hu CW, Chen JL, Hsu YW, Yen CC, Chao MR (2015) Trace analysis of methylated and hydroxymethylated cytosines in DNA by isotope-dilution LC-MS/MS: first evidence of DNA methylation in *Caenorhabditis elegans*. *Biochem J* 465:39-47. doi:10.1042/bj20140844
- Huang C, Sloan EA, Boerkoel CF (2003) Chromatin remodeling and human disease. *Current Opinion in Genetics & Development* 13 (3):246-252. doi:10.1016/s0959-437x(03)00054-6
- Huang J, Yang ML, Zhang XM (2016) The function of small RNAs in plant biotic stress response. *J Integr Plant Biol* 58 (4):312-327. doi:10.1111/jipb.12463
- Huertas D, Sendra R, Munoz P (2009) Chromatin dynamics coupled to DNA repair. *Epigenetics* 4 (1):31-42. doi:10.4161/epi.4.1.7733
- Hunt CR, Ramnarain D, Horikoshi N, Iyengar P, Pandita RK, Shay JW, Pandita TK (2013) Histone Modifications and DNA Double-Strand Break Repair after Exposure to Ionizing Radiations. *Radiat Res* 179 (4):383-392. doi:10.1667/rr3308.2
- Hurem S, Gomes T, Brede DA, Lindbo Hansen E, Mutoloki S, Fernandez C, Mothersill C, Salbu B, Kassaye YA, Olsen AK, Oughton D, Alestrom P, Lyche JL (2017) Parental gamma irradiation induces reprotoxic effects accompanied by genomic instability in zebrafish (*Danio rerio*) embryos. *Environ Res* 159:564-578. doi:10.1016/j.envres.2017.07.053
- Hurem S, Gomes T, Brede DA, Mayer I, Lobert VH, Mutoloki S, Gutzkow KB, Teien HC, Oughton D, Alestrom P, Lyche JL (2018a) Gamma irradiation during gametogenesis in young adult zebrafish causes persistent genotoxicity and adverse reproductive effects. *Ecotoxicol Environ Saf* 154:19-26. doi:10.1016/j.ecoenv.2018.02.031
- Hurem S, Martin LM, Lindeman L, Brede DA, Salbu B, Lyche JL, Alestrom P, Kamstra JH (2018b) Parental exposure to gamma radiation causes progressively altered transcriptomes linked to adverse effects in zebrafish offspring. *Environ Pollut* 234:855-863. doi:10.1016/j.envpol.2017.12.023
- IAEA (1992) Effects of Ionizing Radiation on Plants and Animals at Levels Implied by Current Radiation Protection Standards. INTERNATIONAL ATOMIC ENERGY AGENCY, Vienna
- IAEA (2006) Environmental Consequences of the Chernobyl Accident and their Remediation: Twenty Years of Experience. Radiological Assessment Reports Series, vol 8. INTERNATIONAL ATOMIC ENERGY AGENCY, Vienna
- IAEA (2015) The Fukushima Daiichi Accident. INTERNATIONAL ATOMIC ENERGY AGENCY, Vienna
- Iglesias FM, Cerdan PD (2016) Maintaining Epigenetic Inheritance During DNA Replication in Plants. *Front Plant Sci* 7:8. doi:10.3389/fpls.2016.00038
- Iorio MV, Piovan C, Croce CM (2010) Interplay between microRNAs and the epigenetic machinery: An intricate network. *Biochimica Et Biophysica Acta- Gene Regulatory Mechanisms* 1799 (10-12):694-701. doi:10.1016/j.bbagr.2010.05.005
- Irato P, Santovito G, Cassini A, Piccinni E, Albergoni V (2003) Metal accumulation and binding protein induction in *Mytilus galloprovincialis*, *Scapharca inaequalvis*, and *Tapes*

- philippinarum from the lagoon of Venice. Archives of Environmental Contamination and Toxicology 44 (4):476-484
- Jackson D, Copplestone D, Stone DM (2004) Effects of chronic radiation exposure on small mammals in the Chernobyl exclusion zone. Nucl Energy-J Br Nucl Energy Soc 43 (5):281-287. doi:10.1680/nuen.43.5.281.53137
- Jansen M, Coors A, Vanoverbeke J, Schepens M, De Voogt P, De Schamphelaere KAC, De Meester L (2015) Experimental evolution reveals high insecticide tolerance in *Daphnia* inhabiting farmland ponds. Evolutionary Applications 8 (5):442-453. doi:10.1111/eva.12253
- Jimenez-Chillaron JC, Nijland MJ, Ascensao AA, Sardao VA, Magalhaes J, Hitchler MJ, Domann FE, Oliveira PJ (2015) Back to the future: transgenerational transmission of xenobiotic-induced epigenetic remodeling. Epigenetics 10 (4):259-273. doi:10.1080/15592294.2015.1020267
- Jung I, Kim D (2012) Histone modification profiles characterize function-specific gene regulation. Journal of Theoretical Biology 310:132-142. doi:10.1016/j.jtbi.2012.06.009
- Kadhim MA, Moore SR, Goodwin EH (2004) Interrelationships amongst radiation-induced genomic instability, bystander effects, and the adaptive response. Mutat Res-Fundam Mol Mech Mutagen 568 (1):21-32. doi:10.1016/j.mrfmmm.2004.06.043
- Kamstra JH, Hurem S, Martin L, Lindeman LC, Legler J, Oughton D, Salbu B, Brede DA, Lyche JL, Alestrom P (2018) Ionizing radiation induces transgenerational effects of DNA methylation in zebrafish. Sci Rep 8:13. doi:10.1038/s41598-018-33817-w
- Katz DJ, Edwards TM, Reinke V, Kelly WG (2009) A *C. elegans* LSD1 demethylase contributes to germline immortality by reprogramming epigenetic memory. Cell 137 (2):308-320. doi:<http://dx.doi.org/10.1016/j.cell.2009.02.015>
- Kaup S, Grandjean V, Mukherjee R, Kapoor A, Keyes E, Seymour CB, Mothersill CE, Schofield PN (2006) Radiation-induced genomic instability is associated with DNA methylation changes in cultured human keratinocytes. Mutat Res-Fundam Mol Mech Mutagen 597 (1-2):87-97. doi:10.1016/j.mrfmmm.2005.06.032
- Kille P, Andre J, Anderson C, Ang HN, Bruford MW, Bundy JG, Donnelly R, Hodson ME, Juma G, Lahive E, Morgan AJ, Sturzenbaum SR, Spurgeon DJ (2013) DNA sequence variation and methylation in an arsenic tolerant earthworm population. Soil Biology & Biochemistry 57:524-532. doi:10.1016/j.soilbio.2012.10.014
- Kim HY, Lee MJ, Yu SH, Kim SD (2012a) The individual and population effects of tetracycline on *Daphnia magna* in multigenerational exposure. Ecotoxicology 21 (4):993-1002. doi:10.1007/s10646-012-0853-z
- Kim JM, To TK, Seki M (2012b) An Epigenetic Integrator: New Insights into Genome Regulation, Environmental Stress Responses and Developmental Controls by HISTONE DEACETYLASE 6. Plant Cell Physiol 53 (5):794-800. doi:10.1093/pcp/pcs004
- Klosin A, Casas E, Hidalgo-Carcedo C, Vavouri T, Lehner B (2017) Transgenerational transmission of environmental information in *C-elegans*. Science 356 (6335):316-319. doi:10.1126/science.aah6412
- Kovalchuk I, Abramov V, Pogribny I, Kovalchuk O (2004) Molecular aspects of plant adaptation to life in the Chernobyl zone. Plant Physiol 135 (1):357-363. doi:10.1104/pp.104.040477
- Kovalchuk O, Burke P, Arkhipov A, Kuchma N, James SJ, Kovalchuk I, Pogribny I (2003) Genome hypermethylation in *Pinus silvestris* of Chernobyl - a mechanism for radiation adaptation? Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis 529 (1&2):13-20
- Krivolutsky DA (1996) Soil fauna as bioindicator of radioactive pollution, vol 10. Bioindicator Systems for Soil Pollution. Kluwer Academic Publ, Dordrecht

- Kryshev II, G. ST, Beresford NA (2005) Effects on wildlife. In: Smith J, Beresford NA (eds) Chernobyl Catastrophe and Consequences. Praxis Publisher/Springer, Chichester, UK, pp 267-287
- Kubota Y, Tsuji H, Kawagoshi T, Shiomi N, Takahashi H, Watanabe Y, Fuma S, Doi K, Kawaguchi I, Aoki M, Kubota M, Furuhashi Y, Shigemura Y, Mizoguchi M, Yamada F, Tomozawa M, Sakamoto SH, Yoshida S (2015) Chromosomal Aberrations in Wild Mice Captured in Areas Differentially Contaminated by the Fukushima Dai-ichi Nuclear Power Plant Accident. *Environ Sci Technol* 49 (16):10074-10083. doi:10.1021/acs.est.5b01554
- Kuchma O, Vornam B, Finkeldey R (2011) Mutation rates in Scots pine (*Pinus sylvestris* L.) from the Chernobyl exclusion zone evaluated with amplified fragment-length polymorphisms (AFLPs) and microsatellite markers. *Mutat Res Genet Toxicol Environ Mutagen* 725 (1-2):29-35. doi:10.1016/j.mrgentox.2011.07.003
- Kure EH, Sabo M, Stangeland AM, Hamfjord J, Hytterod S, Heggenes J, Lydersen E (2013) Molecular responses to toxicological stressors: Profiling microRNAs in wild Atlantic salmon (*Salmo salar*) exposed to acidic aluminum-rich water. *Aquatic Toxicology* 138-139:98-104. doi:10.1016/j.aquatox.2013.04.004
- Langdon CJ, Pearce TG, Meharg AA, Semple KT (2003) Inherited resistance to arsenate toxicity in two populations of *Lumbricus rubellus*. *Environmental Toxicology and Chemistry* 22 (10):2344-2348
- Larriba E, del Mazo J (2016) Role of Non-Coding RNAs in the Transgenerational Epigenetic Transmission of the Effects of Reprotoxicants. *Int J Mol Sci* 17 (4):13. doi:10.3390/ijms17040452
- Law JA, Jacobsen SE (2010) Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nature Reviews Genetics* 11 (3):204-220. doi:10.1038/nrg2719
- Levinton JS, Suatoni E, Wallace W, Junkins R, Kelaher B, Allen BJ (2003) Rapid loss of genetically based resistance to metals after the cleanup of a Superfund site. *Proceedings of the National Academy of Sciences of the United States of America* 100 (17):9889-9891. doi:10.1073/pnas.1731446100
- Li ZG, Chen X, Li SP, Wang ZC (2015) Effect of nickel chloride on *Arabidopsis* genomic DNA and methylation of 18S rDNA. *Electronic Journal of Biotechnology* 18 (1):51-57. doi:10.1016/j.ejbt.2014.12.004
- Lim JP, Brunet A (2013) Bridging the transgenerational gap with epigenetic memory. *Trends Genet* 29 (3):176-186. doi:10.1016/j.tig.2012.12.008
- Lindeman L, Kamstra JH, Ballangby J, Hurem S, Martín L, Brede DA, Teien HC, Oughton D, Salbu B, Alestrom P (2019) Gamma radiation induces locus specific changes to histone modification enrichment in zebrafish and Atlantic salmon. *PLoS ONE*:Accepted for publication
- Liu XL, Luo F, Ling M, Lu L, Shi L, Lu XL, Xu H, Chen C, Yang QL, Xue JC, Li J, Zhang AH, Liu QZ (2016) MicroRNA-21 activation of ERK signaling via PTEN is involved in arsenite-induced autophagy in human hepatic L-02 cells. *Toxicology Letters* 252:1-10. doi:10.1016/j.toxlet.2016.04.015
- Loayza-Muro RA, de Baat ML, Palomino EJ, Kuperus P, Kraak MHS, Admiraal W, Breeuwer JAJ (2014) Metals and altitude drive genetic diversity of chironomids in Andean streams. *Freshwater Biology* 59 (1):56-63. doi:10.1111/fwb.12245
- Lourenco J, Mendo S, Pereira R (2016) Radioactively contaminated areas: Bioindicator species and biomarkers of effect in an early warning scheme for a preliminary risk assessment. *J Hazard Mater* 317:503-542. doi:10.1016/j.jhazmat.2016.06.020
- Lyko F (2018) The DNA methyltransferase family: a versatile toolkit for epigenetic regulation. *Nature Reviews Genetics* 19 (2):81-92. doi:10.1038/nrg.2017.80

- Marczylo EL, Jacobs MN, Gant TW (2016) Environmentally induced epigenetic toxicity: potential public health concerns. *Critical Reviews in Toxicology* 46 (8):676-700. doi:10.1080/10408444.2016.1175417
- Margueron R, Trojer P, Reinberg D (2005) The key to development: interpreting the histone code? *Curr Opin Genet Dev* 15 (2):163-176. doi:10.1016/j.gde.2005.01.005
- Marsh AG, Pasqualone AA (2014) DNA methylation and temperature stress in an Antarctic polychaete, *Spiophanes tcherniai*. *Frontiers in Physiology* 5:9. doi:10.3389/fphys.2014.00173
- Matson CW, Rodgers BE, Chesser RK, Baker RJ (2000) Genetic diversity of *Clethrionomys glareolus* populations from highly contaminated sites in the Chernobyl Region, Ukraine. *Environ Toxicol Chem* 19 (8):2130-2135. doi:10.1897/1551-5028(2000)019<2130:gdocgp>2.3.co;2
- Mattick JS, Amaral PP, Dinger ME, Mercer TR, Mehler MF (2009) RNA regulation of epigenetic processes. *Bioessays* 31 (1):51-59. doi:10.1002/bies.080099
- McCarrey JR (2012) The epigenome as a target for heritable environmental disruptions of cellular function. *Mol Cell Endocrinol* 354 (1-2):9-15. doi:10.1016/j.mce.2011.09.014
- Meehan RR, Thomson JP, Lentini A, Nestor CE, Pennings S (2018) DNA methylation as a genomic marker of exposure to chemical and environmental agents. *Curr Opin Chem Biol* 45:48-56. doi:10.1016/j.cbpa.2018.02.006
- Meeks HN, Chesser RK, Rodgers BE, Gaschak S, Baker RJ (2009) UNDERSTANDING THE GENETIC CONSEQUENCES OF ENVIRONMENTAL TOXICANT EXPOSURE: CHERNOBYL AS A MODEL SYSTEM. *Environ Toxicol Chem* 28 (9):1982-1994
- Meeks HN, Wickliffe JK, Hooper SR, Chesser RK, Rodgers BE, Baker RJ (2007) Mitochondrial control region variation in bank voles (*Clethrionomys glareolus*) is not related to Chernobyl radiation exposure. *Environ Toxicol Chem* 26 (2):361-369. doi:10.1897/06-346r.1
- Mendez-Acuna L, Di Tomaso MV, Palitti F, Martinez-Lopez W (2010) Histone post-translational modifications in DNA damage response. *Cytogenet Genome Res* 128 (1-3):28-36. doi:10.1159/000296275
- Meng XZ, Zheng TS, Chen X, Wang JB, Zhang WH, Pan SH, Jiang HC, Liu LX (2011) microRNA expression alteration after arsenic trioxide treatment in HepG-2 cells. *Journal of Gastroenterology and Hepatology* 26 (1):186-193. doi:10.1111/j.1440-1746.2010.06317.x
- Mirbahai L, Chipman JK (2014) Epigenetic memory of environmental organisms: A reflection of lifetime stressor exposures. *Mutat Res Genet Toxicol Environ Mutagen* 764:10-17. doi:10.1016/j.mrgentox.2013.10.003
- Molinier J, Ries G, Zipfel C, Hohn B (2006) Transgenerational memory of stress in plants. *Nature* 442 (7106):1046-1049. doi:10.1038/nature05022
- Mondal S, Go YS, Lee SS, Chung BY, Kim JH (2016) Characterization of histone modifications associated with DNA damage repair genes upon exposure to gamma rays in *Arabidopsis* seedlings. *J Radiat Res* 57 (6):646-654. doi:10.1093/jrr/rrw077
- Moon J, Kwak JI, Kim SW, An YJ (2017) Multigenerational effects of gold nanoparticles in *Caenorhabditis elegans*: Continuous versus intermittent exposures. *Environmental Pollution* 220:46-52. doi:10.1016/j.envpol.2016.09.021
- Morgan WF (2002) Genomic Instability and Bystander Effects: A Paradigm Shift in Radiation Biology? *Military Medicine* 167 (suppl_1):44-45. doi:10.1093/milmed/167.suppl_1.44 %J Military Medicine
- Neeb ZT, Nowacki M (2018) RNA-mediated transgenerational inheritance in ciliates and plants. *Chromosoma* 127 (1):19-27. doi:10.1007/s00412-017-0655-4
- Norouzitallab P, Baruah K, Vanrompay D, Bossier P (2019) Can epigenetics translate environmental cues into phenotypes? *Sci Total Environ* 647:1281-1293. doi:10.1016/j.scitotenv.2018.08.063

- Nota B, de Korte M, Ylstra B, van Straalen NM, Roelofs D (2013) Genetic variation in parthenogenetic Collembolans is associated with differences in fitness and cadmium-induced transcriptome responses. *Environmental Science & Technology* 47 (2):1155-1162. doi:10.1021/es303983z
- Pane EF, McGeer JC, Wood CM (2004) Effects of chronic waterborne nickel exposure on two successive generations of *Daphnia magna*. *Environ Toxicol Chem* 23 (4):1051-1056. doi:10.1897/03-208
- Parisot F, Bourdineaud JP, Plaire D, Adam-Guillermin C, Alonzo F (2015) DNA alterations and effects on growth and reproduction in *Daphnia magna* during chronic exposure to gamma radiation over three successive generations. *Aquat Toxicol* 163:27-36. doi:10.1016/j.aquatox.2015.03.002
- Pastor WA, Pape UJ, Huang Y, Henderson HR, Lister R, Ko M, McLoughlin EM, Brudno Y, Mahapatra S, Kapranov P, Tahiliani M, Daley GQ, Liu XS, Ecker JR, Milos PM, Agarwal S, Rao A (2011) Genome-wide mapping of 5-hydroxymethylcytosine in embryonic stem cells. *Nature* 473 (7347):394-397. doi:10.1038/nature10102
- Paun O, Verhoeven KJF, Richards CL (2019) Opportunities and limitations of reduced representation bisulfite sequencing in plant ecological epigenomics. *New Phytol* 221 (2):738-742. doi:10.1111/nph.15388
- Pecinka A, Mittelsten Scheid O (2012) Stress-induced chromatin changes: a critical view on their heritability. *Plant & cell physiology* 53 (5):801-808. doi:10.1093/pcp/pcs044
- Pernot E, Hall J, Baatout S, Benotmane MA, Blanchardon E, Bouffler S, El Saghire H, Gomolka M, Guertler A, Harms-Ringdahl M, Jeggo P, Kreuzer M, Laurier D, Lindholm C, Mkacher R, Quintens R, Rothkamm K, Sabatier L, Tapio S, de Vathaire F, Cardis E (2012) Ionizing radiation biomarkers for potential use in epidemiological studies. *Mutation Research-Reviews in Mutation Research* 751 (2):258-286. doi:10.1016/j.mrrev.2012.05.003
- Plaire D, Bourdineaud JP, Alonzo A, Camilleri V, Garcia-Sanchez L, Adam-Guillermin C, Alonzo F (2013) Transmission of DNA damage and increasing reprotoxic effects over two generations of *Daphnia magna* exposed to uranium. *Comp Biochem Physiol C-Toxicol Pharmacol* 158 (4):231-243. doi:10.1016/j.cbpc.2013.09.001
- Potters G, Pasternak TP, Guisez Y, Palme KJ, Jansen MAK (2007) Stress-induced morphogenic responses: growing out of trouble? *Trends Plant Sci* 12 (3):98-105. doi:10.1016/j.tplants.2007.01.004
- Pouget JP, Georgakilas AG, Ravanat JL (2018) Targeted and Off-Target (Bystander and Abscopal) Effects of Radiation Therapy: Redox Mechanisms and Risk/Benefit Analysis. *Antioxid Redox Signal* 29 (15):1447-1487. doi:10.1089/ars.2017.7267
- Pray LA (2004) Epigenetics: Genome, meet your environment. *Scientist* 18 (13):14-+
- Putiri EL, Robertson KD (2011) Epigenetic mechanisms and genome stability. *Clin Epigenetics* 2:299-314. doi:10.1007/s13148-010-0017-z
- Qi XZ, Yang X, Chen SY, He XY, Dweep H, Guo MZ, Cheng WH, Xu WT, Luo YB, Gretz N, Dai Q, Huang KL (2014) Ochratoxin A induced early hepatotoxicity: new mechanistic insights from microRNA, mRNA and proteomic profiling studies. *Scientific Reports* 4:14. doi:10.1038/srep05163
- Rapp RA, Wendel JF (2005) Epigenetics and plant evolution. *New Phytologist* 168 (1):81-91. doi:10.1111/j.1469-8137.2005.01491.x
- Rasmann S, De Vos M, Casteel CL, Tian DL, Halitschke R, Sun JY, Agrawal AA, Felton GW, Jander G (2012) Herbivory in the Previous Generation Primes Plants for Enhanced Insect Resistance. *Plant Physiol* 158 (2):854-863. doi:10.1104/pp.111.187831
- Raut VV, Sainis JK (2012) ⁶⁰Co-gamma radiation induces differential acetylation and phosphorylation of histones H3 and H4 in wheat. *Plant Biol* 14 (1):110-117. doi:10.1111/j.1438-8677.2011.00463.x

- Rechavi O, Hourri-Ze'evi L, Anava S, Goh Wee Siong S, Kerk Sze Y, Hannon Gregory J, Hobert O (2014) Starvation-induced transgenerational inheritance of small RNAs in *C. elegans*. *Cell* 158 (2):277-287. doi:<http://dx.doi.org/10.1016/j.cell.2014.06.020>
- Regev A, Lamb MJ, Jablonka E (1998) The role of DNA methylation in invertebrates: Developmental regulation or genome defense? *Molecular Biology & Evolution* 15 (7):880-891
- Reid NM, Proestou DA, Clark BW, Warren WC, Colbourne JK, Shaw JR, Karchner SI, Hahn ME, Nacci D, Oleksiak MF, Crawford DL, Whitehead A (2016) The genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild fish. *Science* 354:1305-1308
- Saenen E, Lecomte C, Bradshaw C, Spurgeon D, Oughton D, Lapied E, Bonzom JM, Beaugelin K, Kamstra JH, Orizaola G, Armant O, Gaschak S, Nanba K, Horemans N (2017) Deliverable D-4.3, Initial Research Activity on transgenerational effects and role of epigenetics: Results and Impact. COMET program, Fission-2012-3.4.1-604794.
- Sahu PP, Pandey G, Sharma N, Puranik S, Muthamilarasan M, Prasad M (2013) Epigenetic mechanisms of plant stress responses and adaptation. *Plant Cell Reports* 32 (8):1151-1159. doi:10.1007/s00299-013-1462-x
- Santos AP, Serra T, Figueiredo DD, Barros P, Lourenco T, Chander S, Oliveira MM, Saibo NJM (2011) Transcription Regulation of Abiotic Stress Responses in Rice: A Combined Action of Transcription Factors and Epigenetic Mechanisms. *Omics* 15 (12):839-857. doi:10.1089/omi.2011.0095
- Saze H (2012) Transgenerational inheritance of induced changes in the epigenetic state of chromatin in plants. *Genes Genet Syst* 87 (3):145-152. doi:10.1266/ggs.87.145
- Schofield PN, Kondratowicz M (2018) Evolving paradigms for the biological response to low dose ionizing radiation; the role of epigenetics. *Int J Radiat Biol* 94 (8):769-781. doi:10.1080/09553002.2017.1388548
- Schott D, Yanai I, Hunter CP (2014) Natural RNA interference directs a heritable response to the environment. *Scientific Reports* 4:10. doi:10.1038/srep07387
- Schultz C, Unrine J, Tsyusko O, Svendsen C, D.J. S (2016) The transgenerational toxicity of Ag ions, nanoparticles and sulphadised nanoparticles across 10 *Caenorhabditis elegans* generations. *Proceedings of the Royal Society B-Biological Sciences* 283:20152911
- Schwindt AR, Winkelman DL, Keteles K, Murphy M, Vajda AM (2014) An environmental oestrogen disrupts fish population dynamics through direct and transgenerational effects on survival and fecundity. *J Appl Ecol* 51 (3):582-591. doi:10.1111/1365-2664.12237
- Scourzic L, Mouly E, Bernard OA (2015) TET proteins and the control of cytosine demethylation in cancer. *Genome Medicine* 7. doi:10.1186/s13073-015-0134-6
- Seong KH, Li D, Shimizu H, Nakamura R, Ishii S (2011) Inheritance of Stress-Induced, ATF-2-Dependent Epigenetic Change. *Cell* 145 (7):1049-1061. doi:10.1016/j.cell.2011.05.029
- Shaw JR, Hampton TH, King BL, Whitehead A, Galvez F, Gross RH, Keith N, Notch E, Jung D, Glaholt SP, Chen CY, Colbourne JK, Stanton BA (2014) Natural selection canalizes expression variation of environmentally induced plasticity-enabling genes. *Molecular Biology and Evolution* 31 (11):3002-3015. doi:10.1093/molbev/msu241
- Singer AC, Shaw H, Rhodes V, Hart A (2016) Review of antimicrobial resistance in the environment and its relevance to environmental regulators. *Frontiers in Microbiology* 7. doi:10.3389/fmicb.2016.01728
- Skinner MK (2016) Epigenetic transgenerational inheritance. *Nat Rev Endocrinol* 12 (2):68-70. doi:10.1038/nrendo.2015.206
- Skinner MK, Guerrero-Bosagna C (2009) Environmental signals and transgenerational epigenetics. *Epigenomics* 1 (1):111-117. doi:10.2217/epi.09.11
- Song C-X, Szulwach KE, Fu Y, Dai Q, Yi C, Li X, Li Y, Chen C-H, Zhang W, Jian X, Wang J, Zhang L, Looney TJ, Zhang B, Godley LA, Hicks LM, Lahn BT, Jin P, He C (2011) Selective chemical labeling reveals the genome-wide distribution of 5-

hydroxymethylcytosine. Nat Biotech 29 (1):68-72. doi:<http://www.nature.com/nbt/journal/v29/n1/abs/nbt.1732.html#supplementary-information>

- Song M-K, Song M, Choi H-S, Ryu J-C (2012) Benzo k fluoranthene-Induced Changes in miRNA-mRNA Interactions in Human Hepatocytes. *Toxicology and Environmental Health Sciences* 4 (3):143-153. doi:10.1007/s13530-012-0129-2
- Stassen JHM, Lopez A, Jain R, Pascual-Pardo D, Luna E, Smith LM, Ton J (2018) The relationship between transgenerational acquired resistance and global DNA methylation in Arabidopsis. *Sci Rep* 8:13. doi:10.1038/s41598-018-32448-5
- Steinhauser G, Brandl A, Johnson TE (2014) Comparison of the Chernobyl and Fukushima nuclear accidents: A review of the environmental impacts. *Sci Total Environ* 470:800-817. doi:10.1016/j.scitotenv.2013.10.029
- Stern S, Snir O, Mizrahi E, Galili M, Zaltsman I, Soen Y (2014) Reduction in maternal Polycomb levels contributes to transgenerational inheritance of a response to toxic stress in flies. *Journal of Physiology-London* 592 (11):2343-2355. doi:10.1113/jphysiol.2014.271445
- Strand P, Aono T, Brown JE, Garnier-Laplace J, Hosseini A, Sazykina T, Steenhuisen F, Battle JVI (2014) Assessment of Fukushima-Derived Radiation Doses and Effects on Wildlife in Japan. *Environ Sci Technol Lett* 1 (3):198-203. doi:10.1021/ez500019j
- Sudan J, Raina M, Singh R (2018) Plant epigenetic mechanisms: role in abiotic stress and their generational heritability. *3 Biotech* 8:12. doi:10.1007/s13205-018-1202-6
- Sun BY, Shi YF, Yang XZ, Zhao T, Duan JC, Sun ZW (2018) DNA methylation: A critical epigenetic mechanism underlying the detrimental effects of airborne particulate matter. *Ecotox Environ Safe* 161:173-183. doi:10.1016/j.ecoenv.2018.05.083
- Szyf M (2007) The dynamic epigenome and its implications in toxicology. *Toxicological Sciences* 100 (1):7-23. doi:10.1093/toxsci/kfm177
- Theodorakis CW, Bickham JW, Lamb T, Medica PA, Lyne TB (2001) Integration of genotoxicity and population genetic analyses in kangaroo rats (*Dipodomys merriami*) exposed to radionuclide contamination at the Nevada Test Site, USA. *Environ Toxicol Chem* 20 (2):317-326. doi:10.1897/1551-5028(2001)020<0317:iogapg>2.0.co;2
- Theodorakis CW, Shugart LR (1997) Genetic ecotoxicology .2. population genetic structure in mosquitofish exposed in situ to radionuclides. *Ecotoxicology* 6 (6):335-354. doi:10.1023/a:1018695231565
- Trijau M, Asselman J, Armant O, Adam-Guillermin C, De Schamphelaere KAC, Alonzo F (2018) Transgenerational DNA Methylation Changes in *Daphnia magna* Exposed to Chronic γ Irradiation. *Environ Sci Technol* 52 (7):4331-4339. doi:10.1021/acs.est.7b05695
- Uller T, English S, Pen I (2015) When is incomplete epigenetic resetting in germ cells favoured by natural selection? *Proc R Soc B-Biol Sci* 282 (1811):8. doi:10.1098/rspb.2015.0682
- van de Walle J, Horemans N, Saenen E, Van Hees M, Wannijn J, Nauts R, van Gompel A, Vangronsveld J, Vandenhove H, Cuypers A (2016) Arabidopsis plants exposed to gamma radiation in two successive generations show a different oxidative stress response. *Journal of Environmental Radioactivity* 165:270-279. doi:10.1016/j.jenvrad.2016.10.014
- Van Hoeck A, Horemans N, Nauts R, Van Hees M, Vandenhove H, Blust R (2017) Lemna minor plants chronically exposed to ionising radiation: RNA-seq analysis indicates a dose rate dependent shift from acclimation to survival strategies. *Plant Sci* 257:84-95. doi:<http://dx.doi.org/10.1016/j.plantsci.2017.01.010>
- Van Straalen NM, Roelofs D (2007) *An Introduction to Ecological Genomics*. Oxford University Press, Oxford, UK
- Vandegheuchte MB, Janssen CR (2011) Epigenetics and its implications for ecotoxicology. *Ecotoxicology* 20 (3):607-624
- Vandegheuchte MB, Janssen CR (2014) Epigenetics in an ecotoxicological context. *Mutat Res Genet Toxicol Environ Mutagen* 764:36-45. doi:10.1016/j.mrgentox.2013.08.008

- Vasseur P, Cossu-Leguille C (2006) Linking molecular interactions to consequent effects of persistent organic pollutants (POPs) upon populations. *Chemosphere* 62 (7):1033-1042. doi:10.1016/j.chemosphere.2005.05.043
- Verhoeven KJF, Jansen JJ, van Dijk PJ, Biere A (2010) Stress-induced DNA methylation changes and their heritability in asexual dandelions. *New Phytol* 185 (4):1108-1118. doi:10.1111/j.1469-8137.2009.03121.x
- Verhoeven KJF, van Gorp TP (2012) Transgenerational Effects of Stress Exposure on Offspring Phenotypes in Apomictic Dandelion. *Plos One* 7 (6):8. doi:10.1371/journal.pone.0038605
- Verhoeven KJF, Vonholdt BM, Sork VL (2016) Epigenetics in ecology and evolution: what we know and what we need to know. *Mol Ecol* 25 (8):1631-1638. doi:10.1111/mec.13617
- Volker C, Boedicker C, Daubenthaler J, Oetken M, Oehlmann J (2013) Comparative Toxicity Assessment of Nanosilver on Three Daphnia Species in Acute, Chronic and Multi-Generation Experiments. *Plos One* 8 (10):10. doi:10.1371/journal.pone.0075026
- Volkova PY, Geras'kin SA, Horemans N, Makarenko ES, Saenen E, Duarte GT, Nauts R, Bondarenko VS, Jacobs G, Voorspoels S, Kudin M (2018) Chronic radiation exposure as an ecological factor: Hypermethylation and genetic differentiation in irradiated Scots pine populations. *Environ Pollut* 232:105-112. doi:10.1016/j.envpol.2017.08.123
- Waddington CH (1939) *An Introduction to Modern Genetics*. George Allen & Unwin., London
- Wang L, Bammler TK, Beyer RP, Gallagher EP (2013) Copper-Induced deregulation of microRNA expression in the Zebrafish olfactory system. *Environmental Science & Technology* 47 (13):7466-7474. doi:10.1021/es400615q
- Wang Y, Liu H, Sun Z (2017) Lamarck rises from his grave: parental environment-induced epigenetic inheritance in model organisms and humans. *Biol Rev Camb Philos Soc*. doi:10.1111/brv.12322
- Whittle CA, Otto SP, Johnston MO, Krochko JE (2009) Adaptive epigenetic memory of ancestral temperature regime in *Arabidopsis thaliana*. *Botany* 87 (6):650-657. doi:10.1139/b09-030
- Wigle DT, Arbuckle TE, Walker M, Wade MG, Liu SL, Krewski D (2007) Environmental hazards: Evidence for effects on child health. *Journal of Toxicology and Environmental Health-Part B-Critical Reviews* 10 (1-2):3-39. doi:10.1080/10937400601034563
- Wirgin I, Roy NK, Loftus M, Chambers RC, Franks DG, Hahn ME (2011) Mechanistic basis of resistance to PCBs in Atlantic Tomcod from the Hudson River. *Science* 331 (6022):1322-1325. doi:10.1126/science.1197296
- Wu CT, Morris JR (2001) Genes, genetics, and epigenetics: A correspondence. *Science* 293 (5532):1103-1105. doi:10.1126/science.293.5532.1103
- Wu H, Zhang Y (2014) Reversing DNA Methylation: Mechanisms, Genomics, and Biological Functions. *Cell* 156 (1-2):45-68. doi:10.1016/j.cell.2013.12.019
- Xu W, Wang T, Xu SY, Xu SX, Wu LJ, Wu YJ, Bian P (2015) Radiation-Induced Epigenetic Bystander Effects Demonstrated in *Arabidopsis Thaliana*. *Radiat Res* 183 (5):511-524. doi:10.1667/rr13909.1
- Xu Y, Wu F, Tan L, Kong L, Xiong L, Deng J, Barbera AJ, Zheng L, Zhang H, Huang S, Min J, Nicholson T, Chen T, Xu G, Shi Y, Zhang K, Shi Yujiang G (2011) Genome-wide Regulation of 5hmC, 5mC, and Gene Expression by Tet1 Hydroxylase in Mouse Embryonic Stem Cells. *Molecular Cell* 42 (4):451-464. doi:<http://dx.doi.org/10.1016/j.molcel.2011.04.005>
- Yoschenko V, Nanba K, Yoshida S, Watanabe Y, Takase T, Sato N, Keitoku K (2016) Morphological abnormalities in Japanese red pine (*Pinus densiflora*) at the territories contaminated as a result of the accident at Fukushima Dai-Ichi Nuclear Power Plant. *Journal of Environmental Radioactivity* 165:60-67. doi:10.1016/j.jenvrad.2016.09.006
- Zaka R, Chenal C, Misset MT (2004) Effects of low doses of short-term gamma irradiation on growth and development through two generations of *Pisum sativum*. *Sci Total Environ* 320 (2-3):121-129. doi:10.1016/j.scitotenv.2003.08.010

- Zelena L, Sorochinsky B, von Arnold S, van Zyl L, Clapham DH (2005) Indications of limited altered gene expression in *Pinus sylvestris* trees from the Chernobyl region. *Journal of Environmental Radioactivity* 84 (3):363-373. doi:10.1016/j.jenvrad.2005.03.008
- Ziv-Gal A, Wang W, Zhou C, Flaws JA (2015) The effects of in utero bisphenol A exposure on reproductive capacity in several generations of mice. *Toxicology and Applied Pharmacology* 284 (3):354-362. doi:10.1016/j.taap.2015.03.003