



HAL
open science

New Perspectives in the Treatment of Tumor Cells by Electromagnetic Radiation at Resonance Frequencies in Cellular Membrane Channels

Emanuele Calabrò, Salvatore Magazù

► **To cite this version:**

Emanuele Calabrò, Salvatore Magazù. New Perspectives in the Treatment of Tumor Cells by Electromagnetic Radiation at Resonance Frequencies in Cellular Membrane Channels. *The Open Biotechnology Journal*, 2019, 13 (1), pp.105-110. 10.2174/187407070190130105 . hal-02538544

HAL Id: hal-02538544

<https://hal.science/hal-02538544>

Submitted on 9 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The Open Biotechnology Journal

Content list available at: <https://openbiotechnologyjournal.com>



RESEARCH ARTICLE

New Perspectives in the Treatment of Tumor Cells by Electromagnetic Radiation at Resonance Frequencies in Cellular Membrane Channels

Emanuele Calabrò^{1,5,*} and Salvatore Magazù^{1,2,3,4,5}

¹Department of Mathematical and Informatics Sciences, Physical Sciences and Earth Sciences of Messina University, Viale Ferdinando Stagno D'Alcontres 31, 98166 Messina, Italy

²Le Studium, Loire Valley Institute for Advanced Studies, Orléans & Tours, France

³Centre de Biophysique Moléculaire (CBM)-CNRS UPR 4301 du CNRS, rue Charles Sadron, 45071 Orleans CEDEX 2 France; Laboratoire Interfaces, Confinement, Matériaux et Nanostructures (ICMN) - UMR 7374 CNRS - Université d'Orléans, 1b rue de la Férollerie, CS 40059, 45071 Orléans cedex 2, France

⁴Istituto Nazionale di Alta Matematica "F. Severi" – INDAM - Gruppo Nazionale per la Fisica Matematica – GNFM, Piazzale Aldo Moro, 500185 Rome, Italy

⁵CISFA "Interuniversity Consortium of Applied Physical Sciences" (Consorzio Interuniversitario di Scienze Fisiche Applicate), Viale Ferdinando Stagno D'Alcontres 31, 98166 Messina, Italy

Abstract:

Background:

The use of electromagnetic fields has been considered as adjuvant therapy for the treatment of cancer given that some clinical trials have shown that the irradiation of cancer cells with electromagnetic fields can slow down the disease progression.

Aims:

We hypothesize that this effect could be amplified by irradiating tumor cells with electromagnetic fields having frequencies close to the natural resonant frequencies of membrane channels in tumor cells, in order to obtain a significant change of the ion flux across tumor cell membrane channels, inducing the largest harmful alteration in their cellular function.

Methods:

Neuronal-like cells were used as a cell model and exposed for 6 h to electromagnetic fields at different frequencies (0, 50 Hz, 900 MHz) at the same intensity of 2 mT. The exposure system was represented by two Helmholtz coils driven by a power amplifier in current mode and an arbitrary function generator. FTIR spectroscopy was used to evaluate the results of the exposure.

Results:

The results of this study showed that the Amide I vibration band increased in intensity with the increase of the frequency, leading us to assume that the displacement of the cell channels α -helices depends on the frequency of the applied electromagnetic fields.

Conclusion:

This preliminary result leads us to plan future research aimed at searching for the natural frequencies of membrane channels in tumor cells using resonant electromagnetic fields in order to damage the cellular functions of tumor cells. Clinical trials are needed to confirm such a hypothesis derived from this physical study.

Keywords: Electromagnetic fields, Tumor cells, Irradiation, Protein α -helices, Resonance Frequency, Exposure system.

Article History

Received: April 30, 2019

Revised: July 02, 2019

Accepted: July 08, 2019

1. INTRODUCTION

One of the most serious diseases that has developed in the modern era is certainly cancer.

At this moment the methods of cancer treatment usually used are surgery, radiosurgery, radiotherapy, chemotherapy and immunotherapy. Unfortunately, the use of these techniques is not always effective mainly due to side effects, because even

normal cells are affected by such therapies, inducing the oncologists to limit the same therapy that was chosen [1 - 7].

In recent years, the use of electromagnetic fields has also been considered as adjuvant therapy for the treatment of cancer. In this regard, some results have shown that the irradiation of cancer cells with electromagnetic fields has slowed the disease progression without inducing significant side effects [8 - 14].

The harmful effects of electromagnetic fields on cancer cells can be easily explained by the results of previous studies showing that protein unfolding and aggregation occur under exposure to electromagnetic fields [15 - 22] so that the irradiation of cancerous cells with an electromagnetic field should alter their cellular functions, causing them to die.

In order to amplify this effect, it is plausible to think of irradiating tumor cells with an electromagnetic field having frequencies equal to (or at least close to) the natural frequencies of the tumor cells that are to be treated, in order to obtain the largest alteration in their cellular function.

This hypothesis derives from the resonance theory, according to which, displacements of molecules are amplified at natural resonant frequencies of the same material.

However, the problem is to search for the natural resonance frequencies of tumor cells. For this aim, some mechanical models have been provided up to now. For instance, natural resonant frequencies of cells were assumed to depend on some mechanical properties (the dimension of the cell, density and elastic modulus), so that the natural resonant frequencies of normal cells are different from the natural resonant frequencies of cancer cells [13]. Particularly, a model was proposed in recent studies, derived from civil engineering principles, for computing the natural resonance frequency of cancer cells in order that an applied external electromagnetic field can interact with the cancer cell at that frequency inducing a resonance, damaging the cancer cells [23].

Another study proposed that measuring Young's modulus and cell shape as a function of the frequency of a mechanical vibration can be used for computing the natural frequency of spreading osteoblast cells [24].

Nevertheless, the models proposed in the studies mentioned above are based on the mechanical properties of cells, so that the possible resonant interactions strictly depend on mechanical properties that cannot be considered as the most important characteristics related to cellular functions. In contrast, a physical-chemical parameter which has an important role in cell viability is represented by ion flux in cellular membrane channels. Indeed, it was shown that ion flux across cell channels has a fundamental role in cellular functions as it can regulate the initiation, progression and proliferation of cancer [25 - 29].

Hence, resonance interaction mechanisms between an applied electromagnetic field and membrane channels in cancer cells can be hypothesized, in order to significantly alter the ion

flux across cell membrane channels, damaging cancerous cells, as described in the following sections.

2. MATERIALS AND METHODS

Protein α -helices are present in all types of cellular membrane channels [30 - 32]. In particular, cellular membrane protein accounts for about 50% of its mass [33]. Taking advantage of this circumstance, a preliminary study was carried out studying protein response to electromagnetic fields at different frequencies by means of Fourier Transform Infrared (FTIR) spectroscopy.

Neuroblastoma cell line SH-SY5Y was purchased from American Type Culture Collections (ATCC) (Rockville, MD) and was used as a cell model in this study. Cells were differentiated following the protocol accurately described in the previous studies [16, 17].

The exposure system was represented by two Helmholtz coils, used to produce a uniform magnetic field at the center of the distance between the coils, following Helmholtz coils theory. The Helmholtz coils were driven by a 7570 AE power amplifier in current mode (Techron, Elkhart, IN) and a Model 75 arbitrary function generator (Wavetek, Plainview, NY), that were used to generate a Direct Current (DC) and an Alternating Current (AC) at 50 Hz frequency. Moreover, microwave sources at frequencies ranging up to 900 MHz were obtained using mobile phone devices. Further technical characteristics of the experimental setup used in this study have already been described [34, 35].

However, contrary to the protocol used in previous studies by the authors, different cell samples were exposed for 6 h to electromagnetic fields at different frequencies (0, 50 Hz, 900 MHz), but the same intensity of 2 mT was used for the three types of electromagnetic fields. This protocol was chosen in order to compare the effects of electromagnetic fields at different frequencies.

Neuronal-like SH-SY5Y cells, grown in 25 cm² culture flasks, were located in the center of the field generated between the coils. The coils and cell samples were placed into an incubator (series 5400-115V models, Thermo Electron, Winchester, VA) with 5% CO₂/95% humidity and at the temperature of (37.0 ± 0.1) °C. Unexposed samples were located in another incubator, at the same physical conditions reported above.

Cell cultures were treated following the protocol of [35, 36] and subjected to infrared (IR) spectroscopic techniques by a FTIR spectrometer Vertex 80v of Bruker Optics. For each spectrum, 128 interferograms were collected with a spectral resolution of 4 cm⁻¹, following the technical procedure described previously [16, 17].

3. RESULTS

Representative FTIR spectra of neuronal-like cells samples after 6 h exposure to static, 50 Hz and 900 MHz electromagnetic fields at 2 mT are reported in Fig. (1A - 1C), respectively. The most evident vibration band appearing in these spectra is represented by an intense Amide I band centered

* Address correspondence to this author Department of Mathematical and Informatics Sciences, University of Messina, Physical Sciences and Earth Sciences of Messina University; Tel: +390906765025, +390906765019; E-mail: e.calabro@yahoo.com

around 1650 cm^{-1} , corresponding to the protein α -helix structure, that is due to C=O stretching and N—H bending

vibrations and a weak Amide II vibration centered at 1545 cm^{-1} , that is due to the N—H bending and C—N stretching vibrations.




Fig. (1). Amide I band in neuronal-like cells after 6 h exposure to a static magnetic field (A), 50 Hz (B) and 900 MHz (C) electromagnetic fields. Spectra of exposed and unexposed samples are represented in the colors red and black, respectively. The increase in the intensity of the Amide I band with an increase of the frequency of the applied electromagnetic field after exposure can be observed, which can be explained assuming that α -helices in cells membrane channels had aligned towards the direction of the applied field.

A significant increase in the intensity of the Amide I band was observed after exposures to 2 mT electromagnetic fields at 0, 50 Hz, 900 MHz, with $p < 0.05$ significance. This result was already explained assuming that protein α -helices had aligned with the applied electromagnetic fields, showing that the exposure of proteins in bidistilled water solution to an applied electromagnetic field causes a torque which induces the alignment of protein α -helices towards the direction of the applied field [34, 37- 40]. As a result, an enlargement in the diameter of cellular membrane channel should occur, inducing a decrease in the resistance of the channel and a consequent increase of ion flux, changing the delicate equilibrium of cellular functions [34].

Interestingly, the integrated area of the Amide I band increased significantly ($p < 0.05$) with the increase of the frequency of the applied electromagnetic field, demonstrating that the displacement of protein α -helices is closely dependent on the frequency of the applied electromagnetic field.

4. DISCUSSION

The results of this study have induced us to assume that tumor cells can be irradiated by an electromagnetic field at a frequency that is very close to the natural resonant frequencies of cancer membrane channels, producing a resonance phenomenon which should induce the maximum displacement of channels α -helices, amplifying ion flux across the same channels following the 2nd law of Ohm, as schematized in a figure [34].

Previous studies on the effects of radiation therapy showed that the use of beams of X-rays, gamma rays or particle radiation (protons or heavy ions) towards targeted areas can also damage normal non-irradiated cells neighboring tumor cells, a phenomenon termed 'bystander effect' [41 - 45]. This effect still represents an important limitation to the use of radiation therapy for the treatment of cancer.

In contrast, by following resonance theory, neighboring normal cells would not be damaged significantly by the irradiation of an electromagnetic field at the natural resonant frequency of cancer cell membrane channels, because the natural resonant frequencies of normal cells should be significantly different from the natural resonant frequencies of tumor cells.

Furthermore, it has been shown that even a minimum alteration of ion flux across cell membrane channels induces relevant cell damage [46, 47], so that the resonant mechanism described above can be successfully used for cancer treatment.

The angular deviation of α -helices in membrane channels of tumor cells can be measured as a function of the frequency of the electromagnetic field to which the tumor tissue is exposed. The maximum displacement of channels α -helices should occur at the natural resonance frequency of the cancerous cells which are being analyzed, inducing their death. To this aim, spectroscopic circular dichroism can be used to measure the angular deviations of α -helices in membrane channels of tumor cells as a function of the frequency of an applied electromagnetic field, using variable frequency power generators.

In addition, patch clamp technique can be used for measuring ionic currents across cellular membrane channels of the tumor tissue as a function of the frequency of an applied electromagnetic field, so that the largest value of ionic current would be measured at the natural resonance frequency of tumor cells. Furthermore, the largest decrease in cellular transmembrane potential in a tumor cell should occur at the natural resonant frequency, following the results obtained in previous studies by the authors of this study [16, 17].

Analogue measurement of the physical-chemical parameters cited above can be carried out in neighboring "non-resonant" normal cells, in order to verify that significantly lower values are obtained in normal cells in comparison with those obtained in cancer cells, in order to confirm the absence of a bystander effect after the irradiation by resonant electromagnetic fields of the targeted areas.

Once the natural resonant frequencies of cellular channels α -helices in typical cancerous tissues have been found, experimental verification of results would be carried out by measuring the disease progression in living beings after irradiation by electromagnetic fields at resonant frequencies.

CONCLUSION

Given that cancerous tissues should have natural frequencies quite different from the natural frequencies of healthy tissues, it can be hypothesized that irradiating tumor tissues by electromagnetic fields at frequencies close to the natural frequencies of tumor cells membrane channels would cause resonant interaction with membrane channels α -helices, thereby altering ion flux across them and damaging cellular functions of cancer cells. However, such a perspective should be confirmed by biochemical analyses and medical approaches.

Such resonant frequencies can be obtained for each tumor tissue by both spectroscopic and biochemical measurements of channels α -helices rotation and ion flux across cell channels as a function of the frequency of electromagnetic fields. In any case, clinical trials are needed to evaluate the possible application of such a result.

DISCLAIMERS

The views expressed in the submitted article are our own and not an official position of the institution or funder. The corresponding author is responsible for statistical analyses on the title page. No financial support was received to carry out this study.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

HUMAN AND ANIMAL RIGHTS

No animals/humans were used for studies that are the basis of this research.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIAL

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

We are grateful to Prof. Riccardo Ientile (Department of Biomedical Sciences and Morpho-functional images, University of Messina, Italy) for his kind and precious collaboration regarding the preparation of SH-SY5Y neuronal-like cells samples. The corresponding author is responsible for statistical analyses on the title page.

REFERENCES

- [1] Ottoman RE, Langdon EA, Rochlin DB, Smart CR. Side-effects of combined radiation and chemotherapy in the treatment of malignant tumors. *Radiology* 1963; 81(6): 1014-7. [http://dx.doi.org/10.1148/81.6.1014] [PMID: 14101708]
- [2] Coates A, Abraham S, Kaye SB, *et al.* On the receiving end-patient perception of the side-effects of cancer chemotherapy. *Eur J Cancer Clin Oncol* 1983; 19(2): 203-8. [http://dx.doi.org/10.1016/0277-5379(83)90418-2] [PMID: 6681766]
- [3] Staritz M, Adler G, Knuth A, Schmiegel W, Schmoll H-J. Side-effects of cancer chemotherapy on the gastrointestinal tract – pathophysiology, prophylaxis and therapy. The Netherlands: Kluwer Academic Publishers 2003.
- [4] Peeters KCMJ, van de Velde CJH, Leer JWH, *et al.* Late side effects of short-course preoperative radiotherapy combined with total mesorectal excision for rectal cancer: Increased bowel dysfunction in irradiated patients: a Dutch colorectal cancer group study. *J Clin Oncol* 2005; 23(25): 6199-206. [http://dx.doi.org/10.1200/JCO.2005.14.779] [PMID: 16135487]
- [5] Kayl AE, Meyers CA. Side-effects of chemotherapy and quality of life in ovarian and breast cancer patients. *Curr Opin Obstet Gynecol* 2006; 18(1): 24-8. [http://dx.doi.org/10.1097/01.gco.0000192996.20040.24] [PMID: 16493256]
- [6] Marijnen CAM, Kapiteijn E, van de Velde CJH, *et al.* Cooperative Investigators of the Dutch Colorectal Cancer Group. Acute side effects and complications after short-term preoperative radiotherapy combined with total mesorectal excision in primary rectal cancer: report of a multicenter randomized trial. *J Clin Oncol* 2002; 20(3): 817-25. [http://dx.doi.org/10.1200/JCO.2002.20.3.817] [PMID: 11821466]
- [7] Nurgali K, Jagoe RT, Abalo R. Adverse effects of cancer chemotherapy: Anything new to improve tolerance and reduce sequelae?. *Front Oncol* 2018; pp. 1-245. [http://dx.doi.org/10.3389/978-2-88945-482-2]
- [8] Barbault A, Costa FP, Bottger B, *et al.* Amplitude-modulated electromagnetic fields for the treatment of cancer: Discovery of tumor-specific frequencies and assessment of a novel therapeutic approach. *J Exp Clin Cancer Res* 2009; 28: 51. [http://dx.doi.org/10.1186/1756-9966-28-51] [PMID: 19366446]
- [9] Costa FP, de Oliveira AC, Meirelles R, *et al.* Treatment of advanced hepatocellular carcinoma with very low levels of amplitude-modulated electromagnetic fields. *Br J Cancer* 2011; 105(5): 640-8. [http://dx.doi.org/10.1038/bjc.2011.292] [PMID: 21829195]
- [10] Kirson ED, Gurvich Z, Schneiderman R, *et al.* Disruption of cancer cell replication by alternating electric fields. *Cancer Res* 2004; 64(9): 3288-95. [http://dx.doi.org/10.1158/0008-5472.CAN-04-0083] [PMID: 15126372]
- [11] Kirson ED, Dbalý V, Tovaryš F, *et al.* Alternating electric fields arrest cell proliferation in animal tumor models and human brain tumors. *Proc Natl Acad Sci USA* 2007; 104(24): 10152-7. [http://dx.doi.org/10.1073/pnas.0702916104] [PMID: 17551011]
- [12] Orel VE, Kudryavets YI, Satz S, *et al.* Effects of mechanochemically activated doxorubicin and 40 MHz frequency irradiation on human A-549 lung carcinoma cells. *Exp Oncol* 2004; 26(4): 271-7. [PMID: 15627058]
- [13] Ronchetto F, Barone D, Cintorino M, *et al.* Natural frequency of cancer cells as a starting point in cancer treatment. *Curr Sci* 2016; 110: 1828-32. [http://dx.doi.org/10.18520/cs/v110/i9/1828-1832]
- [14] Ronchetto F, Barone D, Cintorino M, *et al.* Extremely low frequency-modulated static magnetic fields to treat cancer: A pilot study on patients with advanced neoplasm to assess safety and acute toxicity. *Bioelectromagnetics* 2004; 25(8): 563-71. [http://dx.doi.org/10.1002/bem.20029] [PMID: 15515038]
- [15] Calabrò E, Magazù S, Campo S. Microwave-induced increase of amide I and amide II vibration bands and modulating functions of sodium-chloride, sucrose and trehalose aqueous solutions: The case study of Haemoglobin. *Res J Chem Environ* 2012; 16(4): 59-67.
- [16] Calabrò E, Condello S, Currò M, *et al.* 50 Hz electromagnetic field produced changes in FTIR spectroscopy associated with mitochondrial transmembrane potential reduction in neuronal-like SH-SY5Y cells. *Oxid Med Cell Longev* 2013a.414393:8 [http://dx.doi.org/10.1155/2013/414393] [PMID: 23970948]
- [17] Calabrò E, Condello S, Currò M, *et al.* Effects of low intensity static magnetic field on FTIR spectra and ROS production in SH-SY5Y neuronal-like cells. *Bioelectromagnetics* 2013b; 34(8): 618-29. [http://dx.doi.org/10.1002/bem.21815] [PMID: 24217848]
- [18] Calabrò E, Magazù S. Non-thermal effects of microwave oven heating on ground beef meat studied in the mid-infrared region by FTIR spectroscopy. *Spectrosc Lett* 2014a; 47(8): 649-56. [http://dx.doi.org/10.1080/00387010.2013.828313]
- [19] Calabrò E, Magazù S. Unfolding-induced in haemoglobin by exposure to electromagnetic fields: A FTIR spectroscopy study. *Orient J Chem* 2014b; 30(1): 31-5. [http://dx.doi.org/10.13005/ojc/300104]
- [20] Calabrò E, Magazù S. Fourier self deconvolution analysis of β -sheet contents in the amide I region of haemoglobin aqueous solutions under exposure to 900 MHz microwaves and bioprotective effectiveness of sugars and salt solutions. *Spectrosc Lett* 2015; 48(10): 741-7. [http://dx.doi.org/10.1080/00387010.2015.1011278]
- [21] Magazù S, Calabrò E, Caccamo MT, Cannuli A. The shielding action of disaccharides for typical proteins in aqueous solution against static, 50 Hz and 1800 MHz frequencies electromagnetic fields. *Curr Chem Biol* 2016; 10(1): 57-64. [http://dx.doi.org/10.2174/2212796810666160419153722]
- [22] Calabrò E, Magazù S. Effects of the addition of sodium chloride to a tetrameric protein in water solution during exposure to high frequency electromagnetic field. *Open Biotechnol J* 2017b; 11: 72-80. [http://dx.doi.org/10.2174/1874070701711010072]
- [23] Bhimarao M. Collapsing Cancer Cells. California State Science Fair: Exploiting the elasticity and natural frequency of a cancer cell's cytoskeleton. project n. S1703 2011.
- [24] Wee H, Voloshin A. Modal analysis of a spreading osteoblast cell in culturing. 38th Annual Northeast Bioengineering Conference (NEBEC 2012). 16-18 March 2012; Philadelphia, Pennsylvania: USA. 2012. [http://dx.doi.org/10.1109/NEBC.2012.6207016]
- [25] Becchetti A. Ion channels and transporters in cancer. 1. Ion channels and cell proliferation in cancer. *Am J Physiol Cell Physiol* 2011; 301(2): C255-65. [http://dx.doi.org/10.1152/ajpcell.00047.2011] [PMID: 21430288]
- [26] Lang F, Stourmaras C. Ion channels in cancer: Future perspectives and clinical potential. *Philos Trans R Soc Lond B Biol Sci* 2014; 369(1638):20130108 [http://dx.doi.org/10.1098/rstb.2013.0108] [PMID: 24493756]
- [27] Oosterwijk E, Gillies RJ. Targeting ion transport in cancer. *Philos Trans R Soc Lond B Biol Sci* 2014; 369(1638):20130107 [http://dx.doi.org/10.1098/rstb.2013.0107] [PMID: 24493755]
- [28] Peruzzo R, Biasutto L, Szabó I, Leanza L. Impact of intracellular ion channels on cancer development and progression. *Eur Biophys J* 2016; 45(7): 685-707. [http://dx.doi.org/10.1007/s00249-016-1143-0] [PMID: 27289382]
- [29] Turner KL, Sontheimer H. Cl⁻ and K⁺ channels and their role in primary brain tumour biology. *Philos Trans R Soc Lond B Biol Sci* 2014; 369(1638):20130095 [http://dx.doi.org/10.1098/rstb.2013.0095] [PMID: 24493743]
- [30] Becker L, Bannwarth M, Meisinger C, *et al.* Preprotein translocase of

- the outer mitochondrial membrane: Reconstituted Tom40 forms a characteristic TOM pore. *J Mol Biol* 2005; 353(5): 1011-20. [http://dx.doi.org/10.1016/j.jmb.2005.09.019] [PMID: 16213519]
- [31] Feria Bourrellier AB, Valot B, Guillot A, Ambard-Bretteville F, Vidal J, Hodges M. Chloroplast acetyl-CoA carboxylase activity is 2-oxoglutarate-regulated by interaction of PII with the biotin carboxyl carrier subunit. *Proc Natl Acad Sci USA* 2010; 107(1): 502-7. [http://dx.doi.org/10.1073/pnas.0910097107] [PMID: 20018655]
- [32] Szabo I, Zoratti M. Mitochondrial channels: Ion fluxes and more. *Physiol Rev* 2014; 94(2): 519-608. [http://dx.doi.org/10.1152/physrev.00021.2013] [PMID: 24692355]
- [33] Alberts B, Johnson A, Lewis J, *et al.* *Molecular Biology of the Cell*. 4th ed. NY, USA: Garland Science 2002.
- [34] Calabrò E, Magazù S. Resonant interaction between electromagnetic fields and proteins: A possible starting point for the treatment of cancer. *Electromagn Biol Med* 2018b; 37(3): 155-68. [http://dx.doi.org/10.1080/15368378.2018.1499031] [PMID: 30019948]
- [35] Calabrò E, Magazù S. Infrared spectroscopic demonstration of magnetic orientation in SH-SY5Y neuronal-like cells induced by static or 50 Hz magnetic fields. *Int J Radiat Biol* 2019; 95(6): 781-7. [http://dx.doi.org/10.1080/09553002.2019.1571256] [PMID: 30668250]
- [36] Hammiche A, German MJ, Hewitt R, Pollock HM, Martin FL. Monitoring cell cycle distributions in MCF-7 cells using near-field photothermal microspectroscopy. *Biophys J* 2005; 88(5): 3699-706. [http://dx.doi.org/10.1529/biophysj.104.053926] [PMID: 15722424]
- [37] Calabrò E. Competition between hydrogen bonding and protein aggregation in neuronal-like cells under exposure to 50 Hz magnetic field. *Int J Radiat Biol* 2016; 92(7): 395-403. [http://dx.doi.org/10.1080/09553002.2016.1175679] [PMID: 27173917]
- [38] Calabrò E, Magazù S. Parallel β -sheet vibration band increases with proteins dipole moment under exposure to 1765 MHz microwaves. *Bioelectromagnetics* 2016; 37(2): 99-107. [http://dx.doi.org/10.1002/bem.21956] [PMID: 26833949]
- [39] Calabrò E, Magazù S. The α -helix alignment of proteins in water solution toward a high-frequency electromagnetic field: A FTIR spectroscopy study. *Electromagn Biol Med* 2017a; 36(3): 279-88. [http://dx.doi.org/10.1080/15368378.2017.1328691] [PMID: 28632082]
- [40] Calabrò E, Magazù S. Direct spectroscopic evidence for competition between thermal molecular agitation and magnetic field in a tetrameric protein in aqueous solution. *Phys Lett A* 2018a; 382: 1389-94. [http://dx.doi.org/10.1016/j.physleta.2018.03.038]
- [41] Nagasawa H, Little JB. Induction of sister chromatid exchanges by extremely low doses of alpha-particles. *Cancer Res* 1992; 52(22): 6394-6. [PMID: 1423287]
- [42] Shao C, Folkard M, Michael BD, Prise KM. Targeted cytoplasmic irradiation induces bystander responses. *Proc Natl Acad Sci USA* 2004; 101(37): 13495-500. [http://dx.doi.org/10.1073/pnas.0404930101] [PMID: 15345742]
- [43] Prise KM, Schettino G, Folkard M, Held KD. New insights on cell death from radiation exposure. *Lancet Oncol* 2005; 6(7): 520-8. [http://dx.doi.org/10.1016/S1470-2045(05)70246-1] [PMID: 15992701]
- [44] Munro AJ. Bystander effects and their implications for clinical radiotherapy. *J Radiol Prot* 2009; 29(2A): A133-42. [http://dx.doi.org/10.1088/0952-4746/29/2A/S09] [PMID: 19454811]
- [45] Baskar R, Dai J, Wenlong N, Yeo R, Yeoh KW. Biological response of cancer cells to radiation treatment. *Front Mol Biosci* 2014; 1: 24. [http://dx.doi.org/10.3389/fmolb.2014.00024] [PMID: 25988165]
- [46] Leanza L, Managò A, Zoratti M, Gulbins E, Szabo I. Pharmacological targeting of ion channels for cancer therapy: *In vivo* evidences. *Biochim Biophys Acta* 2016; 1863(6 Pt B): 1385-97. [http://dx.doi.org/10.1016/j.bbamer.2015.11.032] [PMID: 26658642]
- [47] Rao VR, Perez-Neut M, Kaja S, Gentile S. Voltage-gated ion channels in cancer cell proliferation. *Cancers (Basel)* 2015; 7(2): 849-75. [http://dx.doi.org/10.3390/cancers7020813] [PMID: 26010603]