



HAL
open science

The Glycosylphosphatidylinositol-Anchored DFG Family Is Essential for the Insertion of Galactomannan into the β -(1,3)-Glucan-Chitin Core of the Cell Wall of *Aspergillus fumigatus*

Laetitia Muszkieta, Thierry Fontaine, Rémi Beau, Isabelle Mouyna, Marian Samuel Vogt, Jonathan Trow, Brendan Cormack, Lars-Oliver Essen, Grégory Jouvion, Jean-Paul Latge

► To cite this version:

Laetitia Muszkieta, Thierry Fontaine, Rémi Beau, Isabelle Mouyna, Marian Samuel Vogt, et al.. The Glycosylphosphatidylinositol-Anchored DFG Family Is Essential for the Insertion of Galactomannan into the β -(1,3)-Glucan-Chitin Core of the Cell Wall of *Aspergillus fumigatus*. *MSphere*, 2019, 4 (4), pp.e00397-19. <10.1128/mSphere.00397-19>. <hal-02545847>

HAL Id: hal-02545847

<https://hal.science/hal-02545847v1>

Submitted on 17 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



The Glycosylphosphatidylinositol-Anchored *DFG* Family Is Essential for the Insertion of Galactomannan into the β -(1,3)-Glucan–Chitin Core of the Cell Wall of *Aspergillus fumigatus*

Laetitia Muszkietta,^a Thierry Fontaine,^a Rémi Beau,^a Isabelle Mouyna,^a Marian Samuel Vogt,^b Jonathan Trow,^c Brendan P. Cormack,^c Lars-Oliver Essen,^b Gregory Jouvion,^d Jean-Paul Latgé^{a*}

^aUnité des *Aspergillus*, Institut Pasteur, Paris, France

^bFaculty of Chemistry, Philipps-Universität Marburg, Marburg, Germany

^cDepartment of Molecular Biology and Genetics, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

^dHistopathologie humaine et modèles animaux, Institut Pasteur, Paris, France

ABSTRACT The fungal cell wall is a complex and dynamic entity essential for the development of fungi. It is composed mainly of polysaccharides that are synthesized by protein complexes. At the cell wall level, enzyme activities are involved in post-synthesis polysaccharide modifications such as cleavage, elongation, branching, and cross-linking. Glycosylphosphatidylinositol (GPI)-anchored proteins have been shown to participate in cell wall biosynthesis and specifically in polysaccharide remodeling. Among these proteins, the *DFG* family plays an essential role in controlling polar growth in yeast. In the filamentous fungus and opportunistic human pathogen *Aspergillus fumigatus*, the *DFG* gene family contains seven orthologous *DFG* genes among which only six are expressed under *in vitro* growth conditions. Deletions of single *DFG* genes revealed that *DFG3* plays the most important morphogenetic role in this gene family. A sextuple-deletion mutant resulting from the deletion of all *in vitro* expressed *DFG* genes did not contain galactomannan in the cell wall and has severe growth defects. This study has shown that *DFG* members are absolutely necessary for the insertion of galactomannan into the cell wall of *A. fumigatus* and that the proper cell wall localization of the galactomannan is essential for correct fungal morphogenesis in *A. fumigatus*.

IMPORTANCE The fungal cell wall is a complex and dynamic entity essential for the development of fungi. It is composed mainly of polysaccharides that are synthesized by protein complexes. Enzymes involved in postsynthesis polysaccharide modifications, such as cleavage, elongation, branching, and cross-linking, are essential for fungal life. Here, we investigated in *Aspergillus fumigatus* the role of the members of the *Dfg* family, one of the 4 GPI-anchored protein families common to yeast and molds involved in cell wall remodeling. Molecular and biochemical approaches showed that *DFG* members are required for filamentous growth, conidiation, and cell wall organization and are essential for the life of this fungal pathogen.

KEYWORDS *Aspergillus fumigatus*, cell wall, glycobiology

The fungal cell wall is a complex and dynamic entity essential for the development of fungi. It has prominent and dual roles during the growth of fungal pathogens. It allows the pathogen to survive environmental challenges posed by nutrient stress, microbiota, or human cells, and it also is central to polarized growth, which helps the fungus to invade host tissues (1). The cell wall of *Aspergillus fumigatus* is mainly composed of polysaccharides organized in a three-dimensional (3D) network (2). Enzymes involved in the biosynthesis of linear glucan and chitin, the main cell wall


Citation Muszkietta L, Fontaine T, Beau R, Mouyna I, Vogt MS, Trow J, Cormack BP, Essen L-O, Jouvion G, Latgé J-P. 2019. The glycosylphosphatidylinositol-anchored *DFG* family is essential for the insertion of galactomannan into the β -(1,3)-glucan–chitin core of the cell wall of *Aspergillus fumigatus*. *mSphere* 4:e00397-19. <https://doi.org/10.1128/mSphere.00397-19>.

Editor Aaron P. Mitchell, Carnegie Mellon University

Copyright © 2019 Muszkietta et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Thierry Fontaine, thierry.fontaine@pasteur.fr, or Jean-Paul Latgé, jean-paul.latge@pasteur.fr.

* Present address: Jean-Paul Latgé, School of Medicine, University of Crete, Heraklion, Greece.

 Remodeling enzymes in cell wall synthesis are essential for fungal life

Received 5 June 2019

Accepted 25 June 2019

Published 31 July 2019

polysaccharides encountered in the fungal kingdom, have been identified previously (3–5). However, the transglycosidases responsible for the branching and cross-linking of these linear polysaccharides in a 3D rigid skeleton are only beginning to be discovered (6). Early studies have shown the merits of concurrent analyses of putative transglycosylases in different fungal species such as *Saccharomyces cerevisiae* and *Aspergillus fumigatus* with different cell wall compositions but similar central polysaccharide 3D cores. Such analyses has led to the selection of four groups of glycosylphosphatidylinositol (GPI)-anchored proteins common to all fungi which could potentially have a central role in the transglycosylation of the cell wall structural polysaccharides. First, a family of β -(1,3)-glucanoyltransferase activity discovered in *A. fumigatus* and encoded by the *GEL* gene or *GAS* gene is responsible for the elongation of the β -(1,3)-glucans and is essential in the biosynthesis of the cell wall (53, 54, 55). This family contains two subgroups that depend on the presence of a carbohydrate binding domain, which is responsible for a dual form of enzyme activity, β -glucan elongation and branching (6). Second, the Sps2p/Ecm33p family of GPI-anchored proteins has been described previously in both *S. cerevisiae* and *A. fumigatus* (7, 8). Members of the Sps2p family play an essential role in the formation of the ascospore cell wall in *S. cerevisiae*, whereas in *A. fumigatus*, Ecm33p is important for conidial morphogenesis and virulence. However, its enzymatic function remains unknown. Third, the Crhp family in *S. cerevisiae* is composed of three genes which are involved in the linkage of chitin to β -(1,6)-glucan in the cell wall (9, 10). Although also present in *A. fumigatus*, it is obvious that the Crh proteins do not have the same function in *S. cerevisiae* and *A. fumigatus* since there is no β -(1,6)-glucan in the cell wall of *A. fumigatus*. Finally, the two genes in the *DFG* (for “defective in filamentous growth”) family encode two GPI-anchored proteins with redundant activities in both *S. cerevisiae* and *Candida albicans* (11, 12). Although single knockouts of *DFG5* and *DCW1* are viable, a double knockout was synthetically lethal in both *S. cerevisiae* and *C. albicans* (11, 12). Cells depleted in either Dfg5p or Dcw1p released GPI-anchored cell wall proteins (GPI-CWPs) into the medium and showed increased cell volume, suggesting an alteration of the cell wall organization. However, the exact biochemical function of the Dfg proteins in yeast as well as in filamentous fungi remains unknown.

To understand the function of Dfg proteins in *A. fumigatus*, single-deletion and multiple *dfg* mutants targeting deletions of the members of the entire *DFG* family were constructed and analyzed. *DFG* multiple deletion resulted in the total loss of the cell wall galactomannan (GM), which was associated with severe growth phenotypes.

RESULTS

The *DFG* family in *A. fumigatus*. A BLAST query of the *S. cerevisiae* Dfg5p/Dcw1p protein sequences against the *A. fumigatus* genome database (https://fungi.ensembl.org/Aspergillus_fumigatus/a1163/Info/Index) identified seven paralogs. The percentages of identity and similarity ranged between 15% and 42% and between 30% and 60%, respectively (see Table S1 in the supplemental material). *In silico* analysis revealed that all of the proteins encoded by these genes contained a secretory signal peptide at the N-terminal region (identified by the use of the SignalP website [<http://www.cbs.dtu.dk/services/SignalP/>]), a hydrophobic region at the C-terminal region with an ω -site characteristic of GPI-anchored proteins (except Dfg2p) (identified by the use of the BigPI website [http://mendel.imp.ac.at/gpi/fungi_server.html] and the PredGPI website [<http://gpcr.biocomp.unibo.it/predgpi/pred.htm>]), and the glycosyl hydrolase domain GH76 characteristic of Dfg5p/Dcw1p proteins from yeast (see Fig. S1 in the supplemental material). The expression levels of the *DFG* genes seen during different stages of development of *A. fumigatus* (swollen conidia, germinating conidia, and mycelium grown in Sabouraud liquid medium; sporulating mycelium and conidia cultivated on Sabouraud agar medium) were analyzed by quantitative real-time PCR (qRT-PCR) (Fig. 1). With the exception of *DFG6*, all of the *DFG* genes were expressed in the different fungal stages but their relative expression levels were highly dependent on the growth stage considered (Fig. 1). *DFG3* was the gene that was the most highly




FIG 1 Expression profiles of seven *DFG* genes in different stages of *A. fumigatus* development assayed by qRT-PCR. RNA was extracted from freshly harvested resting (0 h), swollen (4 h), or germinated (6 to 8 h) conidia and mycelium grown in Sabouraud liquid medium and sporulating mycelium grown on Sabouraud solid medium. The relative expression levels of individual *DFG* genes were analyzed with the $2^{-\Delta\Delta CT}$ method with the *EF1α* gene used as an internal control for normalization. Microscopy analysis of developmental stages was followed by staining with calcofluor white. (Values represent means and standard deviations of results from three different experiments. Bars: time zero h [0h], 4h, 6h, and 8h, 1 μm ; 16h, 10 μm ; 24h, 20 μm .)

expressed in germinating conidia. Three *DFG* genes (*DFG1*, *DFG2*, and *DFG5*) were highly expressed in conidia and sporulating mycelium, while *DFG4* was highly expressed only in conidia and *DFG7* only in sporulating mycelium.

Construction of Δdfg mutants. Based on the gene expression analysis and to further understand their biological role in fungal life, successive deletions of all members of the multigene *DFG* family in *A. fumigatus* were undertaken. These multiple deletions were carried out by employing the β -rec/six system (13). Single-deletion mutants (except for the nonexpressed *DFG6* gene) and multiple-deletion mutants ($\Delta dfg5/2$; $\Delta dfg5/2/1$; $\Delta dfg5/2/1/4$; $\Delta dfg5/2/1/3$; $\Delta dfg5/2/1/3/4$; $\Delta dfg5/2/1/3/4/7$) were constructed. Strategies for gene replacements and their validation are shown in Fig. S2, and the corresponding mutants are listed in Table S2.

Mycelial growth and hyphal morphology of Δdfg mutants. At 37°C, no growth differences were observed for the single-deletion strains ($\Delta dfg1$, $\Delta dfg2$, $\Delta dfg4$, $\Delta dfg5$, and $\Delta dfg7$) and multiple-deletion strains ($\Delta dfg5/2$, $\Delta dfg5/2/1$, $\Delta dfg5/2/1/4$) in comparison to the parental strain during growth on malt agar medium (Fig. 2A). Only the single-knockout $\Delta dfg3$ mutant and the multiple-knockout mutants containing a *DFG3* deletion ($\Delta dfg5/2/1/3$, $\Delta dfg5/2/1/3/4$, and $\Delta dfg5/2/1/3/4/7$) displayed a significant reduction in vegetative mycelial growth compared to the parental strain and the *DFG3* revertant strain (Fig. 2). In the sextuple-deletion mutant, the absence of *DFG6* expression was verified by qRT-PCR at different stages of development of *A. fumigatus* (not shown). This result indicated that the sextuple-deletion mutant can be considered to be a mutant lacking the entire *DFG* gene family. Even though all mutants were evaluated, only the phenotypic analyses of the $\Delta dfg3$ single-deletion mutant and the $\Delta dfg5/2/1/3/4/7$ sextuple-deletion mutant are reported here. The differences in the amounts of mycelium produced by the Δdfg mutants and the parental strain in Sabouraud liquid medium were much lower than those seen in agar media (Fig. 3B). However, the mycelia of these mutants grown in both solid and liquid media were highly branched (Fig. 2B). In liquid cultures, this hyperbranching phenotype resulted in the production of a myriad of very small and tight fungal balls which looked different from the homogeneous mycelial mass produced by the parental strain (Fig. 3A).




FIG 2 Growth of *DFG* deletion mutant strains on solid medium. (A) Radial growth of the parental and *DFG* deletion mutant strains on malt agar medium (48 h at 37°C). (B) Mycelial morphology of the parental strain and Δdfg mutants grown on malt agar medium. Hyphae were stained with calcofluor white (bar, 10 μ m). (C) Mycelial dry weight of the Δdfg mutants obtained after 48 h of growth on malt agar medium at 37°C. (Values represent means and standard deviations of results from three different experiments; statistically significant differences [$P < 0.001$] are indicated by an asterisk.)

Conidiation, conidial morphology of Δdfg mutants, and susceptibility to drugs.

The capacity of the different mutants to conidiate was assayed on malt agar-containing tubes following incubation for 10 days at room temperature. At 37°C, the slant was entirely covered by the fungus and no conidiation difference was observed for the single-deletion strains ($\Delta dfg1$, $\Delta dfg2$, $\Delta dfg4$, $\Delta dfg5$, and $\Delta dfg7$) and multiple-deletion strains ($\Delta dfg5/2$, $\Delta dfg5/2/1$, and $\Delta dfg5/2/1/4$) compared to the parental strain. Interestingly, the single-deletion $\Delta dfg3$ mutant produced as many conidia as the parental strain whereas the sextuple-deletion $\Delta dfg5/2/1/3/4/7$ mutant displayed a drastic reduction in conidiation (Table 1). This conidiation defect could be compensated for by the addition of 6% KCl (Table 1). Similarly, 52% of the resting conidia from the sextuple-deletion $\Delta dfg5/2/1/3/4/7$ mutant were intracellularly stained with fluorescein isothiocyanate (FITC), whereas the resting conidia from the parental and $\Delta dfg3$ strains were weakly stained by FITC at the cell surface (Fig. S3). Moreover, conidia of the sextuple-deletion $\Delta dfg5/2/1/3/4/7$ mutant germinated faster (~18% germ tubes were formed after 4 h of incubation at 37°C) than those of the $\Delta dfg3$ and parental strains, which showed nearly no germinated conidia after the same 4 h of incubation time (Fig. 4A). While conidia of the parental and mutant strains were similar in size in the resting stage, the swollen conidia of the sextuple-deletion $\Delta dfg5/2/1/3/4/7$ mutant were approximately 1.5 times larger than those of the parental strain (Fig. 4B). In line with what was seen for FITC




FIG 3 Morphology of the Δdfg mutants in liquid medium. (A) Visual aspect of Δdfg cultures in Sabouraud medium (10^6 conidia/50 ml medium) after 24 h at 37°C (bar, 1 cm). (B) Mycelial dry weight of the Δdfg mutants quantified after 24 h of growth in Sabouraud medium at 37°C (10^7 conidia/50 ml medium). (Values represent means and standard deviations of results from four different experiments; statistically significant differences [$P < 0.001$] are indicated by an asterisk. ns, not significant.)

staining, calcofluor white (CFW) staining indicated that the cell walls of sextuple-deletion $\Delta dfg5/2/1/3/4/7$ mutant were more permeable than those of the parental strain, since intracellular labeling of around 50% of the swollen conidia was seen following 5 min of exposure to the stain (Fig. 4B). The single-deletion $\Delta dfg3$ mutant and the sextuple-deletion $\Delta dfg5/2/1/3/4/7$ mutant were more susceptible to voriconazole and itraconazole and to the cell wall-disturbing calcofluor white dye than the parental strain on RPMI medium after 72 h of incubation at 37°C (Fig. 5). All these conidial phenotypes suggested that the cell wall integrity was altered in the $\Delta dfg5/2/1/3/4/7$ mutant and that the alterations were associated with defects in cell wall permeability.

Impact of DFG deletions on cell wall galactomannan (GM). In *A. fumigatus*, GM is either cross-linked to cell wall β -(1,3)-glucans or membrane bound through a GPI anchor (LGM) (14–16). The chemical compositions of the alkali-insoluble (AI) and alkali-soluble (AS) fractions of the *A. fumigatus* cell wall of parental and mutant strains are shown in Table 2. A decrease in the level of cell wall galactomannan (GM) was observed in both the AI and AS fractions in the $\Delta dfg3$ mutant, and its complete absence was noted in the AI fraction of the $\Delta dfg5/2/1/3/4/7$ mutant. Quantification of GM carried out with the Δdfg mutants was confirmed *in situ*, as shown by the absence of labeling of the GM with an antigalactofuran monoclonal antibody at the cell surface of the $\Delta dfg3$ mutant (Fig. 6). The reduction in the level of cell wall GM was primarily compensated for by an increase in the glucose content within the AI fraction and by an increase in the galactosamine content within the AS fraction, corresponding to larger amounts of β -(1,3)-glucan and galactosaminogalactan, respectively (Table 2). The single-deletion $\Delta dfg3$ mutant showed a significant reduction in GM content (40% of the level seen with the AI fraction) compared with the parental strain, whereas no cell wall cross-linked GM was detected in the sextuple-deletion mutant (Fig. 7A). In addition, quantification of the GM in the single-deletion $\Delta dfg3$ mutant and the multiple-deletion

TABLE 1 Conidiation of the Δdfg mutants and the parental strains in the presence or absence of KCl

Strain	No. of conidia grown in ^a :	
	Malt agar (2%)	Malt agar + KCl (6%)
Parental strain	$(2.6 \times 10^8) \pm 0.28$	$1.0 \times 10^8 \pm 0.15$
$\Delta dfg3$ mutant	$(2.4 \times 10^8) \pm 0.10$	ND
$\Delta dfg5/2/1/3/4/7$ mutant	$(2.1 \times 10^5) \pm 0.68$	$4.4 \times 10^7 \pm 0.87$

^aData represent means and standard deviations (SD) of results from three different experiments. ND, not determined.




FIG 4 Conidial germination of the Δdfg strains. (A) Germination kinetics of the parental and Δdfg mutants. (B) Calcofluor white staining of swollen conidia of the parental strain and Δdfg mutants (incubated in liquid Sabouraud medium for 4 h at 37°C). The estimated proportion (50%) of intracellularly fluorescent swollen conidia was determined by counting 200 conidia for the $\Delta dfg5/2/1/3/4/7$ mutant. (Bar, 2 μm .)

$\Delta dfg5/2/1/3$, $\Delta dfg5/2/1/3/4$, and $\Delta dfg5/2/1/3/4/7$ mutants showed an additive effect consisting of a concomitant decrease in the level of cell wall GM content corresponding to an increase of the number of *DFG* deletions.

In contrast to the cell wall GM, the GM content was slightly increased in membrane preparations (Fig. 7B). The amount of GM in the $\Delta dfg3$ mutant was close to 110% of the GM content in the membrane of the parental strain and reached 160% of the parental strain level in the sextuple-deletion mutant (Fig. 7B). In addition, no significant differences were observed in the molar ratios of the hexose content of the membrane-bound GM of the parental and the sextuple-deletion $\Delta dfg5/2/1/3/4/7$ strains. Structural analysis of membrane-bound GM showed the presence of terminal galactofuranose, 5-O-substituted galactofuranose, 2-O-substituted mannose, 6-O-substituted mannose, 2,6-di-O-substituted mannose, and 2,3-di-O-substituted mannose residues, typical of the GM sequence (14, 16). The lipid anchor of LGM was isolated from the sextuple-deletion mutant and analyzed by electrospray-mass spectrometry (ES-MS). The major pseudomolecular ion $[M-H]^-$ at $m/z = 1,086$ characterized the GPI-related anchor of LGM with the presence of a glucosamine residue linked to an inositolphosphoceramide moiety, which was composed of a C_{18} -phytosphingosine and a 2-hydroxy- $C_{24:0}$ fatty acid, as previously described (16) (Fig. S4). Taking into account all these structural data, the level of GM from the $\Delta dfg5/2/1/3/4/7$ mutant is identical to that from the parental




FIG 5 Sensitivity of the parental and Δdfg strains to drugs. The levels of sensitivity of the Δdfg mutants to calcofluor white (CFW [50 $\mu\text{g}/\text{ml}$]) and azole compounds (voriconazole [100 ng/ml] and itraconazole [50 ng/ml]) were determined after 72 to 96 h of growth at 37°C on RPMI agar medium.

TABLE 2 Monosaccharide compositions of cell wall fractions from the parental and Δdfg strains^a

Strain	% indicated monosaccharide in:									
	Alkali-insoluble (AI) fraction					Alkali-soluble (AS) fraction				
	Man	Glc	Gal	GlcN	Man	Glc	Gal	GalN	AI (%)	
Parental strain	10.90 ± 0.35	46.25 ± 0.75	11.26 ± 0.09	31.07 ± 1.11	4.78 ± 1.79	74.42 ± 1.61	10.67 ± 0.04	9.67 ± 1.20	68.91 ± 4.24	
$\Delta dfg3$	6.10 ± 0.17	50.17 ± 0.65	10.43 ± 0.23	32.10 ± 0.82	2.18 ± 0.13	55.81 ± 1.98	8.79 ± 1.10	32.78 ± 3.07	58.61 ± 4.67	
$\Delta dfg5/2/1/3/4/7$	0	61.56 ± 2.11	0	34.70 ± 1.62	0.47 ± 0.09	61.47 ± 4.01	1.85 ± 1.08	35.61 ± 4.99	56.46 ± 8.56	

^aData represent means and standard deviations (SD) of results from three different experiments. Man, mannose; Glc, D-glucopyranose; Gal, D-galactopyranose; GlcN, 2-amino-2-deoxy-D-glucopyranose.

strain, indicating that the intracellular biosynthetic pathway of GM was not altered by *DFG* deletions.

Interestingly, the amount of GM in the culture filtrate of the $\Delta dfg3$ and $\Delta dfg5/2/1/3/4/7$ mutants increased substantially in comparison to the level seen with the parental strain (Fig. 7C). An additive effect with a concomitant increase in the free GM content in the culture filtrate was seen with the increase of the number of *DFG* deletions.

In conclusion, the intracellular biosynthetic pathway of GM was not altered by *DFG* deletions qualitatively or quantitatively. In contrast, the *DFG* deletion resulted in a defect in insertion of this polysaccharide into the cell wall and in associated extracellular release of the GM into the external medium. Even though only the deletion of the *DFG 3* gene resulted in a morphogenetic phenotype among all the single-deletion Δdfg mutants, the comparison of the GM phenotypes of the $\Delta dfg3$ mutant and the $\Delta dfg5/2/1/3/4/7$ multiple-deletion mutant indicated additivity of the functions of these *DFG* proteins in the insertion of GM into the cell wall.

Complementation of the $\Delta dfg5/dcw1$ yeast mutant by *AfDFG3*. A double-knockout $\Delta dcw1/\Delta dfg5$ mutant was shown previously to be synthetically lethal in yeast (11). Functional complementation by *A. fumigatus DFG3* (*AfDFG3*) in *S. cerevisiae* was investigated using a thermosensitive $dcw1^{ts}/\Delta dfg5$ mutant. This mutant was unable to grow on solid minimum medium at 37°C and produced a 15% proportion of enlarged yeast cells in liquid medium at 37°C. The introduction of the *AfDFG3* gene into the mutant strain restored normal growth at 37°C at a level that was similar to that seen with the control *DCW1/\Delta dfg5* mutant strain (Fig. 8), showing that the *AfDFG3* genes and *S. cerevisiae DCW1* (*ScDCW1*) genes share similar biological activities.




FIG 6 Labeling of the galactomannan present at the cell surface of the Δdfg mutant complemented strain and the parental strain by an anti-GalF antibody. Mycelia grown for 16 h at 37°C in Sabouraud medium were fixed with *p*-formaldehyde and subjected to immunolabeling with anti-GalF antibody and anti-rat FITC secondary antibody. (Bar, 100 μ m.) Both fluorescence and bright-field images are shown for the $\Delta dfg3$ mutant.




FIG 7 Impact of *DFG* deletion on galactomannan localization. (A) Galactomannan content in the alkali-insoluble fraction of the cell wall (estimated as the amount of mannose in the alkali-insoluble fraction). (B) Lipogalactomannan content extracted from membrane preparation. (C) Galactomannan levels estimated in the culture supernatant (quantified by sandwich enzyme-linked immunosorbent assay [ELISA] using a Platelia kit). Values represent means and standard deviations of data from three different experiments. Statistically significant differences ($P < 0.05$) are indicated by an asterisk.

DISCUSSION

The phenotypes of Δdfg mutants are different among yeast and mold. In this report, we describe the results of characterization of the *DFG* family in *A. fumigatus*. The deletion of *DFG* genes led to a major reduction in vegetative growth with hyper-branched hyphae. In addition, it was shown that the genes of the *DFG* family have additive biological activities, as shown also in *S. cerevisiae*. Nine *DFG* paralogs have been annotated in the genome of *Neurospora crassa*, and among the single-deletion mu-




FIG 8 Functional complementation of yeast *dcw1^{ts}/Δdfg5* mutant by *AfDFG3*. (A) Growth on YNB plate medium after 3 days at 30°C and 37°C. (B) Calcofluor white staining of the cells after 2 days of culture in YNB liquid medium at 30°C and 37°C. An average of 15% of enlarged yeast cell (>15 μm) were estimated in the *dcw1^{ts}/Δdfg5* mutant by counting 400 cells (bar, 6 μm).

tants, only two have been studied, both of which showed altered morphology (17). Of the two mutants, the $\Delta dfg5$ mutant (NCU03770) showed a reduction in colony size and a hyperbranched mycelial pattern, in similarity to the $\Delta dfg3$ mutant in *A. fumigatus*. The $\Delta dcw1$ mutant (NCU08127) did not show a significant growth phenotype in *Neurospora*, but the growth phenotype of the $\Delta dfg5/\Delta dcw1$ double-deletion mutant of *N. crassa* showed more-severe effects than were seen with the strain with a single $\Delta dfg5$ deletion, with increased susceptibility to cell wall-perturbing agents, such as caspofungin (17). In yeast, single deletion of the paralogous *DFG5* gene or *DCW1* gene also leads to cell wall alterations and growth defects. The *ScΔdfg5* mutant has been reported previously to be defective in filamentous growth, cell polarity, and elongation (18). Mutant *ScΔdcw1* (for “defective in cell wall”) was hypersensitive to zymolyase, a cell wall-digesting enzyme, suggesting a role in cell wall organization (19). In *C. albicans*, a mutant lacking *DFG5* was defective in hyphal formation at alkaline pH whereas a mutant lacking *DCW1* had no obvious phenotype. In both *C. albicans* and *S. cerevisiae*, deletions of *DFG5* and *DCW1* were synthetically lethal (12, 19). In *A. fumigatus*, the entire deletion of all members of the DFG family (since *DFG6* remained unexpressed in the sextuple-deletion mutant) was not lethal. Our data showed, as often reported for orthologous genes of yeast and filamentous fungi, that the phenotypes of their respective mutants were different. Among the single-deletion mutants, only *DFG3* deletion led to a growth defect; the major phenotypic difference was seen in the sextuple-deletion mutant, where additive effects on mycelial morphology, germination

rate, drug sensitivity, cell wall composition, and permeability were observed with increasing numbers of DFG gene deletions.

Function of the Dfg proteins. On the basis of sequence homology, Dfg proteins have been assigned to the GH76 glycoside hydrolase family in the CAZy database (20). Dfg proteins were predicted to act as endo- α -(1,6)-mannanases based on the homologies of the sequence of a single member of this family with that of the *Aman6* protein from *Bacillus circulans* (21, 22). However, to date, there has been no biochemical evidence found to indicate that any fungal Dfg protein acts as an endo- α -(1,6)-mannanase or possesses any hydrolytic activity with respect to cell wall polysaccharides. In our hands, efforts to produce recombinant AfDfg3p were not successful. Since ScDcw1p showed 32% identity with AfDfg3p (see Table S1 in the supplemental material) and the *S. cerevisiae* *dcw1^{ts}/Δdfg5* mutant was functionally complemented by AfDFG3 (Fig. 8), we investigated the enzymatic activity of the recombinant ScDcw1 protein. The glycosylhydrolase activity of the recombinant ScDcw1 protein was assayed with several α -mannosides (see Text S1 in the supplemental material). The transglycosylase activity has been also investigated using galactomannan as a donor and soluble β -(1,3)-glucans or the cell wall AI fraction from *S. cerevisiae* as an acceptor (Text S1). No hydrolase or transfer activity has been detected with any of these substrates, even with the sensitive fluorometry assay used with the bacterial GH76 member (23) (Text S1). The enzyme activity of fungal GH76 family members therefore remains uncharacterized but certainly does not represent that of an α -mannanase.

The increased secretion of GPI-anchored proteins in the *Δdfg5* mutant or the *Δdcw1* mutant in *S. cerevisiae* suggested that these proteins were involved in the cross-linking of GPI-CWPs to the cell wall glucans (17, 24). However, in addition to the absence of any biochemical demonstration of such transglycosylase activity, ascribing a cross-linking function remains controversial. In *C. albicans*, shutoff strains for *DFG5* and *DCW1* were previously shown to exhibit the release of both GPI-anchored and non-GPI cell wall mannoproteins (25). Moreover, in previous studies, secretion of very few GPI proteins, including Acw1p (=ECM33), Gel1p in *N. crassa*, and Cwp1p in *S. cerevisiae* (11, 17), was demonstrated, whereas localization of the major GPI-anchored protein, Gas1p, remained unchanged in both *Dfg5* and *Dcw1* single-deletion mutants in *S. cerevisiae* (26). In *N. crassa*, the same proteins were released in the culture supernatant of mutants with deletions of very different genes, such as those encoding a putative hyaluronic synthase (*CPS1*) or mannosyltransferase (*OCH1*) (27, 28). In *A. fumigatus*, *DFG* deletions led also to a 3-fold increase in the amount of secreted proteins (see Fig. S5 in the supplemental material). However, the six overproduced proteins were mainly non-GPI-anchored proteins such as endopolygalacturonase, endochitinases, and pectate lyase (Fig. S5). Such a modification of the secreted protein pattern has been indeed reported in other *A. fumigatus* mutants after genes coding for proteins with a known function in cell wall biosynthesis were deleted (29, 30). Taking the data together, the results suggest that Dfg proteins are not involved in the cell wall localization of GPI-anchored proteins, at least in filamentous fungi. The modification of the cell wall structure due to the loss of cell wall galactomannan in the *Δdfg3* single-deletion and multiple-deletion mutants leads to facilitation of the release of secreted proteins in transit through the cell wall. These cell wall modifications lead to a change in permeability such as was demonstrated when the *Δdfg3* single-deletion mutant and the *Δdfg5/2/1/3/4/7* multiple-deletion mutant were incubated with CFW or FITC (Fig. 5; see also Fig. S3). Intracellular labeling with FITC and CFW has been repeatedly performed with cell wall mutants of *A. fumigatus* to demonstrate the effect of a gene deletion on cell wall permeability (5, 8, 29–32). However, we have no idea of the nature of the material stained intracellularly by CFW or FITC.

Dfg proteins and the cell wall GM in *A. fumigatus*. GM is a polymer composed of a linear α -(1,2)/ α -(1,6)-mannan chain with short side chains of β -(1,5)-galactofuran (Gal_f) (33). This polysaccharide can be covalently bound to β -(1,3)-glucans in the cell wall, bound to the plasma membrane by a GPI anchor, or present in the extracellular

environment. Despite of its importance in fungal morphogenesis and in the host immune response (31), GM biosynthesis is poorly understood. Previous studies have shown that GM biosynthesis takes place in the Golgi apparatus, into which sugar-donor (UDP-galactofuranose and GDP-mannose) are transported prior to polysaccharide polymerization (34–36). The polymerization of galactofuran was due to the action of a specific galactofuranosyltransferase, GfsA (37, 38). Two Ktr/Mnn2 mannosyltransferases have been recently identified to be essential to the mannan polymerization (31, 39). Interestingly, the absence of Ktr mannosyltransferases led to the absence of cell wall GM and to filamentous growth with an hyperbranched mycelium and a conidiation defect (31). The absence of cell wall cross-linked GM in Δdfg mutants could suggest a putative role of *Dfg* family members in mannan polymerization. However, our biochemical characterization showed that the *dfg* mutants still produced membrane-bound GM with the same chemical structure as that of the wild type (WT), showing that *Dfg* proteins are not involved in GM biosynthesis. Our study results show that these *DFG* proteins play a key role in the cross-linking of GM to β -(1,3)-glucans through an as-yet-undiscovered form of remodelling activity. The study also confirmed that all of the polysaccharide components of the cell wall skeleton [β -(1,3)-glucans, chitin, and galactomannan] are essential for fungal life whereas the alkali amorphous content has a nonstructural function and is mainly involved in the communications with the external milieu (32).

MATERIALS AND METHODS

Culture conditions. Parental and mutant strains of *A. fumigatus* were grown at 37°C in *Aspergillus* minimal medium (AMM; 1% glucose and 5 mM ammonium tartrate), Sabouraud medium (2% glucose, 1% Mycoseptone [Difco]), RPMI medium (Sigma), or 2% malt medium (Cristomalt). Media were either used in liquid form or supplemented with 2% agar. When necessary, 6% KCl was added to the media to enhance conidiation. Conidia were collected from agar medium plates after 10 days of growth at 37°C, using water containing 0.05% Tween 20.

Construction and complementation of the Δdfg mutants. The single-deletion and multiple-deletion mutants were constructed in the CEA17_ \DeltaakuB^{KU80} background (40) using the β -rec/six site-specific recombination system (13). The self-excising β -rec/six blaster cassette containing the hygromycin resistance marker was released from pSK529 plasmid using *FspI* restriction enzyme. The *dfg* replacement cassette containing the marker module flanked by 5' and 3' homologous regions of the target gene was generated by using primers listed in Table S3 in the supplemental material and cloned into the pUC19 vector using GeneArt seamless cloning and assembly (Life Technologies, Carlsbad, CA, USA). The corresponding replacement cassettes were released from the resulting vector via the use of either *EcoRV* or *FspI*, respectively. The CEA17 \DeltaakuB^{KU80} parental strain was transformed with the *dfg* replacement cassettes by electroporation to generate the *dfg* single-deletion mutants. Transformants were analyzed by PCR and Southern blotting using the digoxigenin (DIG) probe protocol (Roche Diagnostics) (see Fig. S2 in the supplemental material). For the construction of multiple-deletion strains, single-deletion mutants were cultivated in the presence of 2% xylose-containing minimal medium, which allows the excision of the selection marker by triggering recombination between the six recognition regions in the β -rec/six cassette. Because deletion of *DFG4* results in resistance to hygromycin, the sextuple-deletion mutant was constructed with the phleomycin resistance marker (41). The deletion cassette was made by fusion PCR with primers listed in Table S3 by using the double-joint PCR method.

Complementation of the $\Delta dfg3$ mutant was carried out by reintroduction of the parental copy of the gene flanked by the hygromycin resistance cassette and a 3' flanking region (Fig. S2; see also Table S3). The complementation cassette was transformed into the cassette-excised $\Delta dfg3$ mutant. The presence of the parental copy of the gene at the *DFG3* locus was confirmed by Southern blot analysis (Fig. S2). All mutants are detailed in Table S2.

Complementation of $Sc\Delta dcw1^{ts}/\Delta dfg5$ mutant by *AfDFG3*. *S. cerevisiae* BY230 strain was a derivative of S288c in which both *DCW1* and *DFG5* chromosomal loci were deleted. Since they are synthetically lethal, the strain therefore also carried p413TEF.*DCW1*, a *URA3* plasmid (42) containing the *DCW1* gene. The *DCW1* deletion in a $\Delta dfg5$ strain was obtained from the haploid BY4741 deletion collection. A *DCW1* deletion plasmid was generated by cloning 5' and 3' flanking regions of *DCW1* into Y1PLAC211 (43). The flanking regions were generated by PCR with primers 1975, 1976, 1977, and 1978 (Table S3). Following digestion with *MluI*, this plasmid was integrated at the *DCW1* locus of the $\Delta dfg5$ strain. This integrant was transformed with a CEN-ARS *HIS3* plasmid, p413TEF.*DCW1*, carrying the *DCW1* open reading frame (ORF) cloned as a BamH1-Xho1 fragment (generated with primers 2309 and 2310 and sequence verified). The genomic locus was then deleted by selection on plates with 5-fluoro-orotic acid followed by screening for the ORF deletion by PCR using flanking primers 2306 and 2307 to generate strain JT347 (*DCW1*/ $\Delta dfg5$). In strain JT346 (*dcw1^{ts}/dfg5), plasmid shuffling was used to replace the p413TEF.*DCW1* plasmid with p413TEF.*dcw1^{ts}*, a temperature-sensitive allele unable to support growth at 37°C (J. Trow and B. P. Cormack, unpublished data). Yeast strain JT346 (*dcw1^{ts}/dfg5*) was used for complementation performed with *AfDFG3*. Yeasts were grown at 30°C and 220 rpm in either a standard*

YEPD medium (10 g/liter yeast extract, 20 g/liter Bacto peptone, 20 g/liter glucose) or in YNB medium (1.7 g/liter yeast nitrogen base without amino acids and ammonium, 5 g/liter ammonium sulfate, 20 g/liter glucose) supplemented with the auxotrophic requirements.

The *AfDFG3* cDNA was synthesized by Life Technology SAS (St Aubin, France). A NotI restriction site and an XhoI restriction site were incorporated at the 5' and 3' ends of the cDNA, respectively, and cloned into pMA plasmid containing an ampicillin resistance marker. The *AfDFG3* cDNA was subcloned in plasmid pREP3-ADH (44) containing a *LEU2* gene to obtain pREP3-*AfDFG3* after digestion by NotI/XhoI. Then, the *dcw1^{ts}/Δdfg5* mutant was transformed with 3 μg of plasmid pREP3-*AfDFG3* following the lithium acetate method (45). As a control, the *dcw1^{ts}/Δdfg5* mutant was transformed with the pREP3-ADH plasmid alone. The transformants were selected on YNB plates without leucine, and the levels of expression of the *AfDFG3* genes in the *S. cerevisiae* mutant were checked by RT-PCR using primer pair *AfDFG3*comp1 and *AfDFG3*comp2 (Table S3). Phenotypes of the JT347 (*DCW1/Δdfg5*), JT346 (*dcw1^{ts}/Δdfg5*), and complemented (*dcw1^{ts}/Δdfg5::AfDFG3*) strains were analyzed on YNB medium at 30°C and 37°C.

Quantitative real-time PCR analysis. Fungal material was disrupted by the use of 0.5-mm-diameter glass beads in 500 μl of saturated phenol (Interchim, Montluçon, France) (pH 4.5), and RNA was isolated as described earlier (5) or by using a Qiagen RNeasy minikit. Quantitative PCR assays were performed as previously described (5). The expression ratios were normalized to *EF1α* expression levels and were calculated according to the $2^{-\Delta\Delta CT}$ (threshold cycle) method (46). The absence of genomic DNA contamination was verified with negative controls without reverse transcriptase. Three independent biological replicates were performed. The specificity of each primer (Table S3) was checked by agarose gel electrophoresis of RT-PCR products.

Phenotype of the Δdfg strains. Mycelial growth of the different strains was measured on 2% malt solid media after 48 h of incubation at 37°C. Agar pieces containing the entire colony of each of the strains were boiled 5 min in water to eliminate medium and extensively washed with hot water, and mycelial dry weight was recorded after overnight incubation at 80°C. The morphology of the mutant strains was determined during a kinetic study of the fungus grown in Sabouraud medium after staining of the fungus with calcofluor white (final concentration of 5 μg/ml). Conidia and mycelium were observed with a fluorescence microscope (Evos; Life Technologies) (excitation delta [λ ex], 357/44 nm; emission delta [λ em], 447/66 nm). To quantify growth, 24-h-old mycelial cultures that had been maintained at 37°C were filtered, washed, and dried overnight at 80°C and the dry weights were recorded. The conidiation rates were estimated following inoculation of conidial suspensions (100 μl, 10⁴/ml) into three tubes containing 2% malt agar (10 ml/tube) or malt agar containing 6% KCl. After 10 days at 25°C, conidia were recovered with aqueous 0.05% Tween 20 solution and counted using a hemocytometer.

The conidial permeability determinations were followed in the presence of FITC (47). Conidia were observed with a fluorescence microscope (Evos; Life Technologies) (λ ex, 470/22 nm; λ em, 510/42 nm).

The susceptibility of strains to antifungal and cell wall-disturbing compounds was estimated by spotting 10-fold serial dilutions of conidia (starting from 2 × 10⁶ spores) onto RPMI plates supplemented with the following drugs: itraconazole [50 ng/ml], voriconazole [100 ng/ml], amphotericin B [250 ng/ml], caspofungin [75 ng/ml] and calcofluor white [50 μg/ml]. Plates were incubated for 72 h at 37°C in a humid atmosphere.

Carbohydrate analysis of the cell wall, membrane fractions, and culture supernatant. Following 24 h of growth in Sabouraud liquid medium at 37°C and 150 rpm, mycelia and culture supernatants were separated by filtration. Macromolecules from the culture medium were precipitated by the use of three volumes of ethanol at 4°C overnight and collected by centrifugation (5 min, 4,000 × g). Cell walls and membranes were obtained after mycelium disruption and centrifugation (5, 16). Lipogalactomannan (LGM) was purified from each membrane preparation as previously described (16). The amount of galactomannan present in the culture filtrate was estimated using a Platelia kit as described by the manufacturer (Bio-Rad, Marnes la Coquette, France) with purified GM as the standard. Prior to the GM assay, proteins from the culture supernatant were eliminated by solid-phase extraction (SPE) on Sep-Pak classic C₁₈ cartridges (Waters) as previously described (48). Polysaccharides from the cell wall were separated based on their alkali solubility (5). Neutral hexoses and osamines were quantified by colorimetric and chromatographic assays (5, 49, 50). Proteins were quantified by use of the bicinchoninic acid (BCA) assay (Thermo Scientific), analyzed by SDS-PAGE, and identified by tandem mass spectrometry (MS/MS) analysis.

Galactomannan analysis. The analysis of glycosidic linkages in LGM was performed by methylation (51). The lipid anchor of LGM was released by mild acid hydrolysis (50 mM HCl, 100°C for 15 h), purified on a silica gel column, and analyzed by electrospray-mass spectrometry (16). Galactomannan (GM) amounts were estimated using a Platelia kit according to the instructions of the manufacturer (Bio-Rad, Marnes la Coquette, France) with purified GM as the standard. GM from the membrane preparation was solubilized with 2% Triton X-100. After trypsin digestion (1 mg trypsin for 10 mg of protein at 37°C for 24 h), GM was purified by SPE on Sep Pak classic C₁₈ cartridges and eluted with 5% propanol-1 (48). Mass spectrometry analysis of the lipid anchor was carried out on a Synapt G2Si instrument (Waters Corp., Milford, MA, USA). The source temperature was set to 80°C. The capillary and cone voltages were set to 1,500 and 40 V. Time of flight (TOF) data were collected between *m/z* 50 and 2,000, at a low level of collision energy (10 eV), in negative mode. Argon was used as the collision gas. Scans were collected for 1 s. An external calibration was done with clusters of NaI, and the mass range of calibration was *m/z* 50 to 2,000. Mass Lynx 4.1 was used for both acquisition and data processing. Samples were dissolved in chloroform/methanol (1/4 [vol/vol]) and introduced in nanoelectrospray mode via the use of a coated, medium-sized nano-electrospray ionization (ESI) capillary (Proxeon).

Fluorescence microscopy. Mycelia of the Δdfg and parental strains were fixed using *p*-formaldehyde (at a concentration of 2.5% in phosphate-buffered saline [PBS]) for one night at 4°C, washed three times with 0.1 M NH₄Cl–PBS and once with PBS, and then incubated with antigalactomannan antibody (52). Galactomannan was labeled with a rat anti-galactofuran monoclonal antibody (EBA2; a kind gift of M. Tabouret, Bio-Rad, Steenvorde) and a secondary FITC-conjugated goat anti-rat (Sigma) antibody. A monoclonal antibody with the same isotype was used as a negative control.

Statistical analysis. At least three biological replicates were performed per experiment; the statistical significance of the results was evaluated by one-way variance analysis using JMP1 software (SAS Institute, Cary, NC, USA).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mSphere.00397-19>.

TEXT S1, DOCX file, 0.1 MB.

FIG S1, PPT file, 0.4 MB.

FIG S2, PPT file, 1.2 MB.

FIG S3, PPTX file, 1.77 MB.

FIG S4, PDF file, 0.03 MB.

FIG S5, PDF file, 0.5 MB.

TABLE S1, DOCX file, 0.01 MB.

TABLE S2, DOCX file, 0.02 MB.

TABLE S3, DOCX file, 0.02 MB.

ACKNOWLEDGMENTS

We thank Magali Tichit for her precious involvement in histological experiments, Christian Malosse for MS experiments, and Catherine Comte and Christine Henry for their precious technical help.

This research was funded by the Aviesan project Aspergillus, the French Government's Investissement d'Avenir program, Laboratoire d'Excellence "Integrative Biology of Emerging Infectious Diseases" (grant ANR-10-LABX-62-IBEID), la Fondation pour la Recherche Médicale (DEQ20150331722 LATGE Equipe FRM 2015), and the NIH (5R21DE017085-02).

We declare that we have no conflict of interest.

REFERENCES

- Latgé J-P, Beauvais A, Chamilo G. 2017. The cell wall of the human fungal pathogen *Aspergillus fumigatus*: biosynthesis, organization, immune response, and virulence. *Annu Rev Microbiol* 71:99–116. <https://doi.org/10.1146/annurev-micro-030117-020406>.
- Gastebois A, Clavaud C, Aïmanianda V, Latgé J-P. 2009. *Aspergillus fumigatus*: cell wall polysaccharides, their biosynthesis and organization. *Future Microbiol* 4:583–595. <https://doi.org/10.2217/fmb.09.29>.
- Beauvais A, Bruneau JM, Mol PC, Buitrago MJ, Legrand R, Latgé JP. 2001. Glucan synthase complex of *Aspergillus fumigatus*. *J Bacteriol* 183:2273–2279. <https://doi.org/10.1128/JB.183.7.2273-2279.2001>.
- Henry C, Latgé J-P, Beauvais A. 2012. α 1,3 Glucans are dispensable in *Aspergillus fumigatus*. *Eukaryot Cell* 11:26–29. <https://doi.org/10.1128/EC.05270-11>.
- Muszkietta L, Aïmanianda V, Mellado E, Gribaldo S, Alcázar-Fuoli L, Szweczyk E, Prevost M-C, Latgé J-P. 2014. Deciphering the role of the chitin synthase families 1 and 2 in the in vivo and in vitro growth of *Aspergillus fumigatus* by multiple gene targeting deletion. *Cell Microbiol* 16:1784–1805. <https://doi.org/10.1111/cmi.12326>.
- Aïmanianda V, Simenel C, Garnaud C, Clavaud C, Tada R, Barbin L, Mouyna I, Heddergott C, Popolo L, Ohya Y, Delepierre M, Latgé J-P. 2017. The dual activity responsible for the elongation and branching of β -(1,3)-glucan in the fungal cell wall. *mBio* 8:e00619-17. <https://doi.org/10.1128/mBio.00619-17>.
- Percival-Smith A, Segall J. 1987. Increased copy number of the 5' end of the SPS2 gene inhibits sporulation of *Saccharomyces cerevisiae*. *Mol Cell Biol* 7:2484–2490. <https://doi.org/10.1128/mcb.7.7.2484>.
- Chabane S, Sarfati J, Ibrahim-Granet O, Du C, Schmidt C, Mouyna I, Prevost M-C, Calderone R, Latgé J-P. 2006. Glycosylphosphatidylinositol-anchored Ecm33p influences conidial cell wall biosynthesis in *Aspergillus fumigatus*. *Appl Environ Microbiol* 72:3259–3267. <https://doi.org/10.1128/AEM.72.5.3259-3267.2006>.
- Cabib E, Farkas V, Kosik O, Blanco N, Arroyo J, McPhie P. 2008. Assembly of the yeast cell wall Crh1p and Crh2p act as transglycosylases in vivo and in vitro. *J Biol Chem* 283:29859–29872. <https://doi.org/10.1074/jbc.M804274200>.
- Rodríguez-Peña JM, Cid VJ, Arroyo J, Nombela C. 2000. A novel family of cell wall-related proteins regulated differently during the yeast life cycle. *Mol Cell Biol* 20:3245–3255. <https://doi.org/10.1128/mcb.20.9.3245-3255.2000>.
- Kitagaki H, Wu H, Shimoi H, Ito K. 2002. Two homologous genes, DCW1 (YKL046c) and DFG5, are essential for cell growth and encode glycosylphosphatidylinositol (GPI)-anchored membrane proteins required for cell wall biogenesis in *Saccharomyces cerevisiae*. *Mol Microbiol* 46:1011–1022. <https://doi.org/10.1046/j.1365-2958.2002.03244.x>.
- Spreghini E, Davis DA, Subaran R, Kim M, Mitchell AP. 2003. Roles of *Candida albicans* Dfg5p and Dcw1p cell surface proteins in growth and hypha formation. *Eukaryot Cell* 2:746–755. <https://doi.org/10.1128/ec.2.4.746-755.2003>.
- Hartmann T, Dümig M, Jaber BM, Szweczyk E, Olbermann P, Morschhäuser J, Krappmann S. 2010. Validation of a self-excising marker in the human pathogen *Aspergillus fumigatus* by employing the beta-rec/six site-specific recombination system. *Appl Environ Microbiol* 76:6313–6317. <https://doi.org/10.1128/AEM.00882-10>.
- Latgé JP, Kobayashi H, Debeauvais JP, Diaquin M, Sarfati J, Wieruszkeski JM, Parra E, Bouchara JP, Fournet B. 1994. Chemical and immunological characterization of the extracellular galactomannan of *Aspergillus fumigatus*. *Infect Immun* 62:5424–5433.
- Fontaine T, Simenel C, Dubreucq G, Adam O, Delepierre M, Lemoine J,

- Vorgias CE, Diaquin M, Latgé JP. 2000. Molecular organization of the alkali-insoluble fraction of *Aspergillus fumigatus* cell wall. *J Biol Chem* 275:27594–27607. <https://doi.org/10.1074/jbc.M909975199>.
16. Costachel C, Coddeville B, Latgé J-P, Fontaine T. 2005. Glycosylphosphatidylinositol-anchored fungal polysaccharide in *Aspergillus fumigatus*. *J Biol Chem* 280:39835–39842. <https://doi.org/10.1074/jbc.M510163200>.
 17. Maddi A, Fu C, Free SJ. 2012. The *Neurospora crassa* *dfg5* and *dcw1* genes encode α -1,6-mannanases that function in the incorporation of glycoproteins into the cell wall. *PLoS One* 7:e38872. <https://doi.org/10.1371/journal.pone.0038872>.
 18. Mösch HU, Fink GR. 1997. Dissection of filamentous growth by transposon mutagenesis in *Saccharomyces cerevisiae*. *Genetics* 145:671–684.
 19. Kitagaki H, Ito K, Shimoi H. 2004. A temperature-sensitive *dcw1* mutant of *Saccharomyces cerevisiae* is cell cycle arrested with small buds which have aberrant cell walls. *Eukaryot Cell* 3:1297–1306. <https://doi.org/10.1128/EC.3.5.1297-1306.2004>.
 20. Levasseur A, Drula E, Lombard V, Coutinho PM, Henrissat B. 2013. Expansion of the enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes. *Biotechnol Biofuels* 6:41. <https://doi.org/10.1186/1754-6834-6-41>.
 21. Maruyama Y, Nakajima T. 2000. The *aman6* gene encoding a yeast mannan backbone degrading 1,6- α -D-mannanase in *Bacillus circulans*: cloning, sequence analysis, and expression. *Biosci Biotechnol Biochem* 64:2018–2020. <https://doi.org/10.1271/bbb.64.2018>.
 22. Nakajima T, Maitra SK, Ballou CE. 1976. An endo- α 1 leads to 6-D-mannanase from a soil bacterium. Purification, properties, and mode of action. *J Biol Chem* 251:174–181.
 23. Striebeck A, Robinson DA, Schüttelkopf AW, van Aalten DMF. 11 September 2013, posting date. Yeast *Mnn9* is both a priming glycosyltransferase and an allosteric activator of mannan biosynthesis. *Open Biol* <https://doi.org/10.1098/rsob.130022>.
 24. Gonzalez M, Goddard N, Hicks C, Ovalle R, Rauceo JM, Jue CK, Lipke PN. 2010. A screen for deficiencies in GPI-anchorage of wall glycoproteins in yeast. *Yeast* 27:583–596. <https://doi.org/10.1002/yea.1797>.
 25. Ao J, Chinnici JL, Maddi A, Free SJ. 2015. The N-linked outer chain mannans and the *Dfg5p* and *Dcw1p* endo- α -1,6-mannanases are needed for incorporation of *Candida albicans* glycoproteins into the cell wall. *Eukaryot Cell* 14:792–803. <https://doi.org/10.1128/EC.00032-15>.
 26. Rolli E, Ragni E, Calderon J, Porello S, Fascio U, Popolo L. 2009. Immobilization of the glycosylphosphatidylinositol-anchored *Gas1* protein into the chitin ring and septum is required for proper morphogenesis in yeast. *Mol Biol Cell* 20:4856–4870. <https://doi.org/10.1091/mbc.e08-11-1155>.
 27. Fu C, Sokolow E, Rupert CB, Free SJ. 2014. The *Neurospora crassa* CPS-1 polysaccharide synthase functions in cell wall biosynthesis. *Fungal Genet Biol* 69:23–30. <https://doi.org/10.1016/j.fgb.2014.05.009>.
 28. Maddi A, Free SJ. 2010. α -1,6-Mannosylation of N-linked oligosaccharide present on cell wall proteins is required for their incorporation into the cell wall in the filamentous fungus *Neurospora crassa*. *Eukaryot Cell* 9:1766–1775. <https://doi.org/10.1128/EC.00134-10>.
 29. Mouyna I, Kniemeyer O, Jank T, Lousset C, Mellado E, Aïmanianda V, Beauvais A, Wartenberg D, Sarfati J, Bayry J, Prévost M-C, Brakhage AA, Strahl S, Huerre M, Latgé J-P. 2010. Members of protein O-mannosyltransferase family in *Aspergillus fumigatus* differentially affect growth, morphogenesis and viability. *Mol Microbiol* 76:1205–1221. <https://doi.org/10.1111/j.1365-2958.2010.07164.x>.
 30. Beauvais A, Bozza S, Kniemeyer O, Formosa C, Formosa C, Balloy V, Henry C, Roberson RW, Dague E, Chignard M, Brakhage AA, Romani L, Latgé J-P. 2013. Deletion of the α -(1,3)-glucan synthase genes induces a restructuring of the conidial cell wall responsible for the avirulence of *Aspergillus fumigatus*. *PLoS Pathog* 9:e1003716. <https://doi.org/10.1371/journal.ppat.1003716>.
 31. Henry C, Li J, Danion F, Alcazar-Fuoli L, Mellado E, Beau R, Jouvion G, Latgé J-P, Fontaine T. 2019. Two KTR mannosyltransferases are responsible for the biosynthesis of cell wall mannans and control polarized growth in *Aspergillus fumigatus*. *mBio* 10:e02647-18. <https://doi.org/10.1128/mBio.02647-18>.
 32. Valsecchi I, Dupres Y, Michel JP, Duchateau M, Matondo M, Chamilos G, Saveanu C, Guijarro JI, Aïmanianda V, Lafont F, Latgé JP, Beauvais A. 2019. The puzzling construction of the conidial outer layer of *Aspergillus fumigatus*. *Cell Microbiol* 21:e12994. <https://doi.org/10.1111/cmi.12994>.
 33. Latgé J-P. 2009. Galactofuranose containing molecules in *Aspergillus fumigatus*. *Med Mycol* 47:S104–S109. <https://doi.org/10.1080/13693780802258832>.
 34. Engel J, Schmalhorst PS, Routier FH. 2012. Biosynthesis of the fungal cell wall polysaccharide galactomannan requires intraluminal GDP-mannose. *J Biol Chem* 287:44418–44424. <https://doi.org/10.1074/jbc.M112.398321>.
 35. Engel J, Schmalhorst PS, Dörk-Bousset T, Ferrières V, Routier FH. 2009. A single UDP-galactofuranose transporter is required for galactofuranosylation in *Aspergillus fumigatus*. *J Biol Chem* 284:33859–33868. <https://doi.org/10.1074/jbc.M109.070219>.
 36. Lamarre C, Beau R, Balloy V, Fontaine T, Hoi JWS, Guadagnini S, Berkova N, Chignard M, Beauvais A, Latgé J-P. 2009. Galactofuranose attenuates cellular adhesion of *Aspergillus fumigatus*. *Cell Microbiol* 11:1612–1623. <https://doi.org/10.1111/j.1462-5822.2009.01352.x>.
 37. Komachi Y, Hatakeyama S, Motomatsu H, Futagami T, Kizjakina K, Sobrado P, Ekino K, Takegawa K, Goto M, Nomura Y, Oka T. 2013. *gfsA* encodes a novel galactofuranosyltransferase involved in biosynthesis of galactofuranose antigen of O-glycan in *Aspergillus nidulans* and *A. fumigatus*. *Mol Microbiol* 90:1054–1073. <https://doi.org/10.1111/mmi.12416>.
 38. Katafuchi Y, Li Q, Tanaka Y, Shinozuka S, Kawamitsu Y, Izumi M, Ekino K, Mizuki K, Takegawa K, Shibata N, Goto M, Nomura Y, Ohta K, Oka T. 2017. *GfsA* is a β 1,5-galactofuranosyltransferase involved in the biosynthesis of the galactofuran side chain of fungal-type galactomannan in *Aspergillus fumigatus*. *Glycobiology* 27:568–581. <https://doi.org/10.1093/glycob/cwx028>.
 39. Onoue T, Tanaka Y, Hagiwara D, Ekino K, Watanabe A, Ohta K, Kamei K, Shibata N, Goto M, Oka T. 2018. Identification of two mannosyltransferases contributing to biosynthesis of the fungal-type galactomannan α -core-mannan structure in *Aspergillus fumigatus*. *Sci Rep* 8:16918. <https://doi.org/10.1038/s41598-018-35059-2>.
 40. da Silva Ferreira ME, Kress M, Savoldi M, Goldman MHS, Härtl A, Heinekamp T, Brakhage AA, Goldman GH. 2006. The *akuBKU80* mutant deficient for nonhomologous end joining is a powerful tool for analyzing pathogenicity in *Aspergillus fumigatus*. *Eukaryot Cell* 5:207–211. <https://doi.org/10.1128/EC.5.1.207-211.2006>.
 41. Mattern I, Punt P, van den Hondel C. 1988. A vector for *Aspergillus* transformation conferring phleomycin resistance. *Fungal Genetics Rep* 35:25. <https://doi.org/10.4148/1941-4765.1533>.
 42. Mumberg D, Müller R, Funk M. 1995. Yeast vectors for the controlled expression of heterologous proteins in different genetic backgrounds. *Gene* 156:119–122. [https://doi.org/10.1016/0378-1119\(95\)00037-7](https://doi.org/10.1016/0378-1119(95)00037-7).
 43. Gietz RD, Akio S. 1988. New yeast-*Escherichia coli* shuttle vectors constructed with in vitro mutagenized yeast genes lacking six-base pair restriction sites. *Gene* 74:527–534. [https://doi.org/10.1016/0378-1119\(88\)90185-0](https://doi.org/10.1016/0378-1119(88)90185-0).
 44. Willer T, Brandl M, Sipiczki M, Strahl S. 2005. Protein O-mannosylation is crucial for cell wall integrity, septation and viability in fission yeast. *Mol Microbiol* 57:156–170. <https://doi.org/10.1111/j.1365-2958.2005.04692.x>.
 45. Ito H, Fukuda Y, Murata K, Kimura A. 1983. Transformation of intact yeast cells treated with alkali cations. *J Bacteriol* 153:163–168.
 46. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25:402–408. <https://doi.org/10.1006/meth.2001.1262>.
 47. Lambou K, Lamarre C, Beau R, Dufour N, Latgé J-P. 2010. Functional analysis of the superoxide dismutase family in *Aspergillus fumigatus*. *Mol Microbiol* 75:910–923. <https://doi.org/10.1111/j.1365-2958.2009.07024.x>.
 48. Morelle W, Bernard M, Debeaupuis J-P, Buitrago M, Tabouret M, Latgé J-P. 2005. Galactomannoproteins of *Aspergillus fumigatus*. *Eukaryot Cell* 4:1308–1316. <https://doi.org/10.1128/EC.4.7.1308-1316.2005>.
 49. DuBois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. 1956. Colorimetric method for determination of sugars and related substances. *Anal Chem* 28:350–356. <https://doi.org/10.1021/ac60111a017>.
 50. Stalhaberger T, Simenel C, Clavaud C, Eijssink VGH, Jourdain R, Delepierre M, Latgé J-P, Breton L, Fontaine T. 2014. Chemical organization of the cell wall polysaccharide core of *Malessezia restricta*. *J Biol Chem* 289:12647–12656. <https://doi.org/10.1074/jbc.M113.547034>.
 51. Fontaine T, Talmont F, Dutton GG, Fournet B. 1991. Analysis of pyruvic acid acetal containing polysaccharides by methanolysis and reductive cleavage methods. *Anal Biochem* 199:154–161. [https://doi.org/10.1016/0003-2697\(91\)90083-6](https://doi.org/10.1016/0003-2697(91)90083-6).
 52. Bayry J, Beauvais A, Dufréne YF, Sharma M, Bansal K, Kniemeyer O, Aïmanianda V, Brakhage AA, Kaveri SV, Kwon-Chung KJ, Latgé J-P, Beauvais A. 2014. Surface structure characterization of *Aspergillus fumigatus* conidia mutated in the melanin synthesis pathway and their

- human cellular immune response. *Infect Immun* 82:3141–3153. <https://doi.org/10.1128/IAI.01726-14>.
53. Mouyna I, Fontaine T, Vai M, Monod M, Fonzi WA, Diaquin M, Popolo L, Hartland RP, Latgé JP. 2000. Glycosylphosphatidylinositol-anchored glucanoyltransferases play an active role in the biosynthesis of the fungal cell wall. *J Biol Chem* 275:14882–14889. <https://doi.org/10.1074/jbc.275.20.14882>.
54. Gastebois A, Fontaine T, Latgé J-P, Mouyna I. 2010. β (1-3)Glucanoyltransferase Gel4p is essential for *Aspergillus fumigatus*. *Eukaryot Cell* 9:1294–1298. <https://doi.org/10.1128/EC.00107-10>.
55. Ragni E, Fontaine T, Gissi C, Latgé JP, Popolo L. 2007. The Gas family of proteins of *Saccharomyces cerevisiae*: characterization and evolutionary analysis. *Yeast* 24:297–308. <https://doi.org/10.1002/yea.1473>.