



HAL
open science

On a Phase Transition of Regret in Linear Quadratic Control: The Memoryless Case

Ingvar Ziemann, Henrik Sandberg

► **To cite this version:**

Ingvar Ziemann, Henrik Sandberg. On a Phase Transition of Regret in Linear Quadratic Control: The Memoryless Case. 2020. hal-02546670v2

HAL Id: hal-02546670

<https://hal.science/hal-02546670v2>

Preprint submitted on 13 May 2020 (v2), last revised 10 Sep 2020 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On a Phase Transition of Regret in Linear Quadratic Control: The Memoryless Case

Ingvar Ziemann, Henrik Sandberg

Abstract— We consider an idealized version of adaptive control of a MIMO system without state. We demonstrate how rank deficient Fisher information in this simple memoryless problem leads to the impossibility of logarithmic rates of regret. This speaks to an open issue concerning the attainability of logarithmic regret rates in linear quadratic adaptive control. Our analysis rests on a version of the Cramér-Rao inequality that takes into account possible ill-conditioning of Fisher information and a perturbation result on the corresponding singular subspaces. This is used to define a sufficient condition, which we term *uniformativeness*, for regret to be at least order square root in the samples.

I. INTRODUCTION

Recently, there has been a revitalization of interest in the adaptive linear quadratic regulator (LQR) as it serves as good theoretically tractable example of reinforcement learning in continuous state and action spaces, [1]. Much progress has been made toward analyzing the statistical convergence rate, the *regret* incurred, of adaptive algorithms. Several works over the past decade, [2], [3] and [4], have been able to prove *upper bounds* on the regret at a rate of approximately \sqrt{T} in the time horizon. However, in some special cases, [5], [6] [7] and [8], the authors have actually been able to prove regret to scale at a rate of $\log T$, ensuring considerably faster convergence. In particular, [6] and [7] show that, suitably modified, the Åström-Wittenmark self-tuning regulator [9] for SISO tracking problems converges at the rate $\log T$. Given these two very different rates, it is thus natural to ask whether regret undergoes a phase transition in its asymptotic scaling. Here, we consider a simplified, and memoryless, version of the linear quadratic problem to verify that such a phenomenon indeed occurs. The point of such an analysis, as presented here, is to isolate the essence of this phase transition. With this in mind, Our goal is to identify and give conditions for when the *lower bound changes* from order $\log T$ to \sqrt{T} .

A. Contribution

As hinted above, the main contribution of this note is to establish a sufficient condition, *uniformativeness*, for regret to necessarily scale on the order \sqrt{T} , Definition 4.4 and Theorem 5.2. We will see that this phase transition depends both on the rank of the optimal linear feedback matrix and

Ingvar Ziemann (ziemann@kth.se) and Henrik Sandberg (hsan@kth.se) are with the School of Electrical Engineering and Computer Science, KTH Royal Institute of Technology, SE-100 44 Stockholm, Sweden.

This work was supported in part by the Swedish Research Council (grant 2016-00861), and the Swedish Foundation for Strategic Research (Project CLAS). The authors are also indebted to three anonymous referees for their helpful comments.

on the excitation of the reference signal. In the uninformative regime, there is an asymptotically non-negligible trade-off between exploration and exploitation. These results partially answer an unresolved question in the literature [10], [11], as to precisely when logarithmic rates are attainable for linear quadratic problems by showing when they are not. On the theoretical side, the method of analysis to provide lower bounds here is novel and rests on comparing the Fisher information of any policy to the Fisher information of the optimal policy having knowledge of the system's parameters. This clarifies some connections between regret and parameter estimation ([10], [12]) via the spectral properties of the Fisher information (matrix). We believe this proof strategy to potentially be of general interest. Moreover, we are among the first to record a \sqrt{T} lower bound for linear quadratic adaptive control problems.

B. Recent Related Work

Indeed, we were only after the initial submission of this manuscript made aware of the results by [13] and [14] which independently arrive at \sqrt{T} lower bounds for similar problems. It is particularly interesting to compare our work to [13] which, as our work, contains an instance specific bound. There, it is shown that regret for an unknown LQR scales as $\dim(\text{state}) \times \dim(\text{inputs})^2 \times \sqrt{T}$. By contrast, we consider a memoryless tracking problem where $\dim(\text{state}) = 0$ and the above results are not applicable. In particular, when there is a reference signal to track, we show that regret can be of order \sqrt{T} even if the dimension of the state is zero. We also remark that our proof is very different from those in [13] and [14] and has the appealing property of generating lower bounds directly depending on an Information quantity, Fisher information.

C. Notation

We use \succeq (and \succ) for (strict) inequality in the matrix positive definite partial order. By $\|\cdot\|$ we denote the standard 2-norm and by $\|\cdot\|_\infty$ the matrix operator norm. Moreover, \otimes , vec and \dagger are used to denote the Kronecker product, vectorization (mapping a matrix into a column vector), and Moore-penrose pseudoinverse, respectively. For two functions f, g , $\limsup |f(t)/g(t)| = 0$, for some norm $|\cdot|$, is written as $f = o(g)$. If instead $\limsup |f(t)/g(t)| \leq C, C > 0$, we write $f = O(g)$. For asymptotic lower bounds we write $f = \Omega(g)$ which means that $\liminf f(t)/g(t) \geq c, c > 0$. In general, these limits will be for large times, usually indexed by t or T . We also write \mathbf{E} for the expectation operator. We use ∇ for gradient or Jacobian. All vectors, including gradients, in this paper are column-vectors.

II. PROBLEM FORMULATION

We consider the memoryless adaptive control problem

$$\begin{cases} \min_{(u_t)} & \sum_{t=1}^T \mathbf{E} \|r_t - y_t\|^2 + \lambda \mathbf{E} \|u_t\|^2, \\ \text{s. t.} & y_t = Bu_t + w_t, \end{cases} \quad (1)$$

where $y_t, r_t, w_t \in \mathbb{R}^n$, $u_t \in \mathbb{R}^m$ and $\lambda > 0$. $B \in \mathbb{R}^{n \times m}$ is assumed unknown in advance. Note that since $\lambda > 0$, a unique solution to (1) of the form $u_t = Kr_t$ always exists, $K = (B^\top B + \lambda I)^{-1} B^\top \in \mathbb{R}^{m \times n}$. Our goal is to investigate whether depending on the parametrization, there may be phase transition in the regret – learning-based performance – any algorithm can attain. We are able to prove that there are two regimes for regret (defined below): one in which regret scales like $\log T$ and one in which it scales like \sqrt{T} . To this end, our Theorem 5.2 gives regret lower bounds for (1) and a sufficient condition for the \sqrt{T} -scaling limit to occur. This is then contrasted with Theorem 5.4 which gives a logarithmic lower bound valid in both regimes.

To qualify this, some further assumptions are necessary. We suppose B has rank at least 1. We write $K = K(B, \lambda) \in \mathbb{R}^{m \times n}$ for the optimal linear feedback law and its Jacobian is $G = \nabla_B \text{vec} K(B, \lambda)$. The reference signal, r_t , is assumed to be known in advance and we will make a standard persistence of excitation assumption, namely that $\sum_{k=1}^t r_k r_k^\top \succ tcI + o(t)$ and that $\|r_t\| > c'$ for some $c, c' > 0$ and sufficiently large t . The noise $w_t \in \mathbb{R}^n$ is assumed to be mean zero, independent and identically distributed, and have density $q(w)$, which admits Fisher information¹. The control $u_t \in \mathbb{R}^m$ is constrained to depend on only past inputs and outputs and is in particular oblivious of the parameter B – it is adaptive. To compare adaptive laws to the optimal law, one introduces the regret.

Definition 2.1: The regret of u_t , $R_T = R_T(\{u_t\})$ is

$$\begin{aligned} R_T &= \sum_{t=1}^T \mathbf{E} \|r_t - Bu_t\|^2 - \sum_{t=1}^T \mathbf{E} \|r_t - BKr_t\|^2 \\ &\quad + \lambda \mathbf{E} \sum_{t=1}^T \|u_t\|^2 - \lambda \mathbf{E} \sum_{t=1}^T \|Kr_t\|^2. \end{aligned} \quad (2)$$

This measures the cumulative difference between the cost incurred by the adaptive law (u_t) and the optimal law Kr_t which uses knowledge of B .

Example 2.2: Consider a scalar system $y_t = bu_t + w_t$ of with variance 1 of w_t . Suppose that r_t is sufficiently rich, say $r_t = 1$ for all time and that $\lambda = 0$. This case is then covered by [6] and [7] (by setting all lag parameters of y to zero), where it is shown that

$$R_T = O(\log T)$$

for a policy based on least squares and certainty equivalence.

¹The density of w_t needs to satisfy certain absolute continuity and mean square differentiability conditions. We prefer not to go into these details and simply assume existence, see Definition 4.1 and consult [15] for details.

III. REGRET DECOMPOSITION

The following result is key, as it directly relates regret to an estimation error.

Lemma 3.1: For any linear policy $u_t = \hat{K}_t r_t$, we have

$$\begin{aligned} R_T &= \sum_{t=1}^T \mathbf{E} \text{tr} \left[\left((I \otimes r_t r_t^\top) + \lambda (I \otimes B)^\top (I \otimes r_t r_t^\top) (I \otimes B) \right) \right. \\ &\quad \left. \times \text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top \right]. \end{aligned} \quad (3)$$

Proof: Setting $u_t = \hat{K}_t r_t$, we see that K also is for each t the minimizer of

$$\|r_t - B\hat{K}_t r_t\|^2 - \|r_t - BKr_t\|^2 + \lambda \left(\|\hat{K}_t r_t\|^2 - \|Kr_t\|^2 \right).$$

Vectorizing, we observe that this is a quadratic expression in $\text{vec} \hat{K}_t$, minimized at $\text{vec} K$, where its value is zero. A straightforward computation shows that the Hessian in vectorized variables is

$$2(I \otimes r_t r_t^\top) + 2\lambda (I \otimes B)^\top (I \otimes r_t r_t^\top) (I \otimes B)$$

Since there are no higher order terms, Taylor expansion around the minimum \vec{K} gives

$$\begin{aligned} \|r_t - B\hat{K}_t r_t\|^2 - \|r_t - BKr_t\|^2 + \lambda \left(\|\hat{K}_t r_t\|^2 - \|Kr_t\|^2 \right) \\ = \text{tr} \left[\left((I \otimes r_t r_t^\top) + \lambda (I \otimes B)^\top (I \otimes r_t r_t^\top) (I \otimes B) \right) \right. \\ \left. \times \text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top \right]. \end{aligned}$$

The result follows by summation and expectation. \blacksquare

This shows that regret is linear in the estimation error, $\text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top$. To relate this to any particular policy, we make the following definitions.

Definition 3.2: The control sequence (u_t) is α -fast convergent if for all B , $u_t = Kr_t + v_t$ with $v_t = o(t^{-\alpha})$ in L^2 .

In particular, this prohibits constant strategies such as selecting K which is optimal for one parametrization but sub-optimal for others. Note also that this definition relies implicitly on the uniqueness of the solution to (1).

Lemma 3.3: Any α -fast convergent policy can be written as $u_t = \hat{K}_t r_t$, for some sequence of matrices $\hat{K}_t = K + o(t^{-\alpha})$.

Proof: Since by assumption $\|r_t\| > c'$ uniformly in time for large t and some constant c' , there exists a linear transformation V_t such that $v_t = V_t r_t$ and $\|V_t\| = o(t^{-\alpha}) / \|r_t\| = o(t^{-\alpha})$. Take $\hat{K}_t = K + V_t$. \blacksquare

Lemma 3.3 shows that for α -fast convergent policies, it suffices to consider linear representations. In the subsequent analysis, we will also need some asymptotic control of the gradient of these linear representations, with respect to the parameter, B .

Definition 3.4: An α -fast convergent policy, $u_t = \hat{K}_t r_t$ is regular if $\nabla_B \mathbf{E} \text{vec} \hat{K}_t = \nabla_B \text{vec} K + o(1)$.

IV. INFORMATION

As indicated by the regret decomposition, Lemma 3.1, our regret analysis will essentially be estimation-theoretic. The following notion is key in the Cramér-Rao bound we will use.

Definition 4.1: For a parametrized family probability densities $\{p_\theta, \theta \in \Theta\}$, $\Theta \subset \mathbb{R}^d$, Fisher information I_θ is

$$I_\theta = \int \nabla_\theta \log p_\theta(x) [\nabla_\theta \log p_\theta(x)]^\top p_\theta(x) dx$$

whenever the integral exists.

Since the density, $q(w)$, of the noise in 1 was assumed sufficiently regular, for the parameter $\theta = B$, the location model induced by $y_k = Bu_k + w_k, k = 1, \dots, t$ admits Fisher information, $I_{t,B}$. Let $J = \int q(w) [\nabla_w q(w)] [\nabla_w q(w)]^\top dw$, then one has that

$$\begin{aligned} I_{t,B} &= \sum_{k=1}^t \int q(y - Bu_k) \\ &\quad \times \nabla_B \log q(y - Bu_k) [\nabla_B \log q(y - Bu_k)]^\top dy \\ &= \sum_{k=1}^t u_k u_k^\top \otimes J, \quad (4) \end{aligned}$$

by the chain rule for Fisher information and change of variables. We note in passing that J is often called Fisher information about location parameter for q . If in addition u_t is α -fast convergent (4) becomes

$$\begin{aligned} I_{t,B} &= \sum_{k=1}^t \mathbf{E} (K r_k r_k^\top K^\top + v_k v_k^\top) \otimes J \\ &= \underbrace{\sum_{k=1}^t \mathbf{E} [K r_k r_k^\top K^\top \otimes J]}_{I_{t,B}^*} + \sum_{k=1}^t \mathbf{E} v_k v_k^\top \otimes J. \quad (5) \end{aligned}$$

where $v_k = o(k^{-\alpha})$ due to α -fast convergence. Above, one recognizes $I_{t,B}^*$ as the Fisher information generated by the optimal trajectory, where $v_k \equiv 0$ and the optimal law is always applied. Observe that unless K has rank m , $I_{t,B}^*$ is degenerate for all t . A degenerate Fisher information has bleak implications for model identifiability. Following the analysis of [16], no finite variance estimator with a given bias exists except for very special circumstances for K if $I_{t,K}$ is degenerate. Fortunately, the term $\sum_{k=1}^t \mathbf{E} v_k v_k^\top \otimes J$ may be chosen to complete rank-deficiency. However, all is not won, since the requirement that u_t is α -fast convergence entails that this term is small.

A. A Multi-Scale Cramér-Rao Bound

We now repeat the general Cramér-Rao analysis of [16]. As we require a minor modification of their result, we state the result and proof in full.

Theorem 4.2 (cf. [16]): Let $\{p_\theta, \theta \in \Theta \subset \mathbb{R}^d\}$, be a family of densities with Fisher information I_θ . Suppose that $\alpha = \alpha(\theta) \in \mathbb{R}^{d'}$ is a vector-valued function of θ and let $\hat{\alpha}$ be any estimate of α . Let I_θ have singular value decomposition

$$I_\theta = Z S Z^\top = \begin{bmatrix} Z_1 & Z_2 \end{bmatrix} \begin{bmatrix} S_1 & 0 \\ 0 & S_2 \end{bmatrix} \begin{bmatrix} Z_1^\top \\ Z_2^\top \end{bmatrix}.$$

Suppose $A = \nabla_\theta \mathbf{E} \hat{\alpha}$ exists and set $\begin{bmatrix} A_1 & A_2 \end{bmatrix} = A \begin{bmatrix} Z_1 & Z_2 \end{bmatrix}$. Then one has that

$$\mathbf{E}[(\hat{\alpha} - \alpha)(\hat{\alpha} - \alpha)^\top] \succeq A_1 S_1^\dagger A_1^\top + A_2 S_2^\dagger A_2^\top. \quad (6)$$

Moreover, if $A \neq A I_\theta I_\theta^\dagger$, the trace of the left hand side in (6) can be made arbitrarily large.

Proof: Let $\Delta = \nabla_\theta \log p_\theta(Y)$ where $Y \sim p_\theta$, $W \in \mathbb{R}^{d \times d'}$ and define

$$\begin{bmatrix} W_1 \\ W_2 \end{bmatrix} = \begin{bmatrix} Z_1^\top \\ Z_2^\top \end{bmatrix} W.$$

Let also $C = \mathbf{E}[(\hat{\alpha} - \alpha)(\hat{\alpha} - \alpha)^\top]$. We have that

$$\begin{aligned} &\mathbf{E} \left[(\hat{\alpha} - \alpha) - W^\top \Delta \right] \left[(\hat{\alpha} - \alpha) - W^\top \Delta \right]^\top \\ &= C - A W - W^\top A + W^\top I_\theta W \succeq 0 \quad (7) \end{aligned}$$

since the score, Δ , has mean zero. We now partition I_θ, W and A as assumed. Substituting this decomposition into (7) yields

$$\begin{aligned} C &\succeq A W + W^\top A - W^\top I_{t,B} W \\ &= (A_1 W_1 + A_2 W_2) + (A_1 W_1 + A_2 W_2)^\top \\ &\quad - (W_1^\top S_1 W_1) - (W_2^\top S_2 W_2) \\ &= A_1 S_1^\dagger A_1 - (W_1 - S_1^\dagger A_1^\top)^\top \Lambda_1 (W_1 - S_1^\dagger A_1^\top) \\ &\quad + A_2 S_2^\dagger A_2 - (W_2 - S_2^\dagger A_2^\top)^\top S_2 (W_2 - S_2^\dagger A_2^\top). \quad (8) \end{aligned}$$

Choosing $W_i = S_i^\dagger A_i^\top, i = 1, 2$ gives

$$C \succeq A_1 S_1^\dagger A_1^\top + A_2 S_2^\dagger A_2^\top$$

which was the first part our claim.

Now, if $A \neq A I_\theta I_\theta^\dagger$, it is clear that I_θ is rank-deficient, and we may take $S_2 = 0$, and Z_2 to be the basis of the corresponding kernel. Further, one may choose $W_2 = \sigma A_2^\top$, for arbitrary $\sigma > 0$. In this case,

$$\text{tr } C \geq \text{tr} \left(A_1 S_1^\dagger A_1^\top + \sigma A_2 A_2^\top \right)$$

which can be arbitrarily large by increasing σ . ■

For our model (1), the following consequence is immediate.

Corollary 4.3: Let \hat{K}_t be any estimator of the optimal policy K of (1) such that $H = \nabla_B \text{vec } \hat{K}_t(B, \lambda)$ exists. Decompose Fisher information $I_{t,B}$ (5) and define H_1, H_2 via

$$I_{t,B} = U \Lambda U^\top = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Lambda_1 & 0 \\ 0 & \Lambda_2 \end{bmatrix} \begin{bmatrix} U_1^\top \\ U_2^\top \end{bmatrix} \quad (9)$$

$$\begin{bmatrix} H_1 & H_2 \end{bmatrix} = H \begin{bmatrix} U_1 & U_2 \end{bmatrix} \quad (10)$$

Then

$$\begin{aligned} &\mathbf{E} \text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top \\ &\quad \succeq H_1 \Lambda_1^{-1} H_1^\top + H_2 \Lambda_2^\dagger H_2^\top. \quad (11) \end{aligned}$$

The notation (9-10) is fixed throughout the rest of the paper.

B. Uninformative Optimal Policies

Observe that the regret decomposition (3) shows that regret depends fundamentally on

$$= \text{tr} \left((I \otimes r_t r_t^\top) \mathbf{E} \text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top \right).$$

Since the estimation error is lower-bounded by (11), degeneracy of Fisher information should be a factor in regret if for the small block, say Λ_2 ,

$$\text{tr} \left((I \otimes r_t r_t^\top) H_2 \Lambda_2^\dagger H_2^\top \right) \neq 0. \quad (12)$$

Now, the Fisher-information, $I_{t,B}$, of an α -fast convergent policy one imagines is close to that of the optimal policy, $I_{t,B}^*$. Apply now instead the Spectral theorem to $I_{t,B}^*$ and assume B does not have full column rank. That is, let

$$I_{t,B}^* = O\Lambda'O^\top = \begin{bmatrix} O_1 & O_2 \end{bmatrix} \begin{bmatrix} \Lambda_1' & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} O_1^\top \\ O_2^\top \end{bmatrix}. \quad (13)$$

Recall that $G = \nabla_B \text{vec } K(B, \lambda)$. This is the Jacobian of the optimal policy with respect to the parameter B . Define $G_2 = GO_2$. We expect regret to scale differently in the following regime.

Definition 4.4: The pair (r_t, B) is said to be γ -uninformative if B does not have full column rank and

$$\text{tr}((I \otimes r_t r_t^\top) G_2 G_2^\top) \geq \gamma \quad (14)$$

for $\gamma > 0$, $G_2 = GO_2$ where $G = \nabla_B \text{vec } K(B, \lambda)$ and O_2 defined through (13).

This definition is meaningful when $I_{t,B}^*$ of (5) does not have full rank and is best understood as a quantitative observability-type condition; the kernel of the optimal policy's information matrix should be visible through the cost (regret). Note also the implicit dependence on B , via G_2 which depends both on the subspace $\ker I_{t,B}^*$ and on the optimal policy K . Geometrically, γ -uninformativeness means that $\ker I_{t,B}^*$ (5), which is spanned by O_2 and $G_2 = GO_2$, is γ -separated from being perpendicular to the reference signal's power, $r_t r_t^\top$, in the geometry of symmetric matrices. A rank drop in $I_{t,B}^*$ of (5) can occur when $\text{rank } K = \text{rank } B < m$. Note that [5] establishes logarithmic rates for a related problem precisely when B is invertible, i.e. full rank. We now give an example illustrating this.

Example 4.5: Suppose that y_t is a scalar given by $y_t = b^\top u_t + w_t$ with $b, u_t \in \mathbb{R}^m$ and b given by the first standard Euclidean basis vector

$$b^\top = [1 \quad 0 \quad \dots \quad 0].$$

In this case the optimal policy is given by br_t and so differentiating $\mathbf{E}\hat{K}_t = b + o(1)$ with respect to b yields $H = I + o(1)$.

Let us also assume that the variance of w_t is 1 and choose r_t such that

$$\sum_{k=1}^t r_k r_k^\top = tI + o(1),$$

This happens if r_k is a sequence of standard Euclidean basis vector e_i cyclically repeating themselves, so that $r_1 = e_1, \dots, r_m = e_m, r_{m+1} = e_1, \dots$ and so on. In this case

$$I_{t,B}^* = t \begin{bmatrix} 1 & 0_{1 \times m-1} \\ 0_{m-1 \times 1} & 0_{m-1 \times m-1} \end{bmatrix} + o(1)$$

and for large $t \notin \{t : r_t = e_1\}$ we have, since $G = I$, that

$$\begin{aligned} \text{tr}(r_t r_t^\top G_2 G_2^\top) &= \text{tr} \left(r_t r_t^\top \begin{bmatrix} 0 & 0_{1 \times m-1} \\ 0_{m-1 \times 1} & I_{m-1 \times m-1} \end{bmatrix} \right) + o(1) \\ &= 1 + o(1). \end{aligned}$$

C. Spectral Information Comparison

The main reason for using G_2 (13) instead of H_2 (10), in Definition 4.4 is that we wish to study how the parametrization of (1) impacts regret. If we had defined uninformativeness in terms of the H_2 , which depends on the choice of algorithm through $H = \nabla_B \text{vec } \hat{K}_t(B, \lambda)$, it would be hard to claim that a phase transition occurs through the parametrization and our lower bound would not be algorithm independent. Of course, this entails that we need to control the gap between the system quantity (14) and the algorithm-dependent quantity (12). To this end, we perform a perturbation analysis of the subspace spanned by G_2 , to relate it to H_2 .

Lemma 4.6: Let H_2 be as in (10) and G_2 be as in (13). Then for any regular and α -fast convergent policy \hat{K}_t , one has that

$$H_2 P_t = G_2 + o(1)$$

for some sequence of orthonormal matrices (P_t) .

Proof: The proof relies on Wedin's $\sin \Theta$ Theorem [17] (quoted in the appendix) which describes the perturbation theory of range and nullspace in the singular value decomposition. By (5) it follows that

$$\frac{1}{t} I_{t,B} = \frac{1}{t} I_{t,B}^* + o(t^{-2\alpha} \log t)$$

using α -fast convergence². Moreover, it is clear that rescaling Fisher informations does not change the singular value decomposition except for rescaled singular values. This gives control of the residuals (1.8) in [17] and for sufficiently large t the separation conditions there are satisfied since $(1/t)\Lambda_2 = o(1)$. We recall also from Lemma 1 of [18] that the $\sin \Theta$ Theorem provides an upper bound on the distance, $d(U_2, O_2) = \min_P \|U_2 P - O_2\|_\infty$, where P is optimized over the orthogonal group. Apply now Wedin's Theorem in combination with Lemma 1 of [18] to conclude that

$$U_2 P_t = O_2 + o(t^{-2\alpha})$$

where P_t optimizes $d(U_2, O_2)$. Finally, by regularity $H = G + o(1)$, and therefore $H_2 P_t \rightarrow G_2$. ■

Remark 4.7: The matrix P_t is necessary to account for the possibly arbitrary ordering of the singular vectors corresponding to $\ker I_{t,B}^*$. It exists by continuity and compactness of the orthogonal group in the standard matrix topology.

V. FUNDAMENTAL LIMITATIONS

We need one more lemma which relates regret to the spectral properties of $I_{t,B}$ and $I_{t,B}^*$.

Lemma 5.1: For any γ -uninformative pair (r_t, B) , and regular and α -fast convergent policy \hat{K}_t with Λ_2 as in (9) and H_2 as in (10), one has that

$$\text{tr} \left((I \otimes r_t r_t^\top) H_2 \Lambda_2^\dagger H_2^\top \right) \geq \gamma \sigma_{\min}(\Lambda_2^\dagger) \times (1 + o(1)).$$

²the appearance of the logarithmic factor is due to the case $\alpha = 1/2$, since $\int t^{-1} \sim \log t \neq O(1)$.

Proof: The trace cyclic property and Lemma 4.6 yields

$$\begin{aligned} & \text{tr} \left((I \otimes r_t r_t^\top) H_2 H_2^\top \right) \\ &= \text{tr} \left((I \otimes r_t r_t^\top) H_2 P_t P_t^\top H_2^\top \right) \\ &= \text{tr} \left(P_t^\top H_2^\top (I \otimes r_t r_t^\top) H_2 P_t \right) \\ &= \text{tr} \left(G_2^\top (I \otimes r_t r_t^\top) G_2 \right) + o(1) \end{aligned}$$

using that P_t from Lemma 4.6 is orthonormal and where G_2 is as in (13). Multiplication by $\Lambda_2^\dagger \succeq 0$ rescales the bound by $\sigma_{\min}(\Lambda_2^\dagger)$. ■

We are now in position to state our main result.

Theorem 5.2: Consider the model (1) and suppose that (r_t, B) is γ -uninformative for at least a constant fraction of indices $t \in \mathbb{N}$, for some $\gamma > 0$. Then for any $\alpha > 0$, any regular α -fast convergent policy has regret asymptotically lower bounded as

$$R_T = \Omega(\sqrt{T}). \quad (15)$$

The central idea of the following proof is to balance the ill-conditioning of Fisher information with the convergence rate, $t^{-\alpha}$, of the estimated policy \hat{K}_t .

Proof: According to Lemma 3.3 we may restrict our attention to linear policies. Let Λ_2 as in (9) and to emphasize its time-dependence, we now write $\Lambda_2 = \Lambda_2(t)$. Let also H_2 be as in (10). Combining the results that we have established this far offers

$$\begin{aligned} R_T &\geq \sum_{t=1}^T \mathbf{E} \text{tr} \left[\left((I \otimes r_t r_t^\top) \text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top \right) \right] \\ &\geq \sum_{t=1}^T \left[\text{tr} \left((I \otimes r_t r_t^\top) H_2 \Lambda_2^\dagger(t) H_2^\top \right) \right] \\ &\geq \sum_{j=1}^{T'(T)} \left[\gamma \sigma_{\min}(\Lambda_2^\dagger(t_j)) (1 + o(1)) \right] \\ &\geq \sum_{j=1}^{\lceil cT \rceil} \left[\gamma \sigma_{\min}(\Lambda_2^\dagger(t_j)) (1 + o(1)) \right] \quad (16) \end{aligned}$$

This merits some explanation: The first inequality in (16) above is obtained by dropping the part pertaining to B , in (3). The second inequality is Corollary 4.3 to the Cramér-Rao bound, Theorem 4.2. In the third inequality, we use the assumption that a subsequence $\{t_j\}$ of time $\{t\}$ is uninformative, and apply Lemma 5.1. Fourth, we observe that the length of this subsequence by assumption is proportional to T , $T'(T) \geq cT$ for some $c \in (0, 1]$. Moreover, we may assume that $\sigma_{\min}(\Lambda_2(t_j)) > 0$ for all t_j of this subsequence, for otherwise the second part of Theorem 4.2 implies infinite variance, and thus infinite regret, and the result holds trivially.

Now, since the policy is α -fast convergent, it clear from (5) that $\sigma_{\max}(\Lambda_2(t)) = o(t^{1-2\alpha})$ and hence $\sigma_{\max}(\Lambda_2^\dagger(t)) = \Omega(t^{2\alpha-1})$. From this, we gather that

$$\sum_{t=1}^T \left[\gamma \sigma_{\min}(\Lambda_2^\dagger(t)) (1 + o(1)) \right] = \Omega(T^{2\alpha}).$$

On the other hand, if the policy is not α -fast convergent, one has that

$$\sum_{t=1}^T \left[\text{tr} \left((I \otimes r_t r_t^\top) \mathbf{E} \left[\underbrace{\text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top}_{\Omega(t^{-2\alpha})} \right] \right) \right] = \Omega(T^{1-2\alpha}). \quad (17)$$

Optimizing these terms by setting $\alpha = 1/4$ finishes the proof of the Theorem. ■

Remark 5.3: If (14) is strengthened to

$$\text{tr} \left((I \otimes B r_t r_t^\top B^\top) G_2 G_2^\top \right) > \gamma$$

the first term of the regret decomposition in Lemma 3.1 can also be lower-bounded by the same arguments and the theorem extends to the case $\lambda = 0$ in this way.

Let us now put Theorem 5.2 into perspective by comparing with what kind of lower bound we can prove if Fisher information has full rank.

Theorem 5.4: Suppose that $\sigma_{\min}(I_{t,B}^*) \geq \delta t + o(t)$, $\delta > 0$ and assume the additional regularity condition $H = \nabla_B \text{vec} \hat{K}_t(B) = \Omega(1)$. Then for any $\alpha > 0$ and any regular α -fast convergent policy, (1) satisfies the regret lower bound

$$R_T = \Omega(\log T). \quad (18)$$

In this regime, there is only very little trade-off between exploration and exploitation. Essentially, an optimal policy which has full rank, and thus full rank Fisher information, will already excite all directions of B and so there is little to be gained by adding extra excitation.

Proof: Write, using (3), and the Cramér-Rao bound, Theorem 4.2, with second block of dimension zero,

$$\begin{aligned} R_T &\geq \sum_{t=1}^T \mathbf{E} \text{tr} \left[\left((I \otimes r_t r_t^\top) \text{vec}(\hat{K}_t - K) (\text{vec}(\hat{K}_t - K))^\top \right) \right] \\ &\geq \sum_{t=1}^T \mathbf{E} \text{tr} \left[\left((I \otimes r_t r_t^\top) \underbrace{H I_{t,B}^\dagger H^\top}_{=\Omega(\delta/(t+o(t)))} \right) \right]. \end{aligned}$$

The result is now immediate since $\sum_{t=1}^T \delta/(t+o(t))$ scales like $\log T$. ■

Theorems 5.2 and 5.4 together provide strong evidence that a phase transition occurs. We now return to Example 2.2 to understand why logarithmic rates are feasible in [7].

Example 5.5: Let us revisit the scalar system $y_t = b u_t + w_t$. Here Fisher information is given by $I_{t,b}^* = \sum_{k=1}^t r_k^2 b^2 = t b^2$, for $r_t = 1$, which, in particular, has no nullspace. Hence the optimal policy is not uninformative for any constant and the singular value condition of Theorem 5.4 is satisfied. Indeed, the lower bound $\Omega(\log T)$ presented above matches (in order) the upper bound due to [7] discussed in Example 2.2.

VI. DISCUSSION

Our analysis here is similar spirit to [12]. The largest difference is that we take a closer look at the Cramér-Rao bound when the information matrix is near degenerate. In this regime learning becomes difficult. Similarly, the proof strategy is here also based on comparing the information ‘‘collected’’

by any algorithm and that of the optimal algorithm. However, degeneracy makes this comparison more difficult and we need to resort to singular space perturbation theory, [17]. It is this difference that allows us to demonstrate the \sqrt{T} -rate.

Using this, we show that if the optimal policy to (1) gives degenerate information in a certain sense, then regret must be super-logarithmic. In this regime, one is forced to introduce supplementary excitation beyond the randomness already present in the algorithm. It is also interesting to note that the lower bound strongly suggests that a certainty equivalent controller perturbed by noise with full rank covariance of order $1/\sqrt{t}$ is a good idea since this corresponds closely to the case for which the lower bound is optimized to $\Omega(\sqrt{T})$. Indeed, this was the strategy pursued by [10] attaining regret of order \sqrt{T} for the full LQR.

We also wish to mention that the concept of α -fast convergence is much inspired by the notion of uniformly fast convergence for the related bandit problems, see [19] and [20]. Indeed, what we have considered here can also be seen as a stochastic contextual bandit with side information y_t and context decided by r_t , see also [21] for an overview of bandits.

A. Directions for Future Work

We remark that our method of proving lower bounds has potential to be of general interest. Essentially, the key ingredients are a representation of regret in terms of an estimation error (a Taylor expansion of a convex potential) and an analysis of the rank of the Fisher information corresponding to the data collected by the optimal policy. If one is able to establish such a representation and asymptotically compare this Fisher information to that of any reasonable policy, our method carries through. In particular, it would be very desirable to extend the present results to dynamic models, such as LQR, and there, establish conditions which ensure a similar phase transition as observed here. In particular, this could help describe necessary and sufficient side information for logarithmic regret, [10]. To this end, [11] shows that under an identifiability condition, logarithmic regret is attainable for LQR and it would be very interesting to compare this notion to uninformative.

Another potential direction is to take a more information-theoretic route toward lower bounds as in [22] and as is traditional for bandits [19]. This was recently done for system identification in [23].

REFERENCES

- [1] N. Matni, A. Proutiere, A. Rantzer, and S. Tu, "From self-tuning regulators to reinforcement learning and back again," *arXiv preprint arXiv:1906.11392*, 2019.
- [2] Y. Abbasi-Yadkori and C. Szepesvári, "Regret bounds for the adaptive control of linear quadratic systems," in *Proceedings of the 24th Annual Conference on Learning Theory*, 2011, pp. 1–26.
- [3] M. K. S. Faradonbeh, A. Tewari, and G. Michailidis, "Optimism-based adaptive regulation of linear-quadratic systems," *arXiv preprint arXiv:1711.07230*, 2017.
- [4] H. Mania, S. Tu, and B. Recht, "Certainty equivalent control of LQR is efficient," *arXiv preprint arXiv:1902.07826*, 2019.
- [5] T. Lai, "Asymptotically efficient adaptive control in stochastic regression models," *Advances in Applied Mathematics*, vol. 7, no. 1, pp. 23–45, 1986.

- [6] T. Lai and C.-Z. Wei, "Extended least squares and their applications to adaptive control and prediction in linear systems," *IEEE Transactions on Automatic Control*, vol. 31, no. 10, pp. 898–906, 1986.
- [7] L. Guo, "Convergence and logarithm laws of self-tuning regulators," *Automatica*, vol. 31, no. 3, pp. 435–450, 1995.
- [8] A. Rantzer, "Concentration bounds for single parameter adaptive control," in *2018 Annual American Control Conference (ACC)*, IEEE, 2018, pp. 1862–1866.
- [9] K. J. Åström and B. Wittenmark, "On self tuning regulators," *Automatica*, vol. 9, no. 2, pp. 185–199, 1973.
- [10] M. K. S. Faradonbeh, A. Tewari, and G. Michailidis, "On adaptive linear-quadratic regulators," *Automatica*, vol. 117, 2020.
- [11] —, "Input perturbations for adaptive control and learning," *Automatica*, vol. 117, 2020.
- [12] I. Ziemann and H. Sandberg, "Regret lower bounds for unbiased adaptive control of linear quadratic regulators," *IEEE Control Systems Letters*, forthcoming, 2020.
- [13] M. Simchowitz and D. J. Foster, "Naive exploration is optimal for online lqr," *arXiv preprint arXiv:2001.09576*, 2020.
- [14] A. Cassel, A. Cohen, and T. Koren, "Logarithmic regret for learning linear quadratic regulators efficiently," *arXiv preprint arXiv:2002.08095*, 2020.
- [15] A. W. Van der Vaart, *Asymptotic statistics*. Cambridge university press, 2000, vol. 3.
- [16] P. Stoica and T. L. Marzetta, "Parameter estimation problems with singular information matrices," *IEEE Transactions on Signal Processing*, vol. 49, no. 1, pp. 87–90, 2001.
- [17] P.-Å. Wedin, "Perturbation bounds in connection with singular value decomposition," *BIT Numerical Mathematics*, vol. 12, no. 1, pp. 99–111, 1972.
- [18] T. T. Cai, A. Zhang, *et al.*, "Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics," *The Annals of Statistics*, vol. 46, no. 1, pp. 60–89, 2018.
- [19] T. L. Lai and H. Robbins, "Asymptotically efficient adaptive allocation rules," *Advances in applied mathematics*, vol. 6, no. 1, pp. 4–22, 1985.
- [20] A. Garivier, P. Ménard, and G. Stoltz, "Explore first, exploit next: The true shape of regret in bandit problems," *Mathematics of Operations Research*, vol. 44, no. 2, pp. 377–399, 2018.
- [21] T. Lattimore and C. Szepesvári, "Bandit algorithms," *preprint*, 2018.
- [22] M. Raginsky, "Divergence-based characterization of fundamental limitations of adaptive dynamical systems," in *2010 48th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, IEEE, 2010, pp. 107–114.
- [23] Y. Jedra and A. Proutiere, "Sample complexity lower bounds for linear system identification," *2019 IEEE Conference on Decision and Control (CDC)*, 2019.

APPENDIX

We require the following version of Wedin's $\sin \Theta$ Theorem: Consider matrices M and $\tilde{M} = M + T$ for some perturbation T , with singular value decompositions $M = V_1 \Gamma_1 W_1^T + V_2 \Gamma_2 W_2^T$, $\tilde{M} = \tilde{V}_1 \tilde{\Gamma}_1 \tilde{W}_1^T + \tilde{V}_2 \tilde{\Gamma}_2 \tilde{W}_2^T$. If for $\delta > 0$, $\eta \geq 0$, $\sigma_{\min}(\tilde{\Gamma}_1) \geq \eta + \delta$ and $\sigma_{\max}(\Gamma_2) \leq \eta$ then

$$\max_P \|\tilde{V}_2 P - V_2\|_{\infty} = O(\|T\|_{\infty})$$

where the maximization is over the orthogonal group.

The result is due to Wedin [17]. The formulation of the $\sin \Theta$ distance as an optimization over the norm $\|\cdot\|_{\infty}$ appears for instance in [18].