



HAL
open science

Trophic transfer of trace elements in a euryhaline fish, the turbot *Scophthalmus maximus*: Contrasting effects of salinity on two essential elements

Simon Pouil, François Oberhänsli, Paco Bustamante, Marc Metian

► To cite this version:

Simon Pouil, François Oberhänsli, Paco Bustamante, Marc Metian. Trophic transfer of trace elements in a euryhaline fish, the turbot *Scophthalmus maximus*: Contrasting effects of salinity on two essential elements. *Marine Pollution Bulletin*, 2020, 154, pp.111065. 10.1016/j.marpolbul.2020.111065 . hal-02547207

HAL Id: hal-02547207

<https://hal.science/hal-02547207>

Submitted on 19 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Trophic transfer of trace elements in a euryhaline fish, the turbot *Scophthalmus maximus*: Contrasting effects of salinity on two essential elements

Simon Pouil^{a,b}, François Oberhänsli^a, Paco Bustamante^{b,c}, Marc Metian^a

^a International Atomic Energy Agency, Environment Laboratories, 4a, Quai Antoine 1er, MC-98000, Principality of Monaco, Monaco

^b Littoral Environnement et Sociétés (LIENSs), UMR 7266, CNRS- La Rochelle University, 2 rue Olympe de Gouges, F-17000 La Rochelle, France

^c Institut Universitaire de France (IUF), 1 rue Descartes 75005 Paris, France

* Corresponding author: Marc Metian

 Radioecology Laboratory

 IAEA Environment Laboratories

 4a Quai Antoine 1er

 MC-98000 Principality of Monaco

 E-mail: m.metian@iaea.org

Highlights:

- Effects of salinity on trophic transfer of trace elements were assessed in turbot
- Salinity has contrasting effects on the trophic transfer of trace elements in fish
- Assimilation of Mn in fish is significantly lower at the highest salinity
- Salinity did not significantly influence Zn trophic transfer in fish

Abstract: Trace elements can be accumulated from coastal environment to the aquatic organisms by their food and be transferred throughout the food webs. Studying the effects of salinity on the trophic transfer of trace elements in euryhaline organisms, able to deal with large variations in salinity, is therefore key to understand their dynamics in aquatic environments. In this context, we investigated the potential influence of salinity on the trophic transfer of two essential elements (Mn and Zn) in the euryhaline fish, the turbot *Scophthalmus maximus* using radiotracer techniques. After acclimation to three salinities (10, 25 and 38), juvenile turbot were fed with radiolabelled pellets (^{54}Mn and ^{65}Zn). Kinetic parameters of depuration were determined after a 21-d period and trophic transfer factors were calculated. Trophic transfer of Mn at the highest salinity was significantly lower than for the other conditions whereas salinity did not significantly influence Zn trophic transfer. Differences in the processes involved in the regulation (homeostasis) of the two tested trace elements may explain the contrasting influence of water salinity for Mn and Zn.

Keywords: Diet; Metals; Teleost; Zinc; Manganese; Salinity

1. Introduction

Environmental conditions can affect trace element bioaccumulation in aquatic organisms such as fish (Luoma and Rainbow, 2005; Phillips and Rainbow, 1993). Among environmental factors, water salinity is known to influence trace element accumulation by causing (1) changes in trace element speciation and therefore on their bioavailability, and (2) modifications of fish physiology, especially regarding the osmoregulation processes (Ni et al., 2005). Salinity affects biokinetic parameters, such as the uptake rate (k_u) or the efflux rate (k_e) of some trace elements such as Cd, Cs, Se, and Zn in fish (Ni et al., 2005; Zhao et al., 2001). Nevertheless, most of the studies that looked at the potential effects of salinity on metal accumulation focused on waterborne trace elements rather than dietary ones (e.g., Zhang and Wang, 2007; Zhao et al., 2001; Webb and Wood, 2000). Thus, only limited information is available about the influence of salinity on trace element trophic transfer while diet is recognized as the major pathway for most of the trace elements bioaccumulated by fish (e.g., Pouil et al., 2018a; Mathews and Fisher, 2009; Xu and Wang, 2002).

The turbot, *Scophthalmus maximus* (Scophthalmidae), is a demersal fish widely distributed in Western European coastal waters. This species inhabits in a wide range of water salinities with its breeding usually occurring in low-salinity waters (Kuhlmann and Quantz, 1980). The euryhaline nature of this species has been confirmed by Waller (1992) who reported that osmoregulatory disturbances only occurred below salinity of 6. This euryhaline species, through its ability to move from brackish water to seawater environments and its trophic ecology, is a potential significant vector of trophic transfer of trace elements from coastal to marine ecosystems. Furthermore, turbot became a few years ago a model species for trace element studies on fish. The turbot was recently considered as a biological model for numerous ecotoxicological studies related to the bioaccumulation of dietary trace elements (e.g., Pouil et al., 2018b, 2017a, 2016, 2015). Studying this species is therefore relevant to assess the effect

of salinity on the trophic transfer of trace elements in fish.

The present study investigated the possible effects of a wide range of salinities on the assimilation efficiency (AE) of two essential trace elements (Mn and Zn) in the euryhaline fish, turbot *S. maximus*. Radiotracer techniques were used to determine depuration parameters and body distribution of the selected dietary trace elements in *S. maximus* exposed to three salinities (10, 25 and 38). Kinetic parameters data were then used to model the influence of salinity on the potential of biomagnification (Trophic Transfer Factor, TTF) of Mn and Zn in the field.

2. Materials and Methods

2.1. Origin and acclimation of fish

Juvenile turbot *S. maximus* were purchased from a fish farm (France Turbot, www.france-turbot.com) and shipped to the International Atomic Energy Agency premises in the Principality of Monaco. The fish were kept in a 700-L aquarium (open circuit, water renewal: 350 L h⁻¹; 0.45µm filtered seawater; salinity: 38; light/dark: 12h/12h). Then, three weeks before the experiment, 24 fish were randomly placed in three 20-L aquaria (n=8) and acclimated to the target salinities (10, 25, 38). During the first days of acclimation, salinities were gradually decreased and then stabilized to the targeted values for 10 days before starting experiment. During the acclimation period, the fish were fed a daily ration of 1.5% of their biomass with 1.1-mm commercial pellets (proteins: 55% and lipids: 12%; Le Gouessant, www.legouessant.com).

Salinity was measured twice a day in each aquarium using a hand-held conductivity/salinity meter, which was calibrated using conductivity standards encompassing the range of the three selected experimental waters. Furthermore, in each aquarium, pH and temperature were

monitored every 15 minutes using a continuous measurement system (IKS ComputerSysteme, www.iks-aqua.com). Values of water parameters are summarized in Table 1.

2.2. Experimental procedures

2.2.1. Radiolabelling of pellets

Fifteen grams of 1.1-mm pellets were radiolabelled for 1 h in 20 mL of seawater spiked with 1.5 kBq mL⁻¹ of ⁵⁴Mn and ⁶⁵Zn. Pellets were then dried for 48 h at 50°C and kept in a dry environment in order to prevent mould growth. In terms of stable metal concentrations in the pellet, these additions of radiotracers corresponded to 10 ng g⁻¹ for Mn, 1.4 µg g⁻¹ for Zn, i.e. concentrations that are lower than the common concentrations of these metals in the potential prey of the fish (Pouil et al., 2016). Preliminary tests were performed to determine the possible leakage into the water of radioisotopes from the pellets during the feeding. When food was provided, acclimated fish consumed the pellets in less than 2 min. Therefore, preliminary tests consisted in pouring radiolabelled dry pellets (100 mg per treatment) for 1, 5 and 10 min in 50 mL seawater and to measure any radioactivity in the seawater (Pouil et al., 2015). The leakage of pellet-radioactivity was under the detection limits even after 10 min immersed in the seawater, respectively. Although these tests confirmed the single-pathway contamination (viz. food) of the fish, one turbot was used in each treatment, as a control to take into account the possibility of ⁵⁴Mn and ⁶⁵Zn recycling through water (see Section 2.3.2).

2.2.2. Exposure of turbot via radiolabelled pellets

A total of 8 acclimatized turbots were randomly selected for each experimental salinity (10: 44.0 ± 2.9g; 25: 40.0 ± 3.8g and 38: 42.6 ± 3.4g). Slits cut into the fins were used to facilitate individual recognition. Each experiment consisted of a single feeding of fish with radiolabelled

pellets, a food commonly used in the literature (e.g. Jacob et al., 2017; Zhou et al., 2017; Pouil et al., 2015) to assess trophic transfer of trace elements in fish. After the labelled feeding, an additional turbot was placed in each aquarium to assess any possible radiotracer recycling from seawater due to leaching from the radiolabelled food or, later on, from fish depuration. Two hours after the 15-min feeding, individual fish were whole-body γ -counted alive and then replaced in the same aquarium to follow subsequent trace element depuration. All the fish (including control individual of each condition) were regularly radioanalysed to follow the radiotracer depuration kinetics over 21 days.

After the depuration period, 4 individuals per condition were dissected in 7 compartments: (1) the digestive tract, (2) the gall bladder, (3) the head (including gills), (4) the kidney, (5) the liver, (6) the 2 axial muscles (without dorsal skin) and (7) the remaining tissues (including ventral skin, skeleton, fins, heart and muscle residues) and were separated, weighed and radioanalysed to determine the radiotracer body distribution.

2.3. Radiotracers and Counting

Radiotracers of high specific activity were purchased from Polatom, Poland (^{54}Mn as MnCl_2 in 0.5M HCl, $t_{1/2} = 312$ days and ^{65}Zn as ZnCl_2 in 0.1M HCl, $t_{1/2} = 244$ days). The radioactivity of the tracers was measured using a high-resolution γ -spectrometer system composed of 5 Germanium - N or P type - detectors (EGNC 33-195-R, Canberra[®] and Eurysis[®]) connected to a multi-channel analyser and a computer equipped with a spectra analysis software (Interwinner 6, Intertechnique[®]). The radioactivity in living organisms and samples was determined by comparison with standards of known activity and of appropriate geometry (calibration and counting). Measurements were corrected for background and physical radioactive decay. Living organisms were placed in counting tubes (diameter: 160 mm, height: 80 mm) filled with 500 mL of clean seawater (at the appropriated conditions of salinity) during the counting period.

The counting time was adjusted to obtain a propagated counting error less than 5% (e.g., Rodriguez y Baena et al., 2006) for a maximum of 20 min. As already described by Pouil et al. (2017b), tests were performed prior to the experiment, where fish were placed in similar counting conditions in order to observe their behaviour, i.e. in a counting box for 20 min in the dark. Dissolved O₂ concentration was monitored throughout these tests and was always > 3 mg L⁻¹. No alteration in organism health or behaviour was observed during the tests and then, the experiment.

2.4. Kinetic parameters and TTF

Depuration of radiotracers was expressed as the percentage of remaining radioactivity (radioactivity at time t divided by the initial radioactivity measured in the organism at the beginning of the depuration period; following methods developed in Warnau et al. (1996). The depuration kinetics of Mn and Zn were best fitted using a two-component exponential model:

$$A_t = A_{0s} \cdot e^{-k_{es}t} + A_{0l} \cdot e^{-k_{el}t} \quad (1)$$

where A_t and A_0 are the remaining activities (%) at time t (d) and 0, respectively; k_e is the depuration rate constant (d⁻¹). “s” and “l” subscripts are related to the short- and long-lived component, respectively. The “s” component represents the depuration of the radiotracer fraction that is weakly associated with the organisms and rapidly eliminated (i.e. proportion associated with the faeces). The “l” component describes the depuration of the radiotracer fraction that is actually absorbed by the organism and eliminated slowly (Whicker and Schultz, 1982; Reichle, 1967; Hubbell et al., 1965). The long-lived component allows estimating the assimilation efficiency (AE) of the radiotracer ingested with food (AE = A_{0l} ; Pouil et al., 2018a; Warnau et al., 1996). For the two components, biological half-life ($T_{b1/2}$) can be calculated from the corresponding depuration rate constant according to the relation $T_{b1/2} = \ln 2/k_e$. Kinetic

parameters were determined using the R freeware 3.5.2 (R Development Core Team, 2018) and the ‘nlstools’ package (Baty and Delignette-Muller, 2015).

To assess the biomagnification potential of Mn and Zn following dietary exposure, trophic transfer factors (TTFs) were also calculated for a specific link in the food chain in which a predator consumes metal in prey as follows:

$$TTF = \frac{AE \times IR}{k_{el}} \quad (2)$$

Where AE is the assimilation efficiency of the ingested metal in the fish, IR is the weight-specific ingestion rate of prey ($\text{g g}^{-1} \text{d}^{-1}$) and k_{el} is the depuration rate constant (d^{-1}) of the radionuclide out of the predator (see Mathews et al., 2008; Zhao et al., 2001). A $TTF > 1$ suggests that biomagnification is possible, and $TTF < 1$ suggests that biomagnification is unlikely (Reinfelder et al., 1998). For these TTF calculations we considered a range of ingestion rates (IR) by fish likely to be encountered under natural conditions (0.02 to $0.10 \text{ g g}^{-1} \text{d}^{-1}$; Zhao et al., 2001).

Statistical comparisons between the three different salinity experiments were conducted using individual depuration kinetics of each element: individual kinetic parameters were obtained using the best fitting model at the global scale to the data of each individual. Then differences between these parameters were tested using Kruskal-Wallis and Siegel and Castellan non-parametric tests (Zar, 1996). The same statistical tests were used to compare Mn and Zn organotropism of turbot under the different salinity conditions. The level of significance was always set at $\alpha = 0.05$.

3. Results

In order to evaluate how salinity affects the assimilation of essential trace elements in the

juvenile turbot *S. maximus*, depuration kinetics of Mn and Zn were followed after a pulse-chase feeding, using radiolabelled pellets. During the whole experimental period (i.e. three weeks of acclimation to the targeted salinity values followed by three weeks of depuration) where the fish were exposed to a gradient of salinities (see the Materials and Methods section), only a limited growth of the individuals was recorded and no mortality occurred. Before the single-feeding, the activity level of Mn and Zn was measured in the pellets: $2202 \pm 158 \text{ Bq } ^{54}\text{Mn g}^{-1}$ and $2394 \pm 167 \text{ Bq } ^{65}\text{Zn g}^{-1}$. During the entire experiment, the exclusive foodborne exposure of the fish to both radiotracers was confirmed (no activity was recorded in the control turbot).

Whole-body depuration kinetics of ^{54}Mn and ^{65}Zn in turbot were best fitted by a two-phase model (R^2 : 0.88-0.98; Fig. 1 and Table 2). A large proportion (57-81%, Table 2) of the ingested radiotracers was associated with the short-term component for both the studied elements. This component was characterized by a very rapid loss ($T_{b1/2s} < 1 \text{ d}$, Table 2). Comparison of short-term depuration rate constants (k_{es}) determined for each individual turbot indicated that there was no significant difference for both studied elements ($p > 0.05$) independently of the salinity conditions. Estimated AEs in turbot ranged from 25% to 43% for Mn whereas Zn was less assimilated ($\text{AE} < 25\%$, Fig. 1 and Table 2). Statistical analyses carried out on individual estimated AEs revealed that salinity affected the trophic transfer of Mn with a significantly lower AE at the highest salinity ($p < 0.05$; Fig. 2). In contrast, no significant effect of the salinity was observed for AEs of Zn ($p > 0.05$; Fig. 1). For Mn, long-term efflux rate constants were 2 times higher ($p < 0.05$, Fig. 2) in the turbot maintained at the highest salinity condition (38) with $T_{b1/2l}$ of 46 d while $T_{b1/2l}$ reached 79-86 d in low salinity conditions (10 and 25).

For Mn and, to a lesser extent, for Zn, TTFs calculated for a range of IR were dependent on the salinity conditions (Fig. 3). Depending on the IR, TTFs ranged from 1.1 to 5.4 and 0.3 to 1.6 for Mn at the lowest and the highest salinity conditions, respectively. Zn TTFs ranged from 0.5 to 4.9 with the lowest TTFs observed at the highest salinity (Fig. 3).

Post-feeding distributions of Mn and Zn in turbot exposed to the gradient of salinity, at the end of the 21-d depuration period, is shown in Fig. 4. Similar patterns of Mn and Zn distribution among compartments were observed in turbot exposed to the three salinities ($p > 0.05$). Distribution among the body compartments systematically ranked according to the following decreasing order (Fig. 4): remaining tissues (i.e. remaining skin, skeleton, fins, heart and remaining parts; 44-50%) > head (33-45%) >> axial muscles (4-10%) > digestive tract (<1-9%) >> liver (<2%) > kidney (<1.4%) >> gall bladder (< 0.2%).

4. Discussion

Salinity is an environmental master factor in coastal marine ecosystems (Smyth and Elliott, 2016). Salinity is known to strongly influence the bioaccumulation of dissolved trace elements in aquatic organisms such as fish through osmoregulatory processes (Wang and Rainbow, 2008; Zhang and Wang, 2007; Ni et al., 2005) and thus affects the dynamics of trace elements in marine environment. Both changes in metal speciation and physiology explained differential trace element uptake rates from water in fish exposed to different salinities (Wang and Rainbow, 2008). Nevertheless, although in their natural environment fish are also exposed to trace elements from the water and eventually from the sediment, previous laboratory studies highlighted that diet is the main pathway of trace element bioaccumulation in fish (Pouil et al., 2018a; Mathews and Fisher, 2009; Xu and Wang, 2002).

Effects of salinity on the trophic transfer of trace elements have been poorly investigated in the literature. Ni et al. (2005) found no significant differences in the Assimilation Efficiency (AE) of Cd, Se and Zn in the mudskipper *Periophthalmus modestus* fed on radiolabeled polychaetes and acclimated to a gradient of salinity from 10 to 30. In contrast, Zhou et al. (2017) found more contrasting results with Cu AEs measured in the white-spotted spinefoot *Siganus canaliculatus* fed on CuSO₄-spiked commercial pellets decreased from a salinity of 33 to 10

but increased at lower salinity. In the present study, we found that Zn AEs determined in juvenile turbot at the three different salinities (10, 25 and 38) were similar but Mn AE was significantly lower in the fish exposed to the highest salinity (38) showing that the effects of salinity on the trophic transfer of trace elements in fish are species- and element- dependent.

Zn is one of the most important essential trace elements for fish due to its structural and catalytic roles in more than 300 proteins and it serves as a cofactor in many enzymatic systems, playing a vital role in lipid, protein, and carbohydrate metabolism (Bury et al., 2003; Watanabe et al., 1997). Thus, this element is directly involved in growth, reproduction, development and immunity in fish (Watanabe et al., 1997; Tacon, 1987). Although the mechanisms of Zn transfer from the gut lumen to the internal compartment (absorption) are not fully elucidated yet, it seems to be dominated by active processes involving specific transporters (Bury et al., 2003). This element is, among other things, accumulated into cells through specific channels (ZIP family; Hogstrand, 2011; Bury et al., 2003). However, as concentrations can easily be toxic, steady-state cytosolic Zn concentration is controlled by an efflux transporter of the ZnT family that transports Zn from the cytosol outside the cells (Hogstrand, 2011; Bury et al., 2003). An excess of Zn can be excreted mainly via the bile, intestinal sloughing (Handy, 1996) or the gills (Hardy et al., 1987). Thus, at both organismal and cellular levels Zn status in fish is actively and tightly controlled. In the present experiment, the ^{65}Zn concentrations remained constant in tissues in the fish exposed to the gradient of salinity with $\sim 2.6\text{-}3.2 \text{ Bq } ^{65}\text{Zn g}^{-1}$ fresh weight (FW) demonstrating the existence of internal Zn homeostasis. This mechanism may explain the consistency observed for the proportion of assimilated dietary Zn in turbot (19-24%; this study) and mudskipper (5-7%; Ni et al., 2005) despite variations of salinity.

Mn is necessary for the normal functioning of brain and for lipid and carbohydrate metabolism. This element has a key role as a cofactor for enzymes and as a structural element of metalloenzymes. As a cofactor or component of several key enzyme systems, Mn is also

directly involved in bone formation, regeneration of red blood cells and reproduction (Watanabe et al., 1997; Tacon, 1987). Interestingly, we demonstrated that for Mn both AE and k_{el} were significantly affected by salinity. Indeed, while the lowest Mn AE was measured in turbot acclimated to the highest salinity (38), in the same condition efflux rate constant (k_{el}) was significantly higher indicating a lesser Mn retention for fish from this treatment (i.e. $T_{b1/2}$ two times lower compared to the other treatments). Such differences can be explained by a less tight homeostasis for Mn compared to Zn as suggested by the differences observed in the whole-body ^{54}Mn concentrations in fish depending on the salinity with values of $\sim 5 \text{ Bq } ^{54}\text{Mn g}^{-1} \text{ FW}$ for the turbot acclimation to the low salinity conditions while ^{54}Mn concentrations decreased to $\sim 3 \text{ Bq } ^{54}\text{Mn g}^{-1} \text{ FW}$ in turbot maintained at a salinity of 38. Nevertheless, the mechanisms of transport and absorption of Mn from food in fish are poorly reported and further investigations are needed to support this assumption.

Our results for Mn and Zn revealed the contrasting effects of the salinity on the trophic transfer of trace elements in fish. We assumed that such differences may be reasonably explained by the physiological mechanisms involved in the homeostasis of essential trace elements in fish. In addition to whole-body kinetic determination, we also performed fish dissection at the end of the 21-d depuration period. Indeed, measurements of the distribution of Mn and Zn provide additional mechanistic information potentially helping in the interpretation of results from whole-body kinetic measurements. Our results show that the distribution of Mn and Zn between the tissues of juvenile turbot did not vary significantly with salinity. More than >98% Mn and >88% of Zn were found in the muscles, the head, and in the remaining tissues. Similar findings were reported in previous studies (Pouil et al., 2017a) where >95% of the Mn and >86% of the Zn were found in the same tissues. The absence of changes in distribution of these element in the body compartments could be related to the fact that the experimental context is reflecting non-polluted conditions (i.e., no excess of Mn and Zn in the diet) and rather reflects normal

physiological processes (Pouil et al, 2017a).

Altogether, the results of this study highlighted the effects of salinity that could potentially lead to a change in the transfer of Mn and Zn within aquatic food webs. Indeed, the estimated TTFs ranging from 1-6 at salinity of 10 were 3 times lower at the highest salinity (38) demonstrating that, salinity, in addition to playing an important role in the bioaccumulation of dissolved trace elements, affects also their transfer from diet. Consequently, attention should be paid on this environmental variable for obtaining a better understanding of the dynamics of trace elements within food webs in coastal marine ecosystems subject to variable inputs of freshwater and trace elements. Nevertheless, care needs to be taken in how to interpret and expand on these results. Indeed, the methodological approach used in this study allowed determining the kinetics of depuration of dietary Mn and Zn in living fish acclimated to a gradient of salinity. However, further investigations are needed to fully understand the influence of salinity on the physiological mechanisms involved in assimilation of the studied trace elements. Furthermore, in our study, turbot were acclimated to stable salinities while salinity changes can occur abruptly in coastal environments (Smyth and Elliott, 2016). These variations are likely to affect the physiology of the organisms that are exposed to them and, consequently, to affect the trophic transfer of the trace elements.

5. Conclusion

In summary, our study showed that salinity differently impacted the AE of Mn and Zn, two essential elements in the juvenile turbot, although this species is euryhaline (i.e. species with a large salinity tolerance). Indeed, Mn AE was higher at lower salinities (10 and 25) than at high salinity (38) while Zn AE was not affected by the salinity conditions. These differences were likely caused by the physiological changes rather than the changes in trace element speciation. After the 21-d depuration period, tissue distributions were similar both for Mn and

Zn in turbot acclimated to the three salinities. Given the evidence that food is the major pathway of trace element bioaccumulation in marine fish, salinity would be one important environmental variable driving the trophic transfer of some trace elements in coastal aquatic ecosystems.

Acknowledgments

The IAEA is grateful for the support provided to its Environment Laboratories by the Government of the Principality of Monaco. The IUF (Institut Universitaire de France) is acknowledged for its support to PB as a Senior Member.

References

- Baty, F., Delignette-Muller, M.L., 2015. nlstools: Tools for Nonlinear Regression Analysis. R package version 1.0-2. <http://CRAN.R-project.org/package=nlstools>.
- Bury, N.R., Walker, P.A., Glover, C.N., 2003. Nutritive metal uptake in teleost fish. *Journal of Experimental Biology* 206, 11-23.
- Handy, R.D., 1996. Dietary exposure to toxic metals in fish. In: Taylor, E.W. (eds.) *Toxicology of aquatic pollution: physiological, molecular and cellular approaches*. Cambridge University Press, Cambridge, United Kingdom, pp. 29-60.
- Hardy, R.W., Sullivan, C.V., Koziol, A.M., 1987. Absorption, body distribution, and excretion of dietary zinc by rainbow trout (*Salmo gairdneri*). *Fish Physiology and Biochemistry* 3, 133-143.
- Hogstrand, C., 2011. Zinc. In: Wood, C., Farrell, A., Brauner, C. (eds.) *Fish physiology: homeostasis and toxicology of essential metals*. Academic Press, Cambridge, UK, pp. 135-200.
- Hubbell, S.P., Sikora, A., Paris, O.H., 1965. Radiotracer, gravimetric and calorimetric studies of ingestion and assimilation rates of an isopod. *Health Physics* 11, 1485-1501.
- Jacob, H., Pouil, S., Lecchini, D., Oberhänsli, F., Swarzenski, P., Metian M., 2017. Trophic transfer of essential elements in the clownfish *Amphiprion ocellaris* in the context of ocean acidification. *PloS ONE* 12, e0174344.
- Kuhlmann, D., Quantz G., 1980. Some effects of temperature and salinity on the embryonic development and incubation time of the turbot, *Scophthalmus maximus* L., from the Baltic Sea. *Meeresforschung* 28, 172-178.
- Luoma, S.N., Rainbow, P.S., 2005. Why is metal bioaccumulation so variable? *Biodynamics*

- as a unifying concept. *Environmental Science & Technology* 39, 1921-1931.
- Mathews, T., Fisher, N.S., 2009. Dominance of dietary intake of metals in marine elasmobranch and teleost fish. *Science of the Total Environment* 407, 5156-5161.
- Mathews, T., Fisher, N.S., Jeffree, R.A., Teyssié, J.-L., 2008. Assimilation and retention of metals in teleost & elasmobranch fishes following dietary exposure. *Marine Ecology Progress Series* 360, 1-12.
- Ni, I.-H., Chan, S.M., Wang, W.-X., 2005. Influences of salinity on the biokinetics of Cd, Se, and Zn in the intertidal mudskipper *Periophthalmus cantonensis*. *Chemosphere* 61, 1607-1617.
- Phillips, D.J.H., Rainbow, P.S., 1993. *Biomonitoring of trace aquatic contaminants*. Elsevier Applied Science, London.
- Pouil, S., Bustamante, P., Warnau, M., Metian, M., 2018a. Overview of trace element trophic transfer in fish through the concept of assimilation efficiency. *Marine Ecology Progress Series* 588, 243-254.
- Pouil, S., Oberhänsli, F., Bustamante, P., Metian, M., 2018b. Investigation of temperature and pH variations on the metal trophic transfer in turbot (*Scophthalmus maximus*). *Environmental Science and Pollution Research* 25, 11219-11225.
- Pouil, S., Oberhänsli, F., Bustamante, P., Metian, M., 2017a. Dietary Zn and the subsequent organotropism in fish: No influence of food quality, frequency of feeding and environmental conditions (pH and temperature). *Chemosphere* 183, 503-509.
- Pouil, S., Teyssié, J.-L., Rouleau, C., Fowler, S.W., Metian, M., Bustamante P., Warnau, M. 2017b. Comparative study of trophic transfer of the essential metals Co and Zn in two tropical fish: A radiotracer approach. *Journal of Experimental Marine Biology and Ecology* 486, 42-51.

- Pouil, S., Warnau, M., Oberhänsli, F., Teyssié, J.-L., Bustamante, P., Metian, M., 2016. Influence of food on the assimilation of essential elements (Co, Mn, and Zn) by turbot *Scophthalmus maximus*. Marine Ecology Progress Series 550, 207-218.
- Pouil, S., Warnau, M., Oberhänsli, F., Teyssié, J.-L., Metian, M., 2015. Trophic transfer of ^{110m}Ag in the turbot *Scophthalmus maximus* through natural prey and compounded feed. Journal of the Environmental Radioactivity 150, 189-194.
- R Development Core Team, 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Reichle, D.E., 1967. Radioisotope turnover and energy flow in terrestrial isopod populations. Ecology 48, 351-366.
- Reinfelder, J.R., Fisher, N.S., Luoma, S.N., Nichols, J.W., Wang; W.-X., 1998. Trace element trophic transfer in aquatic organisms: a critique of the kinetic model approach. Science of the Total Environment 219, 117-135.
- Rodriguez y Baena, A.M., Miquel, J.C., Masqué, P., Povinec, P.P., La Rosa J., 2006. A single vs. double spike approach to improve the accuracy of ^{234}Th measurements in small-volume seawater samples. Marine Chemistry 100, 269-281.
- Smyth, K., Elliott, M., 2016. Effects of changing salinity on the ecology of the marine environment. In; Solan, M., Whiteley, M. (eds.) Stressors in the marine environment: Physiological and ecological responses; societal implications. Oxford University Press, Oxford, United Kingdom, pp. 162-174.
- Tacon, A.G.J., 1987. The nutrition and feeding of farmed fish and shrimp - A training manual 1: The essential nutrient. Food and Agriculture Organization of the United Nations, Roma, Italy.

- Waller, U., 1992. Factors influencing routine oxygen consumption in turbot, *Scophthalmus maximus*. *Journal of Applied Ichthyology* 8, 62-71.
- Wang, W.-X., Rainbow, P.S., 2008. Comparative approaches to understand metal bioaccumulation in aquatic animals. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 148, 315-323.
- Warnau, M., Teyssié, J.-L., Fowler, S.W., 1996. Biokinetics of selected heavy metals and radionuclides in the common Mediterranean echinoid *Paracentrotus lividus*: Sea water and food exposures. *Marine Ecology Progress Series* 141, 83-94.
- Watanabe, T., Kiron, V., Satoh, S., 1997. Trace minerals in fish nutrition. *Aquaculture* 151, 185-207.
- Webb, N.A., Wood, C.M., 2000. Bioaccumulation and distribution of silver in four marine teleosts and two marine elasmobranchs: Influence of exposure duration, concentration, and salinity. *Aquatic Toxicology* 49, 111-129.
- Whicker, F.W., Schultz, V., 1982. Radioecology: Nuclear energy and the environment: Volume 1. CRC Press, Boca Raton, FL.
- Xu, Y., Wang, W.-X., 2002. Exposure and potential food chain transfer factor of Cd, Se and Zn in marine fish *Lutjanus argentimaculatus*. *Marine Ecology Progress Series* 238, 173-186.
- Zar, J.H., 1996. *Biostatistical Analysis*: 3rd edition. Prentice-Hall, Upper Saddle River, NJ.
- Zhang, L., Wang, W.-X., 2007. Waterborne cadmium and zinc uptake in a euryhaline teleost *Acanthopagrus schlegeli* acclimated to different salinities. *Aquatic Toxicology* 84, 173-181.
- Zhao, X., Wang, W.-X., Yu, K.N., Lam, P.K.S., 2001. Biomagnification of radiocesium in a

marine piscivorous fish. *Marine Ecology Progress Series* 222, 227-237.

Zhou, Y., Zhang, W., Guo, Z., Zhang, L., 2017. Effects of salinity and copper co-exposure on copper bioaccumulation in marine rabbitfish *Siganus oramin*. *Chemosphere* 168, 491-500.




Figure 1. Influence of salinity on whole-body depuration of ^{54}Mn and ^{65}Zn in juvenile turbot ($n = 7$; % remaining activities). Parameters and statistics of depuration kinetics are given in Table 2.




Figure 2. Comparison of (A) assimilation efficiencies (AEs) and (B) efflux rate constant (k_{ei}) of Mn calculated for each individual turbot acclimated to three salinity conditions. The best fitting model obtained for the entire set of turbots (see Fig. 1 and Table 2) was applied to individuals. Letters denote significant differences between the salinity conditions.




Figure 3. Trophic transfer factors for juvenile turbot acclimated to three salinity conditions (10, 25 and 38) and fed with radiolabeled pellets at different assimilation efficiencies (AE) and ingestion rates. “High” AE (mean + SD), “intermediate” AE (mean), and “low” AE (mean - SD) values are reported in Table 2.




Figure 4. Distribution (%) of ⁵⁴Mn, and ⁶⁵Zn in juvenile turbot acclimated to three salinity conditions (10, 25 and 38), exposed to the radiotracers during a single-feeding with radiolabelled pellets and then maintained for a 21-d depuration period in unspiked seawater at the given salinity. Values are means ± SD (n=4). *The remaining tissues included remaining skin, skeleton, fins, heart and muscle residues

Table 1. Water parameters during the experiment on the assimilation of essential trace elements in juvenile turbot exposed to different conditions of salinity. Values are Means \pm SD. Letters denote significant differences.

Condition	Salinity measured	Conductivity (mS cm ⁻¹)	pH	Temperature (°C)
Low salinity (10)	10.05 \pm 0.13 ^a	16.99 \pm 0.09 ^a	8.02 \pm 0.04 ^a	19.70 \pm 0.08 ^a
Medium salinity (25)	24.94 \pm 0.16 ^b	39.27 \pm 0.15 ^b	7.99 \pm 0.07 ^b	19.72 \pm 0.05 ^a
High salinity (38)	37.8 \pm 0.05 ^c	56.98 \pm 0.04 ^c	7.98 \pm 0.03 ^c	19.71 \pm 0.12 ^a

Table 2. Estimated depuration kinetic parameters of ^{54}Mn , and ^{65}Zn in turbot acclimated to three salinity conditions (10, 25 and 38; $n = 7$ per treatment) and exposed to the radiotracers during a single-feeding with radiolabelled pellets. After the radiolabelled feeding, turbot were maintained for 21d in unspiked seawater at the given salinity. Depuration parameters: A_{0s} and A_{0l} (=AE): activity (%) lost according to the short- and the long- lived exponential component, respectively; k_e : depuration rate constant (d^{-1}); $T_{b1/2}$: biological half-life (d) [$T_{b1/2}=\ln 2/k_e$]; ASE: asymptotic standard error; R^2 : determination coefficient.

Salinity	Short-term			Long-term			R^2
	$A_{0s} \pm \text{ASE}$	$k_{es} \pm \text{ASE}$	$T_{b1/2s} \pm \text{ASE}$	$A_{0l} \pm \text{ASE}$	$k_{el} \pm \text{ASE}$	$T_{b1/2l} \pm \text{ASE}$	
Mn							
10	$56.93 \pm 3.44^{***}$	$2.14 \pm 0.53^{***}$	0.32 ± 0.08	$43.05 \pm 2.24^{***}$	$0.008 \pm 0.004^{***}$	78.93 ± 37.75	0.88
25	$61.63 \pm 2.20^{***}$	$1.62 \pm 0.18^{***}$	0.43 ± 0.05	$38.35 \pm 1.51^{***}$	$0.008 \pm 0.003^{***}$	85.92 ± 32.86	0.95
38	$75.44 \pm 1.61^{***}$	$2.91 \pm 0.40^{***}$	0.24 ± 0.03	$24.56 \pm 1.03^{***}$	$0.015 \pm 0.004^{***}$	46.24 ± 11.17	0.98
Zn							
10	$77.29 \pm 2.64^{***}$	$2.12 \pm 0.28^{***}$	0.33 ± 0.04	$22.71 \pm 1.68^{***}$	$0.006 \pm 0.006^{\text{NS}*}$	$+\infty$	0.95
25	$75.63 \pm 3.27^{***}$	$1.55 \pm 0.20^{***}$	0.47 ± 0.06	$24.39 \pm 2.20^{***}$	$0.005 \pm 0.007^{\text{NS}*}$	$+\infty$	0.93
38	$81.02 \pm 1.82^{***}$	$2.90 \pm 1.12^{***}$	0.18 ± 0.05	$18.98 \pm 1.12^{***}$	$0.008 \pm 0.007^{\text{NS}}$	$+\infty$	0.98

Probability of the model adjustment: $^{\text{NS}} p > 0.05$, $* p < 0.05$, $** p < 0.01$, $*** p < 0.001$