



A neural network approach to estimate water-column nutrient concentrations and carbonate system variables in the Mediterranean Sea: CANYON-MED

Marine Fourrier, Laurent Coppola, Fabrizio d'Ortenzio, Hervé Claustre,
Raphaëlle Sauzède, Henry C. Bittig, Marta Álvarez

► To cite this version:

Marine Fourrier, Laurent Coppola, Fabrizio d'Ortenzio, Hervé Claustre, Raphaëlle Sauzède, et al.. A neural network approach to estimate water-column nutrient concentrations and carbonate system variables in the Mediterranean Sea: CANYON-MED. EGU General Assembly 2019, Apr 2019, Vienne, Austria. hal-02552409

HAL Id: hal-02552409

<https://hal.science/hal-02552409>

Submitted on 23 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CANYON-MED

Contact:

fournier@obs-vlfr.fr



Marine Fourrier¹, Laurent Coppola¹, Fabrizio D'Ortenzio¹, Hervé Claustre¹, Raphaëlle Sauzède², Henry C. Bittig³, and Marta Álvarez⁴

(1) Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, LOV, F-06230 Villefranche-sur-Mer, France.

(3) Leibniz Institute for Baltic Sea Research, Department for Physical Oceanography and Instrumentation, Rostock-Warnemünde, Germany.

(2) Collecte Localisation Satellites, Ramonville-Saint-Agne, France.

(4) IEO Centro Oceanográfico de A Coruña, Coruña, Spain.

1– INTRODUCTION AND OBJECTIVES

The Mediterranean Sea is characterized by a short residence time (Somot et al., 2006), an elevated **salinity** and is one of the largest oceanic nutrient-depleted area (Ignatiades, 2005) exhibiting an eastward-increasing **oligotrophy** gradient (Pujo-Pay et al., 2011).

In the context of a critically undersampled ocean, the development and intensive use of instrumented *in situ* **autonomous platforms** allows to densify the measurements of some biogeochemical variables (remaining however far from exhaustive).

Recently, deep learning techniques have been developed for the global ocean. The **CANYON** (for Carbonate system and Nutrients concentration from hYdrological properties and Oxygen using a Neural-network; Sauzède et al., 2017) neural network-based method provides estimations of nutrients (i.e. nitrates, phosphates and silicates) and carbonate system parameters (i.e. total alkalinity, dissolved inorganic carbon, pH_T, pCO₂) from systematically measured oceanographic variables. The CANYON method produces **limited results in the Mediterranean Sea**.

Objectives of this study:

- Adapt the existing CANYON method to the Mediterranean Sea: **CANYON-MED neural networks**.
- Verify its accuracy on a **BGC-Argo float equipped with a nitrate sensor**.

3– NEURAL NETWORK METHOD

3.1– TRAINING

These Multi-Layered Perceptrons provide estimations of **nutrients** and **carbonate system variables** from systematically measured oceanographic variables such as *in situ* measurements of pressure, temperature, salinity, and oxygen together with geolocation (latitude, longitude) and date of sampling (day of year (doy), year).

Neural network ensembles comprising the 10 best topologies were created for each retrieved variable to reduce the uncertainties in the outputs.




Figure 2: Schematic representation of the CANYON-MED neural networks.

Adapted from Sauzède et al., 2017.

4 – CONCLUSION

The regional CANYON-MED neural networks constitute a **good downscaling** of CANYON, and satisfactory results are obtained: accuracies of **0.73, 0.043, and 0.63 µmol.kg⁻¹** for **NO₃⁻, PO₄³⁻ and Si(OH)₄** respectively, and **0.014, 17 µmol.kg⁻¹** and **12 µmol.kg⁻¹** for **pH_T, A_T and C_T** respectively. It can successfully be applied on **autonomous platforms**, such as profiling floats, gliders or moorings measuring T, S, O₂ to increase the amount of biogeochemical data in the Mediterranean Sea.

References: de Boyer Montégut, C. (2004). Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology. *Journal of Geophysical Research*, 109(C12). <https://doi.org/10.1029/2004JC002378> // Ignatiades, L. (2005). Scaling the trophic status of the Aegean Sea, eastern Mediterranean. *Journal of Sea Research*, 54(1), 51–57. <https://doi.org/10.1016/j.seares.2005.02.010> // Kessouri, F., Ulises, C., Estourmel, C., Marsaleix, P., Severin, T., Pujo-Pay, M., et al. (2017). Nitrogen and Phosphorus Budgets in the Northwestern Mediterranean Deep Convection Region. *Journal of Geophysical Research: Oceans*, 122(12), 9429–9455. <https://doi.org/10.1002/2016JC012665> // Lavigne, H. (2013). Impact of mixed layer depth seasonal variations on the phytoplankton phenology in the Mediterranean Sea. // Pujo-Pay, M., Conan, P., Oriol, L., Cornet-Barthaux, V., Falco, C., Ghiglione, J.-F., et al. (2011). Integrated survey of elemental stoichiometry (C, N, P) from the western to eastern Mediterranean Sea. *Biogeosciences*, 8(4), 883–899. <https://doi.org/10.5194/bg-8-883-2011> // Sauzède, R., Bittig, H. C., Claustre, O., Pasqueron de Fommervault, O., Gattuso, J.-P., Legendre, L., & Johnson, K. S. (2017). Estimates of Water-Column Nutrient Concentrations and Carbonate System Parameters in the Global Ocean: A Novel Approach Based on Neural Networks. *Frontiers in Marine Science*, 4. <https://doi.org/10.3389/fmars.2017.00128> // Somot, S., Sevault, F., & Déqué, M. (2006). Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model. *Climate Dynamics*, 27(7–8), 851–879. <https://doi.org/10.1007/s00382-006-0167-z>

2– MEDITERRANEAN SEA DATABASE

We compiled a **homogeneous quality-controlled** dataset of *in situ* measurements from 1981 to 2017, including samples of **core** variables such as dissolved oxygen (O₂), temperature (T), salinity (S) as well as **macronutrients** (nitrates: NO₃⁻, phosphates: PO₄³⁻, silicates: Si(OH)₄) and **carbonate system variables** (total alkalinity: A_T, total carbon: C_T, and pH on the total scale: pH_T).




Figure 1: Map of the 34 cruises and the time-series used for the training and validation of the CANYON-MED neural networks. (~7 000 values for NO₃⁻, ~10 000 for PO₄³⁻ and Si(OH)₄, and ~5 000 for A_T and pH_T, ~3 500 values for C_T).

Cruises	
ALMOFRONT	METEOR_51_2
BOUIM	METEOR_84_3
CARBGIB	MILLERO
CASCADE	MOOSE_GE_2010
DEWEX	MOOSE_GE_2011
DYFAMED	MOOSE_GE_2012
GEOSECS_LEG_3	MOOSE_GE_2013
GIFT	MOOSE_GE_2014
HOTMIX	MOOSE_GE_2015
MCXIS	MOOSE_GE_2016
MEDIPROD IV	OTRANTO
MEDSEA_2013	PACIFIC_CELEBES
	POEMXXIS
	PROSPE
	SESAME_IT01
	SESAME_IT_02
	SESAME_IT_04
	SESAME_IT_07
	SESAME_SPI
	SOMBA
	TRANSMED_LEGII

3.2– CANYON-MED IMPROVEMENT

The regional CANYON-MED highly **outperforms** the global CANYON method in the Mediterranean Sea. The difference stems from the difference in salinity, as well as the oligotrophic nutrient concentrations and low pH_T values found in the Mediterranean Sea.




Figure 3: Comparison of the CANYON and CANYON-MED retrieved values for the Mediterranean Sea with the corresponding in situ measurements for: NO₃⁻ (top left), PO₄³⁻ (top right), Si(OH)₄ (middle left), A_T (middle right), C_T (bottom left), pH_T (bottom right). The 1:1 line is shown.

	R ²	RMSE
NO ₃ ⁻ (µmol.kg ⁻¹)	0.77 0.96	1.57 0.64
PO ₄ ³⁻ (µmol.kg ⁻¹)	0.65 0.94	0.097 0.037
Si(OH) ₄ (µmol.kg ⁻¹)	0.4 0.96	2.9 0.6
A _T (µmol.kg ⁻¹)	0.74 0.96	32.4 11.4
C _T (µmol.kg ⁻¹)	0.78 0.95	24.8 10.8
pH _T	0.30 0.92	0.138 0.012

Table 1: Statistics (coefficient of determination (R²) and RMSE (Root Mean Squared Error)) of the regressions between the CANYON and CANYON-MED retrieved values with the corresponding in situ measurements.

3.3– CANYON-MED APPLICATION

CANYON-MED was applied on a BGC-Argo float, deployed in 2014 in the northwestern Mediterranean Sea, measuring T, S, O₂ as well as NO₃⁻.

Deep and surface NO₃⁻ values globally agree.

- Main trends are adequately predicted: e.g. water column mixing and **nutrients replenishment** in surface during the convection in the winter 2014–2015.
- Float and CANYON-MED derived **nitraclines** are in agreement with values established by Kessouri et al., 2017 (~60 m average for stratified periods).




Figure 4: Location of the 6901648 BGC-Argo float. The red star and yellow square represent the location of the 1st (07/07/2014) and last (26/05/2016) profile.




Figure 5: Timeseries of NO₃-float data (top) and CANYON-MED corrected-O₂ based NO₃- estimate (bottom), mixed-layer depth (MLD) (black) using the 0.03 criteria (de Boyer Montégut et al., 2004) and nitracline depth (white) using the 1µM isoline according to Lavigne (2013). NO₃- float data was corrected using the reference deployment profile's deep NO₃- value.