



**HAL**  
open science

## Exceptionally high $\delta^{15}\text{N}$ values in collagen single amino acids confirm Neandertals as high-trophic level carnivores

Klervia Jaouen, Michael P. Richards, Adeline Le Cabec, Frido Welker, William Rendu, Jean-Jacques Hublin, Marie Soressi, Sahra Talamo

### ► To cite this version:

Klervia Jaouen, Michael P. Richards, Adeline Le Cabec, Frido Welker, William Rendu, et al.. Exceptionally high  $\delta^{15}\text{N}$  values in collagen single amino acids confirm Neandertals as high-trophic level carnivores. *Proceedings of the National Academy of Sciences of the United States of America*, 2019, 116 (11), pp.4928-4933. 10.1073/pnas.1814087116 . hal-02552549

**HAL Id: hal-02552549**

**<https://hal.science/hal-02552549>**

Submitted on 23 Apr 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Exceptionally high $\delta^{15}\text{N}$ values in collagen single amino acids confirm Neandertals as high-trophic level carnivores

Klervia Jaouen<sup>a,1</sup>, Michael P. Richards<sup>b</sup>, Adeline Le Cabec<sup>a</sup>, Frido Welker<sup>a,c</sup>, William Rendu<sup>d</sup>, Jean-Jacques Hublin<sup>a,e</sup>, Marie Soressi<sup>f</sup>, and Sahra Talamo<sup>a</sup>

<sup>a</sup>Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany; <sup>b</sup>Department of Archaeology, Simon Fraser University, Burnaby BC V5A 1S6, Canada; <sup>c</sup>Evolutionary Genomics, Natural History Museum of Denmark, University of Copenhagen, DK-1353 Copenhagen, Denmark; <sup>d</sup>De la Préhistoire à l'Actuel: Culture, Environnement et Anthropologie, CNRS-University of Bordeaux, 33615 Pessac, France; <sup>e</sup>Chaire Internationale de Paléanthropologie, Collège de France, 75005 Paris, France; and <sup>f</sup>Faculty of Archaeology, Leiden University, 2311 EZ Leiden, The Netherlands

Edited by Richard G. Klein, Stanford University, Stanford, CA, and approved December 21, 2018 (received for review August 16, 2018)

Isotope and archeological analyses of Paleolithic food webs have suggested that Neandertal subsistence relied mainly on the consumption of large herbivores. This conclusion was primarily based on elevated nitrogen isotope ratios in Neandertal bone collagen and has been significantly debated. This discussion relies on the observation that similar high nitrogen isotopes values could also be the result of the consumption of mammoths, young animals, putrid meat, cooked food, freshwater fish, carnivores, or mushrooms. Recently, compound-specific C and N isotope analyses of bone collagen amino acids have been demonstrated to add significantly more information about trophic levels and aquatic food consumption. We undertook single amino acid C and N isotope analysis on two Neandertals, which were characterized by exceptionally high N isotope ratios in their bulk bone or tooth collagen. We report here both C and N isotope ratios on single amino acids of collagen samples for these two Neandertals and associated fauna. The samples come from two sites dating to the Middle to Upper Paleolithic transition period (Les Cottés and Grotte du Renne, France). Our results reinforce the interpretation of Neandertal dietary adaptations as successful top-level carnivores, even after the arrival of modern humans in Europe. They also demonstrate that high  $\delta^{15}\text{N}$  values of bone collagen can solely be explained by mammal meat consumption, as supported by archeological and zooarcheological evidence, without necessarily invoking explanations including the processing of food (cooking, fermenting), the consumption of mammoths or young mammals, or additional (freshwater fish, mushrooms) dietary protein sources.

compound-specific isotope analyses | diet | late Neanderthals | Paleolithic | stable isotopes

Since the first publication of bulk bone collagen carbon and nitrogen stable isotope ratios of Neandertals, (e.g., refs. 1–3), supported by more recent studies (4–8), an intriguing pattern has been observed: Neandertals are characterized by relatively high nitrogen isotope values, often higher than the carnivores from the same sites (Fig. 1). Nitrogen isotopes are usually employed as a tracer of trophic level (9), and this pattern was therefore, at first, interpreted as indicative of the top-level trophic position (TP) of Neandertals in Paleolithic food webs (2, 3, 10), specifically focused on the consumption of large herbivorous mammals (11). Early modern humans have a wider range of isotopic ratios (especially in carbon), and some individuals exhibit higher N isotope ratios than of the fauna associated with them and above the range generally observed in Neandertals (refs. 10 and 12–14 and Fig. 1). These elevated values in Pleistocene modern humans have been interpreted as the signature of freshwater fish consumption (10, 13) as N isotope ratios ( $\delta^{15}\text{N}$ ) in fish or shell consumers are usually higher than in pure terrestrial carnivores (10). Recently, an exceptionally high N isotope ratio has been measured

in an infant Neandertal (AR-14) from the site of Grotte du Renne (Arcy-sur-Cure, France; *SI Appendix, Supporting Information 1*). This individual had a  $\delta^{15}\text{N}$  value 3–5‰ above the associated carnivores (ref. 6 and Fig. 1). In this case, this very elevated ratio is likely to be due to breastfeeding (15), as the infant was ~1 y old.

Over the last few years, several studies have challenged the interpretation of high Neandertal bone collagen nitrogen isotope values as indicative of large herbivore consumption. It has been suggested that the systematic slightly higher  $\delta^{15}\text{N}$  values observed in Neandertal collagen, especially relative to that of the associated carnivores, could be explained by mammoth consumption (4), as mammoths generally exhibit higher  $\delta^{15}\text{N}$  values than other associated herbivores. Furthermore, recent studies carried on dental calculus highlight the existence of plant consumption (16–20), challenging the interpretation of a purely carnivorous diet of Neandertals. Other factors such as the consumption of putrid meat (21), mushrooms (16, 22, 23), freshwater fish (24), or cooked food (25, 26) have also been suggested as explanations for the elevated  $\delta^{15}\text{N}$  values of Neandertal collagen. The selective hunting of young animals by Neandertals and older animals by carnivores is another factor that would increase  $\delta^{15}\text{N}$  values, but the zooarcheological data generally

## Significance

Identifying past hominin diets is a key to understanding adaptation and biological evolution. Bone collagen isotope studies have added much to the discussion of Neandertal subsistence strategies, providing direct measures of diet. Neandertals consistently show very elevated nitrogen isotope values. These values have been seen as the signature of a top-level carnivore diet, but this interpretation was recently challenged by a number of additional theories. We here apply compound-specific isotope analysis of carbon and nitrogen in bone collagen single amino acids of two Neandertals. These Neandertals had the highest nitrogen isotope ratios of bulk collagen measured so far, and our study confirms that these values can be most parsimoniously explained by a carnivorous diet.

Author contributions: K.J. and S.T. designed research; K.J., A.L.C., F.W., and S.T. performed research; K.J. and F.W. contributed new reagents/analytic tools; K.J., M.P.R., A.L.C., F.W., W.R., M.S., and S.T. analyzed data; and K.J., M.P.R., A.L.C., F.W., W.R., J.-J.H., M.S., and S.T. wrote the paper.


The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

<sup>1</sup>To whom correspondence should be addressed. Email: klervia\_jaouen@eva.mpg.de.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1814087116/-DCSupplemental](https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1814087116/-DCSupplemental).



**Fig. 1.**  $\delta^{15}\text{N}$  values of Neandertals, Pleistocene modern humans, and fauna from different Paleolithic European and Asian sites for which analyses on the associated fauna was performed. Data are from (Feldhofer) Richards and Trinkaus (10), (Paviland) Richards et al. (3) and Devièse et al. (56), (Tianyan) Hu et al. (13), (Jonzac) Richards et al. (8), (Spy) Naito et al. (37), (Scladina) Bocherens et al. (2), (Goyet) Wissing et al. (7), (Marillac) Fizet et al. (1, 57), Bocherens et al. (5), (Oase) Trinkaus et al. (12), (Les Cottés) this study, (Grotte du Renne) Welker et al. (6), and (Buran Kaya III) Drucker et al. (30).

suggest the opposite pattern (27). Bocherens et al. (28) also suggested that the difference between Pleistocene modern humans and Neandertals would not be explained by distinctive diets, but by a climate change during the Middle to Upper Paleolithic transition causing the elevation of the N isotope signature of the local plants, impacting the values in the whole food web and, therefore, temporally distinct populations. Most of the modern humans analyzed so far are indeed from Upper Paleolithic contexts while the Neandertals (24, 28), with the exception of the Grotte du Renne (AR-14) (6) and the Spy Cave individuals (29), were found in Middle Paleolithic contexts. Finally, Drucker et al. (30) demonstrated, using compound-specific isotope analyses (CSIA), that some elevated  $\delta^{15}\text{N}$  values of Paleolithic modern human bone collagen could mainly be explained by mammoth consumption, rather than freshwater fish. The above-mentioned factors (Table 1) are all likely to increase the  $\delta^{15}\text{N}$  of body tissues without dramatically influencing the  $\delta^{13}\text{C}$  values. It is therefore very difficult to estimate which one accounts for the pattern observed in bulk Neandertal bone collagen.

Novel isotope techniques, such as compound-specific isotope analyses performed on single amino acids, could help identify the factor explaining this pattern. The CSIA are a very powerful technique, which consists in establishing the C and N isotope composition of the various amino acids in bone or tooth collagen. Carbon isotope values of individual amino acids can clearly distinguish terrestrial from freshwater food sources (31, 32). Honch et al. (31) demonstrated that animals from food webs based on terrestrial plants usually exhibit similar  $\delta^{13}\text{C}_{\text{Phe}}$  and  $\delta^{13}\text{C}_{\text{Val}}$  (Phe: Phenylalanine, Val: Valine) in their collagen, with higher values in  $\text{C}_4$  plant-based food webs than in  $\text{C}_3$  plant-based ones. Fish consumers have  $\delta^{13}\text{C}_{\text{Val}}$  overlapping with the animals from  $\text{C}_3$  plants (for freshwater environment) and  $\text{C}_4$  plants (for marine environment)-based food webs, but much lower  $\delta^{13}\text{C}_{\text{Phe}}$  values (31). Nitrogen isotope values of individual amino acids reveal TP in more detail than bulk isotopic data, especially using the data obtained on phenylalanine (Phe) and glutamic acid (Glu).  $\delta^{15}\text{N}_{\text{Phe}}$  reflects the local baseline without being really impacted by the trophic level of the animal (trophic  $^{15}\text{N}$ -enrichment of  $0.4 \pm 0.4\text{‰}$ ) (33) whereas the

glutamic acid N isotope ratio ( $\delta^{15}\text{N}_{\text{Glu}}$ ), in addition to the local baseline, is strongly impacted by the TP (trophic  $^{15}\text{N}$ -enrichment of  $8.0 \pm 1.1\text{‰}$ ) (34–36). The combination of the  $\delta^{15}\text{N}$  of these two amino acids therefore allows interpretations free of local baseline bias and allow more precisely assessing the TP of a specimen. For example, Naito et al. (37) documented the  $\delta^{15}\text{N}$  values of amino acids for the Spy Cave Neandertals (Fig. 1), revealing a possible contribution of plants into their diet, up to 20%. The use of CSIA of carbon and nitrogen can therefore potentially provide very detailed information on the diet of Neandertals and directly address the debate on the cause of the elevated nitrogen isotope signatures of their collagen.

Recently, we extracted the collagen of a Neandertal tooth from les Cottés (France; *SI Appendix, Supporting Information 1 and Fig. S1*, and ref. 38) for radiocarbon dating (ref. 39, *SI Appendix, Supporting Information 2*). In the frame of the assessment of collagen preservation, we measured the C and N isotope ratios in bulk collagen and found the second highest N isotope ratio ever seen in a Neandertal sample (*SI Appendix, Table S1*), knowing that the first one was the above mentioned breastfeeding infant from Grotte du Renne, AR-14 (6) (Fig. 1). The  $\delta^{15}\text{N}$  value of the Les Cottés Neandertal is about  $\sim 7\text{‰}$  higher than those of the associated herbivores previously measured in Talamo et al. (40), expected one-step trophic level enrichment (9). As there are a number of suggested reasons for the high Neandertal nitrogen isotope values (we list 10 possibilities in Table 1), we aimed to investigate which ones could account for the elevated value of the Les Cottés Neandertal. The tooth root analyzed belonged to a permanent maxillary lateral incisor, and thus, the dentine formed between the fourth and eighth years of age (*SI Appendix, Supporting Information 3*). If explained by breastfeeding, the  $\delta^{15}\text{N}$  value associated to the Les Cottés Neandertal would indicate a very late age of weaning, which would contradict previous findings (41, 42). At Les Cottés, hyena, wolf, and fox bones with cut-marks have been found in layer 06, layer 04/upper, and layer 02. Also, mammoth ivory transformed into beads have been found into layer 04/upper (43). Carnivore and mammoth consumption are two hypotheses that have to be

**Table 1.** List of factors likely to cause elevated nitrogen isotope ratios in tooth collagen of the Les Cottés Neandertal

Dietary adaptation	Output of the present study
Top-level carnivory	Yes
Mammoth consumption	Unknown
Young animal consumption	No/Minor
Freshwater fish consumption	No
Breastfeeding	Unlikely
Environmental factors triggering an elevation of the whole food web isotope values	No
Mushroom consumption	Unknown
Consumption of putrid meat	Unknown
Cooking	Unknown
Carnivore consumption	No/Minor

The second column lists the relevant factors according to the results of this study. See text for more details.

considered. Fish remains are not present at les Cottés. No putrid meat or mushroom consumption has been reported for this site from the archeological evidence; however, these two types of consumption would unlikely leave any significant archeological traces. Concerning the environmental hypothesis suggested by Bocherens et al. (28), the tooth was dated to 43,740–42,720 cal. y BP (*SI Appendix, Supporting Information 2*) and the Neandertal should thus have lived before the hypothesized increased aridity, which was suggested to have impacted local isotope signatures of the whole food webs coming from transition layers in France according to Bocherens et al. (28).

We conducted N and C isotope analyses on bulk collagen and amino acids of the Les Cottés Neandertal and the associated fauna, which were identified using ZooMs (*Dataset S1, Table S5*), to determine which one of the abovementioned factors (Table 1) accounts for the exceptionally elevated  $\delta^{15}\text{N}$  value of the Les Cottés Neandertal. As no CSIA data are yet available for breastfeeding or suckling specimens, we also performed these analyses on (*i*) animal teeth from Les Cottés formed before the weaning age, and (*ii*) faunal and hominin remains from Grotte du Renne, including the breastfed Neandertal (AR-14). The C and N isotope data on bulk collagen of the Grotte du Renne food web are already published (6). The Grotte du Renne and Les Cottés Neandertals both have similar radiocarbon ages and come from Central France contexts, which reinforce the validity of the comparison of these two Neandertals (6, 39).

Below, we report unprecedented CSIA for carbon on collagen amino acids for Paleolithic collagen samples, as well as the second CSIA application for nitrogen on collagen amino acids for Neandertals. We suggest that the exceptionally high N isotope signature values of the Les Cottés Neandertal can simply be explained by a carnivorous diet, possibly preferentially relying on reindeer (*Rangifer tarandus*). We also confirm that the Grotte du Renne Neandertal (AR-14) was a suckling infant and exclude the possibility of freshwater fish consumption being an influence on the collagen isotope values at both sites.

## Results

The raw data for taxonomic identification, bulk collagen and amino acid isotope analyses, as well as measurements on isotopic standards are given in the Supporting Tables (*Dataset S1, Tables S5–S10*).

**Isotope Analyses of the Bulk Collagen.** Carbon and nitrogen isotope analyses of bulk bone collagen are the classic method to estimate trophic relationship in food webs. C and N isotope analyses on the bulk collagen ( $\delta^{13}\text{C}_{\text{bulk}}$  and  $\delta^{15}\text{N}_{\text{bulk}}$ ) for the AR-14 Neandertal and the associated fauna of Grotte du Renne were previously published in Welker et al. (6). The results for the Les Cottés food web are here presented. The collagen extracted from the Les Cottés samples is well preserved, with collagen yields of more than 1% and C:N ratios ranging from 3.1 to 3.3 (ref. 44 and *Dataset S1, Tables S6 and S7*). The herbivore bones and teeth from Les Cottés have  $\delta^{13}\text{C}_{\text{bulk}}$  ranging between  $-20.6$  and  $-19.8\text{‰}$ , except for three *Rangifer* specimens exhibiting higher values, and the mammoth specimen exhibiting the lowest  $\delta^{13}\text{C}_{\text{bulk}}$  value (Fig. 2). The  $\delta^{15}\text{N}_{\text{bulk}}$  of the herbivores are comparable to what was observed in Grotte du Renne, except for the specimens with teeth, which would have formed during the suckling period, and that have, as expected, higher  $\delta^{15}\text{N}_{\text{bulk}}$  values. The mammoth specimen also has an elevated  $\delta^{15}\text{N}_{\text{bulk}}$  value, higher than in Grotte du Renne, but this pattern is generally observed for this species (7, 28, 30). The only carnivore species analyzed at Les Cottés beside the Neandertal is hyenas. Hyena teeth formed during or after suckling periods reveal a large range of  $\delta^{15}\text{N}_{\text{bulk}}$  values ( $9.0$ – $11.9\text{‰}$ ) and of  $\delta^{13}\text{C}_{\text{bulk}}$  values ( $-19.8$  to  $-18.2\text{‰}$ ). The hyena P3 teeth, possibly formed during the suckling period, do not show higher  $\delta^{15}\text{N}_{\text{bulk}}$  values than the other teeth. One of the hyenas has a  $\delta^{15}\text{N}_{\text{bulk}}$  comparable to that of the *Rangifer* teeth. The Neandertal tooth from Les Cottés falls in the range of that of the carnivore hyenas, both for C and N isotope ratios.


**Nitrogen Isotope Analyses of Amino Acids (Les Cottés and Grotte du Renne) and Associated TP.** Nitrogen isotope analyses of single amino acids were conducted to estimate the TP of hominins and associated animals, and assess which species were hunted by the different carnivores. These analyses can unravel the presence of plants, fish, meat, and suckling in the diet of the different animals. Terrestrial herbivores have a wider range of  $\delta^{15}\text{N}_{\text{Phe}}$  values in Grotte du Renne ( $7.6$ – $14.4\text{‰}$ ) than in Les Cottés ( $7.3$ – $12.4\text{‰}$ ). The  $\delta^{15}\text{N}_{\text{Phe}}$  values of the herbivores do not correlate with the bulk collagen  $\delta^{15}\text{N}$  values, whereas the  $\delta^{15}\text{N}_{\text{Glu}}$  values do. This correlation, both in Grotte du Renne and Les Cottés, indicates that most of the N isotopic variability among herbivores is due to a trophic level effect, likely caused by the sampling of tissues formed during the suckling period or shortly after weaning age. The  $\delta^{15}\text{N}_{\text{Phe}}$  value of the Neandertal in les Cottés is clearly higher than that of the other carnivores, while the  $\delta^{15}\text{N}_{\text{Phe}}$  value of the AR-14 Neandertal falls in the range of the other carnivores (Fig. 3). The TP of all of the animals has been assessed using  $\delta^{15}\text{N}_{\text{Glu}}$ ,  $\delta^{15}\text{N}_{\text{Phe}}$  values and a  $\beta$  factor of  $-8.4\text{‰}$  (refs. 33, 36, and 37 and *Dataset S1, Table S6*). The  $\beta$  factor corresponds to an enrichment factor between the  $\delta^{15}\text{N}$  in glutamic acid and phenylalanine ( $\delta^{15}\text{N}_{\text{Glu}} - \delta^{15}\text{N}_{\text{Phe}}$ ). This factor therefore reflects the vertical shift between each line representing a different TP in Figs. 2 and 3. The uncertainty on the estimation of the TP is probably related to the variability of the  $\beta$  factor, which is around 0.1 (30), and therefore yields range between 1.9 and 2.1 for herbivores and between 2.9 and 3.1 for carnivores. Several herbivores have a TP above 2.1 and one hyena equals 3.1 while the other is above this value, which corresponds to the teeth (P3, M1) and bones possibly formed during the suckling period. However, some herbivores have a TP below 1.9 (two Equidae and one *Rangifer* at Les Cottés, and two Equidae and one cave bear at Grotte du Renne). Moreover, one of the four hyenas of les Cottés has a TP of 2.2, similar to what is expected for a herbivore, and three carnivores at Grotte du Renne (two Pantherinae and a wolf) have, respectively, a TP of 2.5, 2.7, and 2.8 (Figs. 2 and 3 and *SI Appendix, Table S6*). The Les Cottés Neandertal has a TP of 2.9, indicating a pure carnivorous diet. The Grotte du Renne Neandertal has a TP of 3.2, slightly higher than the theoretical carnivore TP and much higher than the carnivores from this site.

**Carbon Isotope Analyses of Amino Acids (Les Cottés and Grotte du Renne).** Carbon isotope analyses of amino acids can help to identify the presence of freshwater or marine resources into the diet of past hominins. The carbon isotope composition of 9–12 amino acids (depending on the derivatization method used) has been measured for all of the specimens of les Cottés and Grotte du Renne, with the exception of the Neandertal from the second site for which not enough material was available. The data are given in *Dataset S1, Table S8* and fully discussed in the *SI Appendix, Supporting Information 4*. The  $\delta^{13}\text{C}_{\text{Phe}}$  ( $-24.4$  to  $-28.3\text{‰}$ ) and  $\delta^{13}\text{C}_{\text{Val}}$  ( $-23.2$  to  $-28.9\text{‰}$ ) range of values are similar at both sites and generally fit with the range observed previously in food webs relying on  $\text{C}_3$  plants (31, 32), but slightly above the average food web values reported in a third study (40). These three publications represent the totality of the existing data on  $\delta^{13}\text{C}_{\text{Phe}}$  and  $\delta^{13}\text{C}_{\text{Val}}$  in mammal collagen. The pattern observed between the two sites is quite different: In Grotte du Renne, the carnivores exhibit similar  $\delta^{13}\text{C}$  values in the phenylalanine and valine, while the herbivores have enriched  $\delta^{13}\text{C}_{\text{Phe}}$  values relative to  $\delta^{13}\text{C}_{\text{Val}}$  (Figs. 3 and 4). In Les Cottés, most herbivores have also enriched  $\delta^{13}\text{C}_{\text{Phe}}$  relative to  $\delta^{13}\text{C}_{\text{Val}}$ , but to a lesser extent (Figs. 2 and 4). In this case, the carnivores have depleted  $\delta^{13}\text{C}_{\text{Phe}}$  relative to  $\delta^{13}\text{C}_{\text{Val}}$ , especially in the case of the two hyena teeth formed during the suckling period (Fig. 4).

## Discussion

**TP Uncertainties.** In Les Cottés, the estimated TPs of the animals are consistent with what is expected, except for three herbivores





**Fig. 2.** Isotope results for the site of Les Cottés. (A) C and N isotope ratios in bulk collagen. (B) N isotope ratios in phenylalanine and glutamic acid. The TP lines (B) are defined according to Chikaraishi et al. (35, 36), (C) C isotope ratios in valine and phenylalanine. The green and yellow areas (C) are defined according to Honch et al. (31).


and one hyena. This could be explained by the consumption of plants with a peculiar  $\beta$  factor by these three herbivores. Most plants (50%) have a  $\beta$  factor of  $-8 \pm 1\text{‰}$ , yet some reach a value of  $-12\text{‰}$  and represent 3% of the reported values (45).

However, only 64  $\beta$  values of plants have been analyzed so far, and there is therefore a need to better document those  $\beta$  factors in plants and the species associated to specific values (e.g., 3‰ in the case of tree flowers). With such a  $\beta$  value, the trophic level of the three herbivores goes up to 2.1–2.2 and to 2.8 for the associated hyena. In Grotte du Renne, this  $\beta$  moves back these herbivores (two Equidae and one Cave bear) as well as the Pantharinae to their expected TP. With the same  $\beta$  factor, the Neandertal from Grotte du Renne would then have a TP of 3.8, one trophic level above the carnivores. Such a TP could easily be explained by breastfeeding.


We observed difference between  $\delta^{13}\text{C}_{\text{Val}}$  and  $\delta^{13}\text{C}_{\text{Phe}}$  values, which can be explained by the existence of a trophic level effect, mostly affecting the  $\delta^{13}\text{C}_{\text{Val}}$  (SI Appendix, Supporting Information 4) rather than freshwater fish consumption. This accounts for the fact that the Neandertal from Les Cottés and the hyenas are falling slightly off the range defined for terrestrial animals. As a result, the difference between  $\delta^{13}\text{C}_{\text{Val}}$  and  $\delta^{13}\text{C}_{\text{Phe}}$  ( $\Delta^{13}\text{C}_{\text{Val-Phe}}$ ) values correlate to the estimated trophic level (Fig. 4), and the  $\Delta^{13}\text{C}_{\text{Val-Phe}}$  is systematically positive in carnivores and negative for herbivores. The report of this trophic level effect is unprecedented, since no other carnivore data are currently available for  $\delta^{13}\text{C}_{\text{Val}}$  and  $\delta^{13}\text{C}_{\text{Phe}}$  in mammal collagen. We also observed it for  $\delta^{13}\text{C}_{\text{Leu}}$  and  $\delta^{13}\text{C}_{\text{Gly}}$  values (Gly = Glycine, Leu = Leucine; SI Appendix, Supporting Information 4). The combined use of carbon and nitrogen single amino acid isotope values could therefore help to account for the uncertainty on the TP estimation based on  $\delta^{15}\text{N}$  values only.

**Neandertal Diets at Les Cottés and Grotte du Renne.** Compound-specific isotope analyses conducted on amino acids provide information on the sources of dietary protein over a number of

years of life with a greater level of detail compared with bulk isotope analysis. However, as these CSIA studies are in their infancy, the exact relationship of  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values of amino acids to TPs is not yet fully understood. Despite these caveats, these analyses shed light on the interpretation of Neandertal diets in Europe. We demonstrate here that exceptionally high  $\delta^{15}\text{N}$  values in bulk collagen of the Les Cottés Neandertal can still be explained by a carnivorous diet solely relying on herbivore meat without any need to call upon additional explanations such as consumption of putrid meat or cooked food. These types of consumption might, however, have existed but would mean that fermentation and cooking equally impact  $\delta^{15}\text{N}_{\text{Phe}}$  and  $\delta^{15}\text{N}_{\text{Glu}}$  values (Table 1). Freshwater fish and carnivore meat were not—or rarely—eaten by these Neandertals. Neandertals at les Cottés could have mostly hunted reindeer from arid environments (showing elevated  $\delta^{15}\text{N}_{\text{Phe}}$  and  $\delta^{13}\text{C}_{\text{bulk}}$ ) or horses (also characterized by elevated  $\delta^{13}\text{C}_{\text{Asp}}$ , SI Appendix, Supporting Information 4), whereas hyenas would have been less specific in terms of their environments and prey. This is a similar interpretation to that proposed by the only other Neandertal CSIA study by Naito et al. (37), which also suggested different ecological niches between hyenas and Neandertals. The  $\delta^{13}\text{C}_{\text{bulk}}$  value of the Neandertal tooth does not support here the specific hunting of mammoths, which would generally yield low  $\delta^{13}\text{C}$  (as observed in Grotte du Renne, although only one individual could be analyzed). According to CSIA analysis, the Spy Neandertals (Neandertals dated circa 41,000 BP; ref. 46) show different hunting strategies: Spy 430-a and Spy 92b hunted reindeer, horse, and bovines, while Spy 92a seemed more opportunistic (37). Neandertal hunting of reindeer, rather than mammoths, is also generally supported by zooarchaeological data (27, 47) including at Grotte du Renne and Les Cottés (38, 48, 49), even if some exceptions exist for Middle Paleolithic sites (47). In Grotte du Renne, both  $\delta^{15}\text{N}_{\text{Phe}}$  and  $\delta^{13}\text{C}$  values of AR-14 bulk collagen



**Fig. 3.** Isotope results for the site of Grotte du Renne (Arcy-sur-Cure). (A) C and N isotope ratios in bulk collagen; data from Welker et al. (6). (B) N isotope ratios in phenylalanine and glutamic acid. The TP lines (B) are defined according to Chikaraishi et al. (35, 36); (C) C isotope ratios in valine and phenylalanine. The green and yellow areas (C) are defined according to Honch et al. (31).



**Fig. 4.** Difference (%) between  $\delta^{13}\text{C}_{\text{Val}}$  and  $\delta^{13}\text{C}_{\text{Phe}}$  ( $\Delta^{13}\text{C}_{\text{Val-Phe}}$ ) in bone and tooth collagen and associated TP. The TP was estimated from the  $\delta^{15}\text{N}_{\text{Phe}}$  and  $\delta^{15}\text{N}_{\text{Val}}$  values. The C isotope ratios in amino acids of the Grotte du Renne Neandertal were not measured because not enough material was available. See text for more details.

falls in the variability of the other carnivores. The TP of the AR-14 Neandertal is clearly above that of the other carnivores, whatever the  $\beta$  factor is. This finding fits with the hypothesis of a suckling infant previously suggested by Welker et al. (6). This infant was therefore not weaned but could possibly already have eaten solid food, as its TP is below 4. According to the isotope data performed on bulk collagen and amino acids, the mother was probably sharing the same ecological niche as the other carnivores.

When we then look at all of the bulk collagen isotope analyses of the 29 Neandertals that have been published so far [ranging in radiometric age from  $\sim 90,000$  BP (Scladina; ref. 46) to 36,840 BP (Grotte du Renne; ref. 6)], we can see a similar and very stable diet over time. This situation could be readily explained by Neandertals being top-level carnivores consuming herbivore meat, and this pattern even continues after the arrival of modern humans in Europe (50). Our work shows that even exceptionally high  $\delta^{13}\text{N}$  bulk collagen values of two late Neandertals still result from a terrestrial meat-based diet, the same diet that was interpreted for the other 27 Neandertals that have had bulk collagen C and N isotope measurements. Despite changes in climate, environment and associated faunal spectra, throughout the Middle Paleolithic, the focus remains on large herbivores.

- Fizet M, et al. (1995) Effect of diet, physiology and climate on carbon and nitrogen stable isotopes of collagen in a late Pleistocene anthropic palaeoecosystem: Marillac, Charente, France. *J Archaeol Sci* 22:67–79.
- Bocherens H, et al. (1999) Palaeoenvironmental and palaeodietary implications of isotopic biogeochemistry of last interglacial Neandertal and mammal bones in Scladina Cave (Belgium). *J Archaeol Sci* 26:599–607.
- Richards MP, et al. (2000) Neandertal diet at Vindija and Neandertal predation: The evidence from stable isotopes. *Proc Natl Acad Sci USA* 97:7663–7666.
- Bocherens H (2011) Diet and ecology of Neandertals: Implications from C and N isotopes. *Neandertal Lifeways, Subsistence and Technology, Vertebrate Paleobiology and Paleoanthropology Series* (Springer, Dordrecht, The Netherlands), pp 73–85.
- Bocherens H, Drucker DG, Billiou D, Patou-Mathis M, Vandermeersch B (2005) Isotopic evidence for diet and subsistence pattern of the Saint-Césaire I Neandertal: Review and use of a multi-source mixing model. *J Hum Evol* 49:71–87.
- Welker F, et al. (2016) Palaeoproteomic evidence identifies archaic hominins associated with the Châtelperronian at the Grotte du Renne. *Proc Natl Acad Sci USA* 113:11162–11167.
- Wissing C, et al. (2016) Isotopic evidence for dietary ecology of late Neandertals in North-Western Europe. *Quat Int* 411:327–345.
- Richards MP, et al. (2008) Isotopic dietary analysis of a Neandertal and associated fauna from the site of Jonzac (Charente-Maritime), France. *J Hum Evol* 55:179–185.
- Hedges REM, Reynard LM (2007) Nitrogen isotopes and the trophic level of humans in archaeology. *J Archaeol Sci* 34:1240–1251.
- Richards MP, Trinkaus E (2009) Isotopic evidence for the diets of European Neandertals and early modern humans. *Proc Natl Acad Sci USA* 106:16034–16039.
- Bocherens H (2009) Neandertal dietary habits: Review of the isotopic evidence. *The Evolution of Hominin Diets, Vertebrate Paleobiology and Paleoanthropology* (Springer, Dordrecht, The Netherlands), pp 241–250.

**Concluding Remarks.** Compound-specific isotope analyses are a very promising method for reconstructing aspects of past diets, especially by combining nitrogen and carbon isotope analyses on single amino acids to clearly distinguish aquatic and terrestrial sources of proteins, and to estimate with more precision the contribution of plants in individual diets. Our application of CSIA to two Neandertals has shown that both were high trophic level consumers, with large herbivores being the main protein source. Therefore, there is no reason to invoke myriad dietary interpretations such as the consumption of mushrooms, putrid animal flesh, mammoths, or freshwater fish to explain their high bulk bone collagen nitrogen isotope values of Neandertals in relation to carnivores from the same site. However, we acknowledge that our isotopic study does not rule out the occasional consumption of these foods. Instead, the differing high N values between Neandertals and associated carnivores is likely due to the consumption of different herbivores from different environments. When we conduct CSIA, which is not environmentally baseline sensitive like bulk collagen N values, Neandertals and carnivores clearly are at the same trophic level.

## Methods

The bones and teeth were taxonomically identified using traditional zooarchaeological methods and ZooMS (6, 49, 51) (Dataset S1, Table S5). The collagen extraction was performed using the protocol outlined in Talamo and Richards (44), and the bulk collagen isotope analyses were conducted at the Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany. The collagen samples were then sent to the University of California, Los Angeles Isotope Lab Facility, where the compound-specific isotope analyses were performed using gas chromatography combustion isotope ratio mass spectrometry. The two first batches of samples were prepared with the following protocol: The amino acids were liberated using acid hydrolysis before the derivatization step to produce methoxycarbonyl methyl esters (MOC) for GC analysis, following the protocol established by Yarnes and Herzog (52). The third batch of samples was prepared using the derivatization protocol (53) recommended by O'Connell and Collins (54), producing N-acetyl isopropyl esters, allowing the comparison of isotope signatures in hydroxyproline and proline (SI Appendix, Fig. S2). An external standard was analyzed with both protocols and exhibit similar isotope signatures for both protocols (Dataset S1, Table S10). All samples were analyzed in duplicate by GC-C-IRMS, and additional measurements were performed when the duplicates fell outside expected measurement error (55). Additional information is available in Dataset S1.

**ACKNOWLEDGMENTS.** We thank Geoffrey Smith, Karen Ruebens, Chris Yarnes, and Nicolas Bourgon for helpful discussions; Lysann Klausnitzer and Sven Steinbrenner for technical help with the collagen extraction and bulk isotope analyses; J.-J. Cleyet-Merle (Musée National de Préhistoire) for providing samples; and Heiko Temming for  $\mu\text{CT}$ -scanning of the tooth prior to destructive sampling. This study was funded by the Max Planck Society.

- Trinkaus E, et al. (2009) Stable isotope evidence for early modern human diet in southeastern Europe: Peștera cu Oase, Peștera Muierii and Peștera Cioclovina Uscată. *Mater Cercet Arheol* 5:4–14.
- Hu Y, et al. (2009) Stable isotope dietary analysis of the Tianyuan 1 early modern human. *Proc Natl Acad Sci USA* 106:10971–10974.
- Fu Q, et al. (2014) Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* 514:445–449.
- Fuller BT, Fuller JL, Harris DA, Hedges RE (2006) Detection of breastfeeding and weaning in modern human infants with carbon and nitrogen stable isotope ratios. *Am J Phys Anthropol* 129:279–293.
- Weyrich LS, et al. (2017) Neandertal behaviour, diet, and disease inferred from ancient DNA in dental calculus. *Nature* 544:357–361.
- Henry AG, Brooks AS, Piperno DR (2011) Microfossils in calculus demonstrate consumption of plants and cooked foods in Neandertal diets (Shanidar III, Iraq; Spy I and II, Belgium). *Proc Natl Acad Sci USA* 108:486–491.
- Henry AG, Brooks AS, Piperno DR (2014) Plant foods and the dietary ecology of Neandertals and early modern humans. *J Hum Evol* 69:44–54.
- Hardy BL, Moncel M-H (2011) Neandertal use of fish, mammals, birds, starchy plants and wood 125–250,000 years ago. *PLoS One* 6:e23768.
- Hardy K, et al. (2012) Neandertal medics? Evidence for food, cooking, and medicinal plants entrapped in dental calculus. *Naturwissenschaften* 99:617–626.
- Speth JD (2017) Putrid meat and fish in the Eurasian Middle and Upper Paleolithic: Are we missing a key part of Neandertal and modern human diet? *Paleoanthropology* 2017:44–72.
- Hardy BL, et al. (2013) Impossible Neandertals? Making string, throwing projectiles and catching small game during marine isotope stage 4 (Abri du Maras, France). *Quat Sci Rev* 82:23–40.

23. O'Regan HJ, Lamb AL, Wilkinson DM (2016) The missing mushrooms: Searching for fungi in ancient human dietary analysis. *J Archaeol Sci* 75:139–143.
24. Pearson J (2007) Hunters, fishers and scavengers. *Farming* 2007:1–16.
25. Royer A, Daux V, Fourel F, Lécuyer C (2017) Carbon, nitrogen and oxygen isotope fractionation during food cooking: Implications for the interpretation of the fossil human record. *Am J Phys Anthropol* 163:759–771.
26. DeNiro MJ, Schoeninger MJ, Hastorf CA (1985) Effect of heating on the stable carbon and nitrogen isotope ratios of bone collagen. *J Archaeol Sci* 12:1–7.
27. Gaudzinski-Windheuser S, Roebroeks W (2011) On Neanderthal subsistence in last interglacial forested environments in Northern Europe. *Neanderthal Lifeways, Subsistence and Technology*, Vertebrate Paleobiology and Paleoanthropology Series (Springer, Dordrecht, The Netherlands), pp 61–71.
28. Bocherens H, Drucker DG, Madelaine S (2014) Evidence for a (15)N positive excursion in terrestrial foodwebs at the Middle to Upper Palaeolithic transition in south-western France: Implications for early modern human palaeodiet and palaeoenvironment. *J Hum Evol* 69:31–43.
29. Semal P, et al. (2009) New data on the late Neandertals: Direct dating of the Belgian Spy fossils. *Am J Phys Anthropol* 138:421–428.
30. Drucker DG, et al. (2017) Isotopic analyses suggest mammoth and plant in the diet of the oldest anatomically modern humans from far southeast Europe. *Sci Rep* 7:6833.
31. Honch NV, McCullagh JS, Hedges RE (2012) Variation of bone collagen amino acid  $\delta^{13}\text{C}$  values in archaeological humans and fauna with different dietary regimes: Developing frameworks of dietary discrimination. *Am J Phys Anthropol* 148:495–511.
32. Colonese AC, et al. (2014) Long-term resilience of late Holocene coastal subsistence system in Southeastern South America. *PLoS One* 9:e93854.
33. Chikaraishi Y, Ogawa NO, Ohkouchi N (2010) Further evaluation of the trophic level estimation based on nitrogen isotopic composition of amino acids. *Earth, Life, Isotopes* (Kyoto Univ Press, Kyoto, Japan), pp 37–51.
34. McClelland JW, Montoya JP (2002) Trophic relationships and the nitrogen isotopic composition of amino acids in plankton. *Ecology* 83:2173–2180.
35. Chikaraishi Y, et al. (2009) Determination of aquatic food-web structure based on compound-specific nitrogen isotopic composition of amino acids. *Limnol Oceanogr Methods* 7:740–750.
36. Chikaraishi Y, et al. (2014) High-resolution food webs based on nitrogen isotopic composition of amino acids. *Ecol Evol* 4:2423–2449.
37. Naito YI, et al. (2016) Ecological niche of Neanderthals from Spy Cave revealed by nitrogen isotopes of individual amino acids in collagen. *J Hum Evol* 93:82–90.
38. Soressi M, et al. (2010) Les Cottés (Vienne). Nouveaux travaux sur l'un des gisements de référence pour la transition Paléolithique moyen/supérieur. *Préhistoire entre Vienne et Charente: Hommes et société du Paléolithique* (Association des Publications Chauvinoises, Chauvigny, France), pp 221–234.
39. Hajdinjak M, et al. (2018) Reconstructing the genetic history of late Neanderthals. *Nature* 555:652–656.
40. Talamo S, Soressi M, Roussel M, Richards M, Hublin J-J (2012) A radiocarbon chronology for the complete Middle to Upper Palaeolithic transitional sequence of Les Cottés (France). *J Archaeol Sci* 39:175–183.
41. Skinner M (1997) Dental wear in immature late Pleistocene European hominines. *J Archaeol Sci* 24:677–700.
42. Smith TM, et al. (2010) Dental evidence for ontogenetic differences between modern humans and Neanderthals. *Proc Natl Acad Sci USA* 107:20923–20928.
43. Rigaud S, et al. (2014) Les pratiques ornementales à l'Aurignacien ancien dans le Centre-Ouest de la France: L'apport des fouilles récentes aux Cottés (Vienne). *Bull Soc Préhist Fr* 111:19–38.
44. Talamo S, Richards M (2011) A comparison of bone pretreatment methods for AMS dating of samples >30,000 BP. *Radiocarbon* 53:443–449.
45. Naito YI, et al. (2017) Reply to "Comment on 'Ecological niche of Neanderthals from Spy Cave revealed by nitrogen isotopes of individual amino acids in collagen.'" [J. Hum. Evol. 93 (2016) 82–90] [J. Hum. Evol. 117 (2018) 53–55]. *J Hum Evol* 117:56–60.
46. Pirson S, Toussaint M, Bonjean D, Di Modica K (2018) Spy and Scladina caves: A Neanderthal's story. *Landscapes and Landforms of Belgium and Luxembourg* (Springer, Berlin), pp 357–383.
47. Smith GM (2015) Neanderthal megafaunal exploitation in Western Europe and its dietary implications: A contextual reassessment of La Cotte de St Brelade (Jersey). *J Hum Evol* 78:181–201.
48. David F, et al. (2001) Le Châtelperronien de la grotte du Renne à Arcy-sur-Cure (Yonne). Données sédimentologiques et chronostratigraphiques. *Bull Soc Préhist Fr* 98:207–230.
49. Welker F, Soressi M, Rendu W, Hublin J-J, Collins M (2015) Using ZooMS to identify fragmentary bone from the late Middle/early Upper Palaeolithic sequence of Les Cottés, France. *J Archaeol Sci* 54:279–286.
50. Higham T, et al. (2011) The earliest evidence for anatomically modern humans in northwestern Europe. *Nature* 479:521–524.
51. Buckley M, Collins M, Thomas-Oates J, Wilson JC (2009) Species identification by analysis of bone collagen using matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry. *Rapid Commun Mass Spectrom* 23:3843–3854.
52. Yarnes CT, Herszage J (2017) The relative influence of derivatization and normalization procedures on the compound-specific stable isotope analysis of nitrogen in amino acids. *Rapid Commun Mass Spectrom* 31:693–704.
53. Styring AK, et al. (2012) Practical considerations in the determination of compound-specific amino acid  $\delta^{15}\text{N}$  values in animal and plant tissues by gas chromatography combustion-isotope ratio mass spectrometry, following derivatisation to their N-acetylisopropyl esters. *Rapid Commun Mass Spectrom* 26:2328–2334.
54. O'Connell TC, Collins MJ (2017) Comment on "Ecological niche of Neanderthals from Spy Cave revealed by nitrogen isotopes of individual amino acids in collagen" [J. Hum. Evol. 93 (2016) 82–90]. *J Hum Evol* 117:53–55.
55. Docherty G, Jones V, Evershed RP (2001) Practical and theoretical considerations in the gas chromatography/combustion/isotope ratio mass spectrometry  $\delta^{13}\text{C}$  analysis of small polyfunctional compounds. *Rapid Commun Mass Spectrom* 15:730–738.
56. Deviese T, et al. (2017) Direct dating of Neanderthal remains from the site of Vindija Cave and implications for the Middle to Upper Paleolithic transition. *Proc Natl Acad Sci USA* 114:10606–10611.
57. Fizet M, Lange-Badre B, Vandermeersch B, Borel JP, Bellon G (1991) Isotopic biogeochemistry ( $^{13}\text{C}$ ,  $^{15}\text{N}$ ) of fossil vertebrate collagen: Application to the study of a past food web including Neanderthal man. *J Hum Evol* 20:481–492.



## Supporting Information for

Exceptionally high  $\delta^{15}\text{N}$  values in collagen single amino acids confirm Neandertals as high-trophic level carnivores

Klervia Jaouen, Michael P. Richards, Adeline Le Cabec, Frido Welker, William Rendu, Jean-Jacques Hublin, Marie Soressi, Sahra Talamo.

Corresponding author: Klervia Jaouen

Email: [klervia\\_jaouen@eva.mpg.de](mailto:klervia_jaouen@eva.mpg.de)



**This PDF file includes:**

Supplementary text

Figs. S1 to S7

Tables S1 to S4

SI reference citations


Data table captions S5 to S10

**Other supplementary materials for this manuscript include the following:**

Data Tables S5 to S10

## Supporting Information 1

### Site location



**Figure S1: Location of the two French Neanderthal sites (Les Cottés and Grotte du Renne at Arcy-sur-Cure) studied in this paper.** The green area represents the geographical extent of Châtelperronian transition sites in France. Map modified from Soressi et al. (1).

## Supporting Information 2

### Radiocarbon dating

The tooth root sample was pretreated at the Department of Human Evolution at the Max Planck Institute for Evolutionary Anthropology (MPI-EVA) in Leipzig, Germany, using the method described in Talamo and Richards (2). The sample passed the collagen evaluation criteria (see Table S1), and was sent to the Mannheim AMS laboratory (Lab Code MAMS) where it was graphitized and dated (3)

MPI Code	Used mg	Yield mg	% Yield	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	%C	%N	C:N	AMS Lab Code	$^{14}\text{C}$ Age	1s Err
R-EVA 1526	461	25.7	5.6	-18.1	13.9	44.0	15.9	3.2	MAMS- 26196	39485	271

**Table S1. Radiocarbon dating (uncalibrated) of Les Cottés Z4-1514 Neandertal tooth and collagen evaluation criteria**

## Supporting Information 3

### 1. Anatomical determination of the Les Cottés tooth (Z4-1514).

Although the crown was entirely worn away - even beyond the cervix - the Les Cottés tooth (CTS 2008 US 04.35 Z4-1514) can be described as a single-rooted tooth, most likely an anterior permanent tooth (i.e., a central or lateral incisor or a canine). An anatomical determination was conducted by comparing the Les Cottés tooth to the large sample of Neanderthal anterior teeth already published by one of the authors of this manuscript (ALC, 4).

The tooth was scanned by Heiko Temming on the 11<sup>th</sup> of June 2012, using the Skyscan 1712 micro-CT scanner in the Department of Human Evolution at MPI-EVA (Leipzig, Germany). The scanning parameters used are as follows: 100 kV, 100  $\mu$ A, Al+Cu filters, 2 frame averaging, over 360 degrees, 885 ms of exposure time, data were saved as 16-bit tiff files. The  $\mu$ CT data were reconstructed with a pixel size of 13.04  $\mu$ m, using N-Recon. The stack was downsampled to 27  $\mu$ m using the Triangle filter (2,2,2) in Avizo 6.3, and then further filtered using a combination of median and Kuwahara filters with a kernel size of 3 (see (4) for details). The dentine and the pulp cavity were segmented in Avizo 6.3 to generate separate 3D models of those dental tissues. Linear, surfacic, and volumetric measurements were taken in Avizo 6.3.

The nine root metrics used in the analysis for the anatomical determination involved the labio-lingual and mesio-distal cervical root diameters, the root length, the root total volume, the root pulp volume, the root surface area, the cervical surface area, and the labial and the lingual root surface areas. The values for Les Cottés tooth and the comparative Neanderthal samples are reported in the following table.

Posterior probabilities based on several root metrics (following 4) classify this tooth as a **permanent maxillary lateral incisor** (after abbreviated as: I<sup>2</sup>) with a 9.4 % chance of misclassification (Table S2).



	Root metrics								
	RMD	RLL	CA	RL	RV	RPV	RSA	RSA-Ling	RSA-Lab
<b>Les Cottés</b>	<b>5.7</b>	<b>8.0</b>	<b>43.1</b>	<b>15.9</b>	<b>415.8</b>	<b>7.3*</b>	<b>305.9</b>	<b>130.7</b>	<b>175.8</b>
<b>Neandertals**</b>									
LI1 (N=18)	4.4 ± 0.4 [3.9-5.7]	7.2 ± 0.4 [6.3-8.0]	26.5 ± 5.0 [20.6-43.1]	17.1 ± 1.9 [13.8-20.9]	335.1 ± 63.6 [200.1-421.7]	12.5 ± 6.7 [5.2-24.9]	300.8 ± 42.3 [195.6-358.9]	147.0 ± 31.3 [70.8-196.3]	153.3 ± 27.1 [92.4-191.2]
LI2 (N=16)	4.6 ± 0.6 [3.3-6.4]	7.5 ± 0.6 [6.1-8.7]	28.4 ± 3.9 [18.0-35.7]	18.0 ± 2.3 [13.4-21.6]	381.4 ± 86.4 [208.9-507.6]	20.7 ± 14.3 [6.6-63.0]	324.4 ± 86.4 [208.9-507.6]	143.0 ± 31.1 [70.3-192.6]	180.3 ± 36.8 [98.5-228.2]
LC (N=16)	5.8 ± 0.5 [4.3-6.4]	8.9 ± 0.7 [7.5-10.5]	42.3 ± 94.5 [25.6-54.5]	20.7 ± 2.9 [16.1-25.6]	595.4 ± 150.5 [339.5-902.3]	31.9 ± 18.8 [11.6-70.8]	410.2 ± 94.5 [221.8-619.7]	190.9 ± 45.9 [78.2-280.3]	236.5 ± 49.1 [150.1-340.8]
UI1 (N=17)	7.0 ± 1.1 [5.4-9.3]	7.6 ± 0.5 [7.0-8.6]	41.9 ± 8.0 [33.2-59.1]	17.2 ± 2.4 [10.5-19.8]	452.0 ± 123.5 [195.8-693.6]	37.2 ± 20.7 [2.0-86.0]	315.3 ± 60.7 [162.9-409.3]	134.8 ± 26.6 [68.5-175.7]	179.6 ± 39.5 [94.5-243.8]
<b>UI2 (N=18)</b>	<b>6.1 ± 0.5</b> <b>[5.4-7.2]</b>	<b>8.0 ± 0.6</b> <b>[7.3-9.3]</b>	<b>39.6 ± 6.1</b> <b>[31.2-53.3]</b>	<b>17.6 ± 1.8</b> <b>[14.7-20.0]</b>	<b>487.2 ± 91.0</b> <b>[310.7-633.2]</b>	<b>31.4 ± 18.1</b> <b>[9.3-81.7]</b>	<b>343.1 ± 42.9</b> <b>[252.8-409.2]</b>	<b>138.4 ± 20.8</b> <b>[97.8-183.0]</b>	<b>204.8 ± 32.3</b> <b>[155.3-269.0]</b>
UC (N=12)	6.2 ± 0.2 [5.8-6.5]	9.3 ± 0.7 [8.7-11.0]	46.2 ± 5.0 [40.0-55.8]	22.5 ± 2.5 [17.7-25.2]	650.6 ± 98.9 [406.9-817.3]	36.7 ± 14.8 [16.4-66.3]	454.3 ± 60.3 [311.2-539.4]	198.3 ± 44.6 [122.2-311.6]	255.4 ± 46.0 [170.4-314.4]

\*The very small value of the Les Cottés root pulp volume can be explained by the older age of the individual (evidenced by the filling of the pulp cavity with secondary dentine that occurs with age).

\*\* : Comparative data from (4). “L” stands for “lower”, “U” for “upper”, “I” for incisor, “C” for canine, “1” for “central” and “2” for lateral, so that the UI2 is the upper lateral incisor. Mean ± standard deviation [minimum-maximum]. RMD: mesio-distal root cervical diameter [mm]; RLL: labio-lingual root cervical diameter [mm]; CA: cervical area [mm<sup>2</sup>]; RL: root length [mm]; RV: total root volume [mm<sup>3</sup>]; RPV: root pulp volume [mm<sup>3</sup>]; RSA: root surface area [mm<sup>2</sup>]; RSA-Ling: lingual root surface area [mm<sup>2</sup>]; RSA-Lab: labial root surface area [mm<sup>2</sup>].

**Table S2: Root metrics of the Les Cottés tooth (Z4-1514) and comparative Neandertal permanent anterior teeth.**

## 2. Estimation for root formation time of the Les Cottés Neandertal tooth (Z4-1514).

This permanent maxillary lateral incisor ( $I^2$ ) belongs to an older individual as evidenced by both the heavy degree of wear that reaches the root in its cervical part (the crown is totally worn off; score 8 after 5) and the significant thickness of hypercementosis covering the root dentine. Additionally, some tertiary dentine is visible at the incisal surface and obliterates the exposed pulp cavity, the diameter of which is reduced by the significant apposition of secondary dentine.

This tooth cannot be used to estimate the individual's age at death as it is far too worn and there is no associated permanent first molar, which would give a time of birth (if unworn). Furthermore, neither physical section nor synchrotron imaging were performed on this tooth.

We are left with estimating the time of its root formation using references from other Neanderthals for which an age at death was estimated histologically. This allows us to provide a range of ages for root initiation and completion, as dentine is sampled for collagen extraction.

The  $I^2$  crown from juvenile Neanderthal Engis 2 (6) was nearly complete when the individual died at 3 years of age. The Gibraltar 2 Neandertal was slightly older at death (4.6 years), and its  $I^2$  had already started the development of its root as a spicule (less than 1 mm on the labial side)(6). The dental development of the Scladina child (Scla I-4A) has been extensively investigated (7). Its  $I^2$  root started growing at the age of 3.8 years (7). These comparative observations suggest that the root of the Les Cottés  $I^2$  had started growing at around 4 years of age, which either corresponds to the actual end of the weaning process or shortly after weaning has been completed.

The Scladina child had completed the root of its  $I^2$  by the time of its death at 8 years of age (7). The Spanish juvenile El Sidrón J1 was estimated to have died at 7.7 years (8). The root of its lateral incisor is described as developed at three-quarters of its estimated final length (score F for modern human boys and girls following 8). The primary dentine of Les Cottés  $I^2$  root has likely achieved its growth around ~8 years of age. Overall, this is consistent with the root formation time calculated for Scladina, the  $I^2$  root would have taken 4 years to achieve its development.

In summary, one can hypothesize that **the dentine of the Les Cottés  $I^2$  root formed between 4 and 8 years of age.**

Yearly annulations of cementum were then deposited onto the surface of the root dentine, after ~8 years of age until death, i.e., during adolescence and adulthood. To avoid surface contaminations, and as is routinely done during isotopic analyses, the cementum was removed from the tooth surface to sample only (primary, secondary, and tertiary) dentine. It is of note that the whole root dentine (from cervical area to root tip) was used in the sampling procedure. The amount of tertiary dentine that may have been sampled is negligible as it is restricted to a very small volume of dental tissue formed at the roof of the pulp chamber on the incisal surface. The amount of secondary dentine filling in the pulp cavity is more important, although our sample is dominated by primary dentine which constitutes the core of the tooth root.

Overall, **the root dentine sampled here represents the post-weaning stages (see section 3) of life of the Les Cottés individual.**

39

### 40 3. Estimation of the weaning age

41 The age of weaning in Neandertals remains intensely debated, and results vary according to the  
42 techniques of investigation used and the lines of evidence retrieved.

43 Skinner (9) argued that attrition starts at 3 years of age in Middle Pleistocene infants, citing the  
44 initiation of the weaning process by the introduction of solid food into the diet. The pattern of microwear  
45 of the deciduous molars show that Gibraltar 2 (dead at 4.6 years) had a diet close to that of Fuegians and  
46 Eskimos, i.e., highly carnivorous (10). Similarly, the Neandertal juvenile of the Trou de l'Abîme Cave at  
47 Couvin (Belgium) was estimated to have died at 5 or 6 years of age (11). The microwear signature on its  
48 lower deciduous second molar suggests that the child was consuming mainly tough food, presumably  
49 mostly meat (9). These two examples suggest that by the age of their death, both Gibraltar 2 and Couvin  
50 were weaned long enough to record the microwear signature of their diet on their tooth enamel surfaces.

51 Linear enamel hypoplasias (LEH) are non-specific stress markers occurring in enamel during  
52 tooth formation. They occur often in Neanderthal teeth, more frequently in the permanent dentition (about  
53 41%) than in the deciduous teeth (~4%) (12). This argues in favor of a strong period of stress around 4  
54 years of age, potentially related with nutritional pressure associated with the weaning process (12).

55 More recently, using the mapping of barium distribution in dental hard tissues, Austin et al. (13)  
56 suggested that breast-feeding had abruptly stopped at 1.2 years of age for the Scladina child. Smith et al.  
57 (14) further proposed an age at weaning of 2.5 years for the Payre 6 French Neandertal juvenile.

58 **Provided that the primary dentine of the Les Cottés I<sup>2</sup> root formed between 4 and 8 years of age,**  
59 **the tooth sampling may involve either the very latest stages of the weaning process or concern solely**  
60 **the post-weaning stages when the child is feeding autonomously.**

61

62


63

## 64 Supporting Information 4

### 65 Additional discussion

#### 66 Quality of the data

67 A recent critique of the Naito et al. (16) article was published by O'Connell and Collins (15), revealing  
68 the importance of verifying the correlation between the  $\delta^{15}\text{N}$  values of proline and hydroxyproline to  
69 ensure the quality of the data. Our first batches of samples were prepared by a derivatization method  
70 (MOC) (17) which does not allow for the analysis of nitrogen isotope ratios of hydroxyproline. However,  
71 our last batch was prepared following the protocol suggested by O'Connell and Collins (15). The data  
72 from this protocol is presented in Figure S2 and shows an excellent correlation between  $\delta^{15}\text{N}$  values of  
73 proline and hydroxyproline. Yarnes and Herszage (17) demonstrated that the two protocols used in this  
74 study produce the same values. We monitored it using an external standard, which is also used for C  
75 isotope ratio analyses of the amino acids. The results of measurements of this standard are given in Table  
76 S3.



77

78 **Figure S2: Correlation between  $\delta^{15}\text{N}$  values of proline and hydroxyproline of the samples of this**  
79 **study**

	$\delta^{15}\text{N}_{\text{Glu}}$	SD	$\delta^{15}\text{N}_{\text{Phe}}$	SD	$\delta^{13}\text{C}_{\text{Phe}}$	SD	$\delta^{13}\text{C}_{\text{Val}}$	SD
Bovine								
Liver 1577b	10.3	0.4	8.2	0.2	-25.1	0.5	-25.2	0.3


80

81 **Table S3: Results for the C and N isotope ratios of amino acids (delta values in ‰) discussed in the**  
82 **main text for the external standard, the reference material Bovine Liver 1577b.**




83

84 **Additional discussion on the isotope compositions of the amino acids.**



85

86 **Figure S3:  $\delta^{15}\text{N}$  of phenylalanine as a function of bulk values. Circles: Les Cottés samples, crosses:**  
87 **Grotte du Renne samples. Herbivores are in green, carnivores in red, and omnivores in yellow.**




88

89 **Figure S4:  $\delta^{15}\text{N}$  of glutamic acid as a function of bulk values. Circles: Les Cottés samples, Crosses:**  
90 **Grotte du Renne samples. Herbivores are in green, carnivores in red, and omnivores in yellow.**

91 Nitrogen isotope ratios are impacted by the trophic level effect for the following amino acids: Thr  
92 (source/metabolic), Val (trophic), Hyp, Leu (trophic), Ala (trophic), and Asp to a lesser extent.


93 The trophic spacing was previously observed by Fuller and Petzke (18) for Thr and Ala but not for Leu  
94 and Ala.

95 The threonine  $\delta^{15}\text{N}$  values correlate with the phenylalanine values within each dietary group - with the  
 96 exception of the animals with  $\delta^{15}\text{N}_{\text{Phe}}$  above 12‰. This suggests that threonine  $\delta^{15}\text{N}$  values are both  
 97 impacted by local environment and trophic level. This also confirms that the Neandertal from Grotte du  
 98 Renne was one trophic level above the carnivores, and therefore probably still breastfeeding at the time of  
 99 its death. The Neandertal from Les Cottés is also different from the other carnivores, but its  $\delta^{15}\text{N}_{\text{Phe}}$  is  
 100 above 12‰, so it is difficult to determine the specific cause of this variation.




101  
 102 **Figure S5:  $\delta^{15}\text{N}_{\text{Phe}}$  as a function of  $\delta^{15}\text{N}_{\text{Thr}}$  values. Circles: Les Cottés samples, crosses: Grotte du**  
 103 **Renne samples. Herbivores are in green, carnivores in red, omnivores in yellow, Neandertals are in**  
 104 **black.**

105 Concerning the carbon isotope values of the amino acids, the details of the correlation are given in Table  
 106 S2. The values seem to be impacted by trophic level for Val, Leu and Gly (Figure S6), with almost no  
 107 overlap between carnivores and herbivores (Figure S7). With each dietary group, the values of  $\delta^{13}\text{C}_{\text{Val}}$   
 108 correlate to that of Asp (Figure S6), suggesting that this other amino acid could reveal some metabolic or  
 109 local environmental information and would explain the rest of the variability of the  $\delta^{13}\text{C}_{\text{Val}}$  values. This  
 110 would suggest that the Les Cottés Neandertal was hunting animals with elevated  $\delta^{13}\text{C}_{\text{Asp}}$  values, such as  
 111 horses.



112

113 **Figure S6:  $\delta^{13}\text{C}_{\text{Leu}}$  as a function of  $\delta^{13}\text{C}_{\text{Val}}$ . Circles: Les Cottés samples, crosses: Grotte du Renne**  
 114 **samples. Herbivores are in green, carnivores in red, omnivores in yellow, and the Neandertal is in**  
 115 **black.**



116

117 **Figure S7:  $\delta^{13}\text{C}_{\text{Val}}$  as a function of  $\delta^{13}\text{C}_{\text{Asp}}$ . Circles: Les Cottés samples, crosses: Grotte du Renne**  
 118 **samples. Herbivores are in green, carnivores in red, omnivores in yellow, the Neandertal is in black.**

	<b>Asp</b>	<b>Glu</b>	<b>Gly</b>	<b>Ile</b>	<b>Leu</b>	<b>Phe</b>	<b>Pro</b>	<b>Val</b>
<b>Ala</b>	NC <sup>1</sup>	NC	NC <sup>2</sup>	NC	NC	NC	NC	NC <sup>3</sup>
<b>Asp</b>		NC	NC <sup>2</sup>	NC <sup>6</sup>	NC <sup>3</sup>	SC <sup>5</sup>	SC <sup>5</sup>	NC <sup>3,4</sup>
<b>Glu</b>			NC <sup>2</sup>	NC <sup>6</sup>	NC <sup>3</sup>	SC <sup>5</sup>	SC <sup>5</sup>	NC <sup>3</sup>
<b>Gly</b>				NC <sup>2,6</sup>	C <sup>2</sup>	NC <sup>2</sup>	NC <sup>2</sup>	C <sup>3</sup>
<b>Ile</b>					NC	NC <sup>6</sup>	NC <sup>6</sup>	NC <sup>3</sup>
<b>Leu</b>						NC <sup>3</sup>	C <sup>3</sup>	C <sup>3</sup>
<b>Phe</b>							SC	NC <sup>3</sup>
<b>Pro</b>								NC <sup>3</sup>
<b>Val</b>								

120 **Table S4: Correlation between  $\delta^{13}\text{C}$  values of amino acids. NC: no correlation, C correlation, SC:**  
 121 **slight correlation**

- 1 The Les Cottés Neandertal is enriched in  $^{13}\text{C}$  for Asp compared to other animals.
- 2 Carnivores from Les Cottés are enriched in  $^{13}\text{C}$  for Gly but not those of Grotte du Renne.
- 3 Carnivores are enriched in  $^{13}\text{C}$  for Val and/or Leu, animals can be grouped depending on their diet regardless of site
- 4 Correlation within dietary group.
- 5 No difference between herbivores and carnivores.
- 6 Herbivores tend to be depleted in  $^{13}\text{C}$  for Ile.

122

123

124

125

126

127

128



129 **Additional data tables S5 to S10 (separate file)**

130 **Table S5: General information on the samples analyzed in this study: layer attribution, dating, sample type, taxonomic identification.**, 1. Talamo et al. (19), 2. Hajdinjak et al (20), 3. Welker et al. (21) ,4.Welker et al. (22)

**Table S6. Summary of isotope data for C and N obtained on bulk collagen ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ), and some amino acids: glutamic acid ( $\delta^{15}\text{N}_{\text{Glu}}$ ), phenylalanine ( $\delta^{13}\text{C}_{\text{Phe}}$ ,  $\delta^{15}\text{N}_{\text{Phe}}$ ) and valine ( $\delta^{13}\text{C}_{\text{Val}}$ ).**

TP= trophic position. Bulk collagen data are from this study, from 1. Talamo et al. (19) and 2. Welker et al. (22)

**Table S7. Bulk collagen data from this study: Collagen yield (%), C/N ratios and isotope data.**

**Table S8. Carbon isotope data on amino acids. Average  $\delta^{13}\text{C}$  values (top part) and standard deviation on the duplicate measurement.**

**Table S9. Nitrogen isotope data on amino acids**

**Table S10. Carbon and nitrogen isotope data on the standard bovine liver 1577b**

131

132

133 **Cited literature:**

- 134 1. Soressi M, et al. (2010) *Les Cottés (Vienne). Nouveaux travaux sur l'un des gisements de référence*  
135 *pour la transition Paléolithique moyen/supérieur* (Association des Publications Chauvinoises).
- 136 2. Talamo S, Richards M (2011) A comparison of bone pretreatment methods for AMS dating of  
137 samples 30,000 BP. *Radiocarbon* 53(03):443–449.
- 138 3. Kromer B, Lindauer S, Synal H-A, Wacker L (2013) MAMS—a new AMS facility at the Curt-  
139 Engelhorn-Centre for Archaeometry, Mannheim, Germany. *Nucl Instrum Methods Phys Res Sect B*  
140 *Beam Interact Mater At* 294:11–13.
- 141 4. Le Cabec A, Gunz P, Kupczik K, Braga J, Hublin J-J (2013) Anterior tooth root morphology and size  
142 in Neandertals: Taxonomic and functional implications. *J Hum Evol* 64(3):169–193.
- 143 5. Smith BH (1984) Patterns of molar wear in hunter-gatherers and agriculturalists. *Am J Phys*  
144 *Anthropol* 63:39–56.
- 145 6. Smith TM, et al. (2010) Dental evidence for ontogenetic differences between modern humans and  
146 Neandertals. *Proc Natl Acad Sci* 107:20923–20928.
- 147 7. Smith TM, Toussaint M, Reid DJ, Olejniczak AJ, Hublin J-J (2007) Rapid dental development in a  
148 Middle Paleolithic Belgian Neandertal. *Proc Natl Acad Sci* 104:20220–20225.
- 149 8. Rosas A, et al. (2017) The growth pattern of Neandertals, reconstructed from a juvenile skeleton from  
150 El Sidrón (Spain). *Science* 357(6357):1282–1287.
- 151 9. Skinner M (1997) Dental Wear in Immature Late Pleistocene European Hominines. *J Archaeol Sci*  
152 24:677–700.
- 153 10. Fox CL, Pérez-Pérez A (1993) The diet of the Neandertal Child Gibraltar 2 (Devil's Tower) through  
154 the study of the vestibular striation pattern. *J Hum Evol* 24:29–41.
- 155 11. Toussaint M, et al. (2010) The Neandertal lower right deciduous second molar from Trou de l'Abîme  
156 at Couvin, Belgium. *J Hum Evol* 58:56–67.
- 157 12. Ogilvie MD, Curran BK, Trinkaus E (1989) Incidence and patterning of dental enamel hypoplasia  
158 among the Neandertals. *Am J Phys Anthropol* 79(1):25–41.
- 159 13. Austin C, et al. (2013) Barium distributions in teeth reveal early-life dietary transitions in primates.  
160 *Nature* 498(7453):216–219.
- 161 14. Smith TM, et al. (2018) Wintertime stress, nursing, and lead exposure in Neandertal children. *Sci Adv*  
162 4(10). doi:10.1126/sciadv.aau9483.
- 163 15. O'Connell TC, Collins MJ (2017) Comment on “Ecological niche of Neandertals from Spy Cave  
164 revealed by nitrogen isotopes of individual amino acids in collagen” [J. Hum. Evol. 93 (2016) 82–  
165 90]. *J Hum Evol*. doi:10.1016/j.jhevol.2017.05.006.
- 166 16. Naito YI, et al. (2016) Ecological niche of Neandertals from Spy Cave revealed by nitrogen isotopes  
167 of individual amino acids in collagen. *J Hum Evol* 93:82–90.

- 168 17. Yarnes CT, Herszage J (2017) The relative influence of derivatization and normalization procedures  
169 on the compound-specific stable isotope analysis of nitrogen in amino acids. *Rapid Commun Mass*  
170 *Spectrom* 31(8):693–704.
- 171 18. Fuller BT, Petzke KJ (2017) The dietary protein paradox and threonine  $^{15}\text{N}$ -depletion: Pyridoxal-5'-  
172 phosphate enzyme activity as a mechanism for the  $\delta^{15}\text{N}$  trophic level effect. *Rapid Commun Mass*  
173 *Spectrom* 31(8):705–718.
- 174 19. Talamo S, Soressi M, Roussel M, Richards M, Hublin J-J (2012) A radiocarbon chronology for the  
175 complete Middle to Upper Palaeolithic transitional sequence of Les Cottés (France). *J Archaeol Sci*  
176 39(1):175–183.
- 177 20. Hajdinjak M, et al. (2018) Reconstructing the genetic history of late Neandertals. *Nature*  
178 555(7698):652.
- 179 21. Welker F, Soressi M, Rendu W, Hublin J-J, Collins M (2015) Using ZooMS to identify fragmentary  
180 bone from the late Middle/Early Upper Palaeolithic sequence of Les Cottés, France. *J Archaeol Sci*  
181 54:279–286.
- 182 22. Welker F, et al. (2016) Palaeoproteomic evidence identifies archaic hominins associated with the  
183 Châtelperronian at the Grotte du Renne. *Proc Natl Acad Sci* 113(40):11162–11167.

184

185

186

187

188

Site	SEVA Id	Id	Layer	Dating	Reference (dating)	Sample	Growth period	Anatomical species determination	Zooms/ADN determination
Les Cottés	1526	Z4-1514	US04lower		this study	tooth, I2	post weaning	Neanderthal	2
Les Cottés	2184	X6-727	US06		-	decidual tooth, I3	in utero	hyena	-
Les Cottés	2185	Z3-688	US06		-	tooth, LP3	possible weaning	hyena	-
Les Cottés	2183	X7-738	US04lower		-	tooth, UP3	possible weaning	hyena	-
Les Cottés	13670	Y5-583	US04upper	-	-	tooth, M3	post weaning	hyena	3
Les Cottés	1645	Z4-3466	US04lower	43680-44320	1	bone		bison	bison this study
Les Cottés	1639	Z4-1462	US04lower	39970-40760	1	bone		equidae	equidae this study
Les Cottés	1641	Z4-1911	US04lower	40870-41360	1	bone		equidae	equidae this study
Les Cottés	1642	Z4-2245	US04lower	40010-40770	1	bone		equidae	equidae this study
Les Cottés	1647	Y4-573	US08	45120-45780	1	tooth, UP3	possibl pre weaning	bison	bison this study
Les Cottés	1648	Y4-425	US08	44200-44750	1	tooth, UPM1/M2	possibl pre weaning	bison	bison this study
Les Cottés	1651	Z3-170	US04lower	38600-38990	1	tooth, P3 or 4	possible weaning	Rangifer	
Les Cottés	1653	Z4-3398	US06	43570-44200	1	tooth, P4	post weaning	equidae	equidae this study
Les Cottés	1650	Y6-1960	US04upper	-	-	tooth, Upper M1/M3	post weaning	Rangifer	reindeer this study
Les Cottés	1652	Z3-781	US04lower	40820-41320	1	tooth, LM2	post weaning	reindeer	reindeer this study
Les Cottés	13665	S6-557	US04lower	38710-39360	1	bone		rangifer	-
Les Cottés	2182	R5-1887	US08		-	tooth, LM2	post weaning	bear	-
Les Cottés	2667	X5-215	US08		-	bone		large herbivore	-
Les Cottés	2669	Y5-1647	US08		-	tooth	?	rhinoceros	-
Les Cottés	2670	Y6-2223	US06		-	tooth	?	mammoth	-
Les Cottés	2671	Z2-76	US08		-	tooth	?	megaceros	-
Grotte du Renne	32908		Xb2					-	pantharinae 4
Grotte du Renne	32909		Xb2					-	pantharinae 4
Grotte du Renne	32912		Xb2					-	canidae 4
Grotte du Renne	32919		Xc					-	hyena 4
Grotte du Renne	32867		Xb					-	Bos/Bison 4
Grotte du Renne	32874		Xb					-	bison 4
Grotte du Renne	32904		Xb2				possible weaning	-	rangifer 4
Grotte du Renne	32906		Xb2					-	rangifer 4
Grotte du Renne	32907		Xb2					-	equidae 4
Grotte du Renne	32915		Xb					-	equidae 4
Grotte du Renne	32911							-	mammoth 4
Grotte du Renne	32913		Xb2					-	mammoth 4
Grotte du Renne	32916		Xc					-	rhino 4
Grotte du Renne	32886		X					-	hominin 4
Grotte du Renne	32917		Xc					-	Ursidae 4

Table S5: General information on the samples analysed in this study: layer attribution, dating, sample type, taxonomic identification. 1. Talamo et al. (19) 2 Hajdinjak et al (20) 3 Welker et al. (21) 4. Welker et al. (22)



Site	SEVA	Species determination	Diet	C/N	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	Ref	$\delta^{15}\text{N}_{\text{Phe}}$	SD	$\delta^{15}\text{N}_{\text{Glu}}$	SD	$\delta^{13}\text{C}_{\text{Phe}}$	SD	$\delta^{13}\text{C}_{\text{Val}}$	SD	TP
Les Cottés	1526	Neanderthal	?	3.0	-18.1	13.9	this study	13.2	0.3	19.5	0.1	-24.5	0.8	-23.2	0.3	2.9
Les Cottés	2184	Hyena	Carnivore	3.1	-19.7	11.9	this study	8.9	0.1	16.7	0.5	-28.3	0.2	-26.3	0.0	3.1
Les Cottés	2185	Hyena	Carnivore	3.1	-18.8	10.2	this study	7.1	0.1	15.4	0.2	-26.2	0.2	-23.1	0.2	3.2
Les Cottés	2183	Hyena	Carnivore	3.1	-18.5	11.7	this study	6.2	0.2	17.3	0.4	-26.8	0.1	-23.4	0.1	3.6
Les Cottés	13670	Hyena	Carnivore	3.2	-18.1	9.0	1	12.1	0.1	13.2	0.3	-26.0	0.8	-24.4	0.9	2.2
Les Cottés	1645	Bison	Herbivore	3.2	-20.3	6.7	this study	7.3	1.1	7.3	1.1	-24.4		-27.1	0.3	2.1
Les Cottés	1639	Equidae	Herbivore	3.2	-20.1	6.4	this study	9.1	0.6	9.2	0.5	-26.4		-27.6		2.1
Les Cottés	1641	Equidae	Herbivore	3.2	-20.5	4.8	this study	11.1	1.4	7.6	0.3	-24.8		-25.5		1.7
Les Cottés	1642	Equidae	Herbivore	3.2	-20.6	4.9	this study	11.1	0.6	7.6	0.8	-25.1	0.5	-25.8	0.7	1.6
Les Cottés	1647	Bison	Herbivore	3.3	-20.3	9.2	this study	11.1	0.4	13.1	0.5	-26.8	0.0	-27.6	0.4	2.4
Les Cottés	1648	Bison	Herbivore	3.2	-20.1	7.8	this study	8.7	0.4	12.0	0.9	-26.8		-27.3	0.2	2.5
Les Cottés	1651	Rangifer		3.2	-18.4	6.3	this study	7.9	0.0	8.3	0.5	-25.9	0.8	-27.0	0.6	2.2
Les Cottés	1653	Equidae	Herbivore	3.3	-20.1	7.5	this study	8.3	0.3	9.8	0.1	-27.5	0.1	-27.9	0.1	2.3
Les Cottés	1650	Rangifer	Herbivore	3.3	-18.6	9.1	this study	11.0	0.4	11.1	0.6	-26.6	0.8	-26.6	0.6	2.1
Les Cottés	1652	Reindeer	Herbivore	3.2	-20.6	5.4	this study	10.6	0.5	6.5	0.5	-27.5	0.1	-27.9	0.1	1.6
Les Cottés	13665	Rangifer	Herbivore	3.2	-19.2	8.5	1	12.4	0.3	11.2	0.1	-24.1	0.3	-26.7	0.6	1.9
Les Cottés	2667	large herbivore	Herbivore	3.1	-19.8	5.3	this study	-	-	-	-	-	-	-	-	-
Les Cottés	2669	rhinoceros	Herbivore	3.1	-19.7	8.3	this study	-	-	-	-	-	-	-	-	-
Les Cottés	2670	mammoth	Herbivore	3.1	-21.5	11.1	this study	-	-	-	-	-	-	-	-	-
Les Cottés	2671	megaceros	Herbivore	3.1	-20.1	5.9	this study	-	-	-	-	-	-	-	-	-
Les Cottés	2182	Bear	Omnivore	3.1	-21.6	4.3	this study	9.7	0.6	8.0	0.1	-26.8	1.1	-26.3	0.9	1.9
Grotte du Renne	32908	Pantherinae	Carnivore	3.2	-18.8	8.9	2	10.9	0.7	14.0	0.9	-25.8	0.1	-25.5	0.0	2.5
Grotte du Renne	32909	Pantherinae	Carnivore	3.2	-18	7.4	2	9.6	1.2	14.0	0.4	-24.8	0.2	-24.6	0.3	2.7
Grotte du Renne	32912	Canidae	Carnivore	3.2	-18.9	9.9	2	12.1	0.6	17.0	0.2	-25.9	0.1	-25.5	0.5	2.8
Grotte du Renne	32919	Hyena	Carnivore	3.2	-19.5	9.5	2	9.4	0.8	15.1	0.2	-25.7	0.1	-25.3	0.2	2.9
Grotte du Renne	32867	Bison	Herbivore	3.2	-20.3	5.9	2	9.4	0.8	8.4	0.9	-26.5	0.1	-27.3	0.1	2.0
Grotte du Renne	32874	Bison	Herbivore	3.2	-19	3.5	2	7.8	0.1	8.6	0.4	-25.5	0.1	-28.2	0.1	2.2
Grotte du Renne	32904	Rangifer	Herbivore	3.2	-19.3	7.8	2	13.7	0.2	12.8	0.2	-26.4	0.4	-27.2	0.6	2.0
Grotte du Renne	32906	Rangifer	Herbivore	3.3	-19.4	5.5	2	12.2	0.7	10.3	1.4	-26.4	0.1	-28.0	0.5	1.9
Grotte du Renne	32907	Equidae	Herbivore	3.2	-20.9	5.2	2	14.4	0.3	10.8	0.3	-27.1	0.1	-28.1	0.5	1.6
Grotte du Renne	32915	Equidae	Herbivore	3.2	-20.8	5.2	2	12.3	0.2	10.2	0.7	-26.1	0.1	-27.9	0.1	1.8
Grotte du Renne	32911	Mammoth	Herbivore	3.3	-21.5	6.8	2	12.0	1.4	13.2	0.5	-26.6	0.4	-28.9	0.1	2.3
Grotte du Renne	32913	Mammoth	Herbivore	3.2	-20.2	4.9	2	9.9	0.2	10.1	0.1	-26.2	0.3	-28.9	0.4	2.1
Grotte du Renne	32916	Rhino	Herbivore	3.2	-18.9	2.4	2	7.6	0.6	7.5	1.3	-25.6	0.1	-28.3	0.4	2.1
Grotte du Renne	32886	Neanderthal	?	3.2	-19.5	14.3	2	9.4	0.6	18.1	0.1					3.2
Grotte du Renne	32917	Ursidae	Omnivore	3.3	-21.1	3.2	2	11.3	0.7	9.2	0.7	-26.7	0.1	-27.7	0.5	1.8

**Table S6. Summary of isotope data for C and N obtained on bulk collagen ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ), and some amino acids : glutamic acid ( $\delta^{15}\text{N}_{\text{Glu}}$ ) Phenylalanine ( $\delta^{13}\text{C}_{\text{Phe}}$ ,  $\delta^{15}\text{N}_{\text{Phe}}$ ) and valine ( $\delta^{13}\text{C}_{\text{Val}}$ ).**

TP= trophic position. Bulk collagen data are from this study, from 1. Talamo et al. (19) and 2. Welker et al. (22)

Site	Species determination	Diet	% Collagen	C/N	%C	%N	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)
Les Cottés	Neanderthal	?	5.6	3.2	44.0	15.9	-18.1	13.9
Les Cottés	Hyena	Carnivore	3.4	3.1	44.1	16.4	-19.7	11.9
Les Cottés	Hyena	Carnivore	7.0	3.1	44.8	16.8	-18.8	10.2
Les Cottés	Hyena	Carnivore	6.0	3.1	44.2	16.4	-18.5	11.7
Les Cottés	Large herbivore	Herbivore	5.7	3.1	44.3	16.6	-19.8	5.3
Les Cottés	Rhinoceros	Herbivore	2.9	3.1	44.0	16.6	-19.7	8.3
Les Cottés	Mammoth	Herbivore	1.2	3.1	41.5	15.7	-21.5	11.1
Les Cottés	Megaceros	Herbivore	4.5	3.1	45.2	17.1	-20.1	5.9
Les Cottés	Bear	Omnivore	1.5	3.1	38.4	14.4	-21.6	4.3

Table S7. Bulk collagen data from this study: Collagen yield (%), C/N ratios and isotope data.

Table S8. Carbon isotope data on amino acids. Average  $\delta^{13}\text{C}$  values (top part) and standard deviation on the duplicate measurement.

	Ala	Asp	Glu	Gly	Ile	Leu	Lys	Met	Phe	Pro	Thr	Val
1639	-23.47	-17.01	-19.65	-16.90	-21.89	-32.55			-26.23	-19.77		-27.80
1641	-20.26	-15.40	-19.04	-12.57	-21.61	-32.01			-24.56	-18.78		-26.05
1642	-20.49	-12.74	-20.67	-10.26	-21.62	-32.01			-25.06	-19.30		-25.75
1645	-23.38	-13.88	-16.50	-14.01	-25.72	-30.85			-23.98	-18.16		-27.07
1647	-23.38	-15.33	-20.22	-13.39	-26.28	-30.37			-26.81	-18.09		-27.29
1648	-24.32	-16.56	-20.86	-15.08	-25.63	-31.14			-26.82	-18.84		-27.63
1650	-22.83	-16.27	-19.57	-11.93	-25.47	-30.57			-26.58	-18.45		-26.60
1651	-23.08	-15.87	-19.66	-12.91	-24.93	-30.96			-25.87	-18.00		-26.99
1652	-24.61	-15.06	-21.78	-14.12	-21.71	-32.84			-25.67	-19.89		-27.05
1653	-25.86	-17.03	-20.85	-16.93	-26.07	-31.17			-27.54	-19.47		-27.94
1526	-22.94	-10.96	-22.06	-9.89	-21.45	-29.51			-24.52	-17.71		-23.23
13665	-23.85	-14.17	-19.59	-13.09	-24.48	-30.57			-24.15	-17.60		-26.67
13670	-21.46	-18.06	-19.45	-10.50	-21.23	-31.00			-26.02	-18.18		-24.39
32867	-24.98	-16.64	-19.89	-17.56	-24.80	-31.30			-26.47	-19.57		-27.34
32874	-24.59	-17.67	-18.05	-13.14	-26.40	-32.01			-25.49	-19.36		-28.23
32904	-21.76	-15.14	-18.16	-11.32	-25.57	-31.17			-26.43	-19.02		-27.19
32906	-24.56	-16.86	-20.26	-12.79	-26.40	-32.56			-26.45	-20.07		-28.02
32907	-24.17	-17.68	-21.25	-15.25	-23.00	-33.50			-27.07	-20.40		-28.10
32908	-23.73	-16.32	-21.42	-11.40	-21.96	-31.22			-25.77	-19.46		-25.46
32909	-23.14	-16.49	-20.20	-10.98	-22.67	-30.01			-24.80	-18.53		-24.58
32911	-25.71	-18.75	-22.90	-17.32	-24.14	-33.98			-26.56	-21.06		-28.93
32912	-22.24	-16.57	-21.07	-14.13	-22.14	-31.37			-25.91	-19.73		-25.52
32913	-24.61	-18.07	-18.16	-16.53	-24.04	-33.80			-26.24	-20.57		-28.85
32915	-24.85	-17.17	-20.67	-17.36	-22.07	-32.82			-26.06	-20.40		-27.93
32916	-24.29	-15.88	-18.32	-14.35	-25.49	-31.40			-25.61	-18.91		-28.25
32917	-24.32	-19.73	-23.56	-15.58	-23.12	-32.83			-26.69	-22.48		-27.70
32919	-25.59	-19.04	-20.54	-16.82	-22.08	-30.73			-25.71	-19.51		-25.25
2182	-25.74	-21.10	-23.86	-11.38	-21.19	-31.93	-21.67	-26.20	-26.76	-22.10	-13.94	-26.28
2183	-25.49	-17.72	-22.00	-9.28	-21.60	-29.64	-21.54	-25.34	-26.76	-18.70	-11.88	-23.45
2184	-25.72	-18.97	-23.12	-8.46	-24.13	-30.93	-23.79	-25.94	-28.32	-20.63	-15.41	-26.34
2185	-20.18	-16.16	-23.03	-5.53	-21.50	-29.59	-20.03	-23.26	-26.25	-18.10	-10.14	-23.11

	Ala	Asp	Glu	Gly	Ile	Leu	Lys	Met	Phe	Pro	Thr	Val
1639	0.06	0.32	0.49	0.30	0.04	0.24			0.18	0.09		0.24
1641	0.81	0.59	0.74	0.48	0.96	0.19			0.27	0.47		0.82
1642	0.80	0.63	0.63	1.96	0.55	0.29			0.51	0.74		0.67
1645	0.21	1.15	0.01	0.13	0.10	0.15			0.55	0.18		0.28
1647	1.05	0.81	0.75	0.98	0.42	0.56			0.52	0.53		0.21
1648	0.16	0.12	0.26	0.39	0.56	0.36			0.04	0.12		0.37
1650	0.01	1.15	0.85	0.24	0.27	0.61			0.81	0.04		0.63
1651	0.00	1.19	0.54	0.05	0.13	0.35			0.80	0.13		0.58
1652	0.41	0.79	0.94	0.26	0.29	0.34			0.78	0.20		0.15
1653	0.13	0.20	0.07	0.35	0.13	0.11			0.10	0.02		0.15
1526	0.12	1.22	0.19	0.41	0.07	0.40			0.77	0.16		0.30
13665	0.12	0.85	0.00	0.48	0.50	0.42			0.30	0.29		0.59
13670	0.42	1.30	1.00	1.19	0.76	0.43			0.81	0.56		0.86
32867	0.09	0.25	0.05	0.63	0.10	0.40			0.11	0.15		0.10
32874	0.43	0.41	0.03	1.58	0.12	0.02			0.06	0.06		0.08
32904	0.10	0.09	0.23	0.25	0.06	0.20			0.40	0.08		0.56
32906	0.01	0.25	0.40	0.79	0.02	0.18			0.08	0.31		0.51
32907	0.13	0.13	0.53	0.34	0.24	0.21			0.09	0.30		0.47
32908	0.17	0.42	0.42	0.32	0.04	0.13			0.11	0.18		0.01
32909	1.00	0.19	0.99	0.70	0.52	0.22			0.21	0.27		0.33
32911	0.10	0.39	0.13	0.57	0.68	0.64			0.36	0.03		0.05
32912	0.31	0.20	0.24	0.48	0.35	0.48			0.15	0.07		0.49
32913	0.41	0.60	0.24	0.55	0.06	0.06			0.30	0.38		0.38
32915	0.47	0.30	0.18	0.28	0.69	0.02			0.13	0.22		0.05
32916	0.01	1.58	0.38	0.99	0.78	0.36			0.12	0.13		0.41
32917	0.61	0.90	0.18	0.27	0.14	0.06			0.10	0.31		0.51
32919	0.69	0.29	0.17	0.17	0.70	0.90			0.08	0.90		0.18
2182	0.15	0.23	0.30	0.11	1.19	1.24	0.27	1.35	1.12	0.54	0.67	0.93
2183	0.12	0.07	0.07	0.01	0.12	0.14	0.19	0.39	0.05	0.07	0.22	0.08
2184	0.22	0.55	0.08	0.58	0.18	0.03	0.80	0.23	0.19	0.29	0.13	0.05
2185	0.31	0.53	0.26	0.35	0.24	0.37	0.30	0.02	0.18	0.42	0.67	0.24

Table S9. Nitrogen isotope data on amino acids

	Ala	Asp	Glu	Gly	Hyp	Leu	Phe	Pro	Ser	Thr	Val	
std-c	10.91	8.98	9.85	12.20			2.91	8.11	13.15	9.37	-6.36	11.10
1526	15.50	18.38	19.52	13.97	18.55	17.60	13.19	18.97	11.23	-15.08	20.32	
32874	4.02	8.85	8.64	1.13	7.62	7.58	7.84	7.57	0.58	-8.32	10.04	
32907	6.68	11.32	10.76	3.42	11.44	6.40	14.41	11.99	3.30	-6.08	12.12	
32912	14.72	16.11	17.04	9.17	15.04	16.41	12.08	15.01	9.60	-10.20	18.22	
32913	7.07	10.20	10.14	2.11	10.52	8.04	9.91	10.45	2.86	-2.97	11.60	
32915	5.74	9.76	10.24	1.29	10.27	6.56	12.28	10.07	4.31	-5.45	11.71	
32919	11.14	14.24	15.13	8.11	13.77	11.20	9.39	13.86	8.36	-16.79	16.41	
2182	9.80	8.20	7.95	-1.39	9.11	6.97	9.73	8.56	-3.00	-13.14	10.55	
2183	18.44	16.60	17.32	9.10	16.63	13.93	6.20	16.33	7.99	-17.47	16.62	
2184	18.00	16.04	16.73	9.15	16.87	16.21	8.91	15.85	7.23	-12.79	19.13	
2185	17.17	13.65	15.44	10.19	14.00	13.73	7.12	14.14	8.06	-15.48	17.91	
32886	17.01	15.90	18.09	14.35	19.66	16.51	9.44	18.96	12.31	-21.36	19.18	

	Ala	Asp	Glu	Gly	Hyp	Leu	Phe	Pro	Ser	Thr	Val	
std-c	0.21	0.65	0.36	0.28			0.08	0.04	0.14	0.81	0.63	0.66
1526	0.09	0.19	0.09	0.15	0.03	1.84	0.27	0.09	1.05	0.20	1.18	
32874	0.03	0.05	0.43	0.15	0.19	1.12	0.07	0.23	0.11	1.12	0.42	
32907	0.64	0.74	0.33	0.01	0.25	0.24	0.26	0.35	0.45	0.65	0.18	
32912	0.44	0.30	0.22	0.37	0.22	0.17	0.56	0.07	0.34	0.14	0.46	
32913	0.12	0.06	0.07	0.09	0.12	0.96	0.22	0.12	0.41	0.08	0.49	
32915	0.67	0.24	0.71	0.98	0.56	0.27	0.25	0.61	0.48	0.72	0.48	
32919	0.12	1.24	0.20	0.13	0.30	1.16	0.76	0.18	0.19	0.63	1.38	
2182	0.33	0.31	0.07	0.58	1.40	0.12	0.62	0.00	0.46	0.64	0.98	
2183	0.99	0.52	0.41	0.31	0.51	0.12	0.19	0.01	0.09	1.20	0.23	
2184	0.86	0.82	0.55	1.30	0.81	0.75	0.05	0.16	0.63	0.47	0.53	
2185	0.92	0.07	0.16	1.32	0.31	1.22	0.10	0.17	0.95	1.60	0.20	
32886	0.65	0.34	0.09	0.19	0.12	0.00	0.62	0.22	0.49	0.30	0.58	

	Ala	Asp	Glu	Gly	Ile	Leu	Lys	Phe	Pro	Thr	Val
1577b-a	9.71	10.61	10.60	9.10	7.68	5.37	3.53	8.36	13.66	-6.19	10.04
1639	6.88	11.13	9.23	2.07	8.29	6.25	2.55	9.07	8.60	-6.89	8.76
1641	4.90	10.61	7.64	0.24	7.19	5.24	1.06	11.05	7.52	-7.80	7.59
1642	5.05	11.04	7.56	-0.02	7.51	5.49	3.29	11.05	7.73	-6.31	7.79
1645	4.83	8.34	7.27	1.39	8.44	5.82	2.30	7.26	6.25	-9.56	8.24
1647	8.43	12.92	11.97	4.83	11.66	9.60	8.95	8.66	9.46	-5.65	12.03
1648	9.64	14.06	13.08	6.59	12.63	9.99	7.05	11.08	10.79	-5.01	12.75
1650	8.95	12.97	11.06	8.97	11.69	9.26	4.88	10.98	10.06	-8.02	12.88
1651	6.15	9.85	8.34	6.32	7.74	5.51	2.71	7.87	6.07	-11.38	9.88
1652	5.43	10.28	6.47	0.63	8.54	5.47	4.44	10.58	7.31	-8.03	7.18
1653	7.31	10.97	9.78	6.18	9.80	7.48	7.93	8.29	7.68	-7.85	10.35
1526	16.84	18.98	18.38	12.12	17.76	15.54	7.00	12.61	16.76	-17.16	17.98
13665	8.98	13.27	11.24	4.94	12.61	10.49	7.24	12.42	10.95	-3.56	13.34
13670	12.54	15.16	13.19	4.70	13.16	11.67	8.05	12.11	11.92	-13.57	14.99
32867	6.18	11.11	8.43	3.26	9.85	7.22	8.14	9.43	6.84	-7.11	9.16
32874	4.45	8.71	8.26	0.57	8.38	6.28	0.92	8.76	6.22	-8.70	9.25
32904	8.87	12.35	12.77	5.55	12.16	10.15	4.52	13.71	10.89	-5.49	13.01
32906	6.45	10.02	10.26	2.84	10.35	8.25	5.27	12.19	9.72	-5.97	12.14
32907	5.62	10.35	10.28	1.53	8.56	6.76	2.90	13.51	9.79	-5.96	9.32
32908	12.54	13.95	13.98	4.81	13.97	12.48	5.79	10.92	14.17	-16.72	14.73
32909	10.98	11.92	14.03	3.74	13.70	11.60	5.14	9.60	12.38	-18.30	13.57
32911	5.29	8.99	13.22	8.14	8.61	7.01	4.05	11.99	10.36		11.00
32912	13.61	15.10	17.22	7.50	15.44	14.18	5.33	11.68	13.25	-12.95	17.23
32913	5.66	9.30	9.62	0.23	8.13	6.59	5.36	10.54	8.51	-4.80	10.41
32915	5.91	10.64	10.60	1.78	7.82	6.31	4.12	12.72	10.32	-5.27	9.75
32916	3.11	6.86	7.45	-0.91	6.32	4.74	-1.78	7.59	5.40	-10.75	9.01
32917	6.95	8.14	9.23	-3.65	8.63	7.21	3.88	11.29	8.03	-5.57	10.05
32919	12.14	15.06	14.93	7.86	13.47	12.54	6.51	8.51	14.26	-16.14	14.86
1577b-b	10.49	11.20	10.46	10.87	8.61	6.57	1.97	8.27	14.30	-4.75	11.18



	Ala	Asp	Glu	Gly	Ile	Leu	Lys	Phe	Pro	Thr	Val
1577b-a	1.04	0.43	0.14	1.40	0.84	0.97	1.00	0.41	1.09	0.41	0.10
1639	0.13	0.48	0.50	0.02	0.03	0.01	0.82	0.55	0.02	0.92	0.04
1641	0.18	0.65	0.34	0.14	0.38	0.12	0.30	1.38	0.10	0.74	0.18
1642	0.05	0.28	0.78	0.15	0.99	0.56	1.49	0.55	0.06	0.52	0.14
1645	0.14	0.05	0.07	0.18	0.51	0.45	0.11	1.07	0.31	0.07	0.35
1647	0.10	0.45	0.88	0.10	0.01	0.06	0.90	0.35	0.03	0.58	0.17
1648	0.03	0.42	0.46	0.33	0.10	0.15	0.13	0.41	0.19	0.57	0.67
1650	0.28	0.17	0.57	0.32	0.14	0.05	0.64	0.44	0.09	0.14	0.41
1651	0.04	0.01	0.46	0.06	0.06	0.17	0.01	0.02	0.01	0.72	0.01
1652	0.05	0.53	0.53	0.10	0.18	0.08	1.62	0.50	0.04	0.75	0.35
1653	0.12	0.36	0.08	0.19	0.22	0.27	2.26	0.35	0.26	0.29	0.48
1526	0.20	0.42	0.91	0.19	0.35	0.25	0.27	1.28	0.05	0.06	0.47
13665	0.24	0.21	0.06	0.33	0.81	0.82	2.51	0.34	0.22	0.05	0.51
13670	0.08	0.06	0.25	0.11	0.52	0.35	1.45	0.11	0.22	0.57	0.17
32867	0.01	0.08	0.88	0.06	0.00	0.16	0.13	0.81	0.12	0.38	0.09
32874	0.14	0.24	0.22	0.18	0.13	0.17	1.10	0.87	0.02	0.36	0.32
32904	0.09	0.10	0.25	0.06	0.68	0.55	0.14	0.21	0.12	0.54	0.27
32906	0.19	0.01	1.37	0.34	0.18	0.14	0.84	0.74	0.33	1.45	1.27
32907	0.41	0.35	0.74	0.47	0.21	0.04	2.24	1.14	0.52	0.43	0.38
32908	0.06	0.07	0.92	0.05	0.14	0.04	2.04	0.67	0.14	0.64	0.18
32909	0.09	0.61	0.43	0.13	0.15	0.22	1.21	1.18	0.33	1.27	0.65
32911	0.14	0.84	0.51	0.09	1.32	1.13	0.81	1.41	0.32		0.98
32912	0.28	0.03	1.34	0.30	1.10	1.02	0.49	0.28	0.19	0.41	0.86
32913	0.23	0.04	0.55	0.13	0.41	0.05	0.76	1.06	0.07	0.49	0.14
32915	0.26	0.07	0.40	0.00	0.79	0.81	2.00	1.29	0.02	0.67	0.17
32916	0.47	1.12	1.29	0.50	0.11	0.09	2.08	0.59	0.62	1.51	0.45
32917	0.27	0.23	0.72	0.20	1.19	0.67	0.94	0.73	0.24	1.05	0.61
32919	0.11	0.02	0.43	0.21	0.20	0.20	1.30	1.41	0.21	0.13	0.35
1577b-b	0.26	0.13	0.23	0.07	0.28	0.33	0.05	0.40	0.10	0.46	0.21

Table S10 Carbon and nitrogen isotope data on the standard bovine liver 1577b

Carbon isotope ratios

	Ala	SD	Asp	SD	Glu	SD	Gly	SD	Ile	SD	Leu	SD	Lys	SD
1577b-a	-19.52	1.48	-13.04	0.52	-16.86	0.35	-8.01	0.04	-24.06	0.61	-29.22	0.22		
1577b-b	-20.36	0.39	-12.55	0.03	-16.10	0.85	-9.30	0.39	-22.45	0.03	-29.29	0.13		
1577b-c	-19.59	0.23	-12.83	0.46	-16.73	0.94	-7.00	0.76	-24.51	0.21	-30.48	0.12	-15.89	1.20

Nitrogen isotope ratios

	Ala	SD	Asp	SD	Glu	SD	Gly	SD	Ile	SD	Leu	SD	Lys	SD
1577b-a	9.71	1.04	10.61	0.43	10.60	0.14	9.10	1.40	7.68	0.84	5.37	0.97	3.53	1.00
1577b-b	10.49	0.26	11.20	0.13	10.46	0.23	10.87	0.07	8.61	0.28	6.57	0.33	1.97	0.05
1577b-c	10.91	0.21	8.98	0.65	9.85	0.36	12.20	0.28			2.91	0.08		

Met	SD	Phe	SD	Pro	SD	Thr	SD	Val	SD
		-25.16	0.00	-18.48	0.40			-25.54	0.91
		-24.51	0.09	-18.86	0.03			-24.86	0.58
-23.05	0.84	-25.55	0.57	-18.69	0.94	-7.23	0.25	-25.28	0.06

Phe	SD	Pro	SD	Ser	SD	Thr	SD	Val	SD
8.36	0.41	13.66	1.09			-6.19	0.41	10.04	0.10
8.27	0.40	14.30	0.10			-4.75	0.46	11.18	0.21
8.11	0.04	13.15	0.14	9.37	0.81	-6.36	0.63	11.10	0.66