

# A new mathematical symbol: the terminial

Claude-Alexandre Simonetti

# ▶ To cite this version:

Claude-Alexandre Simonetti. A new mathematical symbol: the termirial. 2020. hal-02556204v3

# HAL Id: hal-02556204 https://hal.science/hal-02556204v3

Preprint submitted on 16 May 2020 (v3), last revised 3 Jun 2020 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A new mathematical symbol: the terminial

Claude-Alexandre Simonetti<sup>1, 2, 3</sup>

 $^{1}LPC$  Caen  $^{2}EAMEA$  Cherbourg  $^{3}ENSICAEN$  (Dated: 16 mai 2020)

The understanding of probability can be difficult for a few young scientists. Consequently, this new mathematical symbol, related to binomial coefficients, could be helpful to science education, and perhaps to other scientific subjects as well, such as computer science.

#### I. STUDY OF BINOMIAL COEFFICIENTS

#### A. Reminder

In a set of n elements, the number of combinations of parts of p elements is the following:

$$\binom{n}{p} = \frac{n!}{p! \ (n-p)!} \tag{1}$$

With  $\binom{n}{p}$  the binomial coefficient,  $(n,p) \in \mathbb{N}^2$  and  $p \leq n$ . And the exclamation point in equation (1) is named "factorial" and is defined as,  $\forall n \in \mathbb{N}^*$ :

$$n! = n \cdot (n-1) \cdot (n-2) \dots 3 \cdot 2 \cdot 1$$
  
 $n! = \prod_{i=1}^{n} i$  (2)

For n = 0, by definition : 0!=1.

#### B. From factorial to terminial

For its part, the terminial is defined as,  $\forall n \in \mathbb{N}^*$ :

$$n_{+}^{\mid} = n + (n - 1) + \dots + 3 + 2 + 1$$
 $n_{+}^{\mid} = \sum_{i=1}^{n} i$  (3)

The termirial symbol is almost like the factorial symbol, with just a little difference, however: the dot  $(\cdot)$  of the interrogation point, which could be a reminder of a multiplication, is replaced by a "plus" (+) sign. Indeed, instead of multiplying factors of a multiplication – factorial – one adds terms of an addition, hence the name of "termirial". Up to this point, things look quite basic. However, as the termirial is the  $n^{th}$  partial sum of an arithmetic sequence  $(U_n)_{n\in\mathbb{N}^*}$  with  $U_1=1$  as first term and r=1 as common difference, one can notice a first thing,  $\forall n\in\mathbb{N}^*$ :

$$n_{+}^{\mid} = \frac{n \cdot (n+1)}{2} = \binom{n+1}{2} = \binom{n+1}{n-1}$$
 (4)

One will get back to it later, with the "generalized" termirial.

### C. Intellectual path

At the beginning, I was studying the black body radiation (Max Planck's law). In this context, suppose one has n particles to fill up in 2 discrete energy levels determined by quantum mechanics. Suppose that the first energy level named  $m_1$  can contain p particles, and that the second one  $(m_2)$  can contain (n-p) particles. The number of possible combinations is same as equation (1):

$$\binom{n}{p} = \frac{n!}{p! \ (n-p)!} \tag{5}$$

Let us start with a simple example : n=5 and p=2. The figure 1 tries to explain the intellectual path which has led to the termirial. One can see the energy level  $m_1$  containing p=2 particles, with all the ways to fill it up, for particles particules ranked from 1 to 5. In this case, as a reminder, it is an unordered sampling without replacement, the rank order of the particles has no importance :  $\{1\ 2\}$  and  $\{2\ 1\}$  are counted only once. Finally :  $\binom{5}{2}=4 \ | =10$ .

Let us make things a little bit more complicated, with n=5 and p=3. In terms of binomial coefficients, one has the same result as before:  $\binom{5}{3} = \binom{5}{2} = 10$ . But the intellectual path is a bit different, see figure 2.

One can see a kind of "termirial of termirials":

$$n_{+}^{(2)} = n_{+}^{\mid} + (n-1)_{+}^{\mid} + \dots + 3_{+}^{\mid} + 2_{+}^{\mid} + 1_{+}^{\mid}$$

$$n_{+}^{(2)} = \sum_{k=1}^{n} k_{+}^{\mid} = \sum_{k=1}^{n} \sum_{i=1}^{k} i = \sum_{k=1}^{n} \frac{k(k+1)}{2!}$$
(6)

$$n \mid_{+}^{(2)} = \binom{n+2}{3} = \binom{n+2}{n-1} = \frac{n(n+1)(n+2)}{3!} \quad (7)$$

The transition from (6) to (7) can be done with mathematical induction. Indeed, if the following proposition P(n) is true:

$$n_{+}^{(2)} = \sum_{k=1}^{n} \frac{k(k+1)}{2!} = \frac{n(n+1)(n+2)}{3!}$$
 (8)




FIGURE 1. This figure represents the intellectual path from the binomial coefficient  $\binom{5}{2}$  to the terminal of 4, through a classical tree view. Here, only the energy level  $m_1$  is represented, which is sufficient, because one only has 2 energy levels, not more.




FIGURE 2. This figure represents the intellectual path from the binomial coefficient  $\binom{5}{3}$  to a sum of termirials, a kind of "termirial of termirials" or a  $2^{nd}$  termirial. From now, one can generalize to the  $3^{rd}$  termirial, the  $4^{th}$  termirial, etc.

And, for P(1):

$$1 \downarrow 2 = \sum_{k=1}^{1} \frac{k(k+1)}{2} = \frac{1 \cdot (1+1)}{2} = 1$$
$$\binom{3}{3} = \frac{1(1+1)(1+2)}{3!} = 1$$

Finally, with P(n+1):

$$(n+1)_{+}^{(2)} = \sum_{k=1}^{n} \frac{k(k+1)}{2} + \frac{(n+1)(n+2)}{2}$$

$$= \frac{n(n+1)(n+2)}{3!} + \frac{(n+1)(n+2)}{2}$$

$$= \frac{n(n+1)(n+2)}{6} + 3\frac{(n+1)(n+2)}{6}$$

$$(n+1)_{+}^{(2)} = \frac{(n+1)(n+2)(n+3)}{3!}$$
(9)

QED.

From now on, one can generalize to the  $p^{th}$  terminal

#### II. GENERALIZATION OF THE TERMIRIAL

# A. The $p^{th}$ terminal

Obviously, the the  $p^{th}$  terminial will be defined as,  $\forall (n, p) \in \mathbb{N}^* \times \mathbb{N}^*$ :

$$n_{+}^{(p)} = \sum_{m=1}^{n} \sum_{l=1}^{m} \dots \sum_{j=1}^{k} \sum_{i=1}^{j} i$$
 (10)

$$n_{+}^{(p)} = \sum_{k=1}^{n} k_{+}^{(p-1)} = \sum_{k=1}^{n} \frac{1}{p!} \prod_{i=0}^{p-1} (k+i)$$
 (11)

$$n \stackrel{(p)}{\underset{+}{\mid}} = \binom{n+p}{p+1} = \binom{n+p}{n-1} \tag{12}$$

$$n_{+}^{(p)} = \frac{1}{(p+1)!} \prod_{i=0}^{p} (n+i)$$
 (13)

With, for the equation (10), a number of sigmas  $(\sum)$  symbols equal to p.

For p = 0, one can define the  $0^{th}$  terminal as:

$$n \mid n = n = \binom{n}{1}$$

The proof to go from (11) to (12) (or from (11) to (13)) can be done with mathematical induction as well. Indeed, admitting that the following P(n) propositionis true:

$$\sum_{k=1}^{n} \frac{1}{p!} \prod_{i=0}^{p} (k+i) = \frac{1}{(p+1)!} \prod_{i=0}^{p} (n+i)$$
 (14)

On a, pour P(1):

$$1_{+}^{(p)} = \frac{1}{p!} \prod_{i=0}^{p-1} (1+i) = \frac{p!}{p!} = 1$$

$$\frac{1}{(p+1)!} \prod_{i=0}^{p} (1+i) = \frac{(p+1)!}{(p+1)!} = 1$$
(15)

Et, pour P(n+1):

$$(n+1)_{+}^{(p)} = \sum_{k=1}^{n} \frac{1}{p!} \prod_{i=0}^{p-1} (k+i) + \frac{1}{p!} \prod_{i=0}^{p-1} (n+1+i)$$

$$= \frac{1}{(p+1)!} \prod_{i=0}^{p} (n+i) + \frac{(p+1)}{(p+1)!} \prod_{i=1}^{p} (n+i)$$

$$= \frac{(n+p+1)}{(p+1)!} \prod_{i=1}^{p} (n+i)$$

$$(n+1)_{+}^{(p)} = \frac{1}{(p+1)!} \prod_{i=1}^{p+1} (n+i) = \binom{n+p+1}{p+1}$$
 (16)

QED.

#### B. Pascal's rule

Let us remind you on Pascal's rule:

$$\binom{n}{p} + \binom{n}{p+1} = \binom{n+1}{p+1}$$

It is possible to adapt it to terminials:

$$(n+1) \mathop{\downarrow}^{(p)}_{\perp} + (n) \mathop{\downarrow}^{(p+1)}_{\perp} = (n+1) \mathop{\downarrow}^{(p+1)}_{\perp}$$

## III. PRATICAL APPLICATIONS

## A. Computational complexity

In computer science, the terminial can be used to calculate the complexity of computer programs with intricated "for" loops as follows: for example, with 4 "for" loops, with:

- for i from 1 to n=100;
- for j from 1 to i;
- for k from 1 to j;
- for 1 from 1 to k.

The complexity will be:

$$100 \Big|_{+}^{(4-1)} = {100+3 \choose 3+1} = {103 \choose 4} = {103 \choose 99} = 4\ 421\ 275$$

So, approximatey 4,4 millions of operations.

More generally, the complexity of this kind of program will be, according to equation (13) and  $\forall (n, p) \in$ 

$$\mathbb{N}^* \times \mathbb{N}^* : n \Big|_{\perp}^{(p-1)} = \Theta(n^p).$$

#### B. Pseudo-fractal aspect




FIGURE 3. This figure tries to convert  $p^{th}$  termirial of 4 into fractal figures, for p=0, p=1 and p=2. Only the grey squares are counted.

The figure 3 tries to convert  $p^{th}$  terminial of 4 into fractal figures, for p=0, p=1 and p=2. Only the grey

squares are counted. For example, for  $4^{1}_{+}$ , 20 squares

are in grey, which corresponds to  $4 \mid = 20$ . For each incrementation of p, the side of each square is divided by 2, and the same basic pattern is repeated:

- 4 when the 4 previous squares are in grey;
- 3 when the 3 previous squares are in grey;
- 2 when the 2 previous squares are in grey;
- 1 when only 1 previous square is in grey.

But the Hausdorff dimension (D) of this presumed fractal is not constant, and the limit of D as p approaches infinity is 2. Indeed, let a the side length of each square related to the  $p^{th}$  terminial. The total surface  $S_p$  of the squares in grey will be:

$$S_p = a^2 \cdot n \Big|_{+}^{(p)} = a^2 \cdot \frac{1}{(p+1)!} \prod_{i=0}^{p} (n+i)$$

And for the  $S_{p-1}$  surface, related to the  $(p-1)^{th}$  terminial, for a side length of each square doubled in size:

$$S_{p-1} = 4a^2 \cdot n \Big|_{+}^{(p-1)} = 4a^2 \cdot \frac{1}{(1)!} \prod_{i=0}^{p-1} (n+i)$$

Hence, for p > 0:

$$\frac{S_{p-1}}{S_p} = 4 \cdot \frac{p+n}{p+1} = 4 \cdot \frac{1+n/p}{1+1/p}$$

Finally, this surface ratio, as p approaches infinity, is 4:

$$\lim_{p\to\infty}\frac{S_{p-1}}{S_p}=4$$

Consequently, the Hausdorff dimension (D) is 2, as  $2^D = 4$ , when p approaches infinity. To conclude, as D is not constant, even if figure 3 looks interesting, termirials do not seem to be related to fractals.

## IV. CONCLUSION

The terminial is a symbol which could be helpful to scientific students, for the understanding of probability. As well, it could be perhaps helpful in other subjects than science education, that are out of this pré-publication.

<sup>[1]</sup> B. Pascal, Traité du triangle arithmétique (Arvensa Editions, 2019).

<sup>[2]</sup> ISO 80000-2:2009(E), Quantities and units (International Organization for Standardization, 2009).

<sup>[3]</sup> C.-A. Simonetti, "Un nouveau symbole mathématique : la termirielle," (2020), working paper or preprint.