


# Non linear dynamics of premixed flames in a Hele-Shaw burner

**Basile Radisson, Christophe Almarcha, Elias Al Sarraf, Joel Quinard, Bruno Denet**


*Aix-Marseille Université, CNRS, Centrale Marseille, IRPHE Marseille France*

# Premixed flame instabilities in a Hele-Shaw burner (thickness 4.7 mm)

- Joulin and Sivashinsky CST (1994)
- Sharif Abid Ronney (1999)
- Kang Baek and Im CTM (2006)
- Recent works




# Flame propagation in a Hele-Shaw cell



# Flame propagation (methane $\phi=0.88$ )

- Central ignition: small slopes
- Lateral ignition: large slopes
- Same properties in 3D: Almarcha Denet Quinard



# Sivashinsky equation

- Small expansion development, several orders
- Sivashinsky 77, Sivashinsky Clavin 87, Kazakov 2005  
Joulin Cambray 92
- Small slopes
- Non dimensional form: one parameter, inverse of the number of unstable modes

$$\sigma = 4 \frac{\sigma_M}{k_c} \left( |k| - \frac{k^2}{k_c} \right)$$

$$\phi_t + \frac{u_A}{2} \phi_x^2 = \frac{4\sigma_M}{k_c} \left( \frac{\phi_{xx}}{k_c} + I(\phi, x) \right)$$

$$\Phi_\tau + \frac{1}{2} \Phi_X^2 = \nu \Phi_{XX} + I(\Phi, X)$$


# Sivashinsky equation : pole solutions

- Poles in complex conjugate pairs
- Differential equations for the position of poles in the complex plane



$$\phi(x, t) = -\frac{8\sigma_M}{u_A k_c^2} \sum_{n=1}^{2N} \ln \left( \sin \left( \frac{\pi (x - z_n(t))}{\Lambda} \right) \right)$$

$$\dot{z}_n = \frac{4\sigma_M}{k_c} \left[ \frac{2\pi}{\Lambda k_c} \sum_{p \neq n} \cot \left( \frac{\pi}{\Lambda} (z_p - z_n) \right) - i \operatorname{sign}(Im(z_n)) \right]$$



# Pole dynamics



# 2D Bunsen flame at the top of the cell + forcing plate




# Amplitude vs time: growth rate measurement



# Dispersion relation: propane $\phi=0.8$


## parameters $\sigma_M$ $k_c$



$$\sigma = 4 \frac{\sigma_M}{k_c} \left( |k| - \frac{k^2}{k_c} \right)$$


# Printing of perturbations on the flame

- Laser cutting : pole solution for the shape of the forcing plate




# Forcing an initial condition

- Crests correspond to the forcing plate



# Forcing an initial condition




# d2 law

- d distance between close cusps


$$d^2(t) = d_0^2 - \frac{64\sigma_M}{k_c^2} t$$

Propane-air flame  $\phi=0.8$  Pole dynamics



# Repeatability

- Crests evolution, several experiments






# Comparison with the Sivashinsky equation


Propane-air flame  $\phi=0.8$

Sivashinsky equation


Pole dynamics




# Pole dynamics: purely aggregative dynamics




PDE + noise: permanent addition of  
new crests




# Experimental cell size distribution (downward propagating propane flames)




(a)



(b)



(d)



(e)

# Conclusion

- Flame instability in a Hele-Shaw burner:  
forcing an initial condition
- Sivashinsky equation works reasonably for  
short times/ small slopes
- Deterministic evolution of crests  
trajectories for short times (pole  
dynamics)
- Cells creation : some role of noise and  
gravity