

# Chronic wound management in Cameroon: The need to integrate traditional healers contribution and patients mental health status

Lucrèce Eteki, Joel Djatche, Rodrigue Ntone, Estelle Metsago, Daniel Nzedou, Alex Mondomobe, Earnest Njih, Franck Wanda, Yves Hako, Christian Minyem

### ▶ To cite this version:

Lucrèce Eteki, Joel Djatche, Rodrigue Ntone, Estelle Metsago, Daniel Nzedou, et al.. Chronic wound management in Cameroon: The need to integrate traditional healers contribution and patients mental health status. OMS conférence sur l'ulcère de Buruli, Mar 2019, Genève, Switzerland. hal-02560350

HAL Id: hal-02560350

https://hal.science/hal-02560350

Submitted on 1 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# CHRONIC WOUND MANAGEMENT IN CAMEROON: THE NEED TO INTEGRATE

Lucrèce Eteki<sup>1</sup>, Joel Djatche<sup>2</sup>, Rodrigue Ntone<sup>1</sup>, Estelle Metsago<sup>2</sup>, Daniel Nzedou<sup>2</sup>, Alex Mondomobe<sup>3</sup>, Earnest Njih Tabah <sup>4</sup>, Franck Wanda<sup>5</sup>, Yves Hako<sup>3</sup>, Christian Minyem<sup>6</sup>,
Radhika Sundararajan<sup>7</sup>, Yap Boum<sup>1</sup>

¹Epicentre, Yaoundé, Cameroon; ²UNIPSY, Yaoundé, Cameroon; ³Hopital Regional Annexe d'Ayos; ⁴ CNLP2LUB Ministry of Health, Cameroun; ⁵CIRES, Akonolinga, Cameroon; ⁴FAIRMED, Bankim, Cameroon; ¹ Well Cornell Medical College, NY, USA

### INTRODUCTION

### BACKGROUND

Delays in diagnosis and non-adherence to biomedical treatment have been linked to:

Psychological status of patients

Local beliefs in the mystical origin of the disease

Preference/or combination with traditional treatment

### **OBJECTIVES**

To assess perceptions and practices of (Buruli Ulcer) BU management among patients, traditional healers, and Hospital workers

To evaluate psychological state and perceived stigmatization of patients with chronic wounds including BU

### METHODS AND MATERIALS

Bankm

### Study sites and period

- BU endemic zones : Akonolinga, Ayos, Bankim
- between April and December 2018

### Study design

Mixed methods

### **Qualitative Data**

- Sampling
- -Patients local traditional healers (as identified by community rep
- -Patients, local traditional healers (as identified by community representatives ), and HCWs were approached for participation.
- Data collection

FGDs and In-depth interviews were organised around

- ✓ Current understanding about BU causes and treatment
- ✓ patients' engagement with biomedical ressources and/or traditional medicine
- ✓ Potential collaboration between HCWs and traditional healers
- ✓ Subjective experience of patients with chronic wounds
- ✓ Data were collected in clinical and community settings
- Data analysis
  - -Recordings were transcribed and analyzed using a thematic content analysis framework (Krippendorf 2012).
  - -NVivo software V.10 (QSR International, Doncaster, Australia) was used to assist with organization and management of the dataset.

Patient interview with psychologist

Bankim



Patients sufferring from BU



FGDs with traditionnal healers Bankim

### **Quantitative Data**

- Sampling
  - -current and former patients in the study sites were approcahed by community representatives for participation
  - -data were collected within clinical settings
- Data collection
  - -psychological scales M.I.N.I to determine psychological pathologies -psychological scale QSCPGS to determine body satisfaction and self-esteem
- -questionnaires to determine perceived stigmatization
- Data analysis
  - -Data were entered in excel and analyzed in R 3.5.3

### RESULTS

## Perception, experience and practices on BU

- We included 58 participants: 17 patients, 29 traditional healers, 12 HCWs
- Buruli Ulcer infection has a mystical origin and can result from a person casting a spell.
- An important proportion of patients are anxio-depressive
- Patients combine traditional and biomedical therapies to heal faster by addressing both the mystical and infectious components of the disease.
- Healthcare workers and traditional healers both reported that patients resist receiving treatment at health facilities
- Biomedical treatments are usually considered after failure of traditional therapies.
- Traditional healers are open to collaborate with physicians, but physicians are rather hostile to such collaboration


"I think that if we combine the two, healing would be faster ... And when you come back to the hospital now that you have removed the mystical barrier that prevented you from healing, it is easier to treat you. For me, I think we should combine the two medicines." [Akonolinga patient]

"Since it (BU) all started, I am sad, I am useless, I fear that my wife will abandon me..." [Akonolinga patient]

"Here, the hospital is the last resort" [Ayos Physician]

disease.

# Psychological pathologies 35,6 24,7 16,4 11,0 DEPRESSION SUICIDE RISK DYSTHYMIA POST-TRAUMATIC STATE



### DISCUSSION AND CONCLUSION

• Patients resort to the hospital after failure of traditional therapies.

therapies.They prefer a combination of treatment to adress the

• Psychological pathologies are numerous and persist even long after healing

Former patients still feel discriminated

• It is important to include mental health interventions in wound management programs

• It is important to educate communities about the realities of BU to avoid stigmatization and facilitate reintegration of patients

It is necessary to implement patient centered programs that are culturally competent enough to integrate traditional considerations as well as social and psychological aspects

**Acknowledgements**: We are thankful to all the patients that contributed to this study. We appreciate the Ministry of Health that supported and collaborated with us to implement this study. We are thankful for our partners Hopital de District d'Akonolinga, CMA Endom, Hopital de District de Bankim, CMA de SonKolong. This study was funded by Medecins Sans frontières