

## Gradient, scattering and other kinds of longitudinal optical forces exerted by off-axis Bessel beams in the Rayleigh regime in the framework of generalized Lorenz-Mie theory, 246, Paper 196913

Gérard Gouesbet

### ▶ To cite this version:

Gérard Gouesbet. Gradient, scattering and other kinds of longitudinal optical forces exerted by offaxis Bessel beams in the Rayleigh regime in the framework of generalized Lorenz-Mie theory, 246, Paper 196913. Journal of Quantitative Spectroscopy and Radiative Transfer, 2020, 246, pp.106913. 10.1016/j.jqsrt.2020.106913. hal-02562373

### HAL Id: hal-02562373 https://hal.science/hal-02562373

Submitted on 7 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## Gradient, scattering and other kinds of longitudinal optical forces exerted by off-axis Bessel beams in the Rayleigh regime in the framework of generalized Lorenz-Mie theory.

### Gérard Gouesbet

CORIA-UMR 6614- Normandie Université
CNRS-Université et INSA de Rouen
Campus Universitaire du Madrillet
76800, Saint-Etienne-du Rouvray, France.
Corresponding author: gouesbet@coria.fr

February 18, 2020

#### Abstract

After Arthur Ashkin's pioneering work in optical levitation and manipulation, the study of optical forces exerted by laser beams on particles has become an active field of research. The present paper is a contribution to this issue. The interest of Bessel beams is that their intensity does not have any longitudinal gradient along the direction of propagation leading to a trivial separation between gradient and scattering forces. Beside the classical gradient and scattering forces, we shall however exhibit a new kind of optical forces associated with the existence of non zero axicon angles.

Keywords: optical forces; gradient and scattering optical forces; Bessel beams; generalized Lorenz-Mie theory; Rayleigh regime.

### 1 Introduction.

After Arthur Ashkin's pioneering work in optical levitation and manipulation, e.g. a collection of preprints in [1], a great deal of work has been devoted to the study of optical forces exerted by laser beams on particles, e.g. [2], [3], [4], [5], [6], [7], [8], [9], [10] for reviews.

1

<sup>© 2020</sup> published by Elsevier. This manuscript is made available under the CC BY NC user license https://creativecommons.org/licenses/by-nc/4.0/

Many of them focused their attention on the classical gradient and scattering forces discussed by Ashkin according to the following definitions: the gradient force is proportional to the gradient of intensity and points in the direction of the intensity gradient, while the scattering force is proportional to the optical intensity and points in the direction of the incident light [11]. The study of gradient and scattering forces is particularly appealing in the Rayleigh regime where the formal computations are facilitated and are even amenable to analytic formulations, e.g. [12], [13], [14], [15], [16], [17], [18] [19], [20], [21], [22], [23], [24], [25], [26], [27] and some reviews above. I shall particularly rely on a paper by J.A. Lock who, in the framework of a study devoted to radiation trapping forces for laser tweezers, provided an analysis of gradient and scattering forces in the Rayleigh regime, in the framework of generalized Lorenz-Mie theory, in the case of an on-axis Gaussian beam described by using a localized approximation in the weak-beam confinement limit [28].

In the present paper, a similar study is carried out in the more difficult case of off-axis Bessel beams. The opportunity of studying off-axis beams instead of on-axis beams is partly due to the fact that some simplifications occur due to the fact that the gradient of intensity along the axis z of propagation is always zero for Bessel beams, in deep contrast with the case of Gaussian beams even in an on-axis configuration. Furthermore, this fact implies a trivial separation between longitudinal gradient and scattering forces since all longitudinal gradient forces are expected to be zero so that only scattering forces act along the longitudinal direction. Another interest of Bessel beams is that the beam shape coefficients encoding the structure of the beams in the framework of generalized Lorenz-Mie theory (GLMT) are available in closed forms, without the need of using a localized approximation. Furthermore, an unexpected result will be the discovery of a new kind of optical forces related to the existence of axicon angles in Bessel beams, which do not occur in Gaussian beams.

The paper is organized as follows. Section 2 provides a background to be used in the sequel, concerning (i) optical forces in the framework of GLMT, (ii) Poynting vector in the framework of GLMT, (iii) basic mathematical formulae for further use and (iv) the description of Bessel beams. Section 3 deals with optical forces for Rayleigh particles exerted by Bessel beams. In Section 4, the optical forces derived in the previous section are analyzed in terms of gradient and scattering forces, revealing the existence of an unexpected new kind of optical forces, namely scattering axicon forces which do not occur for more usual beams whose axicon angle is zero.

### 2 Physical and mathematical background.

The GLMT *stricto sensu* describes the interaction between an arbitrary (structured) shaped beam and a homogeneous spherical particle defined by its diameter and its complex refractive index, e.g. [29], [30], [31], and references

therein. Also, see reviews [32], [33], in particular for other GLMTs devoted to the interaction between laser beams and other kinds of scatterers. In the present paper, we consider a Rayleigh particle located at the origin  $O_P$  of a Cartesian coordinate system  $O_P xyz$  illuminated by an off-axis Bessel beam propagating along the z-direction.

### 2.1 Optical forces.

In the GLMT framework, we express the optical force components  $F_i$  by using (unnormalized) pressure radiation cross-section components  $C_{pr,i}$  (i = x, y, z). In the present paper, we deal only with the longitudinal cross-section (along z-direction) reading as:

$$C_{pr,z} = E_0 H_0^* \frac{\lambda^2}{4\pi} \sum_{n=1}^{\infty} \sum_{m=-n}^{n} \{ \frac{1}{(n+1)^2} \frac{(n+1+|m|)!}{(n-|m|)!}$$
(1)  

$$\operatorname{Re}[(a_n + a_{n+1}^* - 2a_n a_{n+1}^*) g_{n,TM}^m g_{n+1,TM}^{m*} + (b_n + b_{n+1}^* - 2b_n b_{n+1}^*) g_{n,TE}^m g_{n+1,TE}^{m*}]$$

$$+ m \frac{2n+1}{n^2(n+1)^2} \frac{(n+|m|)!}{(n-|m|)!}$$

$$\operatorname{Re}[i(2a_n b_n^* - a_n - b_n^*) g_{n,TM}^m g_{n,TE}^{m*})] \}$$

with notations being the ones of [29]. However, note that, strictly speaking,  $C_{pr,z}$  of Eq.1 would be a pressure radiation cross-section, i.e. would be homogeneous to an area, if we used the normalization condition  $E_0H_0^*/2 = 1$ . Here, we did not use this condition for the sake of consistency with later expressions concerning the Poynting vector. When using the normalization condition, the forces are related to the cross-sections according to  $F_i = C_{pr,i}/c$  ([34], p.14). The time-dependence of the wave is  $\exp(+i\omega t)$  which is the usual choice in GLMT.

Furthermore, specifically,  $a_n$  and  $b_n$  are the usual Mie coefficients of the usual Lorenz-Mie theory,  $g_{n,TM}^m$  and  $g_{,TE}^m$ , with TM and TE standing for "Transverse Magnetic" and "Transverse Electric" respectively, are the beam shape coefficients (BSCs) encoding the description of the beam, either in terms of scalar potentials, e.g. [31] or in terms of vector spherical wave functions [35], the star denotes a complex conjugation and  $\delta_{nm}$  is the Kronecker symbol. Furthermore, we have reintroduced the normation prefactor  $E_0H_0^*/2$  which was set to 1 in original works, e.g. Eq.(3.106) in [31]. In the present paper, we shall consider only the longitudinal force of Eq.1 because it will lead to simpler computations than for the two other force components along x and y, and, more important, because we expect a natural separation between gradient and scattering forces along the z-direction.

### 2.2 Poynting vector.

For the same reasons, we shall only consider the z-component  $S_z$  of the Poynting vector defined as:

$$S_{z} = \frac{1}{2} \operatorname{Re}(E_{x}H_{y}^{*} - E_{y}H_{x}^{*})$$
(2)

which has been already evaluated to [36], [37]:

$$S_z = \frac{-E_0 H_0^*}{2r^2} \operatorname{Re} \sum_{n=1}^{\infty} \sum_{m=-n}^{+n} \sum_{p=1}^{\infty} \sum_{q=-p}^{+p} i c_n^{pw} c_p^{pw*} e^{i(m-q)\varphi}$$
(3)  
$$(\sin \theta S_{np}^{mq} + \cos \theta C_{np}^{mq})$$

in which  $(r, \theta, \varphi)$  are spherical coordinates attached to the Cartesian coordinates  $(x, y, z), c_n^{pw}$  are prefactors appearing in the Bromwich formulation of the Lorenz-Mie theory [38], reading as:

$$c_n^{pw} = \frac{1}{ik} (-i)^n \frac{2n+1}{n(n+1)} \tag{4}$$

and :

$$S_{np}^{mq} = kr[-g_{n,TM}^{m}g_{p,TM}^{q*}\psi_{p}(\psi_{n}+\psi_{n}^{''})P_{n}^{|m|}\tau_{p}^{|q|} + g_{n,TE}^{m}g_{p,TE}^{q*}\psi_{n}(\psi_{p}+\psi_{p}^{''})P_{p}^{|q|}\tau_{n}^{|m|} + qg_{n,TM}^{m}g_{p,TE}^{q*}\psi_{p}^{'}(\psi_{n}+\psi_{n}^{''})P_{n}^{|m|}\pi_{p}^{|q|} + mg_{n,TM}^{m}g_{p,TE}^{q*}\psi_{n}^{'}(\psi_{p}+\psi_{p}^{''})P_{p}^{|q|}\pi_{n}^{|m|}]$$
(5)

$$C_{np}^{mq} = -g_{n,TM}^{m}g_{p,TM}^{q*}\psi_{p}\psi_{n}^{'}(\tau_{n}^{|m|}\tau_{p}^{|q|} + mq\pi_{n}^{|m|}\pi_{p}^{|q|})$$
(6)  
+ $g_{n,TM}^{m}g_{p,TE}^{q*}\psi_{n}^{'}\psi_{p}^{'}(m\pi_{n}^{|m|}\tau_{p}^{|q|} + q\pi_{p}^{|q|}\tau_{n}^{|m|})$   
- $g_{n,TE}^{m}g_{p,TM}^{q*}\psi_{p}\psi_{n}(m\pi_{n}^{|m|}\tau_{p}^{|q|} + q\pi_{p}^{|q|}\tau_{n}^{|m|})$   
+ $g_{n,TE}^{m}g_{p,TE}^{q*}\psi_{n}\psi_{p}^{'}(mq\pi_{n}^{|m|}\pi_{p}^{|q|} + \tau_{n}^{|m|}\tau_{p}^{|q|})$ 

in which k is the wavenumber,  $P_n^{|m|} = P_n^{|m|}(\cos\theta)$  are associated Legendre functions,  $\tau_n^{|m|} = \tau_n^{|m|}(\cos\theta)$  and  $\pi_n^{|m|} = \pi_n^{|m|}(\cos\theta)$  are generalized Legendre functions, and  $\psi_n = \psi_n(kr)$  are Ricatti-Bessel functions, while a prime denotes a derivative with respect to the argument (and a double prime a second derivative). In Eqs.5-6, arguments are omitted for convenience

### 2.3 Mathematical background.

Associated Legendre functions are defined according to Hobson's notation [39]:

$$P_n^{|m|}(\cos\theta) = (-1)^{|m|}(\sin\theta)^{|m|} \frac{d^{|m|}P_n(\cos\theta)}{(d\cos\theta)^{|m|}}$$
(7)

in which  $P_n(\cos \theta)$  are the usual Legendre polynomials. For the description of Bessel beams, we shall also need associated Legendre functions with a negative superscript. They can be obtained from [40]:

$$P_n^{-m}(\cos\theta) = (-1)^m \frac{(n-m)!}{(n+m)!} P_n^m(\cos\theta)$$
(8)

Associated Legendre functions may then be evaluated according to:

$$\tau_n^m(\cos\theta) = \frac{dP_n^m(\cos\theta)}{d\theta} \tag{9}$$

$$\pi_n^m(\cos\theta) = \frac{P_n^m(\cos\theta)}{\sin\theta} \tag{10}$$

Ricatti-Bessel functions  $\psi_n(kr)$  may be expressed in terms of spherical Bessel functions  $j_n(kr)$  according to:

$$\psi_n(x) = x j_n(x) \tag{11}$$

Furthermore, we have:

$$\psi'_n(x) = (n+1)j_n(x) - xj_{n+1}(x) \tag{12}$$

and:

$$\lim_{x \to 0} j_n(x) = \frac{x^n}{(2n+1)!!} \tag{13}$$

### 2.4 Bessel beams.

Bessel beams have been introduced by Durnin and co-workers [41], [42]. They possess the appealing property of being self-healing and non diffracting, and, more important in the context of the present paper, they furthermore possess a propagation invariance property, namely the intensity of the beam is constant along the direction of propagation. There exist an infinity of kinds of Bessel beams, in particular depending on the value given to an arbitrary function  $g(\alpha_0)$ , in which  $\alpha_0$  is the axicon angle (or half-cone angle) of the beam, with different linear and circular polarizations [43], [44], [45], [46]. All of them generically exhibit the following structure:

$$K_i = k_i(x, y)e^{-ik_z z}, \ i = x, y, z, \ K = E \text{ or } H$$
 (14)

in which  $k_z = k \cos \alpha$  is the longitudinal wavenumber. Among all of them, we shall pay a particular attention to circularly symmetric Bessel beams of arbitrary order whose BSCs in an off-axis configuration read as [46]:

$$g_{n,TM}^{m} = -g(\alpha_{0})e^{ik_{z}z_{0}}[i^{l-m+1}e^{i(l-m+1)\phi_{0}}A_{n}^{m} + i^{l-m-1}e^{i(l-m-1)\phi_{0}}B_{n}^{m}] \quad (15)$$

$$g_{n,TE}^{m} = ig(\alpha_0)e^{ik_z z_0}[i^{l-m+1}e^{i(l-m+1)\phi_0}A_n^m - i^{l-m-1}e^{i(l-m-1)\phi_0}B_n^m]$$
(16)

in which:

$$A_n^m = (-1)^{(m-|m|)/2} \frac{(n-m)!}{(n+|m|)!} J_{l-m+1}(\sigma_0) [\tau_n^m(\cos\alpha_0) + m\pi_n^m(\cos\alpha_0)] \quad (17)$$

$$B_n^m = (-1)^{(m-|m|)/2} \frac{(n-m)!}{(n+|m|)!} J_{l-m-1}(\sigma_0) [\tau_n^m(\cos\alpha_0) - m\pi_n^m(\cos\alpha_0)] \quad (18)$$

in which  $\sigma_0 = k_t \rho_0$ ,  $k_t = k \sin \alpha_0$  is the transverse wavenumber,  $\rho_0 = (x_0^2 + y_0^2)^{1/2}$ ,  $\phi_0 = \tan^{-1}(y_0/x_0)$ ,  $(x_0, y_0, z_0)$  denotes the location of the beam origin with respect to the origin of the particle coordinate system at which the scatterer

is located, and therefore specifies the off-axis location of the beam, and  $J_k(.)$ is the k-order Bessel function of the first kind. When  $g(\alpha_0) = (1 + \cos \alpha_0)/4$ , the beam reduces to a Davis circularly symmetric Bessel beam as discussed in [43], [47]. When  $g(\alpha_0) = 1/2$ , the beam reduces to another kind of Bessel beam discussed in [48], [49], [50]. In the sequel, it will be convenient to introduce the notation  $C = \cos \alpha_0$ .

# 3 Optical forces exerted by Bessel beams on Rayleigh particles.

## 3.1 General expressions of optical forces for Rayleigh particles.

Let us consider Eq.1 which contains an infinite number of Mie coefficients. For Rayleigh particles, the Mie coefficients reduce to [28], and [34], pp. 143-144:

$$a_1 = \frac{2i}{3} \frac{m^2 - 1}{m^2 + 2} \alpha^3 + O(i\alpha^5) + \frac{4}{9} (\frac{m^2 - 1}{m^2 + 2})^2 \alpha^6$$
(19)

$$b_1 = O(i\alpha^5) \tag{20}$$

in which m is the refractive index (here taken to be real) with respect to the surrounding medium and  $\alpha$  the size parameter  $\pi d/\lambda$ . The other coefficients  $a_n$  and  $b_n$  (n > 1) involves still higher powers of  $\alpha$ . Real parts are then proportional to  $\alpha^6$  and imaginary parts are proportional to  $\alpha^3$  while higher powers are discarded and we therefore retain only:

$$\operatorname{Im}(a_1) = \frac{2}{3} \frac{m^2 - 1}{m^2 + 2} \alpha^3 \tag{21}$$

$$\operatorname{Re}(a_1) = \frac{4}{9} \left(\frac{m^2 - 1}{m^2 + 2}\right)^2 \alpha^6 \tag{22}$$

We then start from Eq.1, retain only the (n = 1)-partial wave and the Mie coefficient  $a_1$ , leading to:

$$C_{pr,z} = E_0 H_0^* \frac{3\lambda^2}{8\pi} \operatorname{Re}(a_1 G)$$
(23)

in which:

$$G = g_{1,TM}^{-1} (g_{2,TM}^{-1*} + ig_{1,TE}^{-1*}) + g_{1,TM}^{1} (g_{2,TM}^{1*} - ig_{1,TE}^{1*}) + \frac{1}{3} g_{1,TM}^{0} g_{2,TM}^{0*}$$
(24)

This becomes:

$$C_{pr,z} = E_0 H_0^* \frac{3\lambda^2}{8\pi} [\operatorname{Re}(a_1) \operatorname{Re}(G) - \operatorname{Im}(a_1) \operatorname{Im}(G)]$$
(25)

in which we recall that  $\operatorname{Re}(a_1) \sim \alpha^6$  and  $\operatorname{Im}(a_1) \sim \alpha^3$ . According to the literature, e.g. [12], [13], [20], [25], [28], and many other papers already quoted in the introduction, the  $\alpha^6$ -term and the  $\alpha^3$ -term correspond to the scattering and the gradient forces respectively so that  $C_{pr,z}$  is the sum of two terms according to  $C_{pr,z} = C_{pr,z}^s + C_{pr,z}^g$  in which, using Eqs.21-22, we obtain:

$$C_{pr,z}^{s} = E_0 H_0^* \frac{\lambda^2}{6\pi} (\frac{m^2 - 1}{m^2 + 2})^2 \operatorname{Re}(G) \alpha^6$$
(26)

$$C_{pr,z}^{g} = -E_0 H_0^* \frac{\lambda^2}{4\pi} \frac{m^2 - 1}{m^2 + 1} \operatorname{Im}(G) \alpha^3$$
(27)

in which the subscripts s and g recall that we expect that  $C_{pr,z}^s$  and  $C_{pr,z}^g$  correspond to scattering and gradient forces respectively (this expectation will not be perfectly fulfilled). We shall now specify these expressions to the case of circularly symmetric Bessel beams of arbitrary order whose BSCs in an off-axis configuration are given by Eqs.15-18. A remark is worthwhile to be done here, namely that G occur both in  $C_{pr,z}^s$  and in  $C_{pr,z}^g$ , i.e. that both quantities would be expressed in terms of the same set of BSCs. This suggests that it is in general impossible to express the original GLMT series of fields as a decomposition of sub-fields leading to different kinds of optical forces related to different sets of BSCs, as confirmed in a more general context by [51], [52]. [53].

# 3.2 Term with imaginary part of G for circularly symmetric Bessel beams.

Let us consider Eq.24. It is convenient to rewrite it as:

$$G = G_1 + G_2 + G_3 \tag{28}$$

in which:

$$G_1 = g_{1,TM}^{-1} (g_{2,TM}^{-1*} + ig_{1,TE}^{-1*})$$
(29)

$$G_2 = g_{1,TM}^1 (g_{2,TM}^{1*} - ig_{1,TE}^{1*})$$
(30)

$$G_3 = \frac{1}{3} g^0_{1,TM} g^{0*}_{2,TM} \tag{31}$$

We now evaluate these quantities using Eqs.15-18, and obtain, after a significant amount of computations:

$$G_{1} = -|g(\alpha_{0})|^{2} \{ -A_{1}^{-1}A_{2}^{-1} + A_{1}^{-1}A_{1}^{-1} - B_{1}^{-1}B_{2}^{-1} - B_{1}^{-1}B_{1}^{-1} + e^{2i\phi_{0}}[A_{1}^{-1}B_{2}^{-1} + A_{1}^{-1}B_{1}^{-1}] + e^{-2i\phi_{0}}[B_{1}^{-1}A_{2}^{-1} - B_{1}^{-1}A_{1}^{-1}] \}$$
(32)

$$G_{2} = |g(\alpha_{0})|^{2} \{A_{1}^{1}A_{2}^{1} + A_{1}^{1}A_{1}^{1} + B_{1}^{1}B_{2}^{1} - B_{1}^{1}B_{1}^{1} + e^{2i\phi_{0}}[A_{1}^{1}B_{1}^{1} - A_{1}^{1}B_{2}^{1}] - e^{-2i\phi_{0}}[B_{1}^{1}A_{2}^{1} + B_{1}^{1}A_{1}^{1}]\}$$
(33)

$$G_3 = \frac{1}{3} \left| g(\alpha_0) \right|^2 \left[ A_1^0 A_2^0 + B_1^0 B_2^0 - e^{2i\phi_0} A_1^0 B_2^0 - e^{-2i\phi_0} B_1^0 A_2^0 \right]$$
(34)

Recalling that  $A_n^m$  and  $B_n^m$  are real quantities, we then have:

$$\operatorname{Im}(G_3) = -\frac{1}{3} |g(\alpha_0)|^2 \sin 2\phi_0 [A_1^0 B_2^0 - B_1^0 A_2^0]$$
(35)

Expressing the coefficients  $A_n^m$  and  $B_n^m$  involved in Eq.35 by using Eqs.17-18, we obtain:  $A_1^0 = J_{l+1}\tau_1^0$ ,  $B_2^0 = J_{l-1}\tau_2^0$ ,  $B_1^0 = J_{l-1}\tau_1^0$  and  $A_2^0 = J_{l+1}\tau_2^0$ , in which arguments  $\sigma_0$  and  $C = \cos \alpha_0$  are omitted, immediately implying, without the need to specify the expressions for the  $\tau$ 's:

$$A_1^0 B_2^0 - B_1^0 A_2^0 = 0 (36)$$

and therefore:

$$\operatorname{Im}(G_3) = 0 \tag{37}$$

We now deal with  $(G_1 + G_2)$ , i.e. not with  $G_1$  and  $G_2$  separately. From Eqs.32-33, recalling again that  $A_n^m$  and  $B_n^m$  pertain to **R**, we obtain:

$$Im(G_1 + G_2) = |g(\alpha_0)|^2 \sin 2\phi_0 G_{12}$$
(38)

in which:

$$G_{12} = -A_1^{-1}B_2^{-1} - A_1^{-1}B_1^{-1} + B_1^{-1}A_2^{-1} - B_1^{-1}A_1^{-1} + A_1^{-1}B_1^{-1} - A_1^{-1}B_2^{-1} + B_1^{-1}A_2^{-1} + B_1^{-1}A_1^{-1}$$
(39)

Evaluating the coefficients  $A_n^m$  and  $B_n^m$  involved in Eq.39, we obtain after a bit of algebra, omitting again the arguments:

$$G_{12} = 2J_l J_{l+2} \left[ -\tau_1^{-1} \pi_2^{-1} - \tau_1^{-1} \tau_1^{-1} + \pi_1^{-1} \tau_2^{-1} + \pi_1^{-1} \pi_1^{-1} \right]$$
(40)  
+  $\frac{1}{6} J_l J_{l-2} \left[ 3\tau_1^1 \tau_1^1 - 3\pi_1^1 \pi_1^1 + \tau_1^1 \pi_2^1 - \pi_1^1 \tau_2^1 \right]$ 

The  $\tau$ 's and  $\pi$ 's involved in Eq.40 are then evaluated using the expressions provided in subsection 2.3, leading to:  $P_1^1 = -S$ ,  $P_2^1 = -3SC$ ,  $P_1^{-1} = S/2$ ,  $P_2^{-1} = SC/2$ ,  $\tau_1^1 = -C$ ,  $\pi_1^1 = -1$ ,  $\pi_2^1 = -3C$ ,  $\tau_2^1 = 3(S^2 - C^2)$ ,  $\tau_1^{-1} = C/2$ ,  $\pi_1^{-1} = 1/2$ ,  $\pi_2^{-1} = C/2$ ,  $\tau_2^{-1} = (C^2 - S^2)/2$ , in which we again use  $C = \cos \alpha_0$  and furthermore introduced  $S = \sin \alpha_0$ , leading eventually to:

$$Im(G_1 + G_2) = 0 (41)$$

Therefore, as a whole, we have Im(G) = 0 and:

$$C^g_{pr,z} = 0 \tag{42}$$

# 3.3 Term with real part of G for circularly symmetric Bessel beams.

For  $C_{pr,z}^s$  of Eq.26, calculations run in a similar way. We however make a decomposition of G of Eq.24 which is different from the one of Eq.28, into three terms, according to:

$$G = G^{11} + G^{12} + G^0 \tag{43}$$

in which:

$$G^{11} = i[g_{1,TM}^{-1}g_{1,TE}^{-1*} - g_{1,TM}^{1}g_{1,TE}^{1*}]$$
(44)

$$G^{12} = g_{1,TM}^{-1} g_{2,TM}^{-1*} + g_{1,TM}^{1} g_{2,TM}^{1*}]$$
(45)

$$G^{0} = \frac{1}{3}g^{0}_{1,TM}g^{0*}_{2,TM} \tag{46}$$

 $C^s_{pr,z}$  may then be decomposed into three terms according to:

$$C_{pr,z}^{s} = C_{pr,z}^{s11} + C_{pr,z}^{s12} + C_{pr,z}^{s0}$$
(47)

in which:

$$C_{pr,z}^{s11} = E_0 H_0^* \frac{\lambda^2}{6\pi} (\frac{m^2 - 1}{m^2 + 2})^2 \operatorname{Re}(G^{11}) \alpha^6$$
(48)

$$C_{pr,z}^{s12} = E_0 H_0^* \frac{\lambda^2}{6\pi} (\frac{m^2 - 1}{m^2 + 2})^2 \operatorname{Re}(G^{12}) \alpha^6$$
(49)

$$C_{pr,z}^{s0} = E_0 H_0^* \frac{\lambda^2}{6\pi} (\frac{m^2 - 1}{m^2 + 2})^2 \operatorname{Re}(G^0) \alpha^6$$
(50)

 ${\rm Re}(G^{11})$  is then evaluated in the usual manner by (i) expressing the BSCs in terms of  $A_n^m$ 's and  $B_n^m$ 's leading to:

$$\operatorname{Re}(G^{11}) = -|g(\alpha_0)|^2 \left[ (A_1^{-1})^2 - (B_1^{-1})^2 - (A_1^{1})^2 + (B_1^{1})^2 \right], \tag{51}$$

(ii) afterward expressing the  $A_n^m$ 's and  $B_n^m$ 's involved in Eq.51 according to:  $A_1^{-1} = -J_{l+2}(\tau_1^{-1} - \pi_1^{-1}), B_1^{-1} = -J_l(\tau_1^{-1} + \pi_1^{-1}), A_1^1 = \frac{1}{2}J_l(\tau_1^1 + \pi_1^1), B_1^1 = \frac{1}{2}J_{l-2}(\tau_1^1 - \pi_1^1)$  and (iii) evaluating the  $\tau$ 's and  $\pi$ 's according to:  $\tau_1^{-1} = C/2, \pi_1^{-1} = 1/2, \tau_1^1 = -C, \pi_1^1 = -1$ , to obtain  $\text{Re}(G^{11})$  and eventually:

$$C_{pr,z}^{s11} = -E_0 H_0^* \frac{\lambda^2}{24\pi} (\frac{m^2 - 1}{m^2 + 2})^2 |g(\alpha_0)|^2 \{ (C-1)^2 [(J_{l+2})^2 + (J_{l-2})^2] - 2(C+1)^2 (J_l)^2 \} \alpha^6$$
(52)

We similarly obtain:

$$C_{pr,z}^{s0} = E_0 H_0^* \frac{\lambda^2}{6\pi} (\frac{m^2 - 1}{m^2 + 2})^2 \left| g(\alpha_0) \right|^2 C(1 - C^2) [J_{l+1}^2 + J_{l-1}^2 - 2\cos(2\phi_0)J_{l+1}J_{l-1}] \alpha^6$$
(53)

and:

$$C_{pr,z}^{s12} = E_0 H_0^* \frac{\lambda^2}{6\pi} (\frac{m^2 - 1}{m^2 + 2})^2 |g(\alpha_0)|^2$$

$$[\frac{C^3}{2} (J_{l+2}^2 + J_{l-2}^2 + 2J_l^2) - \frac{3C^2 - 1}{4} (J_{l+2}^2 + J_{l-2}^2 - 2J_l^2) - C(C^2 - 1) (J_l J_{l+2} + J_l J_{l-2}) \cos(2\phi_0)] \alpha^6$$
(54)

### 4 Gradient, scattering and scattering axicon forces.

### 4.1 Gradient force.

On the one hand, we have found that  $C_{pr,z}^g$  is 0, e.g. Eq.42. On the other hand, Eq.14 immediately implies that  $\mathbf{E}.\mathbf{E}^*$  does not depend on z, i.e.  $\bigtriangledown$  $(\mathbf{E}.\mathbf{E}^*)$  is zero. This confirms our expectation that  $C_{pr,z}^g$  must be interpreted as a gradient cross-section. Let us interestingly note that  $[\varepsilon_0 \mathbf{E}.\mathbf{E}^* + \mu_0 \mathbf{H}.\mathbf{H}^*]$  and the Poynting vector  $(\mathbf{E} \times \mathbf{H}^*)$  do not depend as well on z, meaning that their z-gradient components are both 0 as well, i.e. the time-averaged energy density and the Poynting vector as well have no gradient in the z-direction.

### 4.2 Scattering and scattering axicon forces.

We now intend to examine whether  $C_{pr,z}^s$  indeed corresponds to scattering forces as expected, i.e. whether  $C_{pr,z}^s$  is proportional to the intensity of the beam that we shall state as being given by the Poynting vector. More specifically, we have to consider the component  $S_z$  of the Poynting vector taken at the location of the particle. We then start from Eqs.3-6, in order to evaluate Eq.3. This equation is to be evaluated on the axis z (where the particle is located). Then all  $\sin \theta$ -terms are cancelled and we are left with the coefficients  $C_{np}^{mq}$  only. Next, we remember that Eq.26 has been derived from Eq.24 which contains the following combinations of BSCs:  $g_{1,TM}^{-1}g_{2,TM}^{-1*}$ ,  $g_{1,TM}^{-1*}g_{1,TM}^{1*}g_{1,TM}$ ,  $g_{1,TM}^{1}g_{1,TE}^{1*}$  and  $g_{1,TM}^0g_{2,TM}^{0*}$ . Therefore, among the quantities  $C_{np}^{mq}$  of Eq.6, we only have to consider the following quantities:  $C_{12}^{-1-1}$ ,  $C_{21}^{-1-1}$ ,  $C_{11}^{-1-1}$ ,  $C_{12}^{11}$ ,  $C_{11}^{11}$ ,  $C_{12}^{00}$ and  $C_{21}^{00}$ . These quantities, restricted to the combinations of BSCs previously given, read as:

$$\widetilde{C_{12}^{-1-1}} = -g_{1,TM}^{-1}g_{2,TM}^{-1*}\psi_2\psi_1'(\tau_1^1\tau_2^1 + \pi_1^1\pi_2^1)$$
(55)

$$\widetilde{C_{21}^{-1-1}} = -g_{2,TM}^{-1}g_{1,TM}^{-1*}\psi_1\psi_2'(\tau_2^1\tau_1^1 + \pi_2^1\pi_1^1)$$
(56)

$$\widetilde{C_{11}^{-1-1}} = -2(g_{1,TM}^{-1}g_{1,TE}^{-1*}\psi_1'\psi_1' - g_{1,TE}^{-1}g_{1,TM}^{-1*}\psi_1\psi_1)\pi_1^1\tau_1^1$$
(57)

$$\widetilde{C}_{12}^{11} = -g_{1,TM}^{1}g_{2,TM}^{1*}\psi_{2}\psi_{1}^{'}(\tau_{1}^{1}\tau_{2}^{1} + \pi_{1}^{1}\pi_{2}^{1})$$
(58)

$$\widetilde{C_{21}^{11}} = -g_{2,TM}^{1}g_{1,TM}^{1*}\psi_{1}\psi_{2}^{'}(\tau_{2}^{1}\tau_{1}^{1} + \pi_{2}^{1}\pi_{1}^{1})$$
(59)

$$\widetilde{C}_{11}^{11} = -2(g_{1,TE}^{1}g_{1,TM}^{1*}\psi_{1}\psi_{1} - g_{1,TM}^{1}g_{1,TE}^{1*}\psi_{1}'\psi_{1}')\pi_{1}^{1}\tau_{1}^{1}$$
(60)

$$\widetilde{C}_{12}^{00} = -g_{1,TM}^0 g_{2,TM}^{0*} \psi_2 \psi_1' \tau_1^0 \tau_2^0 \tag{61}$$

$$\widetilde{C}_{21}^{00} = -g_{2,TM}^0 g_{1,TM}^{0*} \psi_1 \psi_2' \tau_2^0 \tau_1^0$$
(62)

in which the tilde is meant to insist on the fact that these quantities are given with the mentioned restrictions to the combinations of BSCs.

Next, we remember that the particle is located at the origin of the coordinate system, i.e. at r = 0. The quantities  $C_{np}^{mq}$  above do not occur alone in Eq.3 but as combinations of the form  $C_{np}^{mq}/r$ . We therefore have to consider the limit of these various quantities as  $r \to 0$ . We then recall Eqs.11-13 and establish:

$$\lim_{r \to 0} \frac{\psi_2(kr)\psi_1'(kr)}{r^2} = \lim_{r \to 0} \frac{\psi_1(kr)\psi_2'(kr)}{r^2} = \lim_{r \to 0} \frac{\psi_1(kr)\psi_1(kr)}{r^2} = 0$$
(63)

so that we are left with the terms  $C_{11}^{-1-1}/r^2$  and  $C_{11}^{11}/r^2$  reading as:

$$C_{11}^{-1-1} = -\frac{8k^2}{9}g_{1,TM}^{-1}g_{1,TE}^{-1*}$$
(64)

$$C_{11}^{11} = \frac{8k^2}{9} g_{1,TM}^1 g_{1,TE}^{1*} \tag{65}$$

in which we have used  $[\tau_1^1(\cos\theta)]_{\theta=0} = -1$ ,  $[\pi_1^1(\cos\theta)]_{\theta=0} = -1$  and  $\lim_{r\to 0} \psi_1'(kr)\psi_1'(kr)/r^2 = 4k^2/9$ . Inserting these results into Eq.3 and noting that  $c_1^{pw}c_1^{pw*} = 9/(4k^2)$ , we obtain a reduced expression of  $S_z$ , now denoted as  $\widetilde{S_z}$  reading as:

$$\widetilde{S_z} = E_0 H_0^* \operatorname{Re}[i(g_{1,TM}^{-1}g_{1,TE}^{-1*} - g_{1,TM}^1g_{1,TE}^{1*})]$$
(66)

which is to be compared with Eqs.44, 48 leading to:

$$C_{pr,z}^{s11} = E_0 H_0^* \frac{\lambda^2}{6\pi} (\frac{m^2 - 1}{m^2 + 2})^2 \operatorname{Re}[i(g_{1,TM}^{-1}g_{1,TE}^{-1*} - g_{1,TM}^1g_{1,TE}^{1*})]\alpha^6$$
(67)

showing that  $C_{pr,z}^{s11}$ , otherwise given by Eq.52, is proportional to  $\widetilde{S_z}$ , therefore complying with the definition of a scattering force. It then happens that  $C_{pr,z}^{s0}$  of Eq.53 and  $C_{pr,z}^{s12}$  of Eq.54 are associated with two unexpected extra-kinds of forces. The interpretation of Eq.53 is easy. It is zero when the axicon angle is zero, i.e. when C = 1. The term  $C^{s0}_{{\rm pr},z}$  may then be called an axic on pressure radiation cross-section, associated with an axicon force. It is conjectured that such forces may occur whenever the beam description exhibits an axicon angle, i.e. for Bessel beams whatever the polarization, but also for other kinds of beams including other polarizations of Bessel beams, frozen waves, Mathieu and Lommel beams, which are formed by various superpositions of Bessel beams, e.g. [54], [55], [56], [57], [58], [59], [60],

[61], [62], [63], or Laguerre-Gauss beams focused by a lens, e.g. [64], [65], [66], [67], [68].

The term  $C_{pr,z}^{s12}$  corresponds to another kind of axicon optical forces which, however, are not zero when the axicon angle is zero. However, in this case, we obtain from Eqs.52 and 54:

$$C_{pr,z}^{s11} = C_{pr,z}^{s12} = E_0 H_0^* \frac{\lambda^2}{3\pi} (\frac{m^2 - 1}{m^2 + 2})^2 |g(0)|^2 J_l^2(\sigma_0) \alpha^6$$
(68)

in which we reintroduced the argument of the Bessel function  $J_l$ , so that  $C_{pr,z}^{s12}$ , as well as  $C_{pr,z}^{s11}$ , is proportional to  $\widetilde{S}_z$ , therefore complying as well to the usual definition of a scattering force, regardless of the on-axis ( $\sigma_0 = 0$ ) or off-axis ( $\sigma_0 \neq 0$ ) location of the beam. This makes sense because, when  $\alpha_0 = 0$ , the angular spectrum decomposition of the Bessel beam only contains incident plane waves which behaves identically irrespectively of their off-axis location. Furthermore, for an on-axis location, due to the relation  $J_l(0) = \delta_{l0}$ , we have  $C_{pr,z}^{s11} = C_{pr,z}^{s12} = 0$  except for l = 0.

Next, instead of specifying first  $\alpha_0 = 0$  (no axicon angle) and afterward  $\sigma_0 = 0$  (on-axis location), we may consider an on-axis location straight away. Then, both  $C_{pr,z}^{s11}$  and  $C_{pr,z}^{s12}$  are zero (i.e. are proportional with a constant of proportionality equal to 0) for  $l \neq 0, \pm 2$ . If, furthermore  $\alpha_0 = 0$ , then, for l = 0, both  $C_{pr,z}^{s11}$  and  $C_{pr,z}^{s12}$  are equal and are given by Eq.68, i.e. they are proportional with a constant of proportional with a constant of proportionality equal to 1 and, for  $l = \pm 2$ , both  $C_{pr,z}^{s11}$  and  $C_{pr,z}^{s12}$  are found to be zero, i.e. they are proportional with a constant of proportionality equal to 0. Therefore, in all cases,  $C_{pr,z}^{s12}$  may be viewed as a genuine scattering force, whether it is zero or not, in the on-axis location, while it provides a new kind of scattering axicon forces otherwise. Let us also remember that the dichotomy between the cases ( $l \neq 0, \pm 2$ ) and ( $l = 0, \pm 2$ ) correspond as well to a dichotomy between dark and non-dark beams, i.e. between beams possessing or not a hole of intensity on the axis [69].

### 5 Conclusion.

The study of the decomposition between gradient and scattering forces exerted by laser beams on Rayleigh particles is an issue propitious to the formal understanding of the role of such forces. The case of Bessel beams is of particular interest for the understanding of such issues because such beams have no gradient of intensity along the direction of propagation, therefore implying a natural separation between gradient forces (which should be zero) and scattering forces. Furthermore, although similar studies up to now focused on on-axis configurations, the study of off-axis configurations, although much more intensive from a formal computation point of view, might provide complementary insights. This has been the case for the present study devoted to the study of longitudinal optical forces exerted by off-axis Bessel beams in the case of Rayleigh particles. The most important result of this paper is the occurrence of a new kind of optical forces which may be called scattering axicon forces associated with an axicon propagation term of the form  $\exp(ik\cos\alpha_0 z)$  in which  $\alpha_0$  is an axicon angle, which are zero if the axicon angle is zero. Future works will be devoted to associated complementary studies, such as the systematic study of on-axis configurations for Bessel beams, or to the case of transverse optical forces in off-axis Bessel configurations.

### Acknowledgments.

It is my pleasure to thank James A. Lock (Cleveland State University, USA) and J.J. Wang (Xi'dian University, China) for useful exchanges concerning this work. I furthermore heartly and enthousiastically acknowledge the effort of an anonymous reviewer who, surprised (as I have been) by the existence of axicon forces, has "taken some time to check all calculations".

### References

- A. Ashkin. Optical trapping and manipulations of neutral particles using lasers: A reprint volume with commentaries. World Scientific, Singapore, 2006.
- [2] E.M. Furst. Interactions, structure, and microscopic response : complex fluid rheology using laser tweezers. Soft Materials, 1,2:167–185, 2003.
- [3] V.A. Soifer, V.V. Kotlyar, and S.N. Khonina. Optical microparticle manipulation : Advances and new possibilities created by diffractive optics. *Physics of Particles and Nuclei*, 35,6:733–766, 2004.
- [4] K.C. Neuman and S.M. Block. Optical trapping. Review of Scientific Instruments, 75,9:2787–2809, 2004.
- [5] T.A. Nieminen, G. Knoener, N.R. Heckenberg, and H. Rubinsztein-Dunlop. Physics of optical tweezers. *Laser Manipulation of Cells and Tissues*, 82:207–236, 2007.
- [6] M. Dienerowitz, M. Mazilu, and K. Dholakia. Optical manipulation of nanoparticles: a review. *Journal of nanophotonics*, 2:Paper 021875, 32 pages, 2008.

- [7] A. Jonas and P. Zemanek. Light at work: The use of optical forces for particle manipulation, sorting and analysis. *Electrophoresis*, 29:4813–4851, 2008.
- [8] T.A. Nieminen, N. du Preez-Wilkinson, A.B. Stilgoe, V.L.Y. Loke, A.A.M. Bui, and H. Rubinsztein-Dunlop. Optical tweezers: Theory and modelling. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 146:59–80, 2014.
- [9] A.-I. Bunea and J. Glückstad. Strategies for optical trapping in biological samples: Aiming at microrobotic surgeons. *Laser photonics Rev.*, Paper 1800227:17 pages, 2019.
- [10] G. Gouesbet. Generalized Lorenz-Mie theories and mechanical effects of laser light, on the occasion of Arthur Ashkin's receipt of the 2018 Nobel prize in physics for his pioneering work in optical levitation and manipulation: A review. Journal of Quantitative Spectroscopy and Radiative Transfer, 225:258–277, 2019.
- [11] A. Ashkin and J.M. Dziedzic. Optical levitation by radiation pressure. Applied Physics Letters, 19, 8:283–285, 1971.
- [12] Y. Harada and T. Asakura. Radiation forces on a dielectric sphere in the Rayleigh scattering regime. Optics Communications, 124:529–541, 1996.
- [13] E.R. Dufresne and D.G. Grier. Optical tweezer arrays and optical substrates created with diffractive optics. *Review of scientific instruments*, 69,5:1974–1977, 1998.
- [14] P. Zemanek, A. Jonas, L. Sramek, and M. Liska. Optical trapping of Rayleigh particles using a Gaussian standing wave. *Optics Communications*, 151:273–285, 1998.
- [15] A. Resnick. Design and construction of a space-borne optical tweezer apparatus. *Review of Scientific Instruments*, 72,11:4059–4065, 2001.
- [16] A. Rohrbach and E.H.K. Stelzer. Optical trapping of dielectric particles in arbitrary fields. *Journal of the Optical Society of America A*, 18,4:839–853, 2001.
- [17] Y.K. Nahmias and D.J. Odde. Analysis of radiation forces in laser trapping and laser-guided direct writing applications. *IEEE Journal of Quantum* electronics, 38,2:131–141, 2002.
- [18] P. Zemanek, A. Jonas, and M. Liska. Simplified description of optical forces acting on a nanoparticle in the Gaussian standing wave. *Journal of* the Optical Society of America A, 19,5:1025–1034, 2002.
- [19] N. Malagnino, G. Pesce, A. Sasso, and E. Arimondo. Measurements of trapping efficiency and stiffness in optical tweezers. *Optical Communications*, 214,1-6:15–24, 2002.

- [20] C.H. Chen, P.T. Tai, and W.F. Hsieh. Bottle beam from a bare laser for single-beam trapping. *Applied Optics*, 43,32:6001–6006, 2004.
- [21] P. Zemanek, V. Karasek, and A. Sasso. Optical forces acting on a Rayleigh particle placed into interference field. *Optics Communications*, 240,4-6:401– 415, 2004.
- [22] T. Cizmar, M. Siler, and P. Zemanek. An optical nanotrap array movable over a millimetre range. Applied Physics B, 84:197–203, 2006.
- [23] M. Siler, T. Cizmar, M. Sery, and P. Zemanek. Optical forces generated by evanescent standing waves and their usage for sub-micron particle delivery. *Applied Physics B*, B84:157–165, 2006.
- [24] K.C. Neuman, T. Lionnet, and J.F. Allemand. Single-molecule micromanipulation techniques. Annual Review of Materials Research, 37:33–67, 2007.
- [25] J. Chen, J. Ng, S. Liu, and Z. Lin. Analytical calculation of axial optical force on a Rayleigh particle illuminated by Gaussian beams beyond the paraxial approximation. *Physical Review E*, 80:Paper 026607, 1–9, 2009.
- [26] M.J. Kendrick, D.H. McIntyre, and O. Ostroverkhova. Wavelength dependence of optical tweezers trapping forces on dye-doped polystyrene microspheres. *Journal of the Optical Society of America*, 26, 11:2189–2198, 2009.
- [27] R. Kumar, D.S. Mehta, and C. Shakher. Clustering of optically trapped large diameter plasmonic gold nanoparticles by laser beam of hybrid-TEM\*11 mode. *Journal of Nanophotonics*, 5:Paper 053511, 17 pages, 2011.
- [28] J.A. Lock. Calculation of the radiation trapping force for laser tweezers by use of generalized Lorenz-Mie theory. II. On-axis trapping force. *Applied Optics*, 43,12:2545–2554, 2004.
- [29] G. Gouesbet, B. Maheu, and G. Gréhan. Light scattering from a sphere arbitrarily located in a Gaussian beam, using a Bromwich formulation. *Journal of the Optical Society of America A*, 5,9:1427–1443, 1988.
- [30] G. Gouesbet, G. Gréhan, and B. Maheu. Combustion measurements, edited by N. Chigier, chapter : Generalized Lorenz-Mie theory and applications to optical sizing, pages 339–384. Hemisphere Publishing Corporation, New-York, USA, 1991.
- [31] G. Gouesbet and G. Gréhan. Generalized Lorenz-Mie theories, second edition. Springer International Publishing AG, 2017.
- [32] G. Gouesbet. Latest achievements in generalized Lorenz-Mie theories: A commented reference database. Annalen der Physik, 526, 11-12:461–489, 2014.

- [33] G. Gouesbet. T-matrix methods for electromagnetic structured beams: A commented reference database for the period 2014-2018. Journal of Quantitative Spectroscopy and Radiative Transfer, 230:247–281, 2019.
- [34] H.C. van de Hulst. Light scattering by small particles. Wiley, New York, 1957.
- [35] G. Gouesbet. T-matrix formulation and generalized Lorenz-Mie theories in spherical coordinates. *Optics Communications*, 283, 4:517–521, 2010.
- [36] G. Gouesbet. Partial wave expansions and properties of axisymmetric light beams. Applied Optics, 35,9:1543–1555, 1996.
- [37] G. Gouesbet. Poynting theorem in terms of beam shape coefficients and applications to axisymmetric, dark and non-dark, vortex and nonvortex beams. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 201:184–196, 2017.
- [38] G. Gouesbet and G. Gréhan. Sur la généralisation de la théorie de Lorenz-Mie. Journal of Optics, 13,2:97–103, 1982.
- [39] L. Robin. Fonctions sphériques de Legendre et fonctions sphéroidales. Volumes 1, 2, 3. Gauthier-Villars, Paris, 1957.
- [40] Z.X. Wang and D.R. Guo. Special functions. World Scientific, Singapore, 1989.
- [41] J. Durnin, Jr. J.J. Miceli, and J.H. Eberly. Diffraction-free beams. *Physical Review Letters*, 58, 15:1499–1501, 1987.
- [42] J. Durnin. Exact solutions for nondiffracting beams. I. The scalar theory. Journal of the Optical Society of America A, 4, 4:651–654, 1987.
- [43] J.A. Lock. Angular spectrum and localized model of Davis-type beam. Journal of the Optical Society of America A, 30, 3:489–500, 2013.
- [44] J.J. Wang, T. Wriedt, J.A. Lock, and L. Mädler. General description of circularly symmetric Bessel beams of arbitrary order. *Journal of Quantitative* Spectroscopy and Radiative Transfer, 184:218–232, 2016.
- [45] J.J. Wang, T. Wriedt, J.A. Lock, and Y.C. Jiao. General description of transverse mode Bessel beams and construction of basis Bessel fields. *Jour*nal of Quantitative Spectroscopy and Radiative Transfer, 195:8–17, 2017.
- [46] J.J. Wang, T. Wriedt, L.Mädler, Y.P. Han, and P. Hartmann. Multipole expansion of circularly Bessel beams of arbitrary order for scattering calculations. *Optics Communications*, 387:102–109, 2017.
- [47] S.R. Mishra. A vector wave analysis of a Bessel beam. Optics Communications, 85:159–161, 1991.

- [48] T. Cizmar, V. Kollarova, Z. Bouchal, and P. Zemanek. Sub-micron particle organization by self-imaging of non-diffracting beams. *New Journal of Physics*, 8 (3), 43, 2006.
- [49] J.M. Taylor and G.D. Love. Multipole expansion of Bessel and Gaussian beams for Mie scattering calculations. *Journal of the Optical Society of America A*, 26, 2:278–282, 2009.
- [50] J. Chen, J. Ng, P. Wang, and Z. Lin. Analytical partial wave expansion of vector Bessel beam and its application to optical binding. *Optics Letters*, 35, 10:1674–1676, 2010.
- [51] J. Du, C.-H. Yuen, X. Li, K. Ding, G. Du, Z. Lin, C.T. Chan, and J. Ng. Tailoring optical gradient force and optical scattering and absorption force. *Scientific Reports*, 7:DOI:10.1038/s41598-017-17874-1, 2017.
- [52] H. Zheng, X. Yu, W. Lu, J. Ng, and Z. Lin. Gcforce: Decomposition of optical force into gradient and scattering parts. *Computer Physics Communications*, 237:188–198, 2019.
- [53] N. Wang, X. Li, J. Chen, Z. Lin, and J. Ng. Gradient and scattering forces of anti-reflection-coated spheres in an aplanatic beam. *Scientific Reports*, Article number: 17423, 2018.
- [54] L.A. Ambrosio and M. Zamboni-Rached. Analytical approach of ordinary frozen waves for optical trapping and micromanipulation. *Applied Optics*, 54, 10:2584–2593, 2015.
- [55] L.A. Ambrosio and M. Zamboni-Rached. Optical forces experienced by arbitrary-sized spherical scatterers from superpositions of equal-frequency Bessel beams. *Journal of the Optical Society of America B*, 32, 5:B37–B46, 2015.
- [56] L.A. Ambrosio. Circularly symmetric frozen waves: Vector approach for light scattering calculations. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 204:112–119, 2018.
- [57] L.A. Ambrosio, M. Zamboni-Rached, and G. Gouesbet. Discrete vector frozen waves in generalized Lorenz-Mie theory: linear, azimuthal and radial polarization. *Applied Optics*, 57, 12:3293–3300, 2018.
- [58] L.A. Ambrosio, L.F.M. Votto, G. Gouesbet, and J.J. Wang. Assessing the validity of the localized approximation for discrete superposition of Bessel beams. *Journal of the Optical Society of America B*, 35, 11:2690–2698, 2018.
- [59] L.A. Ambrosio, M. Zamboni Rached, and G. Gouesbet. Zeroth-order continuous vector frozen waves for light scattering: exact multipole expansion in the generalized Lorenz-Mie theory. *Journal of the Optical Society of America B*, 36, 1:81–89, 2019.

- [60] A. Chafiq, Z. Hricha, and A. Belafhal. Paraxial approximation of Mathieu beams through an apertured ABCD optical system. *Optics Communications*, 253:223–230, 2005.
- [61] A. Chafiq and A. Belafhal. Radiation pressure cross section exerted on homogeneous dielectric spherical particle by zeroth order Mathieu beams. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 179:170–176, 2016.
- [62] A. Chafiq, L.A. Ambrosio, G. Gouesbet, and A. Belafhal. On the validity of the integral localized approximation for on-axis zeroth-order Mathieu beams. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 204:27–34, 2018.
- [63] A. Chafiq and A. Belafhal. Scattering of Lommel beams by homogeneous spherical particle in generalized Lorenz-Mie theory. *Optical and Quantum Electronics*, 2/2018, 2018.
- [64] A.S. van de Nes, S.F. Pereira, and J.J.M. Braat. On the conservation of fundamental optical quantities in non-paraxial imaging systems. *Journal* of Modern Optics, 53, 5-6:677–687, 2006.
- [65] A.S. van de Nes and P. Török. Rigorous analysis of spheres in Gauss-Laguerre beams. Optics Express, 15,20:13360–13374, 2007.
- [66] P. Török and P.R.T. Munro. The use of Gauss-Laguerre vector beams in STED microscopy. *Optics Express*, 12, 15:3605–3617, 2004.
- [67] L.A. Ambrosio and G. Gouesbet. On localized approximations for Laguerre-Gauss beams focused by a lens. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 218:100–114, 2018.
- [68] G. Gouesbet, L.A. Ambrosio, and L.F.M. Votto. Finite series expressions to evaluate the beam shape coefficients of a Laguerre-Gauss beam focused by a lens in an on-axis configuration. *Journal of Quantitative Spectroscopy* and Radiative Transfer, 242:Paper 106759, 17 pages, 2019.
- [69] G. Gouesbet and J.A. Lock. A darkness theorem for the beam shape coefficients and its relationship to higher-order non vortex Bessel beams. *Journal* of Quantitative Spectroscopy and Radiative Transfer, 201:229–235, 2017.