



HAL
open science

Discrete-time mean field games with risk-averse agents

J Frédéric Bonnans, Pierre Lavigne, Laurent Pfeiffer

► **To cite this version:**

J Frédéric Bonnans, Pierre Lavigne, Laurent Pfeiffer. Discrete-time mean field games with risk-averse agents. 2020. hal-02563949v1

HAL Id: hal-02563949

<https://hal.science/hal-02563949v1>

Preprint submitted on 5 May 2020 (v1), last revised 14 Jun 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Discrete-time mean field games with risk-averse agents*

J. Frédéric Bonnans[†] Pierre Lavigne[‡] Laurent Pfeiffer[§]

May 5, 2020

Abstract

We propose and investigate a discrete-time mean field game model involving risk-averse agents. The model under study is a coupled system of dynamic programming equations with a Kolmogorov equation. The agents' risk aversion is modeled by composite risk measures. The existence of a solution to the coupled system is obtained with a fixed point approach. The corresponding feedback control allows to construct an approximate Nash equilibrium for a related dynamic game with finitely many players.

Introduction

The class of mean field games problem was introduced by J-M. Lasry and P-L. Lions in [17, 18, 19] and M. Huang, R. Malhamé, and P. Caines in [15], to study interactions among a large population of players. Many developments and applications have been proposed this last decade, in particular in economics modeling and finance; one can refer for example to Y. Achdou and al. [1], O. Guéant, J-M. Lasry and P-L. Lions [13], and P. Cardaliaguet and C.-H. Lehalle [7]. Economic models “à la Cournot”, considering interactions between the agents via a price variable, have recently received particular attention, let us mention the works of A. Bensoussan and P.J. Graber [11], J. F. Bonnans, S. Hadikanloo, and L. Pfeiffer [6], Z. Kobeissi [16], and P. J. Graber, V. Ignazio, and A. Neufeld [12].

The specificity of the mean field game of this article is the risk aversion of the involved agents. Here risk aversion is modeled with the help of composite risk measures (also called dynamic risk measures). Mathematically, a risk measure ρ is a map that assigns to a random variable U a real number, which is usually high when U is very volatile. In this way ρ can be used to model the reluctance of a player to face highly uncertain expenses. We refer to the seminal work by P. Artzner, F. Delbaen, J-M. Eber and D. Heath in [3]. We will make use of composite risk measures, the natural extension of risk measures to a multistage framework, see for example the article of A. Shapiro and A. Ruszczyński [24]; for an application to multistage portfolio selection one can refer to A. Shapiro [26].

Let us describe more precisely our coupled system and the obtained results. The coupled system describes a population of identical agents which all optimize a discrete-time dynamical system (in a continuous state space). In the model, the associated cost function depends on a variable called belief, which is related to the behavior of the whole group, whence a coupling between a single agent and the population. Assuming that the population is very large, one can consider that an isolated representative agent has

*This work was supported by a public grant as part of the Investissement d'avenir project, reference ANR-11-LABX-0056-LMH, LabEx LMH, and by the FIME Lab (Laboratoire de Finance des Marchés de l'Energie), Paris.

[†]Inria and CMAP, Ecole Polytechnique, France. E-mail: frederic.bonnans@inria.fr

[‡]CMAP, Ecole Polytechnique, France. E-mail: pierre.lavigne@polytechnique.edu

[§]Inria and CMAP, Ecole Polytechnique, France. E-mail: laurent.pfeiffer@inria.fr

no impact on the belief. Therefore his/her behavior can be conveniently described by dynamic programming equations (in which the belief is a parameter). Mathematically, the belief is the probability distribution of the states and controls of all agents at the different time steps of the game; it is described via the Kolmogorov equation. Our first result is an existence result, obtained with a standard fixed point approach. In our second result, we show that an optimal feedback control for the mean field game yields an ε -Nash equilibrium for an N -player dynamic game, where $\varepsilon \rightarrow 0$ as $N \rightarrow \infty$. The proof of this result is based on an estimate of the expectation of the Wasserstein distance between the empirical measure of i.i.d. variables and the law of these variables, obtained by N. Fournier and A. Guillin [10, Theorem 1]. The approach that we follow was proposed by M. Huang, P. Caines, and R. Malhamé in [14].

Discrete-time and continuous-space mean field game models have been studied in different works. The framework that we propose in this article is close to the one of N. Saldi, T. Başar and M. Raginsky [25], in particular, we make use of similar weighted spaces. A few works have already investigated the issue of risk aversion. Most of them model risk sensitivity via exponential utility functions, see for example H. Tembine, Q. Zhu and T. Başar [27]. The case of robust mean field games is investigated in problem (P2) in the work of J. Moon and T. Başar [20]. In many economic situations, risk modeling is of interest, in particular in the banking industry [21]. Our approach can also be relevant in situations where mean field games are used to design telecommunication systems or smart grids; see C. Bertucci et al. [5] and C. Alasseur, I. Ben Tahar and A. Matoussi [2]. For example, in the latter reference, it could be interesting to take into account the risk of individual no-energy situations or collective black-out situations via robust control.

The article is structured as follows. In Section 1 we introduce notations, assumptions, and the system of coupled equations. In Section 2 we interpret this system as a mean field game system with risk averse agents. In Section 3 we establish general technical results that will be helpful in Section 4, where we prove the existence of a solution to the coupled system. Finally in Section 5 we investigate the connection between the coupled system and an N -player game.

1 Problem Formulation

1.1 Notations

We set $\mathcal{T} := \{0, \dots, T-1\}$ and $\bar{\mathcal{T}} := \{0, \dots, T\}$ with $T \in \mathbb{N}^*$. For any $t \in \bar{\mathcal{T}}$ and any vector (x_0, \dots, x_t) we denote

$$x_{[t]} := (x_0, \dots, x_t).$$

Functions

Let C -Lip denote the set of Lipschitz functions of modulus C on \mathbb{R}^d . We define the p -polynomially weighted space

$$\mathcal{G}_p^C := \left\{ f: \mathbb{R}^d \rightarrow \mathbb{R}^{d'}, |f(x)| \leq C(|x|^p + 1) \right\},$$

where the dimension d' depends on the context, with associated norm

$$\|f\|_{\mathcal{G},p} := \sup_{x \in \mathbb{R}^d} \frac{|f(x)|}{1 + |x|^p}.$$

Let $\mathcal{Q}_p^C \subset \mathcal{G}_p^C$ denote the set of convex mappings $f: \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying

$$-C \leq f(x) \leq C(1 + |x|^p), \quad \forall x \in \mathbb{R}^d. \quad (1.1)$$

Probability measures

Let $\mathcal{P}(\mathbb{R}^d)$ denote the set of probability measures on \mathbb{R}^d . Given $p \in [1, +\infty)$, we define the set of finite p -th order moment measures

$$\mathcal{P}_p(\mathbb{R}^d) := \left\{ m \in \mathcal{P}(\mathbb{R}^d), \int_{\mathbb{R}^d} |x|^p dm(x) < +\infty \right\},$$

that we endow with the Rubinstein-Kantorovitch distance, defined by

$$d_1(\mu, \nu) := \sup_{\phi \in 1\text{-Lip}} \int_{\mathbb{R}^d} \phi(x) d(\mu - \nu)(x),$$

for any μ and $\nu \in \mathcal{P}_1(\mathbb{R}^d)$ (see [28, Particular case 5.15] for more details). We recall that by the Hölder inequality, $\mathcal{P}_p(\mathbb{R}^d) \subseteq \mathcal{P}_1(\mathbb{R}^d)$ for any $p > 1$. Given $C > 0$, we define

$$\mathcal{P}_p^C(\mathbb{R}^d) := \left\{ m \in \mathcal{P}_p(\mathbb{R}^d), \int_{\mathbb{R}^d} |x|^p dm(x) \leq C \right\}.$$

We also consider the following sets of beliefs

$$\mathcal{B}_2 := (\mathcal{P}_2(\mathbb{R}^{2d}))^T \times \mathcal{P}_2(\mathbb{R}^d), \quad \mathcal{B}_2^C := (\mathcal{P}_2^C(\mathbb{R}^{2d}))^T \times \mathcal{P}_2^C(\mathbb{R}^d),$$

endowed with the Rubinstein-Kantorovitch distances for the product topology, also denoted d_1 .

For any m and $\nu \in \mathcal{P}(\mathbb{R}^d)$, we define the convolution product $\nu * m$ by

$$\int_{\mathbb{R}^d} h(x) d(\nu * m)(x) := \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h(y + z) d\nu(y) dm(z), \quad (1.2)$$

for any bounded Borel map $h \in \mathbb{R}^d \rightarrow \mathbb{R}$. For any $m \in \mathcal{P}(\mathbb{R}^d)$ and for any Borel map $g: \mathbb{R}^d \rightarrow \mathbb{R}^{d'}$, we define the image measure $g\#m \in \mathcal{P}(\mathbb{R}^{d'})$ by

$$\int_{\mathbb{R}^d} (h \circ g)(x) dm(x) = \int_{\mathbb{R}^d} h(y) dg\#m(y), \quad (1.3)$$

for any bounded Borel map $h \in \mathbb{R}^d \rightarrow \mathbb{R}^{d'}$.

1.2 Coupled system

Let us first introduce the data of the problem. We consider

- a congestion function $F: \bar{\mathcal{T}} \times \mathbb{R}^d \times \mathcal{B}_2 \rightarrow \mathbb{R}$
- a price function $P: \mathcal{T} \times \mathcal{B}_2 \rightarrow \mathbb{R}^d$
- an initial distribution $\bar{m} \in \mathcal{P}_2(\mathbb{R}^d)$
- individual noise distributions $(\nu(t))_{t \in \mathcal{T}} \in (\mathcal{P}_2(\mathbb{R}^d))^T$.

The running cost $\ell: \mathcal{T} \times \mathbb{R}^d \times \mathbb{R}^d \times \mathcal{B}_2 \rightarrow \mathbb{R}$ is defined by

$$\ell(t, x, a, b) = \frac{1}{2}|a|^2 + \langle a, P(t, b) \rangle + F(t, x, b).$$

For modeling risk aversion, we consider a family of subsets $(\mathcal{Z}_t)_{t \in \mathcal{T}}$ such that

$$\mathcal{Z}_t \subseteq \left\{ Z \in L^\infty(\mathbb{R}^d), \int_{\mathbb{R}^d} Z(y) d\nu(t, y) = 1, Z \geq 0 \right\}, \quad \forall t \in \mathcal{T}.$$

For any $t \in \mathcal{T}$, we define

$$\mathcal{M}_t := \left\{ \xi \in \mathcal{P}(\mathbb{R}^d), d\xi = Z d\nu(t), Z \in \mathcal{Z}_t \right\}. \quad (1.4)$$

For any $t \in \mathcal{T}$, \mathcal{Z}_t is assumed to be nonempty and convex, thus \mathcal{M}_t is a nonempty and convex subset of $\mathcal{P}(\mathbb{R}^d)$.

We propose to study a *risk averse mean field game* (MFG), taking the form of the following coupled system:

$$\left\{ \begin{array}{l} \text{(i)} \quad \begin{cases} u(t, x) = \inf_{a \in \mathbb{R}^d} \left(\ell(t, x, a, b) + \sup_{\xi \in \mathcal{M}_t} \int_{\mathbb{R}^d} u(t+1, x+a+y) d\xi(y) \right), \\ u(T, x) = F(T, x, b), \end{cases} \\ \text{(ii)} \quad \alpha_t(x) = \arg \min_{a \in \mathbb{R}^d} \left(\ell(t, x, a, b) + \sup_{\xi \in \mathcal{M}_t} \int_{\mathbb{R}^d} u(t+1, x+a+y) d\xi(y) \right), \\ \text{(iii)} \quad \begin{cases} m(t+1) = \nu(t) * [(id + \alpha_t) \# m(t)], \\ m(0) = \bar{m}, \end{cases} \\ \text{(iv)} \quad \mu(t) = (id, \alpha_t) \# m(t), \\ \text{(v)} \quad b := (\mu(0), \dots, \mu(T-1), m(T)), \end{array} \right. \quad (\text{MFG})$$

for any $(t, x) \in \mathcal{T} \times \mathbb{R}^d$. The five unknowns in the above system are

- the value function $u \in (\mathcal{G}_2)^{T+1}$
- the feedback control $\alpha \in (\mathcal{G}_1 \cap 1\text{-Lip})^T$
- the distribution of states $m \in (\mathcal{P}_2)^{T+1}$
- the joint distribution of states and controls $\mu \in (\mathcal{P}_2(\mathbb{R}^{2d}))^T$
- the belief $b \in \mathcal{B}_2$.

Let us describe briefly the coupled system; we will justify it more in detail in Section 2. Equation (MFG,i) is a dynamic programming equation associated with a discrete-time optimal control problem for a representative agent. The belief b appears as a parameter of the equation, since a single agent has no impact on it. The corresponding optimal feedback control α is then given by (MFG,ii). Now, assuming that all agents make use of the feedback control α , the distribution of their state m is described by the Kolmogorov equation (MFG,iii) with initial condition \bar{m} .

Our approach for proving the existence of a solution consists in formulating the system (MFG) as a fixed point equation. For this purpose, we consider two mappings. The first one, that we call dynamic programming mapping, assigns to a belief b the solutions $u^*(b)$ and $\alpha^*(b)$ to equations (MFG,i) and (MFG,ii), respectively. The second one, the Kolmogorov mapping, assigns to a feedback control α the triplet $(m^*(\alpha), \mu^*(\alpha), b^*(\alpha))$, where $m^*(\alpha)$, $\mu^*(\alpha)$, and $b^*(\alpha)$ are the solutions to (MFG,iii), (MFG,iv), and (MFG,v), respectively. These two mappings will be investigated in Section 4. They allow to reformulate the system (MFG) as an equivalent fixed point equation

$$b = b^* \circ \alpha^*(b).$$

1.3 Assumptions

We state now the assumptions on the data of the problem, in force all along the article. Note that for the results of Section 5 (dealing with the N -player dynamic game), we will need a slightly stronger assumption on the mapping F .

We make use of the same constant C to formulate the different assumptions. In the sequel, the constant C denotes a generic constant depending only on those involved in the assumptions and T ; its value can change from an inequality to the next one.

Assumption 1.1. *There exists $C > 0$ such that $\bar{m} \in \mathcal{P}_2^C(\mathbb{R}^d)$ and such that for any $t \in \mathcal{T}$, $\nu(t) \in \mathcal{P}_2^C(\mathbb{R}^d)$.*

Assumption 1.2. *There exists $C > 0$ such that for any $t \in \mathcal{T}$ and for any $Z \in \mathcal{Z}_t$,*

$$\|Z\|_\infty \leq C,$$

and there exists $Z' \in \mathcal{Z}_t$ such that

$$Z' \geq \frac{1}{C} \quad \text{a.e.}$$

Remark 1.3. *Assumption 1.2 implies the existence of $C > 0$ such that*

$$\mathcal{M}_t \subseteq \mathcal{P}_2^C(\mathbb{R}^d), \quad \forall t \in \mathcal{T}. \quad (1.5)$$

The results obtained in Section 4 only require (1.5) to hold. The full Assumption 1.2 will be used in Section 5.

Assumption 1.4. *There exists $C > 0$ such that for any $t \in \mathcal{T}$ and for any b_1 and $b_2 \in \mathcal{B}_2$,*

- (i) $F(t, \cdot, b_1) \in \mathcal{Q}_2^C$,
- (ii) $\|F(t, \cdot, b_1) - F(t, \cdot, b_2)\|_{\mathcal{G},2} \leq C d_1(b_1, b_2)$,
- (iii) $|P(t, b_1) - P(t, b_2)| \leq C d_1(b_1, b_2)$,
- (iv) $|P(t, b_1)| \leq C$.

Remark 1.5. *In economics or in finance, prices typically depend on the aggregated demand or supply. One could consider for example*

$$P(t, b) := \psi \left(t, \int_{\mathbb{R}^{2d}} \alpha d\mu(t, x, \alpha) \right),$$

where $\psi: \mathcal{T} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$. In this case, if ψ is a C -Lipschitz mapping then for any b_1 and $b_2 \in \mathcal{B}_2$, one has that

$$|P(t, b_1) - P(t, b_2)| \leq C \left| \int_{\mathbb{R}^{2d}} \alpha d(\mu_1 - \mu_2)(t, x, \alpha) \right| \leq C d_1(\mu_1, \mu_2) \leq C d_1(b_1, b_2),$$

which implies Assumption 1.4 (iii). Assumption 1.4 (iv) also holds if $|\psi| \leq C$.

2 Interpretation of the coupled system

In Subsection 2.1 we describe the risk averse optimal control problem associated with (MFG,i-ii). In Subsection 2.2 we justify the Kolmogorov equation (MFG,iii).

2.1 Dynamic programming equation

Risk measures

Let X_0 and $(Y_t)_{t \in \mathcal{T}}$ be $(T + 1)$ -independent random variables defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $\mathcal{L}(X_0) = \bar{m}$ and $\mathcal{L}(Y_t) = \nu(t)$. We define the filtration $(\mathcal{F}_t)_{t \in \mathcal{T}}$, where $\mathcal{F}_0 := \sigma(X_0)$ is the sigma-algebra generated by X_0 , and $\mathcal{F}_{t+1} := \sigma(X_0, Y_{[t]})$. We denote for any $t \in \bar{\mathcal{T}}$ and any $p \in [1, +\infty)$

$$\mathbb{L}_t^p(\Omega, \mathbb{R}^{d'}) := L^p(\Omega, \mathcal{F}_t, \mathbb{P}, \mathbb{R}^{d'}),$$

the space of \mathcal{F}_t measurable random variables with finite p -th order moment and value in $\mathbb{R}^{d'}$. When the dimension is $d' = 1$, we simplify the notation: $\mathbb{L}_t^p := \mathbb{L}_t^p(\Omega, \mathbb{R})$.

Definition 2.1. *Given $t \in \mathcal{T}$, we say that a mapping $\rho_t: \mathbb{L}_{t+1}^1 \rightarrow \mathbb{L}_t^1$ is a conditional risk mapping if it satisfies the following conditions:*

- **(M) Monotonicity:** *For any U and $U' \in \mathbb{L}_{t+1}^1$ such that $U \leq U'$, we have*

$$\rho_t(U) \leq \rho_t(U'), \quad \text{a.s.}$$

- **(C) Convexity:** *For any U and $U' \in \mathbb{L}_{t+1}^1$, for any $\alpha \in [0, 1]$, we have*

$$\rho_t(\alpha U + (1 - \alpha)U') \leq \alpha \rho_t(U) + (1 - \alpha)\rho_t(U'), \quad \text{a.s.}$$

- **(TI) Translation Invariance:** *For any $U \in \mathbb{L}_{t+1}^1$ and for any $V \in \mathbb{L}_t^1$, we have*

$$\rho_t(U + V) = \rho_t(U) + V, \quad \text{a.s.}$$

- **(PH) Positive Homogeneity:** *For any $\alpha \geq 0$, for any $U \in \mathbb{L}_{t+1}^1$, we have*

$$\rho_t(\alpha U) = \alpha \rho_t(U), \quad \text{a.s.}$$

Quoting [23], the conditional risk mapping $\rho_t(U_{t+1})$ can be interpreted as a fair one-time \mathcal{F}_t -measurable charge we would be willing to incur at time t instead of the random future cost U_{t+1} .

We fix now a family of conditional risk mappings $(\rho_t)_{t \in \mathcal{T}}$, $\rho_t: \mathbb{L}_{t+1}^1 \rightarrow \mathbb{L}_t^1$, defined by

$$\rho_t(U_{t+1})(x_0, y_{[t-1]}) = \sup_{Z \in \mathcal{Z}_t} \int_{\Omega} U_{t+1}(x_0, y_{[t-1]}, Y_t(\omega)) Z(Y_t(\omega)) d\mathbb{P}(\omega), \quad (2.1)$$

where the random variables U_{t+1} and $\rho_t(U_{t+1})$ are explicitly represented as measurable functions of $(x_0, y_{[t]}) \in \mathbb{R}^{(t+2)d}$ and $(x_0, y_{[t-1]}) \in \mathbb{R}^{(t+1)d}$, respectively. Recalling the definition of \mathcal{M}_t (1.4), we have

$$\rho_t(U_{t+1})(x_0, y_{[t-1]}) = \sup_{\xi \in \mathcal{M}_t} \int_{\mathbb{R}^d} U_{t+1}(x_0, y_{[t-1]}, y_t) d\xi(y_t).$$

We set

$$\mathcal{Q}_{t+1} := \{Q = Z(Y_t) \text{ a.s., } Z \in \mathcal{Z}_t\}$$

so that ρ_t can be expressed in the following form:

$$\rho_t(U_{t+1}) = \sup_{Q_{t+1} \in \mathcal{Q}_{t+1}} \mathbb{E}[U_{t+1} Q_{t+1} | \mathcal{F}_t].$$

Remark 2.2. We could allow Z in (2.1) to depend on $(x_0, y_{[t-1]})$. Then \mathcal{M}_t would also depend on $(x_0, y_{[t-1]})$. This approach would be close to the one developed in [24]. However, in the dynamic programming principle, the state should include x_0 and $y_{[t-1]}$ and therefore its dimension might be too large for practical computations.

Finally we construct the associated composite risk measure $\rho: \mathbb{L}_T^1 \rightarrow \mathbb{R}$,

$$\rho(U) := \mathbb{E}[\rho_0 \circ \cdots \circ \rho_{T-1}(U)],$$

which also satisfies **(M)**, **(C)**, **(TI)**, and **(PH)**.

Remark 2.3. Given a probability space $(\Omega', \mathcal{F}', \mathbb{P}')$ and given $\alpha \in (0, 1]$, the conditional value at risk (also called expected shortfall or average value at risk) of a random variable $U \in L^1(\Omega', \mathcal{F}', \mathbb{P}')$ is defined by

$$\text{CV@R}_\alpha(U) := \inf_{W \in L^1(\Omega', \mathcal{F}', \mathbb{P}')} W + \alpha^{-1} \mathbb{E}[(U - W)_+],$$

where $x_+ = \max\{0, x\}$ denotes the positive part of any $x \in \mathbb{R}$. It has the following dual representation (see [9, Lemma 4.51 and Theorem 4.52]):

$$\text{CV@R}_\alpha(U) = \sup \left\{ \mathbb{E}[UZ] \mid Z \in L^\infty(\Omega', \mathcal{F}', \mathbb{P}'), Z \in [0, \alpha^{-1}] \text{ a.s.}, \mathbb{E}[Z] = 1 \right\}.$$

Therefore, a natural extension of the conditional value at risk to the framework of the article is given by

$$\rho_t(U_{t+1}) = \sup_{Z \in \mathcal{Z}_t} \mathbb{E}[U_{t+1}Z(Y_t) | \mathcal{F}_t],$$

where

$$\mathcal{Z}_t := \left\{ Z \in L^\infty(\mathbb{R}^d) \mid Z \in [0, \alpha^{-1}] \text{ a.e.}, \int_{\mathbb{R}^d} Z(y) d\nu(t, y) = 1 \right\}.$$

This particular definition of \mathcal{Z}_t satisfies Assumption 1.2. We refer to [9, Definition 11.8] and [8, Subsection 2.3.1] for extensions of the conditional value at risk to general filtrations in a discrete-time setting.

Control problem

We consider the following set of controls for any $t \in \mathcal{T}$,

$$\mathcal{A}_t = \mathbb{L}_t^2(\Omega, \mathbb{R}^d), \quad \mathcal{A} := \mathcal{A}_0 \times \cdots \times \mathcal{A}_{T-1}.$$

Given a control $A \in \mathcal{A}$, the evolution of the state of the representative player is given by

$$X_{t+1} = X_t + A_t + Y_t, \quad \forall t \in \mathcal{T}. \quad (\text{C})$$

The initial condition is the random variable X_0 fixed previously. In the notation, we do not make explicit the dependence of $(X_t)_{t \in \mathcal{T}}$ with respect to A , which is always clear from the context. Note that by induction, $X_t \in \mathbb{L}_t^2(\Omega, \mathbb{R}^d)$ for any $t \in \mathcal{T}$.

For a given belief $b \in \mathcal{B}_2$, the risk averse multistage cost of the representative agent is given by

$$\mathcal{J}(A, b) := \rho \left(\sum_{t=0}^{T-1} \ell(t, X_t, A_t, b) + F(T, X_T, b) \right). \quad (2.2)$$

The corresponding problem is

$$\inf_{A \in \mathcal{A}} \mathcal{J}(A, b). \quad (\text{P})$$

As a consequence of the translation invariance property **(TI)**, the problem (P) can be expressed in a nested form

$$\mathbb{E} \left[\inf_{A_0 \in \mathcal{A}_0} \ell(0, X_0, A_0, b) + \inf_{A_1 \in \mathcal{A}_1} \rho_0 \left(\ell(1, X_1, A_1, b) + \dots \right. \right. \\ \left. \left. + \inf_{A_{T-1} \in \mathcal{A}_{T-1}} \rho_{T-2} \left(\ell(T-1, X_{T-1}, A_{T-1}, b) + \rho_{T-1} \left(F(T, X_T, b) \right) \right) \dots \right) \right]. \quad (2.3)$$

The solution of problem (P) can be characterized by a dynamic programming approach, that we briefly describe. We refer to [23] for a rigorous presentation. The dynamic programming approach consists, in short, in investigating a family of control problems of same nature as (P) . The problems are parameterized by their initial time t and initial condition x ; their value is denoted $u(t, x)$. The function u is characterized by equation **(MFG,i)**, which itself derives from the nested form (2.3). Consider now the function α , solution to **(MFG,ii)** (we will justify the existence and uniqueness of the “argmin” in this equation in Lemma 4.1). Let $(\bar{X}_t)_{t \in \bar{\mathcal{T}}}$ be the solution to the closed-loop system

$$\bar{X}_{t+1} = \bar{X}_t + \alpha_t(\bar{X}_t) + Y_t \quad (2.4)$$

and let \bar{A} be defined by

$$\bar{A}_t = \alpha_t(\bar{X}_t). \quad (2.5)$$

Let us briefly justify that $\bar{A} \in \mathcal{A}$. Since \bar{X}_t is adapted to \mathcal{F}_t , we also have that \bar{A}_t is \mathcal{F}_t -measurable. In addition we show in Lemma 4.2 that α_t is 1-Lipschitz. Since the random variables X_0 and $(Y_t)_{t \in \mathcal{T}}$ have a bounded second-order moment, \bar{A}_t also have a bounded second-order moment and thus, $\bar{A} \in \mathcal{A}$.

The following proposition states the optimality of the control \bar{A} .

Proposition 2.4. *We have*

$$\inf_{A \in \mathcal{A}} \mathcal{J}(A, b) = \mathbb{E}[u(0, X_0)] = \int_{\mathbb{R}^d} u(0, x) dm(0, x), \quad (2.6)$$

where u solves the dynamic programming equation **(MFG,i)**. Moreover, the control \bar{A} (defined by (2.4)-(2.5)) is the unique solution to Problem (P) .

2.2 Kolmogorov equation

Lemma 2.5. *Let $\alpha: \mathcal{T} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a continuous vector field. Suppose that the state equation **(C)** is of the feedback form*

$$X_{t+1} = X_t + \alpha_t(X_t) + Y_t.$$

Then for any $t \in \bar{\mathcal{T}}$, $m(t) = \mathcal{L}(X_t) \in \mathcal{P}(\mathbb{R}^d)$ is characterized by the Kolmogorov equation **(MFG,iv)**.

Proof. Let ϕ be a bounded Borel test function. For any $t \in \mathcal{T}$, by independence of X_t and Y_t we have

$$\mathbb{E}[\phi(X_{t+1})] = \mathbb{E}[\phi(X_t + \alpha_t(X_t) + Y_t)] \\ = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \phi(x + \alpha_t(x) + y) dm(t, x) d\nu(t, y).$$

By definition of the push-forward (1.3) we obtain

$$\int_{\mathbb{R}^d} \phi(x + \alpha_t(x) + y) dm(t, x) = \int_{\mathbb{R}^d} \phi(z + y) d(id + \alpha_t)\#m(t, z).$$

By definition of convolution (1.2) we have

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \phi(z+y) d\nu(t,y) d(id + \alpha_t) \# m(t,z) = \int_{\mathbb{R}^d} \phi(x) d(\nu(t) * [(id + \alpha_t) \# m(t)])(x),$$

as was to be proved. \square

3 Technical lemmas

This section contains independent technical lemmas. The reader only interested in the main results of the article can skip it.

Lemma 3.1. *Let $p \in [1, +\infty)$ and let $C > 0$. For any m_1 and m_2 in $\mathcal{P}_p^C(\mathbb{R}^d)$, the probability measure $m_1 * m_2$ lies in $\mathcal{P}_p^{2^p C}(\mathbb{R}^d)$. In addition, given $m_0 \in \mathcal{P}_p^C(\mathbb{R}^d)$, the mapping $\mathcal{P}_p^C(\mathbb{R}^d) \ni m \mapsto m_0 * m$ is non-expansive for the distance d_1 .*

Proof. Let m_1 and m_2 in $\mathcal{P}_p^C(\mathbb{R}^d)$. We have

$$\begin{aligned} \int_{\mathbb{R}^d} |x|^p d(m_1 * m_2)(x) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |y+z|^p dm_1(y) dm_2(z) \\ &\leq \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} 2^{p-1} (|y|^p + |z|^p) dm_1(y) dm_2(z) \leq 2^p C. \end{aligned}$$

Thus $m_1 * m_2 \in \mathcal{P}_p^{2^p C}(\mathbb{R}^d)$. Moreover, given $m_0 \in \mathcal{P}_p^C(\mathbb{R}^d)$, we have

$$\begin{aligned} d_1(m_0 * m_1, m_0 * m_2) &= \sup_{\phi \in 1\text{-Lip}} \int_{\mathbb{R}^d} \phi(x) d(m_0 * m_1 - m_0 * m_2)(x) \\ &= \sup_{\phi \in 1\text{-Lip}} \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \phi(y+z) dm_0(y) \right) d(m_1 - m_2)(z). \end{aligned}$$

Since the mapping $z \mapsto \int_{\mathbb{R}^d} \phi(y+z) d\nu(y)$ is non-expansive, we further obtain that

$$d_1(m_0 * m_1, m_0 * m_2) \leq d_1(m_1, m_2),$$

which concludes the proof. \square

Lemma 3.2. *Let $p \in [1, +\infty)$ and let $C > 0$. For any $m \in \mathcal{P}_p^C(\mathbb{R}^d)$ and for any Borel map $g \in \mathcal{G}_1^C$, the probability measure $g \# m$ lies in $\mathcal{P}_p^q(\mathbb{R}^d)$, with $q = 2^{p-1} C^p (1+C)$. In addition, the inequality*

$$d_1(g_1 \# m_1, g_2 \# m_2) \leq (1+C) \|g_1 - g_2\|_{\mathcal{G},1} + C d_1(m_1, m_2) \quad (3.1)$$

holds for any m_1 and m_2 in $\mathcal{P}_p^C(\mathbb{R}^d)$ and for any Borel maps g_1 and g_2 in $\mathcal{G}_1^C \cap C\text{-Lip}$.

Proof. Let $m \in \mathcal{P}_p^C(\mathbb{R}^d)$ and let $g \in \mathcal{G}_1^C$ be a Borel map. We have

$$\int_{\mathbb{R}^d} |x|^p d(g \# m)(x) = \int_{\mathbb{R}^d} |g(x)|^p dm(x) \leq \|g\|_{\mathcal{G},1}^p \int_{\mathbb{R}^d} (1+|x|)^p dm(x) \leq q.$$

Consider (g_1, m_1) and (g_2, m_2) in $\mathcal{G}_1^C \times \mathcal{P}_p^C(\mathbb{R}^d)$. We have

$$\begin{aligned} d_1(g_1 \# m_1, g_2 \# m_2) &= \sup_{\phi \in 1\text{-Lip}} \int_{\mathbb{R}^d} \phi(x) d(g_1 \# m_1 - g_2 \# m_2)(x) \\ &= \sup_{\phi \in 1\text{-Lip}} \int_{\mathbb{R}^d} (\phi \circ g_1(x) - \phi \circ g_2(x)) dm_2(x) + \int_{\mathbb{R}^d} \phi \circ g_1(x) d(m_1 - m_2)(x) \\ &\leq \|g_1 - g_2\|_{\mathcal{G},1} \int_{\mathbb{R}^d} (1+|x|) dm_2(x) + \sup_{\phi \in 1\text{-Lip}} \int_{\mathbb{R}^d} \phi \circ g_1(x) d(m_1 - m_2)(x) \\ &\leq (1+C) \|g_1 - g_2\|_{\mathcal{G},1} + C \sup_{\phi \in 1\text{-Lip}} \int_{\mathbb{R}^d} C^{-1} \phi \circ g_1(x) d(m_1 - m_2)(x). \end{aligned}$$

Observing that $C^{-1}\phi \circ g_1 \in 1\text{-Lip}$, we deduce inequality (3.1). \square

Given a convex function $u: \mathbb{R}^d \rightarrow \mathbb{R}$, we define the Moreau envelope V_u and the proximal operator prox_u of u as follows:

$$V_u(x) := \min_{y \in \mathbb{R}^d} \frac{1}{2}|x - y|^2 + u(y), \quad \text{prox}_u(x) := \arg \min_{y \in \mathbb{R}^d} \frac{1}{2}|x - y|^2 + u(y). \quad (3.2)$$

In the proofs, we will occasionally consider the map $g_u: \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, defined by

$$g_u(x, y) := \frac{1}{2}|x - y|^2 + u(y).$$

Proposition 3.3. *Let $u: \mathbb{R}^d \rightarrow \mathbb{R}$ be a convex function. Then prox_u and $(\text{id} - \text{prox}_u)$ are non-expansive.*

Proof. Direct consequence of [4, Proposition 12.27]. \square

Lemma 3.4. *Let $R > 0$ and let $u \in \mathcal{Q}_2^R$ (the set was defined in (1.1)). Then $|\text{prox}_u|^2 \in \mathcal{G}_2^{C_1(R)}$ and $|\text{prox}_u| \in \mathcal{G}_1^{(C_1(R))^{1/2}}$, where $C_1(R) := 8R + 2$.*

Proof. Let $u \in \mathcal{Q}_2^R$. By Proposition 3.3, the map prox_u is non-expansive. Thus

$$|\text{prox}_u(x)| \leq |\text{prox}_u(0)| + |x|. \quad (3.3)$$

In addition, from the definition of the proximal operator (3.2), we have

$$\frac{1}{2}|\text{prox}_u(0)|^2 + u(\text{prox}_u(0)) \leq u(0).$$

Since $u \in \mathcal{Q}_2^R$, we deduce that $|\text{prox}_u(0)|^2 \leq 4R$. We further obtain with (3.3) that

$$|\text{prox}_u(x)|^2 \leq 2(|x|^2 + |\text{prox}_u(0)|^2) \leq C_1(R)(1 + |x|^2), \quad (3.4)$$

as was to be proved. Taking the square root of (3.4), we infer that $|\text{prox}_u| \in \mathcal{G}_1^{C_1(R)^{1/2}}$. \square

Lemma 3.5. *Let $R > 0$ and let $u \in \mathcal{Q}_2^R$. Then $V_u \in \mathcal{Q}_2^{C_2(R)}$, where*

$$C_2(R) := (R + 1)(1 + C_1(R)).$$

Proof. Let $u \in \mathcal{Q}_2^R$. Clearly V_u is convex as the infimum with respect to $y \in \mathbb{R}^d$ of the jointly convex map $(x, y) \mapsto g_u(x, y)$. For any $x \in \mathbb{R}^d$, we have

$$V_u(x) = \frac{1}{2}|x - \text{prox}_u(x)|^2 + u(\text{prox}_u(x)),$$

by definition of V_u and prox_u . Since $u \in \mathcal{Q}_2^R$, we further obtain that

$$-R \leq V_u(x) \leq |x|^2 + |\text{prox}_u(x)|^2 + R(1 + |\text{prox}_u(x)|^2).$$

Applying Lemma 3.4, we finally obtain that $V_u \in \mathcal{Q}_2^{C_2(R)}$. \square

Lemma 3.6. *Let $R > 0$. For any u and v in \mathcal{Q}_2^R , the inequality*

$$\|\text{prox}_u - \text{prox}_v\|_{\mathcal{G},1} \leq C_3(R)\|u - v\|_{\mathcal{G},2}^{1/2} \quad (3.5)$$

holds, where $C_3(R) := \sqrt{2(1 + C_1(R))}$.

Proof. Let u and v in \mathcal{Q}_2^R . Observing that g_u and g_v are 1-strongly convex with respect to their second argument, we have

$$\begin{aligned}\frac{1}{2}|\text{prox}_u(x) - \text{prox}_v(x)|^2 &\leq g_u(x, \text{prox}_v(x)) - g_u(x, \text{prox}_u(x)), \\ \frac{1}{2}|\text{prox}_u(x) - \text{prox}_v(x)|^2 &\leq g_v(x, \text{prox}_u(x)) - g_v(x, \text{prox}_v(x)).\end{aligned}$$

Summing up the two inequalities, we obtain that

$$\begin{aligned}|\text{prox}_u(x) - \text{prox}_v(x)|^2 &\leq v(\text{prox}_u(x)) - u(\text{prox}_u(x)) + u(\text{prox}_v(x)) - v(\text{prox}_v(x)) \\ &\leq (2 + |\text{prox}_u(x)|^2 + |\text{prox}_v(x)|^2)\|u - v\|_{\mathcal{G},2}.\end{aligned}\quad (3.6)$$

By Lemma 3.4,

$$2 + |\text{prox}_u(x)|^2 + |\text{prox}_v(x)|^2 \leq 2 + 2C_1(R)(1 + |x|^2) \leq C_3(R)^2(1 + |x|^2).\quad (3.7)$$

Combining (3.6) and (3.7) and taking the square root, we obtain (3.5). \square

Lemma 3.7. *Let $R > 0$. For any u and v in \mathcal{Q}_2^R , we have*

$$\|V_u - V_v\|_{\mathcal{G},2} \leq C_4(R)\|u - v\|_{\mathcal{G},2},\quad (3.8)$$

where $C_4(R) := 1 + C_1(R)$.

Proof. Let u and v in \mathcal{Q}_2^R . Recalling the definitions of g_u and g_v , we have

$$V_u(x) - V_v(x) \leq g_u(x, \text{prox}_v(x)) - g_v(x, \text{prox}_v(x)) = u(\text{prox}_v(x)) - v(\text{prox}_v(x)).$$

Lemma 3.4 yields

$$V_u(x) - V_v(x) \leq (1 + \|\text{prox}_v(x)\|_{\mathcal{G},1}^2)\|u - v\|_{\mathcal{G},2} \leq (1 + C_1(R)(1 + |x|^2))\|u - v\|_{\mathcal{G},2}.$$

Exchanging u and v , we deduce (3.8). \square

Lemma 3.8. *Let $R > 0$. For any $u \in \mathcal{Q}_2^R$ and for any $(x, y) \in \mathbb{R}^d \times \mathbb{R}^d$,*

$$|V_u(x) - V_u(y)| \leq C_5(R)(1 + |x| + |y|)|x - y|,\quad (3.9)$$

where $C_5(R) := 1 + \sqrt{C_1(R)}$.

Proof. Let $u \in \mathcal{Q}_2^R$. We have

$$\begin{aligned}V_u(x) - V_u(y) &\leq g_u(x, \text{prox}_u(y)) - g_u(y, \text{prox}_u(y)) \\ &= \frac{1}{2}|x - \text{prox}_u(y)|^2 - \frac{1}{2}|y - \text{prox}_u(y)|^2 \\ &\leq \frac{1}{2}|x + y - 2\text{prox}_u(y)| \cdot |x - y|.\end{aligned}$$

We further obtain with Lemma 3.4 that

$$|x + y - 2\text{prox}_u(y)| \leq |x| + |y| + 2\sqrt{C_1(R)}(1 + |y|) \leq 2(1 + \sqrt{C_1(R)})(1 + |x| + |y|).$$

Combining the two obtained inequalities and exchanging x and y , we obtain (3.9). \square

Lemma 3.9. *Let $R > 0$ and let \mathcal{M} be a subset of $\mathcal{P}_2^R(\mathbb{R}^d)$. Given $u \in \mathcal{Q}_2^R$, consider the mapping $\Upsilon[u](x)$ defined for any $x \in \mathbb{R}^d$ by*

$$\Upsilon[u](x) := \sup_{\xi \in \mathcal{M}} \int_{\mathbb{R}^d} u(x+y) d\xi(y).$$

Then $\Upsilon[u] \in \mathcal{Q}_2^{C_6(R)}$, where $C_6(R) = 2R(1+R)$. Moreover, the map $\mathcal{Q}_2^R \ni u \mapsto \Upsilon[u]$ is Lipschitz continuous with modulus $2(1+R)$.

Proof. Let $u \in \mathcal{Q}_2^R$. For any $\xi \in \mathcal{M}$, the map $\mathbb{R}^d \ni x \mapsto \int_{\mathbb{R}^d} u(x+y) d\xi(y)$ is convex, as can be easily verified. Thus $\Upsilon[u](x)$ is convex with respect to x , as a supremum of convex maps. Moreover, for any $x \in \mathbb{R}^d$, we have

$$-R \leq \Upsilon[u](x) \leq \sup_{\xi \in \mathcal{M}} \int_{\mathbb{R}^d} 2R(1+|x|^2+|y|^2) d\xi(y) \leq 2R(1+|x|^2+R).$$

This proves that $\Upsilon[u] \in \mathcal{Q}_2^{C_6(R)}$. Consider now $v \in \mathcal{Q}_2^R$. We have

$$\begin{aligned} |\Upsilon[u](x) - \Upsilon[v](x)| &\leq \sup_{\xi \in \mathcal{M}} \left| \int_{\mathbb{R}^d} (u(x+y) - v(x+y)) d\xi(y) \right| \\ &\leq \|u - v\|_{\mathcal{G},2} \left(\sup_{\xi \in \mathcal{M}} \int_{\mathbb{R}^d} (1+|x+y|^2) d\xi(y) \right). \end{aligned} \quad (3.10)$$

For any $\xi \in \mathcal{M}$, we further have

$$\int_{\mathbb{R}^d} (1+|x+y|^2) d\xi(y) \leq 1 + 2|x|^2 + 2R \leq 2(1+R)(1+|x|^2). \quad (3.11)$$

Combining (3.10) and (3.11), we deduce that

$$\|\Upsilon[u] - \Upsilon[v]\|_{\mathcal{G},2} \leq 2(1+R)\|u - v\|_{\mathcal{G},2},$$

as was to be proved. \square

4 Existence result

In this section we prove the main existence result. We first investigate the continuity of the dynamic programming mapping and the continuity of the Kolmogorov mapping introduced in Subsection 1.2.

4.1 Dynamic Programming mapping

Given a belief $b \in \mathcal{B}_2$, let us recall that $u^*(\cdot, \cdot, b)$ is the solution to (MFG,i) and $\alpha^*(\cdot, b)$ the solution to (MFG,ii). For conveniency, we introduce an intermediate mapping $\bar{u}^*(\cdot, \cdot, b)$, defined by

$$\bar{u}^*(t+1, x, b) := \sup_{\xi \in \mathcal{M}_t} \int_{\mathbb{R}^d} u^*(t+1, x+y, b) d\xi(y),$$

for any $t \in \mathcal{T}$. This allows to rewrite equations (MFG,i-ii) as follows:

$$u^*(t, x, b) = \inf_{a \in \mathbb{R}^d} \frac{1}{2}|a|^2 + \langle a, P(t, b) \rangle + F(t, x, b) + \bar{u}^*(t+1, x+a, b), \quad (4.1)$$

$$\alpha_t^*(x, b) = \operatorname{argmin}_{a \in \mathbb{R}^d} \frac{1}{2}|a|^2 + \langle a, P(t, b) \rangle + F(t, x, b) + \bar{u}^*(t+1, x+a, b). \quad (4.2)$$

The first step consists in rewriting these equations in a functional form, with the help of the Moreau envelope and the proximal operator (introduced in (3.2)). In addition, we justify the existence and uniqueness of the minimizer in the right-hand side of (4.2).

Lemma 4.1. *Let $b \in \mathcal{B}_2$. For any $t \in \mathcal{T}$, the map $u^*(t, \cdot, b)$ is convex and for any $x \in \mathbb{R}^d$, the minimizer in (MFG,ii) is unique. Moreover,*

$$u^*(t, x, b) = V_{\bar{u}^*(t+1, \cdot, b)}(x - P(t, b)) + F(t, x, b) - \frac{1}{2}|P(t, b)|^2, \quad (4.3)$$

$$\alpha_t^*(x, b) = (\text{prox}_{\bar{u}^*(t+1, \cdot, b)} - \text{id})(x - P(t, b)) - P(t, b). \quad (4.4)$$

Proof. We proceed by backward induction. The terminal condition $u^*(T, \cdot, b) = F(T, \cdot, b)$ and Assumption 1.4 (i) yield the convexity of $u^*(T, \cdot, b)$. Let $t \in \mathcal{T}$. Suppose that $u^*(t+1, \cdot, b)$ is convex. Then by Lemma 3.9, $\bar{u}^*(t+1, \cdot, b)$ is also convex. By the change of variable

$$y = x + a,$$

the dynamic programming equation (MFG,i) can be written as follows:

$$\begin{aligned} u^*(t, x, b) &= \inf_{y \in \mathbb{R}^d} \left(\frac{1}{2}|y - x|^2 + \langle (y - x), P(t, b) \rangle + F(t, x, b) + \bar{u}^*(t+1, y, b) \right) \\ &= \inf_{y \in \mathbb{R}^d} \left(\frac{1}{2}|y - (x - P(t, b))|^2 + \bar{u}^*(t+1, y, b) \right) + F(t, x, b) - \frac{1}{2}|P(t, b)|^2. \end{aligned} \quad (4.5)$$

This proves (4.3). Moreover, $u^*(t, \cdot, b)$ is convex, as a consequence of Assumption 1.4 (i) and Lemma 3.5. Besides the unique minimizer in (4.5) is

$$y^* := \text{prox}_{\bar{u}^*(t+1, \cdot, b)}(x - P(t, b))$$

and therefore, the unique minimizer in (MFG,ii) is $y^* - x$, which proves (4.4). The lemma is proved. \square

Lemma 4.2. *Let $b \in \mathcal{B}_2$. For any $t \in \bar{\mathcal{T}}$, we have*

$$u^*(t, \cdot, b) \in \mathcal{Q}_2^{C_u}, \quad (4.6)$$

and for any $t \in \mathcal{T}$, we have

$$\alpha_t^*(\cdot, b) \in \mathcal{G}_1^{C_\alpha} \cap 1\text{-Lip}, \quad (4.7)$$

for some positive constants C_α and C_u independent of t and b .

Proof. In the proof, all constants C are independent of b . Let us prove (4.6) by backward induction. The terminal condition $u^*(T, \cdot, b) = F(T, \cdot, b)$ and Assumption 1.4 (i) imply that $u^*(T, \cdot, b) \in \mathcal{Q}_2^C$, for some constant $C > 0$ (independent of b). Let $t \in \mathcal{T}$. Suppose that $u^*(t+1, \cdot, b) \in \mathcal{Q}_2^C$. Then by Lemma 3.9 and relation 1.5, we have $\bar{u}^*(t, \cdot, b) \in \mathcal{Q}_2^C$. Recall that by Lemma 4.1, we have

$$u^*(t, \cdot, b) = V_{\bar{u}^*(t+1, \cdot, b)}(\cdot - P(t, b)) + F(t, \cdot, b) - \frac{1}{2}|P(t, b)|^2. \quad (4.8)$$

By Assumptions 1.4 (i) and (iv), $F(t, \cdot, b) - \frac{1}{2}|P(t, b)|^2 \in \mathcal{Q}_2^C$. Using again Assumption 1.4 (iv) and Lemma 3.5, we obtain that $V_{\bar{u}^*(t+1, \cdot, b)}(\cdot - P(t, b)) \in \mathcal{Q}_2^C$. Therefore, the right-hand side of (4.8) lies in \mathcal{Q}_2^C and finally, $u^*(t, \cdot, b) \in \mathcal{Q}_2^C$, where C is independent of b .

Let us prove (4.7). By Lemma 4.1, we have

$$\alpha_t^*(\cdot, b) = (\text{prox}_{\bar{u}^*(t+1, \cdot, b)} - \text{id})(\cdot - P(t, b)) - P(t, b). \quad (4.9)$$

We already know that $\bar{u}^*(t+1, \cdot, b) \in \mathcal{Q}_2^C$. Moreover, by Assumption 1.4 (iv), $P(t, b)$ is bounded. Therefore, by Lemma 3.4, $\text{prox}_{\bar{u}^*(t+1, \cdot, b)}(\cdot - P(t, b)) \in \mathcal{G}_1^C$. Then it is easy to show that $\alpha_t^*(\cdot, b) \in \mathcal{G}_1^C$, where again, C does not depend on b . Finally, $\alpha^*(t, \cdot, b)$ is non-expansive as a consequence of (4.9) and Proposition 3.3. The lemma is proved. \square

Lemma 4.3. *There exists $C > 0$ such that for any $(t, b_1, b_2) \in \mathcal{T} \times \mathcal{B}_2 \times \mathcal{B}_2$,*

$$\|u^*(t, \cdot, b_1) - u^*(t, \cdot, b_2)\|_{\mathcal{G},2} \leq Cd_1(b_1, b_2), \quad (4.10)$$

$$\|\bar{u}^*(t, \cdot, b_1) - \bar{u}^*(t, \cdot, b_2)\|_{\mathcal{G},2} \leq Cd_1(b_1, b_2). \quad (4.11)$$

Proof. In the proof, all constants C are independent of b_1 and b_2 . We proceed by backward induction. By Assumption 1.4 (iii) and by the terminal condition $u^*(T, \cdot, b) = F(T, \cdot, b)$, inequality (4.10) holds true for $t = T$. Let $t \in \mathcal{T}$. Suppose that

$$\|u^*(t+1, \cdot, b_1) - u^*(t+1, \cdot, b_2)\|_{\mathcal{G},2} \leq Cd_1(b_1, b_2),$$

for some positive constant $C > 0$ independent of b_1 and b_2 . By Assumption 1.3 and Lemma 3.9, we deduce that

$$\|\bar{u}^*(t+1, \cdot, b_1) - \bar{u}^*(t+1, \cdot, b_2)\|_{\mathcal{G},2} \leq Cd_1(b_1, b_2). \quad (4.12)$$

By Lemma 4.1, we have

$$u^*(t, x, b_1) - u^*(t, x, b_2) = a_1(t, x, b_1, b_2) + a_2(t, x, b_1, b_2) + a_3(t, x, b_1, b_2), \quad (4.13)$$

where

$$\begin{aligned} a_1(t, x, b_1, b_2) &:= V_{\bar{u}^*(t+1, \cdot, b_1)}(x - P(t, b_1)) - V_{\bar{u}^*(t+1, \cdot, b_2)}(x - P(t, b_1)), \\ a_2(t, x, b_1, b_2) &:= V_{\bar{u}^*(t+1, \cdot, b_2)}(x - P(t, b_1)) - V_{\bar{u}^*(t+1, \cdot, b_2)}(x - P(t, b_2)), \\ a_3(t, x, b_1, b_2) &:= F(t, x, b_1) - F(t, x, b_2) + \frac{1}{2}(|P(t, b_2)|^2 - |P(t, b_1)|^2). \end{aligned}$$

It remains to bound $a_1(t, \cdot, b_1, b_2)$, $a_2(t, \cdot, b_1, b_2)$, and $a_3(t, \cdot, b_1, b_2)$ in \mathcal{G}_2^C . We deduce from Lemma 3.7, Assumption 1.4 (iv), and estimate (4.12), that

$$\begin{aligned} |a_1(t, x, b_1, b_2)| &\leq \|V_{\bar{u}^*(t+1, \cdot, b_1)} - V_{\bar{u}^*(t+1, \cdot, b_2)}\|_{\mathcal{G},2}(1 + |x - P(t, b_1)|^2) \\ &\leq C\|\bar{u}^*(t+1, \cdot, b_1) - \bar{u}^*(t+1, \cdot, b_2)\|_{\mathcal{G},2}(1 + |x|^2) \\ &\leq Cd_1(b_1, b_2)(1 + |x|^2). \end{aligned}$$

Then by Lemma 3.8 and Assumption 1.4 (iv), we have

$$\begin{aligned} |a_2(t, x, b_1, b_2)| &\leq C(1 + |x - P(t, b_1)| + |x - P(t, b_2)|)|P(t, b_2) - P(t, b_1)| \\ &\leq C(1 + |x|)d_1(b_1, b_2) \\ &\leq C(1 + |x|^2)d_1(b_1, b_2). \end{aligned}$$

Finally by Assumption 1.4 (ii-iv), we have

$$\begin{aligned} |a_3(t, x, b_1, b_2)| &\leq \|F(t, \cdot, b_1) - F(t, \cdot, b_2)\|_{\mathcal{G},2}(1 + |x|^2) + C|P(t, b_1) - P(t, b_2)| \\ &\leq C(1 + |x|^2)d_1(b_1, b_2). \end{aligned}$$

Then combining (4.13) and the three estimates of a_1 , a_2 , and a_3 , we obtain that

$$\|u^*(t, \cdot, b_1) - u^*(t, \cdot, b_2)\|_{\mathcal{G},2} \leq Cd_1(b_1, b_2),$$

which concludes the proof. \square

Lemma 4.4. *There exists $C > 0$ such that for any $(t, b_1, b_2) \in \mathcal{T} \times \mathcal{B}_2 \times \mathcal{B}_2$,*

$$\|\alpha_t^*(\cdot, b_1) - \alpha_t^*(\cdot, b_2)\|_{\mathcal{G},1} \leq C \left(d_1(b_1, b_2)^{1/2} + d_1(b_1, b_2) \right). \quad (4.14)$$

Proof. Let $(t, b_1, b_2) \in \mathcal{T} \times \mathcal{B}_2 \times \mathcal{B}_2$. By Lemma 4.1, we have

$$\alpha_t^*(x, b_1) - \alpha_t^*(x, b_2) = a_4(t, x, b_1, b_2) + a_5(t, x, b_1, b_2), \quad (4.15)$$

where

$$\begin{aligned} a_4(t, x, b_1, b_2) &= \text{prox}_{\bar{u}^*(t+1, \cdot, b_1)}(x - P(t, b_1)) - \text{prox}_{\bar{u}^*(t+1, \cdot, b_2)}(x - P(t, b_1)), \\ a_5(t, x, b_1, b_2) &= \text{prox}_{\bar{u}^*(t+1, \cdot, b_2)}(x - P(t, b_1)) - \text{prox}_{\bar{u}^*(t+1, \cdot, b_2)}(x - P(t, b_2)). \end{aligned}$$

Using successively Lemma 3.6, Assumption 1.4 (iv), and estimate (4.11), we obtain

$$\begin{aligned} |a_4(t, x, b_1, b_2)| &\leq \|\text{prox}_{\bar{u}^*(t+1, \cdot, b_1)} - \text{prox}_{\bar{u}^*(t+1, \cdot, b_2)}\|_{\mathcal{G},1} (1 + |x - P(t, b_1)|) \\ &\leq C \|\bar{u}^*(t+1, \cdot, b_1) - \bar{u}^*(t+1, \cdot, b_2)\|^{1/2} (1 + |x|) \\ &\leq C \|d_1(b_1, b_2)\|^{1/2} (1 + |x|). \end{aligned}$$

Moreover, since $\text{prox}_{\bar{u}^*(t+1, \cdot, b_2)}$ is non-expansive, we have with Assumption 1.4 (iii) that

$$|a_5(t, x, b_1, b_2)| \leq |(x - P(t, b_1)) - (x - P(t, b_2))| \leq d_1(b_1, b_2).$$

Combining the two obtained estimates of a_4 and a_5 with (4.15), we obtain (4.14). \square

4.2 Kolmogorov mapping

We study now the Kolmogorov mapping

$$(\mathcal{G}_1^{C_\alpha} \cap 1\text{-Lip})^T \ni \alpha \mapsto (m^*, \mu^*, b^*)(\alpha),$$

where (m^*, μ^*, b^*) is the solution to (MFG,iii-v).

Lemma 4.5. *There exists $C_b > 0$ such that for any $\alpha \in (\mathcal{G}_1^{C_\alpha} \cap 1\text{-Lip})^T$,*

$$m^*(\alpha) \in (\mathcal{P}_2^{C_b}(\mathbb{R}^d))^{T+1}, \quad \mu^*(\alpha) \in (\mathcal{P}_2^{C_b}(\mathbb{R}^{2d}))^T, \quad \text{and} \quad b^*(\alpha) \in \mathcal{B}_2^{C_b}.$$

In addition the three mappings m^, μ^* and b^* are continuous.*

Proof. Let $\alpha \in (\mathcal{G}_1^{C_\alpha} \cap 1\text{-Lip})^T$. All constants C in the proof are independent of α . Let us first prove by induction that for any $t \in \bar{\mathcal{T}}$, there exists a constant $C > 0$ independent of α such that $m^*(t, \cdot, \alpha) \in \mathcal{P}_2^C(\mathbb{R}^d)$ and such that, $m^*(t, \cdot, \alpha)$ is continuous with respect to α . The claim is clear for $t = 0$, since $m^*(0, \cdot, \alpha) = \bar{m} \in \mathcal{P}_2^C(\mathbb{R}^d)$, by Assumption 1.1. Now, let us assume that the claim holds true for some $t \in \mathcal{T}$. We recall that

$$m^*(t+1, \cdot, \alpha) = \nu(t) * [(id + \alpha_t) \# m^*(t, \cdot, \alpha)].$$

Since $\nu(t) \in \mathcal{P}_2^C(\mathbb{R}^d)$ (by Assumption 1.1) and since $\alpha_t \in \mathcal{G}_1^{C_\alpha} \cap 1\text{-Lip}$, we obtain with Lemma 3.1 and Lemma 3.2 that $m^*(t+1, \cdot, \alpha) \in \mathcal{P}_2^C(\mathbb{R}^d)$ and that $m^*(t+1, \cdot, \alpha)$ is a continuous function of α , by composition.

It remains to justify the boundedness of μ^* and b^* . We recall that for any $t \in \mathcal{T}$,

$$\mu^*(t, \cdot, \alpha) = (id, \alpha_t) \# m^*(t, \cdot, \alpha).$$

We deduce from Lemma 3.2 that $\mu^*(t, \cdot, \alpha) \in \mathcal{P}_2^C(\mathbb{R}^{2d})$ and that $\mu^*(t, \cdot, \alpha)$ is a continuous function of α , by composition. It immediately follows that $b^*(\alpha) \in \mathcal{B}_2^C$ and that b is continuous. \square

4.3 Existence of equilibrium

We are ready to prove the existence of a solution of system (MFG). The proof relies on the Schauder fixed point theorem, that we first recall.

Theorem 4.6. (Schauder) *Let C be a convex and compact set in a Banach space X , and let $T: C \rightarrow C$ be a continuous mapping. Then T has a fixed point, i.e. there exists $x \in C$ such that*

$$T(x) = x.$$

Theorem 4.7. *There exists $(u, \alpha, m, \mu, b) \in (\mathcal{G}_2^{C_u})^T \times (\mathcal{G}_1^{C_\alpha} \cap 1\text{-Lip})^T \times (\mathcal{P}_2^{C_b}(\mathbb{R}^d))^{T+1} \times (\mathcal{P}_2^{C_b}(\mathbb{R}^{2d}))^T \times \mathcal{B}_2^{C_b}$ solution to system (MFG), where C_u, C_α and C_b are the constants obtained in Lemma 4.2 and Lemma 4.5.*

Proof. By Lemma 4.4 and Lemma 4.5, the mapping

$$\mathcal{B}_2^{C_b} \ni b \mapsto b^* \circ \alpha^*(b) \in \mathcal{B}_2^{C_b}$$

is continuous for the distance d_1 . Moreover, $\mathcal{B}_2^{C_b}$ is compact for d_1 , see [22, Lemma 25]. Therefore, by the Schauder fixed point theorem, there exists $\bar{b} \in \mathcal{B}_2^{C_b}$ such that $\bar{b} = b^* \circ \alpha^*(\bar{b})$. Let us set $\bar{u} = u^*(\bar{b})$, $\bar{\alpha} = \alpha^*(\bar{b})$, $\bar{m} = m^*(\bar{\alpha})$, and $\bar{\mu} = \mu^*(\bar{\alpha})$. Then $(\bar{u}, \bar{\alpha}, \bar{m}, \bar{\mu}, \bar{b})$ is solution to (MFG) and lies in the announced set. \square

5 Connection with a finite player game

In this section we establish a connection between the coupled system (MFG) and a dynamic game with N players. More precisely, we fix a solution $(\bar{u}, \bar{\alpha}, \bar{m}, \bar{\mu}, \bar{b})$ of system (MFG) and consider the situation where each of the N players adopts the feedback $\bar{\alpha}$. We show that this situation is an ε -Nash equilibrium for the N -player game and we quantify the rate of convergence of ε to 0 as N goes to infinity.

To show this, the following restriction on Assumption 1.4 (ii) will be required, in particular to prove Lemma 5.12.

Assumption 5.1. *There exists $C > 0$ such that for any $t \in \mathcal{T}$ and for any b_1 and b_2 in \mathcal{B}_2 ,*

- (i) $F(t, \cdot, b_1) \in \mathcal{Q}_1^C$,
- (ii) $\|F(t, \cdot, b_1) - F(t, \cdot, b_2)\|_{\mathcal{G},1} \leq C d_1(b_1, b_2)$.

We have already fixed a solution to system (MFG), now we also fix the number of players N ; all constants C appearing in the sequel are independent of N .

5.1 Formulation of the game

Let $\mathcal{N} := \{1, \dots, N\}$ and let $i \in \mathcal{N}$. For any vector (x^1, \dots, x^N) we denote

$$\begin{aligned} \mathbf{x} &= (x^1, \dots, x^N), \\ \mathbf{x}^{-i} &= (x^1, \dots, x^{i-1}, x^{i+1}, \dots, x^N). \end{aligned}$$

Consider a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $(X_0^i)_{i \in \mathcal{N}}$ be i.i.d. random variables with law $\mathcal{L}(X_0^i) = \bar{m}$. Let $(Y_t^i)_{i \in \mathcal{N}, t \in \mathcal{T}}$ be independent random variables, independent of $(X_0^i)_{i \in \mathcal{N}}$, with law $\mathcal{L}(Y_t^i) = \nu(t)$. We denote $\nu(t) := \bigotimes_{i=1}^N \nu(t)$. We define the filtration $(\mathcal{F}_t)_{t \in \bar{\mathcal{T}}}$ as follows: $\mathcal{F}_0 := \sigma(\mathbf{X}_0)$ is the sigma-algebra generated by \mathbf{X}_0 , $\mathcal{F}_{t+1} := \sigma(\mathbf{X}_0, \mathbf{Y}_{[t]})$. In this section we denote

$$L_t^p(\Omega, \mathbb{R}^{d'}) := L^p(\Omega, \mathcal{F}_t, \mathbb{P}, \mathbb{R}^{d'}),$$

the space of \mathcal{F}_t measurable random variables with finite p -th order moment and value in $\mathbb{R}^{d'}$. When the dimension is $d' = 1$, we simplify the notation $\mathbf{L}_t^p = \mathbf{L}_t^p(\Omega, \mathbb{R})$. For any $t \in \mathcal{T}$, we consider the control set

$$\mathcal{A}_t := \mathbf{L}_t^2(\Omega, \mathbb{R}^d), \quad \mathcal{A} := \mathcal{A}_0 \times \cdots \times \mathcal{A}_{T-1}, \quad \mathcal{A}^N := \prod_{i=1}^N \mathcal{A}.$$

For any $t \in \mathcal{T}$ and for any constant $C > 0$ we denote \mathcal{A}_t^C the set of controls $A \in \mathcal{A}_t$ such that

$$\int_{\Omega} |A(\omega)|^2 d\mathbb{P}(\omega) \leq C$$

and we set $\mathcal{A}^C := \mathcal{A}_0^C \times \cdots \times \mathcal{A}_{T-1}^C$. The control of player $i \in \mathcal{N}$ is an adapted stochastic process $A^i \in \mathcal{A}$, whose associated trajectory $(X_t^i[A^i])_{t \in \bar{\mathcal{T}}}$ is defined by the following state equation

$$X_{t+1}^i = X_t^i + A_t^i + Y_t^i.$$

Remark 5.2. Let $R > 0$. There exists $C > 0$ (depending on R) such that for any $i \in \mathcal{N}$ and for any $A^i \in \mathcal{A}^R$, $\mathbb{E}[|X_t^i[A^i]|^2] \leq C$ for any $t \in \bar{\mathcal{T}}$, since $\mathcal{L}(X_0^i) \in \mathcal{P}_2(\mathbb{R}^d)$ and $\mathcal{L}(Y_t^i) \in \mathcal{P}_2(\mathbb{R}^d)$.

Given $\mathbf{A} \in \mathcal{A}^N$, we define the random empirical measure of the positions and the random empirical joint measure of the positions and actions of players by

$$m_{\mathbf{A}}^N(t) := \frac{1}{N} \sum_{i \in \mathcal{N}} \delta_{X_t^i[A^i]}, \quad \mu_{\mathbf{A}}^N(t) := \frac{1}{N} \sum_{i \in \mathcal{N}} \delta_{(X_t^i[A^i], A_t^i)},$$

where δ denotes the Dirac measure. We set

$$b_{\mathbf{A}}^N := (\mu_{\mathbf{A}}^N(0), \dots, \mu_{\mathbf{A}}^N(T-1), m_{\mathbf{A}}^N(T)).$$

For any $i \in \mathcal{N}$ and for any $t \in \mathcal{T}$, we define the *individual conditional risk measure* $\rho_t^i: \mathbf{L}_{t+1}^1 \rightarrow \mathbf{L}_t^1$,

$$\rho_t^i(U_{t+1})(\mathbf{x}_0, \mathbf{y}_{[t-1]}) = \sup_{Z \in \mathcal{Z}_t} \int_{\Omega} U_{t+1}(\mathbf{x}_0, \mathbf{y}_{[t-1]}, \mathbf{Y}_t(\omega)) Z(Y_t^i(\omega)) d\mathbb{P}(\omega).$$

We define the set

$$\mathbf{Q}_{t+1}^i := \{Q \in \mathbf{L}_{t+1}^\infty, Q = Z(Y_t^i) \text{ a.s.}, Z \in \mathcal{Z}_t\}.$$

Then ρ_t^i can be expressed in the following form:

$$\rho_t^i(U_{t+1}) = \sup_{Q_{t+1} \in \mathbf{Q}_{t+1}^i} \mathbb{E}[U_{t+1} Q_{t+1} | \mathcal{F}_t]. \quad (5.1)$$

In addition we have that

$$\rho_t^i(U_{t+1})(\mathbf{x}_0, \mathbf{y}_{[t-1]}) = \sup_{\xi \in \mathcal{M}_t} \int_{\mathbb{R}^d} \int_{\mathbb{R}^{Nd}} U_{t+1}(\mathbf{x}_0, \mathbf{y}_{[t]}) d\nu^{-i}(t, \mathbf{y}_t^{-i}) d\xi(y_t^i),$$

where $\nu^{-i}(t) := \bigotimes_{j \in \mathcal{N} \setminus \{i\}} \nu^j(t)$. Then $(\rho_t^i)_{t \in \mathcal{T}}$ is a family of conditional risk mappings. We define the associated individual composite risk measure $\rho^i: \mathbf{L}_T^1 \rightarrow \mathbb{R}$,

$$\rho^i(U) := \mathbb{E}[\rho_0^i \circ \cdots \circ \rho_{T-1}^i(U)].$$

Here players are risk averse with respect to their individual noise only. For any $\mathbf{A} \in \mathcal{A}^N$ the cost of the player $i \in \mathcal{N}$ is given by

$$\mathcal{J}^{i,N}(A^i, \mathbf{A}^{-i}) := \rho^i \left(\sum_{t=0}^{T-1} \ell(t, X_t^i[A^i], A_t^i, b_{\mathbf{A}}^N) + F(T, X_T^i[A^i], b_{\mathbf{A}}^N) \right).$$

Definition 5.3. Let $\varepsilon \geq 0$. We say that an N -uplet $\widehat{\mathbf{A}} \in \mathcal{A}^N$ is an ε -Nash equilibrium for the N -player game if for any $i \in \mathcal{N}$,

$$\mathcal{J}^{i,N}(\widehat{A}^i, \widehat{\mathbf{A}}^{-i}) \leq \inf_{A^i \in \mathcal{A}} \mathcal{J}^{i,N}(A^i, \widehat{\mathbf{A}}^{-i}) + \varepsilon. \quad (5.2)$$

For $\varepsilon = 0$, we recover the usual definition of a Nash equilibrium.

5.2 An approximate Nash equilibrium

For any player $i \in \mathcal{N}$, we denote by $(\bar{X}_t^i)_{t \in \mathcal{T}}$ the solution to the closed-loop system

$$X_{t+1}^i = X_t^i + \bar{\alpha}_t(X_t^i) + Y_t^i.$$

We define the control $\bar{A}^i \in \mathcal{A}$ by

$$\bar{A}_t^i = \bar{\alpha}_t(\bar{X}_t^i). \quad (5.3)$$

Since \bar{X}_t^i is adapted to \mathcal{F}_t , the control \bar{A}_t^i is also \mathcal{F}_t -measurable. Moreover, $\bar{\alpha}_t$ is 1-Lipschitz and the random variables X_0 and $(Y_t)_{t \in \mathcal{T}}$ have a bounded second-order moment, thus $\bar{A}^i \in \mathcal{A}$. In addition, by Proposition 2.4, \bar{A}^i minimizes the following cost \mathcal{J}^i :

$$\mathcal{J}^i(A^i, \bar{b}) := \rho^i \left(\sum_{t=0}^{T-1} \ell(t, X_t^i[A^i], A_t^i, \bar{b}) + F(T, X_T^i[A^i], \bar{b}) \right). \quad (5.4)$$

Finally we set $\bar{\mathbf{A}} = (\bar{A}^1, \dots, \bar{A}^N)$. The following result states that $\bar{\mathbf{A}}$ is an ε -Nash equilibrium.

Theorem 5.4. Let $\xi \in (0, 1/2)$. There exists a constant $C > 0$, independent of N , such that the N -uplet $\bar{\mathbf{A}}$ defined above is an ε -Nash equilibrium with

$$\varepsilon = CN^{-\tau(d)/2}, \quad \tau(d) = \begin{cases} 1/2 - \xi & \text{if } d \in \{1, 2\}, \\ 1/d & \text{if } d \geq 3. \end{cases}$$

In addition we have that

$$|\mathcal{J}^{i,N}(\bar{\mathbf{A}}) - \mathcal{J}^i(\bar{A}^i, \bar{b})| \leq CN^{-\tau(d)/2}, \quad (5.5)$$

for any $i \in \mathcal{N}$.

The proof of the theorem can be found at the end of Subsection 5.3 (page 23), which contains technical intermediate lemmas. They rely on the following result.

Theorem 5.5. (Fournier-Guillin) Let $c > 0$, let $\xi \in (0, 1/2)$, and let $\mu \in \mathcal{P}_2^c(\mathbb{R}^d)$. Consider N i.i.d. random variables $(X_i)_{i \in \{1, \dots, N\}}$ in \mathbb{R}^d with law μ and denote by μ_N their empirical measure, defined by

$$\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{X_i}. \quad (5.6)$$

There exists a constant $C > 0$ depending only on c , d , and ξ such that

$$\mathbb{E}[d_1(\mu, \mu_N)] \leq CN^{-\tau(d)}.$$

Proof. The theorem is a direct application of [10, Theorem 1] with $q = \frac{2}{1+2\xi}$ if $d \in \{1, 2\}$ and $q = 2$ if $d \geq 3$. \square

5.3 Proof of Theorem 5.4

We begin with four technical lemmas dealing with the regularity of the individual risk measures ρ^i .

Lemma 5.6. *For any player $i \in \mathcal{N}$ the risk measure ρ^i is subadditive, that is*

$$\rho^i(U + V) \leq \rho^i(U) + \rho^i(V),$$

for any U and V in \mathbf{L}_T^1 .

Proof. Let us define $\pi_T^i(U) = U$ and $\pi_t^i(U) = \rho_t^i \circ \rho_{t+1}^i \dots \circ \rho_{T-1}^i(U)$, for any $U \in \mathbf{L}_T^1$. Note that $\pi_t^i = \rho_t^i \circ \pi_{t+1}^i$, for any $t \in \mathcal{T}$. We prove by backward induction that π_t^i is subadditive for any $t \in \overline{\mathcal{T}}$. The claim is trivial for $t = T$. Let $t \in \mathcal{T}$. Assume that π_{t+1}^i is subadditive, let us prove that π_t^i is subadditive. First we observe that for any U and V in \mathbf{L}_T^1 ,

$$\begin{aligned} \rho_t^i(U + V) &= \sup_{Q \in \mathcal{Q}_{t+1}^i} \mathbb{E}[(U + V)Q | \mathcal{F}_t] \\ &\leq \sup_{Q \in \mathcal{Q}_{t+1}^i} \mathbb{E}[UQ | \mathcal{F}_t] + \sup_{Q \in \mathcal{Q}_{t+1}^i} \mathbb{E}[VQ | \mathcal{F}_t] = \rho_t^i(U) + \rho_t^i(V), \quad \text{a.s.} \end{aligned}$$

It follows with the monotonicity of ρ_t^i that

$$\begin{aligned} \pi_t^i(U + V) &= \rho_t^i \circ \pi_{t+1}^i(U + V) \\ &\leq \rho_t^i(\pi_{t+1}^i(U) + \pi_{t+1}^i(V)) \\ &\leq \rho_t^i \circ \pi_{t+1}^i(U) + \rho_t^i \circ \pi_{t+1}^i(V) = \pi_t^i(U) + \pi_t^i(V), \quad \text{a.s.} \end{aligned}$$

Recalling that $\rho^i(U) = \mathbb{E}[\rho_0^i \circ \dots \circ \rho_{T-1}^i(U)] = \mathbb{E}[\pi_0(U)]$, we conclude that ρ^i is also subadditive. \square

The following result is close to a triangle inequality for risk measures. The difference with the triangle inequality is due to the positive homogeneity of risk measures, while norms are absolutely homogeneous.

Lemma 5.7. *For any $i \in \mathcal{N}$ and for any U and V in \mathbf{L}_T^1 , we have*

$$|\rho^i(U + V) - \rho^i(U)| \leq \rho^i(|V|). \quad (5.7)$$

Proof. By the subadditivity and by the monotonicity of ρ^i , we have

$$\rho^i(U + V) - \rho^i(U) \leq (\rho^i(U) + \rho^i(V)) - \rho^i(U) = \rho^i(V) \leq \rho^i(|V|).$$

Similarly, we have

$$\rho^i(U) - \rho^i(U + V) = \rho^i(U + V - V) - \rho^i(U + V) \leq \rho^i(-V) \leq \rho^i(|V|).$$

Inequality (5.7) follows. \square

Lemma 5.8. *There exists $C > 0$ such that for any $(i, t) \in \mathcal{N} \times \mathcal{T}$ and for any $U \in \mathbf{L}_T^1$,*

$$\frac{1}{C} \mathbb{E}[\|U\|] \leq \rho^i(U) \leq C \mathbb{E}[\|U\|]. \quad (5.8)$$

Proof. All constants C in the proof are independent of U . Recall the definition of π_t^i , introduced in the proof of Lemma 5.6. We prove by backward induction that for any $t \in \bar{\mathcal{T}}$, there exists $C > 0$ such that for any $U \in \mathbf{L}_T^1$,

$$\frac{1}{C} \mathbb{E} [|U| | \mathcal{F}_t] \leq \pi_t^i(U) \leq C \mathbb{E} [|U| | \mathcal{F}_t], \quad \text{a.s.}$$

The claim is trivial for $t = T$. Let $t \in \mathcal{T}$. Assume that the claim holds true for $t + 1$. We first observe that for any $U \in \mathbf{L}_{t+1}^1$,

$$\frac{1}{C} \mathbb{E} [|U| | \mathcal{F}_t] \leq \rho_t^i(U) \leq C \mathbb{E} [|U| | \mathcal{F}_t], \quad \text{a.s.}, \quad (5.9)$$

as a direct consequence of Assumption 1.2. It follows with the monotonicity of ρ_t^i that

$$\begin{aligned} \pi_t^i(U) &= \rho_t^i \circ \pi_{t+1}^i(U) \\ &\leq \rho_t^i(C \mathbb{E} [|U| | \mathcal{F}_{t+1}]) \\ &\leq C \mathbb{E} [C \mathbb{E} [|U| | \mathcal{F}_{t+1}] | \mathcal{F}_t] \leq C \mathbb{E} [|U| | \mathcal{F}_t], \quad \text{a.s.} \end{aligned}$$

Similarly we prove that $\pi_t^i(|U|) \geq \frac{1}{C} \mathbb{E} [|U| | \mathcal{F}_t]$ a.s. Recalling that $\rho^i(U) = \mathbb{E} [\pi_0(U)]$, we finally obtain (5.8). \square

The following lemma is an estimate of the second-order moment of suboptimal controls (for problem (5.2)).

Lemma 5.9. *There exists $C > 0$ such that for any $i \in \mathcal{N}$, if \hat{A}^i satisfies*

$$\mathcal{J}^{i,N}(\hat{A}^i, \bar{\mathbf{A}}^{-i}) \leq \inf_{A^i \in \mathcal{A}} \mathcal{J}^{i,N}(A^i, \bar{\mathbf{A}}^{-i}) + 1, \quad (5.10)$$

then $\hat{A}^i \in \mathcal{A}^C$.

Proof. Let $i \in \mathcal{N}$ and let \hat{A}^i satisfy (5.10). All constants C in the proof are independent of \hat{A}^i . We have

$$\mathcal{J}^{i,N}(\hat{A}^i, \bar{\mathbf{A}}^{-i}) \leq \mathcal{J}^{i,N}(0, \bar{\mathbf{A}}^{-i}) + 1 = \rho^i \left(\sum_{t=0}^T F \left(t, X_t^i[0], b_{(0, \bar{\mathbf{A}}^{-i})}^N \right) \right) + 1.$$

By Assumption 1.4 (i), Lemma 5.8, and Remark 5.2,

$$\rho^i \left(\sum_{t=0}^T F \left(t, X_t^i[0], b_{(0, \bar{\mathbf{A}}^{-i})}^N \right) \right) \leq C \mathbb{E} \left[T + \sum_{t=0}^T |X_t^i[0]|^2 \right] \leq C.$$

Therefore,

$$\mathcal{J}^{i,N}(\hat{A}^i, \bar{\mathbf{A}}^{-i}) \leq C. \quad (5.11)$$

We need now to bound $\mathcal{J}^{i,N}(\hat{A}^i, \bar{\mathbf{A}}^{-i})$ from below. We obtain by using successively Lemma 5.8, Assumptions 1.4 (i) and (iv), and Young's inequality that

$$\mathcal{J}^{i,N}(\hat{A}^i, \bar{\mathbf{A}}^{-i}) \geq \frac{1}{C} \mathbb{E} \left[\sum_{t=0}^{T-1} \left(\frac{1}{2} |\hat{A}_t^i|^2 - C |\hat{A}_t^i| \right) \right] - C \geq \frac{1}{C} \mathbb{E} \left[\sum_{t=0}^{T-1} |\hat{A}_t^i|^2 \right] - C. \quad (5.12)$$

We deduce then from (5.11) and (5.12) that $\mathbb{E} \left[\sum_{t=0}^{T-1} |\hat{A}_t^i|^2 \right] \leq C$, which concludes the proof. \square

In the following we fix a constant $c > 0$ such that the result of Lemma 5.9 holds and such that $\bar{A}^i \in \mathcal{A}^c$ for any $i \in \mathcal{N}$. Let b and b' in \mathcal{B}_2 , for any $(t, t', x) \in \mathcal{T} \times \bar{\mathcal{T}} \times \mathbb{R}^d$ we define

$$\Delta P(t, b, b') := P(t, b) - P(t, b'), \quad \Delta F(t', x, b, b') := F(t', x, b) - F(t', x, b').$$

For any $(x, A) \in \mathbb{R}^{Td} \times \mathcal{A}$ we define

$$\Delta \ell(x, A, b, b') := \sum_{t=0}^{T-1} \langle A_t, \Delta P(t, b, b') \rangle + \sum_{t=0}^T \Delta F(t, x_t, b, b').$$

Remark 5.10. For any $t \in \mathcal{T}$ and for any b and b' in \mathcal{B}_2 , we have

$$\|\Delta F(t, \cdot, b, b')\|_{\mathcal{G},1} \leq 2Cd_1(b, b')^{1/2}.$$

Indeed if $d_1(b, b') \geq 1$, Assumption 5.1 (i) yields

$$\|\Delta F(t, \cdot, b, b')\|_{\mathcal{G},1} \leq 2 \sup_{b \in \mathcal{B}_2} \|F(t, \cdot, b)\|_{\mathcal{G},1} \leq 2C.$$

If $d_1(b, b') \leq 1$, by Assumption 5.1 (ii) we have

$$\|\Delta F(t, \cdot, b, b')\|_{\mathcal{G},1} \leq Cd_1(b, b') \leq Cd_1(b, b')^{1/2}.$$

In the following lemma we study the convergence of the empirical belief to the reference belief $\bar{b} \in \mathcal{B}_2$.

Lemma 5.11. There exists $C > 0$ such that for any $i \in \mathcal{N}$ and for any $A^i \in \mathcal{A}^c$,

$$\mathbb{E} \left[d_1 \left(b_{(A^i, \bar{A}^{-i})}^N, \bar{b} \right) \right] \leq CN^{-\tau(d)}. \quad (5.13)$$

Proof. Let $i \in \mathcal{N}$ and let $A^i \in \mathcal{A}^c$. For any $t \in \mathcal{T}$, we have by the triangle inequality

$$d_1 \left(\mu_{(A^i, \bar{A}^{-i})}^N(t), \bar{\mu}(t) \right) \leq d_1 \left(\mu_{(A^i, \bar{A}^{-i})}^N(t), \mu_{\bar{A}}^N(t) \right) + d_1 \left(\mu_{\bar{A}}^N(t), \bar{\mu}(t) \right). \quad (5.14)$$

Let us consider the first term of the right-hand side. By definition of the distance d_1 ,

$$d_1 \left(\mu_{(A^i, \bar{A}^{-i})}^N(t), \mu_{\bar{A}}^N(t) \right) \leq \frac{1}{N} (|X_t^i[A^i] - \bar{X}_t^i| + |A_t^i - \bar{A}_t^i|), \quad \text{a.s.} \quad (5.15)$$

Since the controls \bar{A}^i and A^i belong to \mathcal{A}^c , the first-order moment of $X_t^i[A^i]$ and \bar{X}_t^i are finite as a consequence of Remark 5.2, thus

$$\mathbb{E} [|X_t^i[A^i] - \bar{X}_t^i| + |A_t^i - \bar{A}_t^i|] \leq C. \quad (5.16)$$

Therefore, by (5.15) and (5.16), we have

$$d_1 \left(\mu_{(A^i, \bar{A}^{-i})}^N(t), \mu_{\bar{A}}^N(t) \right) \leq \frac{C}{N}. \quad (5.17)$$

Let us consider now the second-term of the right-hand side of (5.14). We recall that $\bar{\mu}(t) = (id, \bar{\alpha}_t) \sharp \bar{m}(t)$. Since $\bar{A}_t^i = \bar{\alpha}_t(\bar{X}_t^i)$, we also have $\mu_{\bar{A}}^N(t) = (id, \bar{\alpha}_t) \sharp \bar{m}_{\bar{A}}^N(t)$. We deduce from the Lipschitz continuity of $(id, \bar{\alpha}_t)$ and from Lemma 3.2 that

$$\begin{aligned} d_1 \left(\mu_{\bar{A}}^N(t), \bar{\mu}(t) \right) &= d_1 \left((id, \bar{\alpha}_t) \sharp \bar{m}_{\bar{A}}^N(t), (id, \bar{\alpha}_t) \sharp \bar{m}(t) \right) \\ &\leq Cd_1 \left(\bar{m}_{\bar{A}}^N(t), \bar{m}(t) \right). \end{aligned} \quad (5.18)$$

The random variables \bar{X}_t^j are independent and $\mathcal{L}(\bar{X}_t^j) \sim \bar{m}(t)$. Therefore, Theorem 5.5 applies and yields

$$\mathbb{E} \left[d_1 \left(\bar{m}_{\mathcal{A}}^N(t), \bar{m}(t) \right) \right] \leq CN^{-\tau(d)}. \quad (5.19)$$

Combining (5.14), (5.17), (5.18), and (5.19), we obtain

$$\mathbb{E} \left[d_1 \left(\mu_{(A^i, \bar{\mathcal{A}}^{-i})}^N(t), \bar{\mu}(t) \right) \right] \leq CN^{-\tau(d)}.$$

It is then easy to verify that

$$\mathbb{E} \left[d_1 \left(m_{(A^i, \bar{\mathcal{A}}^{-i})}^N(T), \bar{m}(T) \right) \right] \leq CN^{-\tau(d)}.$$

Estimate (5.13) follows immediately. \square

Lemma 5.12. *There exists $C > 0$ such that for any $i \in \mathcal{N}$, for any $t \in \bar{\mathcal{T}}$, and for any $A^i \in \mathcal{A}^c$, we have*

$$\mathbb{E} \left[\left| \Delta F \left(t, X_t^i[A^i], b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \right| \right] \leq CN^{-\tau(d)/2}.$$

Proof. Let $N \in \mathbb{N}^*$ and let $t \in \bar{\mathcal{T}}$. By Remark 5.10, we have

$$\mathbb{E} \left[\left| \Delta F \left(t, X_t^i[A^i], b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \right| \right] \leq C \mathbb{E} \left[\left(1 + |X_t^i[A^i]| \right) d_1 \left(b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right)^{1/2} \right]. \quad (5.20)$$

Since $A^i \in \mathcal{A}^c$, by Remark 5.2 we have that $\mathbb{E} [|X_t^i[A^i]|^2] \leq C$. We obtain with the Cauchy-Schwarz inequality and Lemma 5.11 that

$$\begin{aligned} \mathbb{E} \left[\left(1 + |X_t^i[A^i]| \right) d_1 \left(b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right)^{1/2} \right] &\leq \mathbb{E} \left[\left(1 + |X_t^i[A^i]| \right)^2 \right]^{1/2} \mathbb{E} \left[d_1 \left(b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \right]^{1/2} \\ &\leq CN^{-\tau(d)/2}. \end{aligned} \quad (5.21)$$

Combining (5.20) and (5.21), we obtain the announced inequality. \square

Lemma 5.13. *There exists $C > 0$ such that for any $i \in \mathcal{N}$, for any $t \in \mathcal{T}$ and for any $A^i \in \mathcal{A}^c$, we have*

$$\mathbb{E} \left[\left| \langle A_t^i, \Delta P \left(t, b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \rangle \right| \right] \leq CN^{-\tau(d)/2}. \quad (5.22)$$

Proof. Let $i \in \mathcal{N}$, let $t \in \mathcal{T}$, and let $A^i \in \mathcal{A}^c$. By the Cauchy-Schwarz inequality, we have

$$\mathbb{E} \left[\left| \langle A_t^i, \Delta P \left(t, b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \rangle \right| \right] \leq C \left(\mathbb{E} \left[\left| \Delta P \left(t, b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \right|^2 \right] \right)^{1/2}. \quad (5.23)$$

We obtain with Assumptions 1.4 (iii-iv) and Lemma 5.11 that

$$\begin{aligned} \mathbb{E} \left[\left| \Delta P \left(t, b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \right|^2 \right] &\leq 2C \mathbb{E} \left[\left| \Delta P \left(t, b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \right| \right] \\ &\leq C \mathbb{E} \left[d_1 \left(b_{(A^i, \bar{\mathcal{A}}^{-i})}^N, \bar{b} \right) \right] \leq CN^{-\tau(d)}. \end{aligned} \quad (5.24)$$

Combining (5.23) and (5.24), we deduce (5.22). \square

We finally prove the main result of the section.

Proof of Theorem 5.4. Let $i \in \mathcal{N}$. We first show that for any $A^i \in \mathcal{A}^c$, the inequality

$$|\mathcal{J}^{i,N}(A^i, \bar{\mathbf{A}}^{-i}) - \mathcal{J}^i(A^i, \bar{b})| \leq CN^{-\tau(d)/2} \quad (5.25)$$

holds for some constant $C > 0$ independent of A^i . This will imply (5.5). For any $A^i \in \mathcal{A}^c$, we can write $\mathcal{J}^{i,N}(A^i, \bar{\mathbf{A}}^{-i}) = \rho^i(U)$ and $\mathcal{J}^i(A^i, \bar{b}) = \rho^i(V)$, where

$$\begin{aligned} U &:= \sum_{t=0}^{T-1} \ell \left(t, X_t^i[A^i], A_t^i, b_{(A^i, \bar{\mathbf{A}}^{-i})}^N \right) + F \left(T, X_T^i[A^i], b_{(A^i, \bar{\mathbf{A}}^{-i})}^N \right), \\ V &:= \sum_{t=0}^{T-1} \ell \left(t, X_t^i[A^i], A_t^i, \bar{b} \right) + F \left(T, X_T^i[A^i], \bar{b} \right). \end{aligned}$$

Lemma 5.7 and Lemma 5.8 yield

$$\begin{aligned} |\mathcal{J}^{i,N}(A^i, \bar{\mathbf{A}}^{-i}) - \mathcal{J}^i(A^i, \bar{b})| &\leq \rho^i(|U - V|) \\ &\leq C\mathbb{E}[|U - V|] = C\mathbb{E} \left[\left| \Delta \ell \left(X^i[A^i], b_{(A^i, \bar{\mathbf{A}}^{-i})}^N, \bar{b} \right) \right| \right]. \end{aligned}$$

We finally obtain (5.25) with Lemma 5.12 and Lemma 5.13.

Let us fix now $\hat{A}^i \in \mathcal{A}$ such that

$$\mathcal{J}^{i,N}(\hat{A}^i, \bar{\mathbf{A}}^{-i}) \leq \left(\inf_{A^i \in \mathcal{A}} \mathcal{J}^{i,N}(A^i, \bar{\mathbf{A}}^{-i}) \right) + \min \left\{ 1, N^{-\tau(d)/2} \right\}. \quad (5.26)$$

By Lemma 5.9, we have $\hat{A}^i \in \mathcal{A}^c$. Thus inequality (5.25) yields

$$\begin{aligned} \mathcal{J}^i(\hat{A}^i, \bar{b}) &\leq \mathcal{J}^{i,N}(\hat{A}^i, \bar{\mathbf{A}}^{-i}) + CN^{-\tau(d)/2} \\ &\leq \left(\inf_{A^i \in \mathcal{A}} \mathcal{J}^{i,N}(A^i, \bar{\mathbf{A}}^{-i}) \right) + CN^{-\tau(d)/2}. \end{aligned} \quad (5.27)$$

We apply again inequality (5.25) to $A^i = \bar{A}^i$. Using also the optimality of \bar{A}^i (with respect to \mathcal{J}^i), we obtain

$$\mathcal{J}^{i,N}(\bar{\mathbf{A}}) - CN^{-\tau(d)/2} \leq \mathcal{J}^i(\bar{A}^i, \bar{b}) \leq \mathcal{J}^i(\hat{A}^i, \bar{b}). \quad (5.28)$$

Finally, combining (5.27) and (5.28) we have

$$\mathcal{J}^{i,N}(\bar{\mathbf{A}}) \leq \left(\inf_{A^i \in \mathcal{A}} \mathcal{J}^{i,N}(A^i, \bar{\mathbf{A}}^{-i}) \right) + CN^{-\tau(d)/2},$$

which shows that $\bar{\mathbf{A}}$ is an ε -Nash equilibrium with $\varepsilon = CN^{-\tau(d)/2}$. \square

6 Conclusion

This paper has studied a mean field game model with risk averse agents, and provided a framework under which an equilibrium holds, for general composite risk measures and congestion terms. The specific structure of the integral cost of the agents has been exploited in order to rewrite the dynamic programming equations in a functional form (using the Moreau envelope and the proximal operator). In that way, the coupled system could be formulated as an equivalent fixed point equation, yielding the existence of a solution. Regularity properties have been obtained for risk averse agents. This has allowed to show that an optimal feedback control (for the mean field game) results in an ε -Nash equilibrium for a related dynamic game with N players. Future work could focus on the uniqueness of the Nash equilibrium with contraction arguments and smallness assumptions on the coupling terms. In this work, agents risk averse with respect to their own noise have been considered; investigating a mean field game model with common noise and risk averse agents would be of particular interest.

References

- [1] Yves Achdou, Francisco J. Buera, Jean-Michel Lasry, Pierre-Louis Lions, and Benjamin Moll. Partial differential equation models in macroeconomics. *Phil. Trans. R. Soc. A*, 372(2028):19 pages, 2014. [1](#)
- [2] Clémence Alasseur, Imen Ben Taher, and Anis Matoussi. An extended mean field game for storage in smart grids. *Journal of Optimization Theory and Applications*, pages 1–27, 2020. [2](#)
- [3] Philippe Artzner, Freddy Delbaen, Jean-Marc Eber, and David Heath. Coherent measures of risk. *Mathematical Finance*, 9(3):203–228, 1999. [1](#)
- [4] Heinz H. Bauschke and Patrick L. Combettes. *Convex analysis and monotone operator theory in Hilbert spaces*. CMS Books in Mathematics. Springer-Verlag New York, first edition, 2011. [10](#)
- [5] Charles Bertucci, Spyridon Vassilaras, Jean-Michel Lasry, Georgios S. Paschos, Merouane Debbah, and Pierre-Louis Lions. Transmit strategies for massive machine-type communications based on mean field games, Aug 2018. [2](#)
- [6] J. Frédéric Bonnans, Saeed Hadikhanloo, and Laurent Pfeiffer. Schauder estimates for a class of potential mean field games of controls. *Applied Mathematics & Optimization*, pages 1–34, 2019. [1](#)
- [7] Pierre Cardaliaguet and Charles-Albert Lehalle. Mean field game of controls and an application to trade crowding. *Mathematics and Financial Economics*, 12(3):335–363, 2018. [1](#)
- [8] Patrick Cheridito and Michael Kupper. Composition of time-consistent dynamic monetary risk measures in discrete time. *International Journal of Theoretical and Applied Finance*, 14(01):137–162, 2011. [7](#)
- [9] Hans Föllmer and Alexander Schied. *Stochastic finance: an introduction in discrete time*. Walter de Gruyter, 2011. [7](#)
- [10] Nicolas Fournier and Arnaud Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162(3):707–738, Aug 2015. [2](#), [18](#)
- [11] P. Jameson Graber and Alain Bensoussan. Existence and uniqueness of solutions for Bertrand and Cournot mean field games. *Applied Mathematics & Optimization*, pages 1–25, 2015. [1](#)
- [12] P. Jameson Graber, Vincenzo Ignazio, and Ariel Neufeld. Nonlocal Bertrand and Cournot mean field games with general nonlinear demand schedule. *arXiv preprint arXiv:2002.11055*, 2020. [1](#)
- [13] Olivier Guéant, Jean-Michel Lasry, and Pierre-Louis Lions. *Mean Field Games and Applications*, pages 205–266. Springer, Berlin, Heidelberg, 2011. [1](#)
- [14] Minyi Huang, Peter E Caines, and Roland P Malhamé. Large-population cost-coupled lqg problems with nonuniform agents: individual-mass behavior and decentralized ε -nash equilibria. *IEEE transactions on automatic control*, 52(9):1560–1571, 2007. [2](#)

- [15] Minyi Huang, Roland P. Malhamé, and Peter E. Caines. Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Communications in Information & Systems*, 6(3):221–252, 2006. [1](#)
- [16] Ziad Kobeissi. On classical solutions to the mean field game system of controls. *arXiv preprint arXiv:1904.11292*, 2019. [1](#)
- [17] Jean-Michel Lasry and Pierre-Louis Lions. Jeux à champ moyen. i–le cas stationnaire. *Comptes Rendus Mathématique*, 343(9):619–625, 2006. [1](#)
- [18] Jean-Michel Lasry and Pierre-Louis Lions. Jeux à champ moyen. ii–horizon fini et contrôle optimal. *Comptes Rendus Mathématique*, 343(10):679–684, 2006. [1](#)
- [19] Jean-Michel Lasry and Pierre-Louis Lions. Mean field games. *Japanese journal of mathematics*, 2(1):229–260, 2007. [1](#)
- [20] Jun Moon and Tamer Başar. Linear quadratic risk-sensitive and robust mean field games. *IEEE Transactions on Automatic Control*, 62(3):1062–1077, March 2017. [2](#)
- [21] Basel Committee on Banking Supervision. Messages from the academic literature on risk measurement for the trading book, 2011. [2](#)
- [22] Laurent Pfeiffer. Optimality conditions in variational form for non-linear constrained stochastic control problems. *Mathematical Control & Related Fields*, 2019. Online first. [16](#)
- [23] Andrzej Ruszczyński. Risk-averse dynamic programming for markov decision processes. *Mathematical Programming*, 125(2):235–261, October 2010. [6](#), [8](#)
- [24] Andrzej Ruszczyński and Alexander Shapiro. Conditional risk mappings. *Mathematics of Operations Research*, 31(3):544–561, 2006. [1](#), [7](#)
- [25] Naci Saldi, Tamer Basar, and Maxim Raginsky. Markov–Nash equilibria in mean-field games with discounted cost. *SIAM Journal on Control and Optimization*, 56(6):4256–4287, 2018. [2](#)
- [26] Alexander Shapiro. Minimax and risk averse multistage stochastic programming. *European Journal of Operational Research*, 219(3):719 – 726, 2012. Feature Clusters. [1](#)
- [27] Hamidou Tembine, Quanyan Zhu, and Tamer Başar. Risk-sensitive mean-field games. *IEEE Transactions on Automatic Control*, 59(4):835–850, April 2014. [2](#)
- [28] Cédric Villani. *Optimal transport: Old and New*. Springer Verlag, 2008. [3](#)