



HAL
open science

Draft Genome Sequences of Two Carbapenemase-Producing *Klebsiella pneumoniae* Strains Isolated from Blood Cultures

Rémi Le Guern, Teddy Grandjean, Karine Faure, Marvin Bauduin, Eric
Kipnis, Rodrigue Dessein

► **To cite this version:**

Rémi Le Guern, Teddy Grandjean, Karine Faure, Marvin Bauduin, Eric Kipnis, et al.. Draft Genome Sequences of Two Carbapenemase-Producing *Klebsiella pneumoniae* Strains Isolated from Blood Cultures. *Microbiology Resource Announcements*, 2018, 7 (8), 10.1128/MRA.01057-18 . hal-02566182

HAL Id: hal-02566182

<https://hal.science/hal-02566182>

Submitted on 31 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Draft Genome Sequences of Two Carbapenemase-Producing *Klebsiella pneumoniae* Strains Isolated from Blood Cultures

Rémi Le Guern,^{a,b} Teddy Grandjean,^a Karine Faure,^{a,c} Marvin Bauduin,^a Eric Kipnis,^{a,d} Rodrigue Dessein^{a,b}

^aRecherche Translationnelle: Relations Hôte Pathogènes, University of Lille, Lille, France

^bInstitut de Microbiologie, CHU Lille, Lille, France

^cUnité de Maladies Infectieuses, CHU Lille, Lille, France

^dService de Réanimation Chirurgicale, CHU Lille, Lille, France

ABSTRACT Carbapenemase-producing *Klebsiella pneumoniae* represents an emerging public health issue. Here, we present the draft whole-genome sequences of *K. pneumoniae* clinical strains KPL0.1 (OXA-48 carbapenemase) and KPL0.2 (NDM-1 carbapenemase). These genome sequences should help in investigating pathophysiological mechanisms of digestive colonization or infection with these highly resistant bacteria.

Carbapenemase-producing *Enterobacteriaceae* (CPE) are resistant to most beta-lactams, including last-line options such as carbapenems, and are an emerging public health issue (1). *Klebsiella pneumoniae* colonizes the human gastrointestinal tract and can persist in the hospital environment, and outbreaks of carbapenemase-producing *K. pneumoniae* are frequently described (2). In France, the most frequently identified carbapenemases are OXA-48 (86% of the CPE) and NDM (9% of the CPE) (3). Two strains of carbapenemase-producing *K. pneumoniae* were isolated in the Lille University Teaching Hospital (CHU Lille, France), KPL0.1 (producing OXA-48 carbapenemase) and KPL0.2 (producing NDM-1 carbapenemase). Both strains were isolated from blood cultures using the Virtuo automated system (bioMérieux, Marcy-l'Étoile, France) and identified by matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry (Bruker Daltonics, Inc., Billerica, MA). KPL0.1 was isolated from a 77-year-old patient following digestive surgery, and KPL0.2 came from an 84-year-old patient with a urinary tract infection.

The DNA of each isolate was extracted using a QIAamp DNA minikit (Qiagen, Hilden, Germany), and quality control was performed using a Qubit v3.0 fluorometer (Thermo Fisher Scientific, Waltham, MA). Libraries were prepared using the Nextera XT DNA Library prep kit (Illumina, San Diego, CA), followed by paired-end (2 × 150-bp) sequencing on a HiSeq platform (Illumina). Genome coverage was about 650× for each isolate. Paired reads were filtered, and Nextera adapters were removed using Trimmomatic on a Galaxy server using default settings (4). Processed reads were *de novo* assembled with Unicycler. Genomes were annotated using the Prokaryotic Genome Annotation Pipeline (PGAP) v4.5 (5).

The draft genome of KPL0.1 consists of 5,825,863 bp and has a mean G+C content of 56.70%. A total of 5,628 protein-coding genes were annotated, including 92 RNA-coding genes and 75 tRNAs, and the remaining genes were annotated as hypothetical proteins. KPL0.1 belongs to sequence type 307 (ST-307) and possesses the KL102 capsule locus (6). The following resistance genes were predicted using ResFinder v3.0 (7): five beta-lactam resistance genes (*bla*_{OXA-48}, *bla*_{SHV-28}, *bla*_{TEM-1B}, *bla*_{CTX-M-15}, *bla*_{OXA-1}), four aminoglycoside resistance genes [*aac*(3)-IIa, *aph*(3'')-Ib, *aph*(6)-Id, *aac*(6')-Ib-cr], five fluoroquinolone resistance genes [*qnrS1*, *qnrB1*, *aac*(6')-Ib-cr, *oqxA*, *oqxB*], and

Received 26 July 2018 Accepted 31 July 2018 Published 30 August 2018

Citation Le Guern R, Grandjean T, Faure K, Bauduin M, Kipnis E, Dessein R. 2018. Draft genome sequences of two carbapenemase-producing *Klebsiella pneumoniae* strains isolated from blood cultures. *Microbiol Resour Announc* 7:e01057-18. <https://doi.org/10.1128/MRA.01057-18>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2018 Le Guern et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Rodrigue Dessein, rodrigue.dessein@chru-lille.fr.

individual antimicrobial resistance genes (fosfomycin, *fosA*; phenicol, *catB4*; sulfonamide, *sul2*; trimethoprim, *dfrA14*; tetracycline, *tetA*).

The draft genome of KPL0.2 consists of 5,741,089 bp with a mean G+C content of 56.94%. A total of 5,570 protein-coding genes were annotated, including 93 RNA-coding genes and 79 tRNAs, and the remaining genes were annotated as hypothetical proteins. KPL0.2 belongs to ST-147 and possesses the KL64 capsule locus. The following resistance genes were predicted: five beta-lactam resistance genes (*bla*_{NDM-1}, *bla*_{SHV-11}, *bla*_{TEM-1B}, *bla*_{CTX-M-15}, *bla*_{OXA-1}), four aminoglycoside resistance genes [*aac(3)-IIa*, *strA*, *aph(6)-Id*, *aac(6')-Ib-cr*], four fluoroquinolone resistance genes [*qnrB1*, *aac(6')-Ib-cr*, *oqxA*, *oqxB*], and individual antimicrobial resistance genes (fosfomycin, *fosA*; phenicol, *catB4*, sulfonamide, *sul1*; trimethoprim, *dfrA1*; tetracycline, *tetA*).

ST-147 was identified as one of the three major STs of carbapenemase-producing *K. pneumoniae* in a worldwide study; ST-147 is usually recovered from India, Italy, and Greece (8). ST-307 is described as highly prevalent in Italy (9). Overall, these two strains are representative of the carbapenemase-producing *K. pneumoniae* isolates in Europe. Further studies are warranted to investigate the pathophysiological mechanisms of digestive colonization or infection with these highly resistant bacteria.

Data availability. These two draft whole-genome shotgun projects have been deposited at DDBJ/ENA/GenBank under the accession numbers [QHMA00000000](https://doi.org/10.1093/mgen.0000000000000000) (KPL0.1) and [PYBH00000000](https://doi.org/10.1093/mgen.0000000000000000) (KPL0.2). The versions described in this paper are QHMA01000000 (KPL0.1) and PYBH01000000 (KPL0.2). The SRA accession number is [SRP155589](https://doi.org/10.1093/mgen.0000000000000000).

REFERENCES

- Nordmann P, Naas T, Poirel L. 2011. Global spread of carbapenemase-producing *Enterobacteriaceae*. *Emerg Infect Dis* 17:1791–1798. <https://doi.org/10.3201/eid1710.110655>.
- Tzouveleki LS, Markogiannakis A, Psychogiou M, Tassios PT, Daikos GL. 2012. Carbapenemases in *Klebsiella pneumoniae* and other *Enterobacteriaceae*: an evolving crisis of global dimensions. *Clin Microbiol Rev* 25:682–707. <https://doi.org/10.1128/CMR.05035-11>.
- Dortet L, Cuzon G, Ponties V, Nordmann P. 2017. Trends in carbapenemase-producing *Enterobacteriaceae*, France, 2012 to 2014. *Euro Surveill* 22:30461. <https://doi.org/10.2807/1560-7917.ES.2017.22.6.30461>.
- Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltmann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res* 15:1451.
- Schwaber MJ, Klarfeld-Lidji S, Navon-Venezia S, Schwartz D, Leavitt A, Carmeli Y. 2008. Predictors of carbapenem-resistant *Klebsiella pneumoniae* acquisition among hospitalized adults and effect of acquisition on mortality. *Antimicrob Agents Chemother* 52:1028–1033. <https://doi.org/10.1128/AAC.01020-07>.
- Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, Holt KE. 2016. Identification of *Klebsiella* capsule synthesis loci from whole genome data. *Microb Genom* 2:e000102. <https://doi.org/10.1099/mgen.0.000102>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
- Lascols C, Peirano G, Hackel M, Laupland KB, Pitout JDD. 2013. Surveillance and molecular epidemiology of *Klebsiella pneumoniae* isolates that produce carbapenemases: first report of OXA-48-like enzymes in North America. *Antimicrob Agents Chemother* 57:130–136. <https://doi.org/10.1128/AAC.01686-12>.
- Bartolini A, Basso M, Franchin E, Menegotto N, Ferrari A, De Canale E, Andreis S, Scaggiante R, Stefani S, Palù G, Parisi SG. 2017. Prevalence, molecular epidemiology and intra-hospital acquisition of *Klebsiella pneumoniae* strains producing carbapenemases in an Italian teaching hospital from January 2015 to September 2016. *Int J Infect Dis* 59:103–109. <https://doi.org/10.1016/j.ijid.2017.04.007>.