



HAL
open science

Biotic formation of methylmercury: A bio–physico–chemical conundrum

Andrea Bravo, Claudia Cosio

► **To cite this version:**

Andrea Bravo, Claudia Cosio. Biotic formation of methylmercury: A bio–physico–chemical conundrum. *Limnology and Oceanography*, 2020, 65 (5), pp.1010-1027. 10.1002/lno.11366 . hal-02571695

HAL Id: hal-02571695

<https://hal.science/hal-02571695v1>

Submitted on 13 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biotic formation of methylmercury: A bio–physico–chemical conundrum

Andrea G. Bravo ,^{1*} Claudia Cosio ^{2*}

¹Department of Marine Biology and Oceanography, Institute of Marine Sciences, Spanish National Research Council (CSIC), Barcelona, Spain

²Université de Reims Champagne Ardennes, UMR-I 02 INERIS-URCA-ULH SEBIO, Unité Stress Environnementaux et BIOSurveillance des milieux aquatiques, Reims, France

Abstract

Mercury (Hg) is a natural and widespread trace metal, but is considered a priority pollutant, particularly its organic form methylmercury (MMHg), because of human's exposure to MMHg through fish consumption. Pioneering studies showed the methylation of divalent Hg (Hg^{II}) to MMHg to occur under oxygen-limited conditions and to depend on the activity of anaerobic microorganisms. Recent studies identified the *hgcAB* gene cluster in microorganisms with the capacity to methylate Hg^{II} and unveiled a much wider range of species and environmental conditions producing MMHg than previously expected. Here, we review the recent knowledge and approaches used to understand Hg^{II} -methylation, microbial biodiversity and activity involved in these processes, and we highlight the current limits for predicting MMHg concentrations in the environment. The available data unveil the fact that Hg^{II} methylation is a bio-physico-chemical conundrum in which the efficiency of biological Hg^{II} methylation appears to depend chiefly on Hg^{II} and nutrients availability, the abundance of electron acceptors such as sulfate or iron, the abundance and composition of organic matter as well as the activity and structure of the microbial community. An increased knowledge of the relationship between microbial community composition, physico-chemical conditions, MMHg production, and demethylation is necessary to predict variability in MMHg concentrations across environments.

The mercury problem

Mercury (Hg) is a natural and ubiquitous trace metal in the environment that might damage the central nervous system and causes tremors, distorted speech, kidney effects, respiratory failure, dizziness, blurred vision, hallucinations, and even death in severely exposed people (Clarkson and Magos 2006). This pollutant is naturally emitted during episodic events such as volcanic eruptions or ubiquitous weathering of Hg-containing rocks in the Earth's crust and geothermal activity. Among anthropogenic Hg sources, artisanal and small-scale gold mining, coal combustion, production of nonferrous metals, cement production, and disposal of wastes containing Hg are of special concern (UNEP 2013). Hg is unique among transition metals due to its high volatility as gaseous elemental Hg (Hg^0), with a residence time in the atmosphere of about 6–12 months, allowing for long-range transport

of Hg. Although both Hg^0 and inorganic divalent Hg (Hg^{II}) are released from many sources through a variety of natural and anthropogenic processes, the reported rise in Hg levels in the biosphere, and in terrestrial and marine systems is a consequence of anthropogenic emissions (Amos et al. 2013; Lamborg et al. 2014; Kocman et al. 2017). In contrast to Hg^0 and Hg^{II} , direct anthropogenic sources of organic Hg, mono-methylmercury (MMHg, i.e., CH_3Hg^+) or dimethylmercury (DMHg, i.e., $[\text{CH}_3]_2\text{Hg}$) are scarce.

The chemical behaviors of the different chemical forms of Hg (i.e., Hg^0 , Hg^{II} , CH_3Hg^+ and $(\text{CH}_3)_2\text{Hg}$) play critical roles in the biogeochemical cycling of Hg. Hg^0 allows for long-range transport (Jackson 1997; Pirrone et al. 2009), Hg^{II} is the dominant reservoir for Hg in soils and aquatic systems (Fleck et al. 2015; Eklöf et al. 2018), and MMHg is bioconcentrated and biomagnified in aquatic food webs, reaching up to 80–100% of the total-Hg (THg) measured in fish muscle (Bloom 1992; Mason et al. 2012; Bravo et al. 2014). As a consequence, MMHg exposure through fish consumption is of special concern for human health. A recent study performed in 175 countries, showed that 38% of studied populations (mainly insular and developing nations) were exposed to doses of MMHg above governmental thresholds (Lavoie et al. 2018). Indeed, concentration of Hg in fish is known to repeatedly overpass environmental quality guidelines even in

*Correspondence: andrea.bravo@icm.csic.es (A.G.B.); claudia.cosio@univ-reims.fr (C.C.)

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Author Contribution Statement: The two authors have equally contributed to this work.

absence of local sources (Depew et al. 2013; Åkerblom et al. 2014; Eagles-Smith et al. 2016).

Comprehensive evaluations of the chemical and physical processes that govern Hg distribution and fate among the major environmental compartments can be found in the literature (Chételat et al. 2015; Sundseth et al. 2015; Kim et al. 2016; Bjørklund et al. 2017; Dranguet et al. 2017; Paranjape and Hall 2017; Klapstein and Driscoll 2018). Briefly, in the water column Hg^{II} can (1) be reduced to Hg^0 and reemitted back to the atmosphere, (2) methylated to the organic form MMHg, or (3) bind to organic matter (OM) as well as inorganic particles and directly deposit to bottom sediments. MMHg formed in aquatic ecosystems can also deposit to sediments, be methylated and/or form DMHg. Part of DMHg might be re-emitted to the atmosphere or again degraded to MMHg, which can also be biotically (Barkay et al. 2003) or abiotically demethylated (Fernández-Gómez et al. 2013). Although some of the MMHg found in aquatic systems might come from the degradation of DMHg to MMHg, i.e., in oceans (Mason et al. 2012), several studies concluded that most of the MMHg measured in ecosystems was formed in situ or in the surrounding catchment (e.g., soils, wetlands, etc.) and subsequently transported into rivers, lakes (Louis et al. 1996; Eklöf et al. 2012; Bravo et al. 2017), and oceans (Schartup et al. 2015). Abiotic methylation of Hg^{II} is possible if suitable methyl donors are present (Celo et al. 2006; Munson et al. 2018). Nevertheless, recent studies have shown that biological Hg^{II} methylation is in most environments performed by a variety of microorganisms, carrying the *hgcA* and *hgcB* gene cluster (Gilmour et al. 2013, 2018; Parks et al. 2013; Yu et al. 2018). An increasing number of recent studies detailed below have intended to evaluate the biotic Hg^{II} methylation by studying the biodiversity and activity of *hgcAB*+ microorganisms (Gionfriddo et al. 2016; Bravo et al. 2018*b,a*; Bowman et al. 2019; Jones et al. 2019; Villar et al. 2019). Nevertheless, it is established that net MMHg production also depends on other concomitant processes, including (1) the composition and activity of the whole microbial community that in turn modulate the activity of *hgcAB*+ microorganisms (Bravo et al. 2018*a*), (2) physico-chemistry that controls Hg^{II} bioavailability (Schaefer and Morel 2009; Jonsson et al. 2012; Chiasson-Gould et al. 2014), and uptake in microorganisms (Schaefer et al. 2011), and (3) biotic and abiotic MMHg demethylation (Du et al. 2019).

Recent reviews critically summarized Hg^{II} uptake and MMHg efflux in methylating anaerobes, methods and equations to analyze Hg methylation rates (Regnell and Watras 2019) as well as chemotrophic and biotic Hg methylation and demethylation processes by anaerobes and phototrophs (Grégoire and Poulain 2018; Du et al. 2019). In this review, we aim to summarize the main findings on mechanisms responsible for the formation of MMHg, focusing in detail on the new knowledge recently gained on microorganisms involved in MMHg formation. We also aim to highlight pitfalls and limitations impeding the progress in the

current understanding, and we propose a road map to overcome these limitations. Indeed, to improve our ability to predict MMHg generation in the environment, a current research priority is to better understand the distribution of methylating populations in the context of the physico-chemical constraints known to affect MMHg production.

Methylmercury formation is widespread in the environment

Recent advances in the methodology to (1) determine in situ Hg^{II} methylation (Jonsson et al. 2014) and (2) identify the organisms involved in this process (Parks et al. 2013; Christensen et al. 2016) have revealed that MMHg can be formed in a wider range of environments than previously identified. Indeed, 30 yr ago, first studies showed that Hg^{II} methylation in aquatic systems occurred mainly in sediments and under anaerobic conditions (Compeau and Bartha 1984; Korthals and Winfrey 1987; Pak and Bartha 1998). In general, sediments and sinking particles are a complex matrix of solid phases including clays, quartz, metal oxides (FeOOH , MnO_2 , AlO_3) carbonates, sulfides and a number of other minerals and OM. They provide various microenvironments and habitats to organism populations notably bacteria, archaea, algae, diverse invertebrates, and so forth. To date, biological Hg^{II} -methylation is known to be mediated by species carrying the *hgcAB* gene cluster (Parks et al. 2013). Because all the identified microorganisms with Hg^{II} -methylating capacity were anaerobes, it was assumed for a long time that MMHg formation was occurring in strictly anoxic environments, i.e., sediments. Nonetheless, several studies revealed that Hg^{II} -methylation can occur in oxygen deficient zones of water column (Eckley et al. 2005; Malcolm et al. 2010), sediments (Drott et al. 2008; Hines et al. 2012; Bouchet et al. 2013; Jonsson et al. 2014; Bravo et al. 2015; Liem-Nguyen et al. 2016) flooded soils, e.g., wetlands (Louis et al. 1996; Tjerngren et al. 2012; Windham-Myers et al. 2014) and ponds (Lehnher et al. 2012; MacMillan et al. 2015; Herrero Ortega et al. 2018). In recent years, Hg^{II} -methylation processes were in addition observed in microenvironments such as periphyton, growing on macrophytes (Cleckner et al. 1999; Mauro et al. 2002; Guimarães et al. 2006; Achá et al. 2011; Hamelin et al. 2011; Bouchet et al. 2018) and settling particles of oxic water columns, including pelagic ocean waters (Monperrus et al. 2007; Cossa et al. 2009; Sunderland et al. 2009; Lehnher et al. 2011; Gascón Díez et al. 2016). Initially, MMHg formation in oxic waters was considered negligible due to high redox and low concentrations of bacteria and nutrients, but studies demonstrated that about 20–40% of the MMHg measured below the surface mixed layer originates from the surface and enters deeper ocean waters (Blum et al. 2013). Similarly, a query of more than 3500 publicly available microbial metagenomes performed by Podar et al. (2015) unveiled the presence of *hgcAB*-like genes in sediments and in previously unsuspected environments, including invertebrate digestive tracts, thawing permafrost soils, coastal “dead zones,” soils and

extreme environments. Moreover, a recent study assessing 243 metagenomes from the *Tara* Oceans expedition reported high abundances of *hgcAB* genes in 77 samples across all oceans (Villar et al. 2019). The progress in genetics (Gilmour et al. 2013; Parks et al. 2013; Podar et al. 2015; Bravo et al. 2018b,a; Liu et al. 2018b; Jones et al. 2019) combined with recent advances in the use of stable isotopes to determine Hg^{II} methylation rate constants in sediments (Monperrus et al. 2007; Jonsson et al. 2012; Bravo et al. 2014, 2015), lakes (Eckley and Hintelmann 2006), water columns and oceans (Munson et al. 2018) as well as in sinking particles of marine and lake waters (Lehnher et al. 2011; Gascón Díez et al. 2016) have demonstrated that the potential for MMHg formation in the environment is widespread across ecosystems.

Toward a better understanding of microbial methylmercury formation

The discovery of *hgcAB*

Fifty years ago, a decade after the first observation of the Minamata disease in Japan, pioneering research pointed to surface sediments, and bacteria activity as responsible of Hg^{II}-methylation (Jensen and Jernelöv 1969). One of the first studies targeting MMHg and bacteria, evaluated the Hg^{II}-methylation capacity of *Pseudomonas fluorescens*, *Mycobacterium phlei*, *Escherichia coli*, *Aerobacter aerogenes*, and *Bacillus megaterium* over a 7-day period in pure cultures (Vonk and Sijpesteijn 1973). In the presence of sublethal amounts of HgCl₂, tested bacteria produced 49 to 169 ng L⁻¹ d⁻¹ of MMHg in aerobic conditions (Vonk and Sijpesteijn 1973). Another decade later, sulfate-reducing bacteria (SRB) were eventually identified as major Hg^{II}-methylator in saltmarsh through inhibition of their activity with sodium molybdate and isolation of *Desulfovibrio desulfuricans* from sediments (Compeau and Bartha 1985).

The relationship between bacterial sulfate reduction and Hg^{II}-methylation was studied adding Hg^{II} to anoxic sediment slurries or lake water overlying intact sediment cores collected in Quabbin Reservoir, MA (Gilmour et al. 1992). Comparable profiles of sulfate reduction and Hg^{II}-methylation in sediment cores were reported, further suggesting that Hg^{II}-methylation was linked to this specific bacterial metabolism. Almost 20 yr ago, a correlation between the Hg^{II}-methylation and sulfate reduction rates in sediment of a saltmarsh was also shown (King et al. 2000, 2001). Sulfate reduction was then accepted as the main metabolic pathway related to Hg^{II}-methylation. In 2006, two studies revealed the role of iron-reducing bacteria (FeRB) on Hg^{II}-methylation in ferruginous conditions (Fleming et al. 2006; Kerin et al. 2006). In 2010, Hamelin et al. further identified methanogens as important Hg^{II}-methylators in lake periphyton. Two laboratory studies later confirmed the efficiency of methanogens in converting Hg^{II} to MMHg (Yu et al. 2013; Gilmour et al. 2018). By culturing and isolating Hg^{II}-methylating strains or by using inhibitors of known Hg^{II}-methylators such as molybdate for sulfate-reduction and BESA for methanogenesis, Hg^{II}-methylation has mainly been then attributed to the action of SRB (Devereux et al. 1996; Pak and

Bartha 1998; Hylander 2000; King et al. 2001; Achá et al. 2011, 2012; Yu et al. 2012; Bravo et al. 2016), and in some cases to FeRB (Fleming et al. 2006; Bravo et al. 2015, 2018b) as well as methanogens (Hamelin et al. 2011; Bravo et al. 2018a).

A recent breakthrough in the understanding of the biological Hg^{II}-methylation pathway was the identification of a two-gene cluster, *hgcAB* -involved in C1 metabolism and the acetyl-CoA pathway (Qian et al. 2016) required for Hg^{II}-methylation (Parks et al. 2013). The gene *hgcA* encodes a corrinoid protein that is essential for the biosynthesis of the folate branch of acetyl-CoA pathway, whereas the gene *hgcB* encodes a ferredoxin-like protein thought to be an electron donor to *hgcA* (Parks et al. 2013). Both provide methyl groups required for Hg^{II} methylation, although it is not clear whether MMHg production is a controlled or an accidental metabolic process (Qian et al. 2016). However, deletion of either gene eliminated Hg^{II} methylation in *Desulfovibrio desulfuricans* ND132 (Parks et al. 2013). By directly measuring Hg^{II} methylation in several bacterial and archaeal strains encoding *hgcAB*, Gilmour et al. (2013) confirmed that Hg^{II}-methylation capability could be predicted by the presence of *hgcAB* in the genome. For the first time, Gilmour et al. (2013) demonstrated Hg^{II}-methylation capability in previously completely unsuspected species including syntrophic, acetogenic, and fermentative *Firmicutes*.

In recent years, the biodiversity of *hgcAB* microorganisms was actively studied in contrasting environments. First biodiversity studies using this newly identified gene cluster were based on classical polymerase chain reaction (PCR) amplification with one pair of primers targeting *hgcA* developed based on the couple of available sequenced genomes of methylating strains at that time (Table 1), followed by cloning and sequencing (Bae et al. 2014; Liu et al. 2014; Schaefer et al. 2014; Smith et al. 2015). In soils of the Florida Everglades, the sequences identified were distributed in diverse phyla, including *Deltaproteobacteria*, *Chloroflexi*, *Firmicutes*, and *Methanomicrobia*; however, *hgcA* clone libraries from all sites were dominated by sequences clustering within the order *Syntrophobacterales* (Bae et al. 2014) (Table 2). By comparing the taxonomically identified *hgcA* sequences with the activity of SRB (mRNA of *dsrB* gene), Bae et al. concluded that *Syntrophobacterales* largely dominated the Hg^{II} methylating microbial community of the Florida Everglades (Bae et al. 2014). In the Three Gorges Reservoir in China, PCR amplification and sequencing of *hgcA* gene resulted in the identification of *δ-Proteobacteria*, methanogens and a *Clostridia* group as putative Hg^{II} methylators in this ecosystem. Authors reported in addition a positive correlation between the abundance of *hgcA* and *dsrB* genes and MMHg concentrations, suggesting SRB as the main group responsible for Hg^{II} methylation in those systems (Luo et al. 2016). In Wanshan Hg mining area of China, the taxonomically annotated sequences were related to *δ-Proteobacteria*, *Firmicutes*, *Chloroflexi*, and *Euryarchaeota* (Liu et al. 2014). In temperate and tropical wetland soils, *hgcA* gene sequences were attributed to *δ-Proteobacteria*, *Chloroflexi*, and *Methanomicrobia* (Schaefer et al. 2014). In nine rice paddy soils sampled in three mining areas in China, *hgcA*+ microbes were dominated by *Proteobacteria* or *Euryarchaeota* in six and three sites,

Table 1. Published studies on *hgcA* diversity and quantification in environmental samples. Sequences and position of primers used, amplicon sizes, targets, and used methods are given.

References	Primer sequences 5' - > 3'	Target	Amplicon size (bp)	Methods
Bae et al. 2014	hgcA_268F GGNRTYAAAY RTNTGGTYGC hgcB_1198R CADGCNCCRCAYTCVATRCA NA	<i>hgcA-hgcB</i>	888 to 945	PCR and cloning
Bowman et al. 2019	As Schaefer et al. 2014			Metagenomics
Bravo et al. 2018a	As Schaefer et al. 2014			PCR and high throughput sequencing
Bravo et al. 2018b	As Schaefer et al. 2014			PCR and high throughput sequencing
Bravo et al. 2016	hgcA_262F GGNRTYAAAYRTNTGGTYGC hgcA_912R GGTGTAGGGGGTGAGCCSGRWARKT	<i>hgcA</i>	650	qPCR
Christensen et al. 2016, 2018	ORNL-HgcAB-uni-268F AAYGTCTGGTYGNGCVGG ORNL-HgcAB-uni-1198R CABGCNCCRCAYTCVATRCA ORNL-Delta-HgcA-181F GCCAACTACAAGMTGASCTWC ORNL-Delta-HgcA-287R CCSGCNCCRACCCAGACRTT ORNL-SRB-firm-HgcA-444F TGGDCCGGTDARAGCWAARGATA	<i>hgcA-hgcB</i> <i>hgcA</i> of SRB <i>hgcA</i> of Firmicutes	818 to 1020 107 167	PCR qPCR qPCR
Dranguet et al. 2017	ORNL-SRB-firm-HgcA-610R AAAAGAGHAYBCCAAAAATCA ORNL-archaea-HgcA-184F AAYTAWCNCTSGYTTYGAYGC	<i>hgcA</i> of Archae	125	qPCR
Du et al. 2017	ORNL-archaea-HgcA-308R TCDGTCCCRABGTSCCYTT			qPCR
Gionfriddo et al. 2016	As Bravo et al. 2016			qPCR, PCR, and cloning
Lei et al. 2019	As Schaefer et al. 2014			Metagenomics
Liu et al. 2014	NA			qPCR
Liu et al. 2018a	As Christensen et al. 2016 hgcA_626F GGNRTYAAAYRTCTGGTYGC hgcA_941R CGCATYCCCTTYBACNCC As Schaefer et al. 2014	<i>hgcA</i>	315	PCR and cloning, qPCR
Liu et al. 2018b	hgcA_515F GTGCCAGCMGCCGCGGTAA hgcA_806R GGACTACHVGGGTWTCTAAT As Christensen et al. 2016	<i>hgcA</i> <i>hgcA</i> <i>hgcA</i> <i>hgcAB</i>	291	PCR and cloning qPCR
Ma et al. 2017	As Bae et al. 2014			qPCR, metagenomics
Ndu et al. 2018	As Christensen et al. 2016			PacBio sequencing
Podar et al. 2015	NA	<i>hgcAB</i>		qPCR
Schaefer et al. 2014	hgcA_261F CGGCATCAAYGTCTGGTYGC hgcA_912R GGTGTAGGGGGTGAGCCSGRWARKT NA	<i>hgcAB</i> <i>hgcA</i>		PCR and cloning, qPCR Metagenomics
Villar et al. 2019	As Christensen et al. 2016			PCR and cloning Metagenomics
Vishnivetskaya et al. 2018	As Schaefer et al. 2014			PCR and cloning, qPCR
Xu et al. 2019				PCR and high throughput sequencing

Table 2. Main characteristics and outcomes of published studies on *hgcA* biodiversity in environmental samples.

Method	Studied environment	Number of sequence/reads	Number of OTUs	Dominant group	Dominant Deltaproteobacteria	Reference
PCR and cloning	Florida Everglades	220	168	<i>Deltaproteobacteria</i>	<i>Syntrophobacteriales</i>	Bae et al. 2014
	Wetlands	108	40	<i>Methanomicrobia</i>	<i>Geobacter</i>	Schaefer et al. 2014
	Three gorges reservoir	151	151	Unidentified	Unidentified	Du et al. 2017
	Rice paddy soils	~ 1800	190	<i>Proteobacteria</i> <i>Euryarchaeota</i>	Unidentified	Liu et al. 2018a
PCR and high throughput sequencing	Laboratory and sediment slurries			Unidentified	Unidentified	Ndu et al. 2018
	Rice paddy soils	5	?	Unidentified		Vishnivetskaya et al. 2018
	Sulfate-impacted lakes	300	174	<i>Geobacteraceae</i> <i>Methanomicrobia</i> <i>Unidentified</i>		Jones et al. 2019
Metagenomics	Boreal lakes	78,642	225	<i>Deltaproteobacteria</i>	Unidentified	Bravo et al. 2018a
	Lake Geneva	741,890	356	<i>Deltaproteobacteria</i>	Unidentified (<i>Geobacter</i>)	Bravo et al. 2018b
Metaproteomics	Rice paddy soils	1,257,577	573	<i>Deltaproteobacteria</i>	Unidentified (<i>Geobacter</i>)	Liu et al. 2018a
	Boreal forest soils	823,000,000		Environmental compartment dependent		Xu et al. 2019
	Global			<i>Nitrospirinae</i>		Podar et al. 2015
	Antarctic Sea-ice and brine	901,610,484		<i>Methanoregula</i> spp.	<i>Geobacter</i>	Gionfriddo et al. 2016
	Rice paddy soils	885,923	27	<i>Deltaproteobacteria</i>		Liu et al. 2018b
Metaproteomics	Sulfate-impacted lakes	381–102		<i>Deltaproteobacteria</i>		Jones et al. 2019
	Eight sites	1				Christensen et al. 2019
Metaproteomics	Tara gene catalogs oceans	111,530,851	10	<i>Nitrospirinae</i>	<i>Desulfovibrionales</i>	Villar et al. 2019
	EFPC and Hinds Creek, TN	15,270–16,852		<i>Deltaproteobacteria</i>		Christensen et al. 2019

respectively. Only nine of the 190 operational taxonomic unit (OTUs) found in these rice paddy soils were common to all sites (Liu et al. 2018a).

Based on a higher number of sequenced microbial genomes now available, Christensen et al. (2016) developed a broad range *hgcAB* primer pair that improved the coverage of prior developed primers by 10% (Bae et al. 2014; Schaefer et al. 2014), and several clade-specific PCR primers to improve amplification of the various members of the Hg^{II}-methylating community (Christensen et al. 2016) (Table 1). In sediments, these clade-specific primers were useful to detect Hg^{II}-methylating *δ-Proteobacteria* and *Archaea* but failed to detect Hg^{II}-methylating *Firmicutes* (Christensen et al. 2017).

Recent studies based on *hgcAB* biodiversity analyzed both *hgcA* (using one single pair of primers, Table 1) and 16S rRNA genes by high-throughput illumina sequencing techniques, allowing a deeper sequencing than cloning-sequencing and hence resulting in a higher number of OTUs. Results evidenced that microbial Hg^{II}-methylating community was composed of members of various clades, including SRB, FeRB, methanogens and syntrophs in temperate and boreal lake sediments (Bravo et al. 2018a) as well as in boreal forest soils (Xu et al. 2019). In boreal lakes, besides the identification of Hg^{II}-methylating methanogens and *Geobacteraceae*, authors further showed thanks to inhibition of sulfate reduction with molybdate that only 40% of MMHg was dependent on SRB (Bravo et al. 2018a). Another study performed in sediments impacted by a sewage treatment plant showed that Hg^{II} methylating *Geobacteraceae* seemed to have an important role in Hg^{II} methylation in sediments showing ferruginous conditions (Bravo et al. 2018b). Importantly, those studies suggested that the differences in the distributions of Hg^{II} methylating taxa among the different sites might derive primarily from different species of the same family having different niche requirements (Bravo et al. 2018a). In particular, the high relative abundance of phytoplankton-derived OM and the presence of specific strains of non-Hg^{II}-methylating bacteria involved in OM decomposition (e.g., *Rhizobiales*, *Fibrobacterales*, *Holophalages*, etc.) seems to be essential in creating a niche that promotes Hg^{II} methylation (Bravo et al. 2018a; Lei et al. 2019). Another study conducted in sulfate-impacted lakes combined cloning-sequencing of *hgcA* with metagenomics targeting *hgcA* gene and genes involved in other metabolic functions (Jones et al. 2019). This approach yielded after in silico assembly of reads with overlapping sequences in a relatively low number of contigs (27), but revealed a high occurrence of *hgcA* genes together with genes involved in sulfate-reduction and fermentation, but also that some abundant *hgcA+* microbes were related to uncultivated microbes, such as *Aminicenantes*, *Kiritimatiellaota*, *Spirochaetes*, as well as completely unidentified microbes. Data showed that potential methylators from uncultivated organisms occurred more abundantly than previously anticipated in these overlooked clades and that they can dominate the methylating community in certain circumstances (Jones et al. 2019).

A recent study conducted in rice paddy soils combining metagenomics illumina sequencing and long-read PacBio sequencing, which allows overcoming the inherent risk of short-reads chimeric assembly, revealed the dominance of *Geobacter* spp. for bacteria and *Methanoregula* spp. for *Archaea* (Liu et al. 2018b). These authors hypothesize a syntrophic interaction between both species and in addition reported a significant correlation between *Geobacter hgcA+* DNA relative abundance and MMHg concentration in soils (Liu et al. 2018b), supporting an important role of this genus for MMHg production in iron-rich paddy soils. A recent study in sediments collected in eutrophic lakes showing cyanobacteria blooms in China also reported a correlation between *Archae hgcA+* DNA relative abundance and MMHg concentration (Lei et al. 2019). However, several studies that tried to correlate the level of expression of *hgcA* mRNA with Hg^{II} methylation rates (Goñi-Urriza et al. 2015; Bravo et al. 2016; Christensen et al. 2019) were mostly unsuccessful. For example, in pure cultures of *Desulfovibrio dechloroacetivorans* BerOc1, the level of expression of *hgcA* was not correlated with Hg^{II} methylation rates (Goñi-Urriza et al. 2015). Similarity, in sediments collected in a river impacted by effluents from a chlor-alkali plant, data suggested that physico-chemistry varied significantly among reservoirs, while functional gene activities, including *hgcA*, were very similar and did not correlate with MMHg concentrations (Bravo et al. 2016). In contrast, in Hg-contaminated paddy soils, the *hgcAB* copy number increased with both increasing THg and MMHg concentrations (Vishnivetskaya et al. 2018).

Despite the differences in primer pairs used in the studies mentioned earlier (Table 1), until now, data globally suggested that in *hgcA+* *δ-Proteobacteria* communities are abundant in surface sediments, but in some sites *hgcA+* methanogens and other *hgcA+* uncultivated groups are prevalent (Christensen et al. 2017, 2019; Vishnivetskaya et al. 2018; Bravo et al. 2018a; Jones et al. 2019). Among *δ-Proteobacteria*, syntrophs and *Geobacter* spp. appear more prevalent in *hgcA+* community than previously expected. Moreover, other groups of *hgcA-* microbes seem to be of high importance for Hg^{II}-methylating species, certainly by providing some kind of dependence or mutualistic relationship in sediments (Bravo et al. 2018a; Liu et al. 2018b). For example, syntrophs have been shown to modulate Hg^{II} methylation of *hgcA* strains in controlled exposures (Yu et al. 2018). Syntrophy between methanogens or propionate utilizing syntrophs and SRB is hypothesized to enhance methylation in environments devoid of sulfate or where the type and concentration of energy sources are limiting.

Studies listed above focused in freshwaters and therefore the microorganisms processing Hg^{II} to MMHg in the ocean are still barely described. Podar et al. (2015) showed that *hgcAB* appeared to be abundant in marine sediments but they rarely found it in pelagic marine water column, as from 138 metagenome samples analyzed, only seven showed evidence of *hgcAB*. A recent analysis of 243 seawater metagenome samples from 68 different sites of the *Tara* Oceans revealed high abundances of *hgcAB* corresponding to taxonomic relatives of known Hg^{II} methylators from

Deltaproteobacteria, *Firmicutes*, and *Chloroflexi* across all oceans, with the exception of the Arctic that was not studied (Villar et al. 2019). More recently, Bowman et al. (2019), combining PCR amplification and shotgun metagenomics, searched the *hgcAB* gene cluster in Arctic Ocean seawater without success. Out of all the *hgcA*-like genes found in the queries of marine metagenomes, the *Nitrospina* phylum, a marine nitrite oxidizing bacteria abundant in oxygen-deficient zones, appeared to be widespread, predominant and likely a key player for MMHg production in the oxic subsurface waters of the global ocean (Villar et al. 2019), including the Arctic (Bowman et al. 2019) as well as Antarctic sea ice-brine-sea water interfaces (Gionfriddo et al. 2016). However, despite metagenomic evidence for the abundance of *Nitrospina* in the global ocean, the few cultured strains harboring a fused *hgcAB*-like gene (*Methanococcoides methylutens* and *Pyrococcus furiosus*) were unable to produce MMHg in experimental conditions (Podar et al. 2015; Gilmour et al. 2018). Moreover, there is yet no experimental or observational report on the expression of *hgcAB*-like genes in *Nitrospina* bacteria. As such, an experimental evidence of the Hg^{II} methylating capacity in *Nitrospina* is awaited to confirm their role as important Hg^{II} methylators in the global ocean. As MMHg has been detected in the water column of every ocean basin, except for the Indian Ocean (Bowman et al. 2019), it is crucial to unveil the role of the microorganisms involved in both MMHg and degradation in seawaters.

These results and observations reveal the diversity of Hg^{II}-methylating microbial communities' structure across ecosystems and point for the need of a thorough investigation of their functioning. Notably further work is necessary to better understand the contribution of overlooked microbial groups in Hg^{II} methylation, highlighted by the high proportion of unidentified OTUs found in recent studies concerning *hgcAB* biodiversity (Table 2). However, it would be useful to agree on a standardized protocol to conduct *hgcAB* biodiversity studies and have an *hgcAB* open-access library, as published studies are currently difficult to directly compare due to differences in methods, including primer pairs, alignment algorithms used and depth of sequencing.

Hg^{II} methylators are part of a complex microbial community

As described in the previous sections, current knowledge established that MMHg net production was linked to biotic and abiotic variables. Microorganisms behave differently from one system to another due to interactions with the physico-chemical variables but also with other organisms of their (micro)environment (Andersson et al. 2014; Bravo et al. 2018a). Studies with one strain can only describe the metabolism of this strain in a batch (Andersson et al. 2014), which is useful for a mechanistic understanding of its potential metabolism. However, it cannot be straightforwardly applied for environmental predictions because its metabolism is likely modified by the activity of other microbial groups and the ambient physico-chemistry. In this sense, one of the most insightful discoveries is the syntrophic Hg^{II} methylation recently described in both laboratory (Kerin

et al. 2006; Ranchou-Peyruse et al. 2009) and field studies (Yu et al. 2018). Syntrophy is just a "proof of concept" illustrating the complexity of microbial communities carrying out Hg^{II} methylation. It is also important to consider that within a microbial community, besides Hg^{II}, some bacteria carrying out the *merB* gene or other genes yet to be discovered, might demethylate MMHg (Barkay et al. 2003).

Electron donors are also essential for Hg^{II} methylation. Different microbial clades are involved in the anaerobic oxidation of OM from complex organic compounds generally that goes through several steps and processes (Gilmour et al. 2013; Bae et al. 2014). For example, an initial hydrolysis of large organic substances is followed by a fermentation of intermediates into smaller organic molecules, such as lactate, propionate, butyrate, acetate, and formate, as well as CO₂ and H₂. These fermentation products might then be used as electron donors for *Geobacterales*, *Desulfovibrionales*, and *Syntrophobacterales* known to host Hg^{II} methylators. Hg^{II} methylators thus likely rely on other microorganisms involved in the degradation of large organic compounds. Furthermore, it was also demonstrated that the specific metabolism of one strain may provoke a new metabolism (unknown) in another strain; this is called the Quorum sensing (Lovley and Chapelle 1995). It is therefore very important to better tackle the complexity of microbial communities and describe the compendium of metabolic processes that can affect directly or indirectly Hg^{II} methylation.

Physico-chemistry plays a pivotal role in Hg^{II} methylation

Besides the presence and diversity of Hg^{II} methylating microbes, Hg^{II} methylation depends on the amount of Hg^{II} bioavailable for methylation (Schaefer et al. 2011; Jonsson et al. 2014), which is determined by chemical speciation of Hg^{II}, solubility of the Hg-S particles (Hsu-Kim et al. 2013; Liem-Nguyen et al. 2017) as well as the availability of electron donors and acceptors for Hg^{II} methylating microorganisms (Desrochers et al. 2015). Hg^{II} methylation is a bio-physico-chemical conundrum because both the amount of Hg^{II} available for methylation and the activity of microorganisms involved in the process are determined by multiple physico-chemical variables such as sulfur (Skylberg et al. 2003; Drott et al. 2007), iron (Bravo et al. 2015) and OM concentration and speciation (Schartup et al. 2013; Bravo et al. 2017) as well as Eh, pH, nutrient availability, and temperature (Ullrich et al. 2001; Paranjape and Hall 2017) (Fig. 1).

Variables affecting Hg^{II} chemical speciation

Salinity, sulfur, iron, and OM affect Hg^{II} chemical speciation. Sea salt anions may also affect Hg^{II} speciation and/or methylation in estuarine and marine environments. In marine waters, Hg^{II} forms compounds with chlorine (HgCl₃⁻ and HgCl₄²⁻) to a greater extent than oxides, that are in turn formed in freshwaters (Mason and Fitzgerald 1993). A lower Hg^{II} methylating activity in marine and estuarine sediments than in freshwater sediments




Fig. 1. Conceptual summary of the biological and chemical interplays affecting Hg^{II} methylation in the environment. Orange boxes and arrows refer to geochemical variables directly affecting microbial activity and Hg speciation. Purple boxes refer to Hg chemical forms. The red arrow indicates the transformation of Hg^{II} to CH_3Hg^+ . Blue refers to a compendium of metabolic processes occurring in the environment, among these processes Hg^{II} methylation carried out by the *hgcAB* gene cluster.

has been attributed to the formation of charged chloride but also sulfide complexes, that undergo a slower methylation processes than other Hg^{II} forms (Gårdfeldt et al. 2003; Jonsson et al. 2012; Gworek et al. 2016).

In the context of the biological and chemical interplays controlling Hg^{II} methylation, sulfur plays a central role by directly affecting Hg^{II} speciation and solubility (Jonsson et al. 2012; Graham et al. 2013; Hsu-Kim et al. 2013; Liem-Nguyen et al. 2017) and consequently its bioavailability (Schaefer and Morel 2009; Chiasson-Gould et al. 2014; Schartup et al. 2015; Mazrui et al. 2016). Reactions between Hg^{II} and sulfide control the formation of the solid phase metacinnabar, $\beta\text{-HgS(s)}$ but also aqueous complexes such as $\text{Hg}(\text{SH})_2^0$, HgS_2H^- , and HgS_2^{2-} or polysulfides $\text{HgS}_n\text{SH}^-(\text{aq})$ ($n = 4-6$) (Liem-Nguyen et al. 2017). Elevated sulfide concentrations, eventually limits Hg^{II} bioavailability for methylation (Drott et al. 2007; Hsu-Kim et al. 2013; Bigham et al. 2017). Conversely low sulfide concentrations might enhance Hg^{II} methylation processes (Benoit et al. 1999; Graham et al. 2012). Natural OM often contain thiols, which sulfhydryl group has a high capacity to complex Hg^{II} and MMHg (Skylberg et al. 2006; Skylberg 2008). Iron plays also a key role on Hg^{II} methylation as its reduced form Fe^{2+} can scavenge sulfide, form stable iron-sulfur compounds (FeS , Fe_3S_4 , or FeS_2) and then might let Hg^{II} bioavailable for methylation (i.e., ferruginous conditions; Bravo et al. 2015). Each of these Hg^{II} complexes, i.e., inorganic sulfides, polysulfides, and OM in aqueous, solid, and adsorbed phases, show different reactivity in the environment. For example, Hg^{II} methylation rate constants in estuarine sediments spanned over two orders of magnitude depending on the chemical form: metacinnabar ($\beta\text{-HgS(s)}$) < cinnabar ($\alpha\text{-HgS(s)}$) < Hg^{II} reacted with mackinawite (FeS-

Hg^{II}) < Hg^{II} bonded to natural OM ($\text{NOM-Hg}^{\text{II}}$) < $(\text{Hg}(\text{NO}_3)_2(\text{aq}))$ that is a typical aqueous tracer (Jonsson et al. 2012).

The effect of natural OM on the sediment and pore water-partitioning coefficient for Hg^{II} (Hammerschmidt et al. 2008; Liem-Nguyen et al. 2016) needs still to be determined. Under low OM and porewater sulfide concentrations, Hg^{II} partitioning coefficient becomes a major factor for methylation (Mitchell and Gilmour 2008; Hollweg et al. 2009). Some studies have shown that increases in natural OM concentrations might also raise partitioning coefficients in sediment and pore water and have shown to decrease Hg^{II} concentration in the pore water, and thus its bioavailability for uptake by methylating microorganisms (Liem-Nguyen et al. 2016). In contrast, other studies concluded that OM content did not explain variations in Hg^{II} partitioning, most likely only a limited fraction of OM was relevant for Hg complexation or because it is also possible that the composition, rather than amount, of OM controls Hg^{II} partitioning (Schartup et al. 2013).

High concentrations of OM decreased Hg^{II} bioavailability in laboratory experiments (Chiasson-Gould et al. 2014) and in marine sediments (Hammerschmidt et al. 2008). Moreover, OM might also be important in determining Hg^{II} bioavailability by stabilizing HgS particles at nanoscale that can be methylated by anaerobic bacteria (Graham et al. 2013; Hsu-Kim et al. 2013). To progress further in the understanding of Hg^{II} methylation in future research, it is of the utmost importance to measure and model Hg^{II} chemical speciation, which is ultimately determining the Hg^{II} availability for methylating bacteria (Fig. 1, violet box).

Variables affecting microbial activity

Sulfur, iron, OM, redox, pH, nutrients, and temperature affect microbial activity. Both sulfur and iron have oxidized forms (SO_4^{2-} and Fe^{3+} , respectively) that can serve as electron acceptor for some Hg^{II} methylating bacteria (e.g., SRB and FeRB). Natural OM, besides its strong capacity to bind Hg^{II} and influence Hg^{II} speciation, controls microbial activity and Hg^{II} methylation (Graham et al. 2013; Hsu-Kim et al. 2013). The molecular composition of OM shows a central role in controlling Hg^{II} methylation (Bravo et al. 2017), notably phytoplankton derived OM and fresh humic matter were associated with both high bacterial activity and Hg^{II} methylation rates in sediments (Graham et al. 2013; Schartup et al. 2013; Mazrui et al. 2016; Bravo et al. 2017; Christensen et al. 2017; Herrero Ortega et al. 2018). Also, an increase in nutrients, associated to an enhanced algae biomass production, increased MMHg formation in sediments (Bravo et al. 2017; Herrero Ortega et al. 2018). Changes in redox conditions might also affect iron, sulfur and Hg^{II} speciation (Liem-Nguyen et al. 2016), and the activity of the microorganisms (Grégoire and Poulain 2018) and consequently Hg^{II} methylation (Fig. 1, orange boxes and arrows). Furthermore, Hg^{II} methylation strongly depends on the activity of the whole microbial community (Weber et al. 2006; DeAngelis et al. 2010). Therefore, all the physico-chemical variables causing increased microbial activity in sediments, such as temperature (Gudasz et al. 2010), might indirectly lead to enhanced Hg^{II} methylation (Bravo et al. 2017; Dijkstra et al. 2011).

From all the studies mentioned earlier, we can conclude that both microbial activity and Hg^{II} chemical speciation control the formation of MMHg in the environment. Therefore, Hg cycling and in particular, the methylation of Hg^{II} is an intricate process regulated by both physico-chemical and biological constraints that needs holistic approaches to be fully understood.

Knowledge gaps and uncertainties

A high number of unidentified putative Hg^{II} methylators

The complexity of microbial communities and its implications for Hg cycling is one of the main current challenges. First biodiversity studies used 16S rRNA gene to investigate the diversity of Hg^{II} methylating microbial communities. But biodiversity of this gene cannot provide reliable and robust identification of Hg^{II} methylator diversity and abundance, because *hgcAB*+ strains are too rare (<1%) and not well identified in 16S rRNA databases (<300 species) (Miller and Bassler 2001; Christensen et al. 2019). Recent studies targeting *hgcAB* are interesting, but these molecular approaches entail several limits. First, as for all microbial biodiversity studies, the DNA extraction protocols need to be well planned to ensure clean subsampling avoiding contamination and may need to be optimized for the efficient recovery of DNA and the elimination of potential inhibitors for PCR and/or sequencing technologies. Indeed, DNA extraction and PCR amplification are known to be prone to artifacts due to any combination of high primer mismatch, low abundance, and/or low

DNA extraction efficiency that can significantly affect results (Bravo et al. 2018a; Epp et al. 2019). Although claimed as “universal,” primers inherently show preferences and limitations in covering all species equally among different environmental samples. More specifically, currently available primer pairs for the gene *hgcA* are predicted to cover 84% (Schaefer et al. 2014) and 94% (Christensen et al. 2016) of the whole biodiversity by PCR-based approaches. However, within amplified sequences of *hgcA* a significant proportion cannot be identified above the clade level because of (1) the lack of identified organisms in the databases, and (2) the low conservation of *hgcA* gene that is not ideal for biodiversity studies. Consequently, a significant proportion of OTUs are attributed to unidentified species. Indeed, identification of *hgcA* sequences from short reads is made by aligning metagenomics data with a set of known gene sequences isolated from cultivated strains. This approach is prone to inaccuracies, especially if the data are evaluated down to the genus level. For example, in recent studies 62% (Bravo et al. 2018b) and 57% (Bravo et al. 2018a) of the identified OTUs could be taxonomically assigned at the order level. More in detail, different algorithms are available to identify OTUs e.g., simple sequence alignment-based algorithms (e.g., Basic Local Alignment Search Tool [BLAST]) or profile hidden Markov model (HMM)-based searches, which are expected to be more sensitive and accurate in identifying homologs. Nonetheless, the HMM model is satisfying for *hgcA* but not for *hgcB* that cannot be confidently differentiates from other ferredoxin-encoding genes due to homology (Christensen et al. 2019). Besides, both primers and algorithms are developed on sequences from cultivated strains and consequently identify those species with a higher efficiency (Podar et al. 2015; Bravo et al. 2018b; Liu et al. 2018b). Moreover, only few hundred sequenced genomes of methylating strains are currently available in databases. Although this number is increasing regularly, databases still do not include reliably microorganisms that cannot be grown in the laboratory. Besides we do not have information on the rates and methylation capabilities of these uncultivated populations and thus we cannot affirm that the genes identified code for MMHg production in these organisms. Further, we cannot evaluate how Hg methylation rates in these organisms compare to other methylators in culture. In sum, available sequences are far from being sufficient to confidently identify all OTUs homologs of *hgcA* and might not allow identifying unknown microbes. This is a current inherent technical limit that has to be considered when interpreting data.

However, studies conducted on DNA do not reflect the activity of the protein or the enzyme performing the process, but only the possibility that a strain could sometime methylate Hg^{II} . Although the use of RNA should theoretically provide more detailed information on the activity of the organisms involved in Hg^{II} , up to now the abundance of *hgcAB* transcripts could not be correlated to Hg^{II} methylation or MMHg concentration (Goñi-Urriza et al. 2015; Bravo et al. 2016; Vishnivetskaya et al. 2018; Christensen et al. 2019), except in one study (Ledeker and De Long 2013). The precise factors regulating *hgcA* gene and protein

expression are not identified yet but it seems that *hgcA* gene could be constitutively expressed or regulated by carbon, metabolism, but Hg does not appear to be a key regulator (Goñi-Urriza et al. 2015; Christensen et al. 2019). Moreover, the correlation between mRNA and protein abundance is not necessarily linear as posttranslational regulations can take place. In addition to the factors regulating *hgcA* gene expression, it is important to consider that net MMHg production also depends on MMHg demethylation activity. Therefore, the absence of correlation between *hgcA* gene and MMHg concentration is not surprising and highlights the complexity of predicting MMHg concentration dynamics in the environment.

Currently, the scientific community lacks a straightforward method to directly measure the activity of *hgcAB* at the protein level. New metaproteomic approaches are currently developed and could in the near future allow a direct analysis of *hgcA* protein abundance in the environment (Meier et al. 2019), but this approach has been tested only once to study *hgcA* biodiversity yet (Christensen et al. 2019). Metagenomics, metatranscriptomics, and metaproteomics offer a deeper sequencing and a wealth of information than earlier but they are cost and time-consuming to conduct analysis of hundreds of samples. This highlights that we still lack of a simple and accurate method to identify Hg^{II}-methylating microbial activity in environmental samples.

Another issue is that there is currently no medium or protocol to reliably culture a whole microbial community including FeRB, SRB, firmicutes, *Clostridia*, and so forth. Even studies with intact sediments in controlled experimental conditions can only approximate interactions of microbial community in situ and need to be confirmed in the field. For this reason, it is difficult to study and explain interactions of microbial communities through controlled experiments using a single strain. For example, microbial strains showing maximum methylation rates in the laboratory may not be as active in complex consortia under real field conditions. Metagenomics, metatranscriptomics, and metaproteomics could help to progress in the elucidation of the biological metabolism behind Hg^{II} methylation in the field. Instead of identifying different taxa, these approaches combine genes coding for all metabolic pathways of the different taxa to describe the potential function of the whole community (metagenomics) or its activity in one compartment (sediment, water, soil, etc.) at the moment of the sampling (metatranscriptomics and metaproteomics). Nonetheless, available metagenomics approaches have seldom identified *hgcA* sequences (i.e., 63 out of 203 metagenome projects revealed *hgcA* occurrence; Podar et al. 2015; and 77 out of 243, Villar et al. 2019) supporting that new study should be conducted in environments relevant for Hg^{II} methylation. Furthermore, metatranscriptomics and metaproteomics are still rarely applied, most likely because they are expensive and time-consuming. However, although we believe that these approaches are promising and their use will increase in coming years, they will need to be coupled with a detailed analysis of environmental conditions (e.g., physico-chemistry analysis) to be fully informative.

Are we measuring realistic Hg^{II} methylation rates?

Last but not least, besides limitations to directly measure biological process responsible for Hg^{II} methylation, there are several pitfalls on the methods used to determine Hg^{II} methylation at environmentally relevant Hg^{II} concentrations. Several approaches have been used to estimate MMHg formation: using labeled Hg^{II} forms with radio-isotopes (Goñi-Urriza et al. 2015; Bravo et al. 2016), stable isotopes (Ramlal et al. 1986), by measuring the percentage of total Hg and MMHg (% MMHg) (Hintelmann et al. 2000) or the change in MMHg concentration over time (Drott et al. 2008). The amendment of standards enriched in Hg^{II} stable isotope tracers to environmental samples is now widely used in different laboratories because they allow determining Hg^{II} methylation rates and MMHg demethylation rates simultaneously. However, it is known that the geochemical form of the Hg^{II} isotope used as tracer determines its reactivity (i.e., methylation rate) in the environment. For example, Hg^{II} methylation rate constants in estuarine sediments spanned over two orders of magnitude from metacinnabar to a typical aqueous tracer such as Hg(NO₃)₂(aq) (Jonsson et al. 2012). Therefore, Hg^{II} methylation rate constants measured using a highly available tracer might result in an overestimation of the in situ Hg^{II} methylation rate, when the tracer is a poor representative of the indigenous Hg^{II} chemical forms. The %MMHg and the change of MMHg concentration overtime represent the net MMHg, accounting for MMHg degradation processes and while it might be useful to predict MMHg concentrations in the environment, its use is limited to provide mechanistic understanding of Hg^{II} methylation processes.

From this literature review, we observe that after 50 yr of efforts to study Hg^{II} methylation through the world, knowledge has greatly increased, but there are still many aspects within the bio-physico-chemistry of MMHg formation that need to be unveiled in the natural environment because: (1) there is a lack of techniques and methods for the measurements of the different Hg^{II} chemical species available for methylation; (2) the physico-chemical factors affecting Hg^{II} chemistry are still only partly understood; (3) the biological mechanisms involved in the whole bio-physico-chemical process of MMHg net production remains to be understood and described more accurately.

How to study methylmercury formation in future research?

Besides current limits inherent to used analysis described earlier, main questions lacking a clear answer concerning the behavior of Hg in environmental systems include: What are all the Hg^{II} chemical species available for methylation? How diverse are *hgcA*⁺ species? How does microbial community consortia impact the amount and speciation of Hg^{II} available for methylation? And what is the impact of the *hgcA*⁻




Fig. 2. Schematic representation of the proposed conceptual iterative strategy for studying Hg^{II} methylation.

microbial species on the activity of Hg^{II} methylators? Do techniques exist to tackle these questions?

Laboratory studies are essential to identify mechanisms driving Hg^{II} methylation but need to include more realistic scenarios and/or be validated under field conditions. Currently, in situ studies to elucidate the biodiversity of Hg^{II} methylators and determine the drivers of their activity need to be conducted more widely and in more diverse environments. For in situ studies, Hg^{II} speciation and Hg^{II} rates need to be determined, and accompanied by the study of the factors controlling the activity of Hg^{II} methylating microorganisms. Determination of the methylating activity of natural microbial assemblages in relation to sediment characteristics, specific environmental conditions and the level of Hg contamination needs to be undertaken to validate laboratory-based measurements and to improve our understanding of Hg cycling in contaminated environments (Fig. 2). In this context, interactions between microbial communities and the physico-chemistry are key to predict Hg^{II} methylation as several studies pointed out that physico-chemistry rather than the microbial community structure were determining Hg^{II} methylation rate constants in lake sediments (Bravo et al. 2016, 2017, 2018a; Liu et al. 2018b). This is likely true but it should not be forgotten that the local physico-chemistry is also the result of the microbial activity (Bravo et al. 2017, 2018b).

We can use existing and future new molecular biology and chemistry techniques to study Hg^{II} methylation, but studies

have now to put more effort on combining metagenomics, metatranscriptomics, and metaproteomics approaches with experiments on bacterial isolates and also complex matrices (i.e., sediments, marine waters, etc.) to explain more in detail environmental mechanisms (Fig. 2). Last, we should keep in




Fig. 3. Proposed model to quantify MMHg in freshwater systems. The model considers formation (k_m , Hg^{II} methylation) and degradation (k_d , MMHg demethylation) and the inputs and outputs of MMHg. Processes are represented with orange arrows, known functional genes are shown in brackets. Transport of MMHg to the system or out the system is represented by blue arrows.

mind that MMHg in aquatic systems is the net result of different processes: (1) formation (Hg^{II} methylation), (2) degradation (MMHg demethylation), and (3) the inputs and outputs of the system (Fig. 3). Therefore, when studying Hg^{II} methylation, we target only a limited part of the whole Hg cycling determining MMHg concentrations in the environment. Future studies need to be invested in studying concomitantly the drivers of MMHg methylation and MMHg demethylation mechanisms in situ as well as the transport of MMHg within the aquatic network, which is indeed highly controlled by hydrological processes (Fig. 3).

References

- Achá, D., H. Hintelmann, and J. Yee. 2011. Importance of sulfate reducing bacteria in mercury methylation and demethylation in periphyton from Bolivian Amazon region. *Chemosphere* **82**: 911–916. doi:10.1016/j.chemosphere.2010.10.050
- Achá, D., H. Hintelmann, and C. A. Pabón. 2012. Sulfate-reducing bacteria and mercury methylation in the water column of the Lake 658 of the experimental lake area. *Geomicrobiol. J.* **29**: 667–674. doi:10.1080/01490451.2011.606289
- Åkerblom, S., A. Bignert, M. Meili, L. Sonesten, and M. Sundbom. 2014. Half a century of changing mercury levels in Swedish freshwater fish. *Ambio* **43**: 91–103. doi:10.1007/s13280-014-0564-1
- Amos, H. M., D. J. Jacob, D. G. Streets, and E. M. Sunderland. 2013. Legacy impacts of all-time anthropogenic emissions on the global mercury cycle. *Global Biogeochem. Cycles* **27**: 410–421. doi:10.1002/gbc.20040
- Andersson, M. G. I., M. Berga, E. S. Lindström, and S. Langenheder. 2014. The spatial structure of bacterial communities is influenced by historical environmental conditions. *Ecology* **95**: 1134–1140. doi:10.1890/13-1300.1
- Bae, H. S., F. E. Dierberg, and A. Ogram. 2014. Syntrophs dominate sequences associated with the mercury methylation-related gene *hgcA* in the water conservation areas of the Florida Everglades. *Appl. Environ. Microbiol.* **80**: 6517–6526. doi:10.1128/AEM.01666-14
- Barkay, T., S. M. Miller, and A. O. Summers. 2003. Bacterial mercury resistance from atoms to ecosystems. *FEMS Microbiol. Rev.* **27**: 355–384. doi:10.1016/S0168-6445(03)00046-9
- Benoit, J. M., C. C. Gilmour, R. P. Mason, and A. Heyes. 1999. Sulfide controls on mercury speciation and bioavailability to methylating bacteria in sediment pore waters. *Environ. Sci. Technol.* **33**: 951–957. doi:10.1021/es9808200
- Bigham, G. N., K. J. Murray, Y. Masue-Slowey, and E. A. Henry. 2017. Biogeochemical controls on methylmercury in soils and sediments: Implications for site management. *Integr. Environ. Assess. Manag.* **13**: 249–263. doi:10.1002/ieam.1822
- Bjørklund, G., M. Dadar, J. Mutter, and J. Aaseth. 2017. The toxicology of mercury: current research and emerging trends. *Environ. Res.* **159**: 545–554. doi:10.1016/j.envres.2017.08.051
- Bloom, N. S. 1992. On the chemical form of mercury in edible fish and marine invertebrate tissue. *Can. J. Fish. Aquat. Sci.* **49**: 1010–1017. doi:10.1139/f92-113
- Bouchet, S., and others. 2013. MMHg production and export from intertidal sediments to the water column of a tidal lagoon (Arcachon Bay, France). *Biogeochemistry* **114**: 341–358. doi:10.1007/s10533-012-9815-z
- Bouchet, S., and others. 2018. Linking microbial activities and low-molecular-weight thiols to Hg methylation in biofilms and periphyton from high-altitude tropical lakes in the Bolivian Altiplano. *Environ. Sci. Technol.* **52**: 9758–9767. doi:10.1021/acs.est.8b01885
- Bowman, K. L., R. E. Collins, A. M. Agather, C. H. Lamborg, C. R. Hammerschmidt, D. Kaul, C. L. Dupont, G. A. Christensen, and D. A. Elias. 2019. Distribution of mercury-cycling genes in the Arctic and equatorial Pacific oceans and their relationship to mercury speciation. *Limnol. Oceanogr.*: 1–11. doi:10.1002/lno.11310
- Bravo, A. G., C. Cosio, D. Amouroux, J. Zopfi, P.-A. Chevalley, J. E. Spangenberg, V.-G. Ungureanu, and J. Dominik. 2014. Extremely elevated methyl mercury levels in water, sediment and organisms in a Romanian reservoir affected by release of mercury from a chlor-alkali plant. *Water Res.* **49**: 391–405. doi:10.1016/j.watres.2013.10.024
- Bravo, A. G., S. Bouchet, S. Guédron, D. Amouroux, J. Dominik, and J. Zopfi. 2015. High methylmercury production under ferruginous conditions in sediments impacted by sewage treatment plant discharges. *Water Res.* **80**: 245–255. doi:10.1016/j.watres.2015.04.039
- Bravo, A. G., J. L. Loizeau, P. Dranguet, S. Makri, E. Björn, V. G. Ungureanu, V. I. Slaveykova, and C. Cosio. 2016. Persistent Hg contamination and occurrence of Hg-methylating transcript (*hgcA*) downstream of a chlor-alkali plant in the Olt River (Romania). *Environ. Sci. Pollut. Res.* **23**: 10529–10541. doi:10.1007/s11356-015-5906-4
- Bravo, A. G., S. Bouchet, J. Tolu, E. Björn, A. Mateos-Rivera, and S. Bertilsson. 2017. Molecular composition of organic matter controls methylmercury formation in boreal lakes. *Nat. Commun.* **8**: 14255. doi:10.1038/ncomms14255
- Bravo, A. G., and others. 2018a. Methanogens and iron-reducing bacteria: the overlooked members of mercury-methylating microbial communities in boreal lakes. *Appl. Environ. Microbiol.* **84**: 0–16. doi:10.1128/AEM.01774-18
- Bravo, A. G., J. Zopfi, M. Buck, J. Xu, S. Bertilsson, J. K. Schaefer, J. Poté, and C. Cosio. 2018b. Geobacteraceae are important members of mercury-methylating microbial communities of sediments impacted by waste water releases. *ISME J.* **12**: 802–812. doi:10.1038/s41396-017-0007-7
- Celo, V., D. R. S. Lean, and S. L. Scott. 2006. Abiotic methylation of mercury in the aquatic environment. *Sci. Total*

- Environ. **368**: 126–137. doi:[10.1016/j.scitotenv.2005.09.043](https://doi.org/10.1016/j.scitotenv.2005.09.043)
- Chételat, J., B. Braune, J. Stow, and S. Tomlinson. 2015. Special issue on mercury in Canada's north: summary and recommendations for future research. *Sci. Total Environ.* **509–510**: 260–262. doi:[10.1016/j.scitotenv.2014.06.063](https://doi.org/10.1016/j.scitotenv.2014.06.063)
- Chiasson-Gould, S. A., J. M. Blais, and A. J. Poulain. 2014. Dissolved organic matter kinetically controls mercury bioavailability to bacteria. *Environ. Sci. Technol.* **48**: 3153–3161. doi:[10.1021/es4038484](https://doi.org/10.1021/es4038484)
- Christensen, G. A., and others. 2016. Development and validation of broad-range qualitative and clade-specific quantitative molecular probes for assessing mercury methylation in the environment. *Appl. Environ. Microbiol.* **82**: 6068–6078. doi:[10.1128/AEM.01271-16](https://doi.org/10.1128/AEM.01271-16)
- Christensen, G. A., and others. 2017. Carbon amendments alter microbial community structure and net mercury methylation potential in sediments. *Appl. Environ. Microbiol.* **84**: 1–14. doi:[10.1128/AEM.01049-17](https://doi.org/10.1128/AEM.01049-17)
- Christensen, G. A., A. C. Somenahally, J. G. Moberly, C. M. Miller, A. J. King, C. C. Gilmour, S. D. Brown, M. Podar, C. C. Brandt, S. C. Brooks, A. V. Palumbo, J. D. Wall, and D. A. Elias, (2018) Carbon amendments alter microbial community structure and net mercury methylation potential in sediments. *Applied and Environmental Microbiology* **84**.
- Christensen, G. A., and others. 2019. Determining the reliability of measuring mercury cycling gene abundance with correlations with mercury and methylmercury concentrations. *Environ. Sci. Technol.* **53**: 8649–8663. doi:[10.1021/acs.est.8b06389](https://doi.org/10.1021/acs.est.8b06389)
- Clarkson, T. W., and L. Magos. 2006. The toxicology of mercury and its chemical compounds. *Crit. Rev. Toxicol.* **36**: 609–662. doi:[10.1080/10408440600845619](https://doi.org/10.1080/10408440600845619)
- Cleckner, L. B., C. C. Gilmour, J. P. Hurley, and D. P. Krabbenhoft. 1999. Mercury methylation in periphyton of the Florida Everglades. *Limnol. Oceanogr.* **44**: 1815–1825. doi:[10.4319/lo.1999.44.7.1815](https://doi.org/10.4319/lo.1999.44.7.1815)
- Compeau, G., and R. Bartha. 1984. Methylation and demethylation of mercury under controlled redox, pH and salinity conditions. *Appl. Environ. Microbiol.* **48**: 1203–1207.
- Compeau, G. C., and R. Bartha. 1985. Sulfate-reducing bacteria : principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* **50**: 498–502.
- Cossa, D., B. Averty, and N. Pirrone. 2009. The origin of methylmercury in open Mediterranean waters. *Limnol. Oceanogr.* **54**: 837–844. doi:[10.4319/lo.2009.54.3.0837](https://doi.org/10.4319/lo.2009.54.3.0837)
- DeAngelis, K. M., W. L. Silver, A. W. Thompson, and M. K. Firestone. 2010. Microbial communities acclimate to recurring changes in soil redox potential status. *Environ. Microbiol.* **12**: 3137–3149. doi:[10.1111/j.1462-2920.2010.02286.x](https://doi.org/10.1111/j.1462-2920.2010.02286.x)
- Depew, D., and others. 2013. An overview of mercury concentrations in freshwater fish species: a national fish mercury dataset for Canada. *Can. J. Fish Aquat. Sci.* **451**: 436–451. doi:[10.1139/cjfas-2012-033](https://doi.org/10.1139/cjfas-2012-033)
- Desrochers, K. A. N., K. M. A. Paulson, C. J. Ptacek, D. W. Blowes, and W. D. Gould. 2015. Effect of electron donor to sulfate ratio on mercury methylation in floodplain sediments under saturated flow conditions. *Geomicrobiol. J.* **32**: 924–933. doi:[10.1080/01490451.2015.1035818](https://doi.org/10.1080/01490451.2015.1035818)
- Devereux, R., M. R. Winfrey, J. Winfrey, and D. A. Stahl. 1996. Depth profile of sulfate-reducing bacterial ribosomal RNA and mercury methylation in an estuarine sediment. *FEMS Microbiol. Ecol.* **20**: 23–31. doi:[10.1111/j.1574-6941.1996.tb00301.x](https://doi.org/10.1111/j.1574-6941.1996.tb00301.x)
- Dijkstra, P., S. C. Thomas, P. L. Heinrich, G. W. Koch, E. Schwartz, and B. A. Hungate. 2011. Effect of temperature on metabolic activity of intact microbial communities: evidence for altered metabolic pathway activity but not for increased maintenance respiration and reduced carbon use efficiency. *Soil Biol. Biochem.* **43**: 2023–2031. doi:[10.1016/j.soilbio.2011.05.018](https://doi.org/10.1016/j.soilbio.2011.05.018)
- Dranguet, P., S. Le Faucheur, and V. I. Slaveykova. 2017. Mercury bioavailability, transformations, and effects on freshwater biofilms. *Environ. Toxicol. Chem.* **36**: 3194–3205. doi:[10.1002/etc.3934](https://doi.org/10.1002/etc.3934)
- Blum, J. D., B. N. Popp, J. C. Drazen, M. W. Johnson, and C. Anela Choy. 2013. Methylmercury production below the mixed layer in the North Pacific Ocean. *Nat. Geosci.* **6**: 879–884. doi:[10.1038/ngeo1918](https://doi.org/10.1038/ngeo1918)
- Drott, A., L. Lambertsson, E. Björn, and U. Skyllberg. 2007. Importance of dissolved neutral mercury sulfides for methyl mercury production in contaminated sediments. *Environ. Sci. Technol.* **41**: 2270–2276. doi:[10.1021/es061724z](https://doi.org/10.1021/es061724z)
- Drott, A., L. Lambertsson, E. Björn, and U. Skyllberg. 2008. Do potential methylation rates reflect accumulated methyl mercury in contaminated sediments? *Environ. Sci. Technol.* **42**: 153–158. doi:[10.1021/es0715851](https://doi.org/10.1021/es0715851)
- Du, H., M. Ma, T. Sun, X. Dai, C. Yang, F. Luo, D. Wang, and Y. Igarashi. 2017. Mercury-methylating genes *dsrB* and *hgcA* in soils/sediments of the three gorges reservoir. *Environ. Sci. Pollut. Res.* **24**: 5001–5011. doi:[10.1007/s11356-016-8213-9](https://doi.org/10.1007/s11356-016-8213-9)
- Du, H., M. Ma, Y. Igarashi, and D. Wang. 2019. Biotic and abiotic degradation of methylmercury in aquatic ecosystems: A review. *Bull. Environ. Contam. Toxicol.* **102**: 605–611. doi:[10.1007/s00128-018-2530-2](https://doi.org/10.1007/s00128-018-2530-2)
- Eagles-Smith, C. A., and others. 2016. Spatial and temporal patterns of mercury concentrations in freshwater fish across the Western United States and Canada. *Sci. Total Environ.* **568**: 1171–1184. doi:[10.1016/j.scitotenv.2016.03.229](https://doi.org/10.1016/j.scitotenv.2016.03.229)
- Eckley, C. S., and H. Hintelmann. 2006. Determination of mercury methylation potentials in the water column of lakes across Canada. *Sci. Total Environ.* **368**: 111–125. doi:[10.1016/j.scitotenv.2005.09.042](https://doi.org/10.1016/j.scitotenv.2005.09.042)
- Eckley, C. S., C. J. Watras, H. Hintelmann, K. Morrison, A. D. Kent, and O. Regnell. 2005. Mercury methylation in the hypolimnetic waters of lakes with and without connection

- to wetlands in northern Wisconsin. *Can. J. Fish. Aquat. Sci.* **411**: 400–411. doi:10.1139/F04-205
- Eklöf, K., J. Fölster, L. Sonesten, and K. Bishop. 2012. Spatial and temporal variation of THg concentrations in run-off water from 19 boreal catchments, 2000–2010. *Environ. Pollut.* **164**: 102–109. doi:10.1016/j.envpol.2012.01.024
- Eklöf, K., K. Bishop, S. Bertilsson, E. Björn, M. Buck, U. Skyllberg, O. A. Osman, R. M. Kronberg, and A. G. Bravo. 2018. Formation of mercury methylation hotspots as a consequence of forestry operations. *Sci. Total Environ.* **613–614**: 1069–1078. doi:10.1016/j.scitotenv.2017.09.151
- Epp, L. S., H. H. Zimmermann, and K. R. Stoof-Leichsenring. 2019. Sampling and extraction of ancient DNA from sediments, p. 31–44. *In* Ancient DNA: Methods and protocols, methods in molecular biology. Springer doi:10.1007/978-1-4939-9176-1_5
- Fernández-Gómez, C., A. Drott, E. Björn, S. Díez, J. M. Bayona, S. Tesfalidet, A. Lindfors, and U. Skyllberg. 2013. Towards universal wavelength-specific photodegradation rate constants for methyl mercury in humic waters, exemplified by a boreal lake-wetland gradient. *Environ. Sci. Technol.* **47**: 6279–6287. doi:10.1021/es400373s
- Fleck, J. A., and others. 2015. Mercury and methylmercury in aquatic sediment across western North America. *Sci. Total Environ.* **568**: 727–738. doi:10.1016/j.scitotenv.2016.03.044
- Fleming, E. J., E. E. Mack, P. G. Green, and D. C. Nelson. 2006. Mercury methylation from unexpected sources: molybdate-inhibited freshwater sediments and an iron-reducing bacterium. *Appl. Environ. Microbiol.* **72**: 457–464. doi:10.1128/AEM.72.1.457
- Gårdfeldt, K., and others. 2003. Evasion of mercury from coastal and open waters of the Atlantic Ocean and the Mediterranean Sea. *Atmos. Environ.* **37**: 73–84. doi:10.1016/S1352-2310(03)00238-3
- Gascón Díez, E., J. L. Loizeau, C. Cosio, S. Bouchet, T. Adatte, D. Amouroux, and A. G. Bravo. 2016. Role of settling particles on mercury methylation in the oxic water column of freshwater systems. *Environ. Sci. Technol.* **50**: 11672–11679. doi:10.1021/acs.est.6b03260
- Gilmour, C. C., E. A. Henry, and R. Mitchell. 1992. Sulfate stimulation of mercury methylation in freshwater sediments. *Environ. Sci. Technol.* **26**: 2281–2287. doi:10.1021/es00035a029
- Gilmour, C. C., and others. 2013. Mercury methylation by novel microorganisms from new environments. *Environ. Sci. Technol.* **47**: 11810–11820. doi:10.1021/es403075t
- Gilmour, C. C., A. L. Bullock, A. Mcburney, and M. Podar. 2018. Robust mercury methylation across diverse methanogenic archaea. *mBio* **9**: 1–13. doi:10.1128/mBio.02403-17
- Gionfriddo, C. M., and others. 2016. Microbial mercury methylation in Antarctic Sea ice. *Nat. Microbiol.* **1**: 16127. doi:10.1038/nmicrobiol.2016.127
- Goñi-Urriza, M., Y. Corsellis, L. Lancelleur, E. Tessier, J. Gury, M. Monperrus, and R. Guyoneaud. 2015. Relationships between bacterial energetic metabolism, mercury methylation potential, and hgcA/hgcB gene expression in *Desulfovibrio dechloroacetivorans* BerOc1. *Environ. Sci. Pollut. Res.* **22**: 13764–13771. doi:10.1007/s11356-015-4273-5
- Graham, A. M., G. R. Aiken, and C. C. Gilmour. 2012. Dissolved organic matter enhances microbial mercury methylation under sulfidic conditions. *Environ. Sci. Technol.* **46**: 2715–2723. doi:10.1021/es203658f
- Graham, A. M., G. R. Aiken, and C. C. Gilmour. 2013. Effect of dissolved organic matter source and character on microbial hg methylation in hg-S-DOM solutions. *Environ. Sci. Technol.* **47**: 5746–5754. doi:10.1021/es400414a
- Grégoire, D. S., and A. J. Poulain. 2018. Shining light on recent advances in microbial mercury cycling. *Facets* **3**: 858–879. doi:10.1139/facets-2018-0015
- Gudasz, C., D. Bastviken, K. Steger, K. Premke, S. Sobek, and L. J. Tranvik. 2010. Temperature-controlled organic carbon mineralization in lake sediments. *Nature* **466**: 478–481. doi:10.1038/nature09383
- Guimarães, J. R. D., J. B. N. Mauro, M. Meili, M. Sundbom, A. L. Haglund, S. A. Coelho-Souza, and L. D. Hylander. 2006. Simultaneous radioassays of bacterial production and mercury methylation in the periphyton of a tropical and a temperate wetland. *J. Environ. Manage.* **81**: 95–100. doi:10.1016/j.jenvman.2005.09.023
- Gworek, B., O. Bemowska-Kałabun, M. Kijeńska, and J. Wrzosek-Jakubowska. 2016. Mercury in marine and oceanic waters—a review. *Water Air Soil Pollut.* **227**: 371. doi:10.1007/s11270-016-3060-3
- Hamelin, S., M. Amyot, T. Barkay, Y. Wang, and D. Planas. 2011. Methanogens: principal methylators of mercury in lake periphyton. *Environ. Sci. Technol.* **45**: 7693–7700. doi:10.1021/es2010072
- Hammerschmidt, C. R., W. F. Fitzgerald, P. H. Balcom, and P. T. Visscher. 2008. Organic matter and sulfide inhibit methylmercury production in sediments of New York/New Jersey Harbor. *Mar. Chem.* **109**: 165–182. doi:10.1016/j.marchem.2008.01.007
- Herrero Ortega, S., and others. 2018. High methylmercury formation in ponds fueled by fresh humic and algal derived organic matter. *Limnol. Oceanogr.* **63**: S44–S53. doi:10.1002/lno.10722
- Hines, M. E., E. N. Poitras, S. Covelli, J. Faganeli, A. Emili, S. Žižek, and M. Horvat. 2012. Mercury methylation and demethylation in Hg-contaminated lagoon sediments (Marano and Grado lagoon, Italy). *Estuar. Coast. Shelf Sci.* **113**: 85–95. doi:10.1016/j.ecss.2011.12.021
- Hintelmann, H., K. Keppel-Jones, and D. Evans. 2000. Constants of mercury methylation and demethylation rates in sediments and comparison of tracer and ambient mercury

- availability. *Environ. Toxicol. Chem.* **19**: 2204–2211. doi:10.1002/etc.5620190909
- Hollweg, T. A., C. C. Gilmour, and R. P. Mason. 2009. Methylmercury production in sediments of Chesapeake Bay and the mid-Atlantic continental margin. *Mar. Chem.* **114**: 86–101. doi:10.1016/j.marchem.2009.04.004
- Hsu-Kim, H., K. H. Kucharzyk, T. Zhang, and M. A. Deshusses. 2013. Mechanisms regulating mercury bioavailability for methylating microorganisms in the aquatic environment: a critical review. *Environ. Sci. Technol.* **47**: 2441–2456. doi:10.1021/es304370g
- Hylander, L. 2000. Relationship of mercury with aluminum, iron and manganese oxy-hydroxides in sediments from the Alto Pantanal, Brazil. *Sci. Total Environ.* **260**: 97–107. doi:10.1016/S0048-9697(00)00544-1
- Jackson, T. a. 1997. Long-range atmospheric transport of mercury to ecosystems, and the importance of anthropogenic emissions—a critical review and evaluation of the published evidence. *Environ. Rev.* **5**: 207. doi:10.1139/er-5-3-4-207
- Jensen, S., and A. Jernelöv. 1969. Biological methylation of mercury in aquatic organisms. *Nature* **223**: 753–754. doi:10.1038/223753a0
- Jones, D. S., G. M. Walker, N. W. Johnson, C. P. J. Mitchell, J. K. Coleman Wasik, and J. V. Bailey. 2019. Molecular evidence for novel mercury methylating microorganisms in sulfate-impacted lakes. *ISME J.* **13**: 1659–1675. doi:10.1038/s41396-019-0376-1
- Jonsson, S., U. Skjällberg, M. B. Nilsson, P. Westlund, A. Shchukarev, E. Lundberg, and E. Björn. 2012. Mercury methylation rates for geochemically relevant HgII species in sediments. *Environ. Sci. Technol.* **46**: 11653–11659. doi:10.1021/es3015327
- Jonsson, S., U. Skjällberg, M. B. Nilsson, E. Lundberg, A. Andersson, and E. Bjo. 2014. Differentiated availability of geochemical mercury pools controls methylmercury levels in estuarine sediment and biota. *Nat. Commun.* **5**: 4624–4634. doi:10.1038/ncomms5624
- Kerin, E. J., C. C. Gilmour, E. Roden, M. T. Suzuki, J. D. Coates, and R. P. Mason. 2006. Mercury methylation by dissimilatory Iron-reducing bacteria. *Appl. Environ. Microbiol.* **72**: 7919–7921. doi:10.1128/AEM.01602-06
- Kim, K.-H., E. Kabir, and S. Ara Jahan. 2016. A review on the distribution of Hg in the environment and its human health impacts. *J. Hazard. Mater.* **310**: 278–279. doi:10.1016/j.jhazmat.2016.02.043
- King, J. K., J. E. Kostka, M. E. Frischer, and F. M. Saunders. 2000. Sulfate-reducing bacteria methylate mercury at variable rates in pure culture and in marine sediments. *Appl. Environ. Microbiol.* **66**: 2430–2437. doi:10.1128/AEM.66.6.2430-2437.2000
- King, J. K., J. E. Kostka, M. E. Frischer, F. M. Saunders, and R. A. Jahnke. 2001. A quantitative relationship that demonstrates mercury methylation rates in marine sediments are based on the community composition and activity of sulfate-reducing bacteria. *Environ. Sci. Technol.* **35**: 2491–2496. doi:10.1021/es001813q
- Klapstein, S. J., and N. J. O. Driscoll. 2018. Methylmercury biogeochemistry in freshwater ecosystems: a review focusing on DOM and photodemethylation. *Bull. Environ. Contam. Toxicol.* **100**: 14–25. doi:10.1007/s00128-017-2236-x
- Kocman, D., S. J. Wilson, H. M. Amos, K. H. Telmer, F. Steenhuisen, E. M. Sunderland, R. P. Mason, P. Outridge, and M. Horvat. 2017. Toward an assessment of the global inventory of present-day mercury releases to freshwater environments. *Int. J. Environ. Res. Public Health* **14**: 138. doi:10.3390/ijerph14020138
- Korthals, E. T., and M. R. Winfrey. 1987. Seasonal and spatial variations in mercury methylation and demethylation in an oligotrophic lake. *Appl. Environ. Microbiol.* **53**: 2397–2404.
- Lamborg, C. H., and others. 2014. A global ocean inventory of anthropogenic mercury based on water column measurements. *Nature* **512**: 65–68. doi:10.1038/nature13563
- Lavoie, R. A., A. Bouffard, R. Maranger, and M. Amyot. 2018. Mercury transport and human exposure from global marine fisheries. *Sci. Rep.* **8**: 1–9. doi:10.1038/s41598-018-24938-3
- Ledeker, B. M., and S. K. De Long. 2013. The effect of multiple primer-template mismatches on quantitative PCR accuracy and development of a multi-primer set assay for accurate quantification of pcrA gene sequence variants. *J. Microbiol. Methods* **94**: 224–231. doi:10.1016/j.mimet.2013.06.013
- Lehnher, I., V. L. S. Louis, H. Hintelmann, and J. L. Kirk. 2011. Methylation of inorganic mercury in polar marine waters. *Nat. Geosci.* **4**: 298–302. doi:10.1038/ngeo1134
- Lehnher, I., V. L. S. Louis, and J. L. Kirk. 2012. Methylmercury cycling in high arctic wetland ponds: controls on sedimentary production. *Environ. Sci. Technol.* **46**: 10523–10531. doi:10.1021/es300577e
- Lei, P., L. M. Nunes, Y. R. Liu, H. Zhong, and K. Pan. 2019. Mechanisms of algal biomass input enhanced microbial Hg methylation in lake sediments. *Environ. Int.* **126**: 279–288. doi:10.1016/j.envint.2019.02.043
- Liem-Nguyen, V., S. Jonsson, U. Skjällberg, M. B. Nilsson, A. Andersson, E. Lundberg, and E. Björn. 2016. Effects of nutrient loading and mercury chemical speciation on the formation and degradation of methylmercury in estuarine sediment. *Environ. Sci. Technol.* **50**: 6983–6990. doi:10.1021/acs.est.6b01567
- Liem-Nguyen, V., U. Skjällberg, and E. Björn. 2017. Thermodynamic modeling of the solubility and chemical speciation of mercury and methylmercury driven by organic thiols and micromolar sulfide concentrations in boreal wetland soils. *Environ. Sci. Technol.* **51**: 3678–3686. doi:10.1021/acs.est.6b04622
- Liu, Y.-R., R.-Q. Yu, Y.-M. Zheng, and J.-Z. He. 2014. Analysis of the microbial community structure by monitoring an Hg methylation gene (hgcA) in paddy soils along an Hg

- gradient. *Appl. Environ. Microbiol.* **80**: 2874–2879. doi:10.1128/AEM.04225-13
- Liu, X., A. Ma, G. Zhuang, and X. Zhuang. 2018a. Diversity of microbial communities potentially involved in mercury methylation in rice paddies surrounding typical mercury mining areas in China. *Microbiology* **7**: e00577. doi:10.1002/mbo3.577
- Liu, Y.-R., A. Johs, L. Bi, X. Lu, H.-W. Hu, D. Sun, J.-Z. He, and B. Gu. 2018b. Unraveling microbial communities associated with methylmercury production in Paddy soils. *Environ. Sci. Technol.* **52**: 13110–13118. doi:10.1021/acs.est.8b03052
- Louis, V. L. S. T., J. W. M. Rudd, C. a. Kelly, K. G. Beaty, R. J. Flett, and N. T. Roulet. 1996. Production and loss of methylmercury and loss of total mercury from boreal forest catchments containing different types of wetlands. *Environ. Sci. Technol.* **30**: 2719–2729. doi:10.1021/es950856h
- Lovley, D. R., and F. H. Chapelle. 1995. Deep subsurface microbial processes. *Rev. Geophys.* **33**: 365–381. doi:10.1029/95rg01305
- Luo, F., C. Yang, T. Sun, Y. Igarashi, X. Dai, D. Wang, M. Ma, and H. Du. 2016. Mercury-methylating genes *dsrB* and *hgA* in soils/sediments of the three gorges reservoir. *Environ. Sci. Pollut. Res.* **24**: 5001–5011. doi:10.1007/s11356-016-8213-9
- Ma, M., H. Du, D. Wang, S. Kang, and T. Sun. 2017. Biotically mediated mercury methylation in the soils and sediments of Nam Co Lake, Tibetan plateau. *Environ. Pollut.* **227**: 243–251. doi:10.1016/j.envpol.2017.04.037
- MacMillan, G. A., C. Girard, J. Chételat, I. Laurion, and M. Amyot. 2015. High methylmercury in arctic and subarctic ponds is related to nutrient levels in the warming eastern Canadian arctic. *Environ. Sci. Technol.* **49**: 7743–7753. doi:10.1021/acs.est.5b00763
- Malcolm, E. G., J. K. Schaefer, E. B. Ekstrom, C. B. Tuit, A. Jayakumar, H. Park, B. B. Ward, and F. M. M. Morel. 2010. Mercury methylation in oxygen deficient zones of the oceans: no evidence for the predominance of anaerobes. *Mar. Chem.* **122**: 11–19. doi:10.1016/j.marchem.2010.08.004
- Mason, R. P., and W. F. Fitzgerald. 1993. The distribution and biogeochemical cycling of mercury in the equatorial Pacific Ocean. *Deep. Res. Part I.* **40**: 1897–1924. doi:10.1016/0967-0637(93)90037-4
- Mason, R. P., A. L. Choi, W. F. Fitzgerald, C. R. Hammerschmidt, C. H. Lamborg, A. L. Soerensen, and E. M. Sunderland. 2012. Mercury biogeochemical cycling in the ocean and policy implications. *Environ. Res.* **119**: 101–117. doi:10.1016/j.envres.2012.03.013
- Mauro, J., J. Guimares, H. Hintelmann, C. Watras, E. Haack, and S. Coelho-Souza. 2002. Mercury methylation in macrophytes, periphyton, and water - comparative studies with stable and radio-mercury additions. *Anal. Bioanal. Chem.* **374**: 983–989. doi:10.1007/s00216-002-1534-1
- Mazrui, N. M., S. Jonsson, S. Thota, J. Zhao, and R. P. Mason. 2016. Enhanced availability of mercury bound to dissolved organic matter for methylation in marine sediments. *Geochim. Cosmochim. Acta* **194**: 153–162. doi:10.1016/j.gca.2016.08.019
- Meier, D. V., and others. 2019. Microbial metal-sulfide oxidation in inactive hydrothermal vent chimneys suggested by metagenomic and metaproteomic analyses. *Environ. Microbiol.* **21**: 682–701. doi:10.1111/1462-2920.14514
- Miller, M. B., and B. L. Bassler. 2001. Quorum sensing in bacteria. *Annu. Rev. Microbiol.* **55**: 165–199. doi:10.1146/annurev.micro.55.1.165
- Mitchell, C. P. J., and C. C. Gilmour. 2008. Methylmercury production in a Chesapeake Bay salt marsh. *J. Geophys. Res. Biogeo.* **113**: G00C04. doi:10.1029/2008jg000765
- Monperrus, M., E. Tessier, D. Amouroux, and A. Leynaert. 2007. Mercury methylation, demethylation and reduction rates in coastal and marine surface waters of the Mediterranean Sea. **107**: 49–63. doi:10.1016/j.marchem.2007.01.018
- Munson, K. M., C. H. Lamborg, R. M. Boiteau, and M. A. Saito. 2018. Dynamic mercury methylation and demethylation in oligotrophic marine water. *Biogeosciences* **15**: 6451–6460. doi:10.5194/bg-15-6451-2018
- Ndu, U., G. A. Christensen, N. A. Rivera, C. M. Gionfriddo, M. A. Deshusses, D. A. Elias, and H. Hsu-Kim. 2018. Quantification of mercury bioavailability for methylation using diffusive gradient in thin-film samplers. *Environ. Sci. Technol.* **52**: 8521–8529. doi:10.1021/acs.est.8b00647
- Pak, K., and R. Bartha. 1998. Mercury methylation and demethylation in anoxic lake sediments and by strictly anaerobic bacteria. *Appl. Environ. Microbiol.* **64**: 1–6.
- Paranjape, A. R., and B. D. Hall. 2017. Recent advances in the study of mercury methylation in aquatic systems, p. 85–119. *In* Recent advances in the study of mercury methylation in aquatic systems. doi:10.1139/facets-2016-0027
- Parks, J. M., and others 2013. The genetic basis for bacterial mercury methylation. *Science* **339**: 1332–1335. doi:10.1126/science.1230667
- Pirrone, N., and others 2009. Global mercury emissions to the atmosphere from natural and anthropogenic sources. *In* Mercury fate and transport in the global atmosphere: Emissions, measurements and models. Springer doi:10.1007/s11259-009-9255-y
- Podar, M., and others. 2015. Global prevalence and distribution of genes and microorganisms involved in mercury methylation. *Sci. Adv.* **1**: e1500675–e1500675. doi:10.1126/sciadv.1500675
- Qian, C., A. Johs, H. Chen, B. F. Mann, X. Lu, P. E. Abraham, R. L. Hettich, and B. Gu. 2016. Global proteome response to deletion of genes related to mercury methylation and

- dissimilatory metal reduction reveals changes in respiratory metabolism in *Geobacter sulfurreducens* PCA. *J. Proteome Res.* **15**: 3540–3549. doi:10.1021/acs.jproteome.6b00263
- Ramlal, P. S., J. W. M. Rudd, and R. E. Hecky. 1986. Methods for measuring specific rates of mercury methylation and degradation and their use in determining factors controlling net rates of mercury methylation. *Appl. Environ. Microbiol.* **51**: 110–114.
- Ranchou-Peyruse, M., M. Monperrus, R. Bridou, R. Duran, D. Amouroux, J. C. Salvado, and R. Guyoneaud. 2009. Overview of mercury methylation capacities among anaerobic bacteria including representatives of the sulphate-reducers: implications for environmental studies. *Geomicrobiol. J.* **26**: 1–8. doi:10.1080/01490450802599227
- Regnell, O., and C. J. Watras. 2019. Microbial mercury methylation in aquatic environments: A critical review of published field and laboratory studies. *Environ. Sci. Technol.* **53**: 4–19. doi:10.1021/acs.est.8b02709
- Schaefer, J. K., and F. M. M. Morel. 2009. High methylation rates of mercury bound to cysteine by *Geobacter sulfurreducens*. *Nat. Geosci.* **2**: 123–126. doi:10.1038/ngeo412
- Schaefer, J. K., S. S. Rocks, W. Zheng, L. Liang, B. Gu, and F. M. M. Morel. 2011. Active transport, substrate specificity, and methylation of Hg(II) in anaerobic bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **108**: 8714–8719. doi:10.1073/pnas.1105781108
- Schaefer, J. K., R.-M. Kronberg, F. M. M. Morel, and U. Skyllberg. 2014. Detection of a key Hg methylation gene, *hgcA*, in wetland soils. *Environ. Microbiol. Rep.* **6**: 441–447. doi:10.1111/1758-2229.12136
- Schartup, A. T., R. P. Mason, P. H. Balcom, T. A. Hollweg, and C. Y. Chen. 2013. Methylmercury production in estuarine sediments: role of organic matter. *Environ. Sci. Technol.* **47**: 695–700. doi:10.1021/es302566w
- Schartup, A. T., U. Ndu, P. H. Balcom, R. P. Mason, and E. M. Sunderland. 2015. Contrasting effects of marine and terrestrially derived dissolved organic matter on mercury speciation and bioavailability in seawater. *Environ. Sci. Technol.* **49**: 5965–5972. doi:10.1021/es506274x
- Skyllberg, U. 2008. Competition among thiols and inorganic sulfides and polysulfides for Hg and MeHg in wetland soils and sediments under suboxic conditions: illumination of controversies and implications for MeHg net production. *J. Geophys. Res.* **113**: 1–14. doi:10.1029/2008JG000745
- Skyllberg, U., J. Qian, W. Frech, K. Xia, and W. F. Bleam. 2003. Distribution of mercury, methyl mercury and organic sulphur species in soil, soil solution and stream of a boreal forest catchment. *Biogeochemistry* **64**: 53–76. doi:10.1023/A:1024904502633
- Skyllberg, U., P. R. Bloom, J. Qian, C.-M. Lin, and W. F. Bleam. 2006. Complexation of mercury(II) in soil organic matter: EXAFS evidence for linear two-coordination with reduced sulfur groups. *Environ. Sci. Technol.* **40**: 4174–4180. doi:10.1021/es0600577
- Smith, S. D., and others. 2015. Site-directed mutagenesis of *hgcA* and *hgcB* reveals amino acid residues important for mercury methylation. *Appl. Environ. Microbiol.* **81**: 3205–3217. doi:10.1128/AEM.00217-15
- Sunderland, E. M., D. P. Krabbenhoft, J. W. Moreau, S. A. Strode, and W. M. Landing. 2009. Mercury sources, distribution, and bioavailability in the North Pacific Ocean: insights from data and models. *Global Biogeochem. Cycles* **23**. doi:10.1029/2008GB003425
- Sundseth, K., J. Pacyna, A. Banel, E. Pacyna, and A. Rautio. 2015. Climate change impacts on environmental and human exposure to mercury in the Arctic. *Int. J. Environ. Res. Public Health* **12**: 3579–3599. doi:10.3390/ijerph120403579
- Tjerngren, I., M. Meili, E. Björn, and U. Skyllberg. 2012. Eight boreal wetlands as sources and sinks for methyl mercury in relation to soil acidity, C/N ratio, and small-scale flooding. *Environ. Sci. Technol.* **46**: 8052–8060. doi:10.1021/es300845x
- Ullrich, S. M., T. W. Tanton, S. A. Abdrashitova, and A. Svetlana. 2001. Mercury in the aquatic environment: a review of factors affecting methylation. *Crit. Rev. Environ. Sci. Technol.* **31**: 241–293. doi:10.1080/20016491089226
- UNEP. 2013. Global mercury assessment 2013: Sources, emissions, releases, and environmental transport. UNEP 42. doi:DTI/1636/GE
- Villar, E., L. Cabrol, and L.-E. Heimbürger-Boavida. 2019. Widespread microbial mercury methylation genes in the global ocean. *bioRxiv*. doi:10.1101/648329
- Vishnivetskaya, T. A., and others. 2018. Microbial community structure with trends in methylation gene diversity and abundance in mercury-contaminated rice paddy soils in Guizhou, China. *Environ. Sci. Process. Impacts* **20**: 673–685. doi:10.1039/C7EM00558J
- Vonk, J. W., and A. K. Sijpesteijn. 1973. Studies on the methylation of mercuric chloride by pure cultures of bacteria and fungi. *In* Studies on the methylation of mercuric chloride by pure cultures of bacteria and fungi. Antonie Van Leeuwenhoek. doi:10.1007/BF02578894
- Weber, K. A., L. A. Achenbach, and J. D. Coates. 2006. Microorganisms pumping iron: Anaerobic microbial iron oxidation and reduction. *Nat. Rev. Microbiol.* **4**: 752–764. doi:10.1038/nrmicro1490
- Windham-Myers, L., and others. 2014. Mercury cycling in agricultural and managed wetlands: A synthesis of methylmercury production, hydrologic export, and bioaccumulation from an integrated field study. *Sci. Total Environ.* **484**: 221–231. doi:10.1016/j.scitotenv.2014.01.033
- Xu, J., and others. 2019. Mercury methylating microbial communities of boreal forest soils. *Sci. Rep.* **9**: 518. doi:10.1038/s41598-018-37383-z
- Yu, R. Q., J. R. Flanders, E. E. MacK, R. Turner, M. B. Mirza, and T. Barkay. 2012. Contribution of coexisting sulfate and iron reducing bacteria to methylmercury production in

freshwater river sediments. *Environ. Sci. Technol.* **46**: 2684–2691. doi:[10.1021/es2033718](https://doi.org/10.1021/es2033718)

Yu, R.-Q., J. R. Reinfelder, M. E. Hines, and T. Barkay. 2013. Mercury methylation by the methanogen *Methanospirillum hungatei*. *Appl. Environ. Microbiol.* **79**: 6325–6330. doi:[10.1128/AEM.01556-13](https://doi.org/10.1128/AEM.01556-13)

Yu, R. Q., J. R. Reinfelder, M. E. Hines, and T. Barkay. 2018. Syntrophic pathways for microbial mercury methylation. *ISME J.* **12**: 1826–1835. doi:[10.1038/s41396-018-0106-0](https://doi.org/10.1038/s41396-018-0106-0)

Acknowledgments

Authors thank Prof. Jeffra Schaeffer (Rutger University, USA) for critical reading of the draft.

AGB acknowledges the European Commission, for the *MER-CURE* (*Using global marine metagenomics to understand MERcury microbial associated processes: finding a CURE for mercury contaminated environments*), an individual Fellowship of the Marie Skłodowska-Curie Actions.

Conflict of Interest

None declared.

Submitted 09 April 2019

Revised 17 September 2019

Accepted 02 October 2019

Associate editor: Vanessa Hatje