

EpiBrainRad An epidemiologic study of the neurotoxicity induced by radiotherapy in high grade glioma patients

Thomas Durand, Sophie Jacob, Laura Lebouil, Hassen Douzane, Philippe Lestaevel, Amithys Rahimian, Dimitri Psimaras, Loïc Feuvret, Delphine Leclercq, Bruno Brochet, et al.

▶ To cite this version:

Thomas Durand, Sophie Jacob, Laura Lebouil, Hassen Douzane, Philippe Lestaevel, et al.. Epi-BrainRad An epidemiologic study of the neurotoxicity induced by radiotherapy in high grade glioma patients. BMC Neurology, 2015, 15 (1), pp.261. 10.1186/s12883-015-0519-6 . hal-02572573

HAL Id: hal-02572573 https://hal.science/hal-02572573

Submitted on 11 Jan2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

STUDY PROTOCOL

Open Access



EpiBrainRad: an epidemiologic study of the neurotoxicity induced by radiotherapy in high grade glioma patients

Thomas Durand^{1,2}, Sophie Jacob³, Laura Lebouil², Hassen Douzane², Philippe Lestaevel³, Amithys Rahimian⁴, Dimitri Psimaras², Loïc Feuvret², Delphine Leclercq⁵, Bruno Brochet⁶, Radia Tamarat³, Fabien Milliat³, Marc Benderitter³, Nicolas Vayatis⁷, Georges Noël⁸, Khê Hoang-Xuan², Jean-Yves Delattre², Damien Ricard^{1,2,9} and Marie-Odile Bernier^{3*}

Abstract

Background: Radiotherapy is one of the most important treatments of primary and metastatic brain tumors. Unfortunately, it can involve moderate to severe complications among which leukoencephalopathy is very frequent and implies cognitive deficits such as memory, attention and executive dysfunctions. However, the incidence of this complication is not well established and the risk factors and process are poorly understood. The main objective of the study is to improve knowledge on radio-induced leukoencephalopathy based on pluridisciplinar approaches combining cognitive, biologic, imagery and dosimetric investigations.

Method/Design: The EpiBrainRad study is a prospective cohort study including newly diagnosed high grade gliomas patients treated by radiotherapy and concomitant-adjuvant temozolomide chemotherapy. Patients are included between their surgery and first day of radio-chemotherapy, and the follow-up lasts for 3 years after treatment. Cognitive functioning assessments, specific blood biomarkers measures and magnetic resonance imagery are performed at different moment during the follow-up, and a specific dosimetric assessment of organs involved in the beam fields is performed. Firstly, leukoencephalopathy incidence rate will be estimated in this population. Secondly, correlations between cognitive impairments and dosimetry, biomarkers ranges and anomalies on imagery will be analyzed in order to better understand the onset and evolution of cognitive decrement associated with radiotherapy. Furthermore, a new cognitive test, quickly and easily performed, will be studied to determine its sensibility to detect leukoencephalopathy decrement.

Discussion: With an original multidisciplinary approach, the EpiBrainRad study aims to improve knowledge on radio-induced leukoencephalopathy in order to improve its early diagnosis and prevention. The main challenge is to preserve quality-of-life after cancer treatments which imply to study the incidence of radiation-induced complications and their associated risk factors.

Trial Registration: NCT02544178

Keywords: Radiotherapy, Cognitive impairments, Neurotoxicity, Leukoencephalopathy, Quality-of-life

* Correspondence: marie-odile.bernier@irsn.fr

³Institut de Radioprotection et de Sûreté Nucléaire (IRSN), PRP-HOM, SRBE, 31 avenue de la Division Leclerc, 92260 Fontenay aux Roses, France Full list of author information is available at the end of the article



© 2015 Durand et al. **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

Background

Radiotherapy is one of the most important treatments of metastatic and primary brain tumors of which 60 % are high grade glioblastomas. With the recent development of modern irradiation techniques, survival of patients has increased and mid- to long-term side effects became more visible, such as neurologic complications [1]. Among these complications, leukoencephalopathy seems to be the most frequent. It is characterized by a progressive and diffuse demyelination, an axonal loss and vascular lesions [2]. Consequences of leukoencephalopathy involve cognitive deficits which dramatically reduce the patient's quality-of-life [3-5]. Thus, since the last few years, the neuropsychological status represents an important issue in clinical trials as well as in individual outcomes [6]. Long-term memory, information speed processing, attention and executive functions are recognized to be the most sensitive functions to be affected by radiations [2, 7-10]. However, there is no consensus to describe the main evolution of cognitive decrement following radiotherapy because of important differences in studies' protocols (assessment time, decrement definitions, material). The radiation-induced leukoencephalopathy incidence is thus difficult to estimate precisely and varies from 30 to 50 % [10, 11] according to the length of follow-up. Moreover, available studies are often retrospective which may induce bias and based on small samples which limits statistical robustness of results.

Although several risk factors of neurotoxic complication have been identified such as patient's age, tumor location, total dose of radiation, fractionation, field size [2, 9, 12], the pathophysiology of radiation-induced neurotoxicity is still poorly understood. It could involve inflammation, blood barrier disruption, vascular lesion, demyelination, radio-necrosis and edema [2, 9]. Until now, it is impossible to precisely estimate at individual level the risk for a patient to develop this neurotoxic complication [13, 14]. Some individual risk factor such as cardio-vascular diseases (hypertension, diabetes), smoking, old age [15, 16] seem implicated in leukoencephalopathy. Assessment of the impact of both individual risk factors and treatment toxicity requires to study a large group of patient with prospective collection of accurate data and follow-up.

From a biological point of view, biological mechanisms of the initiation and progression of cognitive impairements is not well known. However, specific biomarkers predictive of the cognitive impairments in this population would help in screening patients at risk of neurotoxicity. Currently, biomarkers of neurotoxicity have been poorly studied. Thus, several markers deserve to be explored such as the S-100B protein known to be a marker of traumatic cerebral lesion, neurodegenerative diseases and aggressive gliomas [17], specific isoprostanes (as 8,12-*iso*-iPF_{2α}-VI) implicated in the oxidative/nitrate stress in neurodegenerative diseases [18] and homocystein which plasma concentration has been associated with poor memory performances in old individuals [19]. Furthermore, new biomarkers seems to be potentially very interesting. Indeed, micro RNA and microparticules have been identified as potential biomarkers of neurological defects [20, 21]. Then the strategy proposed is to test classical and new biomarkers in order to individualize those, which could be accurate for the screening of patients at risk of cognitive impairments.

At last, assessment of cognitive impairment is difficult because the exploration of cognitive side effects is not systematic and often depends on patients' complaints and clinicians' sensibility to this aspect. A formal neuropsychological assessment can't be performed in routine, as often as it should be for an adapted follow-up of brain tumor patients. Indeed, it is time consuming, needs expert professionals and validated material, and consequently it is not easily implemented in most medical center. A new tool for exploration of cognitive deficits is the Computerized Speed Cognitive Test (Legal deposit: IDDN.FR.001.180018.000.S.P.2014.000.31230) (CSCT) [22]. The CSCT was first developed to assess information speed processing in multiple sclerosis. Because it presents a low learning-effect and focuses on a cognitive characteristic often impaired in neuro-oncologic population, it could be used to detect a cognitive deterioration during repeated medical visits, and thus be used as an alarm to start a more extensive formal assessment. Then, it is suitable for the follow-up of a cohort of patients treated for brain tumor and allows to give quick and reliable information on cognitive impairment, compatible with the usual clinical followup of the patients.

Objectives

The purpose of the EpiBrainRad study is to establish a cohort of patients treated with radiotherapy for a high grade glioma.

The main objective is to estimate the cognitive impairment incidence related to radio-induced leukoencephalopathy in this population.

Secondary objectives are:

- To study the impact of associated risk factors on leukoencephalopathy development, including individual factors and treatment.
- Biologic markers of neurologic degradation will be studied to evaluate the correlations between cognitive impairments and biological abnormalities.
- Specific organs dosimetry (such as hippocampus, temporal lobes, corpus callosum, peri-ventricular white matter, posterior fossa) correlated with radiologic abnormalities on magnetic resonance imagery (MRI)

and clinical symptoms will be analyzed to better understand the evolution of radio-induced leukoencephalopathy.

 Finally, the assessment of the sensibility and specificity of a quick cognitive test, *the Computerised Speed Cognitive Test* [22], to detect a cognitive decrement during the follow up of the patients.

Methods/Design

Study design and patient selection

The EpiBrainRad study is an observational prospective cohort. Patients' recruitment takes place at *Hôpital de la Pitié-Salpêtrière (Paris, France)* and *Centre de Lutte Contre le Cancer Paul Strauss (Strasbourg, France)* for 2 years and each patient is followed during 3 years. The cohort includes newly-diagnosed adults treated for a high grade glioma according to the procedure described by Stupp et al. [23]. According to patient's age, the radiotherapy regiment can be modified to 40 gy in 15 fractions to ensure a better tolerance in elders [24]. Inclusion and exclusion criteria are listed in Table 1. Data are prospectively including clinical features, MRI images and results, blood samples, dosimetry and cognitive assessments during the usual medical follow-up.

Ethical approval

This study has received ethical approval from the *Comité de Protection de la Personne de Paris VI Ile de France* in December 2014 (ID: CPP/132-14) and from National Agency for Medical and Health products Safety (Reference: 2014-A01697-40). The Clinical Trial Registration Information is available at http://www.clinicaltrials.gov (Unique identifier NCT02544178). Participants enrolled in the study provide their written informed consent.

Study endpoints

The primary endpoint is the reduction in performance on cognitive tests defined as follows: After exclusion of local recurrence, decreased detailed neuropsychological assessment score compared to the patient's baseline score before radiotherapy for at least 2 cognitive domains.

Secondary endpoints are:

Paul Strauss (Strasbourg, France)

- Differences in biomarkers' concentration between the measurements before RT and: at the end of the RT, RT + 4 weeks, RT + 12 months and RT + 36 months.
- Changes in MRI images compared to the baseline imaging.
- Correlations between dosimetry, cognitive decay and MRI abnormalities onset after RT.

Procedures

Figure 1 gives an overview of the study flowchart. Patients' inclusion occurs after surgery and before radio/ chemotherapy (Baseline). At this time, individual information including demographic information, medical background and tumor characteristic are collected.

Every 3 months, once the MRI of the patient is done, a medical visit occurs during which a short cognitive assessment is performed (the CSCT).

Cognitive tests, scales and guestionnaires used for the formal assessment are listed in Table 2. An extensive evaluation lasting 2 h is performed with a trained neuropsychologist, followed by self-reported and caregiverreported scales and questionnaires. The explored domains are intellectual efficiency, general functioning, verbal and visual episodic memory, attention, executive functioning, language, visuospatial and visuoconstructive abilities, mood, fatigue, quality-of-live, subjective complaints and autonomy. The formal assessment occurs at baseline and yearly after radiotherapy. In case of CSCT score decrement during follow-up the next yearly formal assessment is performed earlier. CSCT decrement is defined as a loss of 1 standard deviation compared to baseline (given its low learning-effect and good stability in controls [22]), during two consecutive assessments. The necessity of detecting a decrement during two visits permits to insure that a variation in score is not due to a transitory cause like an infection or fatigue, which is reversible and not linked to leukoencephalopathy. This procedure will allow to study the nature, course and incidence of cognitive decrement following radiotherapy in this population. At baseline and during the follow-up, the CSCT will be compared to the formal cognitive assessment to study its sensibility to detect impairments in this context.

Table 1 Inclusion and exclusion criteria				
Inclusion criteria	Exclusion criteria			
 Newly diagnosed grade III or IV glioma Age ≥ 18 Treated by radiotherapy and concomitant-adjuvant Tomozolomide as described by Stupp et al. [23] or with an abbreviated course of radiotherapy in elders patients [24] Treatment and clinical follow-up performed at Hôpital de la 	 History of other CNS tumor History of neurologic and/or psychiatric disease involving cognitive impairments (TBI, Stroke, mood and personality disorders) Insufficient understanding of French language 			
Pitié-Salpêtrière (Paris, France) or Centre de lute contre le cancer				



Fig. 1 Study procedure

Baseline and trimensual MRI during the medical follow-up are used for extra analysis in order to study white matter abnormalities and cortical atrophy. Moreover, tumor response or progression will be assessed by the Response Assessment in Radio-Oncology working group (RANO) Criteria [25]. The white matter lesion quantification will follow the procedure described by Wahlund et al. [26] using a 4 points scale (0: no lesion; 1: focal lesions; 2: beginning confluence of lesion; 3: diffuse involvement of the entire region). The cortical atrophy quantification will follow the procedure described by Pasquier et al. [27] using a 4 points scale (0: absence of atrophy; 1: mild atrophy; 2: moderate atrophy; 3: severe atrophy). White matter lesion and cortical atrophy is assessed for all key region listed in Table 3. A double blind lecture of the MRI will be done in order to quantify these anomalies. Radiological outcomes will then be correlated with cognitive outcomes.

The need for a biological marker of cognitive disorders has proven to be urgent, both for diagnosis and for monitoring of the diseases. Among the advantages that such markers may bring, we emphasize the possibility of early and even preclinical diagnosis of the disease with the subsequent correct treatment of disease by the medical team. Biomarkers for cognitive impairment have been an issue in recent years. The value of a variety of these markers, like amyloid beta or tau, has been evaluated and discussed [28]. Besides proteins, microRNAs have also demonstrated their potential as non-invasive biomarkers from blood and serum for a wide variety of human pathologies [29]. A deregulation of microRNA expression might be involved in neurological dysfunction or neurodegenerative processes. A recent serum profiling of Alzheimer disease patients provided evidence that expression changes of circulating miRNAs may be valuable biomarkers for this neurological disease [20]. In addition, microparticles have been implicated to have pathological roles in many diseases such as Alzheimer's disease. The consensus among recent studies is that increased levels of specific types of microparticles in plasma may represent reliable biological markers for the onset and progression of central nervous system diseases [30].

Consequently, plasmatic classical biomarkers as S-100B protein, isoprostane 8,12-*iso*-iPF_{2 α}-VI, homocysteine and new ones, as micro RNA and microparticules (see Table 4) will be sampled in blood at baseline, 4 weeks, 12 months and 36 months after the end of radiotherapy. These biomarkers could be potential blood predictors of cognitive impairments [17, 19–21] and thus their concentration evolution after radiations will be analyzed. Additional blood sample tubes are stored to constitute a biological collection for future investigations about others biomarkers.

Finally, histograms of dose-volumes (HDV) of specific organs and areas are collected to study the relation between the radiation doses received by these structures and the cognitive decrements observed. Indeed, some authors showed correlations between specific structures and cognitive performances [31, 32] but few data are available and thus these relations need to be explored. More, dosimetry is compared with onset of abnormalities on MRI to improve knowledge about radio-induced leukoencephalopathy process. Structures prospectively investigated are listed in Table 3.

Statistical analysis

Sample size consideration

The main objective being to estimate the incidence of occurrence of neurological complications in the cohort, with an expected proportion P varying from 0.2 to 0.5, for a confidence level of 95 %, with alpha risk of 5 %, the number of subjects to be included varies between 246

Table 2 Tests and questionnaires used for the cognitive, mood and behavioral assessment

Domain	Tests, scales and questionnaires
Cognitive assessment	
Intellectual efficiency	National Adult Reading Test (French version) [38]
General functioning	Mini Mental State Examination [39]
	Montreal Cognitive Assessment [40]
	Dementia Rating Scale-2 [41]
Verbal episodic memory	RLRI-16 (Adaptation of the Grober-Buschke test) [42]
Visual episodic memory	Rey-Osterrieth Complex Figure [43]
Executive functions and Attention	Digit Span forward [44]
	Digit Span backward [44]
	Letter-number sequencing [44]
	Categorical and phonetic word fluency [45, 46]
	Trail Making Test [45, 46]
	Computerized Speed Cognitive Test [22]
	Stroop color-word Test [45, 46]
Langage	Boston Naming Test [47]
	Token Test [48]
Visuospatial and visuoconstructive functions	Rey-Osterrieth Complex Figure [43]
Self-reported Scales	
Mood	Anxiety questionnaire of Goldberg [49]
	Center for Epidemiologic Study Depression Scale [50]
Fatigue	FACIT Fatigue Scale [51]
Memory complaint	MacNair and Khan questionnaire [52, 53]
Quality-of-life	EORTC QLQ-C30 [54]
	EORTC QLQ-BN20 [55]
Care-giver-reported Scales	
Dysexecutive syndrome	ISDC [45, 46]
Autonomy	EIADL [56]

Table 3	Regions	prospectively	explored	on	MRI	and	dosimetric
analysis							

Key regions investigated				
	MRI	Dosimetry		
Frontal lobes	*	*		
Temporal lobes	*	*		
Parietal lobes	*			
Occipital lobes	*			
Cerebellum	*	*		
Basal ganglia	*			
Brain Stem	*	*		
Hippocampus		*		
Corpus Callosum		*		
Periventricular white matter		*		
* anatomical region investigated				

and 384. Given the risk of relatively early death related to the disease, the inclusion of 400 patients will provide good statistical power for this purpose.

Planned analysis

The combination of data from four axes (Clinical Epidemiology, Dosimetry, Biomarkers and Radiobiology) will give the opportunity to precisely analyze the links between cognitive impairments and changes in biomarkers depending on the precise dose absorbed at different brain areas during radiotherapy brain cancer.

Descriptive statistics (means, standard deviations, percentages and 95 % confidence intervals) will be used to describe the distribution of subjects according to each of the terms of the variables studied. Analyses of variance to one or more classification criteria will be used to compare the means of continuous quantitative variables. Chi-square test or Fisher exact test will be used to compare the distribution of nominal discrete variables and /

Table 4 Biomarkers sampled in blood

Classical biomarkers correlated with cognitive impairments	S-100B protein [GenBank : NP_006263]
	8,12-iso-iPF2a-VI isoprostane
	Homocysteine [GenBank : NP_060084]
	microRNAs
New biomarkers	brain-miR-112 [GenBank : AF480510]
	brain-miR-161 [GenBank : AJ535829]
	hsa-let-7d-3p [GenBank : LM380164]
	hsa-miR-5010-3p [GenBank : LM38284]
	hsa-miR-26a-5p [GenBank : LM378769]
	hsamiR-1285-5p [GenBank : LM383022]
	hsa-miR-151a-3p [GenBank : LM379262]
	hsamiR-103a-3p [GenBank : LM378788]
	hsa-miR-107 [GenBank : LM378791]
	hsa-miR-532-5p [GenBank : LM379811]
	hsa-miR-26b-5p [GenBank : LM378770]
	hsa-let-7f-5p [GenBank : LM378754]
	microparticles

or ordinal. All tests will be bilateral with alpha = 5 %. Details of the analyzes are shown below:

Description of the population at baseline (before radiotherapy) This description will include:

- Medical data collected in the questionnaire, including in particular the baseline risk factors of neurological complications.
- All biomarkers
- The MRI abnormalities
- The results of CSCT and complete cognitive assessments.

The results will be presented for the complete sample.

Dose assessment

This reconstruction will be made for each patient based on data from radiotherapy treatment plan. Histograms of dose/volumes will be retrieved for each structures of interest listed in Table 3 in order to study correlations between anatomic regions and neuropsychological dysfunctions.

Analysis of evaluation criteria

Based on primary endpoint definition, the estimate of the incidence of cognitive impairment related to radioinduced leukoencephalopathy in our cohort will be made.

An analysis of the dose–response relationship between radiation dose absorbed by different brain structures and cognitive abnormalities observed after radiotherapy will be made using a multivariate Cox model taking into account the delivered dose to the organ of interest and the time occurrence of cognitive complications. An adjustment on risk factors for cognitive complications will be achieved. Similarly for patients with cognitive impairment before radiotherapy, an analysis of potential associated risk factors will be carried out with the use of a multivariate model.

Secondary judgments criteria, including changes in MRI images relative to the reference imaging and changes in series of biological markers between measurements before RT, 4 weeks, 12 month and 36 months after RT, will be analyzed by parametric distribution comparison tests (Z) or nonparametric if needed (Wilcoxon tests).

In addition, an analysis will be conducted of the sensitivity and specificity of the CSCT to demonstrate cognitive impairment by comparing the CSCT performed before radiotherapy and detailed neuropsychological assessment (gold standard) achieved at the same time. More, CSCT decay during follow-up will be analyzed by comparing its results to formal assessment during years after treatment, to study its sensibility in detecting cognitive decrement after radiotherapy.

Time plan

Initial inclusion data collection began in April 2015 and will continue through 2017. First statistical analysis about cognitive impairment related to radio-induced leukoencephalopathy incidence will be available in the end of 2016.

Discussion

This study should improve our knowledge on neurologic complications of radiotherapy through an original multidisciplinar approach combining cognitive, biologic, imagery and dosimetric investigations. Our primary objective is to precisely explore cognitive impairments present before radiotherapy and to follow their worsening, or the onset of new impairments not present at baseline, during the first years after the treatment. The collection of data from different nature could be very useful to understand the leukoencephalopathy process and risk factors associated. Indeed, the assessment of cognitive defect linked to leukoencephalopathy could be obscured by the progression of the disease. Then, precise assessment of the type of cognitive defect associated with information on dose received by organs of interest and MRI results will be a major issue to study the dose

response relationship between leukoencephalopathy development and radiation exposure to the brain.

Chemotherapy is known to induce cognitive impairments as well and produce a synergistic effect when administrated in combination with RT [2, 12]. Unfortunately, the treatment protocol received by our patient population doesn't allow to specifically study the participation of chemotherapy in cognitive decrement. Nevertheless, chemotherapy related impairment are assumed to be transient [33] and not as strong as radiationrelated impairments. Moreover, neurotoxicity of chemotherapy is not always proved in studies investigating this problem [34], and subjective complaints of patient seems to be more related to emotional distress and fatigue than to formal dysfunctions [35].

One other important goal of our project is to improve the detection of cognitive defects at early stage. Current formal assessment of cognitive status lasts between 1 and 2 h or more, and cannot be performed as often as needed in this population because of an important learning-effect. The Computerised Speed Cognitive Test (CSCT) is a quick cognitive assessment tool validated in a population of patients presenting multiple sclerosis. The CSCT just need few minutes to be performed and appears to be a potential useful tool in the patient follow-up [36]. This test use in the neuro-oncologic context could importantly improve the detection of cognitive impairments and decrement. Our study will help to validate its use in our population by comparing the CSCT to a more complete battery at baseline time before radiotherapy and during the follow-up after treatment. Several biomarkers of neurotoxicity will be also tested in our population in order to try to individualize prognostic biomarkers of leukoencephalopathy.

Prospective methodology of data collection and the choice of a homogenous population will allow us to analyze precisely the selected outcomes, avoiding potential bias linked to retrospective studies. However, one limit of our study is to focus only on patients with high grade of glioblastoma, who are known to have a very short life expectancy, with a median expected survival being less than 18 months [37]. Nevertheless, it seems that leukoencephalopathy could appear as soon as within the 6 first months after radiotherapy. Furthermore, the rather large number of included patients will allow us to characterize cognitive defects at short and medium term.

With a better screening and understanding of neurotoxicity, the treatment regiments could be adapted to risk factors present for each patient in order to reduce complication. More, preventive actions and cares could be developed to reduce the burden of cognitive deficits and preserve the quality-of-life and autonomy for patient at risk.

Conclusion

Leukoencephalopathy is one of the most frequent midto long-term complications of cranial radiotherapy. However, its incidence is not well established and the pathophysiology remains poorly understood. By an original multidisciplinary approach, the study EpiBrainRad aims to improve knowledge about this condition to facilitate the detection and prevention of radio-induced impairments. The main challenge is to preserve qualityof-life after cancer treatments, which involves to study risk factors and development of such complications.

Abbreviations

CSCT: Computerised Speed Cognitive Test.; MRI: Magnetic Resonance Imagery; RT: Radiotherapy.

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

Conception and design of the study: TD, PL, DP, DL, GN, NV, RT, FM, MB, DR, MOB. Data acquisition: TD, LL, AR, DP, LF, GN. Data interpretation: TD, SJ, PL, DP, LF, DL, GN, DR, MOB. Study supervision: TD, DP, GN, DR, MOB. Manuscript drafting: TD. Critical revision of the manuscript: all authors. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by the French *Institut des Neurosciences Translationnelles de Paris (ICM-IHU)* for the collection of data. The authors are grateful to the clinicians of participating hospitals, who take care of the patients included in the study.

Author details

¹UMR CNRS 8257 SSA MD4 Cognition and Action Group, 45 rue des Saints Pères, 75270 Paris CEDEX 06, France. ²Service de neurologie Mazarin, hôpital de la Pitié-Salpêtrière, 47-83 boulevard de l'Hôpital, 75013 Paris, France. ³Institut de Radioprotection et de Sûreté Nucléaire (IRSN), PRP-HOM, SRBE, 31 avenue de la Division Leclerc, 92260 Fontenay aux Roses, France. ⁴Institut du Cerveau et de la Moelle, 47-83 boulevard de l'Hôpital, 75013 Paris, France. ⁵Unité de neuroradiologie diagnostique et fonctionnelle, hôpital de la Pitié-Salpêtrière, 47-83 boulevard de l'Hôpital, 75013 Paris, France. ⁶Unité de neuroradiologie diagnostique et fonctionnelle, hôpital de la Pitié-Salpêtrière, 47-83 boulevard de l'Hôpital, 75013 Paris, France. ⁶Service de Neurologie, groupe hôspitalier Pellegrin, place Amélie Raba-Léon, 33076 Bordeaux, France. ⁷UMR CNRS 8536 Centre de mathématiques et de leurs applications, ENS Cachan, 61 avenue du président Wilson, 94235 Cachan CEDEX, France. ⁸Département de radiothérapie, centre de lutte contre le cancer Paul Strauss, 3 rue de la porte de l'hôpital, 67065 Strasbourg CEDEX, France. ⁹Service de neurologie, hôpital d'instruction des armées du Val-de-Grace, 71 boulevard de Port-Royal, 75005 Paris, France.

Received: 2 October 2015 Accepted: 11 December 2015 Published online: 18 December 2015

References

- Verdecchia A, Baili P, Quaglia A, Kunkler I, Ciampichini R, Berrino F, et al. Patient survival for all cancers combined as indicator of cancer control in Europe. Eur J Public Health. 2008;18:527–32.
- Soussain C, Ricard D, Fike JR, Mazeron JJ, Psimaras D, Delattre JY. CNS complications of radiotherapy and chemotherapy. Lancet. 2009;374:1639–51.
- Li J, Bentzen SM, Li J, Renschler M, Mehta MP. Relationship between neurocognitive function and quality of life after whole-brain radiotherapy in patients with brain metastases. Int J Radiat Oncol Biol Phys. 2008;71:64–70.
- Hahn CA, Dunn RH, Logue PE, King JH, Edwards CL, Halperin RC. Prospective study of neuropsychologic testing and quality-of-life assessment of adults with primary malignant brain tumors. Int J Radiat Oncol Biol Phys. 2003;55:992–9.
- Dirven L, Aaronson N, Heimans JJ, Taphoorn MJ. Health-related quality of life in high grade glioma patient. Chin J Cancer. 2014;33:40–5.

- Meyer CA, Rock EP, Fine HA. Refining endpoints in brain tumors clinical trials. J Neurooncol. 2012;108:227–30.
- Froklage FE, Oosterbaan LJ, Sizoo EM, de Groot M, Bosma I, Sanchez E, et al. Central neurotoxicity of standard treatment in patients with newlydiagnosed high-grade glioma: a prospective longitudinal study. J Neurooncol. 2014;116:387–94.
- Habets EJJ, Taphoorn MJB, Nederend S, Klein M, Delgadillo D, Hoang-Xuan K, et al. Health-related quality-of-life and cognitive functioning in long-term anaplastic oligodendroglioma and oligoastrocytoma survivors. J Neurooncol. 2014;116:161–8.
- 9. Saad S, Wang TJ. Neurocognitive deficits after radiation therapy for brain malignancies. Am J Clin Oncol. 2014. [Epub ahead of print].
- Douw L, Klein M, Fagel SS, van den Heuvel J, Taphoorn MJ, Aaronson NK, et al. Cognitive and radiological effects of radiotherapy in patients with low-grade glioma: long-term follow-up. Lancet Neurol. 2009;8:810–18.
- Hottinger AF, Yoon H, DeAngelis LM, Abrey LE. Neurological outcome of long-term glioblastoma survivors. J Neurooncol. 2009;95:301–5.
- Scoccianti S, Detti B, Cipressi S, Iannalfi A, Franzese C, Biti G. Changes in neurocognitive functioning and quality of life in adult patients with brain tumors treated with radiotherapy. J Neurooncol. 2012;108:291–308.
- Klein M, Heimans JJ, Aaronson NK, van der Ploeg HM, Grit J, Muller M, et al. Effect of radiotherapy and other treatment-related factors on mid-term to long-term cognitive sequelae in low-grade gliomas: a comparative study. Lancet. 2002;360:1361–8.
- Armstrong CL, Hunter JV, Ledakis GE, Cohen B, Tallent EM, Goldstein BH, et al. Late cognitive and radiographic changes related to radiotherapy: initial prospective findings. Neurology. 2002;59:40–8.
- Imperato JP, Paleologos NA, Vick NA. Effects of treatment on long- term survivors with malignant astrocytomas. Ann Neurol. 1990;28:818–22.
- Vigliani M, Duyckaerts C, Delattre J. Radiation-induced cognitive dysfunction in adults. In: Vecht C et al., editors. Handbook of clinical neurology. Amsterdam: Elsevier Science; 1997. p. 371–88.
- 17. Astrand R, Undén J, Romner B. Clinical use of the calcium-binding S100B protein. Methods Mol Biol. 2013;963:373–84.
- Pratico D, Clark CM, Liun F, Rokach J, Lee VY, Trojanowski JQ. Increase of brain oxidative stress in mild cognitive impairment: a possible predictor of Alzheimer's disease. Arch Neurol. 2002;59:972–6.
- Morris MS, Jacques PF, Rosenberg IH, Selhub J. Hyperhomocysteinemia associated with poor recall in the third national health and nutrition examination survey. Am J Clin Nutr. 2001;73:927–33.
- Leidinger P, Backes C, Deutscher S, Schmitt K, Mueller SC, Frese K, et al. A blood based 12-miRNA signature of Alzheimer disease patients. Genome Biol. 2013;14(7):R78.
- Xue S, Cai X, Li W, Zhang Z, Dong W, Hui G. Elevated plasma endothelial microparticules in Alzheimer's disease. Dement Geriatr Cogn Disord. 2012;34:174–80.
- Ruet A, Deloire MS, Charré-Morin J, Hamel D, Brochet B. A new computerised cognitive test for the detection of information processing speed impairment in multiple sclerosis. Mult Scler. 2013;19:1665–72.
- Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J Med. 2005;352:987–96.
- Roa W, Brasher PM, Bauman G, Anthes M, Bruera E, Chan A, et al. Abbreviated course of radiation therapy in older patient with glioblastoma: a prospective randomized controlled trial. J Clin Oncol. 2004;22:1583–8.
- Wen PY, Macdonald DR, Reardon DA, Cloughesy TF, Sorensen AG, Galanis E, et al. Updated response assessment criteria for high-grade gliomas: Response Assessment in Neuro-Oncology working group. J Clin Oncol. 2010;28:1963–72.
- Wahlund LO, Barkhof F, Fazekas L, Bronge L, Augustin M, Sjögren M, et al. A new rating scale for age-related white matter changes applicable to MRI and CT. Stroke. 2001;32:1318–22.
- Pasquier F, Leys D, Weerts JG, Mounier-Vehier F, Barkhof F, Scheltens P. Inter- and intraobserver reproducibility of cerebral atrophy assessment on MRI scans with hemispheric infarcts. Eur Neurol. 1996;36:268–72.
- Maccioni RB, Lavados M, Maccioni CB, Mendoza-Naranjo A. Biological markers of Alzheimer's disease and mild cognitive impairment. Curr Alzheimer Res. 2004;1:307–14.
- 29. Keller A, Leidinger P, Bauer A, Elsharawy A, Haas J, Backes C, et al. Toward the bloodborne miRNome of human diseases. Nat Methods. 2011;8:841–3.

- Verderio C, Muzio L, Turola E, Bergami A, Novellino L, Ruffini F, et al. Myeloid microvesicles are a marker and therapeutic target for neuroinflammation. Ann Neurol. 2012;72:610–24.
- Gondi V, Hermann BP, Mehta MP, Tomé WA. Hippocampal dosimetry predicts neurocognitive function impairment after fractionated stereotactic radiotherapy for benign or low-grade adult brain tumors. Int J Radiat Oncol Biol Phys. 2012;85:348–54.
- Peiffer AM, Leyrer CM, Greene-Schloesser DM. Neuroanatomical target theory as a predictive model for radiation-induced cognitive decline. Neurology. 2013;80:747–53.
- Moore HC. An overview of chemotherapy-related cognitive dysfunction, or 'chemobrain'. Oncology. 2014;28:797–804.
- Hermelink K, Küchenhoff H, Untch M, Bauerfeind I, Lux MP, Bühner M, et al. Two different sides of 'chemobrain': determinants and nondeterminants of self-perceived cognitive dysfunction in a prospective, randomized, multicenter study. Psychooncology. 2010;19:1321–8.
- Schagen SB, Boogerd W, Muller MJ, Huinink WT, Moonen L, Meinhardt W, et al. Cognitive complaints and cognitive impairments following BEP chemotherapy in patients with testicular cancer. Acta Oncol. 2008;47:63–70.
- Taillia H, Bompaire F, Jacob J, Noël G. Cognitive evaluation during brain radiotherapy in adults: a simple assessment is possible. Cancer Radiother. 2013;17:413–8.
- Ricard D, Idbaih A, Ducray F, Lahutte M, Hoang-Xuan K, Delattre JY. Primary brain tumors in adults. Lancet. 2012;379:1984–96.
- Mackinnon A, Mullingan R. Estimation de l'intelligence prémorbide chez les francophones. Encéphale. 2005;31:31–43.
- Kalafat M, Hugonot-Diener L, Poitrenaud J. Standardisation et étalonnage français du "Mini Mental State" (MMS) version GRECO. Rev Neuropsychol. 2003;13:209–36.
- Nasreddine Z, Phillips N, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment (moca): a brief screening tool for middle cognitive impairment. J Am Ger Soc. 2005;53:695–99.
- 41. Mattis M. Dementia Rating Scale-2. Lutz: Psychological assessment resource. Inc; 2001.
- Van Der Linden M, Coyette F, Poitrenaud J, Kalafat M, Calicis F, Wyns C, et al. L'épreuve du rappel libre/rappel indicé à 16 items (RL/RI-16). L'évaluation des troubles de la mémoire. Marseille: Editions Solal; 2004. p. 85–101.
- Rey A. L'examen psychologique dans les cas d'encéphalopathie traumatique. Arch Psychol. 1941;8:286–340.
- Wechsler D. Echelle d'intelligence de Wechsler pour adultes 3e edition ECPA. 2000.
- Meulmans T. La batterie GREFEX. In: Fonctions executives et pathologies neurologiques et psychiatriques. Godefroy et GREFEX, éditeurs. Marseille, Solal; 2008. p. 217–29.
- Roussel M, Godefroy O. La batterie GREFEX: données normative Godefroy O et GREFEX, éditeurs. Marseille, Solal; 2008. p. 217–29.
- 47. Tuillard-Colombo F, Assal G. Boston Naming Test: French-language adaptation and short forms. Eurn Rev Appl Psychol. 1992;42:67–73.
- 48. De Renzi E, Faglioni P. Normative data and screening power of a shortened version of the Token Test. Cortex. 1978;14:41–9.
- Goldberg D, Bridges K, Duncun-Jones P, Grayson D. Detecting anxiety and depression in general medical setting. BMJ. 1988;297:897–9.
- Furher R, Rouillon F. La version française de l'échelle CES-D (Center for Epidemiologic Study Depression Scale). Description et traduction de l'échelle d'auto-évaluation. Psychiat Psychobiol. 1989;4:163–66.
- Cella D, Yount S, Sorensen M, Chartash E, Sengupta N, Grober J. Validation of the functional assessment of chronic illness therapy fatigue relative to other instrumentation in patients with rheumatoid arthritis. J Rheumatol. 2005;32:811–9.
- McNair DM, Kahn RJ. Self assessment of cognitive deficit. In: Crook Th. And Coll. Assessment in geriatric psychopharmacology. Mark Powley Associates Ed, New Canaan-Connecticut; 1983. p. 137–43.
- Poitrenaud J, Israel L, Barrere H, le Roc'h. Version française de l'échelle de difficulties cognitive de Mac-Nair et Khan. In: de la plainte mnésique à la maladie d'Alzheimer. Michel BF, Derouesne C, Gely-Nargeot MC, Eds. Marseille, Solal; 1997.
- Mac Lachlan S, Devins G, Goodwin P. Validation of European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (QLQ-C30) as a measure of psychosocial function in breast cancer patients. Eur J Cancer. 1998;34:510–7.

- Taphoorn MJ, Claasens L, Aaronson NK, Coens C, Mauer M, Osoba D, et al. An international validation study of the EORTC brain cancer module (EORTC QLQ-BN20) for assessing health-related quality of life and symptoms in brain cancer patients. Eur J Cancer. 2010;46:1033–40.
- Israel L. L'IADL. In: L'évaluation Clinique standardisée en psychitrie (tome 2). Guelfi, Ed. Editions médicales Pierre Fabre, 1996. p. 477–80.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at www.biomedcentral.com/submit

