



HAL
open science

RESCALED ENTROPY OF CELLULAR AUTOMATA

David Burguet

► **To cite this version:**

| David Burguet. RESCALED ENTROPY OF CELLULAR AUTOMATA. 2020. hal-02610374v1

HAL Id: hal-02610374

<https://hal.science/hal-02610374v1>

Preprint submitted on 16 May 2020 (v1), last revised 8 Jul 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESCALED ENTROPY OF CELLULAR AUTOMATA

DAVID BURGNET

ABSTRACT. For a d -dimensional cellular automaton with $d \geq 1$ we introduce a rescaled entropy which estimates the growth rate of the entropy at small scales by generalizing previous approaches [1, 9]. We also define a notion of Lyapunov exponent and proves a Ruelle inequality as already established for $d = 1$ in [16, 15]. Finally we generalize the entropy formula for 1-dimensional permutative cellular automata [18] to the rescaled entropy in higher dimensions. This last result extends recent works [17] of Shinoda and Tsukamoto dealing with the metric mean dimensions of two-dimensional symbolic dynamics.

1. INTRODUCTION

In this paper we estimate the dynamical complexity of multidimensional cellular automata. In the following the main results will be stated in a more general setting, but let us focus in this introduction on the following algebraic cellular automaton on $(\mathbb{F}_p)^{\mathbb{Z}^d}$ with p prime given for some finite family $(a_i)_{i \in I}$ in \mathbb{F}_p^* by

$$\forall (x_j)_j \in (\mathbb{F}_p)^{\mathbb{Z}^d}, f((x_j)_j) = \left(\sum_{i \in I} a_i x_{i+j} \right)_j.$$

Let $I' = I \cup \{0\}$. For $d = 1$ the topological entropy of f is finite and equal to $\text{diam}(I') \log p$ where $\text{diam}(I')$ denotes the diameter of I' for the usual distance on \mathbb{R} [18]. However in higher dimensions the topological entropy of f is always infinite unless f is the identity map [13, 10]. Moreover the topological entropy of the \mathbb{Z}^{d+1} -action given by f and the shift vanishes. In this paper we investigate the growth rate of $(h_{\text{top}}(f, \mathcal{P}_{J_n}))_n$ for nondecreasing sequences (J_n) of convex subsets of \mathbb{R}^d where $(\mathcal{P}_{J_n})_n$ denotes the clopen partitions into J_n -coordinates with $J_n := J_n \cap \mathbb{Z}^d$. This sequence appears to increase as the perimeter $p(J_n)$ of J_n . We define the rescaled entropy $h_{\text{top}}^d(f)$ of f as $\limsup_{J_n} \frac{h_{\text{top}}(f, \mathcal{P}_{J_n})}{p(J_n)}$. In [9] another renormalization is used, whereas in [1] the authors only investigate the case of squares $J_n = [-n, n]^2$, $n \in \mathbb{N}$. For $d = 1$ we get $h_{\text{top}}^1(f) = \frac{h_{\text{top}}(f)}{2}$. We generalize the entropy formula for algebraic cellular automata as follows :

Theorem 1. *Let f be an algebraic cellular automaton on $(\mathbb{F}_p)^{\mathbb{Z}^d}$ as above, then*

$$h_{\text{top}}^d(f) = R_{I'} \log p,$$

where $R_{I'}$ denotes the radius of the smallest bounding sphere containing I' .

In fact we establish such a formula for any permutative cellular automaton (see Section 7). In [17] the authors compute, inter alia, the metric mean dimension of the horizontal shift in \mathbb{Z}^2 for some standard distances. These dimensions may be interpreted as the rescaled entropy with respect to some particular sequence of convex sets $(J_n)_n$. In particular we extend these results in higher dimensions for general permutative cellular automata.

Date: June 2018.

2010 *Mathematics Subject Classification.* 37B15, 37A35, 52C07.

We also consider a measure theoretical analogous quantity of the rescaled entropy. In dimension one, a notion of Lyapunov exponent has been defined in [15]. Then Tisseur [16] proved in this case a Ruelle inequality relating this exponent with the Kolmogorov-Sinai entropy. In this paper we also introduce a notion of Lyapunov exponent in higher dimensions, which bounds from above the rescaled entropy of measures.

The paper is organized as follows. In Section 2 we state some measure geometrical properties of convex sets in \mathbb{R}^d . We recall the dynamical background of cellular automata in Section 4 and we introduce then a Lyapunov exponent for multidimensional cellular automata. In Section 5 we define and study the topological and measure theoretical rescaled entropy. The last section is devoted to the proof of the entropy formula for permutative cellular automata.

2. BACKGROUND ON CONVEX GEOMETRY

2.1. Convex bodies, domains and polytopes. For a fixed positive integer d we endow the vector space \mathbb{R}^d with its usual Euclidean structure. The associated scalar product is simply denoted by \cdot and we let \mathbb{S}^d be the unit sphere. For a subset F of \mathbb{R}^d we let \bar{F} , $\text{Int}(F)$ and ∂F be respectively its closure, interior set and boundary. We denote by $\underline{F} = F \cap \mathbb{Z}^d$ the set of integer points in F , i.e. $\underline{F} = F \cap \mathbb{Z}^d$. We also denote by $V(F)$ the d -Lebesgue measure of F (also called the volume of F) when the set F is Borel.

The extremal set of a convex set J is denoted by $\text{ex}(J)$ and the convex hull of $F \subset \mathbb{R}^d$ by $\text{cv}(F)$. A convex body is a compact convex set of \mathbb{R}^d . A convex body containing the origin $0 \in \mathbb{R}^d$ in its interior set is said to be a **convex domain**. The set of convex bodies endowed with the Hausdorff topology is a locally compact metrizable space. In the following we denote by \mathcal{D} , resp. \mathcal{D}^1 , the set of convex domains, resp. with unit perimeter, endowed with the Hausdorff topology. A **convex polytope** (resp. k -polytope with $k \leq d$) in \mathbb{R}^d is a convex body given by the convex hull of a finite set (resp. with topological dimension equal to k). When this set lies inside the lattice \mathbb{Z}^d , the convex polytope is said **integral**. We let $\mathcal{F}(P)$ be the set of faces of a convex polytope P . For a convex body J we denote by \mathbb{J} the integral polytope given by the convex hull of integer points in J , i.e. $\mathbb{J} = \text{cv}(\underline{J})$.

A convex domain J has Lipschitz boundary and finite perimeter $p(J)$. For convex domains the perimeter in the distributional sense of De Giorgi coincides with the $(d-1)$ -Hausdorff measure \mathcal{H}_{d-1} of the boundary. For $J \in \mathcal{D}$ we let $\partial'J$ be the subset of points $x \in \partial J$, where the tangent space $T_x J$ is well defined. The set $\partial'J$ has full \mathcal{H}_{d-1} -measure in ∂J . We let $N^J(x) \in \mathbb{S}^d$ be the unit J -external normal vector at $x \in \partial'J$. For any $x \in \partial'J$ we let $T_x^+ J$ (resp. $T_x^- J$) be the open external (resp. closed internal) semi-space with boundary $T_x J$. With these notations we have $J = \bigcap_{x \in \partial'J} T_x^- J$. For $\epsilon \in \mathbb{R}$ we denote by $T_x^\pm J(\epsilon)$ the semi-planes $T_x^\pm J(\epsilon) = T_x^\pm J + \epsilon N^J(x)$. When J is a convex polytope and $F \in \mathcal{F}(J)$, we write T_F to denote the tangent affine space supporting F , T_F^\pm for the associated semi-spaces and N^F for the unit external normal to F .

The **support function** of a convex body I is the real continuous function h_I on \mathbb{S}^d :

$$\forall x \in \mathbb{S}^d, h_I(x) = \max_{u \in I} u \cdot x.$$

The support function completely characterizes the convex body I . The **area measure** σ_J of a convex domain J is the Borel measure on \mathbb{S}^d given by $N_*^J \mathcal{H}_{d-1}$:

$$\forall B \text{ Borel of } \mathbb{S}^d, \sigma_J(B) = \mathcal{H}_{d-1}((N^J)^{-1}B).$$

If a sequence $(J_n)_n$ in \mathcal{D} is converging to $J_\infty \in \mathcal{D}$ (for the Hausdorff topology), then σ_{J_n} is converging weakly to σ_{J_∞} , in particular the perimeter of J_n goes to the perimeter of J_∞ (see Proposition 10.2 in [7]).

2.2. Convex exhaustions. We consider sequences $\mathcal{J} = (J_n)_{n \in \mathbb{N}}$ of convex domains with $p(J_n) \xrightarrow{n} +\infty$, such that the sets $\tilde{J}_n = p(J_n)^{-\frac{1}{d-1}} J_n \in \mathcal{D}^1$ are converging to a limit $J_\infty \in \mathcal{D}$ in the Hausdorff topology. In particular $\bigcup_n J_n = \mathbb{R}^d$. Moreover the limit J_∞ has unit perimeter.

The sequences $\mathcal{J} = (J_n)_n$ satisfying the above properties are said to be **convex exhaustions**. For $O \in \mathcal{D}^1$ we denote by $\mathcal{E}(O)$ the set of convex exhaustions $\mathcal{J} = (J_n)_n$ with $J_\infty = O$. Moreover for $O \in \mathcal{D}$ we let $\mathcal{J}_O \in \mathcal{E}(p(O)^{-1}O)$ be the convex exhaustion given by $\mathcal{J}_O := (nO)_n$. A convex exhaustion is said integral when J_n is an integral polytope for all n .

Lemma 1. *When $\mathcal{J} = (J_n)_n$ is a convex exhaustion, the associated sequence of integral convex polytopes $(J_n)_n$ defines an integral convex exhaustion with the same limit $J_\infty = J_\infty$ and $p(J_n) \sim^n p(J_\infty)$ when n goes to infinity.*

Proof. As $J_n \subset J_n$ the monotonicity of the perimeter of convex domains implies $p(J_n) \leq p(J_n)$ (see e.g. Chapter 7 in [3]). Then for any $x \in \text{Int}(J_\infty)$ there is a ball $B_x \subset J_\infty$ centered at x such that \tilde{J}_n contains B_x for n large enough. Moreover the rescaled ball $p(J_n)^{\frac{1}{d-1}} B_x \subset J_n$ has a non empty intersection with \mathbb{Z}^d when n goes to infinity. Therefore there is $u_n \in J_n$ with $p(J_n)^{-\frac{1}{d-1}} u_n \in B_x$ for large n . Let $(E_p)_{p \in \mathbb{N}}$ be a sequence of finite subsets of $\text{Int}(J_\infty)$ with $d_H(E_p, J_\infty) < \frac{1}{p}$, where d_H denotes the Hausdorff distance. Fix p . Arguing as above for any $x \in E_p$ with B_x of radius less than $\frac{1}{p}$, we get $d_H(J_\infty \cap p(J_n)^{-\frac{1}{d-1}} J_n, J_\infty) < \frac{2}{p}$ for n large enough. Therefore $J_\infty \cap p(J_n)^{-\frac{1}{d-1}} J_n$ goes to J_∞ , so that we obtain by taking the perimeters in this limit

$$\begin{aligned} \liminf_n \frac{p(J_n)}{p(J_n)} &= \liminf_n p\left(p(J_n)^{-\frac{1}{d-1}} J_n\right), \\ &\geq \lim_n p\left(J_\infty \cap p(J_n)^{-\frac{1}{d-1}} J_n\right), \\ &\geq p(J_\infty) = 1. \end{aligned}$$

Together with $p(J_n) \leq p(J_n)$, we get $p(J_n) \sim^n p(J_n)$. Now we observe that $J_\infty \cap \tilde{J}_n \supset J_\infty \cap p(J_n)^{-\frac{1}{d-1}} J_n$ is also converging to J_∞ . Moreover, it follows from the inclusions $J_n \subset J_n$ that *

$$\begin{aligned} \limsup_n \tilde{J}_n &= \limsup_n p(J_n)^{-\frac{1}{d-1}} J_n, \\ &\subset \lim_n \tilde{J}_n = J_\infty. \end{aligned}$$

Consequently \tilde{J}_n also goes to J_∞ when n goes to infinity. \square

2.3. Internal and external morphological boundary. We recall some terminology of mathematical morphology used in image processing. For two subsets I and J of \mathbb{R}^d , the **dilation** (also known as the Minkowski sum) $J \oplus I$ and the **erosion** $J \ominus I$ of J by I are defined as follows

$$\begin{aligned} J \oplus I &= \{i + j \mid i \in I \text{ and } j \in J\}, \\ J \ominus I &= \{j \in \mathbb{R}^d \mid \forall i \in I, i + j \in J\}. \end{aligned}$$

When the origin 0 belongs to I then we have $J \subset J \oplus I$ and $J \ominus I \subset J$. Assume now moreover that I and J are convex bodies. The dilation $J \oplus I$ is then also a convex body with $\text{ex}(J \oplus I) \subset \text{ex}(I) \oplus \text{ex}(J)$. In particular, when I and J are moreover convex polytopes, then so is $J \oplus I$. We have $J \ominus I = \bigcap_{x \in \partial' J} T_x^- J(h_I(-N^J(x)))$ (also $J \oplus I \subset \bigcap_{x \in \partial' J} T_x^- J(h_I(N^J(x)))$), but this last inclusion may be strict). Consequently $J \ominus I$ is a convex body. When J is a convex polytope, the above intersection is finite, thus $J \ominus I$ is also a convex polytope. The convex bodies given by the erosion $J \ominus I$ and the dilation $J \oplus I$ are also known as the inner and outer parallel bodies of J relative to I . We recall that $h_{J \oplus I} = h_J + h_I$. In particular when $I = \{i\}$ is a singleton, we get $h_{J+i}(x) = h_J(x) + i \cdot x$ for all $x \in \mathbb{S}^d$. In general we only have $h_{J \ominus I} \leq h_J - h_I$.

*For a sequence $(A_n)_n$ of subsets in \mathbb{R}^d , we let $\limsup_n A_n = \bigcap_n \overline{\bigcup_{k \geq n} A_k}$.

The **internal and external (morphological) boundaries** of J relative to I denoted respectively by $\partial_I^- J$ and $\partial_I^+ J$ are given by

$$\begin{aligned}\partial_I^+ J &= (I \oplus J) \setminus J, \\ \partial_I^- J &= J \setminus (J \ominus I).\end{aligned}$$

Clearly we have $\partial_I^\pm J = \partial_{I'}^\pm J$ with $I' = I \cup \{0\}$. When J is a convex domain then we have $\partial_I^- J = \partial_{\text{cv}(I)}^- J$ and $\partial_I^+ J \subset \partial_{\text{cv}(I)}^+ J$. In the following the set I will be fixed so that we omit the index I in the above definitions when there is no confusion.

3. COUNTING INTEGER POINTS IN MORPHOLOGICAL BOUNDARY OF LARGE CONVEX SETS

For a large convex domain J and a fixed integral polytope I we estimate the cardinality of the integer points in the morphological boundaries of J relative to I . We first compare the cardinality of integer points in the internal and external boundaries of J and \mathbb{J} . Recall that \underline{F} denotes the integer points in a subset set F of \mathbb{R}^d and $\mathbb{J} = \text{cv}(\mathbb{J})$.

Lemma 2. *With the above notations we have*

$$\underline{\partial^- \mathbb{J}} = \underline{\partial^- J}$$

and

$$\underline{\partial^+ \mathbb{J}} \subset \underline{\partial^+ J}.$$

In general the last inclusion is strict.

Proof. For any convex domain J , a point u of J belongs to $\partial^- J$ if and only if there is v in $\text{ex}(I)$ such that $u+v$ does not lie in J . As $J \cap \mathbb{Z}^d = \mathbb{J} \cap \mathbb{Z}^d$ and $\text{ex}(I) \subset \mathbb{Z}^d$, we get $\underline{\partial^- J} = \underline{\partial^- \mathbb{J}}$. Similarly if a point $u \in \partial^+ J$ is an integer, then $u \in J \oplus I$ but $u \notin J$. Therefore we get $\underline{\partial^+ \mathbb{J}} \subset \underline{\partial^+ J}$. \square

Lemma 3. *Let J be a convex polytope.*

$$\#\underline{\partial^- J} \leq \#\underline{\partial^+ J}.$$

Proof. We have $\partial^- J \subset \bigcup_{F \in \mathcal{F}(J)} T_F^+ J(-h_I(N^F))$. For $F \in \mathcal{F}(J)$ there exists $u^F \in \text{ex}(I)$ with $h_I(N^F) = u^F \cdot N^F$. Let F_1, \dots, F_N be an enumeration of $\mathcal{F}(J)$. Let $\phi : \partial^- J \rightarrow \partial^+ J$ be the function defined by $\phi(x) = x + u^{F_1}$ for $x \in S_1 := \partial^- J \cap T_{F_1}^+ J(-h_I(N^{F_1}))$ and $\phi(x) = x + u^{F_l}$ for $x \in S_l := \partial^- J \cap T_{F_l}^+ J(-h_I(N^{F_l})) \setminus \bigcup_{k < l} T_{F_k}^+ J(-h_I(N^{F_k}))$ by induction on l .

This map is injective : indeed if $\phi(x) = \phi(y)$ either x and y lie in the same S_l and then $\phi(x) = x + u^{F_l} = y + u^{F_l} = \phi(y)$ clearly implies $x = y$ or $x \in S_k, y \in S_l$ with $k \neq l$. We may assume $k < l$ without loss of generality. Then $y + u^{F_l} \in T_{F_k}^- J$ whereas $x + u^{F_k} \in T_{F_k}^+ J$ and we get thus a contradiction. Finally the map ϕ preserves the integer points since we have $\text{ex}(I) \subset \mathbb{Z}^d$. \square

3.1. First relative quermass integral. Let O be a convex domain and let I be a convex body. For $\rho \in \mathbb{R}$ we let

$$O_\rho = \begin{cases} O \oplus \rho I & \text{when } \rho \geq 0, \\ O \ominus \rho I & \text{when } \rho < 0. \end{cases}$$

Proposition 2.

$$\lim_{\rho \rightarrow 0} \frac{V(O_\rho) - V(O)}{\rho} = \int_{\mathbb{S}^d} h_I d\sigma_O.$$

For $\rho > 0$ the formula follows from Minkowski's formula on mixed volume (see Theorem 6.5 and Corollary 10.1 in [7]). For $\rho < 0$ we refer to [12] (see also Lemma 2 in [4] for the 2-dimensional case).

The quantity $d \int_{\mathbb{S}^d} h_I d\sigma_O$ is known as the **first I -relative quermass integral** of J . In the following we denote by $V_I(O)$ the integral $\int_{\mathbb{S}^d} h_I d\sigma_O$. For convex bodies $I \subset I'$ and $k \in \mathbb{N}$, we have $V_I(O) \leq V_{I'}(O)$ and $V_{kI}(O) = kV_I(O)$ for any convex domain O . The support function h_I being continuous, the first I -relative quermass integral of O is continuous with respect to the Hausdorff topology, i.e. if $(O_n)_n$ is a sequence of convex domains converging to a convex domain O_∞ in the Hausdorff topology, then we have

$$V_I(O_n) \xrightarrow{n \rightarrow +\infty} V_I(O_\infty).$$

We deduce now from Proposition 2 an estimate on the volume of the morphological boundary for large convex sets.

Corollary 3. *Let I be a convex body containing 0 and let $O \in \mathcal{D}$. Then*

$$V(\partial_I^\pm nO) \sim n^{d-1} \int_{\mathbb{S}^d} h_I d\sigma_O.$$

Proof. We only consider the case of the external boundary as one may argue similarly for the internal boundary. For all n we have

$$\begin{aligned} V(\partial_I^+ nO) &= V(nO \oplus I) - V(nO), \\ &= n^d (V(O \oplus n^{-1}I) - V(O)) \end{aligned}$$

According to Proposition 2 we conclude that

$$V(\partial_I^+ nO) \sim n^{d-1} \int_{\mathbb{S}^d} h_I d\sigma_O.$$

□

3.2. Counting integer points in large convex sets. Since Gauss circle problem counting lattice points in convex sets has been extensively investigated. Let $\mathbb{C} = [0, 1]^d$. Clearly for any Borel subset K of \mathbb{R}^d we have always

$$(3.1) \quad \#\underline{K} \leq V(K \oplus \mathbb{C}).$$

In the other hand, Bokowski, Hadwiger and Wills have proved the following general (sharp) inequality for any convex domain O [2] :

$$(3.2) \quad V(O) - \frac{p(O)}{2} \leq \#\underline{O}.$$

There exists precise asymptotic estimate of $\#\underline{xO}$ for large $x > 0$ for convex domains with smooth domain having positive curvature, in particular we have in this case $\#\underline{xO} = V(xO) + o(x^{d-1})$ [8].

3.3. First rough estimate for $\#\partial_I^\pm nO \cap \mathbb{Z}^d$ with $O \in \mathcal{D}$. For a real sequences $(a_n)_n$ and two numbers l and $C > 0$ we write $a_n \sim^C l$ when the accumulation points of $(a_n)_n$ lie in $[l - C, l + C]$.

Lemma 4. *There exists a constant C depending only on d such that we have for any convex domain O and any convex body I of \mathbb{R}^d with $0 \in I$:*

$$\frac{\#\partial_I^\pm nO}{n^{d-1}} \sim^C V_I(O).$$

Proof. We only argue for $\partial_I^+ O$, the other case being similar. We have $\#\partial_I^+ nO = \#\underline{nO \oplus I} - \#\underline{nO}$, and then by combining Equation (3.1) and (3.2) we get :

$$V(nO \oplus I) - \frac{p(nO \oplus I)}{2} - V(nO + \mathbb{C}) \leq \#\partial_I^+ nO \leq V(nO \oplus I \oplus \mathbb{C}) - V(nO) + \frac{p(nO)}{2},$$

After dividing by n^{d-1} , the right (resp. left) hand side term is going to $\int_{\mathbb{S}^d}(h_I - h_C - 1/2) d\sigma_O$ (resp. $\int_{\mathbb{S}^d}(h_I + h_C + 1/2) d\sigma_O$) according to Corollary 3. \square

3.4. Fine estimate of $\#\partial_I^\pm J_n$ for general convex exhaustion $(J_n)_n$ in dimension 2. We compare directly the cardinality of lattice points in the morphological boundary with the first I -relative quermass integral of J_∞ for two-dimensional convex exhaustion. This result will not be used directly in the next sections but is potentially of independent interest.

Proposition 4. *For any convex exhaustion $(J_n)_n$ in \mathbb{R}^2 , we have*

$$\lim_n \frac{\#\partial_I^- J_n}{p(J_n)} = V_I(J_\infty).$$

By Lemma 1 and Lemma 2 we only need to consider integral convex exhaustions. In fact in this case we also show the corresponding statement for the external morphological boundary.

Proposition 5. *For any integral convex exhaustion $(J_n)_n$ in \mathbb{R}^2 , we have*

$$\lim_n \frac{\#\partial_I^\pm J_n}{p(J_n)} = V_I(J_\infty).$$

The rest of this subsection is devoted to the proof of Proposition 5. We start by giving some preliminary lemmas.

We denote by $\angle P$ the minimum of the interior angles at the vertices of a convex polygon $P \subset \mathbb{R}^2$.

Lemma 5. *For any integral convex exhaustion $(J_n)_n$ in \mathbb{R}^2 , we have*

$$\liminf_n \angle J_n > 0.$$

Proof. We have $\angle \tilde{J}_n = \angle J_n$. Moreover the minimal angle is lower semi-continuous for the Hausdorff topology, therefore $\liminf_n \angle J_n \geq \angle J_\infty$. Since J_∞ has non-empty interior, we have $\angle J_\infty > 0$. \square

Lemma 6. *For any integral convex exhaustion $(J_n)_n$ in \mathbb{R}^2 , we have*

$$\#\mathcal{F}(J_n) = o(p(J_n)).$$

Proof. Two integral polytopes are said equivalent when there is a translation (necessarily by an integer) mapping one to the other. For any L the number a_L of equivalence classes of integral 1-polytopes with 1-Hausdorff measure less than L is finite (these polytopes are just line segments with integral endpoints and their 1-Hausdorff measure is just equal to their length). Moreover for a integral convex polytope there are at most two faces in the same class. Therefore

$$\begin{aligned} \#\mathcal{F}(J_n) &\leq 2a_L + \#\{F \in \mathcal{F}(J_n), \mathcal{H}_1(F) \geq L\}, \\ &\leq 2a_L + \frac{p(J_n)}{L}. \end{aligned}$$

This inequality holds for all n and $p(J_n)$ goes to infinity with n so that we conclude $\#\mathcal{F}(J_n) = o(p(J_n))$ as L was arbitrarily fixed. \square

Given two distinct points A, B in \mathbb{R}^2 and $h \neq 0$, the rectangle $R_{AB}(h)$ of basis AB and height $h > 0$ (resp. $h < 0$) is the semi-open rectangle $[AB[\times]A, D[$ oriented as $ABCD$ (resp. $ADCB$)[†] with $|AD| = |h|$. This rectangle is said integral when A, B belong to \mathbb{Z}^2 and the line (CD) has a non-empty intersection with \mathbb{Z}^2 .

[†]We denote a convex polytope with its vertices by respecting the usual orientation of the plane.

Lemma 7. *For any integral rectangle R ,*

$$\#R = V(R).$$

Proof. After a translation by an integer we may assume that the origin is the vertex A of the integral rectangle $R = R_{AB}(h)$. Let (p', q') be an integer on the line segment $[A, B]$ with p', q' relatively prime. By Bezout theorem there is $(u, v) \in \mathbb{Z}^2$ with $up + vq = 1$. Therefore there is a matrix $M \in SL_2(\mathbb{Z})$ with $M(p, q) = (k, 0)$. As the transformation M preserve both the volume and the integer points it is enough to consider the semi-open parallelogram $M(R)$. But there is a piecewise integral translation, which maps $M(R)$ to a semi-open integral rectangle with basis $M([A, B]) \subset \mathbb{R} \times \{0\}$. For such a rectangle the area is obviously equal to the cardinality of its integer points. \square

For $A, B \in \mathbb{R}^2$ and $\epsilon < \frac{|AB|}{2}$ we let A^ϵ and B^ϵ be the points in the line (AB) with Euclidean distance $|\epsilon|$ to A and B respectively, which lie inside $[A, B]$ if $\epsilon > 0$ and outside if not. As the symmetric difference of $R_{A^\epsilon B^\epsilon}(h)$ and $R_{AB}(h)$ is given by the union of two rectangles with sides of length $|\epsilon|$ and $|h|$ we have for some constant $C = C(|\epsilon|, |h|)$

$$(3.3) \quad \left| \#R_{A^\epsilon B^\epsilon}(h) - \#R_{AB}(h) \right| \leq C.$$

This estimate still holds true for $\epsilon \geq |AB|/2$ when choosing the convention $R_{A^\epsilon B^\epsilon}(h) = \emptyset$ for such ϵ .

Fact. *For any convex body I and for any $a > 0$, there exists $\epsilon^+ = \epsilon^+(I) > 0$ and $\epsilon^- = \epsilon^-(I, a) > 0$ such that any convex polytope $J = A_1 \cdots A_n$ with $\angle J \geq a$ satisfies*

$$\partial^+ J \subset \bigcup_{l < n} R_{A_l^{\epsilon^+} A_{l+1}^{-\epsilon^+}}(-h_I(N^{A_l A_{l+1}}))$$

and

$$\partial^- J \supset \bigcup_{l < n} R_{A_l^{\epsilon^-} A_{l+1}^{-\epsilon^-}}(h_I(N^{A_l A_{l+1}})).$$

This fact is illustrated on Figure 1 and its easy proof is left to the reader. We are now in a position to prove Proposition 5.

Proof of Proposition 5. From the above fact and (3.3) there is $\epsilon = \epsilon^-(I, \angle J) > 0$ and $C = C(I, \angle J) > 0$ such that for any convex polytope $J = A_1 \cdots A_n$

$$\begin{aligned} \#\partial^- J &\geq \sum_{l < n} \#R_{A_l(\epsilon) A_{l+1}(\epsilon)}(-h_I(N^{A_l A_{l+1}})), \\ &\geq \sum_{F \in \mathcal{F}(J)} \left[\#R_F(-h_I(N^F)) - C \right]. \end{aligned}$$

Then when J is an integral convex polytope we get by Lemma 7 :

$$\begin{aligned} \#\partial^- J &\geq -C\#\mathcal{F}(J) + \sum_{F \in \mathcal{F}(J)} V(R_F(-h_I(N^F))), \\ &\geq -C\#\mathcal{F}(J) + \int_{\mathbb{S}^d} h_I d\sigma_J. \end{aligned}$$




Figure 1: *The external and internal rectangles associated to a face F of a polygon. The external and internal morphological boundaries are respectively represented by the areas in yellow and green. The rectangles, R_F^+ and R_F^- , given by Fact 3.4 are drawn in blue.*

For an integral convex exhaustion $(J_n)_n$ we obtain finally for large n by using Lemma 6 and Lemma 5

$$\begin{aligned} \# \partial^- J_n &\geq -C(I, \frac{\angle J_\infty}{2}) \cdot \# \mathcal{F}(J_n) + \int_{\mathbb{S}^d} h_I d\sigma_{J_n}, \\ \liminf_n \frac{\# \partial^- J_n}{p(J_n)} &\geq \lim_n \int_{\mathbb{S}^d} h_I d\sigma_{J_n}, \\ &\geq \int_{\mathbb{S}^d} h_I d\sigma_{J_\infty}. \end{aligned}$$

One proves similarly that $\limsup_n \frac{\# \partial^+ J_n}{p(J_n)} \leq \int_{\mathbb{S}^d} h_I d\sigma_{J_\infty}$ and this concludes the proof as we have $\# \partial^+ J_n \geq \# \partial^- J_n$ according to Lemma 3. \square

3.5. Supremum of $O \mapsto V_I(O)$. In this section we investigate the supremum of V_I on \mathcal{D}^1 for a given convex body I of \mathbb{R}^d . We recall that there is a unique sphere S_I containing I with minimal radius, usually called the **smallest bounding sphere** of I . We let R_I and x_I be respectively the radius and the center of S_I . There are at least two distinct points in $S_I \cap I$. Moreover we have the following alternative :

- either there is a subset of $S_I \cap I$ generating an inscribable polytope T with $\text{Int}(T) \ni x_I$ (in particular the interior set of I is non empty),
- or there is a hyperplane H containing x_I such that I lies in an associated semiplane and $S_I \cap H$ is the smallest bounding sphere of $I \cap H$.

The smallest bounding sphere S_I will be said **nondegenerated** (resp. **degenerated**) and an associated polytope T (resp. hyperplane H) is said **generating**. For an inscribable polytope T in \mathbb{R}^d we may define its dual T' as the polytope given by the intersection of the inner semi-spaces tangent to the circumsphere of T at the vertices of T . In the following T' always denote the dual polytope of a generating polytope T with respect to I .

When S_I is degenerated and H is an associated generating hyperplane, we let T'_R , $R > 0$, be the polytope given by the intersection of the square $[-R, R]^d$ with the inner semiplanes tangent to S_I at $H \cap S_I \cap I$. We also denote by $\mathcal{F}_{\parallel}(T'_R)$ the subspace of $\mathcal{F}(T'_R)$ given by subfaces of the square $[-R, R]^d$.

Proposition 6.

$$\sup_{O \in \mathcal{D}^1} V_I(O) = R_I.$$

The supremum of V_I is achieved if and only if S_I is nondegenerated. The supremum is then achieved for $O \in \mathcal{D}^1$ homothetic to the dual polytope T' of a generating polytope T .

Proof. For any $v \in \mathbb{R}^d$ we have

$$\begin{aligned} V_{I+v}(O) &= \int h_{I+v} d\sigma_O, \\ &= \int h_I d\sigma_O + \int_{\mathbb{S}^d} v \cdot u d\sigma_O(u), \\ &= \int h_I d\sigma_O + \int_{\partial O} v \cdot N^O d\mathcal{H}_{d-1}. \end{aligned}$$

By the divergence formula we have $\int_{\partial O} v \cdot N^O d\mathcal{H}_{d-1} = 0$ for any $v \in \mathbb{R}^d$ and $O \in \mathcal{D}^1$. Therefore we may assume $x_I = 0$. With the above notations we have $\max_{i \in I} i \cdot v \leq R_I$ for all $v \in \mathbb{R}^d$ with $\|v\| = 1$ with equality iff v belongs to $R_I^{-1}I$. Therefore $V_I(O) \leq R_I$ for any $O \in \mathcal{D}^1$. Moreover if the equality occurs then for x in a subset E of ∂O with full \mathcal{H}_{d-1} -measure, $h_I(N^O(x)) = \max_{i \in I} i \cdot x = R_I$ and therefore the normal unit vector $N^O(x)$ belongs to $R_I^{-1}I$. But as O is a convex domain, we may find $d+1$ points x_1, \dots, x_{d+1} in E in such a way the origin belongs to the interior of the simplex $T = R_I \text{cv}(N^O(x_1), \dots, N^O(x_{d+1}))$. Thus S_I is nondegenerated and the polytope T is a generating polytope with respect to I . Moreover we have with the above notations

$$\int h_I d\sigma_{T'} = R_I p(T').$$

Therefore the homothetic tetrahedron of T' with unit perimeter achieves the supremum of V_I . We consider now the degenerated case. With the above notations we have $\mathcal{H}_{d-1}\left(\bigcup_{F \in \mathcal{F}_{\parallel}(T'_R)} F\right) = o(p(T'_R))$ when R goes to infinity. Therefore the renormalization $O_R \in \mathcal{D}^1$ of T'_R satisfies

$$V_I(O_R) \xrightarrow{R \rightarrow +\infty} R_I.$$

□

4. CELLULAR AUTOMATA

4.1. Definitions. We consider a finite set \mathcal{A} . We endow the set \mathcal{A} with the discrete topology and $X_d = \mathcal{A}^{\mathbb{Z}^d}$ with the product topology. We consider the \mathbb{Z}^d -shift σ on $\mathcal{A}^{\mathbb{Z}^d}$ defined for $l \in \mathbb{Z}^d$ and $u = (u_k)_k \in X_d$ by $\sigma^l(u) = (u_{k+l})_k$. Any closed subset X of X_d invariant under the action of σ is called a \mathbb{Z}^d -**subshift**. We fix such a subshift X in the remaining of the paper.

For a bounded subset J of \mathbb{R}^d we consider the partition \mathbb{P}_J into J -**cylinders**, i.e. the element \mathbb{P}_J^x of \mathbb{P}_J containing $x = (x_i)_{i \in \mathbb{Z}^d} \in X$ is given by $\mathbb{P}_J^x := \{y = (y_i)_{i \in \mathbb{Z}^d} \in X, \forall i \in J y_i = x_i\}$. In other terms we may define \mathbb{P}_J as the joined partition $\bigvee_{j \in J} \sigma^{-j} \mathbb{P}_0$ with \mathbb{P}_0 being the zero-coordinate partition.

A **cellular automaton** (CA for short) defined on a \mathbb{Z}^d -subshift X is a continuous map $f : X \rightarrow X$ which commutes with the shift action σ . By a famous theorem of Hedlund [14] the cellular automaton f is given by a local rule, i.e. there exists a finite subset I of \mathbb{Z}^d and a map $F : \mathcal{A}^I \rightarrow \mathcal{A}$ such that

$$\forall j \in \mathbb{Z}^d \quad (fx)_j = F((x_{j+i})_{i \in I}).$$

The (minimal) subset I is called the **domain** of the CA. Recall $I' = I \cup \{0\}$.

4.2. Lyapunov exponents for higher dimensional cellular automata. Lyapunov exponent of one-dimensional cellular automata have been defined in [15, 16]. We develop a similar theory in higher dimensions. Let f be a CA on a \mathbb{Z}^d -subshift X with domain I .

Given a convex body J of \mathbb{R}^d and $x \in X$, we let

$$\mathcal{E}_f^+(x, J) := \{K \text{ convex body, } f\mathbb{P}_K^x \subset \mathbb{P}_J^{fx}\}$$

$$\mathcal{E}_f^-(x, J) := \{K \text{ convex body, } f\mathbb{P}_J^x \subset \mathbb{P}_K^{fx}\}$$

The family $\mathcal{E}_f^+(x, J)$ has a minimal element for the inclusion given by $J \oplus f(x) := \bigcap \{K, K \in \mathcal{E}_f^+(x, J)\}$, but a priori this is not the case of $\mathcal{E}_f^-(x, J)$. Observe that $J \oplus I$ and $J \ominus I$ belongs respectively to $\mathcal{E}_f^+(x, J)$ and $\mathcal{E}_f^-(x, J)$. Then we let for all x :

$$\text{gr}_J^+ f(x) := \min\{\#K \setminus J, K \in \mathcal{E}_f^+(x, J)\}$$

$$\text{gr}_J^- f(x) := \min\{\#J \setminus K, K \in \mathcal{E}_f^-(x, J)\}.$$

Observe that the family $\mathcal{E}_f^-(x, J)$ and the function $\text{gr}_J^- f(x)$ are constant on each atom A of \mathbb{P}_J , then we let $\mathcal{E}_f^-(A, J)$ and $\text{gr}_J^- f(A)$ be these quantities. For a convex exhaustion $\mathcal{J} = (J_n)_n$ and a convex domain $O \in \mathcal{D}^1$, we define **the growth** $\text{gr}_{\mathcal{J}}^\pm f$ with respect to \mathcal{J} and O as the following real functions on X :

$$\begin{aligned} \text{gr}_{\mathcal{J}}^\pm f &:= \limsup_n \frac{\text{gr}_{J_n}^\pm f}{p(J_n)}, \\ \text{gr}_O^\pm f &= \sup_{\mathcal{J} \in \mathcal{E}(O)} \text{gr}_{\mathcal{J}}^\pm f. \end{aligned}$$

Lemma 8.

$$\forall O \in \mathcal{D}^1, \text{gr}_O^+ f = \text{gr}_O^- f.$$

Proof. It follows from the definitions that we have for any convex bodies K, J :

$$(4.1) \quad [K \in \mathcal{E}^+(x, J)] \Leftrightarrow [J \in \mathcal{E}^-(x, K)].$$

Then if $\mathcal{J} = (J_n)_n$ is a convex exhaustion and $\mathcal{K}^\pm = (K_n^\pm)_n$ is a sequence of convex bodies K_n which realizes the maximum in the definition of $\text{gr}_{J_n}^\pm f$. As $J \oplus I$ and $J \ominus I$ belong respectively to $\mathcal{E}^\pm(x, J)$ the sequence \mathcal{K}^\pm is a convex exhaustion with $p(K_n^\pm) \sim^n p(J_n)$. By the equivalence (4.1) we get

$$\begin{aligned} \text{gr}_{\mathcal{J}}^+ f &\leq \limsup_n \frac{\#K_n^+ \setminus J_n}{p(J_n)}, \\ &\leq \limsup_n \frac{\#K_n^+ \setminus J_n}{p(K_n^+)}, \\ &\leq \text{gr}_{\mathcal{K}^+}^- f, \end{aligned}$$

therefore we get $\text{gr}_O^+ f \leq \text{gr}_O^- f$ and then $\text{gr}_O^+ f \geq \text{gr}_O^- f$ by following the same lines. \square

In the following we let $\text{gr}_O f = \text{gr}_O^\pm f$.

Lemma 9. *The sequence of functions $(\text{gr}_O f^k)_k$ is subadditive, i.e.*

$$\forall k, l \in \mathbb{N} \quad \forall x \in X, \text{gr}_O f^{k+l}(x) \leq \text{gr}_O f^l(f^k x) + \text{gr}_O f^k(x).$$

Proof. Fix $x \in X$ and $k, l \in \mathbb{N}$. Let $\mathcal{J} = (J_n)_n \in \mathcal{E}(O)$. As already observed, the sequences $\mathcal{J}' = (J'_n)_n := (J_n \oplus f^l(f^k x))_n$ and $\mathcal{J}'' = (J''_n)_n := (J'_n \oplus f^l(x))_n$ both belong to $\mathcal{E}(O)$ and their perimeters are equivalent to $p(J_n)$, when n goes to infinity. Moreover for all positive integers n we have

$$\begin{aligned} f^{k+l} \mathbf{P}_{J''_n}^x &= f^l(f^k \mathbf{P}_{J''_n}^x), \\ &\subset f^l \left(\mathbf{P}_{J'_n}^{f^k x} \right), \\ &\subset \mathbf{P}_{J_n}^{f^{k+l} x}. \end{aligned}$$

Therefore we conclude

$$\begin{aligned} \text{gr}_{J_n}^+ f^{k+l}(x) &\leq \# \underline{J''_n} \setminus J_n, \\ &\leq \# \underline{J''_n} \setminus J'_n + \# \underline{J'_n} \setminus J_n, \\ \text{gr}_{\mathcal{J}}^+ f^{k+l}(x) &\leq \text{gr}_{\mathcal{J}'}^+ f^k(x) + \text{gr}_{\mathcal{J}'}^+ f^l(f^k x), \\ \text{gr}_O f^{k+l}(x) &\leq \text{gr}_O f^k(x) + \text{gr}_O f^l(f^k x). \end{aligned}$$

□

For $d = 2$ the nonnegative function $\text{gr}_O f$ is bounded from above by $V_I(O)$ according to Proposition 4. Therefore the subadditive ergodic theorem applies : for any $\mu \in \mathcal{M}(X, f)$ the sequence $(\frac{1}{n} \text{gr}_O f^n(x))_k$ converge almost everywhere to a f -invariant function χ_O with $\int \chi_O d\mu = \lim / \inf_n \frac{1}{n} \int \text{gr}_O f^n d\mu$. We call the function χ_O the Lyapunov exponent of f with respect to O .

Remark 7. *The exponent χ_O for $O \in \mathcal{D}$ plays some how the role of the sum of the positive Lyapunov exponents in smooth dynamical systems.*

5. RESCALED ENTROPY OF CELLULAR AUTOMATA

5.1. Definition. We let $\mathcal{M}(f)$ (resp. $\mathcal{M}(f, \sigma)$) be the set of invariant Borel probability measures on X which are f -invariant (resp. f - and σ -invariant). For a finite clopen partition \mathbf{P} of X we let $H_{top}(\mathbf{P}) = \log \# \mathbf{P}$ and $H_\mu(\mathbf{P}) = - \sum_{A \in \mathbf{P}} \mu(A) \log \mu(A)$ with $\mu \in \mathcal{M}(f)$. In the following the symbol \ast denotes either $\ast = top$ or $\ast = \mu \in \mathcal{M}(f)$. We let $h_\ast(f, \mathbf{P})$ be the entropy with respect to the clopen partition \mathbf{P} :

$$h_\ast(f, \mathbf{P}) := \lim_n \frac{1}{n} H_\ast \left(\bigvee_{k=0}^{n-1} f^{-k} \mathbf{P} \right).$$

For two partitions \mathbf{P}, \mathbf{Q} of X , we say \mathbf{P} is finer than \mathbf{Q} and we write $\mathbf{P} > \mathbf{Q}$, when any atom of \mathbf{P} is contained in an atom of \mathbf{Q} . The functions $H_\ast(\cdot)$ and $h_\ast(f, \cdot)$ are nondecreasing with respect to this order.

The rescaled entropy with respect to a convex exhaustion $\mathcal{J} = (J_n)_n$ is defined as follows

$$h_\ast^d(f, \mathcal{J}) = \limsup_n \frac{h_\ast(f, \mathbf{P}_{J_n})}{p(J_n)}.$$

In [9] the authors defines a similar notion for the rescaled topological entropy with the renormalization factor $\# \partial_I^- J_n$ (which depends on the domain I of f) rather than $p(J_n)$.

Remark 8. When $J = \bigcup_{i \in I} J_i$ is a finite disjoint union of Jordan domains J_i with Lipschitz boundary, we have

$$\begin{aligned} \frac{h_{top}(f, P_J)}{p(J)} &\leq \frac{\sum_{i \in I} h_{top}(f, P_{J_i})}{\sum_{i \in I} p(J_i)}, \\ &\leq \sup_{i \in I} \frac{h_{top}(f, P_{J_i})}{p(J_i)}. \end{aligned}$$

Moreover for each i , we have $p(J_i) \geq p(\text{cv}(J_i))$ and $P_{\text{cv}(J_i)}$ is finer than P_{J_i} . Therefore

$$\begin{aligned} \frac{h_{top}(f, P_J)}{p(J)} &\leq \frac{\sum_{i \in I} h_{top}(f, P_{J_i})}{\sum_{i \in I} p(J_i)}, \\ &\leq \sup_{i \in I} \frac{h_{top}(f, P_{\text{cv}(J_i)})}{p(\text{cv}(J_i))}. \end{aligned}$$

This inequality justifies somehow that we focus on convex bodies J of \mathbb{R}^d .

We let also

$$h_{top}^d(f) = \sup_{\mathcal{J}} h_*^d(f, \mathcal{J}).$$

For $d = 1$ we have $p(J) = 2$ for any convex subset J . Therefore up to a factor 2 we recover the usual definition of entropy, $2h_*^1(f) = h_*(f)$.

As the CA f commutes with the shift action σ we have for all $k \in \mathbb{Z}^d$ and any subset J of \mathbb{Z}^d $h_{top}(f, P_{J+k}) = h_{top}(f, \sigma^{-k}P_J) = h_{top}(f, P_J)$ and the same holds for the measure theoretical entropy with respect to measures in $\mathcal{M}(f, \sigma)$. In particular we have $h_{top}^d(f, O) = h_{top}^d(f, O + \alpha)$ for any $\alpha \in \mathbb{R}^d$. Also for any μ in $\mathcal{M}(f)$ and $k \in \mathbb{Z}^d$ we have $h_\mu^d(f, O) = h_{\sigma^k \mu}^d(f, O)$ because for $\mathcal{J} = (J_n)_n \in \mathcal{E}(O)$, the sequence $\sigma^k(\mathcal{J}) = (k + J_n)_n$ belongs to $\mathcal{E}(O)$.

Remark 9. (1) The partition P_{J_n} may be written as $\bigvee_{k \in J_n} \sigma^{-k}P_0$ with P_0 being the zero-coordinate partition. Instead of P_0 we could choose another clopen generating partition P , i.e. a partition of X into clopen sets with $\bigvee_{k \in \mathbb{Z}^d} \sigma^{-k}P$ equal to the partition of X into points. But for a finite subset J of \mathbb{Z}^d we have $\bigvee_{k \in J} \sigma^{-k}P > P_0$ and $\bigvee_{k \in J} \sigma^{-k}P_0 > P$ so that in the definition of the rescaled entropy we may replace P_0 by any other generator P of X , i.e. P_{J_n} by $\bigvee_{k \in J_n} \sigma^{-k}P$.

(2) Let X be a zerodimensional compact metrizable space endowed with a expansive \mathbb{Z}^d -action τ . We consider a map f preserving (X, τ) i.e. f is an homeomorphism of X commuting with τ . The triple (X, τ, f) is called a topological \mathbb{Z}^d -expansive preserving system (t.e.p.s. for short). Two t.e.p.s. (Y, ϕ, g) are conjugated when there is a homeomorphism $h : X \rightarrow Y$ such that $h \circ f \circ h^{-1} = g$ and $h \circ \tau \circ h^{-1} = \phi$. We may define the rescaled entropy as we did for a CA and all the previous results hold in this more general setting. Moreover two conjugated t.e.p.s. have the same rescaled entropy. Any t.e.p.s. is conjugated to a CA.

5.2. Link with the metric mean dimension. In a compact metric space (X, d) , the ball of radius $\epsilon \geq 0$ centered at $x \in X$ will be denoted by $B_d(x, \epsilon)$. For a continuous map $f : X \rightarrow X$ we denote by d_n the dynamical distance defined for all $n \in \mathbb{N}$ by

$$\forall x, y \in X, d_n(x, y) = \max\{d(f^k x, f^k y), 0 \leq k < n\}.$$

The metric mean dimension of f is defined as $\text{mdim}(f, d) = \limsup_{\epsilon \rightarrow 0} \frac{h_{top}(f, \epsilon)}{|\log \epsilon|}$ where $h_{top}(f, \epsilon)$ denotes the topological entropy at the scale $\epsilon > 0$:

$$h_{top}(f, \epsilon) := \limsup_n \frac{1}{n} \log \min\{\#C, \bigcup_{x \in C} B_{d_n}(x, \epsilon) = X\}.$$

The topological mean dimension is the infimum of $\text{mdim}(f, \mathbf{d})$ over all distances on X . We refer to [11] for alternative definitions and further properties of mean dimension. The topological mean dimension of a finite dimensional topological system is null.

Here f is a CA on a subshift of \mathbb{Z}^d . In particular it has zero topological mean dimension. For a norm $\|\cdot\|$ of \mathbb{R}^d we may associate a metric $\mathbf{d}_{\|\cdot\|}$ on X_d by letting $\mathbf{d}_{\|\cdot\|}(u, v) = \alpha^{-\min\{\|k\|, k \in \mathbb{Z}^d, u_k \neq v_k\}}$ for all $u = (u_k)_k, v = (v_k)_k \in X_d$. Then for $l \in \mathbb{N}$ the (open) ball $B_{\mathbf{d}_{\|\cdot\|}}(x, 2^{-l})$ with respect to $\mathbf{d}_{\|\cdot\|}$ coincides with the cylinder $\mathbb{P}_{J_l}^x$ with $J_l = B_{\|\cdot\|}(0, l)$.

As there is a correspondence between convex symmetric bodies and unit balls of norms on \mathbb{R}^d , the mean dimension with respect to such distances $\mathbf{d}_{\|\cdot\|}$ are given by $h_{top}^d(f, \mathcal{J}_O)$ for convex symmetric bodies O .

Remark 10. In [17] the authors work with a measure theoretical quantity, called the measure distortion rate dimension and show a variational principle with the metric mean dimension of $\mathbf{d}_{\|\cdot\|}$. Does this quantity coincides with $\mu \mapsto h_\mu^d(f, O)$ with O being the symmetric convex domain associated to the norm $\|\cdot\|$?

5.3. Monotonicity and Power. We investigate now basic properties of the rescaled entropy.

Lemma 10. For any $O \in \mathcal{D}$ and any $\alpha > 0$, we have

$$h_*^d(f, \mathcal{J}_O) = h_*^d(f, \mathcal{J}_{\alpha O}).$$

Proof. For $n \in \mathbb{N}$, we let $k_n = \lceil \frac{n}{\alpha} \rceil$, thus $nO \subset k_n \alpha O$ and $p(nO) \sim^n p(k_n \alpha O)$. Therefore

$$\begin{aligned} h_*^d(f, \mathcal{J}_O) &= \limsup_n \frac{h_*(f, \mathbb{P}_{nO})}{p(nO)}, \\ &\leq \limsup_n \frac{h_*(f, \mathbb{P}_{k_n \alpha O})}{p(nO)}, \\ &\leq \limsup_n \frac{h_*(f, \mathbb{P}_{k_n \alpha O})}{p(k_n \alpha O)}, \\ &\leq h_*^d(f, \mathcal{J}_{\alpha O}). \end{aligned}$$

The other inequality is obtained by considering αO and α^{-1} in place of O and α . \square

Lemma 11. For any $O \in \mathcal{D}^1$ and $O' \in \mathcal{D}$ with $O \subset \text{Int}(O')$, we have

$$h_*^d(f, \mathcal{J}_O) \leq h_*^d(f, O) \leq p(O') h_*^d(f, \mathcal{J}_{O'}).$$

Proof. As $\mathcal{J}_O \in \mathcal{E}(O)$ the inequality $h_*^d(f, \mathcal{J}_O) \leq h_*^d(f, O)$ follows from the definitions. Let now $\mathcal{J} \in \mathcal{E}(O)$. For n large enough we have $\tilde{J}_n \subset \text{Int}(O')$, therefore $J_n \subset k_n O'$ with $k_n = ([p(J_n)] + 1)^{\frac{1}{d-1}}$. Therefore we conclude that

$$\begin{aligned} h_*^d(f, \mathcal{J}) &\leq \limsup_n \frac{p(k_n O')}{p(J_n)} h_*^d(f, \mathcal{J}_{O'}), \\ &\leq p(O') h_*^d(f, \mathcal{J}_{O'}). \end{aligned}$$

\square

For $O \in \mathcal{D}^1$ the origin belongs to $\text{Int}(O)$ so that $\alpha O \in \mathcal{D}$ and $O \subset \text{Int}(\alpha O)$ for any $\alpha > 1$. Moreover we have $h_*^d(f, \mathcal{J}_{\alpha O}) = h_*^d(f, \mathcal{J}_O)$ by Lemma 10. Together with Lemma 11 we get immediately :

Corollary 11.

$$\forall O \in \mathcal{D}^1, \quad h_*^d(f, O) = h_*^d(f, \mathcal{J}_O).$$

Corollary 12.

$$O \mapsto h_*^d(f, O) \text{ is continuous on } \mathcal{D}^1.$$

Convex polytopes are dense in \mathcal{D} . Therefore we get with \mathcal{P} being the collections of convex d -polytopes with the origin in their interior set :

Corollary 13.

$$\sup_{O \in \mathcal{D}^1} h_*^d(f, O) = \sup_{P \in \mathcal{P}} h_*^d(f, \mathcal{J}_P).$$

However we will see that the supremum is not always achieved. We prove now a formula for the rescaled entropy of a power.

Lemma 12.

$$\forall O \in \mathcal{D}^1 \quad \forall k \in \mathbb{N}, \quad h_*^d(f^k, O) = kh_*^d(f, O).$$

Proof. Let $O \in \mathcal{D}^1$ and $\mathcal{J} = (J_n)_n \in \mathcal{E}(O)$. Let $J_n^k = J_n \oplus \underbrace{I \oplus \cdots \oplus I}_{k \text{ times}}$ for all n . The sequence $\mathcal{J}^k = (J_n^k)_n$ belongs also to $\mathcal{E}(O)$. Moreover the partition $\mathbb{P}_{J_n^k}$ is finer than $\bigvee_{l=0}^{k-1} f^{-l} \mathbb{P}_{J_n}$. Therefore

$$h_*(f^k, \mathbb{P}_{J_n}) \leq kh_*(f, \mathbb{P}_{J_n}) = h_* \left(f^k, \bigvee_{l=0}^{k-1} f^{-l} \mathbb{P}_{J_n} \right) \leq h_*(f^k, \mathbb{P}_{J_n^k})$$

and we then obtain

$$h_*^d(f^k, \mathcal{J}) \leq kh_*^d(f, \mathcal{J}) \leq h_*^d(f^k, \mathcal{J}^k).$$

We conclude by taking the supremum in $\mathcal{J} \in \mathcal{E}(O)$. □

5.4. Asymptotic properties of the rescaled entropy. Let (X, f) be a cellular automaton with domain I . We will show that in dimension two, the limsup defining $h_{top}^d(f, \mathcal{J}_O)$ with $O \in \mathcal{D}^1$ is in fact a limit. We first relate the entropy of \mathbb{P}_J with the entropy of $\mathbb{P}_{\partial^\pm J}$ and we prove an upperbound for the rescaled entropy $h_{top}^d(f, O)$ in term of the first relative quermass integral $V_{\mathbb{I}}(O)$ with \mathbb{I} being the convex hull of I' .

Lemma 13. *For any bounded subset J of \mathbb{R}^d , we have*

$$h_*(f, \mathbb{P}_J) = h_*(f, \mathbb{P}_{\partial_I^- J}) \text{ and } h_*(f, \mathbb{P}_J) \leq h_*(f, \mathbb{P}_{\partial_I^+ J}).$$

Proof. The inequality $h_*(f, \mathbb{P}_J) \geq h_*(f, \mathbb{P}_{\partial_I^- J})$ follows directly from the inclusion $\partial^- J \subset J$. By definition of the domain I and the erosion $J \ominus I$, we have $\mathbb{P}_J > f^{-1} \mathbb{P}_{J \ominus I}$. Therefore we get $f^{-1} \mathbb{P}_J \vee \mathbb{P}_J = f^{-1} \mathbb{P}_{\partial^- J} \vee \mathbb{P}_J$ and then by induction $\mathbb{P}_J \vee \bigvee_{l=0}^{k-1} f^{-l} \mathbb{P}_{\partial^- J} = \bigvee_{l=0}^{k-1} f^{-l} \mathbb{P}_J$ for all k . We conclude that :

$$\begin{aligned} h_*(f, \mathbb{P}_J) &= \lim_k \frac{1}{k} H_*(f, \bigvee_{l=0}^{k-1} f^{-l} \mathbb{P}_J), \\ &\leq \lim_k \frac{1}{k} \left(H_*(\mathbb{P}_J) + H_* \left(\bigvee_{l=0}^{k-1} f^{-l} \mathbb{P}_{\partial^- J} \right) \right), \\ &\leq h_*(f, \mathbb{P}_{\partial^- J}). \end{aligned}$$

We also have

$$\mathbb{P}_J \vee \mathbb{P}_{\partial^+ J} > \mathbb{P}_{J \oplus I} > f^{-1} \mathbb{P}_J.$$

Therefore we get now by induction on k

$$\mathbb{P}_J \vee \bigvee_{l=0}^{k-2} f^{-l} \mathbb{P}_{\partial^+ J} > \bigvee_{l=0}^{k-1} f^{-l} \mathbb{P}_J.$$

This implies $h_*(f, \mathbb{P}_{\partial^+ J}) \leq h_*(f, \mathbb{P}_J)$. □

Proposition 14. *For any $O \in \mathcal{D}^1$,*

$$h_{top}^d(f, O) \leq V_{\mathbb{I}}(O) \log \#\mathcal{A}.$$

Proof. Recall that

$$\begin{aligned} h_{top}^d(f, O) &= h_{top}^d(f, \mathcal{J}_O), \\ &= \limsup_n \frac{h_{top}(f, \mathbf{P}_{nO})}{p(nO)}. \end{aligned}$$

Then by applying Lemma 13 we obtain

$$\begin{aligned} h_{top}^d(f, O) &\leq \limsup_n \frac{h_{top}(f, \mathbf{P}_{\partial^\pm nO})}{p(nO)}, \\ &\leq \limsup_n \frac{\#\partial^\pm nO \log \#\mathcal{A}}{p(nO)}. \end{aligned}$$

For all $k \in \mathbb{N} \setminus \{0\}$ we let I_k be the domain of f^k and we denote by \mathbb{I}_k the convex hull of $I'_k = I_k \cup \{0\}$. Clearly we have $I_k \subset \underbrace{I \oplus \cdots \oplus I}_{k \text{ times}}$, therefore $\mathbb{I}_k \subset k\mathbb{I}$. By Lemma 4, we get for

some constant $C = C(d)$:

$$\begin{aligned} h_{top}^d(f^k, O) &\leq (V_{\mathbb{I}_k}(O) + C) \log \#\mathcal{A}, \\ &\leq (V_{k\mathbb{I}}(O) + C) \log \#\mathcal{A}, \\ &\leq (kV_{\mathbb{I}}(O) + C) \log \#\mathcal{A}. \end{aligned}$$

But by Lemma 17 we have $h_{top}^d(f^k, O) = kh_{top}^d(f, O)$, so that we finally conclude when k goes to infinity

$$h_{top}^d(f, O) \leq V_{\mathbb{I}}(O) \log \#\mathcal{A}.$$

□

Lemma 14. *Assume from now $d = 2$. For any integral polygon O there is a sequence $(\epsilon_m)_m$ going to zero such that for all $m \in \mathbb{N}$*

$$\exists N \forall n > N \forall \mu \in \mathcal{M}(f) \exists k \in \mathbb{Z}^d, \frac{h_\mu(f, \mathbf{P}_{nO})}{n} < \frac{h_{\sigma^k \mu}(f, \mathbf{P}_{mO})}{m} + \epsilon_m.$$

Proof. For a fixed face F of O , we let E_F be the band given by $E_F := T_F^+ O(-h_I(N^F)) \cap T_F^- O$, thus $\partial^- nO \subset \bigcup_{F \in \mathcal{F}(O)} E_F$. There is $M > 0$ depending only on O and I such that for $m > M$ we may cover E_F by a collection \mathcal{C}_F of polygons (a priori not pairwise disjoint) such that

- for any $C \in \mathcal{C}_F$, there is $\alpha_C \in \mathbb{Z}^d$ with $C = \alpha_C + mO$,
- for any $C \in \mathcal{C}_F$ we have $\alpha_C + mF \subset T_F$,
- the segments $\alpha_C + mF$ for $C \in \mathcal{C}_F$ are tiling the line T_F .

Then for $1 \ll m \ll n$ the set $\partial^- nO$ may be covered by a subfamily \mathcal{D} of $\bigcup_F \mathcal{C}_F$ with $\#\mathcal{D} \lesssim n/m$. Therefore we get

$$\begin{aligned} h_\mu(\mathbf{P}_n O) &= h_\mu(\mathbf{P}_{\partial^- nO}), \\ &\leq h_\mu\left(\bigvee_{C \in \mathcal{D}} \mathbf{P}_{\alpha_C + mO}\right), \\ &\leq \sum_{C \in \mathcal{D}} h_\mu(\mathbf{P}_{\alpha_C + mO}), \\ &\leq \sum_{C \in \mathcal{D}} h_{\sigma^{\alpha_C} \mu}(\mathbf{P}_m O), \\ h_\mu(\mathbf{P}_n O) &\lesssim \frac{n}{m} \sup_{k \in \mathbb{Z}^d} h_{\sigma^k \mu}(\mathbf{P}_m O). \end{aligned}$$

□

Remark 15. We ignore if the above Lemma also holds in higher dimensions. In the proof we use the fact that any face F tiles the hyperplane H_F . Therefore we may reproduce the proof in higher dimension for polygons satisfying this property, e.g. pallelipeds or hexagonal prisms.

Corollary 16. For any $O \in \mathcal{D}$,

$$h_{top}^d(f, \mathcal{J}_O) = \lim_n \frac{h_{top}(f, \mathbf{P}_n O)}{p(nO)}.$$

Proof. When O is an integral convex polygon, the statement follows directly from Lemma 14 and the variational principle $h_{top}(f, \mathbf{P}) = \sup_{\mu \in \mathcal{M}(f)} h_\mu(f, \mathbf{P})$ for clopen partitions P . Consider now the general case $O \in \mathcal{D}$. Let O' be a convex polygon with rational vertices satisfying $O \subset \text{Int}(O') \subset O' \subset \alpha O$ for $\alpha \ll 1$. For some $m \in \mathbb{N}$, the polygon mO' is integral. Then we get :

$$\begin{aligned} h_*^d(f, \mathcal{J}_O) &\leq p(O') h_{top}^d(f, \mathcal{J}_{O'}), \text{ according to Lemma 11,} \\ &\lesssim h_{top}^d(f, \mathcal{J}_{O'}), \text{ as } p(O') \sim p(O) \text{ by monotonicity of the perimeter,} \\ &\lesssim h_{top}^d(f, \mathcal{J}_{mO'}), \text{ as } p(O') \sim p(O) \text{ by Lemma 10,} \\ &\lesssim \lim_n \frac{h_{top}(f, \mathbf{P}_{nmO'})}{p(nmO')}, \text{ by the formula for integral convex polygons,} \\ &\lesssim \liminf_n \frac{h_{top}(f, \mathbf{P}_{nm\alpha O})}{p(nm\alpha O)}, \\ &\lesssim \liminf_n \frac{h_{top}(f, \mathbf{P}_n O)}{p(nO)} \text{ by applying again Lemma 10.} \end{aligned}$$

□

Similarly we get :

Corollary 17. For any $\mu \in \mathcal{M}(f, \sigma)$ and any $O \in \mathcal{D}$,

$$h_\mu^d(f, \mathcal{J}_O) = \lim_n \frac{h_\mu(f, \mathbf{P}_n O)}{p(nO)}.$$

Remark 18. Clearly we have $h_\mu^d \leq h_{top}^d$ for any $\mu \in \mathcal{M}(f)$ but we ignore if a general variational principle holds true.

6. RUELLE INEQUALITY

Recall (X, σ) denotes a \mathbb{Z}^d -subshift. The topological entropy of σ is defined for any Følner sequence $\mathcal{L} = (L_n)_n$ (see e.g. [19]) as

$$h_{top}(\sigma) = \limsup_n \frac{H_{top}(\mathbf{P}_{L_n})}{\#L_n}.$$

Lemma 15. *Then for all $\epsilon > 0$ there exists $c > 0$ such that we have for any $K \subset J$ convex bodies :*

$$H_{top}(\mathbf{P}_{J \setminus K}) \leq \left(\#J \setminus K + cp(J) \right) \cdot (h_{top}(\sigma) + \epsilon).$$

Proof. Let $\epsilon > 0$. As the sequence of cubes $\mathcal{C} = (C_n)_n$ defined by $C_n = [-n, n]^d \cap \mathbb{Z}^d$ is a Følner sequence, there is a positive integer m such that $\frac{H_{top}(\mathbf{P}_{C_m})}{\#C_m} < h_{top}(\sigma) + \epsilon$. Then for some $c = c(m) > 0$ we may cover $J \setminus K$ by a family \mathcal{F} at most $\frac{\#J \setminus K + cp(J)}{\#C_m}$ disjoint translated copies of C_m . Therefore

$$\begin{aligned} H_{top}(\mathbf{P}_{J \setminus K}) &\leq \left(\#J \setminus K + cp(J) \right) \frac{H_{top}(\mathbf{P}_{C_m})}{\#C_m}, \\ &\leq \left(\#J \setminus K + cp(J) \right) \cdot (h_{top}(\sigma) + \epsilon). \end{aligned}$$

□

We refine now the inequality obtained in Lemma 14 at the level of invariant measures :

Lemma 16.

$$\forall \mu \in \mathcal{M}(f), \quad h_\mu(f, O) \leq h_{top}(\sigma) \int \chi_O d\mu.$$

Proof. For any convex domain J and any $\mu \in \mathcal{M}(f)$ we have

$$\begin{aligned} h_\mu(f, \mathbf{P}_J) &\leq H_\mu(f^{-1}\mathbf{P}_J | \mathbf{P}_J), \\ &\leq \sum_{A \in \mathbf{P}_J} \mu(A) H_{\mu_A}(f^{-1}\mathbf{P}_J). \end{aligned}$$

Fix $\epsilon > 0$ and let c be as in Lemma 15. Then if $(K_A)_{A \in \mathbf{P}_J}$ is a family of convex bodies in $\prod_{A \in \mathbf{P}_J} \mathcal{E}_f^-(A, J)$ we obtain

$$\begin{aligned} h_\mu(f, \mathbf{P}_J) &\leq \sum_{A \in \mathbf{P}_J} \mu(A) H_{\mu_A}(f^{-1}\mathbf{P}_{J \setminus K_A}), \\ &\leq \sum_{A \in \mathbf{P}_J} \mu(A) H_{top}(\mathbf{P}_{J \setminus K_A}), \\ &\leq \sum_{A \in \mathbf{P}_J} \mu(A) \left(\#J \setminus K_A + cp(J) \right) \cdot (h_{top}(\sigma) + \epsilon). \end{aligned}$$

By choosing K_A with $\#J \setminus K_A$ minimal we obtain

$$h_\mu(f, \mathbf{P}_J) \leq (h_{top}(\sigma) + \epsilon) \cdot \left(\int \text{gr}_J^- f d\mu + cp(J) \right).$$

Therefore we have for any convex exhaustion $\mathcal{J} = (J_n)_n \in \mathcal{E}(O)$:

$$\begin{aligned} h_\mu^d(f, \mathcal{J}) &= \limsup_n \frac{h_\mu(f, P_J)}{p(J_n)}, \\ &\leq (h_{top}(\sigma) + \epsilon) \cdot \left(\int \limsup_n \frac{\text{gr}_{J_n}^- f}{p(J_n)} d\mu + c \right), \\ &\leq (h_{top}(\sigma) + \epsilon) \left(\int \text{gr}_{\mathcal{J}}^- f d\mu + c \right). \end{aligned}$$

By taking the supremum over $\mathcal{J} \in \mathcal{E}(O)$ we get

$$h_\mu^d(f, O) \leq (h_{top}(\sigma) + \epsilon) \left(\int \text{gr}_O f d\mu + c \right).$$

By Lemma 12 we have $\frac{h_\mu^d(f^k, O)}{k} = h_\mu^d(f, O)$ for any k . Apply the above inequality to f^k :

$$h_\mu^d(f, O) \leq (h_{top}(\sigma) + \epsilon) \left(\int \frac{\text{gr}_O f^k}{k} d\mu + \frac{c}{k} \right).$$

When k goes to infinity and then ϵ goes to zero, we conclude $h_\mu^d(f, O) \leq h_{top}(\sigma) \int \chi_O d\mu$. \square

7. ENTROPY FORMULA FOR PERMUTATIVE CA

The cellular automaton f is said **permutative** at $i \in \mathbb{Z}^d$ if for all pattern P on $I \setminus \{i\}$ and for all $a \in \mathcal{A}$ there is $b \in \mathcal{A}$ such that the pattern P_b^i on $I \cup \{i\}$ given by the completion of P at i by b satisfies $F(P_b^i) = a$, in particular i belongs to the domain I of f . The CA is said permutative when it is permutative at the nonzero extreme points of the convex hull \mathbb{I} of $I' = I \cup \{0\}$ (these points lie in I). The algebraic CA as described in the introduction are permutative. The sets I' and \mathbb{I} have the same smallest bounding sphere, thus $R_{I'} = R_{\mathbb{I}}$.

Proposition 19. *The topological rescaled entropy of a permutative CA f on X_d is given by*

$$h_{top}^d(f) = R_{I'} \log \#\mathcal{A}.$$

Theorem 1, stated in the introduction, follows from Proposition 19.

Question. For a permutative CA, the uniform measure $\lambda^{\mathbb{Z}^d}$ with λ being the uniform measure on \mathcal{A} is known to be invariant [20]. Does the uniform measure maximize the entropy ?

Recall that for any $k \in \mathbb{N} \setminus \{0\}$ we denote by I_k the domain of f^k and \mathbb{I}_k the convex hull of $I'_k = I_k \cup \{0\}$. In the following we also let $C(P, L)$ be the cylinder associated to the pattern P on $L \subset \mathbb{Z}^d$. We also write $C(P)$ for this cylinder when there is no confusion on J .

Lemma 17. *For any permutative CA f and any $k \in \mathbb{N} \setminus \{0\}$, the CA f^k is also permutative and*

$$\mathbb{I}_k = k\mathbb{I}.$$

Proof. As already observe, the inclusion $\mathbb{I}_k \subset k\mathbb{I}$ holds for any CA (not necessarily permutative). We will show $k \text{ex}(\mathbb{I}) \subset I'_k$, which implies together with $\mathbb{I}_k \subset k\mathbb{I}$ the equality $\mathbb{I}_k \subset k\mathbb{I}$. Let $i \in \text{ex}(\mathbb{I}) \setminus \{0\} \subset I$. For a fixed k we prove by induction on k that f^k is permutative at ki , in particular $ki \in I'_k$. Let P be a pattern on $I_k \setminus \{ki\}$ and let $a \in \mathcal{A}$. Since we have $I_k \subset I_{k-1} \oplus I$, we may complete P by a pattern Q on $(I_{k-1} \oplus I) \setminus \{ki\}$. By induction hypothesis, $(k-1)i$ lies in $\text{ex}(\mathbb{I}_{k-1})$ and i lies in $\text{ex}(\mathbb{I})$, therefore ki does not belong to $I_{k-1} \oplus (I \setminus \{i\})$. Therefore there is a pattern R on $I \setminus \{i\}$ such that $f^{k-1}C(Q, (I_{k-1} \oplus I) \setminus \{ki\})$ is contained in the cylinder $C(R, I \setminus \{i\})$. As f is permutative at i there is $b \in \mathcal{A}$ with $F(R_b^i) = a$ or in other terms

$f(C(R_b^i, I)) \subset C(a, \{0\})$. Since f^{k-1} is permutative at $(k-1)i$, we may find $c \in \mathcal{A}$ with $f^{k-1}(C(Q_c^{ki}, I_{k-1} \oplus I)) \subset C(b, \{i\})$. Therefore we get

$$f^k(C(Q_c^{ki}, I_{k-1} \oplus I)) \subset f(C(R_b^i, I)) \subset C(a, \{0\}).$$

But I_k is the domain of f^k and P is the restriction of Q to $I_k \setminus \{ki\}$, so that we also have $f^k(C(P_c^{ki}, I_k)) \subset C(a, \{0\})$, i.e. f^k is permutative at ki . \square

For a convex polytope J and a face F of J we consider the subset of $\partial_{\mathbb{I}}^- J$ given by $\partial_{\mathbb{I}}^- F := \partial_{\mathbb{I}}^- J \cap T_F^+ J(-h_{\mathbb{I}}(N^F))$. The sets $\partial_{\mathbb{I}}^- F$ for $F \in \mathcal{F}(J)$ are covering $\partial_{\mathbb{I}}^- J$ but do not define a partition in general. For any $F \in \mathcal{F}(J)$ we let $u^F \in \text{ex}(\mathbb{I}) \subset I'$ with $u^F \cdot N^F = h_{\mathbb{I}}(N^F)$ and we also let d_F be the the Euclidean distance to T_F . Then for $j \in \partial_{\mathbb{I}}^- J$ we let F_j be a face of J such that $d_{F_j}(j + u^{F_j}) = -d_{F_j}(j) + u^{F_j} \cdot N^{F_j}$ is maximal among faces F with $j \in \partial_{\mathbb{I}}^- F$. We consider then a total order \prec on $\partial_{\mathbb{I}}^- J$ such that $i \prec j$ if $d_{F_i}(j + u^{F_i}) < d_{F_j}(j + u^{F_j})$. We also let $\mathcal{F}_{\mathbb{I}}(J)$ be the subset of $\mathcal{F}(J)$ given by faces F for which u_F is uniquely defined. We denote by $\partial_{\mathbb{I}}^\perp J$ the subset of $\partial_{\mathbb{I}}^- J$ given by

$$\partial_{\mathbb{I}}^\perp J := \bigcup_{F \in \mathcal{F}_{\mathbb{I}}(J)} \partial_{\mathbb{I}}^- F.$$

Lemma 18. *Let $j \in \partial_{\mathbb{I}}^\perp J$. Then*

$$\forall k \in \mathbb{N}, j + ku^{F_j} \notin \{j' \prec j\} \oplus k\mathbb{I}.$$

Proof. We argue by contradiction : there are $j' \prec j$ and $u \in \mathbb{I}$ with $j + ku^{F_j} = j' + ku$. Observe that

$$\begin{aligned} d_{F_j}(j + ku^{F_j}) &= d_{F_j}(j + u^{F_j}) + (k-1)u^{F_j} \cdot N^{F_j}, \\ d_{F_j}(j' + ku) &= d_{F_j}(j' + u) + (k-1)u \cdot N^{F_j}. \end{aligned}$$

We will show that the equality between these two distances implies $u = u^{F_j}$, therefore $j = j'$. Indeed we have

$$\begin{aligned} d_{F_j}(j' + u) &\leq \sup_{v \in \text{ex}(\mathbb{I})} d_{F_j}(j' + v), & u \cdot N^{F_j} &\leq \sup_{v \in \text{ex}(\mathbb{I})} v \cdot N^{F_j}, \\ &\leq d_{F_{j'}}(j' + u^{F_{j'}}), & &\leq h_{\mathbb{I}}(N^{F_{j'}}), \\ d_{F_j}(j' + u) &\leq d_{F_j}(j + u^{F_j}) & u \cdot N^{F_j} &\leq u^{F_j} \cdot N^{F_j}, \end{aligned}$$

therefore $u \cdot N^{F_j} = u^{F_j} \cdot N^{F_j}$, and finally $u = u^{F_j}$ as j belongs to $\partial_{\mathbb{I}}^\perp J$. \square

For a partition \mathbf{P} of X and a positive integer k , we write \mathbf{P}^k to denote the iterated partition $\bigvee_{l=0}^{k-1} f^{-l}\mathbf{P}$ in order to simplify the notations.

Lemma 19. *Let $O \in \mathcal{P}$ and let k, n be positive integers. For any $A^k \in \mathbf{P}_{nO}^k$ and any pattern P on $\partial_{\mathbb{I}}^\perp nO$, there is $w \in A^k$ such that $f^k w$ belongs to $C(P, \partial_{\mathbb{I}}^\perp nO)$.*

Proof. For any $j \in \partial_{\mathbb{I}}^\perp nO$ we let P_j be the restriction of $P = (p_l)_{l \in \partial_{\mathbb{I}}^\perp nO}$ to $\{j' \prec j\}$. We show now by induction on $j \in \partial_{\mathbb{I}}^\perp nO$ that there is $w \in A^k$ with $f^k w \in C(P_j)$. By Lemma 17 the CA f^k is permutative at ku^{F_j} so that we may change the $(j + ku^{F_j})^{\text{tearith}}$ -coordinate of w to get $w' \in X$ with $(f^k w')_j = p_j$. Moreover the j' -coordinates of $f^k w$ for $j' \prec j$ only depends on the coordinates of w on $\{j' \prec j\} \oplus k\mathbb{I}$ so that by Lemma 18 we still have $f^k w' \in C(P_j, \{j' \prec j\})$, thus $f^k w' \in C(P_{j''})$ with j'' being the successor of j for \prec in $\partial_{\mathbb{I}}^\perp nO$. \square

Lemma 20. *With the notations of Subsection 3.5, we have*

$$\mathcal{F}(T') = \mathcal{F}_{\mathbb{I}}(T')$$

and

$$\forall R > 0, \mathcal{F}(T'_R) \setminus \mathcal{F}_{\parallel}(T'_R) \subset \mathcal{F}_{\mathbb{I}}(T'_R).$$

Proof. Let $F \in \mathcal{F}(T')$ or $F \in \mathcal{F}(T'_R) \setminus \mathcal{F}_{\parallel}(T'_R)$. Such a face F is tangent to $S_{I'}$ at some $u \in \text{ex}(\mathbb{I})$ with $u \cdot N^F = h_{\mathbb{I}}(N^F)$. Then any v with $v \cdot N^F = h_{\mathbb{I}}(N^F)$ belongs to T_F . But $T_F \cap \mathbb{I} \subset T_F \cap S_{I'} = \{u\}$, therefore we have necessarily $u_F = u$. \square

We are now in a position to prove Proposition 19.

Proof of Proposition 19. The inequality $h_{top}^d(f) \leq R_{I'} \log \#\mathcal{A}$ follows immediately from Proposition 14 and Proposition 6. By Lemma 19 we have for any $O \in \mathcal{P}$

$$\forall A^k \in \mathbf{P}_{nO}^k, \#\{A^{k+1} \in \mathbf{P}_{nO}^{k+1}, A^{k+1} \subset A^k\} \geq \#\underline{\partial^{\perp} nO}.$$

Consequently we have

$$\begin{aligned} h_{top}(f, \mathbf{P}_{nO}) &\geq \#\underline{\partial^{\perp} nO} \log \#\mathcal{A}, \\ h_{top}^d(f, \mathcal{J}_O) &\geq \limsup_n \frac{\#\underline{\partial^{\perp} nO}}{n^{d-1}p(O)} \log \#\mathcal{A}. \end{aligned}$$

We first assume that $S_{\mathbb{I}} = S_{I'}$ is nondegenerated. Let T' be the dual polytope of a generating polytope T . By Lemma 20 we have $\mathcal{F}(T') = \mathcal{F}_{\mathbb{I}}(T')$, therefore $\mathcal{F}(nT') = \mathcal{F}_{\mathbb{I}}(nT')$ and $\partial^{\perp} nT' = \partial^{-} nT'$ for all n . Applying then Lemma 4 we get for some constant $C = C(d)$:

$$\begin{aligned} h_{top}^d(f, \mathcal{J}_{T'}) &\geq \limsup_n \frac{\#\underline{\partial^{-} nT'}}{n^{d-1}p(T')} \log \#\mathcal{A}, \\ &\geq \frac{V_{\mathbb{I}}(T')}{p(T')} \log \#\mathcal{A} - C. \end{aligned}$$

Then it follows from Proposition 6 that :

$$h_{top}^d(f, \mathcal{J}_{T'}) \geq R_{\mathbb{I}} \log \#\mathcal{A} - C.$$

For any positive integer k , we have $\mathbb{I}_k = k\mathbb{I}$ according to Lemma 17, so that we get together with the power formula of Lemma 12 :

$$\begin{aligned} h_{top}^d(f, \mathcal{J}_{T'}) &= \frac{h_{top}^d(f^k, \mathcal{J}_{T'})}{k}, \\ &\geq \frac{R_{\mathbb{I}_k}}{k} \log \#\mathcal{A} - \frac{C}{k}, \\ &\geq \frac{R_{k\mathbb{I}}}{k} \log \#\mathcal{A} - \frac{C}{k}, \\ &\geq R_{\mathbb{I}} \log \#\mathcal{A} - \frac{C}{k}, \\ h_{top}^d(f, T') &\geq R_{I'} \log \#\mathcal{A}. \end{aligned}$$

Let us deal now with the degenerated case. Then we have for all $R > 0$ by Lemma 20

$$h_{top}^d(f, \mathcal{J}_{T'_R}) \geq \limsup_n \frac{\#\underline{\partial^{-} nT'_R} - \sum_{F \in \mathcal{F}_{\parallel}(nT'_R)} \#\underline{\partial^{-} F}}{p(nT'_R)} \log \#\mathcal{A}.$$

As the $(d-1)$ -Hausdorff measure of the faces in $\mathcal{F}_{\parallel}(T'_R)$ is small compared to $p(T'_R)$ for large R , we also have

$$\limsup_n \frac{\sum_{F \in \mathcal{F}_{\parallel}(nT'_R)} \#\underline{\partial^{-} F}}{p(nT'_R)} \xrightarrow{R \rightarrow +\infty} 0.$$

We conclude as in the degenerated case. Fix $\epsilon > 0$ and let $k > C\epsilon^{-1}$. We obtain finally

$$\begin{aligned} h_{top}^d(f, \mathcal{J}_{T'_R}) &= \frac{h_{top}^d(f^k, \mathcal{J}_{T'_R})}{k}, \\ &\geq \left(\frac{V_{\mathbb{I}_k}(T'_R)}{kp(T'_R)} - \epsilon - \limsup_n \frac{\sum_{F \in \mathcal{F}_{\parallel}(nT'_R)} \#\partial^- F}{p(nT'_R)} \right) \log \#\mathcal{A}, \\ &\geq \left(\frac{V_{\mathbb{I}}(T'_R)}{p(T'_R)} - \epsilon - \limsup_n \frac{\sum_{F \in \mathcal{F}_{\parallel}(nT'_R)} \#\partial^- F}{p(nT'_R)} \right) \log \#\mathcal{A}, \\ &\xrightarrow{R \rightarrow +\infty} (R_{I'} - \epsilon) \log \#\mathcal{A}. \end{aligned}$$

□

REFERENCES

- [1] F. Blanchard, P. Tisseur, Entropy rate of higher-dimensional cellular automata, 2012. hal-00713029
- [2] Bokowski, J., H. Hadwiger and J.M. Will, Eine Ungleichung zwischen Volumen, Oberflache and Gitterpunktanzahl konvexer Korper im n-dimensionalen euklidischen Raum, Math. Z. 127, 363-364 (1972).
- [3] T. Bonnesen and W. Fenchel, Theory of convex bodies, BCS Associates, Moscow, ID, 1987. Translated from the German and edited by L. Boron, C. Christenson and B. Smith.
- [4] Chakerian, G. D.; Sangwine-Yager, J. R., A generalization of Minkowski's inequality for plane convex sets. Geom. Dedicata 8 (1979), no. 4, 437-444.
- [5] M. Damico, G. Manzini, L. Margara, On computing the entropy of cellular automata, Theoretical Comput. Sci. 290, 1629-1646 (2003).
- [6] Gritzmann, Peter; Wills, Jrg M, Lattice points. Handbook of convex geometry, Vol. A, B, 765-797, North-Holland, Amsterdam, 1993.
- [7] Gruber, Peter M, Convex and discrete geometry. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 336. Springer, Berlin, 2007.
- [8] Hlawka, E., Uber Integrale auf konvexen Korpern. I, II, Monatsh. Math. 54 (1950) 136, 8199
- [9] E. L. Lakshtanov, E. S. Langvagen, Entropy of Multidimensional Cellular Automata Problemy Peredachi Informatsii, 2006, 42:1, 4351
- [10] Lakshtanov, E. L.; Langvagen, E. S., A criterion for the infinity of the topological entropy of multidimensional cellular automata. (Russian) Problemy Peredachi Informatsii 40 (2004), no. 2, 7072; translation in Probl. Inf. Transm. 40 (2004), no. 2, 165-167
- [11] Lindenstrauss, Elon Mean dimension, small entropy factors and an embedding theorem. Inst. Hautes Etudes Sci. Publ. Math. No. 89 (1999), 227-262 (2000)
- [12] Matheron, G., La formule de Steiner pour les rosions. (French) J. Appl. Probability 15 (1978), no. 1, 126-135.
- [13] G. Morris, T. Ward, Entropy bounds for endomorphisms commuting with K actions, Israel J. Math. 106 (1998) 1-12.
- [14] Hedlund, Gustav A., Endomorphisms and Automorphisms of the Shift Dynamical Systems, Mathematical System Theory, 3 (4): 320-375 (1969),
- [15] Shereshevsky M A 1991, Lyapunov exponents for one-dimensional cellular automata, J. Nonlinear Sci. 2 18
- [16] Tisseur, P.(F-CNRS-IML) Cellular automata and Lyapunov exponents. (English summary) Nonlinearity 13 (2000), no. 5, 1547-1560.
- [17] M. Shinoda, M. Tsukamoto, Symbolic dynamics in mean dimension theory, arXiv:1910.00844
- [18] Thomas B. Ward, Additive Cellular Automata and Volume Growth, Entropy 2000, 2, 142-167
- [19] D. Ornstein and B. Weiss Entropy and isomorphism theorems for actions of amenable groups J. d'Anal. Math., 48 (1987), 1141.
- [20] Willson, Stephen J. On the ergodic theory of cellular automata. Math. Systems Theory 9 (1975), no. 2, 132-141.

SORBONNE UNIVERSITE, LPSM, 75005 PARIS, FRANCE
E-mail address: david.burguet@upmc.fr