



HAL
open science

Mec1 Is Activated at the Onset of Normal S Phase by Low-dNTP Pools Impeding DNA Replication

Romain Forey, Ana Poveda, Sushma Sharma, Antoine Barthe, Ismaël Padioleau, Claire Renard, Robin Lambert, Magdalena Skrzypczak, Krzysztof Ginalski, Armelle Lengronne, et al.

► **To cite this version:**

Romain Forey, Ana Poveda, Sushma Sharma, Antoine Barthe, Ismaël Padioleau, et al.. Mec1 Is Activated at the Onset of Normal S Phase by Low-dNTP Pools Impeding DNA Replication. *Molecular Cell*, 2020, 78 (3), pp.396-410.e4. 10.1016/j.molcel.2020.02.021 . hal-02612438

HAL Id: hal-02612438

<https://hal.science/hal-02612438>

Submitted on 27 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Mec1 is activated at the onset of normal S phase
by low dNTP pools impeding DNA replication**

Romain Forey^{1*}, Ana Poveda^{1,2*}, Sushma Sharma³, Antoine Barthe¹, Ismael Padioleau¹,
Claire Renard¹, Robin Lambert¹, Magdalena Skrzypczak⁴, Krzysztof Ginalski⁴, Armelle
Lengronne¹, Andrei Chabes³, Benjamin Pardo^{1#} & Philippe Pasero^{1#}

1. Institut de Génétique Humaine, CNRS, Université de Montpellier, Montpellier, France
2. Instituto de Investigación en Salud Pública y Zoonosis, Facultad de Ciencias Químicas, Universidad Central del Ecuador, Quito, Ecuador
3. Department of Medical Biochemistry and Biophysics and Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå, Sweden
4. Laboratory of Bioinformatics and Systems Biology, Centre of New Technologies, University of Warsaw, Zwirki i Wigury 93, 02-089 Warsaw, Poland

Correspondence to: Benjamin Pardo, benjamin.pardo@igh.cnrs.fr
Philippe Pasero, philippe.pasero@igh.cnrs.fr (lead contact)

*These two authors contributed equally

SUMMARY

The Mec1 and Rad53 kinases play a central role during acute replication stress in budding yeast. They are also essential for viability in normal growth conditions, but the signal that activates the Mec1-Rad53 pathway in the absence of exogenous insults is currently unknown. Here, we show that this pathway is active at the onset of normal S phase because dNTP levels present in G₁ phase **may not be sufficient** to support processive DNA synthesis and impede DNA replication. This activation can be suppressed experimentally by increasing dNTP levels in G₁ phase. Moreover, we show that unchallenged cells entering S phase in the absence of Rad53 undergo irreversible fork collapse and mitotic catastrophe. Together, these data indicate that cells use suboptimal dNTP pools to detect the onset of DNA replication and activate the Mec1-Rad53 pathway, which in turn maintains functional forks and triggers dNTP synthesis, allowing the completion of DNA replication.

KEYWORDS:

DNA replication, S-phase checkpoint, dNTP synthesis, cell cycle, budding yeast

INTRODUCTION

The integrity of the genome is at risk during the S phase of the cell cycle when the cell must duplicate faithfully all of its genetic information before mitosis. This is achieved through the sequential activation of thousands of replication origins, distributed along the chromosomes following a defined replication timing program (Fragkos et al., 2015). A new copy of the DNA is synthesized by replication forks, which progress bidirectionally from the replication origins until they converge with other forks (Burgers and Kunkel, 2017; Dewar and Walter, 2017). During this process, the forks may encounter obstacles that cause them to stall (Gaillard et al., 2015; Techer et al., 2017; Zeman and Cimprich, 2014). This replication fork stalling, commonly referred to as replication stress (RS), activates checkpoint signaling pathways. The best characterized of these signaling pathways is the DNA damage checkpoint, which arrests cell cycle progression in response to DNA breaks until lesions are repaired (Weinert and Hartwell, 1988). Stressed replication forks, by contrast, are detected by the DNA replication checkpoint, which acts in multiple ways to maintain genome integrity (Saldivar et al., 2017). Failure to build an efficient RS response leads to genomic instability, which can fuel tumorigenesis (Kotsantis et al., 2018; Macheret and Halazonetis, 2015).

The mechanism by which cells detect and signal stalled forks has been characterized extensively in budding yeast (Pardo et al., 2017) and it is highly conserved among eukaryotes (Saldivar et al., 2017). Briefly, obstacles to replication on the DNA template result in accumulation of single-stranded DNA (ssDNA) at replication forks. This ssDNA is coated with replication protein A (RPA) and recruits the Ser/Thr kinase Mec1 (known as ATR in vertebrates) through its partner Ddc2 (ATRIP in vertebrates) (Rouse and Jackson, 2002; Zou and Elledge, 2003). Mec1 phosphorylates multiple targets near blocked forks, including histone H2A (H2AX in vertebrates) (Redon et al., 2003; Rogakou et al., 1999). Mec1 also activates the effector kinase Rad53 (the functional homologue of CHK1 in vertebrates) to transduce the checkpoint response (Sanchez et al., 1999) and prevent fork collapse (Desany et al., 1998; Tercero et al., 2003). In addition, the Mec1-Rad53 pathway slows down DNA replication (Bacal et al., 2018; Santocanale and Diffley, 1998; Shirahige et al., 1998), prevents premature entry into mitosis (Clarke et al., 2001) and promotes dNTP synthesis by multiple mechanisms that increase the activity of ribonucleotide reductase (RNR), including the Dun1-dependent degradation of the RNR inhibitor Sml1 (Zhao et al., 1998).

In budding yeast, quantitative mass spectrometry analyses have shown that Mec1 is functional during normal S phase but does not activate Rad53 (Bastos de Oliveira et al., 2015; Lanz et

al., 2018). Rad53-independent roles of Mec1 include activation of replication origins (Randell et al., 2010), *de novo* telomere addition (Zhang and Durocher, 2010) and the response to proteotoxic stress (Corcoles-Saez et al., 2018). Likewise, checkpoint-independent functions of Rad53 have been identified during S phase, including degradation of excess histones (Gunjan and Verreault, 2003). Together, these data suggest that the canonical Mec1-Rad53 checkpoint pathway is not functional during normal DNA replication. Yet, Mec1 and Rad53 are both essential for cell viability (Weinert, 1997) and the lethal phenotype of *mec1* Δ and *rad53* Δ mutants can be suppressed by deleting *SML1* (Zhao et al., 1998). These observations led to the proposal that the Mec1-Rad53 pathway triggers the degradation of Sml1 at the onset of S phase to promote dNTP synthesis (Zhao et al., 2001; Zhao and Rothstein, 2002), but the signal that activates it has remained unknown.

Here, we show that the Mec1-Rad53 pathway is activated by a spontaneous replication stress resulting from limiting dNTP pools upon initiation of DNA replication. This transient activation of the Mec1 pathway triggers dNTP synthesis through the Rad53-dependent degradation of Sml1 and the upregulation of RNR activity. Importantly, this pathway also prevents fork collapse and premature entry into mitosis during this transient dNTP shortage. These data indicate that cells do not anticipate the G₁/S transition by producing dNTPs in late G₁ phase but rather use nucleotide shortage and fork arrest as a physiological signal to activate dNTP synthesis.

RESULTS

The Mec1-Rad53 pathway is active during normal S phase

Replication origins fire sequentially throughout S phase and slight differences in Rad53 activity have major effects on the extent of ‘early’ and ‘late’ replication origin firing (Poli et al., 2012). To determine whether the Mec1-Rad53 pathway is active during normal S phase, we monitored repression of late replication origins as a readout of checkpoint activation (Crabbé et al., 2010). Wild-type, *mec1-1 sml1-1* and *rad53-11 Saccharomyces cerevisiae* cells were arrested in G₁ phase by using α -factor and then released synchronously into S phase in the presence of BrdU to label ongoing DNA synthesis (Fig. 1A). As a positive control, we also treated the cells with 200 mM hydroxyurea (HU) to block elongation. The BrdU-labelled DNA was then immunoprecipitated and hybridized to tiling arrays to analyze origin firing, as described previously (Crabbé et al., 2010). In HU-treated wild-type cells, BrdU incorporation was detected at early origins (open arrowheads), but not at late origins (filled arrowheads; Fig. 1B), as illustrated for chromosome 14, reflecting checkpoint activation of the Mec1-Rad53 pathway. By contrast, BrdU incorporation was detected at both early and late origins in *mec1-1 sml1-1* and *rad53-11* mutants, indicating their inability to repress late replication origins. Of note, the BrdU peaks were wider in *mec1-1 sml1-1* mutants than in wild-type and *rad53-11* cells because the *sml1-1* mutation prevents inhibition of RNR by Sml1 and thus the dNTP pools are larger in these cells in G₁ phase, so the replication forks are able to progress further in the presence of HU (Poli et al., 2012). In untreated wild-type cells, BrdU incorporation was detected at early origins 20 minutes after their release from G₁ phase arrest and at late origins 10 minutes later (Fig. S1B), as previously described (Müller et al., 2014). In *mec1-1 sml1-1* and *rad53-11* mutants, BrdU incorporation at late origins was increased compared to wild-type cells (Fig. 1B). When expressed relative to wild-type levels, the ratio of BrdU incorporation was close to 1 for all early origins, but increased at late origins proportionally to the time of origin activation (Fig. 1C, D). These data indicate that although early origins fire with similar kinetics in the three strains, late origins are activated prematurely in the *mec1-1 sml1-1* and *rad53-11* mutants. This confirms and extends earlier studies that analyzed only two late origins (Santocanale and Diffley, 1998; Shirahige et al., 1998). We obtained similar results by monitoring BrdU incorporation (Fig. S1B–D) and DNA copy number variation (Fig. S1E–G) by deep sequencing (Fang et al., 2017; Müller et al., 2014). Interestingly, we saw no premature activation of late origins in the absence of the DDC mediator Rad9 (Fig. S1B–D), indicating that Rad53 is not activated by DNA damage during

normal DNA replication. Together, these data indicate that the Mec1-Rad53 pathway is active during normal S phase and delays the activation of late origins, independently of the DDC.

Mec1 is transiently activated around early replication origins

To identify regions of the yeast genome that could activate Mec1 during normal S phase, we used chromatin immunoprecipitation coupled to NGS (ChIP-seq) to analyze the genome-wide distribution of the Mec1-binding protein Ddc2 and the histone H2A phosphorylated on serine 129 by Mec1 (γ -H2A). The detection of Ddc2 allows the localization of Mec1, whereas γ -H2A indicates Mec1 activity. To position these signals relative to forks, the distribution of Cdc45, a component of the replication helicase, was also mapped by ChIP-seq and variations in DNA copy number were followed by NGS (Müller et al., 2014). Wild-type cells expressing epitope-tagged Cdc45 or Ddc2 were released from a G₁ arrest, as above, and samples were collected in G₁ phase and at the indicated times after release, during the entire S phase (**Fig. 2A, B**). In G₁ phase, Ddc2 was enriched at tRNA genes and at highly expressed genes enriched in the RNAPII subunit Rpb3 (**Fig. S2A–C**), which corresponds to documented ChIP artefacts (Teytelman et al., 2013). By contrast, γ -H2A was absent from these highly expressed genes (high Rpb3 levels), presumably because of the fast histone turnover at these loci; γ -H2A was detected only at a subset of repressed genes, as reported earlier (Szilard et al., 2010). To focus on S phase-specific profiles, ChIP-seq signals from G₁-phase samples were subtracted from those from S-phase samples. In a typical experiment, Cdc45 peaks were detected at early origins 20 minutes after release from G₁ phase arrest (**Fig. 2C**). At 30 minutes, new peaks were detected at late origins and the Cdc45 peaks at early origins split in two due to bidirectional fork progression, as seen by analyzing the signals over 40 kb regions centered on replication origins (**Fig. 2D**). Interestingly, narrow Ddc2 and γ -H2A peaks (5–10 kb) centered on early origins appeared at 30 minutes and had disappeared 10 minutes later (**Fig. 2C, D**), indicating transient activation and inactivation of Mec1 around 30 minutes after release from G₁ phase arrest. Similar kinetics of Mec1 activation and inactivation can also be seen on heat maps of Ddc2 and γ -H2A ChIP signals at intervals centered on origins and sorted according to their replication time (**Fig. S2D**). Interestingly, Ddc2 and γ -H2A peaks did not overlap with Cdc45 peaks, which is consistent with the accumulation of RPA-coated ssDNA gaps behind forks under replication stress conditions (Bacal et al., 2018; García-Rodríguez et al., 2018). To confirm that Ddc2 enrichment reflects the recruitment of Mec1 to RPA-coated ssDNA upon early origins firing, we also immunoprecipitated RPA and an HA-tagged version of Mec1 (Mallory and Petes, 2000) and monitored their enrichment at four early origins by

ChIP-qPCR in G₁-arrested cells and after release for 30 minutes into S phase (**Fig. S2E**). This analysis confirmed that RPA and Mec1 accumulate near early origins during normal replication to ~15-20% of the levels detected in HU-arrested cells (**Fig. S2F**). Importantly, we also confirmed that H2A phosphorylation at these loci was essentially mediated by Mec1 and not by Tel1 as γ -H2A levels at early origins was not affected by the deletion of *TEL1* (**Fig. S2G and S2H**). Finally, the transient nature of Mec1 activation was confirmed by plotting variations in Ddc2 and γ -H2A levels at all early-, mid- and late-replicating regions of the genome (**Fig. 2E**). Together, these data show that Mec1-Ddc2 is recruited and transiently activated at early replicating regions during a normal S phase.

Mec1 activation does not correlate with highly transcribed regions

To investigate the origin of the stress causing Mec1 activation during normal DNA replication, we determined the percentages of various genome annotations that overlapped with the Ddc2 and γ -H2A peaks (**Fig. 3A**). This analysis showed that both Ddc2 and γ -H2A were enriched at early replication origins and centromeres. Ddc2 was also detected at tRNA genes and γ -H2A was enriched at ORFs. Since highly expressed genes interfere with fork progression (Azvolinsky et al., 2009; Szilard et al., 2010), we hypothesized that Mec1 might be activated by conflicts between replication and transcription. To assess this possibility, we monitored by ChIP-seq of G₁ and S phase cells the distribution of Rpb3 as an indicator of total RNAPII and Rpb1-pS2 as an indicator of active RNAPII and plotted their distributions relative to Ddc2 and γ -H2A. This analysis revealed that the levels of Ddc2 and γ -H2A at origins correlated neither with increased levels of active RNAPII (Rpb1-pS2, $R < 0.1$; **Fig. 3B**) nor with total RNAPII (Rpb3, $R < 0.2$; **Fig. 3B and S3A**). Moreover, the presence of tRNA genes in the vicinity of early origins did not correlate with the enrichment in Ddc2 and γ -H2A (**Fig. 3C**). This result was further confirmed by plotting Ddc2 and γ -H2A enrichment at origins relative to RNAPIII (Rpo31) levels (Cutler et al., 2018) (**Fig. S3B**). Together, these data indicate that the transient Mec1 activation observed at early replicating regions does not correlate with high transcription and is probably not caused by replication-transcription conflicts. These results are fully consistent with a recent study reporting no detectable replication pausing at highly transcribed genes by RNAPII in wild-type budding yeast cells (Osmundson et al., 2017).

Mec1 activation follows entry into S phase with limiting dNTP pools

We compared the distribution of γ -H2A during normal S phase to the pattern observed in cells exposed to HU, a potent RNR inhibitor (Nyholm et al., 1993). Strikingly, we observed that the distribution of γ -H2A (Fig. 4A) and its intensity around early origins were highly similar and correlated in the two cell populations ($R=0.9$; Fig. 4B, C), suggesting that Mec1 may be activated by a transient shortage of dNTP upon entry into S phase. To investigate this possibility, wild-type cells were released synchronously into S phase from a G₁ arrest and were collected every three minutes to measure variations in dNTP levels (Fig. 4D, E). Interestingly, levels of the dNTPs did not increase before the onset of S phase (arrow). Levels of dATP and dGTP rather increased after the beginning of DNA synthesis and Sml1 degradation (arrow), 15–18 minutes after release from the α -factor block (Fig. 4E–G). In particular, dATP levels were relatively low in G₁ phase compared to the other dNTPs (Fig. 4E) and were fully exhausted in HU-treated cells (Fig. S4). Together, these data suggest that cells do not prepare for DNA replication by building dNTP pools before activating early origins, but rather enter S phase with suboptimal dNTP levels. DNA polymerases are probably unable to efficiently utilize dNTPs at these low concentrations and would stall and activate the Mec1-Rad53 pathway.

Mec1 activation in normal S phase is suppressed by increasing dNTP levels

To test whether activation of Mec1 in early S phase is due to limiting amounts of dNTPs, we artificially increased the levels of dNTPs by overexpressing RNR components just before the G₁–S transition. To this end, we constructed a strain in which *SML1* was deleted (*sml1* Δ), to remove the inhibitor of RNR, and additional copies of the four *RNR* genes were expressed under the control of a galactose-inducible promoter. When induced by the addition of 2% galactose, the excess RNR activity in G₁ phase cells prevented entry into S phase (Fig. S5A), as described previously (Chabes and Stillman, 2007); however, by optimizing the conditions of induction, we were able to find conditions in which the cells progressed through S phase with normal kinetics (Fig. 5A, B). The *sml1* Δ mutation alone, before induction of the *RNR* genes, increased the amounts of individual dNTPs by 2–12-fold in G₁ phase cells (Fig. 5C). Induction of the *RNR* genes further increased dNTP pools by 4–35-fold in G₁ phase and in S phase cells 30 minutes after release from G₁ arrest (Fig. 5C). Remarkably, these increased dNTP levels reduced to background levels the accumulation of γ -H2A at four early origins (Fig. 5D) and globally at the genome-wide level (Fig. 5E). It also induced a premature activation of late origins, which is consistent with the fact that cells do not activate Mec1 and

Rad53 (**Fig. S5B**). Finally, the accumulation of γ -H2A in early S phase and its suppression by increased dNTP levels were also observed in the absence of Tel1 (**Fig. S5C and S5D**). Together, these findings are consistent with a model in which low dNTP levels at the onset of normal S phase activate Mec1.

Suboptimal dNTP pools in early S induces a global replication blockage in *rad53* Δ cells

The data presented above suggest that cells enter S phase with low dNTP pools, activate the Mec1-Rad53 pathway upon fork pausing and increase RNR via the Rad53-dependent degradation of the RNR inhibitor Sml1. To determine the fate of *rad53* Δ cells entering S phase in the presence of Sml1, we set out to overexpress *SML1* with a galactose-inducible promoter in *rad53* Δ cells released synchronously from G₁ phase (Zhao et al., 2001). Since *rad53* Δ cells grow more slowly than wild-type cells due to the toxicity of free histones (Gunjan and Verreault, 2003; Zhao et al., 2001), one copy of histone H3 and H4 genes (*hht2* Δ *hhf2* Δ) was deleted to suppress this slow growth phenotype without affecting nucleosome assembly (Gunjan and Verreault, 2003; Maya Miles et al., 2018). Control, *pGAL-SML1* and *rad53* Δ *pGAL-SML1* cells were arrested in G₁ phase with α -factor and *SML1* was overexpressed with the addition of galactose before releasing cells synchronously into S phase (**Fig. 6A**). As expected, *rad53* Δ *pGAL-SML1* cells remained blocked in early S phase because the lack of Rad53 makes them unable to degrade Sml1 (**Fig. 6B, C**). By contrast, Sml1 was rapidly degraded in *pGAL-SML1* cells, which entered G₂ phase 90 minutes after release from G₁ and accumulated again Sml1 in the following cell cycle (**Fig. 6B, C**). Induction of *SML1* in G₁-arrested cells induced a proportional reduction of the four dNTP levels. Interestingly, the relative amounts of dCTP, dTTP and dGTP were also similar upon release into S phase, but the relative amount of dATP was lower in *rad53* Δ *pGAL-SML1* cells (**Fig. 6D**, arrows). These data suggest that dATP levels could become limiting for DNA replication when cells are unable to degrade Sml1 in early S phase.

To characterize the consequences of suboptimal dNTP pools on DNA synthesis in the absence of Rad53, we next measured variations in DNA content in *pGAL-SML1* and *rad53* Δ *pGAL-SML1* cells by NGS as described (Müller et al., 2014). In control cells, DNA content increased at 213 early origins 45 minutes after release from the G₁ arrest, reflecting the initiation of DNA synthesis (**Fig. 6E, S6A**) and forks progressed by 6.3 kb on average (**Fig. 6F**). Local variations in DNA content were no longer detected at 90 minutes (**Fig. S6A**), indicating that replication is complete. In *rad53* Δ *pGAL-SML1* cells, initiation was detected at

179 early origins but elongation progressed more slowly than in control cells (2.2 vs 6.3 kb; **Fig. 6E**). At 90 minutes, forks did not progress further than 4.5 kb from early and late origins (**Fig. 6F**), as observed in HU-arrested *rad53-11* mutants (Poli et al., 2012). Persistent fork arrest in these cells was accompanied by a strong accumulation of γ -H2A 45 and 90 minutes after release from G₁-phase arrest (**Fig. 6E, S6B**), reminiscent of that observed in HU-arrested forks (**Fig. 4A**). By contrast, *pGAL-SML1* cells completed genome duplication in 90 minutes and γ -H2A was only transiently detected in early-replicated regions (**Fig. S6B**), as seen in wild-type cells (**Fig. 2C**). Together, these data indicate that the inability of *rad53Δ* cells to increase dNTP levels at the onset of S phase causes an irreversible arrest of replication forks.

Loss of viability in *rad53Δ* cells is caused by fork collapse and premature entry in mitosis

The degradation of the RNR inhibitor Sml1 depends on Dun1, a protein kinase activated by Rad53 (Andreson et al., 2010; Zhao and Rothstein, 2002). Unlike *MEC1* and *RAD53*, however, *DUN1* is not an essential gene (Zhao and Rothstein, 2002). These observations suggest that Mec1 and Rad53 have other functions during a normal S phase besides regulating dNTP levels. To investigate these other possible functions, we assessed the impact of *SML1* expression on the progress of *rad53Δ*, *mec1Δ* and *dun1Δ* mutants through S phase. To this end, *pGAL-SML1*, *rad53Δ pGAL-SML1* *mec1Δ pGAL-SML1* and *dun1Δ pGAL-SML1* cells were arrested in G₁ phase by using α -factor and were released into S phase after induction of *SML1* expression with galactose (**Fig. 7A**). Samples were collected at the indicated times, plated on glucose-containing medium to turn off *SML1* expression, and the number of colonies was counted as a measure of cell survival (**Fig. 7B**). Overexpression of Sml1 in the *dun1Δ* mutant blocked S phase progression, as in *rad53Δ* and *mec1Δ* cells (**Fig. 7C**) but did not affect cell viability (**Fig. 7B**). We assume that cell viability in *dun1Δ* mutants was mediated by Mec1 and Rad53, the latter being phosphorylated at 120 minutes to the same extent as in HU-arrested cells (**Fig. 7D**). When *SML1* expression was turned off, *dun1Δ pGAL-SML1* cells were able to complete S phase, despite the persistence of detectable Sml1 levels. **This is consistent with the fact that overexpression of a poorly degradable Sml1 derivative, sml-4SA, slows entry into S phase (Andreson et al. 2010).** By contrast, *rad53Δ pGAL-SML1* and *mec1Δ pGAL-SML1* cells remained blocked with a nearly 1C DNA content (**Fig. 7E**), presumably because they were unable to maintain the integrity of forks under low dNTP conditions. These data indicate that unlike *mec1Δ* and *rad53Δ* cells, *dun1Δ* cells can

tolerate a transient depletion of dNTP pools in early S phase and can also complete S phase at a slow rate despite the presence of Sml1, as long as Mec1 and Rad53 are present to protect stalled forks.

To determine whether cells progress through the cell cycle in the absence of Sml1 degradation, we next monitored the levels of the mitotic cyclin Clb2 in *rad53Δ pGAL-SML1*, *mec1Δ pGAL-SML1* and *dun1Δ pGAL-SML1* cells released from G₁ arrest. Despite the fact that they failed to complete DNA replication, *rad53Δ*, *mec1Δ* and *dun1Δ* mutants accumulated Clb2 with similar kinetics as control cells expressing *pGAL-SML1* (**Fig. 7D**). We therefore asked whether these cells would progress through mitosis with partially replicated chromosomes. In budding yeast, there is no checkpoint to prevent cells with unreplicated DNA from entering mitosis (Torres-Rosell et al., 2007). Nevertheless, Rad53 and the mitotic spindle assembly checkpoint (SAC) prevent chromosome segregation in anaphase until the completion of DNA replication (Magiera et al., 2014; Palou et al., 2017). We used fluorescence microscopy to follow the segregation of nuclei during the time-course of this experiment by expressing mCherry-Pus1, a protein that marks the whole nuclear compartment (Siniosoglou et al., 1998). Two hours after the release of the cells from G₁ phase arrest, we saw normal segregation of nuclei only in the control *pGAL-SML1* cells (**Fig. 7F, G**). At the same time point, the *rad53Δ pGAL-SML1* cells had deformed nuclei, including nuclei of unequal sizes in mother and daughter cells and anaphase bridges (**Fig. 7F, G** and **Fig. S7**). These observations indicate that the SAC does not completely impede anaphase completion in *rad53Δ pGAL-SML1* cells. In *dun1Δ pGAL-SML1* cells, by contrast, the chromosomes had not segregated into mother and daughter cells at 120 minutes, consistent with Rad53 preventing the completion of anaphase (**Fig. 7D, F, G**). After inhibiting *SML1* expression in *rad53Δ pGAL-SML1* and *dun1Δ pGAL-SML1* cells by the addition of glucose, only the *dun1Δ pGAL-SML1* cells fully replicated their DNA and correctly segregated their chromosomes (**Fig. 7E–G**); most *rad53Δ pGAL-SML1* cells failed to complete S phase and had severe segregation defects (**Fig. 7E–G**).

We then asked whether the failure of *rad53Δ pGAL-SML1* cells to complete S phase and chromosome segregation might be the cause of the loss of viability observed above (**Fig. 7B**). To this end, we assayed the formation of nuclear foci of Rad52–GFP, as a marker of DNA damage and ongoing DNA repair by recombination. We observed a >10-fold increase in the percentage of cells with Rad52 foci in *rad53Δ pGAL-SML1* cells relative to the control *pGAL-SML1* and *dun1Δ pGAL-SML1* cells after release from G₁ phase arrest (**Fig. 7H**). These foci

were detected as early as 45 minutes after release from G₁, before cells attempted to separate their chromosomes (**Fig. 7C**). This suggests that the formation of Rad52 foci first occurs at blocked forks. Both the fraction of *rad53Δ pGAL-SML1* cells with Rad52 foci and the number of foci per cell further increased after addition of glucose and progression through mitosis, suggesting further chromosome fragmentation during the course of this mitosis (**Fig. 7E, F, H**). Altogether, these data indicate that the essential roles of Rad53 during normal S phase are to maintain the integrity of replication forks and prevent the segregation of partially replicated chromosomes after the initiation of DNA replication with limiting dNTP pools, which occurs spontaneously at the onset of S phase.

DISCUSSION

We used budding yeast as a model to investigate how the Mec1^{ATR} pathway is activated during normal S phase. We show that Mec1 and Rad53 delay the activation of late origins of replication even in the absence of exogenous RS. This function does not depend on the DDC mediator Rad9, indicating that the Mec1-Rad53 pathway detects paused forks in unperturbed cells independently of the presence of DNA damage. To identify regions of the yeast genome that activate the Mec1-Rad53 pathway during normal DNA replication, we mapped the distribution of Ddc2 and γ -H2A on DNA in cells synchronously passing through S phase in the absence of drugs. Time-resolved ChIP-seq experiments revealed that Mec1 is transiently activated at the onset of S phase in short regions (5–10 kb) centered on early replicating origins. These regions contain a high density of tRNA and highly expressed RNAPII genes that have been previously described to cause replication stress because of conflicts between replication and transcription (Azvolinsky et al., 2009; Ivessa et al., 2003; Stirling et al., 2012; Szilard et al., 2010). We found no significant correlation, however, between Mec1 activation and transcription levels at these loci, indicating that the burst of Mec1 activation occurring at the onset of S phase is caused by another type of stress.

The distribution of γ -H2A in untreated cells in early S phase was very similar to that observed in HU-treated cells in which RNR is inhibited and therefore dNTP levels are low. We hypothesized, therefore, that Mec1 might be activated by replication occurring under a suboptimal concentration of dNTPs. Consistent with this hypothesis, we found that cells entered S phase with low dNTP pools, which were not sufficient to sustain the activity of the hundreds of forks generated from early origins. Moreover, the accumulation of γ -H2A at early origins was totally suppressed by artificially increasing dNTP levels before releasing cells into S phase. Finally, impeding dNTP *de novo* synthesis in *rad53 Δ* cells blocked the progression of replication forks shortly after their activation, even though dNTP pools were not exhausted. Together, these data indicate that the Mec1-Rad53 pathway is activated by critically low dNTP levels that do not permit optimal DNA replication, similar to the effect of HU (**Fig. 8**). Thus, bulk dNTP synthesis occurs *after* the activation of early origins.

Our finding that Mec1 is activated at the onset of S phase by low dNTP levels raises the question of how eukaryotic cells coordinate the production of dNTPs with DNA synthesis. Since high dNTP levels are detrimental to the fidelity of DNA replication (Ganai and Johansson, 2016; Pai and Kearsley, 2017), nucleotide pools are maintained at levels that are limiting for the progress of the replication fork (Davidson et al., 2012; Poli et al., 2012;

Stodola and Burgers, 2016). In budding yeast, dNTPs present in late G₁ phase are sufficient only to replicate 10–15% of the genome and are exhausted within minutes upon activation of early origins (Poli et al., 2012). Cells cannot anticipate entry into S phase by producing large dNTP pools in G₁ phase as they interfere with the initiation of replication (Chabes and Stillman, 2007). The production of dNTPs must therefore occur concomitantly with the activation of early origins, but the mechanism responsible for this coordination had remained unclear. Our data shed **new** light on this mechanism by explaining how the Mec1-Rad53 pathway could regulate RNR activity during normal S phase.

RNR is the rate-limiting enzyme for dNTP production (Giannattasio and Branzei, 2017; Sanvisens et al., 2013). Its activity is regulated at the G₁/S transition by various transcriptional and posttranscriptional mechanisms (Bastos de Oliveira et al., 2012; Elledge and Davis, 1990; Niida et al., 2010; Wu et al., 2018). In addition, RNR activity is upregulated by the Mec1-Rad53 pathway in response to DNA damage via the degradation of the RNR repressors Crt1, Dif1 and Sml1 (Huang et al., 1998; Lee et al., 2008; Zhao and Rothstein, 2002). Since deletion of *SML1* or overexpression of *RNR* genes rescues the lethality of *mec1* and *rad53* null alleles (Desany et al., 1998; Zhao et al., 1998), it has been proposed that the Mec1-Rad53 pathway activates RNR during normal S phase by promoting the Dun1-dependent degradation of Sml1 (Zhao et al., 2001; Zhao and Rothstein, 2002). However, the signal that triggers the Mec1–Rad53–Dun1 cascade in the absence of exogenous stress has remained unknown. Our finding that low dNTP pools impede DNA replication at the onset of the S phase provides this missing link and defines a simple and robust mechanism ensuring that dNTPs are produced at the very moment when they are needed.

Our data indicate that the Mec1–Rad53–Dun1 pathway plays a critical role at the onset of S phase by coordinating RNR activity with the firing of early replication origins. Yet, this model does not explain why the *MEC1* and *RAD53* genes are essential for viability, whereas *DUN1* is not. Mec1 and Rad53 should therefore have another important role during normal growth that is not shared by Dun1. We have found that the Mec1-Rad53 pathway delays the activation of late origins during a normal S phase, but this function is not essential for viability (Zegerman and Diffley, 2010).

As a matter of fact, the most important function of the Mec1-Rad53 pathway under acute RS conditions is to maintain the integrity of stalled forks to ensure that they will be able to resume DNA synthesis once the stress is relieved (Tercero et al., 2003). Remarkably, we have found that Rad53, but not Dun1, is also essential to prevent fork collapse in untreated cells.

These data indicate that Rad53 is not only required to stimulate dNTP synthesis at the onset of S phase but also to maintain the ability of cells to recover from dNTP shortage (**Fig. 8**). Unlike in HU-arrested cells, the activation of Rad53 during normal S phase is very transient, explaining why it has remained undetected (Bastos de Oliveira et al., 2015; Hoch et al., 2013). Importantly, we have observed that *rad53Δ* cells bearing under-replicated and damaged chromosomes progress through mitosis. Normally, the spindle assembly checkpoint (SAC) prevents the segregation of chromosomes with unreplicated centromeres, which impedes the correct kinetochores attachment. Since centromeres are linked to early origins in budding yeast (Fang et al., 2017), these observations suggest that cells lacking Rad53 are still able to replicate their centromeres before reaching anaphase, allowing them to bypass the SAC. The incapacity of these cells to properly segregate their chromosomes is causing even more genomic instability. These data illustrate the many critical functions of the Mec1-Rad53 pathway to ensure proper DNA replication and subsequent mitosis in normal growth conditions.

An important question that remains to be addressed is to what extent this mechanism is conserved in vertebrates. Unlike in yeast, dNTP pools do not increase after DNA damage in mammalian cells (Hakansson et al., 2006; Técher et al., 2016; Techer et al., 2017). Yet, ATR promotes the expression of the RNR subunit *RRM2* in late G₁ phase (Buisson et al., 2015) and prevents its degradation in response to DNA damage (D'Angiolella et al., 2012). Moreover, mice with mutations in *ATR* and with an extra copy of the *RRM2* gene show reduced RS at common fragile sites, which is consistent with the conclusion that ATR prevents spontaneous genomic instability by controlling dNTP pools (Lopez-Contreras et al., 2015). It is therefore likely that similar mechanisms operate in all eukaryotes to coordinate dNTP supply, fork stability and DNA synthesis.

ACKNOWLEDGEMENTS

We thank Maria Moriel Carretero, Etienne Schwob and Carol Featherstone for discussions and for critical comments on the manuscript. We thank Alain Verreault for the gift of anti- γ -H2A antibody, Corrado Santocanale for anti-Rad53 antibody, Monica Segurado and Symeon Siniossoglou for plasmids and Tom Petes for strains. We thank the imaging facility MRI, member of the national infrastructure France-BioImaging supported by the French National Research Agency (ANR-10-INBS-04, «Investments for the future»). R.F. thanks the French Ministry of Higher Education, Research and Innovation and Fondation ARC pour la Recherche sur le Cancer for fellowships. A.P., I.P. and B.P. (SPF20121226243) thank Fondation Recherche Médicale for fellowships. Work in the PP lab is supported by grants from the Agence Nationale pour la Recherche (ANR), Institut National du Cancer (INCa), the Ligue Nationale Contre le Cancer (équipe labéllisée) and the Fondation MSDAvenir. Work in the AC lab is supported by the Swedish Cancer Society and the Swedish Research Council. Funding to KG and MS was provided by Foundation for Polish Science (TEAM) and Polish National Science Centre (2015/17/D/NZ2/03711).

AUTHOR CONTRIBUTIONS

B.P., P.P., A.L., R.F., A.P. and A.C. conceived and planned the study. R.F., A.P., S.S., A.B., R.L. and B.P. conducted the experiments. R.F., I.P. and C.R. performed the bioinformatic analyses. M.S. and K.G. performed NGS experiments. B.P. and P.P. wrote the manuscript and all the authors reviewed it.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. The Mec1-Rad53 pathway represses late origins during a normal S phase.

(A) Wild-type (wt), *rad53-11* and *mec1-1 sml1-1* cells grown at 25°C in YPD medium were released synchronously from G₁ phase arrest for 20, 30 or 40 minutes in the presence of BrdU to label nascent DNA. Cells were also released for 60 minutes in medium containing 200 mM HU to block DNA synthesis.

(B) BrdU-labelled DNA was immunoprecipitated and hybridized to Affymetrix tiling arrays. Enrichment of replicated DNA fragments relative to input DNA (signal log ratio) is shown for a representative region on chromosome 14. Open arrowheads, early origins; filled arrowheads, late origins. *, transposon. Bracketed numbers indicate the range of exponents on the y axis.

(C) Scatter plot of BrdU incorporation at all replication origins (+/- 2.5 kb) in *rad53-11* and *mec1-1 sml1-1* mutants relative to wild-type cells, plotted against the time of origin activation (replication time) (Yabuki et al., 2002).

(D) Levels of BrdU incorporation in *rad53-11* and *mec1-1 sml1-1* mutants relative to wild-type cells expressed for groups of origins sorted according to their time of replication. The mean replication time for each bin is indicated on the x axis. ***, P<0.0001; ns: not significant, Mann–Whitney rank sum test.

Figure 2. Mec1 is recruited to early replicating regions and phosphorylates H2A

(A) Wild-type cells were synchronized in G₁ phase with α -factor and then released from G₁ phase arrest by the addition of Pronase to degrade the α -factor. Samples were recovered at the indicated times.

(B) Analysis of DNA content by flow cytometry.

(C) Distribution of Cdc45 (black), Ddc2 (orange) and γ -H2A (blue) was analyzed by ChIP–seq. A representative region on chromosome 15 is shown. Protein signals are expressed as a ratio of IP to input after subtraction of the corresponding signals from G₁ phase cells. Open arrowheads, early origins; filled arrowheads, late origins. Bracketed numbers indicate the range of exponents on the y axis.

(D) Average profiles of Cdc45, Ddc2 and γ -H2A across 40 kb regions centered on annotated replication origins (n=386).

(E) Variations of total Ddc2 and γ -H2A levels at early-, mid- and late-replicating regions during S phase.

Figure 3. Ddc2 and γ -H2A enrichment at early replicating regions does not correlate with high transcription levels.

(A) Percentage of genomic annotations overlapping with Ddc2 (orange) and γ -H2A (blue) signals in early S phase.

(B) Intensity of Ddc2 and γ -H2A signals plotted against the levels of Rpb1-pS2 and Rpb3 across 5 kb intervals centered on 192 early (blue) and 192 late (grey) origins.

(C) Scatter plots of Ddc2 and γ -H2A signals across 5 kb intervals centered on replication origins either containing (grey dots) or not containing (black dots) tRNA genes (tDNA) plotted against replication time (Trep). Statistical analysis of bins of early origins grouped according to their time of replication indicates that the presence of a tDNA (grey bars) does not increase Ddc2 or γ -H2A binding. Ns, not significant, Mann–Whitney rank sum test.

Figure 4. Entry into S phase is followed by a spontaneous reduction of dNTP pools.

(A) ChIP–seq analysis of the distribution of γ -H2A in wild-type cells 30 minutes after release from G₁ phase arrest (S₃₀) or after 60 minutes in medium containing 200 mM HU. A representative region on chromosome 15 is shown. Open arrowheads, early origins; filled arrowheads, late origins. Bracketed numbers indicate the range of exponents on the y axis.

(B) Scatter plot of γ -H2A levels in cells released from G₁ phase arrest in the presence or the absence of HU. Blue, early origins (n=192); grey, late origins (n=192). R, Spearman correlation coefficient.

(C) Average profiles and heat maps of γ -H2A levels across 60 kb regions centered on 192 early (blue) and 192 late (red) origins.

(D) Wild-type cells were arrested in G₁ phase with α -factor and released synchronously into S phase. Samples were collected every 5 minutes after release and DNA content was analyzed by flow cytometry.

(E) Analysis of intracellular concentrations of dNTPs at the indicated times after release from G₁ phase arrest.

(F) Western blot analysis of Sml1 protein levels after releasing cells from G₁ phase arrest. Tubulin is shown as a control for loading.

(G) Variation of DNA content (grey) and Sml1 protein levels (black) after release from G₁ phase arrest. Error bars correspond to the standard deviation of three independent experiments. DNA content is derived from flow cytometry data. Sml1 levels were determined by western blotting and were normalized to tubulin.

Figure 5. Experimental increase of dNTP levels suppresses Mec1 activation.

(A) Wild-type and pGAL-*RNR sml1Δ* cells were synchronized in G₁ phase before induction of *RNR* gene expression with 1% galactose and 1% sucrose. Cells were then released from G₁ phase arrest and collected at the indicated times.

(B) Analysis of DNA content by flow cytometry.

(C) Analysis of intracellular concentrations of dNTPs in G₁ phase before and after *RNR* induction and 30 minutes after release (S₃₀). Mean values and SD for two independent experiments are shown.

(D) ChIP-qPCR analysis of γ -H2A enrichment near four representative early origins in wild-type and pGAL-*RNR1 sml1Δ* cells in G₁ phase and after release from G₁ phase arrest. Means and SD for four independent experiments are shown.

(E) ChIP-seq analysis of γ -H2A levels at 192 early origins in wild-type and pGAL-*RNR sml1Δ* cells collected 30 minutes after release from G₁ phase arrest. Origins have been divided in two bins according to their replication time (Trep). ***: p<0.001, Mann-Whitney rank sum test.

Figure 6. Rad53 absence in normal S phase induces a strong block of replication forks associated with permanent activation of the S phase checkpoint.

(A) Control pGAL-*SML1* and *rad53Δ* pGAL-*SML1* cells in an *hht2Δ hhf2Δ* background were synchronized in G₁ phase and *SML1* was overexpressed by addition of 2% galactose. Cells were released from G₁ phase arrest and samples were taken at the indicated times.

(B) Analysis of DNA content by flow cytometry.

(C) Western blot analysis of Sml1 protein levels. Tubulin was used as a control for loading.

(D) Analysis of intracellular concentrations of dNTPs in G₁ phase before and after *SML1* induction and at the indicated times after release from G₁ phase arrest. Mean values and SD for two independent experiments are shown. Arrows point to reduced dATP levels in *rad53Δ* p*GAL-SML1* cells.

(E) Variations in DNA copy number (ratio of S/G₁ reads) and γ -H2A (ChIP-seq signal) in control p*GAL-SML1* and *rad53Δ* p*GAL-SML1* cells. A representative region on chromosome 15 is shown. Open arrowheads, early origins; filled arrowheads, late origins. Bracketed numbers indicate the range of exponents on the y axis.

(F) Distance covered by individual replication forks in control p*GAL-SML1* and *rad53Δ* p*GAL-SML1* cells released synchronously into S phase for the indicated times. The length of replicated tracks is derived from DNA copy number variations displayed in panel (E). Median distances and numbers of active origins are indicated. n/a indicates that distances could not be determined for control p*GAL-SML1* cells after 90 minutes due to completion of DNA replication.

Figure 7. *SML1* expression in *rad53Δ* cells induces fork collapse and mitotic catastrophe.

(A) p*GAL-SML1*, *rad53Δ* p*GAL-SML1*, *mec1Δ* p*GAL-SML1* and *dun1Δ* p*GAL-SML1* cells were synchronized in G₁ phase and *SML1* was overexpressed by addition of galactose before release from G₁ phase arrest. Glucose was added after two hours to repress *SML1* expression.

(B) Analysis of cell survival after induction of *SML1* for the indicated times.

(C) Analysis of DNA content by flow cytometry.

(D) Western blot analysis of Rad53, Clb2 and Sml1 levels in cells treated as indicated in (a) or after exposure to 200 mM HU for 120 minutes in S phase (HU). Tubulin and Ponceau staining were used as controls for loading.

(E) Analysis of DNA content by flow cytometry of *rad53Δ* p*GAL-SML1*, *mec1Δ* p*GAL-SML1* and *dun1Δ* p*GAL-SML1* cells that were incubated for 120 minutes in galactose-containing medium to induce *SML1* overexpression, then washed and placed in glucose-containing medium to recover from *SML1* overexpression.

(F) Fluorescence microscopy of nucleus segregation at various times after release from G₁ phase arrest of *pGAL-SML1*, *rad53Δ pGAL-SML1* and *dun1Δ pGAL-SML1* cells expressing mCherry–Pus1 (red) as a marker of the nucleus and Rad52–GFP (yellow) as a marker of foci of DNA repair. Representative images are shown for the three strains and for *rad53Δ pGAL-SML1* and *dun1Δ pGAL-SML1* cells released for 90 minutes in glucose-containing medium. Scale bar, 5 μm.

(G) Quantification of normal and abnormal segregation from images as in (f).

(H) Percentage of cells containing Rad52–GFP foci.

Figure 8. Models of dNTP pool regulation during S phase. See text for details.

(A) Classical model: Cells anticipate the requirement of high dNTP pools in S phase by overexpressing RNR subunits in late G₁ and by degrading the RNR inhibitor Sml1 in early S phase. The degradation of Sml1 depends on the Mec1-Rad53 pathway, but the mechanism that activates this pathway during normal DNA replication is unknown.

(B) **Revised** model: Cells enter S phase with suboptimal dNTP pools, which are not sufficient to sustain the activation of hundreds of early origins upon entry into S phase. This leads to the pausing of replication forks, the transient activation of the Mec1-Rad53 pathway and the degradation of Sml1, allowing dNTPs to reach optimal levels.

STAR METHODS

LEAD CONTACT AND MATERIALS AVAILABILITY

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Philippe Pasero (philippe.pasero@igh.cnrs.fr). All unique strains and reagents generated in this study are available from the Lead Contact without restriction. Requests for strains and reagents donated by other laboratories should be directed to the specific laboratory from which they were received.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All *S. cerevisiae* strains used are haploid and are derived from W303 (see Table S1).

METHOD DETAILS

Cell growth and synchronization

Yeast strains used in this study are listed in the Yeast Strains table. Yeast strains were freshly thawed from frozen stocks and grown at 25°C using standard yeast genetics practices. Gene deletion and tagging were performed as described previously (Longtine et al., 1998). Cells were grown in YPD medium at 25°C unless otherwise stated. They were synchronized in G₁ phase with 8 µg/ml α-factor for 180 minutes and were released from G₁ phase arrest by the addition of 75 µg/ml Pronase and 20 mM citrate phosphate buffer (pH 5.6) in the presence or the absence of 200 mM HU. For overexpression of *SML1* and RNR subunits, cells were grown overnight in medium containing 2% raffinose. Three hours after addition of α-factor, the medium was supplemented for 1 hour with 2% galactose (*SML1*) or 1% galactose and 1% sucrose (RNR).

Analysis of DNA content by flow cytometry

Four hundred and fifty µl of culture samples at 10⁷ cells/ml were diluted in 1 ml of 100% ethanol. Cells were centrifuged for 1 minute at 16.000 RCF and resuspended in 50 mM sodium citrate buffer containing 10 µl of RNase A (20 mg/ml, Qiagen 76254) for 2 hours at 50°C. Then, 10 µl of proteinase K (Sigma, P6556) were added for 2 hours at 50°C. Aggregates of cells were dissociated by sonication. Thirty µl of cell suspension were incubated with 170 µl of 50 mM sodium citrate buffer containing 0.5 µM Sytox Green (Invitrogen). Data were acquired on a MACSQuant Analyzer (Miltenyi Biotec) and analyzed

with FlowJo software. Two to four independent biological replicates were performed per sample.

Chromatin immunoprecipitation

Chromatin immunoprecipitation was performed as described (Lengronne et al., 2004) with minor modifications. One billion (1×10^9) cells were crosslinked for 30 minutes with 1% formaldehyde (Sigma F8775) at room temperature on a shaking device. Fixation was quenched by addition of 0.25 M glycine (Sigma G8898) for 5 minutes under agitation. Cells were washed three times with cold TBS1X (4°C). Dry pellets were frozen and stored at -20°C. Cell pellets were resuspended in lysis buffer (50 mM HEPES-KOH pH7.5, 140 mM NaCl, 1 mM EDTA, 1% Triton-X100, 0.1% sodium deoxycholate) supplemented with 1 mM PMSF, phosphatase inhibitor and anti-protease (cOmplete Tablet, Roche, 505649001) and lysed by bead beating (MB400 U, Yasui Kikai, Osaka). The lysate (WCE, Whole Cell Extract) volume was brought to 3 ml with cold lysis buffer and sonicated with a Q500 sonicator (Qsonica; 12 cycles: 15 sec ON, 45 sec OFF, amplitude 50). Forty μ l of input material was saved for qPCR or sequencing analysis. 180 μ l of Dynabeads Protein A (DPA) were washed three times and resuspended in 1 ml of PBS, 0.5% BSA, 0.1% Tween and incubated with specific volumes of antibodies (40 μ l of anti-PK - Anti-V5 tag, AbD Serotec, MCA1360G; 4 μ l of anti-H2A-S128 - AV137; 40 μ l of anti-HA - Santa Cruz, SC-7392), on a rotating wheel for two hours at 4°C. Antibody-coupled Dynabeads were washed three times with 1 ml of PBS, 0.5% BSA, 0.1% Tween, added to 2.7 ml of WCE and incubated on a rotating wheel at 4°C overnight. Beads were then collected on a magnetic rack. They were washed on ice with cold solutions: twice with Lysis buffer (50 mM HEPES-KOH pH7.5, 140 mM NaCl, 1 mM EDTA, 1% Triton-X100, 0.1% sodium deoxycholate), twice with Lysis buffer + 0.36 M NaCl (50 mM HEPES-KOH pH7.5, 360 mM NaCl, 1 mM EDTA, 1% Triton-X100, 0.1% sodium deoxycholate), twice with Wash buffer (10 mM Tris-HCl pH8, 0.25 M LiCl, 0.5% IGEPAL, 1 mM EDTA, 0.1% sodium deoxycholate) and once with TE (10 mM Tris-HCl pH8, 1 mM EDTA). Antibodies were uncoupled from beads with 120 μ l of Elution Buffer (50 mM Tris-HCl pH8, 10 mM EDTA, 1% SDS) for 10 min at 65°C. Eluates were incubated with 120 μ l of TE containing 0.1% SDS at 65°C for 6 hours to de-crosslink then 130 μ l of TE containing 60 μ g RNase A (Sigma, R65-13) were added and the samples were incubated for 2 hours at 37°C. Proteins were digested by addition of 20 μ l of proteinase K (Sigma, P6556) at 20 mg/ml and incubated for 2 hours at 37°C. Fifty μ l of 5M LiCl were added to the DNA before purification by two rounds of extraction with phenol : chloroform :

isoamyl alcohol (25:24:1; Sigma, P2069) and precipitation by addition of 100 mM sodium acetate (Sigma, S2889), 26 µg/ml of glycogen (Roche, 10901393001) and two volumes of 100% ethanol overnight at -20°C. Samples were centrifuged for 45 min at 16,000 RCF at 4°C, washed with cold 70% ethanol and centrifuged again for 15 min at 16,000 RCF at 4°C. DNA pellets were dried and resuspended in 300 µl of H₂O prior to qPCR reactions or in 25 µl prior to deep-sequencing. The qPCR reactions were performed in a LightCycler480 (Roche). IP/Input ratios were calculated and qPCR results were normalized on negative zones: for yH2A ChIP-qPCR Dop1.

BrdU-IP-chip, BrdU-IP-seq, ChIP-seq and NGS-based analysis of DNA copy number

Immunoprecipitation and processing of BrdU-labelled DNA in BrdU-IP-chip and BrdU-IP-seq experiments were performed as described previously (Poli et al., 2012; Yoshida et al., 2014). For NGS-based assays, sequencing libraries were prepared by using the ThruPLEX DNA-seq Kit (Rubicon Genomics). NGS was performed on a HiSeq4000 sequencing system (Illumina). Single-end reads of 50 bp were aligned to the *S. cerevisiae* genome (2011) with Bowtie2, allowing only perfect matches. Log ratio on input were generated with Bamcompare and displayed using IGB v8.2 (Nicol et al., 2009). Average profiles and heat maps were prepared with Deeptools2 on 60 kb regions centered on replication origins. ChIP-seq scores expressed as RPKM (reads per kilobase per million mapped reads) were calculated by using Bedtools on 2 kb windows centered on replication origins.

Protein extracts and western blotting

TCA precipitation was performed as described previously (Longhese et al., 1997). Extracts were resolved by SDS-PAGE (Biorad) and then transferred to nitrocellulose membranes by using an Invitrogen system. The Rad53 shift due to phosphorylation was detected with a rabbit polyclonal antibody (gift of C. Santocanale, Galway) and tubulin was detected with the YOL1/34 antibody (Abcam; #ab6161). Sml1 was detected with a rabbit polyclonal antibody (Agrisera AB, Sweden, #AS10 847). Clb2 was detected with a rabbit polyclonal antibody (Santa Cruz Biotechnology, y-180). Rnr1 was detected with a rabbit polyclonal antibody (Agrisera, AS16 3639).

Survival test

One hundred cells were spread onto YPD plates and incubated at 30°C for 2 days. Survival rates were determined after 2 days by counting colonies.

Microscopy

Five hundred μl of cells were centrifuged for 1 minute at 16,000 RCF then resuspended in 10 μl of YPD medium. Four μl of the cell suspension were spread between slide and coverslip and used for microscopy analysis. Pictures were taken using a Zeiss AxioImager Z2 fluorescence microscope with a 63x objective. One hundred cells per sample were counted with the FIDJ plugin cell counter to determine the percentage of cells with Rad52 foci and chromosome segregation defects.

Analysis of intracellular dNTP pools

Thirty-five ml of cells at 1×10^7 cells/ml were harvested on nitrocellulose membranes and suspended immediately in an ice-cold mixture of 12% TCA and 15 mM MgCl_2 . The cells were vortex-mixed for 15 min at 4°C and then centrifuged at 14,000 rpm for 1 min at 4°C . The supernatant was neutralized with a freon–trioctylamine mix and analyzed as described previously (Jia et al., 2015).

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis

All statistical tests and numbers of biological replicates are listed in the figure legends. To compare statistical significance between mean values of biological replicates, two-tailed unpaired t tests were used. To compare statistical significance between two distributions, the Mann-Whitney rank-sum test was used. All statistical tests were performed with GraphPad Prism 7.

DATA AND CODE AVAILABILITY

The sequencing data generated in this study have been deposited with the Gene Expression Omnibus <https://www.ncbi.nlm.nih.gov/geo/> with the accession number GSE136605. Unprocessed gel images presented in this manuscript can be found at DOI:10.17632/4n27v8xkpg.2

REFERENCES

- Andreson, B.L., Gupta, A., Georgieva, B.P., and Rothstein, R. (2010). The ribonucleotide reductase inhibitor, Sml1, is sequentially phosphorylated, ubiquitylated and degraded in response to DNA damage. *Nucleic Acids Res* 38, 6490-6501.
- Azvolinsky, A., Giresi, P.G., Lieb, J.D., and Zakian, V.A. (2009). Highly transcribed RNA polymerase II genes are impediments to replication fork progression in *Saccharomyces cerevisiae*. *Mol Cell* 34, 722-734.
- Bacal, J., Moriel Carretero, M., Pardo, B., Barthe, A., Sharma, S., Chabes, A., Lengronne, A., and Pasero, P. (2018). Mrc1 and Rad9 cooperate to regulate initiation and elongation of DNA replication in response to DNA damage. *EMBO J* . 37, e99319
- Bastos de Oliveira, F.M., Harris, M.R., Brazauskas, P., de Bruin, R.A., and Smolka, M.B. (2012). Linking DNA replication checkpoint to MBF cell-cycle transcription reveals a distinct class of G1/S genes. *EMBO J* 31, 1798-1810.
- Bastos de Oliveira, Francisco M., Kim, D., Cussiol, José R., Das, J., Jeong, Min C., Doerfler, L., Schmidt, Kristina H., Yu, H., and Smolka, Marcus B. (2015). Phosphoproteomics Reveals Distinct Modes of Mec1/ATR Signaling during DNA Replication. *Molecular Cell* 57, 1124-1132.
- Buisson, R., Boisvert, J.L., Benes, C.H., and Zou, L. (2015). Distinct but Concerted Roles of ATR, DNA-PK, and Chk1 in Countering Replication Stress during S Phase. *Mol Cell* 59, 1011-1024.
- Burgers, P.M.J., and Kunkel, T.A. (2017). Eukaryotic DNA Replication Fork. *Annu Rev Biochem* 86, 417-438.
- Chabes, A., and Stillman, B. (2007). Constitutively high dNTP concentration inhibits cell cycle progression and the DNA damage checkpoint in yeast *Saccharomyces cerevisiae*. *PNAS* 104, 1183-1188.
- Clarke, D.J., Segal, M., Jensen, S., and Reed, S.I. (2001). Mec1p regulates Pds1p levels in S phase: complex coordination of DNA replication and mitosis. *Nat Cell Biol* 3, 619-627.
- Corcoles-Saez, I., Dong, K., Johnson, A.L., Waskiewicz, E., Costanzo, M., Boone, C., and Cha, R.S. (2018). Essential Function of Mec1, the Budding Yeast ATM/ATR Checkpoint-Response Kinase, in Protein Homeostasis. *Dev Cell* 46, 495-503 e492.
- Crabbé, L., Thomas, A., Pantesco, V., De Vos, J., Pasero, P., and Lengronne, A. (2010). Analysis of replication profiles reveals key role of RFC-Ctf18 in yeast replication stress response. *Nature Structural & Molecular Biology* 17, 1391-1397.
- Cutler, S., Lee, L.J., and Tsukiyama, T. (2018). Chromatin Remodeling Factors Isw2 and Ino80 Regulate Chromatin, Replication, and Copy Number of the *Saccharomyces cerevisiae* Ribosomal DNA Locus. *Genetics* 210, 1543-1556.
- D'Angiolella, V., Donato, V., Forrester, F.M., Jeong, Y.T., Pellacani, C., Kudo, Y., Saraf, A., Florens, L., Washburn, M.P., and Pagano, M. (2012). Cyclin F-mediated degradation of ribonucleotide reductase M2 controls genome integrity and DNA repair. *Cell* 149, 1023-1034.
- Davidson, M.B., Katou, Y., Keszthelyi, A., Sing, T.L., Xia, T., Ou, J., Vaisica, J.A., Thevakumaran, N., Marjavaara, L., Myers, C.L., *et al.* (2012). Endogenous DNA replication stress results in expansion of dNTP pools and a mutator phenotype. *EMBO J* 31, 895-907.

- Desany, B.A., Alcasabas, A.A., Bachant, J.B., and Elledge, S.J. (1998). Recovery from DNA replicational stress is the essential function of the S-phase checkpoint pathway. *Genes Dev* 12, 2956-2970.
- Dewar, J.M., and Walter, J.C. (2017). Mechanisms of DNA replication termination. *Nat Rev Mol Cell Biol* 18, 507-516.
- Elledge, S.J., and Davis, R.W. (1990). Two genes differentially regulated in the cell cycle and by DNA-damaging agents encode alternative regulatory subunits of ribonucleotide reductase. *Genes Dev* 4, 740-751.
- Fang, D., Lengronne, A., Shi, D., Forey, R., Skrzypczak, M., Ginalski, K., Yan, C., Wang, X., Cao, Q., Pasero, P., *et al.* (2017). Dbf4 recruitment by forkhead transcription factors defines an upstream rate-limiting step in determining origin firing timing. *Genes & Development* 31, 2405-2415.
- Fragkos, M., Ganier, O., Coulombe, P., and Mechali, M. (2015). DNA replication origin activation in space and time. *Nat Rev Mol Cell Biol* 16, 360-374.
- Gaillard, H., García-Muse, T., and Aguilera, A. (2015). Replication stress and cancer. *Nature Reviews Cancer* 15, 276.
- Ganai, R.A., and Johansson, E. (2016). DNA Replication-A Matter of Fidelity. *Mol Cell* 62, 745-755.
- García-Rodríguez, N., Morawska, M., Wong, R.P., Daigaku, Y., and Ulrich, H.D. (2018). Spatial separation between replisome- and template-induced replication stress signaling. *EMBO J* 37, e98369.
- Giannattasio, M., and Branzei, D. (2017). S-phase checkpoint regulations that preserve replication and chromosome integrity upon dNTP depletion. *Cell Mol Life Sci* 74, 2361-2380.
- Gunjan, A., and Verreault, A. (2003). A Rad53 kinase-dependent surveillance mechanism that regulates histone protein levels in *S. cerevisiae*. *Cell* 115, 537-549.
- Hakansson, P., Hofer, A., and Thelander, L. (2006). Regulation of mammalian ribonucleotide reduction and dNTP pools after DNA damage and in resting cells. *J Biol Chem* 281, 7834-7841.
- Hoch, N.C., Chen, E.S.-W., Buckland, R., Wang, S.-C., Fazio, A., Hammet, A., Pelliccioli, A., Chabes, A., Tsai, M.-D., and Heierhorst, J. (2013). Molecular basis of the essential S phase function of the Rad53 checkpoint kinase. *Molecular and cellular biology*, 3202-3213.
- Huang, M., Zhou, Z., and Elledge, S.J. (1998). The DNA replication and damage checkpoint pathways induce transcription by inhibition of the Crt1 repressor. *Cell* 94, 595-605.
- Ivessa, A.S., Lenzmeier, B.A., Bessler, J.B., Goudsouzian, L.K., Schnakenberg, S.L., and Zakian, V.A. (2003). The *Saccharomyces cerevisiae* helicase Rrm3p facilitates replication past nonhistone protein-DNA complexes. *Mol Cell* 12, 1525-1536.
- Jia, S., Marjavaara, L., Buckland, R., Sharma, S., and Chabes, A. (2015). Determination of Deoxyribonucleoside Triphosphate Concentrations in Yeast Cells by Strong Anion-Exchange High-Performance Liquid Chromatography Coupled with Ultraviolet Detection. In *DNA Replication: Methods and Protocols*, S. Vengrova, and J. Dalgaard, eds. (New York, NY: Springer New York), pp. 113-121.

- Kotsantis, P., Petermann, E., and Boulton, S.J. (2018). Mechanisms of Oncogene-Induced Replication Stress: Jigsaw Falling into Place. *Cancer Discovery* 8, 537-555.
- Lanz, M.C., Oberly, S., Sanford, E.J., Sharma, S., Chabes, A., and Smolka, M.B. (2018). Separable roles for Mec1/ATR in genome maintenance, DNA replication, and checkpoint signaling. *Genes & Development* 32, 822-835.
- Lee, Y.D., Wang, J., Stubbe, J., and Elledge, S.J. (2008). Dif1 is a DNA-damage-regulated facilitator of nuclear import for ribonucleotide reductase. *Mol Cell* 32, 70-80.
- Lengronne, A., Katou, Y., Mori, S., Yokobayashi, S., Kelly, G.P., Itoh, T., Watanabe, Y., Shirahige, K., and Uhlmann, F. (2004). Cohesin relocation from sites of chromosomal loading to places of convergent transcription. *Nature* 430, 573-578.
- Longhese, M.P., Paciotti, V., Frascini, R., Zaccarini, R., Plevani, P., and Lucchini, G. (1997). The novel DNA damage checkpoint protein ddc1p is phosphorylated periodically during the cell cycle and in response to DNA damage in budding yeast. *EMBO J* 16, 5216-5226.
- Longtine, M.S., McKenzie, A., 3rd, Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen, P., and Pringle, J.R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* 14, 953-961.
- Lopez-Contreras, A.J., Specks, J., Barlow, J.H., Ambrogio, C., Desler, C., Vikingsson, S., Rodrigo-Perez, S., Green, H., Rasmussen, L.J., Murga, M., *et al.* (2015). Increased Rrm2 gene dosage reduces fragile site breakage and prolongs survival of ATR mutant mice. *Genes Dev* 29, 690-695.
- Macheret, M., and Halazonetis, T.D. (2015). DNA Replication Stress as a Hallmark of Cancer. *Annual Review of Pathology: Mechanisms of Disease* 10, 425-448.
- Magiera, M.M., Gueydon, E., and Schwob, E. (2014). DNA replication and spindle checkpoints cooperate during S phase to delay mitosis and preserve genome integrity. *J Cell Biol* 204, 165-175.
- Mallory, J.C., and Petes, T.D. (2000). Protein kinase activity of Tel1p and Mec1p, two *Saccharomyces cerevisiae* proteins related to the human ATM protein kinase. *Proc Natl Acad Sci U S A* 97, 13749-13754.
- Maya Miles, D., Penate, X., Sanmartin Olmo, T., Jourquin, F., Munoz Centeno, M.C., Mendoza, M., Simon, M.N., Chavez, S., and Geli, V. (2018). High levels of histones promote whole-genome-duplications and trigger a Swe1(WEE1)-dependent phosphorylation of Cdc28(CDK1). *Elife* 7.
- Müller, C.A., Hawkins, M., Retkute, R., Malla, S., Wilson, R., Blythe, M.J., Nakato, R., Komata, M., Shirahige, K., de Moura, A.P.S., *et al.* (2014). The dynamics of genome replication using deep sequencing. *Nucleic Acids Research* 42, e3.
- Nicol, J.W., Helt, G.A., Blanchard, S.G., Jr., Raja, A., and Loraine, A.E. (2009). The Integrated Genome Browser: free software for distribution and exploration of genome-scale datasets. *Bioinformatics* 25, 2730-2731.
- Niida, H., Katsuno, Y., Sengoku, M., Shimada, M., Yukawa, M., Ikura, M., Ikura, T., Kohno, K., Shima, H., Suzuki, H., *et al.* (2010). Essential role of Tip60-dependent recruitment of ribonucleotide reductase at DNA damage sites in DNA repair during G1 phase. *Genes Dev* 24, 333-338.

- Nyholm, S., Thelander, L., and Graslund, A. (1993). Reduction and loss of the iron center in the reaction of the small subunit of mouse ribonucleotide reductase with hydroxyurea. *Biochemistry* *32*, 11569-11574.
- Osmundson, J.S., Kumar, J., Yeung, R., and Smith, D.J. (2017). Pif1-family helicases cooperatively suppress widespread replication-fork arrest at tRNA genes. *Nat Struct Mol Biol* *24*, 162-170.
- Pai, C.C., and Kearsley, S.E. (2017). A Critical Balance: dNTPs and the Maintenance of Genome Stability. *Genes (Basel)* *8*, e57
- Palou, R., Palou, G., and Quintana, D.G. (2017). A role for the spindle assembly checkpoint in the DNA damage response. *Curr Genet* *63*, 275-280.
- Pardo, B., Crabbé, L., and Pasero, P. (2017). Signaling pathways of replication stress in yeast. *FEMS Yeast Research* *17*, fow101.
- Poli, J., Tsaponina, O., Crabbe, L., Keszthelyi, A., Pantesco, V., Chabes, A., Lengronne, A., and Pasero, P. (2012). dNTP pools determine fork progression and origin usage under replication stress. *EMBO J* *31*, 883-894.
- Randell, J.C.W., Fan, A., Chan, C., Francis, L.I., Heller, R.C., Galani, K., and Bell, S.P. (2010). Mec1 Is One of Multiple Kinases that Prime the Mcm2-7 Helicase for Phosphorylation by Cdc7. *Molecular Cell* *40*, 353-363.
- Redon, C., Pilch, D.R., Rogakou, E.P., Orr, A.H., Lowndes, N.F., and Bonner, W.M. (2003). Yeast histone 2A serine 129 is essential for the efficient repair of checkpoint-blind DNA damage. *EMBO Rep* *4*, 678-684.
- Rogakou, E.P., Boon, C., Redon, C., and Bonner, W.M. (1999). Megabase chromatin domains involved in DNA double-strand breaks in vivo. *J Cell Biol* *146*, 905-916.
- Rouse, J., and Jackson, S.P. (2002). Lcd1p recruits Mec1p to DNA lesions in vitro and in vivo. *Mol Cell* *9*, 857-869.
- Saldivar, J.C., Cortez, D., and Cimprich, K.A. (2017). The essential kinase ATR: ensuring faithful duplication of a challenging genome. *Nature Reviews Molecular Cell Biology* *18*, 622.
- Sanchez, Y., Bachant, J., Wang, H., Hu, F., Liu, D., Tetzlaff, M., and Elledge, S.J. (1999). Control of the DNA damage checkpoint by chk1 and rad53 protein kinases through distinct mechanisms. *Science* *286*, 1166-1171.
- Santocanale, C., and Diffley, J.F. (1998). A Mec1- and Rad53-dependent checkpoint controls late-firing origins of DNA replication. *Nature* *395*, 615-618.
- Sanvisens, N., de Llanos, R., and Puig, S. (2013). Function and regulation of yeast ribonucleotide reductase: Cell cycle, genotoxic stress, and iron bioavailability. *Biomed J* *36*, 51-58.
- Shirahige, K., Hori, Y., Shiraishi, K., Yamashita, M., Takahashi, K., Obuse, C., Tsurimoto, T., and Yoshikawa, H. (1998). Regulation of DNA-replication origins during cell-cycle progression. *Nature* *395*, 618-621.
- Siniossoglou, S., Santos-Rosa, H., Rappsilber, J., Mann, M., and Hurt, E. (1998). A novel complex of membrane proteins required for formation of a spherical nucleus. *EMBO J* *17*, 6449-6464.

- Stirling, P.C., Chan, Y.A., Minaker, S.W., Aristizabal, M.J., Barrett, I., Sipahimalani, P., Kobor, M.S., and Hieter, P. (2012). R-loop-mediated genome instability in mRNA cleavage and polyadenylation mutants. *Genes & Development* 26, 163-175.
- Stodola, J.L., and Burgers, P.M. (2016). Resolving individual steps of Okazaki-fragment maturation at a millisecond timescale. *Nat Struct Mol Biol* 23, 402-408.
- Szilard, R.K., Jacques, P.-É., Laramée, L., Cheng, B., Galicia, S., Bataille, A.R., Yeung, M., Mendez, M., Bergeron, M., Robert, F., *et al.* (2010). Systematic identification of fragile sites via genome-wide location analysis of γ -H2AX. *Nature Structural & Molecular Biology* 17, 299-305.
- Técher, H., Koundrioukoff, S., Carignon, S., Wilhelm, T., Millot, Gaël A., Lopez, Bernard S., Brison, O., and Debatisse, M. (2016). Signaling from Mus81-Eme2-Dependent DNA Damage Elicited by Chk1 Deficiency Modulates Replication Fork Speed and Origin Usage. *Cell Reports* 14, 1114-1127.
- Techer, H., Koundrioukoff, S., Nicolas, A., and Debatisse, M. (2017). The impact of replication stress on replication dynamics and DNA damage in vertebrate cells. *Nat Rev Genet* 18, 535-550.
- Tercero, J.A., Longhese, M.P., and Diffley, J.F. (2003). A central role for DNA replication forks in checkpoint activation and response. *Mol Cell* 11, 1323-1336.
- Teytelman, L., Thurtle, D.M., Rine, J., and van Oudenaarden, A. (2013). Highly expressed loci are vulnerable to misleading ChIP localization of multiple unrelated proteins. *Proc Natl Acad Sci U S A* 110, 18602-18607.
- Torres-Rosell, J., Sunjevaric, I., De Piccoli, G., Sacher, M., Eckert-Boulet, N., Reid, R., Jentsch, S., Rothstein, R., Aragon, L., and Lisby, M. (2007). The Smc5-Smc6 complex and SUMO modification of Rad52 regulates recombinational repair at the ribosomal gene locus. *Nat Cell Biol*.
- Weinert, T. (1997). A DNA damage checkpoint meets the cell cycle engine. *Science* 277, 1450-1451.
- Weinert, T.A., and Hartwell, L.H. (1988). The RAD9 gene controls the cell cycle response to DNA damage in *Saccharomyces cerevisiae*. *Science* 241, 317-322.
- Wu, X., An, X., Zhang, C., and Huang, M. (2018). Clb6-Cdc28 Promotes Ribonucleotide Reductase Subcellular Redistribution during S Phase. *Mol Cell Biol* 38.
- Yabuki, N., Terashima, H., and Kitada, K. (2002). Mapping of early firing origins on a replication profile of budding yeast. *Genes Cells* 7, 781-789.
- Yoshida, K., Bacal, J., Desmarais, D., Padioulet, I., Tsaponina, O., Chabes, A., Pantesco, V., Dubois, E., Parrinello, H., Skrzypczak, M., *et al.* (2014). The Histone Deacetylases Sir2 and Rpd3 Act on Ribosomal DNA to Control the Replication Program in Budding Yeast. *Molecular Cell* 54, 691-697.
- Zegerman, P., and Diffley, J.F.X. (2010). Checkpoint-dependent inhibition of DNA replication initiation by Sld3 and Dbf4 phosphorylation. *Nature* 467, 474-478.
- Zeman, M.K., and Cimprich, K.A. (2014). Causes and consequences of replication stress. *Nat Cell Biol* 16, 2-9.
- Zhang, W., and Durocher, D. (2010). De novo telomere formation is suppressed by the Mec1-dependent inhibition of Cdc13 accumulation at DNA breaks. *Genes Dev* 24, 502-515.

- Zhao, X., Chabes, A., Domkin, V., Thelander, L., and Rothstein, R. (2001). The ribonucleotide reductase inhibitor Sml1 is a new target of the Mec1/Rad53 kinase cascade during growth and in response to DNA damage. *EMBO J* 20, 3544-3553.
- Zhao, X., Muller, E.G., and Rothstein, R. (1998). A suppressor of two essential checkpoint genes identifies a novel protein that negatively affects dNTP pools. *Mol Cell* 2, 329-340.
- Zhao, X., and Rothstein, R. (2002). The Dun1 checkpoint kinase phosphorylates and regulates the ribonucleotide reductase inhibitor Sml1. *PNAS* 99, 3746-3751.
- Zou, L., and Elledge, S.J. (2003). Sensing DNA Damage Through ATRIP Recognition of RPA-ssDNA Complexes. *Science* 300, 1542-1548.