



HAL
open science

Positively Frequency-Dependent Interference Competition Maintains Diversity and Pervades a Natural Population of Cooperative Microbes

Olaya Rendueles, Michaela Amherd, Gregory Velicer

► **To cite this version:**

Olaya Rendueles, Michaela Amherd, Gregory Velicer. Positively Frequency-Dependent Interference Competition Maintains Diversity and Pervades a Natural Population of Cooperative Microbes. *Current Biology - CB*, 2015, 25 (13), pp.1673-1681. 10.1016/j.cub.2015.04.057 . hal-02613099

HAL Id: hal-02613099

<https://hal.science/hal-02613099>

Submitted on 8 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Current Biology

Positively Frequency-Dependent Interference Competition Maintains Diversity and Pervades a Natural Population of Cooperative Microbes

Highlights

- Positively frequency-dependent interference competition is common in *M. xanthus*
- Positively frequency-dependent selection maintains microbial diversity
- Interference competition appears to be contact dependent and is often lethal

Authors

Olaya Rendueles, Michaela Amherd, Gregory J. Velicer

Correspondence

olaya.rendueles@env.ethz.ch

In Brief

Rendueles et al. show that bacterial strains that are severely inferior competitors at intermediate frequency are competitively dominant at high frequency. Such positive frequency dependence pervades a natural population of *Myxococcus xanthus*, promotes diversity in patchily distributed populations, and is mediated by contact-dependent mechanisms.



Positively Frequency-Dependent Interference Competition Maintains Diversity and Pervades a Natural Population of Cooperative Microbes

Olaya Rendueles,^{1,*} Michaela Amherd,¹ and Gregory J. Velicer¹

¹Institute for Integrative Biology, Department of Environmental Sciences, ETH Zürich, 16 Universitätstrasse, 8092 Zürich, Switzerland

*Correspondence: olaya.rendueles@env.ethz.ch

<http://dx.doi.org/10.1016/j.cub.2015.04.057>

SUMMARY

Positively frequency-dependent selection is predicted from theory to promote diversity in patchily structured populations and communities, but empirical support for this prediction has been lacking. Here, we investigate frequency-dependent selection among isolates from a local natural population of the highly social bacterium *Myxococcus xanthus*. Upon starvation, closely related cells of *M. xanthus* cooperate to construct multicellular fruiting bodies, yet recently diverged genotypes co-residing in a local soil population often antagonize one another during fruiting-body development in mixed groups. In the experiments reported here, both fitness per se and strong forms of interference competition exhibit pervasive and strong positive frequency dependence (PFD) among many isolates from a centimeter-scale soil population of *M. xanthus*. All strains that compete poorly at intermediate frequency are shown to be competitively dominant at high frequency in most genotype pairings during both growth and development, and strongly so. Interference competition is often lethal and appears to be contact dependent rather than mediated by diffusible compounds. Finally, we experimentally demonstrate that positively frequency-dependent selection maintains diversity when genotype frequencies vary patchily in structured populations. These results suggest that PFD contributes to the high levels of local diversity found among *M. xanthus* social groups in natural soil populations by reinforcing social barriers to cross-territory invasion and thereby also promotes high within-group relatedness. More broadly, our results suggest that potential roles of PFD in maintaining patchily distributed diversity should be investigated more extensively in other species.

INTRODUCTION

Recently developed sequencing technologies have revealed previously unfathomed degrees of microbial diversity in all habitable environments, including soil [1, 2]. Ecological factors

expected to promote diversity [3, 4] include spatial structure per se [5] and variable abiotic [4, 6, 7] and biotic [4, 8] environmental features, but the actual contributions of these factors to natural microbial diversity often remain obscure. Among biotic factors, cooperative and competitive social interactions can each affect levels of intra-specific diversity, as can both resource and interference competition [4, 7, 9–11].

Many competitive interactions are frequency dependent, either negatively or positively [12–15]. Negative frequency dependence (NFD) of fitness has long been recognized to promote diversity when the fitness ranks of competitors reverse at a threshold frequency and rarity confers a fitness advantage [14, 16–19]. Fitness-rank-reversing NFD has been documented in many organisms [20, 21], including experimentally evolved populations of bacteria [7, 19, 22]. In contrast, positive frequency dependence (PFD) was predicted by early models to reduce diversity due to the systematic loss of rare types [23–26]. This prediction was supported by experiments with bacteria that engage in toxin-mediated interference competition. When cells of *Escherichia coli* that produce an anti-competitor toxin were mixed with non-producers, producers either increased toward fixation or decreased toward extinction, contingent on their starting frequency and the competition environment [27]. However, more recently developed theory has predicted that PFD should actually maintain diversity, rather than reduce it, when populations patchily occupy spatially structured habitats [18, 28]. In this view, spatial variation in competitor frequencies generates local priority effects such that different genotypes (or species, in community scenarios) numerically—and therefore also competitively—dominate distinct local territory patches.

Interference competition is common among microbes [29, 30] and evolves readily in experimental populations [31]. It is thus likely to play a major role in shaping microbial diversity [9, 11, 32]. However, the prevalence of interference competition among conspecific neighbors in local natural populations, the effect of frequency on such interference-competition fitness relationships, and the implications of any such frequency effects in the maintenance of microbial diversity remain little explored.

Myxococcus xanthus is a soil bacterium that engages in several cooperative behaviors, including motility [33, 34], predation [35, 36], and starvation-induced development into multicellular fruiting bodies [37]. *M. xanthus* soil populations are highly diverse, both genotypically [38] and phenotypically [34, 35, 39, 40], even at very small spatial scales. For example, in one centimeter-scale soil population of *M. xanthus* in Tübingen, Germany, dozens of socially distinct genotypes were found to coexist among only 78 independent isolates [41]. Social interactions

during codevelopment when *M. xanthus* natural isolates are forcibly mixed are often severely antagonistic, even among very closely related strains [42, 43].

Here we analyze the frequency dependence and modes of competitive interactions among isolates from the Tübingen population [41, 43]. All of the examined isolates produce large and similar numbers of spores during starvation-induced development in pure culture [43]. In contrast, in many pairwise 1:1 mixes among these strains, codevelopmental fitness is highly asymmetric, with inferior competitors suffering large reductions in spore production due to the presence of the superior competitor. Pairwise codevelopment fitness ranks among nine isolates were previously found to be strongly hierarchical, thus highlighting the question of how so many genotypes can coexist in a local population [43, 44].

Using these same Tübingen isolates, we tested whether relative fitness between antagonistic strains is frequency dependent during both vegetative growth and codevelopment. We further performed experiments to demonstrate how PFD can maintain diversity in patchy environments. Finally, we tested whether observed antagonisms can be explained by diffusible secretions or rather involve contact-dependent mechanisms.

RESULTS

Rarity Is Strongly Disadvantageous Even for Strains that Are Competitively Dominant at Intermediate Frequencies

We tested the relationship between frequency and fitness in pairwise competition mixes among three focal isolates (A23, A47, and A96) during both vegetative growth on high-nutrient agar (without swarm expansion; see [Supplemental Experimental Procedures](#)) and starvation-induced development. For both types of competition, we employed a relative fitness parameter, W_{ij} , that was originally formulated for codevelopment competitions (see [Supplemental Experimental Procedures](#)) [42, 43]. Positive values of W_{ij} indicate that a focal strain i has higher relative fitness than its competitor j , and negative values indicate the opposite. For all three possible pairs, the competitor that was previously shown to be inferior in 1:1 developmental competitions (hereafter always referred to as the $IC_{0.5}$ competitor) [42, 43] was allowed to start competitions in the majority (9:1 and 99:1), as well as at a 1:1 ratio.

At 1:1 mixing ratios, fitness ranks conformed to previous developmental competitions, with A47 winning over both other strains and A96 winning over A23 ([Figure 1A](#); [Table S1](#)) [42, 43]. During both growth and development, relative fitness of the $IC_{0.5}$ competitor exhibited extremely strong PFD. Indeed, PFD was so strong that the fitness ranks of the paired competitors reversed strikingly as a function of frequency in all competitions ([Figure 1A](#); [Table S1](#)). Thus, among these isolates, strains that are rare within groups are unable to invade common strains during either growth or development, even when the rare strain has a strong fitness advantage at intermediate frequencies.

We also tested for strain-interaction effects on the absolute performance of each competitor in mixed groups by calculating the parameter $C_i(j)$, which quantifies the effect of mixing two strains i and j on the viable population size of strain i , relative to pure culture controls [42]. Negative values of $C_i(j)$ indicate

that strain i had lower spore production during development (or population yield during growth) in the presence of j than in pure culture, and positive values indicate the opposite. In all cases, competitive dominance by the majority strain (positive W_{ij}) was caused solely or predominantly by strongly negative effects of mixing on the performance of the rare competitor, rather than by positive responses to mixing by the majority competitor ([Figures 1B](#) and [1C](#)). In other words, during both growth and development, the absolute performance of the rare strain was always much worse in the presence of the majority competitor than in pure culture (usually by multiple orders of magnitude). This pattern indicates the presence of antagonism mechanisms that operate as a function of majority status, irrespective of which strain dominates when strain frequencies are similar.

PFD Is Widespread in a Natural Centimeter-Scale Soil Population

We also analyzed eleven other pairings among the nine Tübingen centimeter-scale isolates that exhibited large fitness asymmetries in a previous study [43] for PFD of both relative fitness (as reflected by W_{ij}) and antagonistic interactions (as reflected by negative $C_i(j)$ values) during codevelopment. On average across all 14 developmental competition pairs (including pairings among A23, A47, and A96), $IC_{0.5}$ competitors that (by definition) lose at intermediate frequency (average $W_{ij} = -4.45$ in 1:1 mixes) win strongly when they start at high frequency (average $W_{ij} = 0.71$ and 2.54 in 9:1 and 99:1 mixes, respectively; [Figure 2A](#)). Among individual competition pairs, fitness rank reversed as a function of frequency in all but one case, with the exact shape of the frequency-fitness relationship varying across strains ([Figure S1A](#); [Table S1](#)).

Correspondingly, the average sporulation efficiency of $IC_{0.5}$ partners was greatly reduced by their paired competitors in the 1:1 mixes, but not when the former started at high frequencies ([Figure 2B](#); $IC_{0.5}$ mean $C_i(j)$ values of -3.73 , -1.02 , and 0.06 in 1:1, 9:1, and 99:1 mixes, respectively). Inversely, the absolute performance of strains that win at intermediate frequency was unaffected by $IC_{0.5}$ strains in the 1:1 mixes (mean $C_i(j) = 0.10$) but was strongly reduced by the presence of $IC_{0.5}$ competitors when the latter started at high frequencies (mean $C_i(j)$ values of -1.9 and -2.61 in 9:1 and 99:1 mixes, respectively; [Figure 2B](#)). Although the average overall relationship between frequency and unidirectional mixing effects is extremely strong and clear ([Figure 2](#)), the frequency interval across which the directionality of antagonism reversed varied across competition pairs ([Table S1](#); [Figure S1](#)). In most mixes, significant antagonistic interactions occurred primarily in one direction, as chimerism in these 14 pairwise mixes significantly reduced total group productivity only at the 9:1 mixing ratio (on average across all 14 pairs), and in that case only to a relatively small degree ([Table S2](#)).

Strong PFD antagonisms are specifically caused by interactions between distinct natural isolates and are not due to either marker or density effects on the ability of each to strain to develop or compete. In competition experiments between differentially marked variants of the same natural isolate, no significantly negative effects of mixing on spore production were found for any strain at any frequency ([Figure S2](#)). Additionally, although total cell density was constant across all sporulation assays, it is also clear that antagonism patterns cannot be explained by




Figure 1. Fitness and Inter-strain Antagonisms among Three Natural Isolates Are Highly Frequency Dependent during both Growth and Development

(A) Relative fitness values (W_{ij}) of the competitor that is inferior in 1:1 mixes ($IC_{0.5}$) during vegetative growth (white circles) and codevelopment (black circles). (B and C) The effects of mixing two strains on the \log_{10} -transformed population size of each after co-culture during growth on an agar surface (B) and codevelopment (C) are given as $C_i(j)$, with i representing the focal strain of interest. Black and white circles represent the inferior and superior competitors, respectively, in 1:1 mixes. Maximally negative $C_i(j)$ values are ~ -8.5 for vegetative growth and ~ -7.0 for codevelopment. Error bars represent 95% confidence intervals.

genotype-specific density effects. If mere genotype-specific density were responsible for spore production patterns, any one genotype should show the same pattern across mixing frequencies regardless of which competitor it is paired with, but this is clearly not the case. For example, frequency-dependent patterns of spore production exhibited by A47 (Figure 1C, left and right panels; Figure S1G), A96 (Figure 1C, middle graph; Figures S1I–S1K), and A75 (Figures S1B, S1C, and S1E) differ greatly as a function of competitor genotype.

Positive Frequency Dependence Maintains Diversity in Patchy Environments

Theory predicts that PFD can maintain diversity in patchily distributed populations and communities due to local priority effects [18, 28]. We tested this prediction experimentally by performing developmental competitions with two pairs of natural *M. xanthus* isolates (A30 versus A96 and A96 versus A47) in which overall initial competitor frequency was invariantly 0.5 across each of two treatments, but in which the spatial distribu-

tion of competitor frequencies was homogeneous in one treatment and heterogeneous in another, with local majority status varying across patchy subpopulations (Figure 3). More specifically, in the spatially homogeneous treatments, each competition arena consisted of four subpopulation patches across which the two competitors were evenly distributed at a 1:1 ratio. In the spatially heterogeneous treatments, each subpopulation patch was initiated with one competitor in a 99:1 majority, with the majority competitor alternating patch-wise to give an overall 1:1 competitor ratio across all four patches. Subpopulation patches were placed in close proximity on a common agar plate so that patches could migrate toward one another and, in the absence of any mechanisms of kin discrimination that operate at the inter-patch level, merge. Arena populations were allowed to compete over two complete cycles of development and one growth phase between the developmental cycles (see Supplemental Experimental Procedures). To maintain population spatial structure that was not eliminated by motility-driven inter-patch migration, patch subpopulations were harvested




Figure 2. Average Fitness and Mixing Effects across 14 Natural Isolate Pairs during Developmental Competitions Are Highly Frequency Dependent

(A) Average fitness values of the IC_{0.5} competitors across all 14 competition pairings at three initial IC_{0.5} frequencies.

(B) Average mixing effects ($C_i(j)$) on spore production across all 14 competition pairings when inoculated at three initial IC_{0.5} frequencies. Values for competitors that are inferior (IC_{0.5}) and superior (SC_{0.5}) at a 1:1 mixing ratio are represented with black and white circles, respectively.

Error bars represent 95% confidence intervals.

separately at the end of the first round of development and cultured separately during the growth phase before being placed in the same relative configuration on a common plate for the second round of development.

During both developmental cycles, patches in the spatially homogeneous treatment merged until patch boundaries were not visible. In contrast, in the spatially heterogeneous treatment, inter-patch demarcation lines were clearly evident, indicating the operation of kin discriminatory social incompatibilities between the distinct majority genotypes in neighboring patches (data not shown) and suggesting that such pre-existing social barriers reduced migration across patches.

We used the Shannon index, which is determined by both genotype richness (here maximally two) and genotype frequencies [45], as our measure of diversity. After two developmental cycles, diversity was entirely or nearly eliminated in all replicates of the spatially homogeneous treatment for both competition pairs (Figure 3), with the IC_{0.5} competitor driven to complete or near extinction in all cases. In arenas in which the IC_{0.5} was still present, it was reduced to extremely low numbers (fewer than 350 cells from initial populations of 10^9) and thus should have gone extinct had the competitions been continued through a third round.

In striking contrast, not only were both competitors maintained in all replicates in the spatially heterogeneous treatment, but they were maintained at nearly equal frequencies in all arenas (Fig-

ure 3). As expected from the outcomes of non-patchy competition arenas (Figures 1 and 2), diversity was maintained almost exclusively at the inter-patch level in this treatment, as the minority strain went completely or nearly extinct within all patch sub-populations (30/48 and 18/48 patches, respectively), with each competitor dominating the two patches in which it started in the majority. Thus, although inter-patch migration was allowed, it either did not occur at all due to pre-existing social barriers or was insufficient to allow competitors that win at intermediate frequencies to significantly reduce IC_{0.5} frequencies in patches in which the latter started in the majority. In conclusion, PFD maintained biological diversity at nearly maximal levels under conditions in which it would be rapidly eliminated in the absence of PFD.

Diffusible Toxins Were Not Detected

Frequency-dependent fitness-rank reversals cannot be explained by intrinsic differences between competitors that are expressed in pure culture but rather imply strong interactions between strains. We investigated whether the antagonistic interactions documented here are mediated by diffusible molecules or rather are contact dependent, first by asking whether conditioned supernatant from vegetative cultures of each strain affected the growth rate of other isolates for seven strain pairs. No significant effects were found across or within pairs (Figure S3). Because development triggers massive changes in gene expression [37], we also tested whether diffusible secretions might be produced specifically during development. To do so, we assessed the ability of three focal isolates that show similar spore production in pure culture (A23, A47, and A96; one-way ANOVA test for variation between strains, $p = 0.806$) to develop in the presence of liquid buffer that had been pre-conditioned by each of the other isolates. First, we prepared cell-free supernatant from development cultures submerged in a liquid buffer (MC7). We then qualitatively assessed fruiting-body formation and quantitatively estimated spore production of each strain in presence of MC7 conditioned by each isolate and fresh, unconditioned buffer (see Supplemental Experimental Procedures).

After 5 days, cultures supplemented with pre-conditioned supernatant from any source strain exhibited greater developmental maturity (darker and larger fruiting bodies) than did controls treated with fresh MC7 (Figure 4). Interestingly, development by A23 and A47 in media that had been pre-conditioned by any strain yielded significantly more spores than did development in unconditioned MC7 alone (Figure 4; two-way ANOVA, $F = 13.82$, $p < 0.0001$). This result demonstrates the presence of a diffusible cue to which both A23 and A47 respond that appears to be produced similarly by all three strains. Importantly, no differences were observed between the effects of different pre-conditioned supernatants from the three source strains on either fruiting-body formation (qualitative visual assessment) or spore production for any of the three recipient strains (Figure 4). This result strongly suggests that the severe developmental antagonisms between these strains are not caused by diffusible antagonistic compounds secreted during development.

In theory, a diffusible secretion might hinder developmental performance by an effect other than direct toxicity, such as




Figure 3. Positive Frequency Dependence Maintains Diversity in Patchily Distributed Populations

(A and B) Each competition arena consisted of four subpopulation patches across which the overall initial competitor frequency in each arena was invariantly 0.5. In homogeneous arenas (A), the two competitors were evenly distributed at a 1:1 ratio in all four patches. In the spatially heterogeneous arenas (B), each subpopulation patch was initiated with one competitor in a 99:1 majority. Subpopulation patches were placed 2 cm apart from each other so that patches could migrate toward one another and, in the absence of any mechanisms of kin discrimination that operate at the inter-patch level, merge.

(C and D) Total spore counts of each natural isolate after two cycles of competition for two pairwise competitions, A96-A30 (C) and A96-A47 (D). For each pairing, the $IC_{0.5}$ is represented with white bars. Error bars represent 95% confidence intervals.

(E and F) In competition treatments in which competitors were distributed homogeneously across subpopulation patches, the $IC_{0.5}$ competitor (white circles) is driven to extinction in both competition pairs A30-A96 (E) and A47-A96 (F), whereas both competitors are maintained at high levels when initial competitor frequencies vary across subpopulation patches (black circles). Dashed lines represent the maximum diversity level possible in this experimental system (both strains present at equal frequency). T0: beginning of competition; D1 and D2: first and second rounds of codevelopment; G1: intermediate liquid growth phase.

A96 across three environments: nutrient-rich liquid, nutrient-rich agar plates, and during development at 1:1 mixing ratios (Table 1).

Within each competitor pairing, the same strain was antagonized in all three environments (Table 1), suggesting that similar or identical mechanisms of antagonism may operate across environments. In the liquid competitions, the winning

hindrance of cell-cell adhesion necessary for effective development [37]. Using the same three focal strains, we tested whether a diffusible factor secreted during vegetative growth alters the ability of competitor cells to adhere, but no such effect was found (Figure S3). Taken together, our results indicate that developmental antagonisms are not caused by obligately produced diffusible secretions.

Antagonism during Vegetative Growth Is Often Lethal and Appears to Be Contact Dependent

If antagonisms are contact dependent, we predicted that they would be more pronounced in populations growing on solid surfaces than in populations dispersed in liquid due to lower density in the latter. We thus compared antagonistic mixing effects on strain performance among pairings between A23, A47, and

strain in all three pairings significantly reduced cell counts of the inferior competitor by more than a factor of ten relative to expectations from pure culture growth (i.e., C_i/j was always lower than -1.0 ; Table 1). Even more strikingly, in two competitions (A23-A47 and A96-A47) these antagonisms caused significant decreases in the absolute size of the $IC_{0.5}$ subpopulation (73% and 69% for A23 and A96, respectively; $p < 0.05$ for differences between initial and final subpopulation sizes, one-sample t test), indicating that antagonisms were lethal to many cells even in liquid.

Although antagonisms during liquid growth were large in their population-level effects, such effects were much greater in competitions on high-nutrient agar surfaces (Table 1). In this environment, the vast majority of cells of victimized competitors were clearly killed by the winning strain. This was evidenced by lack




Figure 4. Diffusible Secretions Enhance Rather Than Inhibit Sporulation across Antagonistic Strains

(A) Cultures of A23 were allowed to develop while submerged in MC7 starvation buffer that was either unconditioned (MC7) or previously conditioned by A23 (self-self control, MC7-A23), A47 (MC7-A47), or A96 (MC7-A96). Pictures were taken after 5 days of development (magnification $\sim 50\times$).

(B) Average total spore production by three focal strains after development while submerged in either fresh MC7 media (white bars) or in MC7 preconditioned by each of the three strains (gray bars). * $p < 0.05$; *** $p < 0.001$, post hoc tests with Bonferroni correction for differences from fresh MC7 treatment. Error bars represent 95% confidence intervals.

of regrowth of the antagonized strains from post-competition cultures on selective plates containing antibiotic to which only the winning strain is sensitive (Figure S4). In most replicates across all three pairings for all three frequencies tested, the entire losing subpopulation was completely killed (38/56), and in the remaining 18 cases it was greatly reduced. Overall, the absence of negative supernatant effects combined with increased degrees of cell death on surfaces relative to liquid cultures strongly suggests that antagonisms are contact dependent.

We also conducted a fourth type of competition experiment in which 1:1 competition cultures were allowed to swarm outward on high-nutrient agar surfaces. After one week of swarming on both hard and soft CTT agar, samples from four locations evenly distributed around each swarm perimeter were transferred onto CTT hard agar selecting for each of two competing isolates. Although the swarming rates of these three strains are indistinguishable in pure culture ([34] and unpublished data), in the mixed competitions the inferior competitor was completely absent from the swarming edge in all replicates. Thus, inclusion of population-level swarming as a component of fitness allowed antagonisms to even more severely eliminate competitors.

DISCUSSION

In concordance with recently developed theory [18, 28], we have experimentally demonstrated that PFD of fitness can maintain, rather than purge, diversity in patchily structured populations in which the majority genotype varies across local patches. Although PFD has been documented in other microbial systems, the role of PFD in maintaining natural microbial diversity, whether

Table 1. Antagonisms Have Smaller Population-Level Effects in Liquid Than on Solid Surfaces

Strain Pair	$C_i(j)$ Development	$C_i(j)$ Liquid Growth	$C_i(j)$ Surface Growth	p Values Liquid versus Surface
A23-A47	-4.02 ± 1.74	-2.32 ± 1.01	-8.24 ± 0.43	<0.001
A23-A96	-4.79 ± 1.46	-1.65 ± 0.74	-8.60 ± 0.24	<0.001
A47-A96	-4.44 ± 1.55	-1.68 ± 0.67	-8.26 ± 0.43	<0.001
$C_i(j)$ max	~ -7	~ -7	~ -8.5	

$C_i(j)$ values for inferior competitors in pairwise competitions between the three focal strains in three different environments (development, liquid, and surface growth) are shown. Data represent mean \pm 95% confidence intervals. The inferior competitor (i.e., the strain with a negative W_{ij}) in 1:1 mixes ($IC_{0.5}$) is underlined. p values are based on post hoc Bonferroni testing. Theoretically maximally negative $C_i(j)$ values are also shown.

at the population or community level, remains largely unexplored. The pervasiveness of extremely strong PFD in a centimeter-scale population of *M. xanthus* documented here suggests that similar patterns may be found among other microbial species. More broadly, PFD also occurs among plants [13] and animals [46], including in human economic systems [47]. Our empirical results suggest that the potential for PFD to help maintain biological, cultural, and technological diversity in spatially structured systems of all types should be more broadly investigated.

PFD traits are expected to generate self-organized population structure in which distinct areas differ in their most common genotype [28]. PFD reinforces genotype-territory boundaries by disfavoring migrants that are locally rare after crossing such boundaries. In our experiments, we showed that patches in which $IC_{0.5}$ competitors were at high initial frequency (0.99) were not overrun from neighboring patches by the alternative competitor (that dominates in 1:1 competitions), thus allowing diversity to be maintained across patches. Moreover, PFD-mediated diversity is expected to increase with the degree to which populations are spatially fragmented by uninhabited patches [28].

Myxococcus cells live in patchily distributed kin groups in the soil [41, 44]. Groups that form fruiting bodies on soil particles often include detectable genetic diversity, but that diversity is usually of endemic origin and tends to be low relative to diversity across kin groups at millimeter and centimeter scales (S. Kraemer, S. Wielgoss, and G.J.V., unpublished data). Laboratory experiments suggest that kin discriminatory traits often prevent genetically distinct neighboring kin groups from freely merging upon encounter in the soil. Such kin discrimination during heterotypic colony encounters is prevalent among the centimeter-scale isolates examined here [43] and reduces co-aggregation into chimeric fruiting bodies at the interface of oncoming colonies formed by the three focal strains A23, A47, and A96 (unpublished data). These colony-merger incompatibilities alone are predicted to increase kin-group structure in soil populations (above that expected from simply clonal reproduction on a structured surface by motile cells [43, 48]). Indeed, in our patchy competition experiments, clearly visible social barriers were established between neighboring patches dominated by different strains that reduced or entirely prevented patch merger, whereas such merger occurred readily when patches were homogenous with respect to initial competitor frequency.

The severe PFD among centimeter-scale neighbors documented here should reinforce fragmented kin-group structure and local diversity generated by colony-merger incompatibilities. If some such incompatibilities were to only reduce, but not entirely prevent, cell-cell encounters between members of distinct kin groups, PFD interference competition should nonetheless reduce the frequency of successful cross-patch invasion events. Cells that migrate into territory numerically dominated by a distinct social type will suffer the PFD disadvantage of being rare. Together, then, as shown in Figure 3, colony-merger incompatibilities [43, 49, 50] and PFD acting against inter-group migration can promote the maintenance of structured diversity by benefiting genotypes that would be severely inferior if local migration were unlimited and genotype frequencies were spatially homogenized. Also, although all of the strains examined here are highly proficient at development in clonal groups [39, 43], PFD antagonisms may also function together with colony-merger incompatibilities to limit the spread of socially defective cheaters across kin groups and thus increase equilibrium levels of within-group cooperation [51].

Of course, our results do not explain how, if many genotypes have an advantage only at high frequency, those genotypes became established in local territory patches in the first place. In one scenario, a new mutant might arise at or near the edge of a patch subpopulation and migrate to occupy an unoccupied patch. There appear to be many soil micro-patches unoccupied by *M. xanthus* at any given time, as a substantial fraction of soil samples do not yield fruiting bodies, even from areas in which *M. xanthus* is common ([41] and unpublished data). Alternatively, patch subpopulations might first diverge in evolutionary isolation at traits that affect developmental competitiveness before subsequently encountering one another due to migration.

The role of PFD in interference competition on structured surfaces has been previously examined for unidirectional antagonisms mediated by obligately produced diffusible toxins [27, 52]. Working with toxin-producing bacteria that exhibit PFD, Chao and Levin showed that toxin production remained beneficial even at very low producer frequencies [27]. In contrast, working with yeast, Greig and Travisano found that the cost of toxin production can outweigh benefits derived from harm to competitors when both overall cell density and toxin-producer frequency are sufficiently low [52]. Indeed, not only was an outright advantage to toxin production lost when both total density and producer frequency were very low, but producers were at a slight disadvantage (a few percent). This result showed that fitness ranks in a PFD relationship can reverse in a frequency-dependent manner, at least when overall population density is sufficiently low.

Our results differ greatly from those of Greig and Travisano. First and foremost, the *M. xanthus* antagonisms documented here are, in all but one case, bidirectional between competitors. In Greig and Travisano's yeast experiments, toxin non-producers exhibited only a slight advantage over producers when the latter were at very low density and frequency. Moreover, this advantage by non-producers was due to resource competition (differential metabolic costs) rather than interference competition [52]. In our experiments with bacteria, fitness-rank reversals were caused by bidirectional interference competition (Figures 2A and S1A). In almost every competition pair, the strain

that strongly antagonized its partner at a 1:1 mixing ratio itself became a victim of strong antagonism when its frequency was reduced. Second, fitness-rank reversals in *M. xanthus* competitions do not require either extreme frequency fluctuations—reversal thresholds were at $IC_{0.5}$ frequencies between either 0.5 and 0.9 (nine cases; Figures S1D, S1F–S1L, 2B, and 2C) or between 0.9 and 0.99 (four cases; Figures 2B, S1B, S1C, and S1E)—or low population densities.

In our experiments, antagonistic interactions during both development and growth were quantitatively severe and in many cases caused outright extinction of the losing subpopulation. Multiple mechanisms of lethal contact-dependent interference competition by bacteria have been reported, including those mediated by type VI secretion systems [53], *rhs*-like proteins [54], and *cdi*-like contact-dependent inhibition systems [55]. The *M. xanthus* genome carries homologs to several such systems (*MXAN_4800–MXAN_4815*, *MXAN_5799*, *MXAN_6679*), some of which are involved in motility and development [56, 57]. Future work is required to address whether any of these homologs are involved in the pervasive antagonistic interactions reported here, whether the molecular mechanisms causing PFD are functionally linked to mechanisms causing colony-merger incompatibilities [43], and the evolutionary mechanisms by which PFD interference competition first evolved [58].

A novel instance of PFD was previously shown to evolve rapidly and de novo in an experimental laboratory population of *M. xanthus*. Through the effect of a single mutation, a socially defective cheater that exhibited a negatively frequency-dependent fitness relationship toward an ancestral cooperator [59, 60] was transformed into a derived, socially proficient cooperator that exhibited positively frequency-dependent fitness dominance over the ancestor [59, 60]. This reversal of NFD into PFD by one mutation suggests that new forms of PFD may evolve easily in natural populations and thereby increase PFD-mediated diversity.

SUPPLEMENTAL INFORMATION

Supplemental Information includes four figures, two tables, and Supplemental Experimental Procedures and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2015.04.057>.

AUTHOR CONTRIBUTIONS

O.R. and G.J.V. designed the experiments, analyzed the data, and wrote the manuscript. O.R. and M.A. performed the experiments.

ACKNOWLEDGMENTS

We are grateful to Michiel Vos and two anonymous reviewers for helpful comments on the manuscript. This work was funded in part by an EMBO Long-Term Fellowship to O.R.

Received: January 12, 2015

Revised: March 31, 2015

Accepted: April 24, 2015

Published: June 4, 2015

REFERENCES

1. Roesch, L.F., Fulthorpe, R.R., Riva, A., Casella, G., Hadwin, A.K.M., Kent, A.D., Daroub, S.H., Camargo, F.A.O., Farmerie, W.G., and Triplett, E.W.

- (2007). Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME J.* 1, 283–290.
2. Young, I.M., and Crawford, J.W. (2004). Interactions and self-organization in the soil-microbe complex. *Science* 304, 1634–1637.
 3. Cordero, O.X., and Polz, M.F. (2014). Explaining microbial genomic diversity in light of evolutionary ecology. *Nat. Rev. Microbiol.* 12, 263–273.
 4. Vos, M., Wolf, A.B., Jennings, S.J., and Kowalchuk, G.A. (2013). Micro-scale determinants of bacterial diversity in soil. *FEMS Microbiol. Rev.* 37, 936–954.
 5. Rainey, P.B., and Travisano, M. (1998). Adaptive radiation in a heterogeneous environment. *Nature* 394, 69–72.
 6. Bressan, M., Mougel, C., Dequiedt, S., Maron, P.A., Lemanceau, P., and Ranjard, L. (2008). Response of soil bacterial community structure to successive perturbations of different types and intensities. *Environ. Microbiol.* 10, 2184–2187.
 7. Rainey, P.B., Buckling, A., Kassen, R., and Travisano, M. (2000). The emergence and maintenance of diversity: insights from experimental bacterial populations. *Trends Ecol. Evol.* 15, 243–247.
 8. Crawford, J.W., Deacon, L., Grinev, D., Harris, J.A., Ritz, K., Singh, B.K., and Young, I. (2012). Microbial diversity affects self-organization of the soil-microbe system with consequences for function. *J. R. Soc. Interface* 9, 1302–1310.
 9. Bürger, R., and Gimelfarb, A. (2004). The effects of intraspecific competition and stabilizing selection on a polygenic trait. *Genetics* 167, 1425–1443.
 10. Chesson, P. (2000). General theory of competitive coexistence in spatially-varying environments. *Theor. Popul. Biol.* 58, 211–237.
 11. Rosenzweig, M.L. (1978). Competitive speciation. *Biol. J. Linn. Soc. Lond.* 10, 275–289.
 12. Ross-Gillespie, A., Gardner, A., West, S.A., and Griffin, A.S. (2007). Frequency dependence and cooperation: theory and a test with bacteria. *Am. Nat.* 170, 331–342.
 13. Eppstein, M.J., Bever, J.D., and Molofsky, J. (2006). Spatio-temporal community dynamics induced by frequency dependent interactions. *Ecol. Modell.* 197, 133–147.
 14. Levin, B.R. (1988). Frequency-dependent selection in bacterial populations. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 319, 459–472.
 15. Christiansen, F.B. (1988). Frequency dependence and competition. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 319, 587–600.
 16. Bever, J.D. (1999). Dynamics within mutualism and the maintenance of diversity: inference from a model of interguild frequency dependence. *Ecol. Lett.* 2, 52–62.
 17. Lenski, R.E., and Hattingh, S.E. (1986). Coexistence of two competitors on one resource and one inhibitor: a chemostat model based on bacteria and antibiotics. *J. Theor. Biol.* 122, 83–93.
 18. Molofsky, J., and Bever, J.D. (2002). A novel theory to explain species diversity in landscapes: positive frequency dependence and habitat suitability. *Proc. Biol. Sci.* 269, 2389–2393.
 19. Velicer, G.J., Kroos, L., and Lenski, R.E. (2000). Developmental cheating in the social bacterium *Myxococcus xanthus*. *Nature* 404, 598–601.
 20. Gigord, L.D.B., Macnair, M.R., and Smithson, A. (2001). Negative frequency-dependent selection maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza sambucina* (L.) Soo. *Proc. Natl. Acad. Sci. USA* 98, 6253–6255.
 21. Takahashi, Y., and Kawata, M. (2013). A comprehensive test for negative frequency-dependent selection. *Popul. Ecol.* 55, 499–509.
 22. Rozen, D.E., and Lenski, R.E. (2000). Long-term experimental evolution in *Escherichia coli*. VIII. Dynamics of a balanced polymorphism. *Am. Nat.* 155, 24–35.
 23. Levins, R. (1974). Discussion paper: the qualitative analysis of partially specified systems. *Ann. N Y Acad. Sci.* 237, 123–138.
 24. May, R. (1974). *Stability and Complexity in Model Ecosystems*. (Princeton University Press).
 25. Frank, S.A. (1994). Spatial polymorphism of bacteriocins and other allelopathic traits. *Evol. Ecol.* 8, 369–386.
 26. Kimura, M., and Ohta, T. (1971). *Theoretical Aspects of Population Genetics*. (Princeton University Press).
 27. Chao, L., and Levin, B.R. (1981). Structured habitats and the evolution of anticompensator toxins in bacteria. *Proc. Natl. Acad. Sci. USA* 78, 6324–6328.
 28. Molofsky, J., Bever, J.D., and Antonovics, J. (2001). Coexistence under positive frequency dependence. *Proc. Biol. Sci.* 268, 273–277.
 29. Iwasa, Y., Nakamaru, M., and Levin, S.A. (1998). Allelopathy of bacteria in a lattice population: Competition between colicin-sensitive and colicin-producing strains. *Evol. Ecol.* 12, 785–802.
 30. Riley, M.A., and Wertz, J.E. (2002). Bacteriocins: evolution, ecology, and application. *Annu. Rev. Microbiol.* 56, 117–137.
 31. Lemonnier, M., Levin, B.R., Romeo, T., Garner, K., Baquero, M.R., Mercante, J., Lemichez, E., Baquero, F., and Blázquez, J. (2008). The evolution of contact-dependent inhibition in non-growing populations of *Escherichia coli*. *Proc. Biol. Sci.* 275, 3–10.
 32. Kerr, B., Riley, M.A., Feldman, M.W., and Bohannan, B.J.M. (2002). Local dispersal promotes biodiversity in a real-life game of rock-paper-scissors. *Nature* 418, 171–174.
 33. Mauriello, E.M.F., Mignot, T., Yang, Z., and Zusman, D.R. (2010). Gliding motility revisited: how do the myxobacteria move without flagella? *Microbiol. Mol. Biol. Rev.* 74, 229–249.
 34. Vos, M., and Velicer, G.J. (2008). Natural variation of gliding motility in a centimetre-scale population of *Myxococcus xanthus*. *FEMS Microbiol. Ecol.* 64, 343–350.
 35. Morgan, A.D., MacLean, R.C., Hillesland, K.L., and Velicer, G.J. (2010). Comparative analysis of myxococcus predation on soil bacteria. *Appl. Environ. Microbiol.* 76, 6920–6927.
 36. Berleman, J.E., and Kirby, J.R. (2009). Deciphering the hunting strategy of a bacterial wolfpack. *FEMS Microbiol. Rev.* 33, 942–957.
 37. Rajagopalan, R., Sarwar, Z., Garza, A.G., and Kroos, L. (2013). Developmental gene regulation. In *Myxobacteria: Genomics, Cellular and Molecular Biology*, Z. Yang, and P.I. Higgs, eds. (Caister Academic Press).
 38. Kraemer, S.A., and Velicer, G.J. (2011). Endemic social diversity within natural kin groups of a cooperative bacterium. *Proc. Natl. Acad. Sci. USA* 108 (Suppl 2), 10823–10830.
 39. Kraemer, S.A., Toups, M.A., and Velicer, G.J. (2010). Natural variation in developmental life-history traits of the bacterium *Myxococcus xanthus*. *FEMS Microbiol. Ecol.* 73, 226–233.
 40. Krug, D., Zurek, G., Revermann, O., Vos, M., Velicer, G.J., and Müller, R. (2008). Discovering the hidden secondary metabolome of *Myxococcus xanthus*: a study of intraspecific diversity. *Appl. Environ. Microbiol.* 74, 3058–3068.
 41. Vos, M., and Velicer, G.J. (2006). Genetic population structure of the soil bacterium *Myxococcus xanthus* at the centimeter scale. *Appl. Environ. Microbiol.* 72, 3615–3625.
 42. Fiegna, F., and Velicer, G.J. (2005). Exploitative and hierarchical antagonism in a cooperative bacterium. *PLoS Biol.* 3, e370.
 43. Vos, M., and Velicer, G.J. (2009). Social conflict in centimeter- and global-scale populations of the bacterium *Myxococcus xanthus*. *Curr. Biol.* 19, 1763–1767.
 44. Velicer, G.J., and Vos, M. (2009). Sociobiology of the myxobacteria. *Annu. Rev. Microbiol.* 63, 599–623.
 45. Magurran, A.E. (1988). *Ecological Diversity and Its Measurements*. (Princeton University Press).
 46. Allen, J.A., and Weale, M.E. (2005). Anti-apostatic selection by wild birds on quasi-natural morphs of the land snail *Cepaea hortensis*: a generalised linear mixed models approach. *Oikos* 108, 335–343.

47. Faber, A., and Frenken, K. (2009). Models in evolutionary economics and environmental policy: Towards an evolutionary environmental economics. *Technol. Forecast Soc. Change* *76*, 462–470.
48. Hallatschek, O., Hersen, P., Ramanathan, S., and Nelson, D.R. (2007). Genetic drift at expanding frontiers promotes gene segregation. *Proc. Natl. Acad. Sci. USA* *104*, 19926–19930.
49. Dienes, L. (1946). Reproductive processes in *Proteus* cultures. *J. Bacteriol.* *51*, 585.
50. Munson, E.L., Pfaller, M.A., and Doern, G.V. (2002). Modification of dienes mutual inhibition test for epidemiological characterization of *Pseudomonas aeruginosa* isolates. *J. Clin. Microbiol.* *40*, 4285–4288.
51. Strassmann, J.E., Gilbert, O.M., and Queller, D.C. (2011). Kin discrimination and cooperation in microbes. *Annu. Rev. Microbiol.* *65*, 349–367.
52. Greig, D., and Travisano, M. (2008). Density-dependent effects on allelopathic interactions in yeast. *Evolution* *62*, 521–527.
53. Ho, B.T., Dong, T.G., and Mekalanos, J.J. (2014). A view to a kill: the bacterial type VI secretion system. *Cell Host Microbe* *15*, 9–21.
54. Koskiniemi, S., Lamoureux, J.G., Nikolakakis, K.C., t'Kint de Roodenbeke, C., Kaplan, M.D., Low, D.A., and Hayes, C.S. (2013). Rhs proteins from diverse bacteria mediate intercellular competition. *Proc. Natl. Acad. Sci. USA* *110*, 7032–7037.
55. Aoki, S.K., Pamma, R., Hernday, A.D., Bickham, J.E., Braaten, B.A., and Low, D.A. (2005). Contact-dependent inhibition of growth in *Escherichia coli*. *Science* *309*, 1245–1248.
56. Youderian, P., and Hartzell, P.L. (2007). Triple mutants uncover three new genes required for social motility in *Myxococcus xanthus*. *Genetics* *177*, 557–566.
57. Konovalova, A., Petters, T., and Søgaard-Andersen, L. (2010). Extracellular biology of *Myxococcus xanthus*. *FEMS Microbiol. Rev.* *34*, 89–106.
58. Velicer, G.J., Mendes-Soares, H., and Wielgoss, S. (2014). Whence social diversity? Ecological and evolutionary analysis of the myxobacteria. In *Myxobacteria: Genomics, Cellular and Molecular Biology*, Z. Yang, and P.I. Higgs, eds. (Caister Academic Press).
59. Fiegna, F., Yu, Y.T.N., Kadam, S.V., and Velicer, G.J. (2006). Evolution of an obligate social cheater to a superior cooperator. *Nature* *441*, 310–314.
60. Yu, Y.T.N., Yuan, X., and Velicer, G.J. (2010). Adaptive evolution of an sRNA that controls *Myxococcus* development. *Science* *328*, 993.

Current Biology

Supplemental Information

**Positively Frequency-Dependent Interference
Competition Maintains Diversity and Pervades
a Natural Population of Cooperative Microbes**

Olaya Rendueles, Michaela Amherd, and Gregory J. Velicer

SUPPLEMENTAL FIGURES




Figure S1. Frequency-dependent interactions among eleven natural isolate pairs during development for both fitness (A) and unidirectional mixing effects (B-L). (Related to Figure 2). Graph features are the same as for Fig. 1, except all results are from co-development experiments. Panel A shows W_{ij} values for the same eleven strain pairs represented in panels B-L. Error bars show 95% confidence intervals. See Table S1 for associations between values and specific strain pairs.




Figure S2. Mixing effect ($C_i(j)$) during codevelopment between variants of the same natural isolate marked with different antibiotic resistance. (Related to Figure 1 and 2). $C_i(j)$ are shown for the three focal strains at the initial frequencies of 0.5 (A), 0.99 (B) and 0.01 (C). * $p < 0.05$, one sample t -test for difference from zero. ‘K’ = kanamycin resistant, ‘R’ = rifampicin resistant. Error bars show 95% confidence intervals.




Figure S3. Conditioned supernatant from growing antagonist strains does not affect growth rate or agglutination patterns. (Related to Figure 4). **A** Natural isolates were grown in only fresh CTT media (black squares) and in media conditioned by pure-culture supernatant from their own genotype (grey squares) or from antagonistic strains (empty squares). Error bars represent 95% confidence intervals. Natural isolates are ordered according to their fitness ranking during co-development as described by Vos & Velicer [S1] (strongest overall competitor (A75) on left, weakest (A9) on right). **B-D.** Exponentially grown cultures of natural isolates supplemented with unconditioned buffer (TPM, filled black), self-conditioned supernatant (filled grey circles) and supernatant conditioned by antagonistic strains (open circles) for A23 (**B**), A47 (**C**) and A96 (**D**). Reduction of absorbance over time reflects the degree of cell-cell adhesion. Error bars are not shown to allow data visibility but are available upon request.

1. Grow strains independently and standardize to 2.5×10^7 CFU/ml.
2. Grow cocultures statically on an agar surface.
3. Recover all cells and deposit on selective plates.
4. Compare swarming rate to standard curves.




Figure S4. Procedure for non-swarming competitions on high-nutrient surfaces.

(Related to Table 1). Two differentially marked competitors are grown independently in liquid culture, diluted to the same density ($\sim 2.5 \times 10^7$ cells/ml), mixed at specified ratios and spread (in a $200 \mu\text{l}$ aliquot) across the entire surface of ten ml of hard agar within a 50 ml flask. Flasks are incubated without shaking for 24 hours at 32°C , 90% rH. After the competition period, cells are harvested and resuspended in $50 \mu\text{l}$ CTT liquid. The whole population is then spotted onto the center of a selective agar plate that contains antibiotic (Ab) that allowed growth of only one competing strain. This experiment is performed in duplicate within each independent biological replicate to allow selection of both competing strains.

SUPPLEMENTAL TABLES

Table S1. Relative fitness during co-development. The difference between \log_{10} -transformed spore production by paired competitors (W_{ij}) [S2] is shown for all competition pairs at all three mixing ratios. W_{ij} values (\pm 95% confidence interval) of the left competitor (IC_{0.5}, the inferior competitor at an initial 1:1 ratio) are shown. Dark grey background indicates mixes in which the IC_{0.5} strain has significantly higher fitness than its competitor. The one competition pair for which qualitative fitness rank did not reverse at even the 99:1 ratio (A30 : A75) is italicized.

	1 : 1	9 : 1	99 : 1
A9 : A23	-2.72 \pm 0.67	1.90 \pm 1.28	2.60 \pm 0.51
A23 : A75	-4.85 \pm 1.22	-2.06 \pm 0.90	1.20 \pm 1.43
A30 : A75	-6.00 \pm 0.11	-6.67 \pm 0.51	<i>-1.15 \pm 1.06</i>
A30 : A41	-4.76 \pm 1.60	4.59 \pm 0.42	4.05 \pm 0.12
A85 : A41	-1.66 \pm 2.04	2.83 \pm 2.92	2.83 \pm 1.04
A9 : A75	-6.11 \pm 0.17	-2.66 \pm 1.97	0.7 \pm 2.47
A23 : A47	-4.68 \pm 1.54	-5.23 \pm 2.05	1.01 \pm 1.22
A23 : A96	-4.72 \pm 1.83	-4.12 \pm 1.05	1.79 \pm 0.32
A96 : A47	-4.21 \pm 1.65	3.33 \pm 3.19	4.45 \pm 0.24
A9 : A96	-5.27 \pm 0.89	0.54 \pm 1.61	2.97 \pm 0.57
A30 : A96	-2.87 \pm 2.01	5.10 \pm 0.44	4.23 \pm 0.32
A30 : A94	-5.45 \pm 0.29	5.16 \pm 0.28	4.06 \pm 0.23
A85 : A96	-5.77 \pm 0.83	2.07 \pm 0.19	2.59 \pm 0.31
A30 : A47	-3.15 \pm 0.97	5.15 \pm 0.23	4.27 \pm 0.34
Grand mean	-4.45 \pm 0.71	0.71 \pm 2.16	2.54 \pm 0.87
(n = 14)			

Table S2. Effect of mixing on total group sporulation (B_{ij}) at three mixing ratios.

(Related to Figure 2). The left number of each ratio corresponds to the frequency of the left strain of each pair, which is the inferior competitor at an initial 1:1 mixing ratio (Table S1). Errors correspond to 95% confidence intervals. Means that differ significantly from zero ($p < 0.05$, one-sample t test) are shown in bold and the one significantly positive effect is italicized.

	1 : 1	9 : 1	99 : 1
A9 : A23	0.28 ± 0.20	0.02 ± 0.26	0.14 ± 0.20
A23 : A75	0.04 ± 0.08	- 0.34 ± 0.45	- 0.23 ± 0.36
A30 : A75	- 0.14 ± 0.29	- 0.17 ± 0.14	0.08 ± 0.26
A30 : A41	0.09 ± 0.69	- 0.21 ± 0.43	- 0.08 ± 0.32
A85 : A41	<i>0.46 ± 0.09</i>	- 0.09 ± 0.33	0.05 ± 0.27
A9 : A75	0.16 ± 0.09	0.16 ± 0.44	0.07 ± 0.79
A23 : A47	0.08 ± 0.18	- 0.16 ± 0.46	- 0.20 ± 0.56
A23 : A96	0.17 ± 0.44	- 0.64 ± 0.41	- 0.27 ± 0.46
A96 : A47	- 0.48 ± 0.25	- 0.15 ± 0.25	- 0.09 ± 0.30
A9 : A96	0.27 ± 0.35	- 0.27 ± 0.19	0.24 ± 0.53
A30 : A96	- 0.04 ± 0.14	0.13 ± 0.15	0.13 ± 0.19
A30 : A94	- 0.57 ± 0.42	0.05 ± 0.46	0.12 ± 0.46
A85 : A96	- 0.21 ± 0.52	- 0.19 ± 0.20	- 0.17 ± 0.21
A30 : A47	<i>-0.54 ± 0.09</i>	-0.04 ± 0.04	0.14 ± 0.28
Grand mean ($n = 14$)	- 0.03 ± 0.168	- 0.136 ± 0.108	- 0.006 ± 0.086

SUPPLEMENTAL EXPERIMENTAL PROCEDURES

Bacterial strains, growth conditions and plate preparation. (i) *Strains.* The *M. xanthus* strains used here were isolated from a 16 x 16 cm soil plot in Tübingen, Germany as described previously [S1] and are the same nine strains previously characterized for developmental fitness at 1:1 mixing ratios [S2]. All antibiotic resistant clones exhibited proper fruiting body formation, high levels of spore production and no significant marker effects on pure culture spore production were detected [S2]. Previously reported kanamycin-resistant strain variants of seven strains (A9, A23, A30, A41, A47, A85 and A96) and rifampicin-resistant variants of eight strains (A9, A23, A30, A41, A47, A75, A94 and A96) were used [S3]. Whenever possible (most pairings), competitions were performed with both reciprocal antibiotic-resistance combination for each pair (i.e. strain *i*-rifampicin with strain *j*-kanamycin and strain *i*-kanamycin with strain *j*-rifampicin) and averaged. (ii) *Growth conditions.* After inoculation from frozen stocks, strains were grown on CTT (10 mM Tris pH 8.0, 8 mM MgSO₄, 10g/L casitone, 1 mM KPO₄) [S4] hard (1.5%) agar plates for three or four days at 32 °C, 90% rH. Prior to experiments, strains were grown in eight ml CTT liquid for 24 hours at 32 °C with constant shaking (300 rpm) until they reached mid exponential phase (OD₆₀₀ ~0.5 (~5 x 10⁸ cells/ml)). Cell biovolume densities of exponential-phase cultures were estimated with a TECAN Genios™ plate reader. Antibiotics were used at 40 µg/ml for kanamycin and 5 µg/ml for rifampicin. (iii) *Plate preparation.* Swarming assay plates were prepared one day prior to assays by pouring 20 ml of CTT soft (0.5%) or hard (1.5%) agar into medium-sized (9 cm) Petri dishes and allowing them to solidify and in a laminar flow hood without lids for 15-20 minutes, after which they were capped and stored overnight at room temperature. For developmental assays, 10 ml of TPM (10 mM Tris pH 8.0, 8 mM MgSO₄, 1 mM KPO₄) [S5] hard (1.5% agar) was poured into small (5 cm) Petri dishes and subsequently prepared as above.

Effect of conditioned supernatant on growth rate. (i) *Preparation of conditioned supernatant.* Cultures were grown in 40 ml of CTT liquid for 24 h until and were then centrifuged for 15 min at 5000 g and supernatant was filtered through 0.22 μm filters. (ii) *Growth rate assays.* All strains were inoculated in four ml of CTT liquid media at an initial concentration of $\sim 5 \times 10^7$ cells/ml, to which was added four ml of either fresh CTT (control) or supernatant conditioned by either the same genotype or a different strain. Cultures were incubated at 32 °C with constant shaking (300 rpm) for 50 hours. One hundred (100) μl samples were withdrawn at multiple time points (after 4, 18, 21, 24, 27, 30, 33, 41, 44, 47, and 50 hours) and absorbance (OD_{600}) using a spectrophotometer (TECAN Genios™ plate reader) was measured. (iii) *Calculations.* Maximum growth rates were estimated as the average slope of $\ln(\text{absorbance})$ across replicates for the 6 h interval spanning the fastest average growth. Experiments were performed in five independent replicates for the three focal strains (A23, A47 and A96) and four for all other strains.

Agglutination assay. One ml of preconditioned supernatant (see above) or unconditioned media (control) was added to three ml of exponentially grown cultures and were then transferred to static test tubes. Absorbance (OD_{600}) of 100 μl samples from each tube was measured repeatedly over a 24-hour period on a Tecan GENios plate reader. Samples were removed from the uppermost layer of tube cultures, which remained static. Decreasing absorbance represents the settling of agglutinated cell clumps.

Development in submerged cultures. Development in submerged cultures was carried out as described previously [S6]. (i) *Preparation of cell-free conditioned developmental supernatant.* Cells were grown to mid-exponential phase, centrifuged and re-suspended to a

final density of $\sim 5 \times 10^9$ cells/ml in MC7 (10mM MOPS, 1mM CaCl₂, adjusted to a pH of 6.8) buffer. Fifty microliters of each suspension was spotted into the center of each well of a 24-well microtiter plate and allowed to dry under laminar flow before 500 μ l of MC7 was added to each well. One entire 24-well plate was inoculated with each strain. Plates were incubated at 32°C, 90%rH for five days, at which time conditioned MC7 from each well was pooled and filtered through a 0.44 μ m to remove fruiting bodies and cells. (ii) *Spore production in submerged cultures.* To assess the effect of preconditioned developmental supernatant, fresh cultures were grown as described above and spotted into 24-well microtiter plates. After drying, 500 μ l of either fresh or preconditioned MC7 were added. Plates were incubated at 32 °C, 90% rH for five days and production of heat-resistant myxospores was then quantified as follows. Microtiter plates were sealed with parafilm and incubated at 50 °C for 2.5 hours to select for viable spores. Then, 500 μ l of MC7 was added to reach a final volume of one ml per well. Each well was sonicated by microtip and spores were diluted and plated into CTT soft agar (0.5% agar). This experiment was performed in three independent replicates with three technical replicates each.

Developmental competition assays. Cultures were grown to mid exponential phase, centrifuged and resuspended with TPM (10 mM Tris pH 8.0, 8 mM MgSO₄, 1 mM KPO₄) liquid to $\sim 5 \times 10^9$ cells/ml. For pure-culture assays, 50 μ l of resuspended culture were spotted in the center of TPM plates (1.5% agar). For paired competition assays, pure resuspended cultures from paired competitors were mixed at the appropriate frequencies (1:1, 9:1 and 99:1) and 50 μ l of the mix were spotted on the center of TPM plates. Thus, the total initial cell number and density were the same in competitive mixtures and clonal treatments. Developmental plates were incubated for five days and then cell populations were harvested with a scalpel, transferred into one ml of double-distilled H₂O, and heated at 50 °C for two

hours to select for viable spores. Samples were then sonicated by microtip, diluted and dilution-plated into CTT soft agar (0.5% agar) with the appropriate antibiotic treatment. In cases of mixed competitions where no colonies grew at our lowest dilution factor (10^1), a value of nine spores/assay was entered for data analysis. In each competition mixture, the paired strains had distinct antibiotic-resistance markers (described previously [S2]), allowing quantification of each competitor's spore production. This was also the case for all other types of direct competition experiments. Appropriate controls mixing the same strain with different antibiotic variants were performed at 1:1, 1:99 and 99:1 ratios. No negative effects of mixing were observed for any strain at any initial frequency (Fig. S2).

Competitions during growth on high-nutrient agar surfaces.

Competitions without swarming. Because *M. xanthus* cells cohere during growth on agar plates, changes in the relative frequencies in mixed competitions without swarming (or lack of any change) were quantified by a method that did not involve dilution plating of harvested post-competition cultures. Because *M. xanthus* swarm expansion on agar plates is dependent on inoculum density [S7] we indirectly assessed competitor subpopulation sizes at the end of 24-h mixed competition experiments by measuring the extent of subsequent swarm expansion by competitor subpopulations placed on selective agar medium specific to each competitor and allowed to swarm outward for four days (Fig. S4). Note that no swarming occurred during the competitions *per se* (which were carried out over the entire surface of agar with glass flasks), but only after the end of the competitions as a means to assess the relative size of post-competition subpopulations with reference to independently generated standard curves of inoculum density vs. swarm expansion, as described in greater detail below.

To initiate non-swarming competitions on agar surfaces, 200 μL of pairwise mixes of differentially marked strains (or of clonal culture controls) at $\sim 2.5 \times 10^7$ cells/ml were spread

across the entire surface of ten ml of hard agar within a 50 ml flask. Flasks were incubated without shaking for 24 hours at 32 °C, 90% rH (Fig. S4). After the competition period, cells were harvested in one ml of CTT liquid by adding approximately ten glass beads to the liquid and shaking at 300 rpm for 15 minutes. The entire volume of suspended liquid was then centrifuged and the resulting pellet was resuspended in 50 μ L CTT liquid, which was then spotted onto the center of a selective agar plate that allowed growth of only one of the two competing strains. Plating of the total harvested population instead of an aliquot (and instead of dilution plating) avoids possible error due to cell-cell adhesion.

In parallel, pure cultures of each strain were inoculated onto agar plates at four initial densities ranging from $\sim 10^6$ - $\sim 10^9$ cells/ml for generation of standard curves that allowed estimation of competitor subpopulation sizes (at the end of competition periods) (Fig. S4). Mean swarming rates between 24 and 96 hours after inoculation of the pure cultures plates were used to generate a standard curve of swarm size vs. initial density that allowed inference of relative competitor subpopulation sizes after 24 hours of competition based on subsequent swarm sizes. Swarming rates of competitor subpopulations were also calculated as the mean extent of swarm expansion between 24 and 96 hours after inoculation. Two technical replicates within each of three independent replicates were performed for each competition.

Growth competitions with swarming. The fitness advantage of superior competitors relative to inferior competitors was demonstrated qualitatively by scoring the presence or absence of each competitor at the leading edge of swarms that had initially been mixed at a 1:1 ratio and spotted (10 μ L) onto the center of plates prepared as described above. Swarm perimeters were marked after one and four days of incubation and swarming rates were calculated for both pairwise competition mixes of different isolates and mixes of differentially marked variants of the same (controls). To test whether marked strains were present at the leading edge of swarming populations, samples from four locations evenly

distributed around each swarm perimeter were harvested with a sterile pipette tip and transferred onto CTT hard agar plates with different antibiotics selecting for either subpopulation. The presence or absence of growth on each plate was scored after one week of incubation.

Competitions in patchy arenas. The maintenance of diversity by PFD was demonstrated by competing two pairs of natural isolates (A96-A30 and A96-A47) over two cycles of development on buffered TPM agar plates in two different competition-arena treatments. In both treatments, the overall initial frequency of each competitor in each plate arena was 0.5 and cells were spotted in four subpopulation patches, but the treatments differed in the spatial distribution of genotypes across patches (Fig. 3). One treatment was spatially homogeneous, with pairwise strain mixes inoculated in 1:1 ratios in all four patches, whereas in the second spatially heterogeneous treatment strains were in a 99:1 ratio, with the majority strain alternating patches so that same total number of cells of each strain was inoculated in each four-patch arena. For each pairing of natural isolates, competitions with both possible reciprocal combinations of antibiotic-resistance marker states were performed within each of three independent replicates and data from the reciprocal marker combinations were pooled within replicates.

To initiate the experiment, cultures were grown to mid exponential phase, centrifuged and resuspended in starvation buffer to $\sim 5 \times 10^9$ cells/ml. Pure resuspended cultures were mixed at the appropriate frequencies (1:1, 1:99 and 99:1) and patch subpopulations were initiated with 50 μ l aliquots of the relevant mix ratio. The four patches in each arena were configured in a square and neighboring patches were initially separated by two centimeters. Developmental plates were incubated for five days at 32°C, after which each patch was then harvested with a scalpel independently, transferred into one ml of double-distilled H₂O,

heated at 50 °C for two hours to select for viable spores and sonicated. An aliquot of each sonicated population (0.5 ml) was used for dilution plating into CTT soft agar (0.5% agar) selective plates and subsequent CFU counting of each competitor type. The other half of the sonicated population was inoculated into nutrient-rich CTT liquid media and allowed to germinate. After two days, vegetative cells were grown to high cell density and were respotted onto fresh TPM plates as described above. After five days, each patch was harvested, sonicated and the size of each competitor subpopulation determined by dilution plating. Dilution plating was also performed at the end of the liquid growth phase between developmental cycles to allow assessment of frequency changes specific to the growth phase.

Calculations. Calculations of B_{ij} , $C_i(j)$ and W_{ij} , employed the equations defining these parameters reported by Fiegna & Velicer [S3]. (i) *Unidirectional mixing-effect parameter* $C_i(j)$. The effect of mixing two strains i and j on the productivity of focal strain i (total spore production during development or population yield during growth) was quantified by the one-way mixing effect parameter $C_i(j)$. To calculate this parameter, the expected log-transformed spore production of strain i based on pure-culture performance (corrected for the frequency at which strain i was added) was subtracted from its actual log-transformed spore production during competition with strain j . Positive $C_i(j)$ values indicate that strain i produced more spores (or grew to a higher population size in the growth experiments) in the presence of strain j than in pure culture, whereas a negative value indicates that mixing with j negatively affected spore production (or growth yield) of i . (ii) *Bidirectional mixing effect parameter* B_{ij} . B_{ij} is the difference between the actual total (log-transformed) group spore production (or cell count) in a mix of strains i and j and the value expected from pure culture performance of the same strains (corrected for initial frequencies of each strain). Positive and negative B_{ij} values indicate that total productivity is higher or lower, respectively, than

expected from pure culture performance. (iii) *Relative spore production*. The relative fitness (W_{ij}) of two strains i and j during development is calculated as the difference between the log-transformed number of spores produced by each strain in mixed competition, with positive values indicating that i produced more spores than j and negative values indicating the opposite. (iv) *Diversity indexes*. Shannon diversity index values were calculated as: $-\sum_i (n_i/N) * \log_2(n_i/N)$.

Statistics. All experiments were performed in at least three temporarily independent replicate blocks. Statistical tests were performed using Prism v5.0 (GraphPad Software). To estimate the significance of the negative overall mean of B_{ij} , the average value of B_{ij} for all fourteen pairs was calculated. These fourteen means were used as statistical units in a one-sample t -test of the difference between the grand mean and the null expectation of zero. Similarly, to calculate the mean of $C_i(j)$ the mean value of the focal competitor class was calculated across all fourteen pairs and this was used as the statistical unit to estimate the significance of the average one-way effect of mixing. All experiments were performed at least three times.

SUPPLEMENTAL REFERENCES

- S1. Vos, M., and Velicer, G.J. (2006). Genetic population structure of the soil bacterium *Myxococcus xanthus* at the centimeter scale. *Appl Env Microbiol* 72, 3615-3625.
- S2. Vos, M., and Velicer, G.J. (2009). Social conflict in centimeter-and global-scale populations of the bacterium *Myxococcus xanthus*. *Curr Biol : CB* 19, 1763-1767.
- S3. Fiegna, F., and Velicer, G.J. (2005). Exploitative and hierarchical antagonism in a cooperative bacterium. *PLoS Biol* 3, e370.
- S4. Kaiser, D. (1979). Social gliding is correlated with the presence of pili in *Myxococcus xanthus*. *Proc Nat Acad Sci USA* 76, 5952-5956.
- S5. Bretscher, A.P., and Kaiser, D. (1978). Nutrition of *Myxococcus xanthus*, a fruiting myxobacterium. *J Bacteriol* 133, 763-768.
- S6. Kuner, J.M., and Kaiser, D. (1982). Fruiting body morphogenesis in submerged cultures of *Myxococcus xanthus*. *J Bacteriol* 151, 458-461.
- S7. Kaiser, D., and Crosby, C. (1983). Cell movement and its coordination in swarms of *Myxococcus xanthus*. *Cell Mot* 3, 227-245.