



HAL
open science

Selenoproteins and Protection against Oxidative Stress: Selenoprotein N as a Novel Player at the Crossroads of Redox Signaling and Calcium Homeostasis

Sandrine Arbogast, Ana Ferreiro

► **To cite this version:**

Sandrine Arbogast, Ana Ferreiro. Selenoproteins and Protection against Oxidative Stress: Selenoprotein N as a Novel Player at the Crossroads of Redox Signaling and Calcium Homeostasis. *Antioxidants and Redox Signaling*, 2010. hal-02613577

HAL Id: hal-02613577

<https://hal.science/hal-02613577v1>

Submitted on 20 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Selenoproteins and Protection against Oxidative Stress: Selenoprotein N as a Novel Player at the Crossroads of Redox Signaling and Calcium Homeostasis

Sandrine Arbogast^{1,2} and Ana Ferreiro¹⁻⁴

Abstract

Healthy cells continually produce low levels of reactive oxygen species (ROS), which are buffered by multiple antioxidant systems. Imbalance between ROS production and elimination results in oxidative stress, which has been implicated in aging and in numerous human diseases, including cancer and diabetes. Selenoproteins are a family of proteins that contain the amino acid selenocysteine, encoded by an in-frame UGA. Those selenoproteins whose function is identified are catalytically active in redox processes, representing one of the main enzymatic antioxidant systems and important mediators of the beneficial role of selenium in human health. Nevertheless, the function of most selenoproteins remains unknown; this included Selenoprotein N (SelN), the only selenoprotein directly associated with a human genetic disease. Mutations of the SelN gene cause SEPNI-related myopathy, a particular early-onset muscle disorder. Recent studies have identified SelN as a key protein in cell protection against oxidative stress and redox-related calcium homeostasis. Furthermore, an effective *ex vivo* treatment of SelN deficiency has been identified, paving the way to a clinical therapy. In this review we discuss the physiological and pathophysiological role of SelN and the interest of SEPNI-related myopathy as a model paradigm to understand and target therapeutically other selenoproteins involved in human health and disease.

Introduction

LIVING CELLS CONTINUALLY PRODUCE LOW LEVELS of reactive oxygen species (ROS) and nitric oxide (NO) derivatives. Both ROS and NO influence cellular function by modulating excitation–contraction coupling (1, 82, 89, 101), glucose uptake (46), mitochondrial respiration (56), and gene expression (63). Furthermore, it has been recently demonstrated that ROS are part of the intracellular signaling cascade (45). The net activity of ROS or NO depends on the overall balance between synthesis and buffering; a correct balance is necessary to keep cells in a state of redox equilibrium and thus preserve normal functioning. When the equilibrium between ROS production and removal is disrupted, oxidative stress occurs. Increasing understanding of the importance of redox signaling pathways in the last years has modified the definition of oxidative stress, from ‘imbalance between oxidants and antioxidants’ to the more comprehensive ‘disruption of redox signaling and control’ (98). Oxidative stress has multi-

ple targets, can cause cellular dysfunction or death, and is accompanied by numerous biochemical indexes: lipid peroxidation, protein carbonyl formation, glutathione oxidation or depletion, and even DNA damage. Its involvement in the pathogenesis of various chronic diseases, including muscle and cardiovascular disorders, diabetes, neurological diseases, and cancer, is increasingly documented (11).

To neutralize these reactive species, cells possess several antioxidant systems with different chemistries and intracellular localizations (84, 107). Nonenzymatic antioxidants such as glutathione, vitamin E, ascorbate, and carotenoids, directly neutralize (scavenge) ROS, being destroyed upon oxidation. The main enzymatic antioxidant systems include catalase, the mitochondrial and cytosolic isoforms of superoxide dismutase (MnSOD and CuZnSOD, respectively) and, importantly, selenoproteins; indeed, the selenoenzymes glutathione peroxidases (GPxs) are one of the major cell protective systems (84). Superoxide dismutase (SOD) dismutates superoxide anions to H₂O₂, which is rapidly converted by catalase into H₂O and

¹Inserm, U787, Institut de Myologie, Paris, France.

²UPMC Univ Paris 06, UMR_S787, IFR14, Paris, France.

³AP-HP, Groupe Hospitalier Pitié-Salpêtrière, Centre de Référence des Maladies Neuromusculaires Paris-Est, Paris, France.

⁴AP-HP, Hôpital Raymond Poincaré, Service de Pédiatrie, Centre de Référence des Maladies Neuromusculaires GNMH, Paris, France.

molecular oxygen; selenoproteins such as glutathione peroxidase and thioredoxin reductase contribute, among others, to regenerate glutathione and thioredoxin oxidized by free radicals. A schematic overview of oxidant sources and main cell antioxidant systems is depicted in Fig. 1.

The human selenoproteome includes 25 characterized selenoproteins; each of them contains at least one selenocysteine, the biological form of selenium, encoded by a UGA codon and cotranslationally incorporated in response to a complex mechanism (9, 81) (reviewed by Donovan *et al.* in this issue). Selenoproteins are essential for mammals, as demonstrated by the fact that deletion of the selenocysteine tRNA gene, which controls the expression of the entire selenoproteome, induces early embryonic lethality in mice (15). Although the function of most selenoproteins remains unknown, the few which have been fully characterized are enzymes with oxidoreductase functions (81). Their catalytic group within the active site includes the selenocysteine residue, whose biochemical properties render it more reactive than thiols in nucleophilic reactions at physiological pH (43). Thus, selenocysteine is more actively oxidized than cysteine (79) and has the potential to repair oxidative damage in proteins by reducing tyrosyl radicals with higher efficiency (99). Selenium is a key factor in regulation of selenoprotein biosynthesis; selenium depletion in food supply has revealed a hierarchy in expression among different selenoproteins and tissues, GPx-1 being one of the first selenoproteins whose enzymatic activity and protein level decrease during selenium deprivation (17).

In the last years, significant progress has been made towards elucidating the functional importance of specific selenoproteins and their implication in human diseases, and large-scale selenium supplementation trials have been launched (64). However, only one selenoprotein, selenoprotein N (SelN), has been directly associated with a genetic disorder. This review will provide a synthetic overview of the antioxidant and redox signaling properties of known human selenoproteins, and will then focus on SelN, whose physiological and pathophysiological role has been recently elucidated, paving the way to the development of a pharmacological therapy for an infantile inherited disease.

Antioxidant Properties of Human Selenoproteins

The human selenoproteome consists of 17 selenoprotein families. Three major classes, the glutathione peroxidases (GPxs), thioredoxin reductases (TrxRs), and iodothyronine deiodinases (DIOs), were the first discovered and are extensively studied. DIOs are involved in thyroid hormone metabolism (8). GPxs and TrxRs have an antioxidant role and, glutathione and thioredoxin being involved in the cellular redox pathways, they modulate cell function by modifying redox status (104). Regulatory and functional interactions between both families of selenoenzymes have been observed (Table 1).

Glutathione peroxidases

Reduced glutathione (GSH) is the most abundant antioxidant; in muscle cells GSH is present in millimolar concentra-




FIG. 1. Schematic diagram depicting the different sites of ROS and NO generation by cells (*italic font*) and the main antioxidant enzymes present in a majority of human cell types (*shadowed*). Superoxide anions (O₂⁻), generated by mitochondria, NAD(P)H oxidase, 5-lipoxygenase, cyclooxygenase, or xanthine oxidase, are the parent molecule of the ROS cascade (*gray*), giving rise to oxygen-based derivatives including hydrogen peroxide (H₂O₂), and hydroxyl radical (OH). Antioxidant enzymes are shaded in *dark gray* (selenoproteins) or *light gray* (nonselenoprotein enzymes). Cat, catalase; CuZn SOD, copper/zinc superoxide dismutase; ecSOD, extracellular superoxide dismutase; ETC, electron transport chain; Mn SOD, manganese superoxide dismutase; mt NOS, mitochondrial nitric oxide synthase; nNOS, neuronal nitric oxide synthase; PLA2, phospholipase A2; reticulum, endoplasmic/sarcoplasmic reticulum; XO, xanthine oxidase; GPx, glutathione peroxidase; TrxR, thioredoxin reductase.

TABLE 1. OVERVIEW OF THE EXPERIMENTALLY-DOCUMENTED FUNCTIONS OF HUMAN SELENOPROTEINS

Maintenance of selenoproteins	
Transport and storage of Se	SeIP
Selenocysteine synthesis	SPS2
Modulation of GPx activity and expression	SeIH
Thyroid hormone metabolism	DIO 1-3
Transcription factor	SeIH
Ethanolamine phosphotransferase	Sell
Development	GPx-4, TrxR-1, TrxR-2, SeIW, SeIN?
Cell proliferation growth factor	TrxR-1
Inflammatory response	SPS1
Calcium homeostasis	SeIW, SeIT, SeIN
Redox signaling	
Regulation of transcription factors (NF-kB, AP-1, p53)	TrxR-1
Detoxification; antioxidant	
Reduction of H ₂ O ₂	GPx-1 and 3, TrxRs, SeIW, SeIK, SeIN?
Reduction of various organic peroxides	GPxs
Protein reduction	
Methionine sulfoxide reduction	TrxR-1, TrxR-2, SeIR
Thiol disulfide isomerase	TrxR-3, Sep15, SeIM
Reduction of sulfoxymethyl group	SeIR
Unknown enzymatic activity (TrxR-like?)	SeIN
Protein folding	Sep15, SeIM, SeIS, SeIN?
Apoptosis	TrxR-1, TrxR-2

Probable SeIN functions requiring further investigation are indicated by a question mark.

tions and is the most important biologically (88). GSH reacts nonspecifically with various oxidants to yield an oxidized glutathione dimer (GSSG) in a reaction facilitated by the selenoenzyme glutathione peroxidase (GPx). Glutathione is recycled from GSSG by a second enzyme, glutathione reductase; the ratio of GSH and GSSG indicates the cellular redox status.

There are at least five selenocysteine-containing glutathione peroxidases (GPx-1 to -4 and GPx-6) in mammalian cells. Human GPxs are selenoenzymes with different substrate specificity and tissue-specific expression (4). GPx-1 and the phospholipid hydroperoxide GPx-4 are ubiquitously expressed proteins, GPx-3 is secreted to plasma, GPx-2 is specific of the gastrointestinal epithelium, and GPx-6 is found in the olfactory epithelium and embryonic tissues (81). All except GPx-3 are localized in the cytosol; GPx-4 presents also in mitochondrial and nuclear isoforms. Although each GPx is a distinct selenoprotein, they all have in common a catalytic site formed by selenocysteine, glutamine, and tryptophan residues in their active center (67). Their antioxidant role is based on their capacity to reduce potentially damaging ROS (such as hydrogen peroxide and organic hydroperoxides) to harmless products like H₂O and alcohols, coupling their reduction with the oxidation of GSH (17).

In vitro and *in vivo* approaches have provided increasing proof of the importance of GPxs in cell homeostasis and survival. Cells overexpressing GPx-1 or GPx-4 are more resistant to oxidative stress elicited by different triggers of cell death (72, 111). These observations were corroborated using transgenic mice. Overexpression of either GPx-1 or GPx-4 resulted in increased survival of mice to the prooxidants paraquat and diquat (26, 86). Paradoxically, GPx-1 had a conflicting role in diquat- and peroxynitrite-induced oxidative injury, protecting against diquat-induced oxidative stress (attenuating NADH, NADPH, protein, and lipid oxidation) but promoting ni-

trostatic stress under peroxynitrite (41). This raises interesting questions about the specificity of oxidant and antioxidant actions; antioxidant protection might not be a general property for a given enzyme such as GPx-1, but rather depend on the specific nature of oxidants (26, 40, 41). Although initial observations did not show any phenotype (48), genetically modified mice lacking GPx-1 have later been shown to have growth retardation, increased mitochondrial oxidative damage (32) and susceptibility to toxicity following administration of pro-oxidants (60). GPx-1 also is associated with protection against viral infection pathogenicity. Thus, selenium and particularly GPx-1 deficiency favor redox-related mutations of benign coxsackievirus, rendering it pathogenic and leading to the viral-induced myocardial lesions characteristic of Keshan disease (7, 59). On the other hand, GPx-4 is a key component of the mammalian antioxidant network (93, 112) and has an indispensable role in cell differentiation during mouse development (49).

Although less well known, other GPxs have also antioxidant capacities. Thus, GPx-2 plays a critical role in protecting mammal gastrointestinal tract from toxicity by ROS and from cancer development, and GPx-3 is an efficient plasma antioxidant (17, 81).

Thioredoxin reductases (TrxRs)

TrxRs, thioredoxin (Trx), and NADPH form the thioredoxin system, a major cellular redox signaling in almost every living cell. TrxRs are involved in protection against reactive oxygen species through control of the redox state of the central molecule thioredoxin, using NADPH as a reducing agent. This system can also directly reduce numerous substrates, including hydroperoxides (13) or ubiquinone-10 (110). Thus, many cellular processes rely on the activity of these enzymes, including notably regulation of gene expression via redox control of transcription factors such as NF-kB or AP-1

(81). The three human TrxR isoenzymes contain one selenocysteine residue in their C-terminus. TrxR-1 is ubiquitously expressed in the cytosol (91, 94); TrxR-2 is localized in mitochondria and involved in control of mitochondrial redox processes; TrxR-3 is the testis-specific thioredoxin-glutathione reductase (44, 81). Like GPx-4, TrxR-1 and TrxR-2 are essential for mouse embryogenesis (28, 50). TrxR-1 has also the capacity of regulating the induction of the antioxidant enzyme heme oxygenase 1 (HO-1) (106).

Selenoprotein R (SelR)

Selenoprotein R (SelR), also known as MsrB1 or selenoprotein X (SelX), is a member of the methionine sulfoxide reductases family, one of the rare systems able to repair mammalian oxidized proteins by reducing methionine sulf-oxide residues (22, 54). SelR, located in the cytosol and nucleus, plays a major role in protein homeostasis during aging (22), thus contributing to delaying the aging process. Overexpression of SelR and other Msr activities leads to an increase in resistance to oxidative stress and life span (96). Furthermore, SelR and cysteine-containing isoforms have been demonstrated to have an important role in lens cell viability and protection against induced oxidative stress (70), potentially playing a role in cataract formation. A recent KO mouse model showed the strongest phenotype of SelR reduction and oxidative stress markers in liver and kidney, establishing an important contribution of SelR to the redox control in these organs (39).

Selenoprotein P

SelP (selenoprotein P), identified early through biochemical studies, is primarily secreted from liver cells and acts as transporter protein, supplying extrahepatic tissues with selenium (20, 78, 95). However, it also acts as a ROS-detoxifying enzyme (21). SelP is one of the key effectors of redox regulation and cell viability in myofibroblasts (53), reduces phospholipid hydroperoxide, provides protection against *Trypanosoma congolense* infection through its N-terminal (redox) domain (16, 92), and protects human astrocytes from induced oxidative stress (77, 97, 98).

Other selenoproteins

Other selenoproteins have been less extensively investigated, although experimental data support their antioxidant capacity. **Selenoprotein W (SelW)** (108) shares the redox motif with the GPx family and has a glutathione-dependant antioxidant activity (10, 51, 65). Its expression is closely modulated by selenium; nutritional selenium deprivation leads to severely reduced SelW expression as observed in white muscle disease (108). SelW is highly and preferentially expressed in proliferating myoblasts, protecting this and other cell types from oxidative stress (65). This selenoprotein has been proposed as a good marker to monitor immediate oxidation and toxic stress (81). **Selenoproteins localized in the ER** such as SelS, H, K, and Sep15 appear to be involved in regulation of intracellular ROS levels and protection from oxidative stress in cultured mammalian cells (for review, see Shchedrina, *et al.* in this issue). **SelS**, known for increasing production of inflammatory cytokines, is induced by ER stress and protects this cell compartment from oxidative stress (42).

SelH significantly reduces superoxide production in neuronal cells challenged with UVB irradiation (12). Furthermore, SelH may act as a transcription factor, upregulating expression of other selenoproteins in response to stress and increasing glutathione levels and GPx activity (12). **Sep15** is a 15 kDa selenoprotein characterized by a glutathione peroxidase-like enzymatic activity (57). Furthermore, in physiological conditions, Sep15 and **SelM** act as thiol-disulfide oxidoreductases (34). Both proteins, localized in the ER, are also involved in protein folding (34). Other ER proteins such as selenoproteins K, S, and T, have largely unknown functions. Only for **SelK** have antioxidant properties in cardiomyocytes been proposed (66).

Selenoprotein N, a Novel Key Player in Antioxidant Protection of Human Cells

Selenoprotein N is the only selenoprotein implicated so far in a human genetic disorder

In 1998, a peculiar and reportedly very rare form of congenital muscular dystrophy with early rigidity of the spine (rigid spine muscular dystrophy or RSMD1) was linked to chromosome 1p35-36 (38, 74). A few months later, *SEPN1*, the selenoprotein N gene localized at 1p36, was identified *in silico* (61). The previously described association between selenium deficiency and striated muscle dysfunction in livestock (25, 62) and humans (6, 14, 19, 59) immediately pointed *SEPN1* as a positional candidate for RSMD1. Indeed, homozygous or compound heterozygous *SEPN1* mutations were rapidly identified in most (75) although not all (76) the typical RSMD1 patients.

Surprisingly, mutations of the same gene, and often the same precise mutations (37), have thereafter been associated with three other early-onset muscle disorders, including two forms of congenital myopathy (the classical form of multi-minicore disease (37) and rare cases of congenital fibre type disproportion (27)) and a myopathy with protein aggregates (desmin-related myopathy with Mallory body-like inclusions (35)). Actually, careful retrospective reassessment showed that these four autosomal recessive muscle conditions share so much clinical and molecular identity that they are best understood as the same unique disorder, now termed SEPNI-related myopathy (SEPNI-RM) (35, 37, 68, 103). Independently of the initial diagnosis, all patients carrying *SEPN1* mutations share a very homogeneous clinical phenotype. This phenotype is marked by severe weakness of neck and trunk muscles from infancy, which leads to major scoliosis and life-threatening respiratory insufficiency in childhood or adolescence and contrasts with relatively preserved limb strength and ambulation. Conversely, the pathological presentation of the disease is unusually heterogeneous. Most SEPNI-RM muscle biopsies show small focal areas of mitochondria depletion and sarcomere disorganization (short core lesions or "minicores") in muscle fibers; necrosis and regeneration and/or protein aggregates are less frequent but may also be present. Since muscle disorders are classically defined according with their pathological manifestations, the large myopathological spectrum of SEPNI-RM explained the unexpected overlap between different categories of myopathies, led to a revision of the established classification in the field and suggested a novel, polymorphic pathophysiological mechanism.

Currently, more than 180 patients with *SEPN1* mutations have been identified worldwide, for the most part in Europe (AF, personal communication), suggesting that this emerging disease is not as rare as once thought. The *SEPN1* gene contains 13 exons and produces a 4.5-kb transcript with an open reading frame encoding a 590-amino acid protein; the selenocysteine residue with predicted catalytic activity is encoded by exon 10 (75). Mutations are distributed along the whole gene, excepting exon 3 which corresponds to an Alu cassette, contains a second in-frame selenocysteine codon and is mostly spliced out in humans (83). Most of them are nonsense mutations, microdeletions or insertions leading to frame-shifts, as well as splice-site mutations leading to aberrant pre-mRNA splicing (reviewed in (62)). This includes a nonsense mutation which converts the UGA selenocysteine (Sec) codon into a UAA stop codon, generating a premature termination codon and drastic mRNA decay (75). Missense mutations are relatively more common around or at the potential catalytic site. Interestingly, several mutations affect the *cis* sequences required for selenocysteine insertion. Thus, a homozygous point mutation in the 3' UTR SECIS element abolishes binding of the SBP2 factor, thereby preventing redefinition of the UGA Sec and leading to a premature termination codon and severe *SEPN1* mRNA reduction (87). One of the mutations localized in the Sec codon redefinition element (SRE) induces a mismatch in the conserved SRE stem-loop structure, weakens the mRNA structure and reduces Sec insertion efficiency (68). Notably, all the heterozygous carriers of a *SEPN1* mutation are healthy, attesting that 50% normal SelN suffices to ensure its function.

The lack of function phenotype associated to SelN has focused attention on this protein. Yet its functional characterization has lagged behind, hindered among others by SelN weak expression in postnatal tissues and the lack of specific technical tools. Thus, no biochemical activity has been attributed to SelN yet, the pathogenesis of SEPNI-RM remained undetermined for a long time and no specific treatment was available for this potentially lethal disorder. However, recent studies have shed light on SelN and brought exciting progress both on the scientific and clinical fields.

SelN is an ER glycoprotein with an early developmental expression and a putative but unclear role in muscle development

SelN is a 65-kDa transmembrane glycoprotein localized within the endoplasmic reticulum and containing a transmembrane addressing site close to an EF-hand motif (83). The topology of the protein has been determined recently: the N-terminal extremity is facing the cytoplasm, whilst the majority of the protein, including the predicted catalytic site and the C-terminal end, is located in the lumen of the ER (62).

Whilst the phenotype in SEPNI-RM involves exclusively skeletal muscles, human SelN is a ubiquitous protein (61) whose mRNA is detected in most fetal and, at a lower level, adult tissues (61, 75). Its expression is increased in proliferating cells such as fibroblasts and myoblasts, progressively decreasing during differentiation from myoblasts to myotubes (83).

An early developmental expression pattern of *SEPN1* was observed in zebrafish embryos in somites and notochord, which are the precursors of muscle and spinal cord, respec-

tively (105). Recent studies have confirmed that the murine *Sepp1* transcript is also expressed early in somites, appears restricted to the myotome, the subectodermal mesenchyme and the dorsal root ganglia at mid-gestation (with levels peaking at E12.5) and then strongly decreases until birth. An additional fall in protein levels, not correlated with transcript modifications, was observed in the perinatal period, suggesting that post-transcriptional mechanisms contribute to SelN regulation (24).

This embryonic pattern of expression evoked a potential role of SelN in muscle and spinal cord development. The consequences of SelN absence on muscle development have been analyzed in zebrafish embryos (29, 52) and in *Sepp1* deficient mouse embryos (Rederstorff *et al.*, unpublished observations; 24), throwing interesting although somewhat conflicting results. In zebrafish embryos, inhibition of the *SEPN1* gene by injection of antisense morpholinos did not prevent muscle formation but caused reduced embryo motility (29). Muscle ultrastructure disclosed defects in the sarcomeric organization (29, 52) and/or myofibril attachment (29), suggesting a role of SelN at early steps of zebrafish embryogenesis and in muscle maintenance. In contrast, *Sepp1* deficiency in mouse embryos did not alter somitogenesis nor expression of myogenic factors, suggesting that SelN is dispensable for these processes in this mammal model currently under validation (24). The relatively normal morphological appearance of skeletal muscles at birth in most SEPNI-RM patients suggests that the latter conclusion may also apply to humans. No study has yet reported data on a potential implication of SelN in muscle regeneration or in cell proliferation, a key point that merits to be addressed.

SelN cellular function: At the crossroads of redox signaling, cell stress, and calcium homeostasis

SelN structure predicts a putative although uncharacterized enzymatic activity. The biochemical activity of SelN remains unknown. To some extent, selenoproteins activity can be deduced from the sequence context of the selenocysteine residue, which constitutes a landmark of the catalytic center. SelN harbors a SCUG predicted catalytic site, reminiscent of the thioredoxin reductase GCUG motif (61), which suggests a putative reductase activity. However, classical thioredoxin reductases contain two other functionally important domains, the FAD and the NADPH-binding domains, which are absent in SelN, although this might be compensated by interactions with yet unidentified SelN partners (61). In addition, the highly accessible localization of the selenolate active site at the C-terminus of the thioredoxin reductases, supposed to confer them a broad range of substrates, contrasts with the localization of the active site in the central part of SelN; this lesser accessibility might reflect a higher selectivity of SelN for its substrate(s) (61).

SEPN1 expression is potentially regulated by cell stress signals. Analysis of the *SEPN1* promoter region predicts regulation of its expression by stress signals, supporting a potential implication of SelN in cell stress response pathways. NF- κ B is one of the main transcription factors involved in cellular responses to exogenous insults such as free radicals or cytokines. Stoytcheva *et al.* revealed that all 25 human selenoprotein genes contain putative NF- κ B response elements,

although only for DIO2, GPx-4, and SelS has regulation by NF- κ B been experimentally proven (100). Using the MatInspector software (Genomatix Software GmbH, Munich, Germany) (23), we identified five potential NF- κ B binding sites in the promoter region (2500 basepairs upstream of the transcription start site) of *SEPN1*, as well as a putative ERSE (ER stress response) (113) element (-572/-554, CCAATCCCCA TAACCCATG) (SA and AF, unpublished results). In addition, the *SEPN1* promoter contains one putative CpG island (5, 100), suggesting a possible role of methylation in expression of this gene (Fig. 2). Putative binding sites for the redox-sensitive transcription factor AP-1 have also been found in *SEPN1* (100). Presence of binding sites for multiple redox-sensitive transcription factors in the *SEPN1* promoter suggests that SelN might be regulated by different cell stressors (ROS, ER stress, and/or cytokines) in a cooperative or synergistic manner. Interestingly, the predicted sequences of the ERSE element and of the first NF- κ B putative binding site overlap (Fig. 2), suggesting a mutually exclusive, balanced regulation of *SEPN1* expression at this site. Experimental studies of *SEPN1* regulation are necessary to clarify this interesting point.

SelN and the pathophysiology of muscle disorders: SelN absence is associated with protein oxidation, Ca^{2+} handling abnormalities, and increased susceptibility to oxidative stress. Muscle disorders associated with selenoprotein N deficiency constitute a valuable disease model, providing further indications of a potential relation between SelN and redox homeostasis. Thus, axial muscles and especially the diaphragm, which are predominantly affected in *SEPN1*-related myopathy, are particularly sensitive to oxidative stress (3, 80). Moreover, there is a morphological and in some cases clinical overlap between *SEPN1*-RM and core myopathy associated with *RYR1* mutations (36). *RYR1* encodes the ryanodine receptor type 1 (RyR1), a sarcoplasmic reticulum (SR) Ca^{2+} release channel which plays a key role in excitation contraction (EC) coupling. Some primary *RYR1* mutations causing core myopathies have been associated with Ca^{2+} leaking (30, 31). Since both *RYR1* and *SEPN1* mutations are associated to minicore lesions, one can hypothesize that the SR proteins encoded by these two genes might be implicated

in common physiopathological pathways. The most obvious one would be redox homeostasis, since RyR1 is emerging as a paradigm of redox-sensor ion channel (71). Due to its important number of reactive cysteines (*i.e.*, thiols susceptible to redox-based modifications), RyRs are directly modulated by intracellular oxidants. NO and derivatives, via the nitrosylation of determined RyR thiol groups (33), may induce Ca^{2+} release from the SR into the cytosol. Along these lines, in livestock muscle disorders due to selenium deficiency (white muscle disease, rigid lamb syndrome), altered ability of the SR membranes to retain calcium has been described (25, 62). Conversely, Ca^{2+} overload and/or leaking could lead to oxidative/nitrosative stress (55) by activating calcium-dependant oxidant sources.

Recently, analysis of an *ex vivo* model of SelN deficiency allowed us to confirm that SelN plays an important role in cell defence against oxidative stress and Ca^{2+} homeostasis in human skeletal muscle cells (3). Primary cultured myotubes from patients with *SEPN1* null mutations showed a significant increase in basal intracellular oxidant activity compared to controls, with a significant contribution of NO; in contrast, mutant fibroblast oxidant activity remained unaltered. Oxidized protein content, a hallmark of oxidative stress, was increased both in SelN-devoid fibroblasts and myotubes; in the latter, excessive oxidation of the contractile proteins actin and myosin heavy chain II was observed, potentially contributing to mechanical dysfunction (3, 58). Furthermore, SelN-devoid human myotubes had an increased resting cytosolic Ca^{2+} concentration, together with reduced SR Ca^{2+} load and caffeine-induced Ca^{2+} release. This suggests that, in the absence of SelN, the ROS/NO generated by muscle cells regulate intracellular Ca^{2+} concentration via modulation of Ca^{2+} channels (RyR1 and/or SR Ca^{2+} ATPase), favoring Ca^{2+} release or leaking. Interestingly, independent studies using C2C12 myotubes knocked-down for *SEPN1* showed consistent results, namely increase of oxidant activity and protein carbonylation (73). In addition, zebrafish studies support the role of SelN as a RyR modifier. In zebrafish, SelN is physically associated with the ryanodine receptors; both types of proteins are required for the same cellular differentiation events and are needed for normal calcium fluxes in the embryo (52). Furthermore, in the absence of SelN, ryanodine receptors from zebrafish embryos or human diseased muscle have altered biochemical properties and have lost their normal sensitivity to redox conditions (52). Remarkably, *RYR* mutant and *SEPN1* mutant zebrafish have in common the disruption of slow muscle fiber formation and show a significant decrease in slow muscle fibers (29, 52). This contrasts with the human situation, stressing the important differences in redox signaling between zebrafish and mammals; indeed, in *SEPN1*-RM slow (type 1) muscle fibers are typically more abundant than fast fibers.

On the other hand, there is evidence to suggest that SelN is not exclusively a "redox chaperone" of the ryanodine receptors, but is also involved in modulation of redox defense and cell survival pathways. In the *ex vivo* cell model of SelN deficiency mentioned above, fibroblasts and muscle cells devoid of SelN showed a higher susceptibility to H_2O_2 -induced oxidative stress, manifested by a severely increased cell death rate (3). Interestingly, no compensatory upregulation of MnSOD or catalase was observed either in basal conditions (despite high oxidant activity) or after exogenous oxidant




FIG. 2. Schematic structure of the SelN promoter showing the positions of five putative NF- κ B binding sites, the ERSE element, and the GC-rich region. The numbers indicate the nucleotide positions relative to the transcription start site.

challenge, suggesting that SelN absence is associated with permanent abnormalities in antioxidant defense capacity (3). It has been previously reported that addition of H₂O₂ to mammalian cultured cells activates a series of growth arrest, cell death, and survival signaling pathways (63). In SelN-devoid cells treated with H₂O₂, cell death pathways (*e.g.*, p53 and NF-κB) might be more active than cell survival pathways (such as Akt, ERK, JAK/STAT, or phospholipase C-g1). Further studies are needed to dissect further these pathways, which could point towards potential therapeutic targets for selenoprotein N defects and other related disorders.

The current knowledge and our working model of SelN antioxidant role in human muscle cells (3) is summarized in Fig. 3. Absence of SelN is associated with oxidative/nitrosative stress, suggesting that this selenoprotein is involved in buffering and/or preventing overproduction of oxidants. Presence of the SelN selenocysteine in the lumen of the SR could protect proteins from excessive and irreversible oxidation/nitrosylation in at least two ways. First, since selenocysteine is more reactive than cysteine (62), SelN could react passively with oxidants more quickly than other cysteine-containing SR proteins (including the most cysteine-rich RyRs). Second, SelN could enzymatically exchange hydroxyl groups with oxidized proteins, hence allowing these to return to the reduced state and to pursue normal trafficking through the SR. Thus, SelN potentially protects general maturation and trafficking of proteins, preventing excessive protein carbonylation and degradation. In addition, SelN can protect specific target proteins of particular relevance for muscle function. This is the case of several contractile proteins whose function is known to be affected by oxidation [such as myosin heavy chain (2, 18), myosin light chain (102), actin (47, 58), troponin (85), or tropomyosin (109)]. Also, by limiting nitrosylation of RyR1, SelN would prevent excessive Ca²⁺ leakage from the SR into the cytosol. In the absence of SelN, Ca²⁺ leak reduces the amount of SR Ca²⁺ available for EC coupling (potentially contributing to contractile dysfunction and weakness) and leads to cytosolic Ca²⁺ overload, which in turn can increase oxidative/nitrosative stress (55) by activating calcium-dependant oxidant sources. Finally, SelN can contribute to activate a network of redox-sensitive transcrip-

tion factors that modulate the expression of antioxidant enzymes (predictedly including SelN itself) and determine the fate of the cell between survival and death.

This model would account for the variety of lesions in muscles from SEPNI-RM patients (Ca²⁺-related core lesions, protein aggregates, and/or necrosis and regeneration). More importantly, it shows SelN as a key protein at the crossroads of cell stress, redox signalling, and Ca²⁺ homeostasis pathways.

Antioxidants as a therapeutic approach to selenoprotein defects: Lessons from SEPNI-RM

SEPNI-RM is the only inherited disease due to a selenoprotein defect and the only structural myopathy primarily due to oxidative stress and antioxidant protection failure. As such, it represents an useful model paradigm for more complex, multifactorial disorders (such as cancer, cardiovascular disorders, diabetes, or aging) in which selenoproteins implication has been suggested (11). Therefore, finding a specific pharmacological treatment for this potentially-lethal, early-onset disease could also open interesting perspectives to human health at large.

The pathophysiological data described above suggest that the degree of muscle dysfunction in SEPNI-RM could be both free radical- and calcium-dependent; therefore, both pathophysiological pathways represent potential therapeutic targets for this disorder. Primary cultured cells from patients with *SEPNI* mutations show a quantifiable susceptibility to H₂O₂-induced oxidative stress, manifested by a significantly increased cell death rate (3). We used this cell phenotype to evaluate the *ex vivo* effect of antioxidant drugs targeting the primary pathogenic mechanism of the disease. Pretreatment of these SelN-devoid cells with the flavonoid fisetin or the carotenoid astaxanthin prior to H₂O₂ exposure had a partial or null protective effect. Interestingly, only pretreatment with N-acetyl cysteine (NAC) rescued this cell phenotype, significantly improving myoblast survival and rendering the cell death rate of SelN-devoid fibroblasts identical to that of control unchallenged cells. In addition, NAC normalized the levels of oxidized proteins in SEPNI-mutant myotubes (3). These findings demonstrate that NAC is an effective *ex vivo* treatment of SelN deficiency. Remarkably, they also establish that not all types of antioxidant drugs have the potential for therapeutic replacement of a particular selenoprotein function.

The specificity of NAC consists in its cysteine-donor activity, which could partially replace the hypothetical thiol exchanger activity of the selenocysteine in SelN. But this molecule has also other numerous biological roles (Fig. 4). NAC can act as a direct ROS scavenger and is a precursor for glutathione (GSH), an important nonenzymatic antioxidant. In addition, NAC increases the expression of glutathione peroxidase, the enzymatic antioxidant that oxidizes glutathione (GSH) to glutathione-disulfide (GSSH), thereby reducing H₂O₂ to H₂O and protecting cell components from oxidative damage. It has also been proposed that NAC downregulates the expression of NF-κB, which is a major mediator of inflammatory responses and controls expression of a large variety of genes (63). Therefore, NAC multifactorial antioxidant effect can potentially and specifically address at several levels the pathological consequences of SelN deficiency, and we propose that this antioxidant might be of particular interest to target defects of other selenoproteins in humans.




FIG. 3. Putative role of SelN in redox homeostasis of human muscle cells.




FIG. 4. Mechanisms of action and pathophysiological targets of N-acetyl cysteine (NAC) in SEP1N-related myopathy. ECC: excitation-contraction coupling; GPx: glutathione peroxidase; TFs: transcription factors.

NAC is one of the rare antioxidant drugs approved for human use, has been shown to inhibit muscle fatigue in healthy adults, and has no serious side effects when administered at the established dosage (69, 90). This, together with its *ex vivo* efficiency on SeIN-devoid cells, paves the way to a first therapeutic trial of SEP1N-RM in human patients, which is currently under preparation.

Conclusions, Bottlenecks, and Open Roads

In conclusion, important progress has been made in understanding the complex relation between selenoproteins and oxidative stress. Contribution of this particular protein family to different major cellular pathways—protection against oxidative stress, intracellular signaling, regulation of calcium homeostasis, or links with ER stress—represents an emerging and exciting field of research. With regards to SeIN, recent results demonstrate that this selenoprotein plays a significant role in human cell defense against oxidative stress and is implicated in redox control of several key processes in skeletal muscle cells, including EC coupling and possibly transcriptional control of antioxidant enzymes. Thus, SEP1N-RM represents the first structural muscle disorder primarily related with oxidative/nitrosative stress, and can be used as a model paradigm to study other disorders in which oxidative stress or selenoproteins are involved.

Despite these significant advances, many central questions remain to be answered. The function of most selenoproteins is unknown; even for those which have been more extensively characterized, our understanding of their multiple actions, partners, and potential roles in human pathogenesis is still incomplete. The complex mechanisms involved in modulation of expression, localization, and splicing of selenoproteins are far from fully understood. Some selenoproteins have a potential dual oxidant/antioxidant action that must be carefully investigated prior to considering therapeutic interventions. Along these lines, identification or development of compounds that could modify either the whole selenoproteome or specific selenoproteins would be a valuable tool for therapeutic studies. In the case of SeIN, two key points that need to be addressed regard its enzymatic function and its protein interactions, particularly whether RyR is its sole substrate. Contrary to RyR1, RyR3 shows a developmental expression pattern similar to that of *SEP1N* (52); although this

RyR isoform is not thought to be relevant in postnatal skeletal muscle, its potential interaction with SeIN deserves further investigation.

In the last years, the development and characterization of cellular or animal models, which allow connecting the molecular selenoprotein dysfunction and the physiological defects, have represented a significant contribution to the field. On the other hand, the differences in basal oxidant activity observed in different SeIN-devoid human cell types (fibroblasts *versus* muscle cells) stress the complexity of antioxidant defense mechanisms that show species, time, and tissue specificities. Redox homeostasis and signaling are certainly not identical in mice and men; therefore, research using human material can provide invaluable information regarding human health and disease. Most importantly, characterization of these pathways has opened avenues of investigation for the design of therapeutic approaches, of which SEP1N-RM is a good example. Further studies are in progress to determine a safe dosage of NAC, to establish its clinical efficiency in this condition, and to identify other drugs. Because in this unique condition antioxidants target the primary pathogenic defect, the planned trial with NAC would represent the first specific pharmacological treatment for a congenital myopathy and provide an important proof-of-concept regarding the efficiency of antioxidant drugs in a genetic disorder.

Acknowledgments

We thank Mrs. C. Serreri and Mrs. C. Ramahefasolo for help with writing of the article, Dr. P. Richard for diagnostic analysis of *SEP1N*, and Dr. A. Lescure for interesting discussions. The financial support of the Agence National de la Recherche (ANR), the Assistance Publique-Hôpitaux de Paris (AP-HP) and the Institut National de la Santé et la Recherche Médicale (Contrat d'Interface INSERM-Hôpital, AF) is gratefully acknowledged.

References

1. Abraham RZ, Kobzik L, Moody MR, Reid MB, and Stamler JS. Cyclic GMP is a second messenger by which nitric oxide inhibits diaphragm contraction. *Comp Biochem Physiol A Mol Integr Physiol* 119: 177–183, 1998.
2. Ajtai K and Burghardt TP. Fluorescent modification and orientation of myosin sulfhydryl 2 in skeletal muscle fibers. *Biochemistry* 28: 2204–2210, 1989.
3. Arbogast S, Beuvin M, Fraysse B, Zhou H, Muntoni F, and Ferreira A. Oxidative stress in SEP1N-related myopathy: From pathophysiology to treatment. *Ann Neurol* 65: 677–686, 2009.
4. Arthur JR. The glutathione peroxidases. *Cell Mol Life Sci* 57: 1825–1835, 2000.
5. Baylin SB, Herman JG, Graff JR, Vertino PM, and Issa JP. Alterations in DNA methylation: A fundamental aspect of neoplasia. *Adv Cancer Res* 72: 141–196, 1998.
6. Beck MA. Selenium and vitamin E status: Impact on viral pathogenicity. *J Nutr* 137: 1338–1340, 2007.
7. Beck MA, Esworthy RS, Ho YS, and Chu FF. Glutathione peroxidase protects mice from viral-induced myocarditis. *FASEB J* 12: 1143–1149, 1998.
8. Beckett GJ and Arthur JR. Selenium and endocrine systems. *J Endocrinol* 184: 455–465, 2005.
9. Behne D and Kyriakopoulos A. Mammalian selenium-containing proteins. *Annu Rev Nutr* 21: 453–473, 2001.

10. Beilstein MA, Vendeland SC, Barofsky E, Jensen ON, and Whanger PD. Selenoprotein W of rat muscle binds glutathione and an unknown small molecular weight moiety. *J Inorg Biochem* 61: 117–124, 1996.
11. Bellingier FP, Raman AV, Reeves MA, and Berry MJ. Regulation and function of selenoproteins in human disease. *Biochem J* 422: 11–22, 2009.
12. Ben Jilani KE, Panee J, He Q, Berry MJ, and Li PA. Overexpression of selenoprotein H reduces Ht22 neuronal cell death after UVB irradiation by preventing superoxide formation. *Int J Biol Sci* 3: 198–204, 2007.
13. Bjornstedt M, Hamberg M, Kumar S, Xue J, and Holmgren A. Human thioredoxin reductase directly reduces lipid hydroperoxides by NADPH and selenocystine strongly stimulates the reaction via catalytically generated selenols. *J Biol Chem* 270: 11761–11764, 1995.
14. Boldery R, Fielding G, Rafter T, Pascoe AL, and Scalia GM. Nutritional deficiency of selenium secondary to weight loss (bariatric) surgery associated with life-threatening cardiomyopathy. *Heart Lung Circ* 16: 123–126, 2007.
15. Bosl MR, Takaku K, Oshima M, Nishimura S, and Taketo MM. Early embryonic lethality caused by targeted disruption of the mouse selenocysteine tRNA gene (Trsp). *Proc Natl Acad Sci USA* 94: 5531–5534, 1997.
16. Bosschaerts T, Williams M, Noel W, Herin M, Burk RF, Hill KE, Brys L, Raes G, Ghassabeh GH, De Baetselier P, and Beschin A. Alternatively activated myeloid cells limit pathogenicity associated with African trypanosomiasis through the IL-10 inducible gene selenoprotein P. *J Immunol* 180: 6168–6175, 2008.
17. Brigelius-Flohe R. Tissue-specific functions of individual glutathione peroxidases. *Free Radic Biol Med* 27: 951–965, 1999.
18. Brooke MH and Kaiser KK. Three “myosin adenosine triphosphatase” systems: The nature of their pH lability and sulfhydryl dependence. *J Histochem Cytochem* 18: 670–672, 1970.
19. Brown MR, Cohen HJ, Lyons JM, Curtis TW, Thunberg B, Cochran WJ, and Klish WJ. Proximal muscle weakness and selenium deficiency associated with long term parenteral nutrition. *Am J Clin Nutr* 43: 549–554, 1986.
20. Burk RF and Hill KE. Selenoprotein P: An extracellular protein with unique physical characteristics and a role in selenium homeostasis. *Annu Rev Nutr* 25: 215–235, 2005.
21. Burk RF, Hill KE, Awad JA, Morrow JD, Kato T, Cockell KA, and Lyons PR. Pathogenesis of diquat-induced liver necrosis in selenium-deficient rats: assessment of the roles of lipid peroxidation and selenoprotein P. *Hepatology* 21: 561–569, 1995.
22. Cabreiro F, Picot CR, Friguet B, and Petropoulos I. Methionine sulfoxide reductases: Relevance to aging and protection against oxidative stress. *Ann NY Acad Sci* 1067: 37–44, 2006.
23. Cartharius K, Frech K, Grote K, Klocke B, Haltmeier M, Klingenhoff A, Frisch M, Bayerlein M, and Werner T. MatInspector and beyond: Promoter analysis based on transcription factor binding sites. *Bioinformatics* 21: 2933–2942, 2005.
24. Castets P, Maugey S, Gartioux C, Rederstorff M, Krol A, Lescure A, Tajbakhsh S, Allamand V, and Guicheney P. Selenoprotein N is dynamically expressed during mouse development and detected early in muscle precursors. *BMC Dev Biol* 9: 46, 2009.
25. Chariot P and Bignani O. Skeletal muscle disorders associated with selenium deficiency in humans. *Muscle Nerve* 27: 662–668, 2003.
26. Cheng WH, Ho YS, Valentine BA, Ross DA, Combs GF, Jr., and Lei XG. Cellular glutathione peroxidase is the mediator of body selenium to protect against paraquat lethality in transgenic mice. *J Nutr* 128: 1070–1076, 1998.
27. Clarke NF, Kidson W, Quijano-Roy S, Estournet B, Ferreiro A, Guicheney P, Manson JI, Kornberg AJ, Shield LK, and North KN. SEPN1: associated with congenital fiber-type disproportion and insulin resistance. *Ann Neurology* 59: 546–552, 2006.
28. Conrad M, Jakupoglu C, Moreno SG, Lippl S, Banjac A, Schneider M, Beck H, Hatzopoulos AK, Just U, Sinowatz F, Schmahl W, Chien KR, Wurst W, Bornkamm GW, and Brielmeier M. Essential role for mitochondrial thioredoxin reductase in hematopoiesis, heart development, and heart function. *Mol Cell Biol* 24: 9414–9423, 2004.
29. Deniziak M, Thisse C, Rederstorff M, Hindelang C, Thisse B, and Lescure A. Loss of selenoprotein N function causes disruption of muscle architecture in the zebrafish embryo. *Exp Cell Res* 313: 156–167, 2007.
30. Dirksen RT and Avila G. Altered ryanodine receptor function in central core disease: Leaky or uncoupled Ca(2+) release channels? *Trends Cardiovasc Med* 12: 189–197, 2002.
31. Ducreux S, Zorzato F, Ferreiro A, Jungbluth H, Muntoni F, Monnier N, Muller CR, and Treves S. Functional properties of ryanodine receptors carrying three amino acid substitutions identified in patients affected by multi-minicore disease and central core disease, expressed in immortalized lymphocytes. *Biochem J* 395: 259–266, 2006.
32. Esposito LA, Kokoszka JE, Waymire KG, Cottrell B, MacGregor GR, and Wallace DC. Mitochondrial oxidative stress in mice lacking the glutathione peroxidase-1 gene. *Free Radic Biol Med* 28: 754–766, 2000.
33. Eu JP, Sun J, Xu L, Stamler JS, and Meissner G. The skeletal muscle calcium release channel: Coupled O₂ sensor and NO signaling functions. *Cell* 102: 499–509, 2000.
34. Ferguson AD, Labunskyy VM, Fomenko DE, Arac D, Chelliah Y, Amezcua CA, Rizo J, Gladyshev VN, and Deisenhofer J. NMR structures of the selenoproteins Sep15 and SelM reveal redox activity of a new thioredoxin-like family. *J Biol Chem* 281: 3536–3543, 2006.
35. Ferreiro A, Ceuterick-de Groote C, Marks JJ, Goemans N, Schreiber G, Hanefeld F, Fardeau M, Martin JJ, Goebel HH, Richard P, Guicheney P, and Bonnemann CG. Desmin-related myopathy with Mallory body-like inclusions is caused by mutations of the selenoprotein N gene. *Ann Neurol* 55: 676–686, 2004.
36. Ferreiro A, Monnier N, Romero NB, Leroy J-P, Bönemann C, Straub V, Haenggeli C-A, Voss WD, Nivoche Y, Jungbluth H, Lemainque A, Voit T, Lunardi J, Fardeau M, and Guicheney P. A recessive form of central core disease, transiently presenting as multi-minicore disease, is associated with a homozygous mutation in the ryanodine receptor type 1 gene. *Ann Neurol* 51: 750–759, 2002.
37. Ferreiro A, Quijano-Roy S, Pichereau C, Moghadaszadeh B, Goemans N, Bonnemann C, Jungbluth H, Straub V, Villanova M, Leroy JP, Romero NB, Martin JJ, Muntoni F, Voit T, Estournet B, Richard P, Fardeau M, and Guicheney P. Mutations of the selenoprotein N gene, which is implicated in rigid spine muscular dystrophy, cause the classical phenotype of multimicore disease: Reassessing the nosology of early-onset myopathies. *Am J Hum Genet* 71: 739–749, 2002.

38. Flanigan KM, Kerr L, Bromberg MB, Leonard C, Tsuruda J, Zhang P, Gonzalez-Gomez I, Cohn R, Campbell KP, and Leppert M. Congenital muscular dystrophy with rigid spine syndrome: A clinical, pathological, radiological, and genetic study. *Ann Neurol* 47: 152–161, 2000.
39. Fomenko DE, Novoselov SV, Natarajan SK, Lee BC, Koc A, Carlson BA, Lee TH, Kim HY, Hatfield DL, and Gladyshev VN. MsrB1 (methionine-R-sulfoxide reductase 1) knock-out mice: Roles of MsrB1 in redox regulation and identification of a novel selenoprotein form. *J Biol Chem* 284: 5986–5993, 2009.
40. Fu Y, Cheng WH, Porres JM, Ross DA, and Lei XG. Knockout of cellular glutathione peroxidase gene renders mice susceptible to diquat-induced oxidative stress. *Free Radic Biol Med* 27: 605–611, 1999.
41. Fu Y, Sies H, and Lei XG. Opposite roles of selenium-dependent glutathione peroxidase-1 in superoxide generator diquat- and peroxynitrite-induced apoptosis and signaling. *J Biol Chem* 276: 43004–43009, 2001.
42. Gao Y, Feng HC, Walder K, Bolton K, Sunderland T, Bishara N, Quick M, Kantham L, and Collier GR. Regulation of the selenoprotein SelS by glucose deprivation and endoplasmic reticulum stress—SelS is a novel glucose-regulated protein. *FEBS Lett* 563: 185–190, 2004.
43. Gladyshev VN and Hatfield DL. Selenocysteine-containing proteins in mammals. *J Biomed Sci* 6: 151–160, 1999.
44. Gladyshev VN, Jeang KT, and Stadtman TC. Selenocysteine, identified as the penultimate C-terminal residue in human T-cell thioredoxin reductase, corresponds to TGA in the human placental gene. *Proc Natl Acad Sci USA* 93: 6146–6151, 1996.
45. Goldstein BJ, Mahadev K, Wu X, Zhu L, and Motoshima H. Role of insulin-induced reactive oxygen species in the insulin signaling pathway. *Antioxid Redox Signal* 7: 1021–1031, 2005.
46. Higaki Y, Hirshman MF, Fujii N, and Goodyear LJ. Nitric oxide increases glucose uptake through a mechanism that is distinct from the insulin and contraction pathways in rat skeletal muscle. *Diabetes* 50: 241–247, 2001.
47. Hinshaw DB, Burger JM, Beals TF, Armstrong BC, and Hyslop PA. Actin polymerization in cellular oxidant injury. *Arch Biochem Biophys* 288: 311–316, 1991.
48. Ho YS, Magnenat JL, Bronson RT, Cao J, Gargano M, Sugawara M, and Funk CD. Mice deficient in cellular glutathione peroxidase develop normally and show no increased sensitivity to hyperoxia. *J Biol Chem* 272: 16644–16651, 1997.
49. Imai H, Hirao F, Sakamoto T, Sekine K, Mizukura Y, Saito M, Kitamoto T, Hayasaka M, Hanaoka K, and Nakagawa Y. Early embryonic lethality caused by targeted disruption of the mouse PHGPx gene. *Biochem Biophys Res Commun* 305: 278–286, 2003.
50. Jakupoglu C, Przemeczek GK, Schneider M, Moreno SG, Mayr N, Hatzopoulos AK, de Angelis MH, Wurst W, Bornkamm GW, Brielmeier M, and Conrad M. Cytoplasmic thioredoxin reductase is essential for embryogenesis but dispensable for cardiac development. *Mol Cell Biol* 25: 1980–1988, 2005.
51. Jeong D, Kim TS, Chung YW, Lee BJ, and Kim IY. Selenoprotein W is a glutathione-dependent antioxidant *in vivo*. *FEBS Lett* 517: 225–228, 2002.
52. Jurynek MJ, Xia R, Mackrill JJ, Gunther D, Crawford T, Flanigan KM, Abramson JJ, Howard MT, and Grunwald DJ. Selenoprotein N is required for ryanodine receptor calcium release channel activity in human and zebrafish muscle. *Proc Natl Acad Sci USA* 105: 12485–12490, 2008.
53. Kabuyama Y, Oshima K, Kitamura T, Homma M, Yamaki J, Munakata M, and Homma Y. Involvement of selenoprotein P in the regulation of redox balance and myofibroblast viability in idiopathic pulmonary fibrosis. *Genes Cells* 12: 1235–1244, 2007.
54. Kim HY and Gladyshev VN. Methionine sulfoxide reductases: Selenoprotein forms and roles in antioxidant protein repair in mammals. *Biochem J* 407: 321–329, 2007.
55. Klee C and Means A. Keeping up with calcium: Conference on calcium-binding proteins and calcium function in health and disease. *EMBO Rep* 3: 823–827, 2002.
56. Kobzik L, Stringer B, Balligand JL, Reid MB, and Stamler JS. Endothelial type nitric oxide synthase in skeletal muscle fibers: Mitochondrial relationships. *Biochem Biophys Res Commun* 211: 375–381, 1995.
57. Kyriakopoulos A, Bukalis K, Roethlein D, Hoppe B, Graebert A, and Behne D. Prevention against oxidative stress of eukaryotic cell membranes by selenium compounds of the rat. *Ann NY Acad Sci* 1030: 458–461, 2004.
58. Lassing I, Schmitzberger F, Bjornstedt M, Holmgren A, Nordlund P, Schutt CE, and Lindberg U. Molecular and structural basis for redox regulation of beta-actin. *J Mol Biol* 370: 331–348, 2007.
59. Lei C, Niu X, Wei J, Zhu J, and Zhu Y. Interaction of glutathione peroxidase-1 and selenium in endemic dilated cardiomyopathy. *Clin Chim Acta* 399: 102–108, 2009.
60. Lei XG. Glutathione peroxidase-1 gene knockout on body antioxidant defense in mice. *Biofactors* 14: 93–99, 2001.
61. Lescure A, Gautheret D, Carbon P, and Krol A. Novel selenoproteins identified *in silico* and *in vivo* by using a conserved RNA structural motif. *J Biol Chem* 274: 38147–38154, 1999.
62. Lescure A, Rederstorff M, Krol A, Guicheney P, and Allamand V. Selenoprotein function and muscle disease. *Biochim Biophys Acta* 1790: 1569–1574, 2009.
63. Li YP, Chen Y, Li AS, and Reid MB. Hydrogen peroxide stimulates ubiquitin-conjugating activity and expression of genes for specific E2 and E3 proteins in skeletal muscle myotubes. *Am J Physiol Cell Physiol* 285: C806–812, 2003.
64. Lippman SM, Klein EA, Goodman PJ, Lucia MS, Thompson IM, Ford LG, Parnes HL, Minasian LM, Gaziano JM, Hartline JA, Parsons JK, Bearden JD, 3rd, Crawford ED, Goodman GE, Claudio J, Winkquist E, Cook ED, Karp DD, Walther P, Lieber MM, Kristal AR, Darke AK, Arnold KB, Ganz PA, Santella RM, Albanes D, Taylor PR, Probstfield JL, Jagpal TJ, Crowley JJ, Meyskens FL, Jr., Baker LH, and Coltman CA, Jr. Effect of selenium and vitamin E on risk of prostate cancer and other cancers: The Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA* 301: 39–51, 2009.
65. Loflin J, Lopez N, Whanger PD, and Kiousi C. Selenoprotein W during development and oxidative stress. *J Inorg Biochem* 100: 1679–1684, 2006.
66. Lu C, Qiu F, Zhou H, Peng Y, Hao W, Xu J, Yuan J, Wang S, Qiang B, Xu C, and Peng X. Identification and characterization of selenoprotein K: An antioxidant in cardiomyocytes. *FEBS Lett* 580: 5189–5197, 2006.
67. Maiorino M, Aumann KD, Brigelius-Flohe R, Doria D, van den Heuvel J, McCarthy J, Roveri A, Ursini F, and Flohe L. Probing the presumed catalytic triad of selenium-containing peroxidases by mutational analysis of phospholipid hydroperoxide glutathione peroxidase (PHGPx). *Biol Chem Hoppe Seyler* 376: 651–660, 1995.

68. Maiti B, Arbogast S, Allamand V, Moyle MW, Anderson CB, Richard P, Guicheney P, Ferreiro A, Flanigan KM, and Howard MT. A mutation in the SEPN1 selenocysteine redefinition element (SRE) reduces selenocysteine incorporation and leads to SEPN1-related myopathy. *Hum Mutat* 30: 411–416, 2009.
69. Mant TG, Tempowski JH, Volans GN, and Talbot JC. Adverse reactions to acetylcysteine and effects of overdose. *Br Med J (Clin Res Ed)* 289: 217–219, 1984.
70. Marchetti MA, Pizarro GO, Sagher D, Deamicis C, Brot N, Hejtmancik JF, Weissbach H, and Kantorow M. Methionine sulfoxide reductases B1, B2, and B3 are present in the human lens and confer oxidative stress resistance to lens cells. *Invest Ophthalmol Vis Sci* 46: 2107–2112, 2005.
71. Meissner G. Regulation of mammalian ryanodine receptors. *Front Biosci* 7: 2072–2080, 2002.
72. Mirault ME, Tremblay A, Beaudoin N, and Tremblay M. Overexpression of seleno-glutathione peroxidase by gene transfer enhances the resistance of T47D human breast cells to clastogenic oxidants. *J Biol Chem* 266: 20752–20760, 1991.
73. Moghadaszadeh B, Aracena-Parks P, Ronan M, Gasmí H, Agrawal PB, Hamilton S, and Beggs A. SEPN1-related myopathy: a defect in redox regulation. In *12th International Congress of the World Muscle Society*, Giardini Naxos-Taormina, Italy, 17–20 October 2007. *Neuromusc Disord* 17: 899, 2007.
74. Moghadaszadeh B, Desguerre I, Topaloglu H, Muntoni F, Pavsek S, Sewry C, Mayer M, Fardeau M, Tomé FMS, and Guicheney P. Identification of a new locus for a peculiar form of congenital muscular dystrophy with early rigidity of the spine on chromosome 1p35-36. *Am J Hum Genet* 62: 1439–1445, 1998.
75. Moghadaszadeh B, Petit N, Jaillard C, Brockington M, Quijano Roy S, Merlini L, Romero N, Estournet B, Desguerre I, Chaigne D, Muntoni F, Topaloglu H, and Guicheney P. Mutations in SEPN1 cause congenital muscular dystrophy with spinal rigidity and restrictive respiratory syndrome. *Nat Genet* 29: 17–18, 2001.
76. Moghadaszadeh B, Topaloglu H, Merlini L, Muntoni F, Estournet B, Sewry C, Naom I, Barois A, Fardeau M, Tome FM, and Guicheney P. Genetic heterogeneity of congenital muscular dystrophy with rigid spine syndrome. *Neuromuscul Disord* 9: 376–382, 1999.
77. Moskovitz J. Prolonged selenium-deficient diet in MsrA knockout mice causes enhanced oxidative modification to proteins and affects the levels of antioxidant enzymes in a tissue-specific manner. *Free Radic Res* 41: 162–171, 2007.
78. Motsenbocker MA and Tappel AL. A selenocysteine-containing selenium-transport protein in rat plasma. *Biochim Biophys Acta* 719: 147–153, 1982.
79. Nauser T, Dockheer S, Kissner R, and Koppenol WH. Catalysis of electron transfer by selenocysteine. *Biochemistry* 45: 6038–6043, 2006.
80. Nethery D, DiMarco A, Stofan D, and Supinski G. Sepsis increases contraction-related generation of reactive oxygen species in the diaphragm. *J Appl Physiol* 87: 1279–1286, 1999.
81. Papp LV, Lu J, Holmgren A, and Khanna KK. From selenium to selenoproteins: Synthesis, identity, and their role in human health. *Antioxid Redox Signal* 9: 775–806, 2007.
82. Perkins WJ, Han YS, and Sieck GC. Skeletal muscle force and actomyosin ATPase activity reduced by nitric oxide donor. *J Appl Physiol* 83: 1326–1332, 1997.
83. Petit N, Lescure A, Rederstorff M, Krol A, Moghadaszadeh B, Wewer UM, and Guicheney P. Selenoprotein N: An endoplasmic reticulum glycoprotein with an early developmental expression pattern. *Hum Mol Genet* 12: 1045–1053, 2003.
84. Powers SK and Hamilton K. Antioxidants and exercise. *Clin Sports Med* 18: 525–536, 1999.
85. Putkey JA, Dotson DG, and Mouawad P. Formation of inter- and intramolecular disulfide bonds can activate cardiac troponin C. *J Biol Chem* 268: 6827–6830, 1993.
86. Ran Q, Liang H, Gu M, Qi W, Walter CA, Roberts LJ, 2nd, Herman B, Richardson A, and Van Remmen H. Transgenic mice overexpressing glutathione peroxidase 4 are protected against oxidative stress-induced apoptosis. *J Biol Chem* 279: 55137–55146, 2004.
87. Rederstorff M, Allamand V, Guicheney P, Gartioux C, Richard P, Chaigne D, Krol A, and Lescure A. Ex vivo correction of selenoprotein N deficiency in rigid spine muscular dystrophy caused by a mutation in the selenocysteine codon. *Nucleic Acids Res* 36: 237–244, 2008.
88. Reid MB and Durham WJ. Generation of reactive oxygen and nitrogen species in contracting skeletal muscle: Potential impact on aging. *Ann NY Acad Sci* 959: 108–116, 2002.
89. Reid MB, Khawli FA, and Moody MR. Reactive oxygen in skeletal muscle. III. Contractility of unfatigued muscle. *J Appl Physiol* 75: 1081–1087, 1993.
90. Reid MB, Stokic DS, Koch SM, Khawli FA, and Leis AA. N-acetylcysteine inhibits muscle fatigue in humans. *J Clin Invest* 94: 2468–2474, 1994.
91. Rubartelli A, Bajetto A, Allavena G, Wollman E, and Sita R. Secretion of thioredoxin by normal and neoplastic cells through a leaderless secretory pathway. *J Biol Chem* 267: 24161–24164, 1992.
92. Saito Y, Sato N, Hirashima M, Takebe G, Nagasawa S, and Takahashi K. Domain structure of bi-functional selenoprotein P. *Biochem J* 381: 841–846, 2004.
93. Seiler A, Schneider M, Forster H, Roth S, Wirth EK, Culmsee C, Plesnila N, Kremmer E, Radmark O, Wurst W, Bornkamm GW, Schweizer U, and Conrad M. Glutathione peroxidase 4 senses and translates oxidative stress into 12/15-lipoxygenase dependent- and AIF-mediated cell death. *Cell Metab* 8: 237–248, 2008.
94. Soderberg A, Sahaf B, and Rosen A. Thioredoxin reductase, a redox-active selenoprotein, is secreted by normal and neoplastic cells: Presence in human plasma. *Cancer Res* 60: 2281–2289, 2000.
95. Speckmann B, Walter PL, Alili L, Reinehr R, Sies H, Klotz LO, and Steinbrenner H. Selenoprotein P expression is controlled through interaction of the coactivator PGC-1 α with FoxO1 α and hepatocyte nuclear factor 4 α transcription factors. *Hepatology* 48: 1998–2006, 2008.
96. Stadtman ER. Protein oxidation and aging. *Free Radic Res* 40: 1250–1258, 2006.
97. Steinbrenner H, Alili L, Bilgic E, Sies H, and Brenneisen P. Involvement of selenoprotein P in protection of human astrocytes from oxidative damage. *Free Radic Biol Med* 40: 1513–1523, 2006.
98. Steinbrenner H and Sies H. Protection against reactive oxygen species by selenoproteins. *Biochim Biophys Acta* 1790: 1478–1485, 2009.
99. Steinmann D, Nauser T, Beld J, Tanner M, Gunther D, Bounds PL, and Koppenol WH. Kinetics of tyrosyl radical reduction by selenocysteine. *Biochemistry* 47: 9602–9607, 2008.

100. Stoytcheva ZR and Berry MJ. Transcriptional regulation of mammalian selenoprotein expression. *Biochim Biophys Acta* 1790: 1429–1440, 2009.
101. Supinski GS, Stofan D, Ciuffo R, and DiMarco A. N-acetylcysteine administration and loaded breathing. *J Appl Physiol* 79: 340–347, 1995.
102. Sweeney HL, Bowman BF, and Stull JT. Myosin light chain phosphorylation in vertebrate striated muscle: Regulation and function. *Am J Physiol* 264: C1085–1095, 1993.
103. Tajsharghi H, Darin N, Tulinius M, and Oldfors A. Early onset myopathy with a novel mutation in the selenoprotein N gene (SEPN1). *Neuromuscul Disord* 15: 299–302, 2005.
104. Tamura T and Stadtman TC. Mammalian thioredoxin reductases. *Methods Enzymol* 347: 297–306, 2002.
105. Thisse C, Degraeve A, Kryukov GV, Gladyshev VN, Obrecht-Pflumio S, Krol A, Thisse B, and Lescure A. Spatial and temporal expression patterns of selenoprotein genes during embryogenesis in zebrafish. *Gene Expr Patterns* 3: 525–532, 2003.
106. Trigona WL, Mullarky IK, Cao Y, and Sordillo LM. Thioredoxin reductase regulates the induction of haem oxygenase-1 expression in aortic endothelial cells. *Biochem J* 394: 207–216, 2006.
107. Urso ML and Clarkson PM. Oxidative stress, exercise, and antioxidant supplementation. *Toxicology* 189: 41–54, 2003.
108. Vendeland SC, Beilstein MA, Chen CL, Jensen ON, Barofsky E, and Whanger PD. Purification and properties of selenoprotein W from rat muscle. *J Biol Chem* 268: 17103–17107, 1993.
109. Williams DL, Jr. and Swenson CA. Disulfide bridges in tropomyosin. Effect on ATPase activity of actomyosin. *Eur J Biochem* 127: 495–499, 1982.
110. Xia L, Nordman T, Olsson JM, Damdimopoulos A, Bjorkhem-Bergman L, Nalvarte I, Eriksson LC, Arner ES, Spyrou G, and Bjornstedt M. The mammalian cytosolic selenoenzyme thioredoxin reductase reduces ubiquinone. A novel mechanism for defense against oxidative stress. *J Biol Chem* 278: 2141–2146, 2003.
111. Yagi K, Komura S, Kojima H, Sun Q, Nagata N, Ohishi N, and Nishikimi M. Expression of human phospholipid hydroperoxide glutathione peroxidase gene for protection of host cells from lipid hydroperoxide-mediated injury. *Biochem Biophys Res Commun* 219: 486–491, 1996.
112. Yant LJ, Ran Q, Rao L, Van Remmen H, Shibata T, Belter JG, Motta L, Richardson A, and Prolla TA. The selenoprotein GPX4 is essential for mouse development and protects from radiation and oxidative damage insults. *Free Radic Biol Med* 34: 496–502, 2003.
113. Yoshida H, Haze K, Yanagi H, Yura T, and Mori K. Identification of the cis-acting endoplasmic reticulum stress response element responsible for transcriptional induction of mammalian glucose-regulated proteins. Involvement of basic leucine zipper transcription factors. *J Biol Chem* 273: 33741–33749, 1998.

Address correspondence to:

Dr. Ana Ferreiro

INSERM U787

GH Pitié-Salpêtrière

105 Bd. de l'Hôpital

75651 Paris

France

E-mail: ana.b.ferreiro@gmail.com

Abbreviations Used

Cat = catalase
 Cu/Zn SOD = copper/zinc superoxide dismutase
 DIOs = iodothyronine deiodinases
 ecSOD = extracellular superoxide dismutase
 ERSE = endoplasmic reticulum response element
 GPxs = glutathione peroxidases
 Mn SOD = manganese superoxide dismutase
 mt NOS = mitochondrial nitric oxide synthase
 nNOS = neuronal nitric oxide synthase
 NO = nitric oxide
 PLA = phospholipase A2
 ROS = reactive oxygen species
 SECIS = selenocysteine insertion sequence
 SEPN1-RM = SEPN1-related myopathy
 SOD = superoxide dismutase
 SR = sarcoplasmic reticulum
 SRE = Sec codon redefinition element
 TrxRs = thioredoxin reductases
 XO = xanthine oxidase