

# Differentiated Neogene bauxitization of volcanic rocks (Western Cameroon): Morpho-geological constraints on chemical erosion

Mathieu Nouazi Momo, Anicet Beauvais, Paul Tematio, Martin Yemefack

# ► To cite this version:

Mathieu Nouazi Momo, Anicet Beauvais, Paul Tematio, Martin Yemefack. Differentiated Neogene bauxitization of volcanic rocks (Western Cameroon): Morpho-geological constraints on chemical erosion. CATENA, 2020, 194 (104685), pp.1-13. 10.1016/j.catena.2020.104685. hal-02617017

# HAL Id: hal-02617017 https://hal.science/hal-02617017

Submitted on 25 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

| 1  | Differentiated Neogene bauxitization of volcanic rocks (Western   |
|----|---|
| 2  | Cameroon): Morpho-geological constraints on chemical erosion  |
| 3  |   |
| 4  | Mathieu Nouazi Momo <sup>1</sup> *, Anicet Beauvais <sup>2</sup> , Paul Tematio <sup>3</sup> , Martin Yemefack <sup>4</sup> |
| 5  | <sup>1</sup> Institute of Geological and Mining Research (IRGM), P.O. Box 4110 Yaoundé, Cameroon                            |
| 6  | <sup>2</sup> Aix-Marseille Univ, CNRS, IRD, INRAE, Coll France, CEREGE, BP 80, 13545, Aix-en-                               |
| 7  | Provence, Cedex 4, France   |
| 8  | <sup>3</sup> University of Dschang, Faculty of Science, Department of Earth Sciences. P.O. Box 67                           |
| 9  | Dschang - Cameroon  |
| 10 | <sup>4</sup> International Institute of Tropical Agriculture (IITA), ASB Partnership REALU Project,                         |
| 11 | P.O. Box 2008 (Messa) Yaoundé, Cameroon   |
| 12 | *Corresponding author Email: nouazimat@yahoo.fr   |
| 13 |   |
| 14 |   |
| 15 |   |
| 16 | Catena  |
| 17 | Revised version submitted May 13, 2020  |
| 18 | Accepted May 16, 2020   |
| 19 |   |
| 20 | DOI: https://doi.org/10.1016/j.catena.2020.104685   |
| 21 |   |
| 22 |   |
| 23 |   |
| 24 |   |
| 25 |   |

26 Abstract

Lateritic weathering of Miocene volcanic rocks from western Cameroon highlands formed 27 duricrusted bauxitic profiles. Two weathering profiles on ca. 14 Ma basalt and ca. 16 Ma 28 29 trachyte were studied using geochemical mass balance functions. Less mobile elements Ti and Zr were used as references to quantify volumetric change (strain,  $\varepsilon$ ), element transfer 30 rate  $(\tau)$  and geochemical mass transfers during the bauxitization process of basalt and 31 trachyte. Conversion of parent rocks to kaolinite and goethite rich saprolites evolved to Al-32 Fe rich bauxites, mostly composed of gibbsite and iron oxy-hydroxides (goethite and 33 hematite). However, formation of Al-Fe bauxitic profiles required higher Si leaching on 34 trachyte than on basalt. Our results document that chemical weathering of a larger thickness 35 of trachyte than basalt has been required to form a unit meter of weathering profile, 36 implying differential rates of rock chemical erosion and topographic decay of landscapes. 37 Rates of chemical erosion and formation of lateritic weathering profiles in western 38 Cameroon have been mostly controlled by drainage conditions and volcanic rocks 39 40 composition (mostly SiO<sub>2</sub> content differences), that also resulted in contrasted landscapes evolution during the Neogene. 41 42 Keywords: bauxite; lateritic weathering; geochemical mass balance; chemical erosion; 43 volcanic rocks; Cameroon 44

45

46 <u>Highlights</u>:

47 Differential bauxitization of trachyte and basalt in western Cameroon

48 Mass transfers document differences in chemical weathering and landscapes evolution

49 Drainage and rocks composition control chemical erosion and bauxitic weathering

50

### 1. Introduction

Lateritic regoliths of tropical shields generally have formed by long-term rock 52 weathering under warm / humid and seasonally contrasted climates (Tardy, 1997). In 53 tropical landscapes, lateritic paleosurfaces bearing bauxites and ferricretes have been 54 relatively well preserved over geological times scale (Beauvais and Chardon, 2013). 55 Formation of bauxite or ferricrete depends on climatic and morpho-tectonic conditions that 56 control solute leaching processes and aluminum or iron accumulations in lateritic 57 weathering profiles formed on erosional landforms (Bardossy and Aleva, 1990; Thomas, 58 1994; Valeton, 1999), including duricrusted Fe-cemented pediments (Chardon et al., 2018). 59 In Africa, bauxites have formed by intense lateritic weathering of various lithologies 60 61 (Boulangé and Millot, 1988) mostly from ca. 59 to ca. 45 Ma (Beauvais et al., 2008) upon the African surface that extends from West to East and South-East Africa (Beauvais and 62 Chardon, 2013; Burke and Gunnell, 2008; see also Pérello et al., 2020; De Putter and 63 Ruffet, 2020). Likewise, thick bauxitic profiles later formed from weathering of 14 Ma to 64 23 Ma old trachyte and basalt are well preserved at altitudes of 1500 to 1800 m in the 65 eastern Nigeria (Schwarz, 1997). With the mid-Miocene climatic optimum, Lateritic 66 weathering intensified in Africa (e.g., Beauvais et al., 2008; De Putter and Ruffet, 2020), 67 and more generally on continents of the inter-tropical belt (Retallack, 2010) even at 68 relatively high altitudes, e.g., upon elevated continental margins (Jean et al., 2020), where 69 transported-limited erosion regimes have potentially controlled landscapes evolution over 70 71 the Cenozoic.

Previous studies established the geochemical variability of bauxitic profiles with regard
to underlying parent rocks, as well as the link with local morphoclimatic conditions that
prevailed during lateritic weathering, as exemplified in West Africa (Boulangé, 1984;
Valeton and Beiβner, 1986), Asia (Zarasvandi et al., 2012), Eastern Australia (Eggleton et

al., 1987), Southern America (Boulangé and Colin, 1994; Soler and Lasaga, 2000; Soares
de Oliveira et al., 2011), and India (Babechuk et al., 2014). Chemical weathering of basalt
and volcanic rocks has been recently considered into the frame of major surface processes
that govern the Critical Zone (Brantley and Lebedeva, 2011; Babechuk et al., 2015; Wille et
al., 2018).

In situ formed bauxitic profiles are suitable to quantify geochemical mass transfers 81 during long-term rock weathering. Since the pioneering introduction from Millot and 82 Bonifas (1959), and mathematical formalization by Brimhall and Dietrich (1987) and 83 Brimhall et al. (1988), the geochemical mass balance approach was commonly used for 84 quantifying the accumulation or depletion of major and trace elements during weathering 85 86 processes (e.g., Colin et al., 1992; Beauvais and Colin, 1993; Braun et al., 1993; Tematio and Olson, 1997; Temgoua et al., 2003, among others). However, few studies have 87 addressed elemental geochemical mass balance of lateritic bauxitic profiles (e.g., Boulangé 88 and Colin, 1994; Eggleton et al., 2008). In western Cameroon region and on the Adamaoua 89 plateau above the Central Cameroonian shear zone (CCSZ, Fig.1), bauxitic regolith formed 90 on high altitude surfaces (up to ~1900 m) from lateritic weathering of mid- to lower 91 Miocene volcanic rocks of the Cameroon volcano-tectonic line. Here, we have quantified 92 element mass transfers (gains and losses of major chemical elements) in lateritic weathering 93 profiles formed upon mid-Miocene basalt and trachyte from the western highlands of 94 Cameroon. Our results document differential morpho-geological controls on the rate of 95 rock chemical erosion, formation of bauxitic regolith and landscapes evolution in western 96 Cameroon over the Neogene. 97

# 2. Geological and morphological setting

| 100 | In the western highlands of Cameroon, wide Cenozoic volcanic complexes (e.g.                   |
|-----|--|
| 101 | Mounts Bambouto and Bamenda; Fig. 1) including mafic and felsic plutons have been              |
| 102 | extruded through the Neoproterozoic granito-gneissic basement (Kwekam et al., 2010).           |
| 103 | Tertiary to recent strato-volcanoes culminate up to 3000 m a.s.l., above low-relief lateritic  |
| 104 | landsurfaces and grabens (Deruelle et al., 2007; Nkouathio et al., 2008), i.e., Mbo (ca 700    |
| 105 | m a.s.l.), Noun (ca 1100 m a.s.l.) and Tikar (ca 750 m a.s.l.) plains (Fig. 1b).               |
| 106 | Volcanic activity has begun in Cameroon from the Eocene with the early establishment           |
| 107 | of the Bamoun plateau between 51.8 and 46.7 Ma (Moundi et al., 2007) and Mount Bangou          |
| 108 | between 44.7 and 43.1 Ma (Fosso et al., 2005). Peaks of magmatic activity also generated       |
| 109 | (1) Oligo-Miocene basaltic flows (31 to 14 Ma) of little differentiated silicic magmas from    |
| 110 | an asthenosphere source, and (2) most evolved Mio-Quaternary (15 Ma to 4 Ma) silicic           |
| 111 | magmas from amphibole bearing lithospheric mantle source (Marzoli et al., 1999; Marzoli        |
| 112 | et al., 2000).   |
| 113 | Between longitudes 09°50" and 11°00" and latitudes 05°10" and 06° 40" (Fig. 1b),               |
| 114 | volcanic rocks have been intensively weathered since mid- Miocene. Studied lateritic           |
| 115 | weathering profiles are mostly bauxitic and formed upon trachyte of ca. 16 Ma at Fongo         |
| 116 | Tongo and on basalt of ca. 14 Ma at Doumbouo (Nkouathio et al., 2008) in the                   |
| 117 | southernmost part of the western Cameroon highlands (Fig. 1b and Fig. 2). Sparse pre-          |
| 118 | Neogene lateritic profiles topped by ferricretes also overly gneissic basement.                |
| 119 | On trachyte around Fongo Tongo (Figs. 2a-b), a low-relief lateritic landscape with             |
| 120 | smooth hilly landforms at altitudes ranging from ca. 1570 m to ca. 1650 m is drained by a      |
| 121 | first order dense alveolar drainage network ca. hundred meters meshed, and bounded by          |
| 122 | incisions of ca. 150 m to ca. 200 m deep from West to East (Fig. 2a-b and supplementary        |
| 123 | Fig. DR1a). On basalt near Doumbouo (Fig. 2b), relatively flat lateritic plateaus at altitudes |

between ca. 1545 m to ca. 1585 m are drained by a first order linear drainage network and
incised from ca. 100 to 120 m deep (Fig. 2b and supplementary Fig. DR1b). High altitude
grasslands cover the two landscapes that are both subject to sub-equatorial climate (Kengni
et al., 2009) with ~1600 mm mean annual rainfall and 18.1°C of mean annual temperature.

- 128
- 129

## **3. Materials and Methods**

Two pits were dug through thick lateritic mantles representative of the two landscapes 130 (Fig. 2), on the basis of a previous geomorphological study in the region (Momo Nouazi et 131 al., 2016). The first profile (5° 31' 51.16''N / 9° 58' 34.99''E) formed upon trachyte at 132 133 Fongo-Tongo, while the second one (5° 25' 19.36''N / 10° 05' 46.98''E) formed upon 134 basalt at Doumbouo (Figs. 2a-b; see also Figs. DR1a-b). After detailed macroscopic descriptions along side pits walls, twenty samples ( $\sim 2$  kg) representative of ca. one cubic 135 meter in the different horizons of profiles were collected based on their petrographic 136 structures. Fresh trachyte was sampled in nearby guarry excavation, and basalt under the 137 saprolite of its weathering profile. 138

Petrographic studies were carried out on thin sections of fresh rocks and weathered 139 samples to describe textures of weathering products and their mineral composition. 140 141 Samples were crushed to a sieve size  $< 63 \mu m$  for mineralogical and geochemical analysis. Mineralogical analyses were performed using XRD (X-Ray Diffractometry), with an X'Pert 142 Panalytical diffractometer using CuKα radiation (1.79 Å) in the range of 2° to 78° 2θ and at 143 a speed of  $0.2^{\circ}$ /min (10s counting time every  $0.033^{\circ}$ ). Minerals were identified upon 144 diffraction spectra using High Score X'Pert Plus software (Version 3.0) and compared with 145 known standards (American Mineralogist Crystal Structure Database, 2016). A Rietveld 146 Refinement process (Rietveld, 1969; Ufer et al., 2008) was also applied for semi-147 quantifying the proportion (%) of each mineral. Each Mineral proportion is obtained by the 148

ratio between the surface areas of specific XRD peaks of each mineral to the total surface
of the sample XRD pattern. Geochemical analyses were obtained using ICP methods from
pulverized samples fraction. Major and trace elements were analyzed by ICP-AES
(Inductively coupled plasma-Atomic emission spectrometry).
3.1. Element mass transfer function

The element mass transfer function (Brimhall and Dietrich, 1987) was applied as it 155 accounts for the chemical composition and bulk density of weathered horizons and parent 156 157 rocks, allowing to estimate the porosity and volume change or strain ( $\epsilon$ ) across the weathering profile. This function also requires a reference element, which is generally 158 included in resistant minerals and considered as less mobile during weathering. The 159 reference elements commonly used are Zr, Ti, and Th (Brimhall and Dietrich, 1987; 160 Beauvais and Colin, 1993; Boulangé and Colin, 1994; Braun et al., 1993; White et al., 161 2001; Tematio et al., 2009). Finally this mass transfer function allows quantifying the 162 accumulation or depletion of elements during weathering. 163

164 Physical parameters of weathering horizons are first measured to establish:

 $\phi = 1 - \frac{\rho_b}{\rho_g} \tag{1}$ 

166φ is the porosity, while  $\rho_b$  and  $\rho_g$  in g.cm<sup>-3</sup> refer to bulk density and grain density of each167sample representative of a given weathering horizon. Bulk density,  $\rho_b$ , of dried samples was168estimated after weighing and coating them with molten paraffin wax, followed by their169immersion in water to measure displaced volume. Grain density,  $\rho_g$ , was measured using170water picnometer. Both densities were measured with an accuracy of ~ ± 1%.171Strain, ε, is then defined by:

$$\varepsilon_{i,w} = \left(\frac{\rho_p \times C_{i,p}}{\rho_w \times C_{i,w}}\right) - 1$$
(2)

173 Where *i* refer to the element considered as the less mobile during weathering,  $\rho$  is the bulk 174 density and C the concentration of the reference element in g/100g in weathered sample (*w*) 175 and parent rock (*p*). Measurement of strain,  $\varepsilon$ , allows quantification of the volumetric 176 change across weathering profile owing to its collapse (Brimhall and Dietrich, 1987; see 177 also Chadwick et al., 1990).

178 Then, transfer rate,  $\tau$ , (gain or loss) of a mobile element *j* either by relative or absolute 179 accumulation (including physical translocation) or solute depletion is defined from:

180

$$\tau_{j,w} = \left[\frac{\rho_{w} \times C_{j,w}}{\rho_{p} \times C_{j,p}} \times (\varepsilon_{i,w} + 1)\right] - 1$$
(3)

181 182

Quantification of the mass of element *j*,  $m_{j,w}$ , in kg/m<sup>3</sup> transferred through a unit volume (1 m<sup>3</sup>) of profile during both saprolite formation and bauxitization can also be obtained by:

 $\boldsymbol{m}_{j,w} = \left(\frac{\boldsymbol{C}_{j,p}}{100} \times \boldsymbol{\rho}_{p} \times \boldsymbol{V}_{p}\right) \times \boldsymbol{\tau}_{j,w}$ (4)

187

The total mass of element *j*,  $M_{j,w}$  in kg, transferred through a cross section of weathering profile during the chemical weathering of a given height or thickness of parent rock is derived by integrating  $\tau_{j,w}$  as a function of depth. However, knowledge of initial height *z* of parental rock (or protolith layer thickness) that has been chemically weathered to form the investigated profile of thickness  $\Delta z_w$  is required. According to Egli and Fitze (2000) this is obtained by relating  $\Delta z_w$  to  $\Delta z$  using the strain,  $\varepsilon_{i,w}$ , such as:  $\Delta z = \Delta z_w (1/\varepsilon_{i,w} + 1)$ . Therefore, the higher is  $\varepsilon_{i,w}$  the higher is the thickness of protolith layer that has been chemically weathered to form a given profile. The masses of elements  $M_{j,w}$  (gain or loss) transferred through each profile horizon during the chemical weathering of a corresponding thickness of parent rock is then integrated upon the total profile depth, *D*, as established by Egli and Fitze (2000):

$$\boldsymbol{M}_{j,\boldsymbol{w}}(\boldsymbol{Z}_{\boldsymbol{w}}) = \left(\frac{\boldsymbol{C}_{j,\boldsymbol{\rho}}}{100} \times \boldsymbol{\rho}_{\boldsymbol{\rho}}\right) \times \left(\frac{1}{\varepsilon_{j,\boldsymbol{w}}+1}\right) \times \int_{\boldsymbol{Z}_{\boldsymbol{w}}=0}^{\boldsymbol{Z}_{\boldsymbol{w}}=\boldsymbol{D}} \tau_{j,\boldsymbol{w}} \times \boldsymbol{d}\boldsymbol{Z}_{\boldsymbol{w}}$$
(5)

200

199

## **4. Results**

## 202 4.1. Lateritic weathering on trachyte

203 Trachyte outcrops on steep slopes and in talwegs, and is deeply weathered with bauxite formed on top of flat to gently sloping interfluves. Trachyte has a microlitic texture with no 204 or few porphyric minerals (Fig. 3a), and made up of an assemblage of microcrystalline 205 plagioclases (Pl), clinopyroxenes (cpx), Fe-Ti oxides, and scarce Cpx phenocrysts (up to 1 206 mm) (Figs. 3b and 3c). These minerals also include anorthoclase (An<sub>25-11</sub>Ab<sub>70-54</sub>Or<sub>22-19</sub>) and 207 alkali feldspars, i.e., sanidine to Na-sanidine (Or<sub>15-55</sub>Ab<sub>35-85</sub>An<sub>0-6</sub>), sodic Cpx (En<sub>3-4</sub>Aeg<sub>33-</sub> 208 35Fs<sub>62-65</sub>) or hedenbergite (Wo<sub>45-47</sub>En<sub>26-30</sub>Fs<sub>40-44</sub>) and titanomagnetite (94-25 mol% 209 210 ulvöspinel) (see Nkouathio et al. 2008). 211 Lateritic bauxitic profiles have been formed in-situ on trachyte, whose the structure is coarsely preserved in saprolite, but transported materials may also be reworked on 212 pediments established around the bauxitic hilly landforms at lower altitudes (See Momo 213

Nouazi et al., 2019). The weathering profile includes a saprolite (C), a blocky bauxitic

- duricrust (B5), a massive bauxitic duricrust (B3) and a loose nodular bauxitic horizon on
- top (B1) (Fig. 4a, see Momo Nouazi et al., 2019). The saprolite is porous (Tab. 1), with

217 inter-connected voids resulting form eluviation process of less cohesive clays (Fig. 5a). It is made up of a grey brown matrix of kaolinite and gibbsite (61% and 16%, respectively), and 218 it also contains goethite formed by pseudomorphosis of Px, Cpx and Fe-Ti oxides, and 219 some anatase and maghemite. The horizon B5 shows decimeter-size blocks of porous 220 reddish bauxitic duricrust mixed with centimeter size nodules embedded in a loose matrix. 221 222 Compared to saprolite, the mineralogical composition (Table DR1a, see Momo Nouazi et al., 2019) shows a total conversion of kaolinite into gibbsite (57%), and goethite 223 coating/infilling (29% to 58%) in bauxitic blocks. The massive bauxite duricrust is made up 224 of a reddish brown matrix mixing gibbsite (up to 68%) and goethite (up to 33%) and 225 preserving some residual features of parent minerals (e.g., pyroxene). It is crossed by 226 227 micrometer to millimeter voids coated with thick Al- and Fe-rich accumulations (Fig. 5b), while showing centimeter size sub-horizontal laminated iron cementations at its bottom 228 (Fig. 4a). The surface horizon (B1) is made up of a loose matrix embedding fragments of 229 bauxite duricrusts and bauxitic nodules whose petrographic structures are quite similar to 230 that of the underlying massive bauxite. The horizon B1 has more kaolinite (22%) than the 231 massive bauxite (B3) and less gibbsite and goethite (31 and 18%, respectively), with up to 232 18% of quartz of detrital origin. 233

Main geochemical trends highlight relative enrichment of Al<sub>2</sub>O<sub>3</sub>, Fe<sub>2</sub>O<sub>3</sub>, compared to 234 parental trachyte (Fig. 4a and Fig. 6), while SiO<sub>2</sub> with alkaline/alkaline-earth elements are 235 depleted (see also Table DR2a, and Momo Nouazi et al., 2019). Noted that locally soils and 236 upper parts of profiles formed also on trachyte from the same area (Tematio et al., 2009) 237 may be more siliceous but less rich in Al<sub>2</sub>O<sub>3</sub> and/or Fe<sub>2</sub>O<sub>3</sub> than our samples (Fig. 6). The 238 SiO<sub>2</sub> content decreases from the trachyte (59.5 wt.%) to its minimum content in the massive 239 bauxitic duricrust (0.73 wt.%), and increases toward the surface to 19.25 wt.% (Fig. 4a). 240 The Al<sub>2</sub>O<sub>3</sub> and Fe<sub>2</sub>O<sub>3</sub> contents increase from the trachyte (15.5 wt.% and 8.4 wt.%) to their 241

| 242 | maximum contents in the massive bauxite duricrust (51.8 wt.% and 25.9 wt.%),  |
|-----|---|
| 243 | respectively, and then decrease toward the surface (Fig. 4a). Note the increase in Fe <sub>2</sub> O <sub>3</sub> and   |
| 244 | decrease in $Al_2O_3$ at the transition between B3 and B5 horizons where ca. 1 m thick iron   |
| 245 | rich accumulations occurred (Fig. 4a). The $TiO_2$ content increases from 0.62 wt.% in the  |
| 246 | trachyte to 3.39 wt.% in the massive bauxitic duricrust, while $P_2O_5$ increases in the  |
| 247 | saprolite and then slightly decreases in the upper duricrusted bauxitic horizons (Fig. 7a and   |
| 248 | Table DR2a). Along the profile, trace elements, Nb, Zr, Ga, Cu, Ni, Co, Cr, V, As, Pb, Th,  |
| 249 | Hf, U and Ta contents are higher, while Y, Sr, Rb, contents are lower than those in trachyte  |
| 250 | (see also Momo Nouazi et al., 2019 for more details; and Tab. DR3a).  |
| 251 |   |
| 252 | 4.2. Lateritic weathering on basalt   |
| 253 | Porphyritic basalt is deeply weathered and occurs as boulders on slopes. Phenocrysts  |
| 254 | are clearly identified on field samples (Fig. 3d), and represent up to 30% of the sample  |
| 255 | (Figs. 3e and 3f). Parent minerals of this basalt also include large olivine (Fo <sub>66-87</sub> ),  |
| 256 | diopsides (Wo <sub>45-48</sub> En <sub>43-52</sub> Fs <sub>8-17</sub> ), augites (Wo <sub>38-44</sub> En <sub>43-53</sub> Fs <sub>10-23</sub> ), scarce phenocrysts and |
| 257 | microcrystalline plagioclases (An <sub>46-72</sub> ) and Fe-Ti oxydes (71-19 mol% ulvöspinel) (see  |
| 258 | Nkouathio et al. 2008).   |
| 259 | The lateritic profile is located on top of a flat interfluve limited by incisions (Fig. 2b  |
| 260 | and Fig. DR1b). From the bottom to the top, the profile includes a saprolite (C), a mottled-  |
| 261 | clays layer (B4), a massive bauxitic duricrust (B3), a nodular bauxitic duricrust (B2) and a  |
| 262 | loose nodular horizon on top (B1) (see Fig. 4b). The saprolite is more porous than on   |
| 263 | trachyte (Tab. 1), and composed of a greyish-brown kaolinite matrix (57%) dotted with   |
| 264 | residual porphyric olivines/pyroxenes. It is studded by millimeter sized dissolution voids  |
| 265 | preserving the original shapes of ol and cpx phenocrysts (Fig. 5c). It also shows dark-brown  |
| 266 | goethite (30%) and maghemite (9%). The mottled-clays have lost the basalt structure and   |

| 267 | are composed of a brown argillic frame of kaolinite (46%), goethite (24%), and minor  |
|-----|---|
| 268 | amount of maghemite (5%); the mottled clays also have centimeter size gray volumes of   |
| 269 | microcrystalline gibbsite (18% of the total horizon), and Al-Fe cutans fillings voids and   |
| 270 | cracks. The structure of the massive bauxitic duricrust is quite similar to mottled-clays, but  |
| 271 | the brown argillic plasma bears microcrystalline gibbsite. Clay-ferruginous materials refill  |
| 272 | voids and later differentiate into Fe-rich cutans (Fig. 5d). The bulk mineral composition of  |
| 273 | this bauxitic duricrust shows lesser kaolinite (4%) than mottled-clays, more gibbsite (41%)   |
| 274 | and goethite (27%) and up to 15% hematite (see also Table DR1b). Centimeter size nodules  |
| 275 | of the nodular bauxitic duricrust represent 75% of the horizon. They are composed of a  |
| 276 | brown argillic matrix of goethite and kaolinite locally preserving the basalt structure and   |
| 277 | are rimed with millimeter size dark brown cortex. The reddish inter-nodular matrix is   |
| 278 | mostly composed of goethite and microcrystalline gibbsite filling millimeter size cracks.   |
| 279 | The bulk mineral composition of the horizon shows more kaolinite (10%), gibbsite (46%)  |
| 280 | and goethite (38%), but lesser hematite (> 5%) than the massive bauxite duricrust. The  |
| 281 | surface nodular horizon (B1) is composed of a mixture of fine earth and centimeter size   |
| 282 | nodules, with an internal structure similar to that of massive and nodular duricrusts.  |
| 283 | Kaolinite content increases to 15%, while gibbsite and goethite decrease to 35 and 23%,   |
| 284 | respectively (Table DR1b). However, this horizon contains also up to 15% quartz, which is   |
| 285 | probably, as on profile of trachyte, of detrital origin.  |
| 286 | Main geochemical trends first show higher Al <sub>2</sub> O <sub>3</sub> and Fe <sub>2</sub> O <sub>3</sub> compared to basalt, while |
| 287 | $SiO_2$ is depleted from the profile (Figs. 4b and 6), with $P_2O_5$ and alkaline/alkaline-earth                                      |
| 288 | elements (see Table DR2b). The SiO <sub>2</sub> content decreases from the basalt (42.0 wt.%) to its                                  |
| 289 | minimum content in the massive bauxitic duricrust (1.77 wt.%), and increases toward the   |
| 290 | surface to 18.35 wt.% (Fig. 4b). The $Al_2O_3$ and $Fe_2O_3$ contents increase from the basalt  |

291 (13.0 and 14.25 wt.%) to maximum contents in the duricrusted horizons (36.0 and 34.2

| 292 | wt.%), respectively, and then decrease toward the surface (Fig. 4b). The $TiO_2$ content also  |
|-----|--|
| 293 | increases from 3.69 wt.% in basalt to 8.83 wt.% in the mottled clays (Fig. 7a and Table        |
| 294 | DR2b). Compared to basalt, trace elements Sc, V, Cr, Zn, Ga, Zr, Nb, Hf, Ta, Pb, Th and U      |
| 295 | have higher contents, while Ni, Cu, As, Rb, Sr, Y and Ba contents are lower along the          |
| 296 | profile (see Tab. DR3b).   |
| 297 |  |
| 298 | 4.3. Element mass transfers during weathering and bauxitization                                |
| 299 | Because samples from the upper part of profiles may integrate some reworked                    |
| 300 | materials of potentially allochthonous origin (e.g., colluvia), which include detrital quartz  |
| 301 | and anomalous silica content (Fig. 6; see also Tables DR1 and DR2) and possibly detrital       |
| 302 | heavy minerals (ex: zircon on basalt, Fig. 7a), mass balance calculations have been only       |
| 303 | applied on ten and five samples in the profiles formed in-situ on trachyte and basalt,         |
| 304 | respectively (Fig. 4).   |
| 305 |  |
| 306 | 4.3.1. Characterization of reference elements  |
| 307 | Reference elements, Zr and Ti, are characterized by their relatively low mobility during       |
| 308 | bauxitization. Their distribution across the weathering profiles compared to their amount in   |
| 309 | parent rocks allows documenting changes in chemical and physical processes in different        |
| 310 | horizons of profiles. Here, Zr and Ti show a differentially correlated distribution across the |
| 311 | weathering profiles (Figs. 7a), as previously observed by Schwarz (1997) and Tematio et        |
| 312 | al., (2009) in comparable morphogeological settings (Fig. 7b). The distributions of Zr and     |
| 313 | Ti in the different profiles are comparable (Figs. 7a-b). These two elements show parallel     |

better expressed than Zr and most stable on basalt (Figs. 7d and 7e). However, departures

314

behavior along the profiles (Fig. 7c), but Zr varies less than Ti on trachyte, while Ti is

from stoichiometric distributions owe to possible physical enrichments of Ti and Zr bearing-heavy minerals in different horizons of profiles (Figs. 7a and 7b). 317

Despite this, trachyte petrologic pattern is coarsely preserved in different horizons, 318 319 although eluviation processes produce 20% to 30% porosity while illuviation forms secondary crystallizations of gibbsite and clay-Fe cutans cementations in duricrusted 320 321 horizons (see Fig. 5). This results in grain density  $\rho_g$  of weathering samples close to that of the fresh trachyte (see Tab. 1a and Figs. 8a-b). These physical changes are all indicative of 322 noticeable volumetric variations across the profile (Fig. 8c; see also Tab. 1). However, the 323 overall variation of  $\varepsilon_{Zr}$  is ~13% less than that of  $\varepsilon_{Ti}$ , with a mean value of -51% (against -324 64 % for  $\varepsilon_{Ti}$  in the profile (Fig. 8c), supporting that Zr is the best reference element for 325 mass balance calculations on trachyte. On the other hand, in the profile on basalt where 326 porosity may increase to ca. 40 % (Fig. 9b),  $\varepsilon_{Ti}$  varies much less than  $\varepsilon_{Zr}$  (Fig. 9c) and a 327 328 better correlation between volumetric change  $\varepsilon_{Ti}$  and measured physical parameters validate Ti as most suitable reference element for mass balance calculation in the profile on basalt 329 330 (see Table. 1b and Fig. 9d).

In the next sections, the element mass transfers involved during formation of the two 331 weathering profiles are reported in the tables 2 and 3 and synthetized in the figure 10. The 332 table 2 reports masses  $(m_{i,w})$  of elements transferred through a unit volume  $(1 \text{ m}^3)$ 333 calculated from eq. 4. The table 3 shows masses transferred  $M_{i,w}$ , of these elements 334 calculated from eq. 5, and integrated over the investigated profiles depth related to the 335 corresponding height or thickness of chemically weathered rock. 336

337

338

## 4.3.2. *Element mass transfers in weathering profile on trachyte*

Ti shows a transfer rate of 35% in the saprolite relative to the trachyte (Tab. 1a and Fig. 339 8d) that results in 3.3 kg/m<sup>3</sup> or 11 kg of Ti concentrated in 1.5 m thick saprolite (Tab. 2a) 340

| 341 | and 3a, and Fig. 10a). Above the saprolite, the blocky bauxite (B5) and the massive bauxite                 |
|-----|---|
| 342 | (B3) horizons together show transfer rates of Ti from 21% to 56% that corresponds to ca. 2                  |
| 343 | to 8 kg/m <sup>3</sup> (Table 2a), or ca. 64 kg of Ti concentrated in ca. 10 m thick bauxitic duricrusted   |
| 344 | horizons (Table 3a and Fig. 10a). Al is depleted by 44% in the saprolite (Tab. 1a; Fig. 8d),                |
| 345 | corresponding to ca. 92 kg/m <sup>3</sup> or ca. 308 kg of Al exported from 1.5 m thick saprolite (Tabs     |
| 346 | 2a and 3a, and Fig. 10a). However, during formation of lateritic bauxitic duricrusts, the                   |
| 347 | transfer rate of Al ranges from - 33 % to + 35 % corresponding to transferred mass of ca. –                 |
| 348 | 68 to + 72 kg/m <sup>3</sup> (Tabs. 1a-2a), resulting in concentration of ca. 147 kg Al integrated over     |
| 349 | ca. 10 m thick lateritic profile (Tab. 3a and Fig. 10a). Fe is also depleted by 48% in the                  |
| 350 | saprolite (Tab. 1), with a loss of - 72 kg/m <sup>3</sup> implying that ca. 240 kg of Fe is released during |
| 351 | the formation of 1.5 m thick saprolite (Tabs. 2a and 3a and Fig. 10a). The transfer rate of Fe              |
| 352 | ranges from $-40\%$ to $+97\%$ in the bauxitic duricrusts corresponding to transferred masses               |
| 353 | of $-58$ to $+144$ kg/m <sup>3</sup> and at least ca. 708 kg Fe exported by the formation of 10 m thick     |
| 354 | bauxitic duricrusted profile, despite concentration of ca. 140 kg/m <sup>3</sup> at the base of             |
| 355 | duricrusted horizons (Tab. 2a and 3a and Fig. 10a).   |
| 356 | In summary, Ti is the most enriched element in the trachyte profile, followed by Al, while                  |
| 357 | Si, and P are the most depleted elements along the profile (Tab. 1a; Fig. 8d). Formation of                 |
| 358 | 1.5 m thick saprolite resulted in export of a total mass of ca. 2084 kg of Si at a rate of - 625            |
| 359 | $kg/m^3$ , while a mass of ca. 15898 kg was exported by formation of 10 m thick lateritic                   |
| 360 | duricrusted horizons (Tab. 2a and Fig. 10a). Mass changes calculation of trace elements                     |
| 361 | highlights enrichment of V, Co, Ni, As, Nb, Hf, Ta, Pb, U and Th, while Ba, Ga, Y, Sr, and                  |
| 362 | Rb are depleted. Compared to major elements, their exported/accumulated masses are very                     |
| 363 | low, with mean values of $\pm$ 0.001 to 0.1 kg, except Ba released up to -141 kg (see Tab.                  |
| 364 | DR4a).  |

366 4.3.3. Element mass transfers in weathering profile on basalt

| 367 | Al is gradually enriched throughout the profile from 6% in the saprolite to 134%  |
|-----|---|
| 368 | integrated in lateritic horizons $(B2 + B3 + B4)$ , despite slight depletion of 16% in the                              |
| 369 | mottled clays (Tab. 1b; Fig. 9d). This corresponds to ca. 20 kg Al concentrated in ca. 2 m                              |
| 370 | thick saprolite (C) at a rate of $11 \text{ kg/m}^3$ , while ca. 680 kg of Al concentrated in ca. 8 m                   |
| 371 | thick lateritic horizons (Tabs. 2b and 3b and Fig. 10b). Fe is depleted by 8% to 16% in the                             |
| 372 | saprolite and the mottled clays, respectively, and enriched up to 90% in the nodular bauxite                            |
| 373 | (Tab. 1b; Fig. 9d). These result in ca. 41 kg of Fe exported during saprolite formation at a                            |
| 374 | rate of 22.5 kg/m <sup><math>3</math></sup> , while a total mass of ca. 414 kg of Fe concentrated in bauxitic lateritic |
| 375 | horizons (Tabs. 2b and 3b, and Fig. 10b).   |
| 376 | In summary, Al is the most enriched element in the basalt profile, followed by Fe, while Si,                            |
| 377 | Mn and P are mostly depleted (Tabs. 2b and 3b; Figs. 9d and 10b). Si is strongly leached                                |
| 378 | with a total mass of ca. 570 kg exported by the formation of 2 m thick saprolite, while 4867                            |
| 379 | kg of Si has been released by that of 8 m thick duricrusted lateritic horizons (Tab. 3b and                             |
| 380 | Fig. 10b). The overall mass changes in trace elements show depletions in Rb, Sr, Y, Ba and                              |
| 381 | transitions elements, Co, Ni, Cu and Zn along the profile, while Nb, Zr, Ga, Cr, V, Th, Hf,                             |
| 382 | U and Ta are enriched in the upper 6 m of profile (Tab. DR4b). Most mass transfer values                                |
| 383 | are very low except that of Cr (5.1 kg) and Sr (-10.8 kg) in the upper meter of profile                                 |
| 384 | (nodular bauxite) (see Tab DR4b).   |
| 385 |   |

# **5. Discussion**

5.1. Differential bauxitic weathering and chemical erosion of volcanic rocks
The two bauxitic duricrusted profiles show different weathering and geochemical paths
from saprolite to lateritic duricrusted horizons. Owing to strong leaching of silica and
alkaline/alkaline earth elements, intense lateritization process resulted in gibbsite

| 391 | replacement of kaolinite and formation of iron oxy-hydroxides (goethite and hematite)            |
|-----|--|
| 392 | under redox conditions favorable to Al and Fe concentration (see Beauvais and Colin,             |
| 393 | 1993; Schellmann, 1994; Eggleton et al., 2008; Beauvais, 2009), which are illustrated by         |
| 394 | secondary gibbsite crystallizations, and clay-Fe coatings or cutans (Fig. 5b and 5d; see also    |
| 395 | Momo Nouazi et al., 2019).   |
| 396 | Element mass transfers point out important Al and Fe concentrations in the profile               |
| 397 | formed on basalt, while Al and Fe were seemingly exported from some bauxitic horizons of         |
| 398 | the profile formed on trachyte (Tab. 2 and Fig. 10) correlatively to the anomalous increase      |
| 399 | in reference elements Zr and Ti in these horizons (see Figs. 7c-d, Fig. 8 and Table DR2a).       |
| 400 | Enrichment of Zr and Ti, along with irregular Zr/Ti ratios during formation of trachyte          |
| 401 | profile may owe to their heterogeneity in the fresh trachyte (Brimhall et al., 1991; Schwarz,    |
| 402 | 1997). Mechanical reworking of weathering-resistant carrier heavy minerals (zircon,              |
| 403 | ilmenite and rutile) may also occur during long-term complex evolution of lateritic profiles     |
| 404 | (e.g., Brimhall et al., 1991; Colin et al., 1992; see also Beauvais, 2009). Given translocation  |
| 405 | marks in many lateritic weathering profiles (e.g., Fig. 5d), physical release of small size      |
| 406 | fractions of Zr and Ti carrier minerals (mostly zircon and titanomagnetite), which migrated      |
| 407 | downward into the compacted residual weathering profile, was also previously observed            |
| 408 | (e.g., Fig. 5d; see also Schwarz, 1997; Brimhall et al., 1988, 1991; Colin et al., 1992).        |
| 409 | Hence, this complex and differential behavior of Zr and Ti during the formation and              |
| 410 | evolution of residual lateritic profiles has resulted in differential strains implying the       |
| 411 | chemical weathering of protoliths (parent-rocks) of different thickness (see eq. 5).             |
| 412 | Given the ages of parent rocks (16 Ma for trachyte and 14 Ma for basalt),                        |
| 413 | quantification of their initial thickness before onset of weathering that produced the lateritic |
| 414 | profiles allows estimating average minimum rates of rock chemical erosion over the               |
| 415 | Neogene. Upon basalt, calculations with Ti give an integrated chemical rock-weathering           |

| 416 | rate of ca. 0.84 m/my over the last 14 Myr. A basalt thickness of ca. 12 m is required to                                |
|-----|--|
| 417 | produce the investigated residual ca. 10 m thick weathering profile at a rate of 0.70 m/my.                              |
| 418 | This also implies a topographic erosion of only ca. 2 m owing to congruent rock chemical                                 |
| 419 | erosion at a rate of ca. 0.14 m/my during the same period that is consistent with an almost                              |
| 420 | isovolumetric basalt weathering (average $\varepsilon_{Ti,w}$ = - 0,07; Tab. 1). Congruent chemical                      |
| 421 | weathering of $2m^3$ thick basalt ( $\rho = 2.74 \text{ t/m}^3$ ) corresponds to net exports of ca. 1076 kg Si,          |
| 422 | 377 kg Al and 546 kg Fe. Upon trachyte, calculations with Zr give an integrated chemical                                 |
| 423 | rock-weathering rate of c. 1.625 m/my. Here, chemical weathering of ca. 26 m thick                                       |
| 424 | trachyte led to formation of the 12 m thick studied profile consistently with an average                                 |
| 425 | strain value of $\varepsilon_{Zr,w}$ = - 0,51 (Tab. 1). This implies a topographic decay of similar height of            |
| 426 | 14 m corresponding to the thickness of rock layer, which has been chemically eroded                                      |
| 427 | congruently over the last 16 Myr at a rate of 0.875 m/my, while the residual weathering                                  |
| 428 | profile formed at rate of ca. 0.75 m/my that is comparable with the rate of profile formation                            |
| 429 | upon the basalt. Upon formation of the 12 m thick lateritic profile, the congruent chemical                              |
| 430 | erosion of 14 m <sup>3</sup> trachyte ( $\rho = 2.53 \text{ t/m}^3$ ) resulted in net exports of ca. 9849 kg Si, 2904 kg |
| 431 | Al and 2083 kg Fe.   |
| 432 | Therefore, differential chemical erosion of trachyte and basalt is primarily dependent                                   |
| 433 | on differences of strain ( $\varepsilon_{i,w}$ ) in profiles (Egli et Fitze, 2000). This implies formation of an         |
| 434 | equal 1 m thick bauxitic profile from the chemical weathering of a ca. 2.2 m thick layer of                              |
| 435 | trachyte against 1.2 m thick basalt layer over 1 My and 1.8 My, respectively. However,                                   |
| 436 | such differences call further discussion on parent-rocks geochemical differences and                                     |
| 437 | potential geomorphological constraints.  |



Under similar climatic and floristic constraints, differential chemical erosion and
bauxitic weathering potentially depend on parent rocks composition differences, but also
from drainage constraints upon profiles and landscape owing to local differential tectonics
related to the volcanic dynamics of Cameroon.

Although olivines are known to be more weathering-reactive than plagioclase and 444 pyroxenes (Eggleton et al., 1987; Schwarz, 1997), these minerals present grossly similar 445 weathering rates (Nesbitt and Wilson, 1992), no matter the grain size (Israeli and 446 Emmanuel, 2018). This suggests no direct (or little) influence of differential mineral 447 sensitivity on weathering rates and geochemical transfers on different volcanic rocks. 448 Nonetheless, Basalt carry mainly Fe-Mg minerals (olivine and clinopyroxene), while 449 450 trachyte are mostly Si-Al rich with dominant plagioclases and pyroxenes. However, the total mass of Si released during rock weathering and formation of evolved bauxitic profiles 451 (sum of element masses exported as synthesized in Fig. 10) is a few more than thrice on Si-452 rich trachyte than on basalt (17982 kg against 5437 kg), whose the lateritic bauxitic profile 453 has accumulated a few four time more Al than on trachyte (Tab. 3b and Fig. 10b). 454 According to Braun et al. (2016), chemical weathering rate is primarily controlled by 455 groundwater capacity to export solutes (e.g., dissolved silica, see also Schopka and Derry, 456 2012) far from the reacting solid-fluid interface to impede saturation of the weathering 457 system. This requires continuous and efficient drainage to sustaining propagation of the 458 weathering front into the bedrock (Braun et al., 2012; Prestrud Anderson et al., 2017, and 459 references there-in). Weathering in profile on trachyte may have been enhanced by 460 interconnected porosity (Momo Nouazi et al., 2019) that was formed by eluviation process 461 of less cohesive clayey matrix (Fig. 5a). On the other hand, compact and Fe-coated clay 462 matrix upon basalt has slowed rock weathering (Fig. 5c). This resulted here in rates lower 463

| 464 | than overall estimates in different geological and climatic contexts (Australia, Northern and                                    |
|-----|--|
| 465 | Southern America; see Stockmann et al., 2014), despite high alterability of basalt.  |
| 466 | However, rock-weathering rates, geochemical transfers and residual profile growth are  |
| 467 | primarily dependent on silica content of parent rock to be exported by groundwater to  |
| 468 | streams (Schopka and Derry, 2012). Higher Si abundance in trachyte relative to basalt  |
| 469 | (Figs. 4 and 6) is thus a major geochemical constraint to the growth of weathering profiles.                                     |
| 470 | Optimal drainage upon weathering front was instrumental for rapid dissolution of abundant  |
| 471 | Si carrier minerals (plagioclases) in trachyte, which turned into weathering rate-controlling                                    |
| 472 | minerals. A volcanic-related tectonic instability during Miocene (and later) has potentially                                     |
| 473 | led to rapid deepening of base level and landscape incision in Mount Bambouto (location in                                       |
| 474 | Fig. 2; see also Figs. DR1-a-b). This resulted in landscape rejuvenation, and optimal  |
| 475 | drainage that sustained eluviation processes in soils and weathering profiles, thus favoring                                     |
| 476 | silica export. Hence, saprolite production rate has been higher on trachyte than on basalt                                       |
| 477 | extruded at lower altitude, and where lateritic plateaus are less incised further away from                                      |
| 478 | tectonic influences (e.g., Prestud Anderson et al., 2007; Brantley et al., 2007; Schopka and                                     |
| 479 | Derry, 2012). Furthermore, differences in geomorphic processes resulted in differential  |
| 480 | evolution of profiles over Neogene, with an almost iso-volumetric formation on basalt  |
| 481 | ( $\varepsilon_{Ti,w}$ = - 0,07), while the profile on trachyte underwent substantial collapse ( $\varepsilon_{Zr,w}$ = - 0,51). |
| 482 | Therefore, geochemical differences in volcanic rocks compositions (mostly silica   |
| 483 | content) and geomorphological constraints potentially controlled by differential tectonics                                       |
| 484 | have resulted together in differentiation of chemical erosion and bauxitic weathering in   |
| 485 | highland western Cameroon (see also Samonil et al., 2020). The current bauxitic plateaus   |
| 486 | stand as long-lived elevated but differentially incised landforms (Figs. DR1a-b; see also  |
| 487 | Momo Nouazi et al., 2016). This implies the protolith-regolith interface has seemingly   |

489

moved downward non-uniformly with respect to differentially eroded/incised geomorphic surfaces on trachyte and basalt over the last 15 My.

490

## 491 **6.** Conclusion

492 Comparative geochemical mass balances of lateritic weathering of Mid-Miocene aged trachyte and basalt in highland western Cameroon document different processes of 493 chemical erosion and weathering profile formation. Reference elements Zr and Ti have 494 been used from their relative stability for calculating element mass transfers in weathering 495 profiles. Despite some geochemical anomalies in Zr or Ti owing to possible heterogeneity 496 497 in parent photoliths and physical enrichments of carrier heavy minerals in residual profiles, 498 element mass transfers indicate higher Si depletion and lower Al-Fe concentrations on trachyte than on basalt. Our results document the formation of 1 m thick bauxitic profile 499 500 from lateritic weathering of 1.2 m of basalt or 2.2 m of trachyte, implying chemical erosion and weathering rates lower on basalt than on trachyte. Different strain pattern in the 501 investigated weathering profiles imply a congruent dissolution (net chemical erosion) of 502 initial protolith layer larger on trachyte (14 m) than on basalt (2 m). Differential chemical 503 weathering rates are mostly controlled by drainage conditions at the regolith-photolith 504 505 interface and the Si content of volcanic rocks to be released to form bauxitic regolith. Finally, our results point out combined influence of differential local tectonics, and 506 geochemical differences of volcanic rocks on chemical erosion and bauxitic weathering, 507 with implications on the evolution of western Cameroon landscapes over the Neogene. 508

509

Acknowledgments - Financial support was provided by the 'Service de Coopération et
d'Action Culturelle au Cameroun (SCAC)' from the French Embassy (791779A). The
mineralogical analyses have been carried out at the Centre Européen de Recherche et

| 513 | d'Enseignement des Géosciences de l'Environnement (CEREGE), Aix Marseille University,       |
|-----|---|
| 514 | OSU Pytheas, France. The geochemical analysis has been carried out at ALS Laboratory        |
| 515 | Group SL-Spain. MNM was supported by a 6 months fellowship from the 'Institute of           |
| 516 | Research for Development' (IRD, France). We acknowledge Dr. Nkouathio for                   |
| 517 | geochronological data and assistance on volcanic model of the mount Bambouto. Jean Paul     |
| 518 | Ambrosi is thanked for technical assistance and discussion of some results. We are grateful |
| 519 | to Dr. R.A. Eggleton, an anonymous reviewer, and Pr. Egli, for their respective remarks     |
| 520 | and suggestions that together contributed to clarify and improve our manuscript.            |
| 521 |   |
| 522 | References  |
| 523 | American Mineralogist Crystal Structure Database. http://rruff.geo.arizona.edu/AMS/         |
| 524 | amcsd. php, Accessed date: 8 June 2017.   |
| 525 | Babechuk, M.G., Widdowson, M., Kamber, B.S., 2014. Quantifying chemical intensity and       |
| 526 | trace element release from two contrasting profiles, Decan Traps, India. Chem. Geol.        |
| 527 | 365, 56–75.   |
| 528 | Babechuk, M.G., Widdowson, M., Murphy, M., Kamber, B.S., 2015. A combined Y/Ho,             |
| 529 | high field strength element (HFSE) and Nd isotope perspective on basalt weathering,         |
| 530 | Deccan Traps, India. Chemical Geology 396, 25-41.   |
| 531 | Bardossy, G.Y, Aleva, G.J.J., 1990. Lateritic bauxites. <i>Elsevier</i> , Amsterdam, 624 p. |
| 532 | Beauvais, A., 2009. Ferricrete biochemical degradation on the rainforest savannas boundary  |
| 533 | of Central African Republic. Geoderma 150, 379-388.   |
| 534 | Beauvais, A. Colin, F., 1993. Formation and transformation processes of iron duricrust      |
| 535 | systems in tropical humid environment. Chem. Geol. 106, 77-151.                             |

| 536 | Beauvais, A., Chardon, D., 2013. Modes, tempo, and spatial variability of Cenozoic                   |
|-----|--|
| 537 | cratonic erosion: the West African example. Geochem. Geophys. Geosyst. 14, 1590-                     |
| 538 | 1608. https://doi.org/10.1002/ggge.20093.  |
| 539 | Beauvais, A., Ruffet, G., Hénocque, O., Colin, F. 2008. Chemical and physical erosion                |
| 540 | rhythms of the West African Cenozoic morphogenesis: The <sup>39</sup> Ar- <sup>40</sup> Ar dating of |
| 541 | supergene K-Mn oxides. J. Geophys. Res. 113, F04007, doi:10.1029/2008JF000996.                       |
| 542 | Boulangé, B. Millot, G., 1988. La distribution des bauxites sur le craton ouest-africain. Sci.       |
| 543 | Géol., Bull. 41, 113-123.  |
| 544 | Boulangé, B., 1984. Les formations bauxitiques latéritiques de Côte d'Ivoire. Les faciès,            |
| 545 | leur transformation, leur distribution et l'évolution du modelé. In: Thèse et Mémoire                |
| 546 | ORSTOM 363 pp.   |
| 547 | Boulangé, B., Colin, F., 1994. Rare earth element mobility during conversion of nepheline            |
| 548 | syenite into lateritic bauxite at Passa Quatro, Minais Gerais, Brazil. Applied                       |
| 549 | Geochemistry 9, 701-711.   |
| 550 | Brantley S.L., Goldhaber M.B., Ragnarsdottir, K.V., 2007. Crossing disciplines and scales            |
| 551 | to understand the critical zone. Elements 3, 307-314.  |
| 552 | Brantley, S.L., Lebedeva, M., 2011. Learning to read the chemistry of regolith to                    |
| 553 | understand the critical zone. Annual Review of Earth Planetary Science 39, 387-416.                  |
| 554 | Braun, J.J., Mercier, J., Guillocheau, F., Robin, C., 2016. A simple model for regolith              |
| 555 | formation by chemical weathering. Journal of Geophysical Research: Earth Surface                     |
| 556 | 121, 2140-2171, doi:10.1002/2016JF003914.  |
| 557 | Braun, J. J., Pagel, M., Herbillon A., Rosin C., 1993. Mobilization and redistribution of            |
| 558 | REEs and thorium in syenitic lateritic profile: A mass balance study. Geochim.                       |
| 559 | Cosmochim. Acta 57, 4419-4434.   |

| 560 | Braun, J.J., Marechal, J.C., Riotte, J., Boeglin, J.L., Bedimo Bedimo, J.P., Ndam            |
|-----|--|
| 561 | Ngoupayou, J.R., Nyeck, B., Robain, H., Sekhar, M., Audry, S., Viers, J., 2012.              |
| 562 | Elemental weathering fluxes and saprolite production rate in a Central African               |
| 563 | lateritic terrain (Nsimi, South Cameroon). Geochimica et Cosmochimica Acta 99,               |
| 564 | 243-270.   |
| 565 | Brimhall, G. H., Dietrich, W. E., 1987. Constitutive mass balance relations between          |
| 566 | chemical composition, volume, density, porosity, and strain in metasomatic                   |
| 567 | hydrochemical systems: Results on weathering and pedogenesis. Geochim.                       |
| 568 | Cosmochim. Acta 51, 567-587.   |
| 569 | Brimhall, G. H., Lewis, C. J., Ague, J. J., Dietrich, W. E., Hampel, J., Teague, T., Rix P., |
| 570 | 1988. Metal enrichment in bauxites by deposition of chemically mature aeolian dust.          |
| 571 | Nature 333, 819-824.   |
| 572 | Brimhall, G.H., Lewis, C.J., Ford, C., Bratt, J., Taylor, G., Warin, O., 1991. Quantitative  |
| 573 | geochemical approach to pedogenesis: importance of parent material reduction,                |
| 574 | volumetric expanxion and eolian influx on lateritization. Geoderma 51, 51-91.                |
| 575 | Burke, K., Gunnell, Y., 2008. The African Erosion Surface: A Continental-Scale Synthesis     |
| 576 | of Geomorphology, Tectonics, and Environmental Change over the Past 180 Million              |
| 577 | Year. Geological Society of America, Memoir 201, 66 p.                                       |
| 578 | Chadwick, O.A., Brimhall, G.H., Hendricks , D.M., 1990. From a black to a gray box-A         |
| 579 | mass balance interpretation of pedogenesis. Geomorphology 3, 369-390.                        |
| 580 | Chardon, D., Grimaud, J. L., Beauvais, A., Bamba, O., 2018. West African lateritic           |
| 581 | pediments: Landform-regolith evolution processes and mineral exploration pitfalls.           |
| 582 | Earth- Sci. Rev. 179, 124-146.   |
| 583 | Colin, F., Brimhall, G. H., Nahon, D., Baronnet, A., Kathy, D., 1992. Equatorial rainforest  |
| 584 | lateritic mantles: a geomembrane filter. Geology 20, 523-526.                                |

| 585 | De Putter, T., Ruffet, G., 2020. Supergene manganese ore records 75 Myr-long Campanian         |
|-----|--|
| 586 | to Pleistocene geodynamic evolution and weathering history of the Central African              |
| 587 | Great Lakes Region - Tectonics drives, climate assists. Gondwana Research 83, 96-              |
| 588 | 117.   |
| 589 | Déruelle, B, Ngounouno, I, Demaiffe, D., 2007. The 'Cameroon Hot Line' (CHL): a unique         |
| 590 | example of active alkaline intraplate. structure in both oceanic and continental               |
| 591 | lithospheres. C.R. Geoscience 339, 589-600.  |
| 592 | Eggleton, R. A., Taylor, G., Le Gleuher, M., Foster, L. D., Tilley, D. B., Morgan C. M.,       |
| 593 | 2008. Regolith profile, mineralogy and geochemistry of the Weipa Bauxite, northern             |
| 594 | Australia. Australian Journal of Earth Sciences 55, 17-43. Doi                                 |
| 595 | .org/10.1080/08120090802438233.  |
| 596 | Eggleton, R.A., Foudoulis, C., Varkevisser, D., 1987. Weathering of basalt: changes in rock    |
| 597 | chemistry and mineralogy. Clays and Clay Minerals, 35(3), 161-169.                             |
| 598 | Egli, M., Fitze, P., 2000. Formulation of pedologic mass balance based on immobile             |
| 599 | elements: a revision. Soil Science 165, 437-443.   |
| 600 | Fosso, J., Ménard, J. J., Bardintzeff, J. M., Wandji, P., Tchoua, F. M., Bellon, H., 2005. Les |
| 601 | laves du mont Bangou : une première manifestation volcanique éocène, à affinité                |
| 602 | transitionnelle, de la Ligne du Cameroun. C.R. Geosciences 337, 315-325.                       |
| 603 | Israeli, Y., Emmanuel, S., 2018. Impact of grain size and rock composition on simulated        |
| 604 | rock weathering. Earth Surface Dynamics 6, 319-327.  |
| 605 | Jean, A., Beauvais, A., Chardon, D., Arnaud, N., Jayanada M., Mathe, P. E., 2020.              |
| 606 | Weathering history and landscape evolution of Western Ghats (India) from                       |
| 607 | 40Ar/39Ar dating of supergene K-Mn oxides. Journal of the Geological Society,                  |
| 608 | London. In press, doi: 10.1144/jgs2019-048.  |

| 609 | Kengni, L., Tekoudjou, H., Tematio, P., Tedonkeng, E.P., Mubeteneh Tankou, C., 2009.                      |
|-----|---|
| 610 | Rainfall Variability along the Southern Flank of the Bambouto Mountain (West-                             |
| 611 | Cameroon). Journal of the Cameroon academy of sciences 8, 45-52.  |
| 612 | Kwékam, M., Liégeois, J.P., Njonfang, E., Affaton, P., Hartmann, G., 2010. Nature, origin                 |
| 613 | and significance of the Fomopéa Pan-African high-K calc-alkaline plutonic complex                         |
| 614 | in the Central African fold belt (Cameroon). J. Afr. Earth Sci. 57, 79-95.                                |
| 615 | Marzoli, A., Piccirillo, E. M., Renne, P. R., Bellieni, G., Iacumin, M., Nyobe, J. B.,                    |
| 616 | Tongwa, A. T., 2000. The Cameroon Volcanic Line revisited: Petrogenesis of                                |
| 617 | continental basaltic magmas from lithospheric and asthenospheric mantle sources. J                        |
| 618 | Petrol 41:87-109.   |
| 619 | Marzoli, A., Renne, P. R., Piccirillo, E. M., Francesca, C., Bellieni, G., Mel, A. J., Nyobe,             |
| 620 | J. B., N'ni, J., 1999. Silicic magmas from the continental Cameroon Volcanic Line                         |
| 621 | (Oku, Bambouto and Ngaoundéré): <sup>40</sup> Ar/ <sup>39</sup> Ar dates, petrology, Sr-Nd-O isotopes and |
| 622 | their petrogenetic significance. Contrib Mineral Petrol 135, 133-150.                                     |
| 623 | Millot, G., Bonifas, M., 1959. Transformations isovolumétriques dans les phénomènes de                    |
| 624 | latéritisation et de bauxitisation. Bull. Serv. Carte Géol. Alsace. Lorraine 8, 3-10.                     |
| 625 | Momo Nouazi, M., Beauvais, A., Tematio, P., Ambrosi, J-P., Yemefack, M., Yerima,                          |
| 626 | P.K.B., Yongue-Fouateu, R., 2019. Lateritic weathering of trachyte, and bauxite                           |
| 627 | formation in West Cameroon: Morphological and geochemical evolution. Journal of                           |
| 628 | geochemical exploration 205, 106324. doi.org/10.1016/j.gexplo.2019.06.006.                                |
| 629 | Momo Nouazi, M., Yemefack, M., Tematio, P., Beauvais, A., Ambrosi, JP., 2016.                             |
| 630 | Distribution of duricrusted bauxites and laterites on the Bamiléké plateau (West                          |
| 631 | Cameroon): Constraints from GIS mapping and geochemistry. Catena 140, 15-23.                              |
| 632 | Moundi, A., Wandji, P., Bardintzeff, J. M., Ménard, J. J., Okomo Atouba, L. C.,                           |
| 633 | Mouncherou, O. F., Reusser, E., Bellon, H., Tchoua, F. M., 2007. Les basaltes                             |

| 634 | éocènes à affinité transitionnelle du plateau Bamoun, témoins d'un réservoir                               |
|-----|--|
| 635 | mantellique enrichi sous la Ligne Volcanique du Cameroun. C.R. Geosciences 339,                            |
| 636 | 396-406  |
| 637 | Nesbitt, H. W., Wilson, R. E., 1992. Recent chemical weathering of basalt. Am. J. Sci. 292,                |
| 638 | 740-777.   |
| 639 | Nkouathio, D. G., Kagou Dongmo, A., Bardintzeff, J. M., Wandji, P., Bellon, H., Pouclet,                   |
| 640 | A., 2008. Evolution of volcanism in graben and horst structures along the Cenozoic                         |
| 641 | Cameroon Line (Africa): implications for tectonic evolution and mantle source                              |
| 642 | composition. Mineral. Petrol. 94, 287-303.   |
| 643 | Perelló, J., Brockway, H., Garcia, A., 2020. A minimum Thanetian (Paleocene) age for the                   |
| 644 | African Surface in the Eritrean highlands, Northeast Africa. Journal of African Earth                      |
| 645 | Sciences 164, 103782.  |
| 646 | Prestrud Anderson, S., Von Blanckenburg, F., White, A.F., 2007. Physical and chemical                      |
| 647 | control on the critical zone. Elements 3, 315-319.   |
| 648 | Retallack, G.J. 2010. Lateritization and Bauxitization Events. <i>Economic Geology</i> , <b>105</b> , 655- |
| 649 | 667.   |
| 650 | Rietveld, H.M., 1969. A profile refinement method for nuclear and magnetic structures. J.                  |
| 651 | Appl. Crystallogr. 2, 65-71.   |
| 652 | Samonil, P., Philipps, J., Danek, P., Benes V., Pawlik, L., 2020. Soil, regolith, and                      |
| 653 | weathered rock: Theoretical concepts and evolution in old-growth temperate forests,                        |
| 654 | Central Europe. Geoderma 368, 114261. doi:10.1016/j.geoderma.2020.114261.                                  |
| 655 | Schellmann, W., 1994. Geochemical differenciation in laterite and bauxite formation. In                    |
| 656 | Lateritization Processes and Supergene Ore Formation. Catena Spec. Issue (eds T.                           |
| 657 | Schwarz and K. Germann), 21, 141-143. Elsevier.  |

| 658 | Schopka, H.H., Derry, L.A., 2012. Chemical weathering fluxes from volcanic islands and       |
|-----|--|
| 659 | the importance of groundwater: The Hawaiian example. Earth and Planetary Science             |
| 660 | Letters 339-340, 67-78.  |
| 661 | Schwarz, T., 1997. Distribution and genesis of bauxite on the Mambilla plateau. SE           |
| 662 | Nigeria. Applied Geochemistry 12, 119-131.   |
| 663 | Soares de Oliveira, F., Chicarino Varajão, A. F. D., Chicarino Varajão, C. A., Boulangé, B., |
| 664 | Vieira Soares, C. C., 2013. Mineralogical, micromorphological and geochemical                |
| 665 | evolution of the facies from the bauxite deposit of Barro Alto, Central Brazil. Catena       |
| 666 | 105, 29-39.  |
| 667 | Soler, J. M., Lasaga, A. C., 2000. The Los Pijiguaos bauxite deposit (Venezuela): A          |
| 668 | compilation of field data and implications for the bauxitization process. Journal of         |
| 669 | South American Earth Sciences 13, 47-65.   |
| 670 | Stockmann, U., Minasny, B., McBratney, A.B., 2014. How fast does soil grow? Geoderma         |
| 671 | 216, 48-61.  |
| 672 | Tardy, Y., 1997. Petrology of Laterites and Tropical Soils. Balkema, Rotterdam, 408 pp.      |
| 673 | Tematio, P., Fritsch, E., Hodson, M.E., Lucas, Y., Bitom, D., Bilong, P., 2009. Mineral and  |
| 674 | geochemical characterization of a leptic aluandic soil and a thapto aluandic-ferralsol       |
| 675 | developed on trachyte in Mount Bambouto (Cameroon volcanic line). Geoderma 152,              |
| 676 | 314-323.   |
| 677 | Tematio, P., Olson, K.R., 1997. Characterization of two phases of encrustment in a           |
| 678 | sequence of ferrallitic soils in the South Cameroon and its effects on landscape             |
| 679 | evolution. Soil Sci. 162 (10), 758-766.  |
| 680 | Temgoua, E., Pfeifera, H-R., Bitom, D., 2003. Trace element differentiation in ferruginous   |
| 681 | accumulation soil patterns under tropical rainforest of southern Cameroon, the role of       |
| 682 | climatic change. The Science of the Total Environment 303, 203-214.                          |

| 683 | Thomas, M. F., 1994. Geomorphology in the tropics: a study of weathering and denudation    |
|-----|--|
| 684 | in low altitudes. John Wiley and Sons, Chichester, 460 p.                                  |
| 685 | Ufer, K., Stanjek, H., Roth, G., Dohrmann, R., Kleeberg, R., Kaufhold, S., 2008.           |
| 686 | Quantitative phase analysis of bentonites by the Rietveld method. Clay Clay Miner.         |
| 687 | 56 (2), 272-282.   |
| 688 | Valeton, I., 1999. Saprolite-bauxite facies of ferralitic duricrusts on palaeosurfaces of  |
| 689 | former Pangea. Sp. Pub. Int. Ass. Sedim. 27, 153-188.                                      |
| 690 | Valeton, I., Beißner, H., 1986. Geochemistry and mineralogy of the lower tertiary in situ  |
| 691 | laterites of the Jos plateau. Journal of African Earth Sciences 5, 535-550.                |
| 692 | Wille, M., Babechuk, M.G., Kleinhanns, I.C., Steigmaier, C., Suhr, N., Widdowson, M.,      |
| 693 | Kamber, B.S., Shoenboerg, R., 2018. Silicon and chromium stable isotopic                   |
| 694 | systematics during basalt weathering and lateritisation: A comparison of variably          |
| 695 | weathered basalt profiles in the Deccan Traps, India. Geoderma 314, 190-204.               |
| 696 | White, A. F., Bullen, T. D., Schultz, M. S., Blum A. E., Huntington, T. G., Peters, N. E., |
| 697 | 2001. Differential rates of feldspar weathering in granitic regoliths. Geochimica et       |
| 698 | Cosmochimica Acta 65, 847-869.   |
| 699 | Zarasvandi, A., Carranza, E. J. M., Ellahi, S. S., 2012. Geological, geochemical and       |
| 700 | mineralogical characteristics of the Deh-Now bauxite deposits, Zagros fold belt, Iran.     |
| 701 | Ore Geol. Rev. 48, 125-138.  |
| 702 |  |
| 703 |  |
| 704 |  |
| 705 |  |

### FIGURES CAPTION

Figure. 1. a) Location of the Western Cameroon Highlands. The main geological features 707 (Adapted from Nkouathio et al., 2008) are superimposed to the digital elevation model of 708 Cameroon. CCSZ=Central Cameroonian shear zone; b) Morphology of the Western 709 710 Cameroon Highlands and location of the study area with main occurrences of bauxite and ferricrete bearing surfaces. 711 Figure. 2. a) Geological map of the Western Cameroon Highland and mapped bauxitic 712 713 surfaces (Bamileke plateau, see insert in Figure 1b). b) Morpho-geological section showing the topographic location of studied bauxitic profiles. See section trace on Figure 2a. 714 Figure. 3. a) Macroscopic view of trachyte sample; b and c) Crossed-polarized microscope 715 photomicrographs of the porphyric trachyte. d) Macroscopic view of basalt sample; d) and 716 717 e) Crossed-polarized microscope photomicrographs of the porphyric basalt. 1= plagioclases; 2= olivine; 3= titanomagnetite; 4= clinopyroxenes. 718 Figure. 4. Petrographic structures, location of samples collected (squares), and variations 719 of major oxides content in weathering profiles formed on a) trachyte, and b) basalt. Grey 720 squares= samples analyzed for mass transfer balance. 721 722 Figure. 5. Transmitted natural light microscopic photomicrographs of a) saprolite and b) bauxitic duricrusts from lateritic profile on trachyte; and from lateritic profile on basalt, c) 723 saprolite showing weathering of primary plagioclases and cpx phenocrystal; and d) 724 bauxitic duricrust. 1= primary gibbsite crystallizations; 2= Ferruginous argillic matrix; 3= 725 726 void; 4= primary gibbsite with Fe-coating; 5= goethite rich Fe clay cutan; 6= Clay 727 ferruginous infill; 7= secondary gibbsite crystallizations; 8= gey-argillic matrix with primary gibbsite and Fe-coating; 9= translocated heavy minerals. 728

| 729 | Figure. 6. Distribution of major oxides composition of parent rocks, saprolite and  |
|-----|---|
| 730 | duricrusted horizons in profiles formed on trachyte and basalt.   |
| 731 | Figure. 7. Variations in the content of reference elements Zr and Ti in weathering profiles   |
| 732 | formed on trachyte and basalt. a) and b) Diagrams Zr vs. Ti (see Figure 6 for explanations  |
| 733 | of symbols; Black straight lines indicate stoichiometric distributions); c) Zr/Ti vertical  |
| 734 | variations along both profiles; Zr and Ti variations along profile on d) trachyte, and e)   |
| 735 | basalt.   |
| 736 | Figure. 8. Variation of physical parameters along the profile on trachyte. a) Densities; b)   |
| 737 | Porosity; c) Strain; and, d) Transfer rates of major elements.  |
| 738 | Figure. 9. Variation of physical parameters along the profile on basalt. a) Densities; b)   |
| 739 | Porosity; c) Strain; and, d) Transfer rates of major elements.  |
| 740 | Figure. 10. Comparative transferred masses of major elements over integrated thicknesses  |
| 741 | of protolith layers chemically weathered to form the saprolite (yellow) and bauxitic  |
| 742 | duricrusted horizons (pink) in the weathering profile formed on a) trachyte and b) basalt.  |
| 743 | (Masse calculated from equation 5; see also Table 3).   |
| 744 |   |
| 745 | TABLES CAPTION  |
| 746 | <b>Table. 1</b> . Variations of bulk and grain density ( $\rho_b$ and $\rho_g$ in g.cm <sup>-3</sup> ), porosity, $\phi$ , strain,        |
| 747 | $\boldsymbol{\epsilon}$ , and elements transfer rates, $\tau$ , in weathering profiles formed upon (a) trachyte ( $\boldsymbol{\epsilon}$ |
| 748 | calculated with reference element Zr), and (b) basalt ( $\epsilon$ calculated with reference element                                      |
| 749 | Ti). Calculated from the equations 1 to 3, respectively. The true depth of trachyte being   |
| 750 | unknown we simply mentioned a depth in parentheses for graphical easiness of figures.   |
| 751 | <b>Table. 2</b> . Elements mass transfers in kg per unit volume $(1 \text{ m}^3)$ in weathering profiles                                  |
| 752 | formed on (a) trachyte and (b) basalt. Calculated from the equation 4.  |

**Table. 3.** Transferred masses of elements integrated over the investigated profile horizonsthickness ( $\Delta z_w$ ), related to the corresponding height or thickness of chemically weatheredrock layer ( $\Delta z$ ), in profile on a) trachyte, and b) basalt. Calculated from the equation 5.

| (a)       | Horizon | Depth (m) | $\rho_b$ | $\rho_{g}$ | φ    | $\epsilon_{Zr}$ | $\mathbf{\tau}_{\mathrm{Si}}$ | $\mathbf{\tau}_{\mathrm{Al}}$ | $\mathbf{\tau}_{\mathrm{Fe}}$ | $\mathbf{\tau}_{\mathrm{Ti}}$ | $\mathbf{\tau}_{Ca}$ | $\mathbf{\tau}_{\mathrm{Mg}}$ | $\mathbf{\tau}_{\mathrm{Na}}$ | $\mathbf{\tau}_{\mathrm{K}}$ | $\mathbf{\tau}_{Mn}$ | $\boldsymbol{\tau}_{\mathrm{P}}$ |
|-----------|---------|-----------|----------|------------|------|-----------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|----------------------|-------------------------------|-------------------------------|------------------------------|----------------------|----------------------------------|
| P2-FO-2   | B3      | 2.50      | 1.73     | 2.28       | 24.0 | -0,48           | -0.99                         | 0.17                          | -0.40                         | 0.32                          | -1.00                | -0.98                         | -1.00                         | -1.00                        | -0.95                | -0.27                            |
| P2-FO-3   | В3      | 4.50      | 1.93     | 2.50       | 22.7 | -0,68           | -1.00                         | -0.30                         | -0.26                         | 0.36                          | -1.00                | -1.00                         | -1.00                         | -1.00                        | -0.96                | -0.46                            |
| P2-FO-4   | В3      | 6.00      | 1.90     | 2.32       | 18.4 | -0,54           | -1.00                         | 0.09                          | -0.16                         | 0.21                          | -1.00                | -1.00                         | -1.00                         | -1.00                        | -0.96                | -0.44                            |
| P2-FO-5   | В3      | 8.00      | 1.69     | 2.47       | 31.7 | -0,65           | -1.00                         | -0.33                         | -0.30                         | 0.27                          | -0.99                | -1.00                         | -1.00                         | -1.00                        | -0.96                | -0.44                            |
| P2-FO-6   | B3      | 9.00      | 1.87     | 2.43       | 23.3 | -0,50           | -0.99                         | 0.22                          | -0.24                         | 0.26                          | -1.00                | -1.00                         | -1.00                         | -1.00                        | -0.92                | -0.63                            |
| P2-FO-7   | B3      | 10.0      | 1.83     | 2.35       | 22   | -0,44           | -0.99                         | 0.35                          | -0.16                         | 0.33                          | -1.00                | -1.00                         | -1.00                         | -1.00                        | -0.92                | -0.32                            |
| P2-FO-7.1 | B3      | 10.5      | 2.76     | 2.82       | 2.06 | -0,30           | -0.98                         | 0.17                          | 0.97                          | 0.36                          | -1.00                | -1.00                         | -1.00                         | -1.00                        | -0.67                | 1.55                             |
| P2-FO-8   | В5      | 11.0      | 1.74     | 2.46       | 29.2 | -0,46           | -0.99                         | 0.19                          | -0.10                         | 0.56                          | -1.00                | -1.00                         | -1.00                         | -1.00                        | -0.93                | -0.42                            |
| P2-FO-8.1 | В5      | 11.5      | 1.64     | 2.42       | 32.3 | -0,49           | -0.97                         | -0.16                         | 0.03                          | 0.52                          | -0.99                | -0.91                         | -1.00                         | -1.00                        | -0.91                | -0.15                            |
| P3-FO-3   | С       | 12.0      | 1.41     | 2.04       | 30.9 | -0,55           | -0.89                         | -0.44                         | -0.48                         | 0.35                          | -1.00                | -0.99                         | -1.00                         | -1.00                        | -0.95                | -0.14                            |
| Trachyte  |         | (13.0)    | 2.53     | 2.55       | 0.78 | 0,00            | 0.00                          | 0.00                          | 0.00                          | 0.00                          | 0.00                 | 0.00                          | 0.00                          | 0.00                         | 0.00                 | 0.00                             |
| (b)       |         |           |          |            |      | ε <sub>Ti</sub> |                               |                               |                               |                               |                      |                               |                               |                              |                      |                                  |
| P1-SA     | B2      | 1.00      | 2.33     | 2.58       | 9.69 | 0.14            | -0.78                         | 1.68                          | 0.90                          |                               | -1.00                | -0.99                         | -1.00                         | -0.98                        | -0.83                | -0.50                            |
| P1-SA-2   | B3      | 4.00      | 2.06     | 2.78       | 25.9 | -0.39           | -0.98                         | 0.25                          | 0.10                          |                               | -1.00                | -0.99                         | -1.00                         | -1.00                        | -0.88                | -0.91                            |
| P1-SA-3   | B4      | 6.50      | 1.36     | 2.21       | 38.6 | 0.00            | -0.76                         | 0.05                          | -0.08                         |                               | -1.00                | -0.98                         | -1.00                         | -0.98                        | -0.74                | -0.80                            |
| P1-SA-4   | B4      | 8.00      | 1.66     | 2.51       | 34   | -0.31           | -0.84                         | -0.16                         | -0.16                         |                               | -1.00                | -0.97                         | -1.00                         | -1.00                        | -0.78                | -0.49                            |
| P1-SA-5   | С       | 9.50      | 1.38     | 2.25       | 38.8 | 0.21            | -0.58                         | 0.06                          | -0.08                         |                               | -0.99                | -0.91                         | -1.00                         | -0.95                        | -0.44                | -0.42                            |
| Basalt    |         | 10.5      | 2.74     | 2.74       | 0.0  | 0.00            | 0.00                          | 0.00                          | 0.00                          |                               | 0.00                 | 0.00                          | 0.00                          | 0.00                         | 0.00                 | 0.00                             |

Table. 1

| (a)       | Horizon | Depth (m) | $m_{Si}$ | $m_{Al}$ | m <sub>Fe</sub> | $m_{\mathrm{Ti}}$ | $m_{Ca}$ | $m_{Mg}$ | m <sub>Na</sub> | $m_K$  | $m_{Mn}$ | $m_P$ |
|-----------|---------|-----------|----------|----------|-----------------|-------------------|----------|----------|-----------------|--------|----------|-------|
| P2-FO-2   | B3      | 2.5       | -695.3   | 34.6     | -58.8           | 3.0               | -51.5    | -3.2     | -81.7           | -106.5 | -7.2     | -0.5  |
| P2-FO-3   | В3      | 4.5       | -700.4   | -62.4    | -39.1           | 3.3               | -51.4    | -3.3     | -81.7           | -106.7 | -7.3     | -0.8  |
| P2-FO-4   | В3      | 6.0       | -700.6   | 18.3     | -24.4           | 1.9               | -51.5    | -3.3     | -81.7           | -106.7 | -7.3     | -0.8  |
| P2-FO-5   | В3      | 8.0       | -701.4   | -68.0    | -44.1           | 2.6               | -51.3    | -3.3     | -81.7           | -106.7 | -7.3     | -0.8  |
| P2-FO-6   | В3      | 9.0       | -695.8   | 45.8     | -35.4           | 2.4               | -51.5    | -3.3     | -81.7           | -106.7 | -6.9     | -1.1  |
| P2-FO-7   | В3      | 10.0      | -697.3   | 72.2     | -24.2           | 3.1               | -51.5    | -3.3     | -81.7           | -106.7 | -7.0     | -0.6  |
| P2-FO-7.1 | В3      | 10.5      | -688.7   | 35.4     | 144.3           | 3.4               | -51.5    | -3.3     | -81.7           | -106.6 | -5.1     | 2.7   |
| P2-FO-8   | В5      | 11.0      | -696.6   | 39.9     | -15.1           | 5.3               | -51.6    | -3.3     | -81.7           | -106.6 | -7.1     | -0.7  |
| P2-FO-8.1 | В5      | 11.5      | -680.6   | -32.8    | 4.1             | 4.9               | -51.2    | -3.0     | -81.7           | -106.3 | -6.9     | -0.3  |
| P3-FO-3   | С       | 12.0      | -625.3   | -92.3    | -72.0           | 3.3               | -51.4    | -3.2     | -81.7           | -106.7 | -7.2     | -0.2  |
| Trachyte  |         | (13.0)    | 0.00     | 0.00     | 0.00            | 0.00              | 0.00     | 0.00     | 0.00            | 0.00   | 0.00     | 0.00  |
| (b)       |         |           |          |          |                 |                   |          |          |                 |        |          |       |
| P1-SA     | B2      | 1.0       | -421.7   | 316.6    | 246.0           |                   | -177.2   | -182.8   | -42.7           | -21.2  | -4.1     | -3.3  |
| P1-SA-2   | В3      | 4.0       | -527.4   | 46.4     | 28.5            |                   | -177.5   | -183.1   | -42.7           | -21.7  | -4.3     | -5.9  |
| P1-SA-3   | B4      | 6.5       | -408.0   | 9.6      | -22.7           |                   | -177.1   | -180.7   | -42.7           | -21.2  | -3.7     | -5.3  |
| P1-SA-4   | B4      | 8.0       | -453.1   | -29.8    | -43.5           |                   | -176.9   | -179.5   | -42.7           | -21.6  | -3.9     | -3.2  |
| P1-SA-5   | С       | 9.5       | -313.7   | 10.8     | -22.5           |                   | -176.4   | -168.6   | -42.7           | -20.5  | -2.2     | -2.8  |
| Basalt    |         | 10.5      | 0.0      | 0.0      | 0.0             |                   | 0.0      | 0.0      | 0.0             | 0.0    | 0.0      | 0.0   |

Table 2.

| (a)       | Horizon | Depth (m) | $\Delta z(m)$ | $\Delta z_{w}\left(m ight)$ | $M_{Si}$ | $M_{Al}$ | $M_{\rm Fe}$ | $M_{Ti}$ | $M_{Ca}$ | $M_{\text{Mg}}$ | $M_{\mathrm{Na}}$ | $M_{K}$ | $M_{Mn} \\$ | $M_P$ |
|-----------|---------|-----------|---------------|-----------------------------|----------|----------|--------------|----------|----------|-----------------|-------------------|---------|-------------|-------|
| P2-FO-2   | B3      | 2.5       | 4.8           | 2.5                         | -3349    | 167      | -283         | 15       | -248     | -16             | -394              | -513    | -35         | -2    |
| P2-FO-3   | В3      | 4.5       | 3.1           | 1.0                         | -2165    | -193     | -121         | 10       | -159     | -10             | -253              | -330    | -23         | -3    |
| P2-FO-4   | B3      | 6.0       | 6.0           | 2.75                        | -4216    | 110      | -147         | 12       | -310     | -20             | -492              | -642    | -44         | -5    |
| P2-FO-5   | B3      | 8.0       | 2.9           | 1.00                        | -2017    | -196     | -127         | 7        | -147     | -9              | -235              | -307    | -21         | -2    |
| P2-FO-6   | В3      | 9.0       | 2.5           | 1.25                        | -1747    | 115      | -89          | 6        | -129     | -8              | -205              | -268    | -17         | -3    |
| P2-FO-7   | В3      | 10.0      | 1.3           | 0.75                        | -937     | 97       | -33          | 4        | -69      | -4              | -110              | -143    | -9          | -1    |
| P2-FO-7.1 | В3      | 10.5      | 0.7           | 0.50                        | -495     | 25       | 104          | 2        | -37      | -2              | -59               | -77     | -4          | 2     |
| P2-FO-8   | В5      | 11.0      | 0.9           | 0.50                        | -641     | 37       | -14          | 5        | -48      | -3              | -75               | -98     | -7          | -1    |
| P2-FO-8.1 | В5      | 11.5      | 0.5           | 0.25                        | -331     | -16      | 2            | 2        | -25      | -1              | -40               | -52     | -3          | 0     |
| P3-FO-3   | С       | 12.0      | 3.3           | 1.50                        | -2084    | -308     | -240         | 11       | -171     | -11             | -272              | -356    | -24         | -1    |
| Trachyte  |         | (13.0)    | 26.0          |                             | 0        | 0        | 0            | 0        | 0        | 0               | 0                 | 0       | 0           | 0     |
| (b)       |         |           |               |                             |          |          |              |          |          |                 |                   |         |             |       |
| P1-SA     | B2      | 1.0       | 1.3           | 1.50                        | -556     | 417      | 324          |          | -234     | -241            | -56               | -28     | -5          | -4    |
| P1-SA-2   | B3      | 4.0       | 6.0           | 3.70                        | -3188    | 281      | 172          |          | -1073    | -1107           | -258              | -131    | -26         | -36   |
| P1-SA-3   | B4      | 6.5       | 1.5           | 1.55                        | -630     | 15       | -35          |          | -274     | -279            | -66               | -33     | -6          | -8    |
| P1-SA-4   | B4      | 8.0       | 1.1           | 0.75                        | -492     | -32      | -47          |          | -192     | -195            | -46               | -24     | -4          | -4    |
| P1-SA-5   | С       | 9.5       | 1.8           | 2.20                        | -570     | 20       | -41          |          | -321     | -306            | -78               | -37     | -4          | -5    |
| Basalt    |         | 10.5      | 12.0          |                             | 0        | 0        | 0            |          | 0        | 0               | 0                 | 0       | 0           | 0     |

Table 3.



Fig. 1 (Catena10553)





FIG. 3 (Catena10553)





Fig. 5 (Catena10553)





Fig. 7 (Catena 10553)



FIG. 8 (Catena10553)



Fig. 9 (Catena10553)





FIG. 10 (Catena10553R2)

Supplementary information for the article:

# Differentiated Neogene bauxitization of volcanic rocks (Western Cameroon): Morpho-geological constraints on chemical erosion

Mathieu Nouazi Momo, Anicet Beauvais, Paul Tematio, Martin Yemefack

Supplementary information supplies a figure of the landscapes encompassing the study sites in our article (Fig. DR1a-b). We also provide tables of mineralogical (Table DR1a-b) and geochemical (Table DR2a-b and DR3a-b) compositions of each horizon composing the lateritic weathering profiles formed on trachyte and basalt from the highland western Cameroon. Trace elements transfer rates and transferred masses are also provided (Table DR4a-b) based on calculations from equations (3), (4) and (5) in the associated manuscript.

Figure DR1. Topographic attributes of landscapes on (a) trachytes around Fongo Tongo and (b) basalts near Doumbouo in highlands western Cameroon (Google Earth, image Landsat/ Copernicus Image © 2020 Maxar Technologies).

Table DR1. X ray diffraction mineralogical composition of samples collected in weathering profiles formed on (a) trachyte and (b) basalts. (-) = Mineral present but below calculation limits (< 5%); Ka = kaolinite; Gi = gibbsite; Go = Goethite; An = Anatase; He = Hematite; Ma = Maghemite; Fl = Florencite; Qz =quartz.

Table DR2. Major elements composition of samples collected in weathering profile formed on (a) trachyte and (b) basalt. Major elements in wt.%; LOI=Loss Of Ignition; (-) = below detection limit (0.01 wt.%). The true depth of trachyte being unknown we simply mentioned a depth in parentheses for graphical easiness of figures in the associated article.

Table DR3. Trace elements composition of samples collected in weathering profile formed on (a) trachyte and (b) basalt. Detection limits are indicated.

Table DR4. Trace elements transfer rates, trace elements mass transfers in kg per unit volume (1 m<sup>3</sup>), and transferred masses of elements integrated over the investigated profile horizons

depth ( $\Delta z_w$ ) related to the corresponding height or thickness of chemically weathered rock layer ( $\Delta z$ ), in profile on a) trachyte, and b) basalt.

Supplementary material for on-line publication only Click here to download Supplementary material for on-line publication only: Figure. DR1.jpg

Supplementary material for on-line publication only Click here to download Supplementary material for on-line publication only: DR1\_Catena10553R1.doc

Supplementary material for on-line publication only Click here to download Supplementary material for on-line publication only: DR2\_Catena10553R2.doc

Supplementary material for on-line publication only Click here to download Supplementary material for on-line publication only: DR3\_Catena10550R2.doc

Supplementary material for on-line publication only Click here to download Supplementary material for on-line publication only: DR4\_Catena10553R2.doc