

Mechanistic and ecophysiological study of carbon 14 transfer in fish



Supervisor : Jean-Christophe Poggiale (MIO)
Tutor : Frederic Alonzo (IRSN)

PSE-ENV/SRTE/LECO

SOULOUMIAC Audrey


Context



- Natural sources: reaction of cosmic rays on nitrogen neutrons in the high atmosphere → carbon dating...
- With ${}^3\text{H}$, ${}^{14}\text{C}$ is one of the radionuclides rejected in greatest amounts by Nuclear Power Plant in French rivers.




- Assimilated ${}^{14}\text{C}$ contributes in a major part to estimated – small – doses received by local populations, essentially through fish ingestion.



Need to better assess the environmental (both health and ecological) risks

Problem: Current models used to predict ^{14}C transfer in aquatic ecosystems are very simplistic:


- Unability to explain the variability in ^{14}C activity observed in fish samples collected from different rivers.
- Great variations in ^{14}C releases → No equilibrium in concentration (Sheppard et al., 2006 ; Smith, 2006).
- Models of fish do not take account of the potential influence of fish physiology and environmental factors, depending on species.



→ DEB model is a physiology-based dynamic model = more adequate to describe variation in ^{14}C activity in different compartments of the aquatic ecosystem.

DIB model: Dynamic isotope budget


- Used in ecology to study the dynamics and structure of trophic networks and trace elements fluxes in ecosystems. (Post, David M. 2002. « Using Stable Isotopes to Estimate Trophic Position: Models, Methods, and Assumptions ». *Ecology* 83 (3): 703-18.).




→ Increase in stable isotopes with trophic level

→ DIB model is used for stable isotope analyses

- Aim to quantify dynamics of four biogenic elements (C, H, N and O) and their isotopes within organism by coupling fluxes of elements with those of molecules involved in metabolism.



- 1) Mobilisation : Mobilisation of atoms from a pool of atoms. **Does not change isotopic ratio**



- 2) Molecular selection

Selection for anabolic or catabolic routes. Probability depends on fate of atoms (heavy for catabolic route).



- 3) Molecular reshuffling


The concept of atom reshuffling recognizes that molecules are not completely disassembled into elements during chemical reactions.



Molecular reshuffling



Molecular selection






Molecular reshuffle



Molecular selection





\dot{J}_{EG}






Molecular
reshuffle
Molecular
selection



Structure turnover



Theoretical values of DIB parameters are available for stable isotopes ^{15}N and ^{13}C and are validated in *Crassostrea gigas* (Pacific oyster).

- Parameterize values for the case of ^{14}C in two fish species of fish
- Improvement of DEB model for Common carp (literature search)
- Validation of predictive capacity in Zebrafish



Cyprinus carpio




Danio rerio

Measurements of ^{14}C assimilation, incorporation and elimination kinetics depending the biochemical composition of ^{14}C sources.

Different sources of ^{14}C


Amino acid: Arginine



Glycerol



Glucose





PROTEIN METABOLISM

LIPID METABOLISM


CARBOHYDRATE
METABOLISM

Conception of contaminated food




Preliminary experiment

Contamination duration: range of **1 to 4 days**
Dissection **24h** after last food supply




Results

Mass-specific activity of ^{14}C




Results

Distribution of ^{14}C activity among organs




Proportion of ^{14}C activity among organs




Experimental plan

- ^{14}C kinetics in Common carp




- Model validation on a simple trophic chain




CO₂ traps

Objectif : Quantification of ¹⁴C released through respiration

- To obtain ¹⁴C complete budget among all metabolic processes.
- CO₂ gas traps will be set up with air bubbling in NaOH solution (1M).




Thank for attention!



Scientific communication

- IRSN PhD congress 2018 (Poster) and 2019 (platform)
- Congress of Doctoral school (Aix-Marseille University)(Poster)
- Sixth International Symposium on DEB theory (Brest)



DEB2019 1-12 April 2019 / Brest (France)

Sixth International Symposium and Thematic School
on DEB theory for metabolic organization

Bq/kg C

| Espèces | Min | Max | Moyenne |
|--------------------|-----|------|---------|
| Silure glane | 237 | 334 | 286 |
| Perche | 234 | 931 | 466 |
| Lamproie marine | 247 | | |
| Hotu | 202 | 212 | 207 |
| Grande alose | 249 | | |
| Gardon | 231 | 1262 | 453 |
| Chevesne | 200 | 2000 | 465 |
| Carpe commune | 559 | 581 | 570 |
| Brochet | 234 | 340 | 270 |
| Brème commune | 201 | 2830 | 1138 |
| Barbeau fluviatile | 204 | 2300 | 528 |
| Anguille | 213 | 1393 | 587 |

Catabolique appauvrissement et anabolique enrichissement

The dimensionless reshuffling coefficients (alpha)


1) specify what fraction of N atoms in substrate s ends up in product p in transformation k ($k \frac{1}{4} A, G, L1, L2$).

- Among four compartments: food (X), Feces (P), Reserve (E), structure (V)
- One elemental composition each
 - Stoichiometric equilibrium

Two reactions at a time: Energy for anabolism is brought by catabolism : We detail the catabolic and anabolic routes of each chemical transformation to allow selection of substrate molecules depending on their isotopic composition. The catabolic route of a transformation uses substrates to produce energy for the anabolic route of the transformation. The anabolic route of a transformation uses this energy and substrates as a source of building blocks to produce a given compound. In the growth transformation for instance, reserve is used as a source of energy as well as building blocks to produce structure.

Different sources of ^{14}C


Amino acid: Arginine



Glycerol



Glucose




PROTEIN METABOLISM

LIPID METABOLISM

CARBOHYDRATE METABOLISM

Structure (collagen)



Contraction (myosin)

Transport (albumin)



Immune system
(immunoglobulin)

Mono-, Di- and Tri-glycerids, Phospholipids

Glycogen or
triglycerid
(muscle, liver)