

## Bayesian contributions to radiation dose estimation in biological retrospective dosimetry.

Sophie Ancelet<sup>1</sup>, Gaëtan Gruel<sup>2</sup>, Eric Grégoire<sup>2</sup>, Aurélie Vaurijoux<sup>2</sup>,  
Laurence Roy<sup>3</sup>, Joan Francesco Barquinero Estruch<sup>4</sup>

<sup>1</sup>IRSN, PSE-SANTE/SESANE/LEPID, France (sophie.ancelet@irsn.fr)

<sup>2</sup>IRSN, PSE-SANTE/SERAMED/LRAcc, France

<sup>3</sup>IRSN, PSE-SANTE/SESANE, France

<sup>4</sup>Universitat Autònoma de Barcelona, Facultat de Biociències, Spain

Journée "Big Ideas for Small Data", 13th May 2019

- 1 Introduction
- 2 Data
- 3 Standard approaches
- 4 Bayesian contributions
- 5 Conclusion & Perspectives

# Context

- **Accidents** leading to unplanned exposure of humans to ionizing radiation (IR) have occurred many times
    - **overexposure** in radiotherapy services or occupational settings
    - large-scale **nuclear accidents**
  - **Unclear** radiation exposure scenarios and/or **inconsistent** findings
    - workers at risk of exposure **may not wear** their obligatory personal dosimeter
    - workers at risk of exposure **may not store it** correctly after use.
- Estimation of the **absorbed radiation dose** received by an exposed or suspected exposed individual may be crucial to:
    - Optimize patient-centered care
    - Predict the derived health consequences for both early and late effects
    - Perform rapid triage of exposed versus non-exposed persons
    - Clarify unclear radiation exposure scenarios
    - Appease the "worried well" persons

Dose assessment ⇒ Proof of exposure by court and professional associations










# Biological retrospective dosimetry

- It offers the only possibility to estimate the individual absorbed dose
  - even weeks or months after a potential exposure (Kulka et al. (2018)).
  - when a direct measurement of IR exposure is not or no longer possible

## Main goal

Estimation of the individual absorbed radiation dose from microscope counting of radiation-related **chromosomal anomalies**

- Radiation exposure causes **chromosomal DeoxyriboNucleic Acid (DNA) lesions** like double-strand breaks
- The broken fragments may repair incorrectly ⇒ **Chromosome aberrations**

Atteinte						
Type d'aberrations chromosomiques						
	Restauration	Fragment	Anneau	Inversion	Dicentrique	Translocation

# The dicentric chromosome assay (DCA)

- Dicentrics have a **low naturally occurring background** frequency
- Frequencies of **dicentrics increase** with the absorbed dose  
⇒ Well-established and highly specific **biological marker** of radiation exposure
- Scoring dicentrics in peripheral human blood lymphocytes : **"gold standard" biological method** for retrospective dose estimation (IAEA (2011)).



Photo: Olivier Seignette/Mikaël Lafontan/Médiathèque IRSN



# Main questions

Given the number of dicentrics per cell observed in blood lymphocytes:

## Question Q1

Can it be stated that a **strictly positive radiation dose** has been received by :

- 1 **all of the** analyzed cells (whole-body irradiation)?
- 2 **only a fraction** of the analyzed cells (partial irradiation)?
- 3 **none** of the analyzed cells ? (Relevant for **unclear exposure scenarios**)

# Main questions

Given the number of dicentrics per cell observed in blood lymphocytes:

## Question Q1

Can it be stated that a **strictly positive radiation dose** has been received by :

- 1 **all of the** analyzed cells (whole-body irradiation)?
- 2 **only a fraction** of the analyzed cells (partial irradiation)?
- 3 **none** of the analyzed cells ? (Relevant for **unclear exposure scenarios**)

## Question Q2

What is the **estimated absorbed dose** and the **uncertainty associated** to this estimation?



- 1 Introduction
- 2 Data**
- 3 Standard approaches
- 4 Bayesian contributions
- 5 Conclusion & Perspectives








# 4 real radiation accident victims (2006-2013)

## In-vivo data provided by IRSN/LRAcc



Id	Circumstances of accident	Clinical signs	Physical dosimetry
06-11	<u>Exposure to <math>\gamma</math>-rays</u>	Vomiting (4h30), nausea, hair loss, Lymphocytes: $0.8 \times 10^{-3}$	No
11-08	<u>Medical context; 10 minutes located next to a <math>\gamma</math>-source (Co 60)</u>	<u>Hematopoietic syndrome 7 days after exposure</u>	No
08-03	Put the $\gamma$ -source (Ir) in his hand then in his pocket (10 minutes to 1 hour) -> Hand burn	lymphocytes: $1.05 \times 10^{-3}$	0.25 Sv
05-03	Exposure head and chest : 15-30 seconds Shoulders 5cms away from the X source Neck 20cms away from the X source	Erythema (collarbone) Lymphocytes: $2.39 \times 10^{-3}$	0.045 Sv



# 8 suspected exposed individuals (2006-2013)

## In-vivo data provided by IRSN/LRAcc

Justifier le texte




## 4 real suspected individuals (2006-2013)

### From IRSN/LRAcc

Id	Circumstances of accident	Clinical signs	Physical dosimetry
06-63	Exposure to $\gamma$ -rays (10-15 minutes)	No	No
06-70	Spent the night 25 centimeters away from a $\gamma$ -source	No	No
06-13	Colleague of 06-11	No	No
06-15	Colleague of 06-11	No	No

### Conventional cytogenetics








For some of them, no dicentric was observed...

# Calibration data (Cobalt 60) - In-vitro data provided by IRSN/LRAcc

In-vitro irradiation of blood samples - various healthy donors - different doses

Number of analyzed cells	Dose (Gray)	Number of dicentric
19194	0	21
1676	0.05	3
1552	0.10	6
481	0.15	3
1057	0.24	11
1768	0.30	38
1187	0.33	18
2919	0.50	83
1538	0.80	100
869	1	90
1525	1.6	269
1844	2	545
352	2.31	122
784	3	482
534	4	521
341	4.70	381
94	5.77	143



- 1 Introduction
- 2 Data
- 3 Standard approaches**
- 4 Bayesian contributions
- 5 Conclusion & Perspectives

## Dose-response model $\mathcal{M}_A$ for in-vivo data

### Exposed and suspected exposed individuals

Let's consider a given individual with  $n_0$  analyzed cells:

- $D_0$  : **Unknown** absorbed dose (in Gray) received by each cell
- $R_k$ : Number of dicentrics observed in each cell  $k$  ( $k = 1, \dots, n_0$ )

#### In case of LOW-LET radiation and homogeneous irradiation

$$(\mathcal{M}_A) \quad R_k \sim^{i.i.d} \text{Poisson}(\lambda_0)$$

$$\lambda_0 = A + \alpha D_0 + \beta D_0^2$$

- $\theta = (A, \alpha, \beta)$ : **unknown parameters** with  $A > 0$ ,  $\beta > 0$ ,  $\alpha > -2\sqrt{A\beta}$
- $A$ : background expected number of dicentrics per cell at dose  $D_0 = 0$
- $Y_0 = \sum_{k=1}^{n_0} R_k \sim \text{Poisson}(n_0 \lambda_0)$

**Non-identifiable model**  $\Rightarrow$  External data required to estimate  $\theta = (A, \alpha, \beta)$

## Dose-response model $\mathcal{M}_C$ for calibration data

Let's consider a given experimental (in-vitro) irradiation  $i \in \{1, \dots, I\}$

- $D_i$ : **Fixed** absorbed dose (in Gray) received by each cell
- $Z_{i,l}$ : Number of dicentrics observed in **each cell**  $l \in \{1, \dots, n_i\}$  at dose  $D_i$

In case of LOW-LET radiation and homogeneous irradiation

At a given dose  $D_i$ :

$$(\mathcal{M}_C) \quad Z_{i,l} \sim^{i.i.d} \text{Poisson}(\lambda_i)$$

$$\lambda_i = A + \alpha D_i + \beta D_i^2$$

$$\Rightarrow Y_i = \sum_{l=1}^{n_i} Z_{i,l} \sim \text{Poisson}(n_i \lambda_i)$$

where  $Y_i$  is the total number of dicentrics observed at dose  $D_i$  and  $n_i$  the total number of analyzed cells

## Answering $Q_2$ - Estimation of the dose

- Fit  $\mathcal{M}_C$  to **calibration data** using maximum likelihood estimation
- Plug  $\hat{\theta} = (\hat{A}, \hat{\alpha}, \hat{\beta})$  into  $\mathcal{M}_A$
- Derive point estimate  $\hat{D}_0$  of the absorbed dose  $D_0$  (inverse regression)

$$\hat{D}_0 = g(\hat{A}, \hat{\alpha}, \hat{\beta}) = \frac{-\hat{\alpha} + \sqrt{\hat{\alpha}^2 + 4\hat{\beta}(\hat{\lambda}_0 - \hat{A})}}{2\hat{\beta}}$$

where  $\hat{\lambda}_0 = \frac{Y_0}{n_0}$

# Answering $Q_2$ - Estimation of the dose

Id	Circumstances of accident	MLE for the dose $D_0$	Id	Circumstances of accident	MLE for the dose $D_0$
06-11	Exposure to $\gamma$ -rays	4.40	06-13	Colleague of 06-11	0.02
11-08	Medical context; 10 minutes located next to a $\gamma$ -source (Co 60)	1.88	06-14	Colleague of 06-11	0.02
08-03	Put the $\gamma$ -source (Ir) in his hand then in his pocket (10 minutes to 1 hour) -> Hand burn	0.23	06-15	Colleague of 06-11	-0.03
05-03	Exposure head and chest : 15-30 seconds Shoulders 5cms away from the X source Neck 20cms away from the X source	0.11	06-16	Colleague of 06-11	0.02
06-63	Exposure to $\gamma$ -rays (10-15 minutes)	0.15	04-14	Positive dosimeter	-0.03
06-70	Spent the night 25 centimeters away from a $\gamma$ -source	0.25	13-09	Positive dosimeter	-0.03

## Potential drawbacks:

- If  $\hat{\lambda}_0 = \frac{Y_0}{n_0} = 0$  then  $\hat{D}_0 < 0$  (Context: Small signal in the data)
- Prior information on the dose not accounted for
- Modular approach : Disjoint estimation of  $\theta$  and  $D_0$



## Answering $Q_2$ - Derive a 95% confidence interval on $\hat{D}_0$

- Approach 1: **Multivariate delta-method**

$$\begin{aligned} \sigma_{\hat{D}_0}^2 &= \sigma_{\hat{A}}^2 \left( \frac{\partial g}{\partial A} \right)_{A=\hat{A}}^2 + \sigma_{\hat{\alpha}}^2 \left( \frac{\partial g}{\partial \alpha} \right)_{\alpha=\hat{\alpha}}^2 + \sigma_{\hat{\beta}}^2 \left( \frac{\partial g}{\partial \beta} \right)_{\beta=\hat{\beta}}^2 + \sigma_{\hat{\lambda}_0}^2 \left( \frac{\partial g}{\partial \lambda_0} \right)_{\lambda_0=\frac{Y_0}{n_0}}^2 \\ &+ 2 \left( \frac{\partial g}{\partial A} \right)_{A=\hat{A}} \left( \frac{\partial g}{\partial \alpha} \right)_{\alpha=\hat{\alpha}} \text{cov}(\hat{A}, \hat{\alpha}) + 2 \left( \frac{\partial g}{\partial \alpha} \right)_{\alpha=\hat{\alpha}} \left( \frac{\partial g}{\partial \beta} \right)_{\beta=\hat{\beta}} \text{cov}(\hat{\alpha}, \hat{\beta}) \\ &+ 2 \left( \frac{\partial g}{\partial A} \right)_{A=\hat{A}} \left( \frac{\partial g}{\partial \beta} \right)_{\beta=\hat{\beta}} \text{cov}(\hat{A}, \hat{\beta}) \end{aligned}$$

⇒ **Asymptotical 95% confidence interval** on dose estimate:  $\hat{D}_0 \pm 1.96\hat{\sigma}_{D_0}$

- Approach 2: **Bootstrap**

### Potential drawbacks:


- Is the asymptotic assumption correct?
- Bootstrap ⇒ Strong data redundancy if small signal in data
- Uncertainty on the dose estimation may depend on the statistical method used to compute the confidence interval

# Answering Q1 - Strictly positive absorbed dose received?

**Hypothesis testing:**  $H_0 : D_0 = 0$  vs  $H_1 : D_0 = d_1$  (with  $d_1 > 0$ )

- Test statistic:  $Y_0 = \sum_{k=1}^{n_0} R_k$
- Under  $H_0$ ,  $Y_0 \sim \text{Poisson}(n_0 A)$
- Critical region:  $[y_0^*, +\infty]$  with  $y_0^* = 0.95$  quantile of  $\text{Poisson}(n_0 \hat{A})$ 
  - $y_0^*$  is called "Decision threshold"
- If  $y_0^{obs} > y_0^*$ ,  $H_0$  is rejected with error (of the first kind) = 0.05
- Statistical power:  $1 - \text{Frd}_{H_1}(y_0^*)$  where  $\text{Frd}_{H_1}$  cumulative distribution function of a Poisson distribution with intensity =  $n_0(\hat{A} + \hat{\alpha}d_1 + \hat{\beta}d_1^2)$ 
  - Detection Limit: The smallest value of dose  $d_1$  from which the statistical power of the test is greater or equal to 0.95

# Answering Q1 - Strictly positive absorbed dose received?



# Answering Q1 - Strictly positive absorbed dose received?

Id	$n_0$	$y_0$	$y_0^*$	DL		Id	$n_0$	$y_0$	$y_0^*$	DL
06-11	139	155	1	0.51		06-13	602	1	2	0.19
11-08	451	112	2	0.25		06-14	500	1	2	0.23
08-03	1024	13	3	0.14		06-15	505	0	2	0.22
05-03	500	3	2	0.23		06-16	508	1	2	0.22
06-63	500	4	2	0.23		04-14	503	0	2	0.23
06-70	356	5	2	0.30		13-09	507	0	2	0.22

DL = Detection Limit

## Potential drawbacks:

- Binary answer to  $Q_1$ : Rejection of  $H_0$  or not
- $D_0$  is unknown ! : Statistical power?
- **The statistical power** may be very small for small doses  $D_0$ ...
- Uncertainty on the estimation of the background expected number of dicentrics per cell  $A$  not accounted for
- Does not allow to test if **only a fraction** of the analyzed cells have received a strictly positive radiation dose

## Aim of the work

- Can **Bayesian statistical methods** offer relevant alternative answers to questions  $Q_1$  and  $Q_2$  in biological retrospective dosimetry ?
- To account for **expert knowledge** when assigning a prior distribution on the unknown absorbed dose  $D_0$
- To propose a **unique, flexible and coherent framework** allowing to simultaneously answer to questions  $Q_1$  and  $Q_2$




- 1 Introduction
- 2 Data
- 3 Standard approaches
- 4 Bayesian contributions**
- 5 Conclusion & Perspectives

# Which model?

Approach 1: the one previously described....

**Directed Acyclic Graph of the full model ( $\mathcal{M}_A + \mathcal{M}_C$ )**



- $\theta = (A, \alpha, \beta)$ : shared parameters
- Possibility for the in-vivo data to be accounted for when fitting  $A, \alpha, \beta$
- The Bayesian framework allows fitting this model in one step

# The prior distributions

- $A \sim Unif[0, +\infty[$
- $\alpha \sim Unif[-2\sqrt{A\beta}, +\infty[$
- $\beta \sim Unif[0, +\infty[$
- Prior probability distribution on  $D_0$ 
  - $D_0 \sim Unif(0, 10) \Rightarrow$  Vague prior
  - $D_0 \sim Gamma(a, b) \Rightarrow$  Informative prior



# Using expert knowledge to define an informative Gamma prior $D_0$


- Hyperparameters  $a$  and  $b$  of the Gamma prior may be fixed by expert knowledge given the accident scenario

Id	Circumstances of accident	Clinical signs	Physical dosimetry	Prior distribution on $D_0$
06-11	Exposure to $\gamma$ -rays	Vomiting (4h30), nausea, hair loss, Lymphocytes: $0.8 \times 10^{-3}$	No	$D_0$ .median=2.5 $D_0$ max = 10 (q99-10) <b><math>D_0</math>-Gamma(<math>a=1.98</math> , <math>b=0.66</math>)</b>
11-08	Medical context; 10 minutes located next to a $\gamma$ -source (Co 60)	Hematopoetic syndrom 7 days after exposure	No	$D_0$ .median=2.5 $D_0$ max = 10 (q99-10) <b><math>D_0</math>-Gamma(<math>a=1.98</math> , <math>b=0.66</math>)</b>
08-03	Put the $\gamma$ -source (Ir) in his hand then in his pocket (10 minutes to 1 hour) -> Hand burn	lymphocytes: $1.05 \times 10^{-3}$	0.25 Sv	$D_0$ .median=0.25 $D_0$ max = 5 (q99-5) <b><math>D_0</math>-Gamma(<math>a=0.4</math> , <math>b=0.6</math>)</b>
05-03	Exposure head and chest : 15-30 seconds Shoulders 5cms away from the X source Neck 20cms away from the X source	Erythema (collarbone) Lymphocytes: $2.39 \times 10^{-3}$	0.045 Sv	$D_0$ .median=0.045 $D_0$ max = 5 (q99-5) <b><math>D_0</math>-Gamma(<math>a=0.2</math> , <math>b=0.44</math>)</b>

For individuals for which no clinical sign was observed:  $D_0 \sim Unif(0, 2)$


⇒ **Not enough informative ! To improve!**

# Using expert knowledge to define an informative Gamma prior on $D_0$









# Answering $Q_2$ - Bayesian estimation of the dose

MCMC algorithm - Package R "rjags"



GUM= Multivariate Delta-Method

# Answering $Q_2$ - Bayesian estimation of the dose



# BUT...

Given the prior distribution assigned to  $D_0$ , we are assuming that  $D_0 > 0$

⇒ **Is this assumption relevant for all the considered individuals?**



# Answering $Q_1$ and $Q_2$ under the Bayesian framework

## Question $Q_1$

Can it be stated that a **strictly positive radiation dose** has been received by :

- 1 **all of the** analyzed cells (whole-body irradiation)?
- 2 **only a fraction** of the analyzed cells (partial irradiation)?
- 3 **none** of the analyzed cells ? (Relevant for **unclear exposure scenarios**)










The **above sub-questions 1 and 3** can be formalized as :


## A Bayesian model selection problem


$$\mathcal{M}_0 : R_k \sim^{i.i.d} \text{Poisson}(A) \quad \text{vs} \quad \mathcal{M}_A : R_k \sim^{i.i.d} \text{Poisson}(A + \alpha D_0 + \beta D_0^2)$$

given in-vivo data **and** calibration data following model  $\mathcal{M}_C$  ( $D_0 > 0$ )

# Answering $Q_1$ and $Q_2$ under the Bayesian framework



  $Poisson(A)$

  $Poisson(A + \alpha D_0 + \beta D_0^2)$  with  $D_0 > 0$



Bayesian inference		
	Post.mean	Credible interval at 95%
A	0.001	[0,0006; 0,0015]
$\alpha$	0.040	[0,032; 0,048]
B	0.048	[0,044; 0,052]

# Answering $Q_1$ and $Q_2$ under the Bayesian framework

⇒ A Bayes factor (Jeffreys, 1939) can be efficiently approximated (e.g., Monte-Carlo estimate)

But what about **sub-question 2 about partial irradiation?**



## Idea: using a mixture model (Kamary et al. (2014) - arXiv)

Let's consider a given individual - potentially exposed - with  $n_0$  analyzed cells:


- $p_0$ : **unknown** probability for each cell to have received a dose  $> 0$
- $D_0$ : **unknown** absorbed dose (in Gray) received by each irradiated cell

### A mixture model for in-vivo data (LOW LET + homogeneous irradiation)

$$\mathcal{M}_{mix} : R_k \sim^{i.i.d} (1 - p_0)Poisson(A) + p_0Poisson(A + \alpha D_0 + \beta D_0^2)$$

- $D_0 > 0$  and  $p_0 \in [0, 1]$
- $\theta = (A, \alpha, \beta)$ : unknown parameters with  $A > 0$ ,  $\beta > 0$ ,  $\alpha > -2\sqrt{A\beta}$
- $A$ : **common parameter** shared by both mixture components
- $p_0$  can also be interpreted as the proportion of irradiated cells
- $D_0$  and  $p_0$  assumed to be identical for each irradiated cell
- $\mathcal{M}_0$  and  $\mathcal{M}_A$  are very special cases of the mixture model

# Directed Acyclic Graph of the full model ( $\mathcal{M}_{mix} + \mathcal{M}_C$ )



- $\theta = (A, \alpha, \beta)$ : shared parameters
- The Bayesian framework allows fitting this model in one step



## Answering to $Q_1$ and $Q_2$ with $\mathcal{M}_{mix}$ (1/2)

- If  $p_0 = 0$ , model  $\mathcal{M}_0$  is selected given the available count data
  - $\Rightarrow$  Response to  $Q_1$  is NO= "There is no evidence that a strictly positive radiation dose has been received".
- If  $p_0 = 1$ , model  $\mathcal{M}_A$  is selected given the available count data
  - $\Rightarrow$  Response to  $Q_1$  is YES= "A strictly positive radiation dose has been received **by all the analyzed cells**".
- If  $p_0 \in ]0, 1[$ , neither model  $\mathcal{M}_0$  nor model  $\mathcal{M}_A$  is selected given the available count data
  - $\Rightarrow$  Response to  $Q_1$  is YES= "A strictly positive radiation dose has been received **BUT only by a fraction of the analyzed cells**" (**partial body exposure**).
  - The fraction of the body irradiated is defined as (IAEA report 2001):

$$F_0 = \frac{p_0 \times \exp(D_0/\tilde{D})}{(1 - p_0) + p_0 \times \exp(D_0/\tilde{D})} \quad \tilde{D} \sim Unif(2.7, 3.5)$$

## Answering to $Q_1$ and $Q_2$ with $\mathcal{M}_{mix}$ (2/2)

- Posterior distribution on  $p_0 \Rightarrow$  Probabilistic answer to  $Q_1$
- $\Rightarrow$  Decision criterion to define the range of acceptance, rejection and indecision conclusions



- Let's  $c_1, c_2, U$  be fixed decision thresholds (to calibrate by simulation)
- Compute  $\pi_1 = P(p_0 > c_1 | Y_i, R_k)$  and  $\pi_2 = P(p_0 < c_2 | Y_i, R_k)$ 
  - If  $\pi_1 > U \Rightarrow$  YES= "There is **strong evidence** that a strictly positive radiation dose has been received **by all of the analyzed cells**".
  - If  $\pi_2 > U \Rightarrow$  NO= "There is no evidence that a strictly positive radiation dose has been received".
  - Else YES= "A strictly positive radiation dose has been received **BUT only by a fraction of the analyzed cells**" (partial body exposure).

# The prior distributions

- $A \sim Unif[0, +\infty[$
  - $\alpha \sim Unif[-2\sqrt{A\beta}, +\infty[$
  - $\beta \sim Unif[0, +\infty[$
  - $D_0 \sim Gamma(a, b)$  or  $D_0 \sim Unif(0, 10)$
  - $p_0 \sim Beta(c, d)$
- 
- Hyperparameters a,b,c,d may be fixed by expert knowledge given the accident scenario
  - Default choice (Rousseau and Mengersen (2011)):  $c=0.5, d=0.5$

# Bayesian inference

## Adaptive Metropolis-Hastings algorithm

- Block updating for  $(A, \alpha, \beta)$  using a Gaussian random walk (20% acceptance rate)
- Gaussian random walk for  $D_0$  (40% acceptance rate)
- For the mixture weight  $p_0$ :
  - Iteration  $t$ : Independent proposal  $\Rightarrow p_0^{cand} \sim \text{Beta}(0.5, 0.5)$
  - Iteration  $t+1$ : Random walk  $\Rightarrow p_0^{cand} \sim \text{Beta}(1 + p_0^t, 2 - p_0^t)$
  - 40% acceptance rate
- Implemented in Python (2.7.10) (100000 iterations = 30 seconds)

## Asymptotic consistency of the proposed mixture testing procedure

- Proved by Kamary et al. (2014) in the specific case of embedded mixture components
  - "If one model is indeed correct, the posterior medians of the corresponding weight in the mixture settles very quickly near the boundary values of 1 as the sample size increases"

## Remark

- Equivalent formulation of  $\mathcal{M}_{mix}$  pointing out the latent allocation variables






$$\mathcal{M}_{mix} : \quad R_k \sim^j \text{Poisson}(\lambda_k) \quad \text{with} \quad \lambda_k = A + \alpha D_{0k} + \beta D_{0k}^2$$

$$D_{0k} = \gamma_k \times D_0 \quad \text{with} \quad \gamma_k \sim \text{Bern}(p_0)$$

- Easy implementation in WinBUGS or JAGS but inefficient Gibbs sampler!!!

# Convergence diagnostics on the weight $p_0$

Gibbs sampler (Left) vs Adaptive Metropolis-Hastings (Right)











# Posterior statistics, Bayes factor and posterior probability of $\mathcal{M}_1$

	Bayesian Mixture approach					Bayes Factor $\mathcal{M}_1$ vs $\mathcal{M}_0$  (Kass & Raftery (1995))	$P(\mathcal{M}_1   y)$
	<i><math>D_0</math> posterior median 95%CI</i>	<i><math>p_0</math> posterior median 95%CI</i>	<i><math>F_0</math> posterior median 95%CI</i>	<i><math>P(p_0 &gt; 0.8)</math></i>	<i><math>P(p_0 &lt; 0.2)</math></i>		
06-11	4.61 [4.14; 5.19]	0.91 [0.76; 1.00]	0.97 [0.90; 1.00]	0.93	0.0	$+\infty$ (very strong)	1 [1.0; 1.0]
11-08	2.09 [1.76; 2.69]	0.84 [0.56; 1.00]	0.90 [0.69; 1.00]	0.60	0.0	1.75 <sup>c</sup> +185 (very strong)	1 [1.0; 1.0]
08-03	0.32 [0.15; 1.25]	0.67 [0.10; 1.00]	0.69 [0.11; 1.00]	0.39	0.11	$>10^7$ (very strong)	1 [1.0; 1.0]
05-03	0.13 [0.0002; 1.29]	0.54 [0.011; 1.0]	0.55 [0.01; 1.0]	0.31	0.25	4 (Positive)	0.67 [0.63; 0.70]
06-63	0.47 [0.08; 1.84]	0.23 [0.02; 0.99]	0.26 [0.02; 0.99]	0.16	0.46	8.3 (Positive)	0.86 [0.83; 0.88]
06-70	0.55 [0.16; 1.84]	0.36 [0.04; 1.00]	0.40 [0.06; 1.00]	0.21	0.33	303.03 (Very Strong)	1.00 [1.0; 1.0]




# Comparison of dose estimations



Posterior medians + 95% credible intervals  
 ISO2014 = Multivariate Delta Method

# Comparing prior and posterior probability distributions on $p_0$

Prior probability distribution on  $p_0$  : Beta(0.5,0.5)




From left to right : Victims 06-11 (Estimated dose: 4.61 Gy), 08-03  
(Estimated dose: 0.32Gy), 05-03 (Estimated dose: 0.13Gy)


- Weak influence of the prior choice on  $D_0$  (results not shown)
- Lack of information in the data to infer  $p_0$  especially when dose is small  
⇒ **More data needed to infer  $p_0$  (and then answer  $Q_1$ )?**

# Sensitivity to the prior choice on $p_0$

- Informative Beta priors defined from expert knowledge
- Jeffrey's prior Beta(0.5,0.5)



Posterior distribution on the dose  $D_0$



Posterior distribution on the weight  $p_0$

⇒ Sensitivity is clearly present but should naturally vanish as the number of analyzed blood lymphocytes increases

- 1 Introduction
- 2 Data
- 3 Standard approaches
- 4 Bayesian contributions
- 5 Conclusion & Perspectives**

# Conclusions

- **First fully Bayesian approach** proposed to simultaneously answer to two main questions of interest in biological retrospective dosimetry
  - ⇒ New insights to the **European Radiation Dosimetry (EURADOS)** Working Group 10, task 10.6
- Using the proposed mixture model  $\mathcal{M}_{mix}$  allows to get **rich probabilistic answers** to questions  $Q_1$  and  $Q_2$ 
  - ⇒ Relevant input data for **decision-making** in the contexts of clinical management of patients, rapid triage after large-scale radiation incident, reassuring the 'worried-well'...
- In case of low suspected dose, the number of analyzed blood lymphocytes **should be higher** to obtain more precise answers to question  $Q_1$

# Perspectives

- **Simulation studies** to validate the whole methodology and calibrate the decision thresholds  $(c_1, c_2, U)$
- Validate the whole methodology from new experimental data for which  $D_0$  and  $p_0$  are known
- Bayesian optimal design to define the number of analyzed cells  $n_0$  required to optimally answer to question  $Q_1$  and  $Q_2$  under budget constraint
- **Extend** the proposed approach to other chromosome aberrations
- Provide **operational tools** to dosimetrists

# Bibliography

- [1] IAEA 2011 report (2011) Cytogenetic dosimetry: applications in preparedness and response to radiation emergencies. International Atomic Energy Agency: Vienna.
- [2] Merkle W. (1983) Statistical Methods in Regression and Calibration Analysis of Chromosome Aberration Data. Radiat Environ Biophys. 1:217-233
- [3] Ainsbury et al. (2014) Review of Bayesian statistical analysis methods for cytogenetic radiation biodosimetry with a practical example. Radiation Protection Dosimetry. 162(3):185-96
- [4] Higuera et al. (2016) A new Bayesian model applied to cytogenetic partial body irradiation estimation. Radiation Protection Dosimetry. 168(3):330-6
- [5] Kamary K., Mengersen K., Robert CP., Rousseau J. (2014) Testing hypotheses via a mixture estimation model. ArXiv:1412.2044v2