



HAL
open science

A linearly polarized electromagnetic wave as a swarm of photons half of which have spin -1 and half of which have spin $+1$

Gerrit Coddens

► To cite this version:

Gerrit Coddens. A linearly polarized electromagnetic wave as a swarm of photons half of which have spin -1 and half of which have spin $+1$. 2020. hal-02636464v2

HAL Id: hal-02636464

<https://hal.science/hal-02636464v2>

Preprint submitted on 1 Jun 2020 (v2), last revised 9 Jun 2020 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A linearly polarized electromagnetic wave as a swarm of photons half of which have spin -1 and half of which have spin $+1$

Gerrit Coddens

Laboratoire des Solides Irradiés,
Institut Polytechnique de Paris, UMR 7642, CNRS-CEA- Ecole Polytechnique,
28, Route de Saclay, F-91128-Palaiseau CEDEX, France

27th May 2020

Abstract. We illustrate our solution for the particle-wave duality and the paradox of the cat of Schrödinger on the example of the photons which occur in a linearly polarized electromagnetic wave. The $+1$ and -1 spin states correspond to the left and right circular polarizations, as pointed out by Raman [1].

PACS. 03.65.-w, 03.65.Ca Quantum Mechanics

1 Schrödinger's cat and the particle-wave duality

A stubborn problem in Quantum Mechanics ¹ is the particle-wave duality, whereby it is proposed that a physical object like a photon or an electron can be both a particle and a wave, and that how it behaves depends on the experimental set-up. This is frankly incomprehensible, such that it has been often qualified as a quantum mystery. As this way of interpreting quantum mechanics is not satisfactory I thought we should find something more rational.²

In this document we will abandon the geometrical interpretation that electrons and photons would be both particles and waves and replace it by something that is more simple and free of mysteries. We will also provide a different geometrical interpretation for a wave function which is a superposition state of the type which occurs in the paradox of the cat of Schrödinger. Again, this will be much easier to make sense of.

The example of the double-slit experiment (e.g. for electrons) shows that the electrons reach the detector one by one as particles. The detector could e.g. be a photographic plate and one sees dot after dot appearing on the plate [2]. Therefore electrons are definitely particles. Gradually, after zillions of electrons have travelled through the double-slit the interference pattern appears. Contrary to what you have been told, this does not show that electrons are waves. It shows that the large number of electrons that have travelled through the set-up behaves like a wave. You should just sit down and reflect calmly

¹ In order to avoid all possible confusion I want to clarify the working assumptions underlying the present document. First of all we are fully convinced of the fact that quantum mechanics and its algebraic rules to make the calculations are perfectly correct. To explain the fundamental idea behind my endeavour it is perhaps good to consider an analogy. In algebraic geometry we define a correspondence between algebra and geometry, whereby everything in the geometry can be translated in a one-to-one fashion into algebra and vice versa. In quantum mechanics we perfectly know the algebra but the question how we should interpret it has not been settled. We do not know the geometry. In our analogy, we are looking for a geometry that would correspond to a more intuitive interpretation of quantum mechanics. We are thus not questioning quantum mechanics but only the way how it should be geometrically interpreted.

² One could compare this particle-wave duality to what is claimed in the Christian religion: There is a Father, a Son and a Holy Ghost, but there is only one God and this is a mystery. Dogmatic mystery is unacceptable in science. In fact, introducing such mysteries is a form of brainwashing. The person confronted with the contradictory "information" is startled and plunged into cognitive dissonance, because it is impossible to make sense of the contradiction which is imposed with authority as true. Taking advantage of this destabilization and the associated feeling of alienation, it becomes then easy to push him over into surrender mode. All this is typical of how authoritarian regimes install domination. In physics, the argument that a theory agrees with experiment is abusive and authoritarian when that theory is mathematically or logically ramshackle. Authority serves here to silence the questions the teacher himself cannot answer, because the problem of the meaning of quantum mechanics has never been settled. The teacher is now controlling the student's mind and the indoctrination can begin. Difficult challenging questions will no longer be asked, because the student has been made feel insecure and stupid. His only salvation to pass the examination will be to learn the stuff by heart and repeat it like a parrot. It is the better students who suffer from this the most.

about what I am telling you here to realize how you have been fooled with a leading argument. It is not the single electrons which are waves, it is the ensemble of electrons you have measured which looks like a wave. There is no need for theory to see that. All we need is serene unprejudiced observation. The conclusion is thus that electrons are particles and that it is the statistical ensemble of particles you have used in the experiment that behaves like a wave. This wave comes about without any interaction between the electrons, because one can assure that there is only one electron traveling through the set-up at the time. This is in marked contrast with a water wave, where the water molecules are also particles and not waves, but the macroscopic wave comes about by the interactions between these molecules. There must thus exist a phase relation between those electrons due to their initial conditions because you assume such a coherence for the wave.³ There is then no longer a mystery. A wave function does thus never describe a single particle. It describes a statistical ensemble of particles. The wave function is a probability amplitude and probabilities are defined by considering the outcome of the experiment for a statistical ensemble of a large number of particles.⁴

It is therefore time to stop being cruel with cats. The wave function of Schrödinger's cat does not describe a cat that is half alive and half dead. There are no such cats, because the wave function does not describe a single cat. It describes a statistical ensemble of cats. In this ensemble of cats, half of them are alive and half of them are dead. There is even a group theoretical justification for this solution of the paradox. There is a well-known argument due to Ehrenfest that a superposition state $\sum_j c_j \psi_j$ describes an ensemble of particles whereby $|c_j|^2$ are in the state ψ_j . We will now give a group-theoretical argument to justify this for electrons. We propose to describe the spinning motion of the electron by spinors of SU(2) as described in [4]. In a group G , only one operation \circ is defined, which one most of the time calls the product. In SU(2) the group is non-abelian and its 2×2 matrices represent three-dimensional rotations. The product of two group elements is defined, the sum is not. This is also true for rotations and especially for their representation matrices in SU(2). But in group theory, one nevertheless constructs all-commuting operators of a non-abelian group G . One considers then all operators of a same conjugacy class of elements $h' = g \circ h \circ g^{-1}$. Here h' and h are of the same conjugacy class, and g runs through G . Members of the same conjugacy class C are operations of the same geometrical type. E.g. they are rotations over the same angle but about different axes. The similarity transformation corresponds to a change of axis. An all-commuting operator is then defined as $S = \sum_{h \in C} h$. This is purely formal as the sum of group elements is not defined. And in textbooks one argues then $\forall k \in G : k \circ S = S \circ k$. It is a kind of annoying that this is purely formal, but it can be justified by proposing that $k \circ S = S \circ k$ stands in reality for $k \circ C = C \circ k$, because the reasoning that occurs in both proofs is exactly the same. The latter identity is fully defined ($k \circ C = \{k \circ g \mid g \in C\}$) and exact.

This shows that a sum of group elements can be interpreted as representing a set \mathcal{S} of group elements (in the example this is the conjugacy class C). By isomorphism, one can transpose this to the representation matrices $\mathbf{D}(g)$. Then the sum $\sum_{g \in \mathcal{S}} \mathbf{D}(g)$ stands for $\{\mathbf{D}(g) \mid g \in \mathcal{S}\}$. And we will have $\mathbf{D}(k) [\sum_{g \in \mathcal{S}} \mathbf{D}(g)] = [\sum_{g \in \mathcal{S}} \mathbf{D}(g)] \mathbf{D}(k)$, when $k \circ \mathcal{S} = \mathcal{S} \circ k$. One can now do the same with two electrons described by spinors ψ_1 and ψ_2 of SU(2). The 2×1 spinors ψ_j are stenographic notations for the 2×2 representation matrices $\mathbf{D}(g_j)$ of group elements g_j [5]. Hence we replace $\mathbf{D}(g_1)|\psi_1$ and $\mathbf{D}(g_2)|\psi_2$, but we can then no longer multiply to the right. And we have now $\psi_1 + \psi_2$ as a representation of $\{g_1, g_2\}$. If there are 3 electrons that can be described by g_1 and 2 electrons that can be described by g_2 , then $\{g_1, g_1, g_1, g_2, g_2\}$ (whereby the equal elements are tagged) would describe this set of electrons and thus also $3\psi_1 + 2\psi_2$. But in quantum mechanics there is now a twist, based on the

³ Because that is what you do when you are solving a wave equation and impose the boundary conditions. These boundary conditions impose coherence. In the main text we will consider a situation that is different from the one in this footnote, which addresses the solution of wave equations, where one sets the phase at the source everywhere zero. We could e.g. consider plane waves impinging on a set-up, and then everywhere on the plane of the source, we would attribute the same phase to the electron by the boundary condition. This is not self-evident because the source may be incoherent. This shows that in this kind of problem the phase itself is not important and that it is the phase difference built up starting from the source which counts. That is then the reason why we can act as though the source were coherent. It is this phase relation imposed at the source which applies then to all electrons. In fact, we can put all initial phases equal to zero and keep in mind the error this induces. Then we can correct for this error again at the end. (If this were to be incorrect because e.g. the electron has undergone a spin flip during its trajectory, then the coherence of the wave solution would have been lost anyway). As what we finally measure is $|\psi|^2 = \psi^\dagger \psi$ or $|\psi|^2 = \psi^* \psi$ this will not change the experimental results. What counts in the end is the amplitude of the wave, not its phase. This may sound absurd, because when you have two particles with wave functions ψ_1 and ψ_2 then correcting their phases by $\psi_1 \rightarrow \psi_1 e^{i\alpha_1}$ and $\psi_2 \rightarrow \psi_2 e^{i\alpha_2}$, might destroy the interference in $\psi_1 + \psi_2$: $|\psi_1 + \psi_2|^2 \neq |\psi_1 e^{i\alpha_1} + \psi_2 e^{i\alpha_2}|^2$. However, the idea that you should add up two wave functions $\psi_1 + \psi_2$ and then square the result, is wrong and never applies. It belongs to the brainwash (see the following part about group theory in the main text). Each particle must be counted in its own right, because the particles appear as dots on the screen one by one. Hence we must always determine the quantities of single particles according to $|\psi_1|^2 + |\psi_2|^2$, never according to $|\psi_1 + \psi_2|^2$. When there is an interference pattern the wave functions ψ_1 and ψ_2 of two particles obey and comply already with the number of particles implied by the amplitude of the interference term $|\psi_L + \psi_R|^2$ that you would write down according to the textbook rule, where R and L refer to the left and the right slit, see Eq. 4 of Ref. [3]. E.g. if there were destructive interference $\psi_L(\mathbf{r}) + \psi_R(\mathbf{r}) = 0$ there would be just no particles with wave function ψ_1 and ψ_2 present in \mathbf{r} , such that the coefficients $c_1(\mathbf{r})$ and $c_2(\mathbf{r})$ in $c_1\psi_1 + c_2\psi_2$ would be zero. If the particles are electrons and the wave functions are spinors, the phases are rather changed by some rotations $\psi_1 \rightarrow \mathbf{R}_1\psi_1$ and $\psi_2 \rightarrow \mathbf{R}_2\psi_2$, but the rest of the argument remains analogous because for the SU(2) matrices \mathbf{R}_j we have $\mathbf{R}_j^\dagger \mathbf{R}_j = \mathbb{1}$. As said, what we are discussing here in this footnote for wave equations is different from the situation we are studying in the main text, which is that of a coherent electromagnetic beam.

⁴ A full solution for the paradox of the double slit experiment is proposed in Ref. [3].

fact that $\forall g \in G : [\psi(g)]^\dagger \psi(g) = 1$. The spinors are normalized according to $\psi^\dagger \psi = 1$ because they represent SU(2) matrices. Hence counting a number of electrons cannot be done by adding quantities $\psi(g)$ but by counting quantities $[\psi(g)]^\dagger \psi(g)$. Each count $[\psi(g)]^\dagger \psi(g) = 1$ counts one electron, the electron that tags g .

We must thus count $3\psi_1^\dagger \psi_1 + 2\psi_2^\dagger \psi_2$. And therefore we must start from the set of quantities $\{\sqrt{3}\psi_1, \sqrt{2}\psi_2\}$, or $\sqrt{3}\psi_1 + \sqrt{2}\psi_2$. Therefore, we come to the conclusion that $\sum_j c_j \psi_j$ represents a set of particles whereby $|c_j|^2$ particles are in the state ψ_j , just as Ehrenfest has told us. We have explained this in more details in [5].

For photons the argument is different from the one for electrons. Here we use the idea that $E^2 + c^2 B^2$ represents the total energy. If the wave is monochromatic, all photons have the same energy and therefore $E^2 + c^2 B^2$ is then a measure for the number of photons. In the Eq.1 below we have $\psi_x^\dagger \psi_x = E^2 + c^2 B^2$, which represents the energy, and therefore also the number of photons that contribute to this energy. Hence in both cases, the numbers of the particles in the states are proportional to $\psi^\dagger \psi$. This can be considered as a derivation of the Born rule and the prescription of Ehrenfest. In the next Section we will work out further the idea that photons are particles and that an electromagnetic wave must be a large collection of photons. The fact that we can succeed in working out this idea is in my opinion evidence for its validity.

2 Interpreting a linearly polarized electromagnetic wave as a swarm of photons




Fig. 1. Electromagnetic wave corresponding to Eq. 2.

We are used to discuss about polarizations of electromagnetic waves, but much less so about the spin of photons. Text-books tell us that photons have spin 1, which is puzzling, because this should lead to three substates with quantum numbers $-1, 0, 1$. But we only have linear polarizations -1 and $+1$. Where is the substate with quantum number zero? We will here explain the relationship between polarization and spin. It is important to have this connection, because in interactions we must be able to apply conservation laws. The basic quantity must therefore be the spin, not the polarization. We may note here that the discussion of the conservation law for spin and angular momentum is often tacitly omitted in text books of nuclear physics, which teach us e.g. the photoelectric effect or the Compton effect [6]. Only the conservation laws for energy and momentum are worked out. We are loosing some interesting information this way. Taking into account the conservation of spin and angular momentum shows that an electron must flip its spin when it absorbs a photon. This renders the abstract conservation law much more visual.

A linear polarization can be obtained as the superposition of two circular polarizations of opposite sign. If we accept that photons are particles with a spin then we must conclude that they must be circularly polarized such that a single photon can never have a linear polarization. It is only an ensemble of photons that can have a linear polarization. One should thus not try to derive Malus' law in terms of photons that have a linear polarization. Just presenting photons with polarizations + or - as is done in the derivation of the Bell inequalities is also wrong (but this will not change the inequalities).

We will now try to understand how photons build an electromagnetic wave. We will consider first an electromagnetic wave that is linearly polarized along the x -axis and that propagates along the positive z -axis. The electric field and the magnetic field of a linearly polarized wave are in phase. Therefore we can have for the linear polarization along the x -axis in the Oxy plane:

$$\psi_x = \begin{bmatrix} E_x \\ cB_y \end{bmatrix} = \begin{bmatrix} E \\ \imath cB \end{bmatrix} \cos(\omega t) \quad (1)$$

We are expressing here the polarization with the aid of complex numbers. The y -direction corresponds then to the pre-factor \imath . By rotating this over $\pi/2$ we obtain a linear polarization along the y -axis:

$$\psi_y = \begin{bmatrix} E_y \\ cB_x \end{bmatrix} = \begin{bmatrix} \imath E \\ -cB \end{bmatrix} \sin(\omega t) \quad (2)$$

The electric and magnetic field are again in phase, but we are taking here sine waves in order to end up with dependences $e^{\pm i\omega t}$ later on.⁵ Let us now construct the spin-up and spin-down states. This is just like a change of basis, but we will not discuss the normalization of the basis vectors for the moment. We will just introduce some normalization here, and we will justify our choice later on. By summing we obtain:

$$\psi_L = \frac{1}{\sqrt{2}}(\psi_x + \psi_y) = \frac{1}{\sqrt{2}} \begin{bmatrix} E_x + E_y \\ cB_x + cB_y \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} E \\ \imath cB \end{bmatrix} e^{i\omega t} \quad (3)$$

Now we add the factor that expresses the propagation along the positive z -axis. It must always have an argument with the combination $k_z z - \omega t$. But as we want to end up with a wave in $e^{i\omega t}$ we must thus take here the argument $-(k_z z - \omega t)$.

$$\psi_L = \frac{1}{\sqrt{2}}(\psi_x + \psi_y) = \frac{1}{\sqrt{2}} \begin{bmatrix} E_x + E_y \\ cB_x + cB_y \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} E \\ \imath cB \end{bmatrix} e^{-i(k_z z - \omega t)} \quad (4)$$

By subtracting we obtain the other basis vector (again not discussing its normalization):

$$\psi_R = \frac{1}{\sqrt{2}}(\psi_x - \psi_y) = \frac{1}{\sqrt{2}} \begin{bmatrix} E_x - E_y \\ cB_y - cB_x \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} E \\ \imath cB \end{bmatrix} e^{-i\omega t} \quad (5)$$

We can finally add the terms that express the propagation along the positive z -axis:

$$\psi_R = \frac{1}{\sqrt{2}}(\psi_x - \psi_y) = \frac{1}{\sqrt{2}} \begin{bmatrix} E \\ \imath cB \end{bmatrix} e^{i(k_z z - \omega t)} \quad (6)$$

Now if we want to go back to the original electromagnetic wave which is polarized along the x -axis, we have to take:

$$\psi_x = \frac{1}{\sqrt{2}}(\psi_L + \psi_R) \quad (7)$$

We assume now that each photon has the same internal dynamics as expressed in ψ_L or ψ_R . This way we can express the electromagnetic wave as a swarm of photons where half of them have a left circular polarization and the other half a right circular polarization. These left and right circular polarizations correspond to the spin states -1 and $+1$. We can now see the reason why we normalized with $\frac{1}{\sqrt{2}}$. In fact, when we have a superposition state $c_L \psi_L + c_R \psi_R$, then the numbers of the photons in the states ψ_L and ψ_R are $|c_L|^2$ and $|c_R|^2$. It is therefore that we must use the linear combinations:

$$\psi_x = \frac{1}{\sqrt{2}}(\psi_L + \psi_R). \quad (8)$$

$$\psi_y = \frac{1}{\sqrt{2}}(\psi_L - \psi_R). \quad (9)$$

This can be written as a transformation:

⁵ In fact, as the polarization vector of ψ_y is just \imath times the polarization vector of ψ_x , taking a cosine wave again would not lead to an independent mode. Change from cosines to sines ensures independence.

$$\begin{bmatrix} \psi_x \\ \psi_y \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \psi_L \\ \psi_R \end{bmatrix} \quad (10)$$

The inverse transformation is then indeed:

$$\begin{bmatrix} \psi_L \\ \psi_R \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \psi_x \\ \psi_y \end{bmatrix} \quad (11)$$

This development solves a number of riddles. First of all we can now clearly see that the electromagnetic wave is quantized. Secondly we can understand how an electromagnetic wave can propagate in vacuum. In principle there are no charges in vacuum. There is thus nothing the wave could jiggle to propagate itself. The wave propagates in the absence of any medium.⁶ But by considering the wave as a swarm of photons, each of which carries a small part of the field this conceptual problem disappears. This is perfectly analogous with the electrons in the double-slit experiment which traverse the set-up one by one without any interaction between them. They also do not need any medium to do their stuff like a sound or water wave. All they need is some clock of internal dynamics. Another problem that disappears is that of the collapse of the wave function, as in our approach the particle is always in a well-defined place. Finally we would like to point out that also the electron wave function in the Dirac equation is a mixed state whereby the two components are obtained one from another by reflection (see [4], pp. 162-166).

If we want now a total of N photons in the wave at a certain spot \mathbf{r} we must take $\sqrt{N}\psi_x(\mathbf{r})$. One may wonder why a photon with spin 1 has only two substates -1 and $+1$. The reason for it will become clear in Section 3. It is because the linearly polarized electromagnetic wave has no zero-frequency component. As we can see on a drawing of such a wave in Fig. 1, it has nodes. The numbers -1 , 0 , and $+1$ do not have anything to do with some directions in space. They just correspond to frequencies (or actually to the prefactor m in $e^{im\phi}$ within the spherical harmonics $Y_{\ell m}(\theta, \phi)$, where we put $\phi = \omega t$ in order to be able to use the mathematical tool to describe dynamics, just like we did for the electron in the Rodriguez formula). The waves are transverse. All the oscillations are inside a plane perpendicular to the direction of propagation. Therefore the group we must consider is the one of the rotations and reversals within the polarization plane. Rotations are obtained from an even number of reflections, while reversals are obtained from an odd number of reflections. These operations are abstract because they are not taking place in space-time but in the plane of electromagnetic polarization vectors.

If you consider a rotation as the product of an even number of reflections, then the state obtained by complex conjugation from it is the product of an odd number of reflections. In fact $i \rightarrow -i$ in $e^{i\phi} \rightarrow e^{-i\phi}$ is a reflection with respect to the x -axis. The angular momenta are real, they can provide an optical torque and the effect of such torques has been measured[7]. They also play a rôle in selection rules for atomic transitions [1].

The 2×1 representations we developed above permit to calculate the numbers of the particles. Actually, a real wave would be described with a hull function $\sqrt{\rho(z)}$, where ρ is the probability density, because the linearly polarized electromagnetic wave will not have an infinite extent. It will rather be a wave train, with a coherence length. Here $\sqrt{\rho(z)}$ will only be non-zero in some finite (moving) z -interval $[z_1, z_2]$. And you have then to normalize according to $\int_{z_1}^{z_2} \int_{x_1}^{x_2} \int_{y_1}^{y_2} \psi^\dagger \psi dx dy dz = N\hbar\omega$. N would then be the total number of photons in the wave within the interval $[x_1, x_2] \times [y_1, y_2] \times [z_1, z_2]$, and $\rho = |\psi|^2$ its density. And the true wave will then be a kind of photon burst. We may note that the expressions for $\psi_L(\mathbf{r}, t)$ and $\psi_R(\mathbf{r}, t)$ do not contain the variables x and y . The x and y coordinates can therefore be chosen at will. For the density this implies a homogeneous density throughout the planes $z = K$, where K is constant. Our wave function describes thus really a plane wave.

Let us remind that according to the group theory performing the sum $\psi_1 + \psi_2$ of two spinors actually does not make sense (See e.g. Ref [5]). An example is the sum of a spinor $\psi(\mathbf{s}, 2\pi)$ for a rotation around an axis \mathbf{s} over an angle 2π and a spinor $\psi(\mathbf{s}, 4\pi)$ for a rotation around the same axis over an angle 4π . This sum is exactly zero and has no meaning whatsoever. However when we have two photons, the wave functions contain the electric and magnetic fields of the photons and the fields of the photons would add up, if they were at exactly the same position at the same time. In general, this will not happen and the photons will be in different positions.

A number of properties, like Malus' law, are rather easy to derive from a wave picture but much less so from a particle picture. Conceptually that made it difficult to solve the particle-wave duality for electromagnetic waves along the same line as for electrons. Interpreting the waves in terms of particles as we have done may be instrumental in figuring out a derivation based on the particle picture.

3 Fourier analysis

In the previous Section we have found the decomposition by a lot of guessing and intuition. When we go to more complicated situations that intuition may fail us. We therefore would like to replace it by a more algebraic approach. Let us try to find the decomposition by Fourier analysis. Let us first introduce the notation:

⁶ We consider virtual particles in Quantum Electrodynamics as mere mathematical tools.

$$\mathbf{a} = \begin{bmatrix} E_x \\ cB_y \end{bmatrix}, \quad (12)$$

for the polarization vector of ψ_x . The Fourier coefficient for the $e^{+i\phi}$ component of ψ_x becomes then:

$$\frac{1}{2\pi} \int_0^{2\pi} \psi_x e^{i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} \cos(\phi) e^{i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} \cos^2(\phi) d\phi + \frac{i\mathbf{a}}{2\pi} \int_0^{2\pi} \cos\phi \sin\phi d\phi = \frac{1}{2} \mathbf{a}. \quad (13)$$

The Fourier coefficient for the $e^{-i\phi}$ component of ψ_x becomes then:

$$\frac{1}{2\pi} \int_0^{2\pi} \psi_x e^{-i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} \cos(\phi) e^{-i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} \cos^2(\phi) d\phi - \frac{i\mathbf{a}}{2\pi} \int_0^{2\pi} \cos\phi \sin\phi d\phi = \frac{1}{2} \mathbf{a}. \quad (14)$$

The Fourier coefficient for the $e^{0i\phi} = 1$ component of ψ_x becomes then:

$$\frac{1}{2\pi} \int_0^{2\pi} \psi_x d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} \cos(\phi) d\phi = \mathbf{0}. \quad (15)$$

The polarization vector of ψ_y is:

$$\mathbf{b} = i\mathbf{a} = i \begin{bmatrix} E_x \\ cB_y \end{bmatrix} = i \begin{bmatrix} E \\ i cB \end{bmatrix} = \begin{bmatrix} iE \\ -cB \end{bmatrix}. \quad (16)$$

We obtain for the Fourier component $e^{+i\phi}$ of ψ_y :

$$\frac{1}{2\pi} \int_0^{2\pi} \psi_y e^{i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} i \sin(\phi) e^{i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} i \cos(\phi) \sin(\phi) d\phi - \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} \sin^2 \phi d\phi = -\frac{1}{2} \mathbf{a}, \quad (17)$$

for the Fourier component $e^{-i\phi}$ of ψ_y :

$$\frac{1}{2\pi} \int_0^{2\pi} \psi_y e^{-i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} i \sin(\phi) e^{-i\phi} d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} i \cos(\phi) \sin(\phi) d\phi + \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} \sin^2 \phi d\phi = \frac{1}{2} \mathbf{a}, \quad (18)$$

and for the Fourier component $1 = e^{0i\phi}$ of ψ_y :

$$\frac{1}{2\pi} \int_0^{2\pi} \psi_y d\phi = \frac{\mathbf{a}}{2\pi} \int_0^{2\pi} i \sin(\phi) d\phi = \mathbf{0}. \quad (19)$$

This justifies that photons with a quantum number 0 do not occur. As already stated, the electromagnetic wave has nodes, and therefore has no constant component. Finally, there is a renormalization to be done, because we must make the intensities comply with the rule that they are $|c_j|^2$ rather than c_j . We obtain then the same results as before. Independence of the basis vectors corresponds here to $\int \psi_1^\dagger \psi_2 d\phi = 0$ and normalization to $\int \psi_1^\dagger \psi_1 d\phi = 1$.

4 A remark on gravitons and gravitational waves

We think it could be useful to apply our solution of the particle-wave duality also to gravitons and gravitational waves even if we are not likely to be confronted with the issues of quantum mechanics in gravitation any soon. The currently accepted notion is that gravitational waves are rippling the fabric of space-time. The point we want to make is that also gravitational waves do not need a medium. All they need is some periodic internal dynamics within the gravitons that can be used as a clock. These internal dynamics would mirror the dynamics you attribute to the wave. It could be done the same way by analyzing the polarizations of the waves like we did for electromagnetism.

Considering gravitational waves as deformations of space-time has one merit: It is convenient. Since Euclid, geometry has become a part of mathematics, even if in the beginning it came from observation and was thus a kind of physics. The perfection of the theory was so great that it became considered as pre-existing any physics. It had become abstract and dematerialized. And in Newtonian physics it became a pre-existing playground which God had visited to lay down his laws of physics in it. We even thought at some point further back in time that the surface of the Earth was an infinite Euclidean plane, because our observations were too crude to notice the curvature. Only to find out later on that this infinite plane without curvature does not correspond to the physical reality and had been preconceived. The mathematics of this Euclidean plane were completely logical and contradiction-free, but they were not the mathematics that corresponded to the real situation. With the theory of relativity came the shock of realizing again that the Euclidean metric we had chosen for the vector space \mathbb{R}^4 by which we presented space and time was perfectly logic and error-free but not the right one. But after that, the new geometry, Minkowski's space-time with its Riemannian pseudo-metric evolved again conceptually to the status of a perfectly

logical pre-existing mathematical playground. But the lesson that we should have learned from special relativity is that space-time is created by electromagnetism. We are seeing straight lines because the atoms in the matter we use to visualize them are kept in place by electromagnetic interactions. It is the electromagnetic interactions which conjure up our perceptions of space and time. Space-time of special relativity is thus electromagnetic space-time. It is the space-time that corresponds to the invariance of the electromagnetic interactions. There is a deep connection between physics and the geometry we adopt to describe it and it is just not “geometry first”.

We must warn against the strongly suggestive character of geometrical theories that make them look as a piece of mathematics that pre-exists the physics. We can see that space-time cannot be the vector space \mathbb{R}^4 we imagine it to be because the Universe is expanding. Another idiosyncrasy introduced by this concept of \mathbb{R}^4 as a pre-existing mathematical playground for the laws of physics is the idea that the future would already exist. This takes the mathematical tool of the space-time vector space \mathbb{R}^4 again really too serious. To define an infinite space-time vector space \mathbb{R}^4 we would need to have clocks that are synchronized up to infinite distance. As Einstein synchronization must be carried out by sending light rays to and fro two observers, there is no way we can possibly achieve this. It is this synchronization procedure which is responsible for the superluminal phase velocity of the de Broglie waves. We must thus really keep in mind that space-time is a part of the physics, that it is created by the expansion of the Universe, starting from the big bang, and that it has to be a manifold rather than a vector space. General relativity states that space-time is curved. Our bodies, meter sticks or our tight ropes that we use to visualize a straight line would become curved if we were swirling around a very heavy object. It is then very practical to use the curves the objects are taking as geodesics in our calculations. The tight rope will still be as straight a thing as it gets. The curved geometry is really the best reference geometry we can use to describe our observations.

Describing things this way is really pragmatic. What it really acknowledges for is that two types of forces are shaping now our geometrical stage, rather than one. We can make the identification and claim that it is space-time which is curved. But this relies tacitly on the principle of equivalence, that all matter reacts the same way to gravity. If there were some type of matter that violates the equivalence principle, the universality of the geometry would break down. This shows that there is a tacit assumption underlying our philosophy. One can push this idea of universality to the limit where we consider that gravitational waves are deforming space-time, despite the fact that our local pragmatic choice of geometry ought to be flat because it is dominated by electromagnetism. That becomes then analogous to conceiving phonons in a crystal as deforming space-time, which is not truly convincing an idea. Especially it conjures up the impression that space-time would be a medium like the aether, an elastic medium whose vibrations are the gravitational waves. That aether is now called the fabric of space-time. It could be wrong as many times before. We therefore think that it is better not to decide for the moment if this is true or otherwise. What we are arguing here is not a new theory, it is just a flashing warning light. One could wonder e.g. what our geometry would become when the four fundamental forces were acting at equal strength, or when gravitation overwhelms electromagnetism entirely like in a black hole. The way we cannot stop ourselves from falling into a black hole resembles in this respect very much our notion of the passage of time. For the moment our choices do not have any practical incidence on our physics. They are only changing the way we are telling the story. But we must be aware of the tacit assumptions that are underlying them. They could break down at some new stage of knowledge and then become an intellectual barrier against understanding the physics, just like Euclidean geometry was a conceptual barrier to be overcome in formulating relativity.

References

1. C.V. Raman and S. Bhagavantam, *Ind. J. Phys.* **6**, 353-366 (1931).
2. A. Tonomura, J. Endo, T. Matsuda and T. Kawaski, *Am. J. Phys.* **57**, 647 (1986).
3. G. Coddens, <https://hal.archives-ouvertes.fr/cea-01459890v3>
4. G. Coddens, in *From Spinors to Quantum Mechanics*, Imperial College Press, London (2015).
5. G. Coddens, <https://hal.archives-ouvertes.fr/cea-01572342v1>.
6. P. Marmier and E. Sheldon in *Physics of Nuclei and Particles* (three volumes), Academic Press, New York, (1969).
7. Li He, Huan Li and Mo Li, *Science Advances*, **2**(9), e1600485 (2016).