



HAL
open science

Analysis of variational formulations and low-regularity solutions for time-harmonic electromagnetic problems in complex anisotropic media

Damien Chicaud, Patrick Ciarlet, Axel Modave

► To cite this version:

Damien Chicaud, Patrick Ciarlet, Axel Modave. Analysis of variational formulations and low-regularity solutions for time-harmonic electromagnetic problems in complex anisotropic media. SIAM Journal on Mathematical Analysis, 2021, 53 (3), pp.2691-2717. hal-02651682v1

HAL Id: hal-02651682

<https://hal.science/hal-02651682v1>

Submitted on 29 May 2020 (v1), last revised 17 Dec 2021 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Analysis of variational formulations and low-regularity solutions for time-harmonic electromagnetic problems in complex anisotropic media

Damien Chicaud

*POEMS, CNRS, INRIA, ENSTA Paris, Institut Polytechnique de Paris, 828 Bd des Maréchaux
91762 Palaiseau Cedex, France
damien.chicaud@ensta-paris.fr*

Patrick Ciarlet Jr.

*POEMS, CNRS, INRIA, ENSTA Paris, Institut Polytechnique de Paris, 828 Bd des Maréchaux
91762 Palaiseau Cedex, France
patrick.ciarlet@ensta-paris.fr*

Axel Modave

*POEMS, CNRS, INRIA, ENSTA Paris, Institut Polytechnique de Paris, 828 Bd des Maréchaux
91762 Palaiseau Cedex, France
axel.modave@ensta-paris.fr*

We consider the time-harmonic Maxwell's equations with physical parameters, namely the electric permittivity and the magnetic permeability, that are complex, possibly non-hermitian, tensor fields. Both tensor fields verify a general ellipticity condition. In this work, the well-posedness of formulations for the Dirichlet and Neumann problems (*i.e.* with a boundary condition on the electric field or its curl, respectively) is proven using well-suited functional spaces and Helmholtz decompositions. For both problems, the *a priori* regularity of the solution and the solution's curl is analysed. The regularity results are obtained by splitting the fields and using shift theorems for second-order divergence elliptic operators. Finally, the discretization of the formulations with a $\mathbf{H}(\mathbf{curl})$ -conforming approximation based on edge finite elements is considered. An *a priori* error estimate is derived and verified thanks to numerical results with an elementary benchmark.

Keywords: Maxwell's Equations; Wave Propagation; Anisotropic media; Well-posedness; Regularity Analysis; Edge Finites Elements

AMS Subject Classification: 35J57, 65N30, 78M10

1. Introduction

The study of linear differential models for time-harmonic electromagnetic wave propagation is a popular field. The mathematical analysis of these models has been performed for isotropic materials and certain classes of anisotropic materials. In particular, the well-posedness of boundary value problems and the regularity of their solutions have been studied with strong assumptions on the electric permittivity tensor field $\underline{\epsilon}$ and the magnetic permeability tensor field $\underline{\mu}$ (*e.g.* real-valued isotropic tensors⁸ and symmetric positive definite tensors^{16,4,6}). Nevertheless, to the authors' knowledge, little is known when the coefficients defining the model are complex non-hermitian tensor fields.

The main aim of this work is to provide a detailed analysis of time-harmonic electromagnetic boundary value problems for a general class of material tensors: the electric permittivity and

the magnetic permeability are assumed to be complex tensor fields, possibly non-hermitian, that fulfill a general ellipticity condition. We consider variational formulations with the electric field as unknown, with a boundary condition on the field itself or on its curl, which correspond to the so-called Dirichlet and Neumann cases, respectively. The well-posedness and the *a priori* regularity of the solution and the solution's curl are studied for both cases. In our framework, the regularity estimates depend crucially on the regularity of the data.

The numerical solution of electromagnetic problems are naturally performed with edge finite element methods, which lead to $\mathbf{H}(\mathbf{curl})$ -conforming approximations. While the comprehensive numerical analysis of the approximate problems is out of the scope of this paper, a preliminary analysis is proposed. In the coercive case, an *a priori* error estimate is derived by leveraging the regularity results obtained for the exact problems. A numerical illustration with a simple manufactured benchmark is presented as well.

This paper is organized as follows. In section 2, we extend classical results of functional analysis required for the analyses. In sections 3 and 4, we propose the analysis of the problems supplemented by Dirichlet or Neumann boundary conditions, respectively. In section 5, we address the discretization with edge finite elements. A conclusion and extensions are proposed in section 6.

Notation and hypotheses

Vector fields are written in boldface character, and tensor fields are written in underlined bold characters. Given a non-empty open set \mathcal{O} of \mathbb{R}^3 , we use the notation $(\cdot)_{0,\mathcal{O}}$ (resp. $\|\cdot\|_{0,\mathcal{O}}$) for the $L^2(\mathcal{O})$ and the $\mathbf{L}^2(\mathcal{O}) := (L^2(\mathcal{O}))^3$ scalar products (resp. norms). More generally, $(\cdot)_{\mathbf{s},\mathcal{O}}$ and $\|\cdot\|_{\mathbf{s},\mathcal{O}}$ (resp. $|\cdot|_{\mathbf{s},\mathcal{O}}$) denote the scalar product and the norm (resp. semi-norm) of the Sobolev spaces $H^{\mathbf{s}}(\mathcal{O})$ and $\mathbf{H}^{\mathbf{s}}(\mathcal{O}) := (H^{\mathbf{s}}(\mathcal{O}))^3$ for $\mathbf{s} \in \mathbb{R}$ (resp. for $\mathbf{s} > 0$). The index $_{\text{zmv}}$ denotes functional spaces with zero-mean-value fields. It is assumed that the reader is familiar with the function spaces related to Maxwell's equations, such as $\mathbf{H}(\mathbf{curl}; \mathcal{O})$, $\mathbf{H}_0(\mathbf{curl}; \mathcal{O})$, $\mathbf{H}(\text{div}; \mathcal{O})$ and $\mathbf{H}_0(\text{div}; \mathcal{O})$. A priori, $\mathbf{H}(\mathbf{curl}; \mathcal{O})$ is endowed with the "natural" norm $\mathbf{v} \mapsto (\|\mathbf{v}\|_{0,\mathcal{O}}^2 + \|\mathbf{curl} \mathbf{v}\|_{0,\mathcal{O}}^2)^{1/2}$. We refer to the monographs of Monk¹⁷ and Assous *et al.*² for further details.

The symbol C is used to denote a generic positive constant which is independent of the mesh and the fields of interest; C may depend on the geometry, or on the coefficients defining the model. We use the notation $A \lesssim B$ for the inequality $A \leq CB$, where A and B are two scalar fields, and C is a generic constant.

In this work, Ω is assumed to be an open, connected, bounded domain of \mathbb{R}^3 , with a Lipschitz-continuous boundary $\partial\Omega$. In addition, the boundary is assumed to be of class \mathcal{C}^2 in the sections dedicated to the regularity analysis (in subsections 3.2, 3.3, 4.2 and 4.3). The unit outward normal vector field to $\partial\Omega$ is denoted \mathbf{n} . We recall the classical integration by parts formula (see Eq. (35) in Ref. 5):

$$(\mathbf{u} \mathbf{curl} \mathbf{v})_{0,\Omega} - (\mathbf{curl} \mathbf{u} | \mathbf{v})_{0,\Omega} = \gamma \langle \gamma^T(\mathbf{u}), \pi^T(\mathbf{v}) \rangle_{\pi}, \quad \forall \mathbf{u}, \mathbf{v} \in \mathbf{H}(\mathbf{curl}; \Omega), \quad (1.1)$$

where $\gamma^T : \mathbf{u} \mapsto (\mathbf{u} \times \mathbf{n})|_{\partial\Omega}$ denotes the tangential trace operator, $\pi^T : \mathbf{v} \mapsto \mathbf{n} \times (\mathbf{v} \times \mathbf{n})|_{\partial\Omega}$ denotes the tangential components trace operator, and $\gamma \langle \gamma^T(\mathbf{u}), \pi^T(\mathbf{v}) \rangle_{\pi}$ expresses duality between the *ad hoc* trace spaces. We denote $\{\Gamma_k\}_{0 \leq k \leq K}$ the maximal connected components of $\partial\Omega$. Topologically speaking, the domain Ω can be trivial or non-trivial¹³ under assumptions given in Appendix A. Whenever this knowledge is needed, we use the notation $(\mathbf{Top})_I$, with $I = 0$ for a topologically trivial domain and $I > 0$ for a topologically non-trivial domain.

2. Model and extended functional framework

We consider the time-harmonic Maxwell's equation, posed in Ω :

$$\mathbf{curl}(\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E}) - \omega^2 \underline{\boldsymbol{\varepsilon}} \mathbf{E} = \mathbf{f},$$

where our unknown is the electric field \mathbf{E} , which we assume *a priori* to belong to $\mathbf{H}(\mathbf{curl}; \Omega)$; $\underline{\boldsymbol{\varepsilon}}$ and $\underline{\boldsymbol{\mu}}$ are respectively the electric permittivity tensor and the magnetic permeability tensor; $\omega > 0$ is the angular frequency, and \mathbf{f} is a volume data. The problem shall be supplemented by appropriate boundary conditions on $\partial\Omega$.

In this work, the material tensors $\underline{\boldsymbol{\varepsilon}}$ and $\underline{\boldsymbol{\mu}}$ are assumed to satisfy an ellipticity condition. This condition is defined in the following subsection, and useful technical properties are given. The remaining of the section is dedicated to the introduction of the framework that derives from the ellipticity condition, and the extension of classical results (*i.e.* for elliptic, hermitian tensor fields) given in Ref. 2: Helmholtz decompositions, Weber inequalities, and compact embedding results.

2.1. Ellipticity condition

Definition 2.1. We say that a complex-valued tensor field $\underline{\boldsymbol{\xi}} \in \underline{\mathbf{L}}^\infty(\Omega)$ is elliptic if and only if

$$\exists \theta_\xi \in [0, 2\pi[, \exists \xi_- > 0, \forall \mathbf{z} \in \mathbb{C}^3, \xi_- |\mathbf{z}|^2 \leq \Re[e^{i\theta_\xi} \cdot \mathbf{z}^* \underline{\boldsymbol{\xi}} \mathbf{z}] \quad \text{almost everywhere in } \Omega. \quad (2.1)$$

Additionally, we will use the notation $\xi_+ := \|\underline{\boldsymbol{\xi}}\|_{\underline{\mathbf{L}}^\infty(\Omega)}$.

This condition means that, almost everywhere in Ω , the eigenvalues of $\underline{\boldsymbol{\xi}}(\mathbf{x})$ are contained in a fixed open half-plane of \mathbb{C}^3 ; in other words, there exists a "coercivity direction" for $\underline{\boldsymbol{\xi}}$ in the complex plane. On the other hand, we do not assume that $\underline{\boldsymbol{\xi}}$ is hermitian: as a consequence, the mapping $(\mathbf{v}, \mathbf{w}) \mapsto (\underline{\boldsymbol{\xi}} \mathbf{v} | \mathbf{w})_{0,\Omega}$ is, in general, not a scalar product in $\mathbf{L}^2(\Omega)$; orthogonality properties are lateralized, in the sense that $(\underline{\boldsymbol{\xi}} \mathbf{v} | \mathbf{w})_{0,\Omega} = 0$ is not equivalent to $(\underline{\boldsymbol{\xi}} \mathbf{w} | \mathbf{v})_{0,\Omega} = 0$.

Assumption. In the remaining of Section 2, the tensor $\underline{\boldsymbol{\xi}}$ belongs to $\underline{\mathbf{L}}^\infty(\Omega)$ and satisfies the ellipticity condition.

Proposition 2.1. One has $\underline{\boldsymbol{\xi}}^{-1} \in \underline{\mathbf{L}}^\infty(\Omega)$ with $(\xi^{-1})_+ \leq (\xi_-)^{-1}$. Moreover, $\underline{\boldsymbol{\xi}}^{-1}$ satisfies the ellipticity condition with $\theta_{\xi^{-1}} = -\theta_\xi$, and $(\xi^{-1})_- := \xi_- (\xi_+)^{-2}$.

Proposition 2.2. For any $\mathbf{v} \in \mathbf{L}^2(\Omega)$, one has the following inequalities:

$$\xi_- \|\mathbf{v}\|_{0,\Omega}^2 \leq \Re \left[e^{i\theta_\xi} (\underline{\boldsymbol{\xi}} \mathbf{v} | \mathbf{v})_{0,\Omega} \right] \leq \left| (\underline{\boldsymbol{\xi}} \mathbf{v} | \mathbf{v})_{0,\Omega} \right| \leq \xi_+ \|\mathbf{v}\|_{0,\Omega}^2. \quad (2.2)$$

Then, the ellipticity condition implies that the inverse of the second-order divergence elliptic operators, with Dirichlet or Neumann boundary condition, is well-defined. Equivalently, this implies the well-posedness of the problems below, called either Dirichlet problem, or Neumann problem, from now on.

Theorem 2.1. Under assumption (2.1), the Dirichlet problem

$$\begin{cases} \text{Find } p \in H_0^1(\Omega) \text{ such that} \\ (\underline{\boldsymbol{\xi}} \nabla p | \nabla q)_{0,\Omega} = \ell(q), \quad \forall q \in H_0^1(\Omega), \end{cases} \quad (2.3)$$

is well-posed for all ℓ in $H^{-1}(\Omega) = (H_0^1(\Omega))'$, that is

$$\exists C > 0, \forall \ell \in (H_0^1(\Omega))', \exists! p \text{ solution to (2.3), with } \|p\|_{1,\Omega} \leq C \|\ell\|_{(H_0^1(\Omega))'}.$$

Similarly, under assumption (2.1), the Neumann problem

$$\begin{cases} \text{Find } p \in H_{\text{zmv}}^1(\Omega) \text{ such that} \\ (\underline{\xi} \nabla p | \nabla q)_{0,\Omega} = \ell(q), \forall q \in H_{\text{zmv}}^1(\Omega), \end{cases} \quad (2.4)$$

is well-posed for all in $(H_{\text{zmv}}^1(\Omega))'$, that is

$$\exists C > 0, \forall \ell \in (H_{\text{zmv}}^1(\Omega))', \exists! p \text{ solution to (2.4), with } \|p\|_{1,\Omega} \leq C \|\ell\|_{(H_{\text{zmv}}^1(\Omega))'}.$$

Proof. One uses Lax-Milgram's theorem. All assumptions are fulfilled, and in particular coercivity of the sesquilinear form follows from relations (2.2) and Poincaré, resp. Poincaré-Wirtinger, inequalities. \square

2.2. Helmholtz decompositions

Definition 2.2. We introduce the following function spaces:

$$\begin{aligned} \mathbf{H}(\text{div } \underline{\xi}; \Omega) &:= \{v \in \mathbf{L}^2(\Omega), \underline{\xi} v \in \mathbf{H}(\text{div}; \Omega)\}, \\ \mathbf{H}_0(\text{div } \underline{\xi}; \Omega) &:= \{v \in \mathbf{L}^2(\Omega), \underline{\xi} v \in \mathbf{H}_0(\text{div}; \Omega)\}, \\ \mathbf{H}(\text{div } \underline{\xi} 0; \Omega) &:= \{v \in \mathbf{H}(\text{div } \underline{\xi}; \Omega), \text{div } \underline{\xi} v = 0\}, \\ \mathbf{H}_0(\text{div } \underline{\xi} 0; \Omega) &:= \mathbf{H}_0(\text{div } \underline{\xi}; \Omega) \cap \mathbf{H}(\text{div } \underline{\xi} 0; \Omega), \end{aligned}$$

as well as

$$\begin{aligned} \mathbf{X}_N(\underline{\xi}; \Omega) &:= \mathbf{H}_0(\mathbf{curl}; \Omega) \cap \mathbf{H}(\text{div } \underline{\xi}; \Omega), \\ \mathbf{X}_T(\underline{\xi}; \Omega) &:= \mathbf{H}(\mathbf{curl}; \Omega) \cap \mathbf{H}_0(\text{div } \underline{\xi}; \Omega), \\ \mathbf{K}_N(\underline{\xi}; \Omega) &:= \mathbf{H}_0(\mathbf{curl}; \Omega) \cap \mathbf{H}(\text{div } \underline{\xi} 0; \Omega), \\ \mathbf{K}_T(\underline{\xi}; \Omega) &:= \mathbf{H}(\mathbf{curl}; \Omega) \cap \mathbf{H}_0(\text{div } \underline{\xi} 0; \Omega). \end{aligned}$$

The function spaces $\mathbf{X}_N(\underline{\xi}; \Omega)$, $\mathbf{X}_T(\underline{\xi}; \Omega)$, $\mathbf{K}_N(\underline{\xi}; \Omega)$ and $\mathbf{K}_T(\underline{\xi}; \Omega)$ are endowed with the graph norm $v \mapsto (\|v\|_{\mathbf{H}(\mathbf{curl}; \Omega)}^2 + \|\underline{\xi} v\|_{\mathbf{H}(\text{div}; \Omega)}^2)^{1/2}$. When $\underline{\xi}$ is equal the identity tensor \mathbb{I}_3 , we choose the simpler notation $\mathbf{X}_N(\Omega)$ instead of $\mathbf{X}_N(\mathbb{I}_3; \Omega)$, etc.

As a first noticeable consequence of (2.1), one can prove the Helmholtz decompositions below.

Theorem 2.2. *Under assumption (2.1), one has the first-kind Helmholtz decompositions:*

$$\mathbf{L}^2(\Omega) = \nabla H_0^1(\Omega) \oplus \mathbf{H}(\text{div } \underline{\xi} 0; \Omega); \quad (2.5)$$

$$\mathbf{H}_0(\mathbf{curl}; \Omega) = \nabla H_0^1(\Omega) \oplus \mathbf{K}_N(\underline{\xi}; \Omega). \quad (2.6)$$

In addition, these sums are continuous.

Proof. Let $v \in \mathbf{L}^2(\Omega)$. The Dirichlet problem

$$\begin{cases} \text{Find } p \in H_0^1(\Omega) \text{ such that} \\ (\underline{\xi} \nabla p | \nabla q)_{0,\Omega} = (\underline{\xi} v | \nabla q)_{0,\Omega}, \forall q \in H_0^1(\Omega), \end{cases}$$

is well-posed by theorem 2.1. Let $\mathbf{v}_T = \mathbf{v} - \nabla p \in \mathbf{L}^2(\Omega)$. Then $\operatorname{div} \underline{\xi} \mathbf{v}_T = \operatorname{div} \underline{\xi} \mathbf{v} - \operatorname{div} \underline{\xi} \nabla p = 0$, i.e. $\mathbf{v}_T \in \mathbf{H}(\operatorname{div} \underline{\xi} \mathbf{0}; \Omega)$.

Additionally, the sum is direct: indeed, let $\mathbf{v} \in \nabla H_0^1(\Omega) \cap \mathbf{H}(\operatorname{div} \underline{\xi} \mathbf{0}; \Omega)$, then $\mathbf{v} = \nabla p$ for some $p \in H_0^1(\Omega)$, and $\operatorname{div} \underline{\xi} \nabla p = 0$. But the Dirichlet problem is well-posed, so $p = 0$.

Finally, we note that, thanks to theorem 2.1, ∇p depends continuously on \mathbf{v} : $\|\nabla p\|_{0,\Omega} \lesssim \|\mathbf{v}\|_{0,\Omega}$.

It is also the case for \mathbf{v}_T by the triangle inequality: $\|\mathbf{v}_T\|_{0,\Omega} \leq \|\nabla p\|_{0,\Omega} + \|\mathbf{v}\|_{0,\Omega} \lesssim \|\mathbf{v}\|_{0,\Omega}$.

The second proof is similar, with bounds in $\|\cdot\|_{\mathbf{H}(\operatorname{curl}; \Omega)}$ -norm. \square

Theorem 2.3. *Under assumption (2.1), one has the second-kind Helmholtz decompositions:*

$$\mathbf{L}^2(\Omega) = \nabla H_{\operatorname{zmv}}^1(\Omega) \oplus \mathbf{H}_0(\operatorname{div} \underline{\xi} \mathbf{0}; \Omega); \quad (2.7)$$

$$\mathbf{H}(\operatorname{curl}; \Omega) = \nabla H_{\operatorname{zmv}}^1(\Omega) \oplus \mathbf{K}_T(\underline{\xi}; \Omega). \quad (2.8)$$

In addition, these sums are continuous.

Proof. Let $\mathbf{v} \in \mathbf{L}^2(\Omega)$. The Neumann problem

$$\begin{cases} \text{Find } p \in H_{\operatorname{zmv}}^1(\Omega) \text{ such that} \\ (\underline{\xi} \nabla p | \nabla q)_{0,\Omega} = (\underline{\xi} \mathbf{v} | \nabla q)_{0,\Omega}, \quad \forall q \in H_{\operatorname{zmv}}^1(\Omega), \end{cases}$$

is well-posed by theorem 2.1. We pose $\mathbf{v}_T = \mathbf{v} - \nabla p \in \mathbf{L}^2(\Omega)$. Noting that the formulation is still valid for all $q \in H^1(\Omega)$, and taking $q \in H_0^1(\Omega)$, there holds $\langle \operatorname{div} \underline{\xi} \mathbf{v}_T, q \rangle_{H_0^1(\Omega)} = -(\underline{\xi} \mathbf{v}_T | \nabla q)_{0,\Omega} = 0$. Hence $\operatorname{div} \underline{\xi} \mathbf{v}_T = 0$ and $\mathbf{v}_T \in \mathbf{H}(\operatorname{div} \underline{\xi} \mathbf{0}; \Omega)$. Moreover, $\forall q \in H^1(\Omega)$, $\langle \underline{\xi} \mathbf{v}_T \cdot \mathbf{n}, q \rangle_{H^{1/2}(\partial\Omega)} = (\underline{\xi} \mathbf{v}_T | \nabla q)_{0,\Omega} + (\operatorname{div} \underline{\xi} \mathbf{v}_T | q)_{0,\Omega} = 0$. Hence $\mathbf{v}_T \in \mathbf{H}_0(\operatorname{div} \underline{\xi} \mathbf{0}; \Omega)$.

Additionally, the sum is direct: indeed, let $\mathbf{v} \in \nabla H_{\operatorname{zmv}}^1(\Omega) \cap \mathbf{H}_0(\operatorname{div} \underline{\xi} \mathbf{0}; \Omega)$, then $\mathbf{v} = \nabla p$ for some $p \in H_{\operatorname{zmv}}^1(\Omega)$, and fulfills $\operatorname{div} \underline{\xi} \nabla p = 0$ and $\underline{\xi} \nabla p \cdot \mathbf{n}|_{\partial\Omega} = 0$. As the Neumann problem is well-posed, $p = 0$. The fact that the sum is continuous is derived as previously.

The second proof is similar. \square

Remark 2.1. We recall that the notion of orthogonality no longer applies, as the mapping $(\underline{\xi} \cdot | \cdot)_{0,\Omega}$ is not automatically a scalar product. For instance, for $\mathbf{v} \in \mathbf{H}(\operatorname{div} \underline{\xi} \mathbf{0}; \Omega)$, $q \in H_0^1(\Omega)$, it always holds $(\underline{\xi} \mathbf{v} | \nabla q)_{0,\Omega} = 0$ by integration by parts. On the other hand, $(\underline{\xi} \nabla q | \mathbf{v})_{0,\Omega}$ may not vanish.

2.3. The function space $\mathbf{X}_N(\underline{\xi}; \Omega)$

Let us begin with an extension of the first Weber inequality, cf. Theorem 6.1.6 in Ref. 2.

Theorem 2.4. *Under assumption (2.1), one has the Weber inequality*

$$\begin{aligned} \exists C_W > 0, \quad \forall \mathbf{y} \in \mathbf{X}_N(\underline{\xi}; \Omega) \\ \|\mathbf{y}\|_{0,\Omega} \leq C_W \left(\|\operatorname{curl} \mathbf{y}\|_{0,\Omega} + \|\operatorname{div} \underline{\xi} \mathbf{y}\|_{0,\Omega} + \sum_{1 \leq k \leq K} |\langle \underline{\xi} \mathbf{y} \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)}| \right). \end{aligned} \quad (2.9)$$

Proof. We proceed by contradiction. Let us assume there exists (\mathbf{y}_m) a sequence of $\mathbf{X}_N(\underline{\xi}; \Omega)$ such that, $\forall m$, $\|\mathbf{y}_m\|_{0,\Omega} = 1$ and $\|\operatorname{curl} \mathbf{y}_m\|_{0,\Omega} + \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega} + \sum_{1 \leq k \leq K} |\langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)}| \leq \frac{1}{m+1}$.

Step 1. Consider the solution to the Dirichlet problem

$$\begin{cases} \text{Find } q_m^0 \in H_0^1(\Omega) \text{ such that} \\ (\underline{\xi} \nabla q_m^0 | \nabla q)_{0,\Omega} = (\underline{\xi} \mathbf{y}_m | \nabla q)_{0,\Omega}, \quad \forall q \in H_0^1(\Omega), \end{cases} \quad (2.10)$$

By theorem 2.1, this problem is well-posed. Moreover, taking $q = q_m^0$, one gets

$$\left| (\underline{\xi} \nabla q_m^0 | \nabla q_m^0)_{0,\Omega} \right| = \left| (-\operatorname{div} \underline{\xi} \mathbf{y}_m | q_m^0)_{0,\Omega} \right| \leq \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega} \|q_m^0\|_{0,\Omega}.$$

Using the relation (2.2) on the left-hand side and Poincaré inequality on the right-hand side, one gets $\|\nabla q_m^0\|_{0,\Omega}^2 \lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega} \|\nabla q_m^0\|_{0,\Omega}$, so

$$\|\nabla q_m^0\|_{0,\Omega} \lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega}. \quad (2.11)$$

One gets that $\|\nabla q_m^0\|_{0,\Omega} \rightarrow 0$.

Step 2. Let $\mathbf{x}_m := \mathbf{y}_m - \nabla q_m^0 \in \mathbf{K}_N(\underline{\xi}; \Omega)$ (this is the Helmholtz decomposition (2.6) of \mathbf{y}_m). Consider the finite-dimensional space $Q_N(\underline{\xi}; \Omega) := \{q \in H^1(\Omega) \mid \operatorname{div} \underline{\xi} \nabla q = 0 \text{ in } \Omega, q|_{\Gamma_0} = 0, q|_{\Gamma_k} = cst_k, 1 \leq k \leq K\}$, where cst_k is a constant field on Γ_k , and the solution to the problem

$$\begin{cases} \text{Find } q_m^\Gamma \in Q_N(\underline{\xi}; \Omega) \text{ such that} \\ (\underline{\xi} \nabla q_m^\Gamma | \nabla q)_{0,\Omega} = (\underline{\xi} \mathbf{x}_m | \nabla q)_{0,\Omega}, \quad \forall q \in Q_N(\underline{\xi}; \Omega). \end{cases} \quad (2.12)$$

This problem is also well-posed, following the proof of theorem 2.1 with the Poincaré inequality in $\{q \in H^1(\Omega), q|_{\Gamma_0} = 0\}$. Taking $q = q_m^\Gamma$ and integrating by parts, one has

$$\begin{aligned} \left| (\underline{\xi} \nabla q_m^\Gamma | \nabla q_m^\Gamma)_{0,\Omega} \right| &= \left| \langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, q_m^\Gamma \rangle_{H^{1/2}(\partial\Omega)} \right| \\ &= \left| \sum_{1 \leq k \leq K} \langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, q_m^\Gamma \rangle_{H^{1/2}(\Gamma_k)} \right| \\ &= \left| \sum_{1 \leq k \leq K} \overline{q_m^\Gamma|_{\Gamma_k}} \langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)} \right|. \end{aligned}$$

As $Q_N(\underline{\xi}; \Omega)$ is a finite-dimensional vector space, all the norms are equivalent, and among them, $\|\nabla \cdot\|_{0,\Omega}$ and $\max_k |\cdot|_{\Gamma_k}$. Then, using additionally the relation (2.2), there holds

$$\|q_m^\Gamma\|_{Q_N(\underline{\xi})}^2 \lesssim \left(\|q_m^\Gamma\|_{Q_N(\underline{\xi})} \sum_{1 \leq k \leq K} |\langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)}| \right).$$

Besides, $\langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)} = \langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)} - \langle \underline{\xi} \nabla q_m^0 \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)}$, and, using the continuity of the normal trace and recalling that $\operatorname{div} \underline{\xi} \nabla q_m^0 = \operatorname{div} \underline{\xi} \mathbf{y}_m$ as well as relation (2.11), one obtains

$$\left| \langle \underline{\xi} \nabla q_m^0 \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)} \right| \lesssim \|\underline{\xi} \nabla q_m^0 \cdot \mathbf{n}\|_{H^{-1/2}(\Gamma_k)} \lesssim \|\nabla q_m^0\|_{\mathbf{H}(\operatorname{div} \underline{\xi}; \Omega)} \lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega}.$$

Hence

$$\|q_m^\Gamma\|_{Q_N(\underline{\xi})} \lesssim \left(\sum_{1 \leq k \leq K} |\langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)}| + \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega} \right), \quad (2.13)$$

and one gets that $\|\nabla q_m^\Gamma\|_{0,\Omega} \rightarrow 0$. Observe that $\nabla q_m^\Gamma \times \mathbf{n}|_{\partial\Omega} = 0$, so $\nabla q_m^\Gamma \in \mathbf{H}_0(\mathbf{curl}; \Omega)$.

Step 3. Let $\mathbf{z}_m := \mathbf{y}_m - \nabla q_m^0 - \nabla q_m^\Gamma$. It belongs to $\mathbf{X}_N(\underline{\xi}; \Omega)$ and, additionally, $\mathbf{curl} \mathbf{z}_m = \mathbf{curl} \mathbf{y}_m$, $\operatorname{div} \underline{\xi} \mathbf{z}_m = 0$, and $\langle \underline{\xi} \mathbf{z}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)} = 0$. Indeed, introducing q_k the unique element of $Q_N(\underline{\xi}; \Omega)$ such that $q_k|_{\Gamma_l} = \delta_{k,l}$ for $1 \leq k \leq K$, one has by integration by parts

$$\langle \underline{\xi} \mathbf{z}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)} = \langle \underline{\xi} \mathbf{z}_m \cdot \mathbf{n}, q_k \rangle_{H^{1/2}(\partial\Omega)} = (\underline{\xi} \mathbf{z}_m | \nabla q_k)_{0,\Omega} + (\operatorname{div} \underline{\xi} \mathbf{z}_m | q_k)_{0,\Omega}.$$

However, $(\underline{\xi} \mathbf{z}_m | \nabla q_k)_{0,\Omega} = (\underline{\xi} \mathbf{x}_m | \nabla q_k)_{0,\Omega} - (\underline{\xi} \nabla q_m^\Gamma | \nabla q_k)_{0,\Omega} = 0$ by definition of q_m^Γ ; and $\operatorname{div} \underline{\xi} \mathbf{z}_m = 0$. It follows that $\langle \underline{\xi} \mathbf{z}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)} = 0$. With the help of the vector potential Theorem 3.4.1², there exists $\mathbf{w}_m \in \mathbf{H}^1(\Omega)$ such that $\underline{\xi} \mathbf{z}_m = \mathbf{curl} \mathbf{w}_m$, and $\|\mathbf{w}_m\|_{1,\Omega} \lesssim \|\underline{\xi} \mathbf{z}_m\|_{0,\Omega} \lesssim \|\mathbf{z}_m\|_{0,\Omega}$.

Furthermore, we have by integration by parts $(z_m | \underline{\xi} z_m)_{0,\Omega} = (z_m | \mathbf{curl} \mathbf{w}_m)_{0,\Omega} = (\mathbf{curl} z_m | \mathbf{w}_m)_{0,\Omega} = (\mathbf{curl} \mathbf{y}_m | \mathbf{w}_m)_{0,\Omega}$. Using again relation (2.2), there holds

$$\|z_m\|_{0,\Omega}^2 \lesssim \|\mathbf{curl} \mathbf{y}_m\|_{0,\Omega} \|\mathbf{w}_m\|_{0,\Omega} \lesssim \|\mathbf{curl} \mathbf{y}_m\|_{0,\Omega} \|z_m\|_{0,\Omega},$$

and so $\|z_m\|_{0,\Omega} \lesssim \|\mathbf{curl} \mathbf{y}_m\|_{0,\Omega}$. One gets that $\|z_m\|_{0,\Omega} \rightarrow 0$.

Finally, as $\mathbf{y}_m = z_m + \nabla q_m^0 + \nabla q_m^\Gamma$, we have $\|\mathbf{y}_m\|_{0,\Omega} \rightarrow 0$, which contradicts $\|\mathbf{y}_m\|_{0,\Omega} = 1$ for all m . \square

Similarly, one can also extend the compact embedding result of Theorem 7.5.1 in Ref. 2.

Theorem 2.5. *Under assumption (2.1), the embedding of $\mathbf{X}_N(\underline{\xi}; \Omega)$ into $\mathbf{L}^2(\Omega)$ is compact.*

Proof. Let (\mathbf{y}_m) be a bounded sequence of $\mathbf{X}_N(\underline{\xi}; \Omega)$. As in the previous proof (Steps 1., 2., 3.), we introduce $q_m^0 \in H_0^1(\Omega)$, $q_m^\Gamma \in Q_N(\underline{\xi}; \Omega)$, and $\mathbf{w}_m \in \mathbf{H}^1(\Omega)$, such that $\mathbf{y}_m = \underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_m + \nabla q_m^0 + \nabla q_m^\Gamma$. Additionally, there holds (see previous proof):

$$\begin{aligned} \|\nabla q_m^0\|_{0,\Omega} &\lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega}; \\ \|\nabla q_m^\Gamma\|_{0,\Omega} &\lesssim \left(\|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega} + \sum_{1 \leq k \leq K} |\langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{H^{1/2}(\Gamma_k)}| \right); \\ \|\mathbf{w}_m\|_{1,\Omega} &\lesssim \|\mathbf{curl} \mathbf{y}_m\|_{0,\Omega}. \end{aligned}$$

Let us begin with (q_m^Γ) : it is a bounded sequence of the finite-dimensional vector space $Q_N(\underline{\xi}; \Omega)$, so it admits a subsequence which converges (in particular in $\|\cdot\|_{H^1(\Omega)}$ -norm).

Besides, (q_m^0) and (\mathbf{w}_m) are bounded sequences of $H^1(\Omega)$ (resp. $\mathbf{H}^1(\Omega)$). Then, by Rellich's theorem, they admit subsequences (still denoted with the same indices) which converge in $L^2(\Omega)$ (resp. $\mathbf{L}^2(\Omega)$). It remains to prove that both subsequences (∇q_m^0) and $(\mathbf{curl} \mathbf{w}_m)$ converge in $\mathbf{L}^2(\Omega)$.

By definition of q_m^0 , for any q in $H_0^1(\Omega)$, there holds by integration by parts

$$(\underline{\xi} \nabla q_m^0 | \nabla q)_{0,\Omega} = (\underline{\xi} \mathbf{y}_m | \nabla q)_{0,\Omega} = -(\operatorname{div} \underline{\xi} \mathbf{y}_m | q)_{0,\Omega}.$$

Using the notation $v_{mn} := v_m - v_n$ and taking $q = q_{mn}^0$, one has

$$\left| (\underline{\xi} \nabla q_{mn}^0 | \nabla q_{mn}^0)_{0,\Omega} \right| \leq \|\operatorname{div} \underline{\xi} \mathbf{y}_{mn}\|_{0,\Omega} \|q_{mn}^0\|_{0,\Omega}.$$

Then, by relation (2.2),

$$\|\nabla q_{mn}^0\|_{0,\Omega}^2 \lesssim 2 \sup_m (\|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega}) \|q_{mn}^0\|_{0,\Omega}.$$

Thus (∇q_m^0) is a Cauchy sequence of $\mathbf{L}^2(\Omega)$, hence converges in this Hilbert space.

We recall that $\underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_m \in \mathbf{X}_N(\underline{\xi}; \Omega)$ (cf. Step 3. of previous proof) and $\mathbf{curl}(\underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_m) = \mathbf{curl} \mathbf{y}_m$. Then, still with the same notations, and by integration by parts,

$$(\underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_{mn} | \mathbf{curl} \mathbf{w}_{mn})_{0,\Omega} = (\mathbf{curl}(\underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_{mn}) | \mathbf{w}_{mn})_{0,\Omega} = (\mathbf{curl} \mathbf{y}_{mn} | \mathbf{w}_{mn})_{0,\Omega}.$$

As $\underline{\xi}^{-1}$ also satisfies the ellipticity condition (proposition 2.1), we get

$$\|\mathbf{curl} \mathbf{w}_{mn}\|_{0,\Omega}^2 \lesssim \|\mathbf{curl} \mathbf{y}_{mn}\|_{0,\Omega} \|\mathbf{w}_{mn}\|_{0,\Omega} \leq 2 \sup_m (\|\mathbf{curl} \mathbf{y}_m\|_{0,\Omega}) \|\mathbf{w}_{mn}\|_{0,\Omega},$$

which proves that $(\mathbf{curl} \mathbf{w}_m)$ is a Cauchy, hence converging, sequence of $\mathbf{L}^2(\Omega)$.

As $\mathbf{y}_m = \underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_m + \nabla q_m^0 + \nabla q_m^\Gamma$, we conclude that the subsequence (\mathbf{y}_m) converges in $\mathbf{L}^2(\Omega)$. \square

2.4. The function space $X_T(\underline{\xi}; \Omega)$

Let us begin with an extension of the second Weber inequality, cf. Theorem 6.2.5 in Ref. 2. Some knowledge on the topology of the domain Ω is required, see the Appendix for details and notations.

Theorem 2.6. *Assume that $(\mathbf{Top})_I$ holds. Under assumption (2.1), one has the Weber inequality*

$$\begin{aligned} & \exists C'_W > 0, \forall \mathbf{X}_T(\underline{\xi}; \Omega) \\ & \|\mathbf{y}\|_{0,\Omega} \leq C'_W \left(\|\mathbf{curl} \mathbf{y}\|_{0,\Omega} + \|\operatorname{div} \underline{\xi} \mathbf{y}\|_{0,\Omega} + \sum_{1 \leq i \leq I} |\langle \underline{\xi} \mathbf{y} \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}| \right). \end{aligned} \quad (2.14)$$

Proof. The proof follows a similar structure as in theorem 2.4. By contradiction, we assume there exists (\mathbf{y}_m) a sequence of $\mathbf{X}_T(\underline{\xi}; \Omega)$ such that, $\forall m$, $\|\mathbf{y}_m\|_{0,\Omega} = 1$ and $\|\mathbf{curl} \mathbf{y}_m\|_{0,\Omega} + \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega} + \sum_{1 \leq i \leq I} |\langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}| \leq \frac{1}{m+1}$.

Step 1. Consider the solution to the Neumann problem

$$\begin{cases} \text{Find } q_m^0 \in H_{\text{zmv}}^1(\Omega) \text{ such that} \\ (\underline{\xi} \nabla q_m^0 | \nabla q)_{0,\Omega} = (\underline{\xi} \mathbf{y}_m | \nabla q)_{0,\Omega}, \forall q \in H_{\text{zmv}}^1(\Omega). \end{cases} \quad (2.15)$$

The problem is well-posed by theorem 2.1. Taking $q = q_m^0$ and integrating by parts, one gets, as $\mathbf{y}_m \in \mathbf{H}_0(\operatorname{div} \underline{\xi}; \Omega)$:

$$\left| (\underline{\xi} \nabla q_m^0 | \nabla q_m^0)_{0,\Omega} \right| = \left| (-\operatorname{div} \underline{\xi} \mathbf{y}_m | q_m^0)_{0,\Omega} \right| \leq \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega} \|q_m^0\|_{0,\Omega}.$$

Using the relation (2.2) on the left-hand side, as well as the Poincaré-Wirtinger inequality on the right-hand side, leads to

$$\|\nabla q_m^0\|_{0,\Omega} \lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega}, \quad (2.16)$$

so $\|\nabla q_m^0\|_{0,\Omega} \rightarrow 0$.

Step 2. With the help of the second-kind Helmholtz decomposition (2.8) of \mathbf{y}_m , we define $\mathbf{x}_m := \mathbf{y}_m - \nabla q_m^0 \in \mathbf{K}_T(\underline{\xi}; \Omega)$. Note that $\nabla q_m^0 \in \mathbf{H}_0(\operatorname{div} \underline{\xi}; \Omega)$. Consider the finite-dimensional space

$$Q_T(\underline{\xi}; \dot{\Omega}) := \{ \dot{q} \in H_{\text{zmv}}^1(\dot{\Omega}) \mid \operatorname{div} \underline{\xi} \widetilde{\nabla} \dot{q} = 0 \text{ in } \Omega, \underline{\xi} \widetilde{\nabla} \dot{q} \cdot \mathbf{n}|_{\partial\Omega} = 0, [\dot{q}]_{\Sigma_i} = cst_i, 1 \leq i \leq I \},$$

where cst_i is a constant field on Σ_i . Let us introduce the problem

$$\begin{cases} \text{Find } \dot{q}_m^\Sigma \in Q_T(\underline{\xi}; \dot{\Omega}) \text{ such that} \\ (\underline{\xi} \nabla \dot{q}_m^\Sigma | \nabla \dot{q})_{0,\dot{\Omega}} = (\underline{\xi} \mathbf{x}_m | \nabla \dot{q})_{0,\dot{\Omega}} \quad \forall \dot{q} \in Q_T(\underline{\xi}; \dot{\Omega}). \end{cases} \quad (2.17)$$

This problem is well-posed, adapting the proof of theorem 2.1, and using Poincaré-Wirtinger inequality in $H_{\text{zmv}}^1(\dot{\Omega})$. Taking $\dot{q} = \dot{q}_m^\Sigma$ and integrating by parts (A.1), one has, as $\operatorname{div} \underline{\xi} \mathbf{x}_m = 0$,

$$(\underline{\xi} \nabla \dot{q}_m^\Sigma | \nabla \dot{q}_m^\Sigma)_{0,\dot{\Omega}} = \sum_{1 \leq i \leq I} \left\langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, [\dot{q}_m^\Sigma]_{\Sigma_i} \right\rangle_{\Sigma_i} = \sum_{1 \leq i \leq I} \overline{[\dot{q}_m^\Sigma]_{\Sigma_i}} \langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}.$$

As $Q_T(\underline{\xi}; \dot{\Omega})$ is a finite-dimensional vector space, all the norms are equivalent, and among them, $\|\widetilde{\nabla} \cdot\|_{0,\Omega}$ and $\max_i |[\cdot]_{\Sigma_i}|$. Then, using additionally relation (2.2), there holds

$$\|\dot{q}_m^\Sigma\|_{Q_T(\underline{\xi}; \dot{\Omega})}^2 \lesssim \left(\|\dot{q}_m^\Sigma\|_{Q_T(\underline{\xi}; \dot{\Omega})} \sum_{1 \leq i \leq I} |\langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}| \right).$$

Besides, $\langle \underline{\xi} \mathbf{x}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i} = \langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i} - \langle \underline{\xi} \nabla q_m^0 \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}$. For $1 \leq i \leq I$, let \dot{q}_i be the unique element of $Q_T(\underline{\xi}; \dot{\Omega})$ such that $[\dot{q}_i]_{\Sigma_j} = \delta_{i,j}$ for $1 \leq j \leq I$. If one recalls that $\nabla q_m^0 \in \mathbf{H}_0(\operatorname{div} \underline{\xi}; \Omega)$, then there holds, by integration by parts (A.1),

$$\begin{aligned} |\langle \underline{\xi} \nabla q_m^0 \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}| &= \left| \sum_{1 \leq j \leq I} \langle \underline{\xi} \nabla q_m^0 \cdot \mathbf{n}, [\dot{q}_i]_{\Sigma_j} \rangle_{\Sigma_j} \right| = \left| \left(\underline{\xi} \nabla q_m^0 | \nabla \dot{q}_i \right)_{0, \dot{\Omega}} + (\operatorname{div} \underline{\xi} \nabla q_m^0 | \dot{q}_i)_{0, \dot{\Omega}} \right| \\ &\lesssim \|\nabla q_m^0\|_{0, \dot{\Omega}} \|\nabla \dot{q}_i\|_{0, \dot{\Omega}} + \|\operatorname{div} \underline{\xi} \nabla q_m^0\|_{0, \dot{\Omega}} \|\dot{q}_i\|_{0, \dot{\Omega}} \\ &\lesssim \|\nabla q_m^0\|_{\mathbf{H}(\operatorname{div} \underline{\xi}; \Omega)} \\ &\lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0, \Omega}, \end{aligned}$$

the latter because of (2.16) and $\operatorname{div} \underline{\xi} \nabla q_m^0 = \operatorname{div} \underline{\xi} \mathbf{y}_m$ in Ω . Hence,

$$\|\dot{q}_m^\Sigma\|_{Q_T(\underline{\xi}; \dot{\Omega})} \lesssim \left(\sum_{1 \leq i \leq I} |\langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}| + \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0, \Omega} \right), \quad (2.18)$$

which implies that $\|\widehat{\nabla \dot{q}_m^\Sigma}\|_{0, \Omega} = \|\dot{q}_m^\Sigma\|_{Q_T(\underline{\xi}; \dot{\Omega})} \rightarrow 0$.

Step 3. Let $\mathbf{z}_m := \mathbf{y}_m - \nabla q_m^0 - \widehat{\nabla \dot{q}_m^\Sigma} \in \mathbf{X}_T(\underline{\xi}; \Omega)$. There holds $\operatorname{curl} \mathbf{z}_m = \operatorname{curl} \mathbf{y}_m$, $\operatorname{div} \underline{\xi} \mathbf{z}_m = 0$, and additionally $\langle \underline{\xi} \mathbf{z}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i} = 0$ for $1 \leq i \leq I$. First, one has by integration by parts (A.1)

$$\langle \underline{\xi} \mathbf{z}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i} = \sum_{1 \leq j \leq I} \langle \underline{\xi} \mathbf{z}_m \cdot \mathbf{n}, [\dot{q}_i]_{\Sigma_j} \rangle_{\Sigma_j} = (\underline{\xi} \mathbf{z}_m | \nabla \dot{q}_i)_{0, \dot{\Omega}} + (\operatorname{div} \underline{\xi} \mathbf{z}_m | \dot{q}_i)_{0, \dot{\Omega}}.$$

Then, $(\underline{\xi} \mathbf{z}_m | \nabla \dot{q}_i)_{0, \dot{\Omega}} = (\underline{\xi} \mathbf{x}_m | \nabla \dot{q}_i)_{0, \dot{\Omega}} - (\underline{\xi} \nabla \dot{q}_m^\Sigma | \nabla \dot{q}_i)_{0, \dot{\Omega}} = 0$ by definition of \dot{q}_m^Σ , while $\operatorname{div} \underline{\xi} \mathbf{z}_m = 0$ by definition of \mathbf{z}_m .

One has also $\mathbf{z}_m \in \mathbf{H}_0(\operatorname{div} \underline{\xi}; \Omega)$ so, according to the vector potential Theorem 3.5.1²: there exists $\mathbf{w}_m \in \mathbf{X}_N(\Omega)$ such that $\underline{\xi} \mathbf{z}_m = \operatorname{curl} \mathbf{w}_m$ and $\operatorname{div} \mathbf{w}_m = 0$ in Ω , and $\|\mathbf{w}_m\|_{\mathbf{H}(\operatorname{curl}; \Omega)} \lesssim \|\underline{\xi} \mathbf{z}_m\|_{0, \Omega}$; in particular, $\|\mathbf{w}_m\|_{\mathbf{H}(\operatorname{curl}; \Omega)} \lesssim \|\mathbf{z}_m\|_{0, \Omega}$.

We have by integration by parts $(\mathbf{z}_m | \underline{\xi} \mathbf{z}_m)_{0, \Omega} = (\mathbf{z}_m | \operatorname{curl} \mathbf{w}_m)_{0, \Omega} = (\operatorname{curl} \mathbf{z}_m | \mathbf{w}_m)_{0, \Omega} = (\operatorname{curl} \mathbf{y}_m | \mathbf{w}_m)_{0, \Omega}$. Using again the relation (2.2), we find

$$\|\mathbf{z}_m\|_{0, \Omega}^2 \lesssim \|\operatorname{curl} \mathbf{y}_m\|_{0, \Omega} \|\mathbf{w}_m\|_{0, \Omega} \lesssim \|\operatorname{curl} \mathbf{y}_m\|_{0, \Omega} \|\mathbf{z}_m\|_{0, \Omega},$$

so that $\|\mathbf{z}_m\|_{0, \Omega} \lesssim \|\operatorname{curl} \mathbf{y}_m\|_{0, \Omega}$. It follows that $\|\mathbf{z}_m\|_{0, \Omega} \rightarrow 0$.

As $\mathbf{y}_m = \mathbf{z}_m + \nabla q_m^0 + \widehat{\nabla \dot{q}_m^\Sigma}$, we conclude that $\|\mathbf{y}_m\|_{0, \Omega} \rightarrow 0$, but that contradicts $\|\mathbf{y}_m\|_{0, \Omega} = 1$ for all m . \square

One can also extend the compact embedding result of Theorem 7.5.3 in Ref. 2.

Theorem 2.7. *Assume that $(\mathbf{Top})_I$ holds. Under assumption (2.1), the embedding of $\mathbf{X}_T(\underline{\xi}; \Omega)$ into $L^2(\Omega)$ is compact.*

Proof. Let (\mathbf{y}_m) be a bounded sequence of $\mathbf{X}_T(\underline{\xi}; \Omega)$. As in the previous proof (Steps 1., 2., 3.), we introduce $q_m^0 \in H_{\operatorname{zmv}}^1(\Omega)$, $\dot{q}_m^\Sigma \in Q_T(\underline{\xi}; \dot{\Omega})$, and $\mathbf{w}_m \in \mathbf{X}_N(\Omega)$, such that $\mathbf{y}_m = \underline{\xi}^{-1} \operatorname{curl} \mathbf{w}_m + \nabla q_m^0 + \widehat{\nabla \dot{q}_m^\Sigma}$. Additionally, there holds (see previous proof):

$$\begin{aligned} \|\nabla q_m^0\|_{0, \Omega} &\lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0, \Omega}; \\ \|\widehat{\nabla \dot{q}_m^\Sigma}\|_{0, \Omega} &\lesssim \left(\|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0, \Omega} + \sum_i |\langle \underline{\xi} \mathbf{y}_m \cdot \mathbf{n}, 1 \rangle_{\Sigma_i}| \right); \\ \|\mathbf{w}_m\|_{\mathbf{H}(\operatorname{curl}; \Omega)} &\lesssim \|\operatorname{curl} \mathbf{y}_m\|_{0, \Omega}. \end{aligned}$$

Let us begin with (\hat{q}_m^Σ) : it is a bounded sequence of the finite-dimensional vector space $Q_T(\underline{\xi}; \hat{\Omega})$, so it admits a converging subsequence (in particular in $\|\cdot\|_{H^1(\Omega)}$ -norm).

Besides, (q_m^0) is a bounded sequence of $H^1(\Omega)$ so, by Rellich's theorem, it admits a converging subsequence (denoted with the same index) in $L^2(\Omega)$. Similarly, \mathbf{w}_m is a bounded sequence of $\mathbf{X}_N(\Omega)$, so by theorem 2.5, it admits a converging subsequence in $\mathbf{L}^2(\Omega)$. It remains to prove that the subsequences (∇q_m^0) and $(\mathbf{curl} \mathbf{w}_m)$ converge in $\mathbf{L}^2(\Omega)$.

By definition of q_m^0 , for any q in $H_0^1(\Omega)$, there holds by integration by parts

$$(\underline{\xi} \nabla q_m^0 | \nabla q)_{0,\Omega} = (\underline{\xi} \mathbf{y}_{mn} | \nabla q)_{0,\Omega} = -(\operatorname{div} \underline{\xi} \mathbf{y}_{mn} | q)_{0,\Omega}.$$

Taking $q = q_m^0$, one gets by property (2.1)

$$\|\nabla q_m^0\|_{0,\Omega}^2 \lesssim \|\operatorname{div} \underline{\xi} \mathbf{y}_{mn}\|_{0,\Omega} \|q_m^0\|_{0,\Omega} \leq 2 \sup_m (\|\operatorname{div} \underline{\xi} \mathbf{y}_m\|_{0,\Omega}) \|q_m^0\|_{0,\Omega}.$$

Thus (∇q_m^0) is a Cauchy sequence of $\mathbf{L}^2(\Omega)$, hence it converges in this Hilbert space.

By integration by parts (recall that $\mathbf{w}_{mn} \in \mathbf{X}_N(\Omega)$),

$$(\underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_{mn} | \mathbf{curl} \mathbf{w}_{mn})_{0,\Omega} = (\mathbf{curl}(\underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_{mn}) | \mathbf{w}_{mn})_{0,\Omega} = (\mathbf{curl} \mathbf{y}_{mn} | \mathbf{w}_{mn})_{0,\Omega}.$$

As $\underline{\xi}^{-1}$ satisfies the ellipticity condition (proposition 2.1), we find

$$\|\mathbf{curl} \mathbf{w}_{mn}\|_{0,\Omega}^2 \lesssim \|\mathbf{curl} \mathbf{y}_{mn}\|_{0,\Omega} \|\mathbf{w}_{mn}\|_{0,\Omega} \leq 2 \sup_m (\|\mathbf{curl} \mathbf{y}_m\|_{0,\Omega}) \|\mathbf{w}_{mn}\|_{0,\Omega},$$

which proves that $(\mathbf{curl} \mathbf{w}_m)$ is also a Cauchy, hence converging, sequence of $\mathbf{L}^2(\Omega)$.

As $\mathbf{y}_m = \underline{\xi}^{-1} \mathbf{curl} \mathbf{w}_m + \nabla q_m^0 + \widehat{\nabla} \hat{q}_m^\Sigma$, the subsequence (\mathbf{y}_m) converges in $\mathbf{L}^2(\Omega)$. \square

3. Analysis of the Dirichlet problem

In this section, we supplement the time-harmonic Maxwell's equation by a Dirichlet boundary condition on $\partial\Omega$:

$$\begin{cases} \text{Find } \mathbf{E} \in \mathbf{H}(\mathbf{curl}; \Omega) \text{ such that} \\ \mathbf{curl}(\underline{\mu}^{-1} \mathbf{curl} \mathbf{E}) - \omega^2 \underline{\varepsilon} \mathbf{E} = \mathbf{f} \text{ in } \Omega, \\ \mathbf{E} \times \mathbf{n} = \mathbf{g} \text{ on } \partial\Omega, \end{cases} \quad (3.1)$$

where \mathbf{g} is a boundary data.

We assume that the tensors $\underline{\varepsilon}, \underline{\mu} \in \mathbf{L}^\infty(\Omega)$ are elliptic. We also assume that the volume data \mathbf{f} belongs to $\mathbf{L}^2(\Omega)$, and that the surface data \mathbf{g} is the tangential trace of a field $\mathbf{E}_d \in \mathbf{H}(\mathbf{curl}; \Omega)$, that is $\mathbf{g} = \mathbf{E}_d \times \mathbf{n}|_{\partial\Omega}$. Further assumptions will be made to obtain the extra-regularity results.

3.1. Variational formulation and well-posedness

Let us derive the variational formulation of problem (3.1). In order to deal with a problem with homogeneous boundary condition, we introduce the new unknown $\mathbf{E}_0 := \mathbf{E} - \mathbf{E}_d$. It belongs to $\mathbf{H}_0(\mathbf{curl}; \Omega)$ and, additionally, satisfies

$$\mathbf{curl}(\underline{\mu}^{-1} \mathbf{curl} \mathbf{E}_0) - \omega^2 \underline{\varepsilon} \mathbf{E}_0 = \mathbf{f} - \mathbf{curl}(\underline{\mu}^{-1} \mathbf{curl} \mathbf{E}_d) + \omega^2 \underline{\varepsilon} \mathbf{E}_d \quad \text{in } \Omega.$$

By standard techniques, we get the equivalent variational formulation for \mathbf{E}_0 :

$$\begin{cases} \text{Find } \mathbf{E}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega) \text{ such that} \\ (\underline{\mu}^{-1} \mathbf{curl} \mathbf{E}_0 | \mathbf{curl} \mathbf{F})_{0,\Omega} - \omega^2 (\underline{\varepsilon} \mathbf{E}_0 | \mathbf{F})_{0,\Omega} = \ell_{\mathbf{D},0}(\mathbf{F}), \quad \forall \mathbf{F} \in \mathbf{H}_0(\mathbf{curl}; \Omega). \end{cases} \quad (3.2)$$

Above, $\ell_{D,0} : \mathbf{F} \mapsto (\mathbf{f} + \omega^2 \underline{\boldsymbol{\varepsilon}} \mathbf{E}_d | \mathbf{F})_{0,\Omega} - (\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E}_d | \mathbf{curl} \mathbf{F})_{0,\Omega}$ belongs to $(\mathbf{H}_0(\mathbf{curl}; \Omega))'$. In addition, we observe that

$$\|\ell_{D,0}\|_{(\mathbf{H}_0(\mathbf{curl}; \Omega))'} \lesssim \|\mathbf{f}\|_{0,\Omega} + \|\mathbf{E}_d\|_{\mathbf{H}(\mathbf{curl}; \Omega)}. \quad (3.3)$$

Remark 3.1. In terms of the total field \mathbf{E} , the variational formulation to be solved is

$$\begin{cases} \text{Find } \mathbf{E} \in \mathbf{H}(\mathbf{curl}; \Omega) \text{ such that} \\ (\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E} | \mathbf{curl} \mathbf{F})_{0,\Omega} - \omega^2 (\underline{\boldsymbol{\varepsilon}} \mathbf{E} | \mathbf{F})_{0,\Omega} = \ell_D(\mathbf{F}), \quad \forall \mathbf{F} \in \mathbf{H}_0(\mathbf{curl}; \Omega), \\ \mathbf{E} \times \mathbf{n} = \mathbf{g} \text{ on } \partial\Omega, \end{cases} \quad (3.4)$$

where $\ell_D : \mathbf{F} \mapsto (\mathbf{f} | \mathbf{F})_{0,\Omega}$ belongs to $(\mathbf{H}_0(\mathbf{curl}; \Omega))'$. This formulation can be used to solve the non-homogeneous Dirichlet problem numerically, see section 5.

To study the well-posedness of the formulation (3.2), it is useful to introduce an equivalent problem with the help of the previously derived first-kind Helmholtz decomposition (2.6).

Lemma 3.1. *The formulation (3.2) can be equivalently recast as: set $\mathbf{E}_0 = \tilde{\mathbf{E}} + \nabla p$, with $p \in H_0^1(\Omega)$ and $\tilde{\mathbf{E}} \in \mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega)$, respectively governed by*

$$\begin{cases} \text{Find } p \in H_0^1(\Omega) \text{ such that} \\ -\omega^2 (\underline{\boldsymbol{\varepsilon}} \nabla p | \nabla q)_{0,\Omega} = \ell_{D,0}(\nabla q)_{0,\Omega}, \quad \forall q \in H_0^1(\Omega), \end{cases} \quad (3.5)$$

and

$$\begin{cases} \text{Find } \tilde{\mathbf{E}} \in \mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega) \text{ such that} \\ (\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \tilde{\mathbf{E}} | \mathbf{curl} \tilde{\mathbf{F}})_{0,\Omega} - \omega^2 (\underline{\boldsymbol{\varepsilon}} \tilde{\mathbf{E}} | \tilde{\mathbf{F}})_{0,\Omega} \\ = \omega^2 (\underline{\boldsymbol{\varepsilon}} \nabla p | \tilde{\mathbf{F}})_{0,\Omega} + \ell_{D,0}(\tilde{\mathbf{F}}), \quad \forall \tilde{\mathbf{F}} \in \mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega). \end{cases} \quad (3.6)$$

Proof. Direct. Let us introduce the first-kind Helmholtz decomposition (2.6) of \mathbf{E}_0 : $\mathbf{E}_0 = \tilde{\mathbf{E}} + \nabla p$, with $p \in H_0^1(\Omega)$ and $\tilde{\mathbf{E}} \in \mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega)$. Taking $\mathbf{F} = \nabla q$ for any $q \in H_0^1(\Omega)$ in (3.2), we get

$$-\omega^2 (\underline{\boldsymbol{\varepsilon}} (\tilde{\mathbf{E}} + \nabla p) | \nabla q)_{0,\Omega} = \ell_{D,0}(\nabla q)$$

and, since $\tilde{\mathbf{E}} \in \mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega)$, it holds that, $\forall q \in H_0^1(\Omega)$, $(\underline{\boldsymbol{\varepsilon}} \tilde{\mathbf{E}} | \nabla q)_{0,\Omega} = 0$, so p is governed by (3.5). On the other hand, for $\tilde{\mathbf{E}} = \mathbf{E}_0 - \nabla p$, one has

$$(\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \tilde{\mathbf{E}} | \mathbf{curl} \mathbf{F})_{0,\Omega} - \omega^2 (\underline{\boldsymbol{\varepsilon}} \tilde{\mathbf{E}} | \mathbf{F})_{0,\Omega} = \omega^2 (\underline{\boldsymbol{\varepsilon}} \nabla p | \mathbf{F})_{0,\Omega} + \ell_{D,0}(\mathbf{F})$$

for any $\mathbf{F} \in \mathbf{H}_0(\mathbf{curl}; \Omega)$, hence in particular for any $\tilde{\mathbf{F}} \in \mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega)$: $\tilde{\mathbf{E}}$ is governed by (3.6).

Reverse. By summation one has, $\forall q \in H_0^1(\Omega)$, $\forall \tilde{\mathbf{F}} \in \mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega)$:

$$(\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \tilde{\mathbf{E}} | \mathbf{curl} \tilde{\mathbf{F}})_{0,\Omega} - \omega^2 (\underline{\boldsymbol{\varepsilon}} \tilde{\mathbf{E}} | \tilde{\mathbf{F}})_{0,\Omega} - \omega^2 (\underline{\boldsymbol{\varepsilon}} \nabla p | \nabla q)_{0,\Omega} = \omega^2 (\underline{\boldsymbol{\varepsilon}} \nabla p | \tilde{\mathbf{F}})_{0,\Omega} + \ell_{D,0}(\tilde{\mathbf{F}}) + \ell_{D,0}(\nabla q).$$

One can add the vanishing terms $(\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \nabla p | \mathbf{curl} \tilde{\mathbf{F}})_{0,\Omega}$, $(\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} (\tilde{\mathbf{E}} + \nabla p) | \mathbf{curl} \nabla q)_{0,\Omega}$ and $-\omega^2 (\underline{\boldsymbol{\varepsilon}} \tilde{\mathbf{E}} | \nabla q)_{0,\Omega}$ to the left-hand side. Introducing $\mathbf{E}_0 := \tilde{\mathbf{E}} + \nabla p \in \mathbf{H}_0(\mathbf{curl}; \Omega)$, one gets, after simple rearrangements:

$$(\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E}_0 | \mathbf{curl} (\tilde{\mathbf{F}} + \nabla q))_{0,\Omega} - \omega^2 (\underline{\boldsymbol{\varepsilon}} \mathbf{E}_0 | \tilde{\mathbf{F}} + \nabla q)_{0,\Omega} = \ell_{D,0}(\tilde{\mathbf{F}} + \nabla q).$$

Finally, as $\tilde{\mathbf{F}}$ and q span respectively $\mathbf{K}_N(\underline{\boldsymbol{\varepsilon}}; \Omega)$ and $H_0^1(\Omega)$, the sum $\tilde{\mathbf{F}} + \nabla q$ spans the whole $\mathbf{H}_0(\mathbf{curl}; \Omega)$, thanks to (2.6): \mathbf{E}_0 is governed by (3.2). \square

Remark 3.2. The term $\omega^2 (\underline{\varepsilon} \nabla p | \tilde{\mathbf{F}})_{0,\Omega}$ in formulation (3.6) vanishes automatically *only* if $\underline{\varepsilon}$ is a hermitian tensor field.

Then, we have the following results:

Lemma 3.2. *The formulation (3.5) is well-posed. Moreover, one has the bound $\|p\|_{1,\Omega} \lesssim \|\mathbf{f}\|_{0,\Omega} + \|\mathbf{E}_d\|_{0,\Omega}$.*

Proof. Well-posedness is an immediate consequence of theorem 2.1. Since $\ell_{D0} \in (\mathbf{H}_0(\mathbf{curl}; \Omega))'$, and ∇ is a continuous mapping from $H_0^1(\Omega)$ to $\mathbf{H}_0(\mathbf{curl}; \Omega)$, one has $\ell_{D,0} \circ \nabla \in (H_0^1(\Omega))'$. The bound on $\|p\|_{1,\Omega}$ is a straightforward consequence of the expression $\ell_{D,0}(\nabla q) = (\mathbf{f} + \omega^2 \underline{\varepsilon} \mathbf{E}_d | \nabla q)_{0,\Omega}$. \square

Recall that $\mathbf{K}_N(\underline{\varepsilon}; \Omega)$ is equipped with the norm $\|\cdot\|_{\mathbf{H}(\mathbf{curl}; \Omega)}$.

Lemma 3.3. *The formulation (3.6) fits the coercive + compact framework:*

- *either the problem (3.6) admits a unique solution $\tilde{\mathbf{E}}$ in $\mathbf{K}_N(\underline{\varepsilon}; \Omega)$, which depends continuously on the data \mathbf{f} and \mathbf{E}_d :*

$$\|\tilde{\mathbf{E}}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \lesssim \|\mathbf{f}\|_{0,\Omega} + \|\mathbf{E}_d\|_{\mathbf{H}(\mathbf{curl}; \Omega)};$$

- *or, the problem (3.6) has solutions if, and only if, \mathbf{f} and \mathbf{E}_d satisfy a finite number of compatibility conditions; in this case, the space of solutions is an affine space of finite dimension. Additionally, the component of the solution which is orthogonal (in the sense of the $\mathbf{H}_0(\mathbf{curl}; \Omega)$ scalar product) to the corresponding linear vector space, depends continuously on the data \mathbf{f} and \mathbf{E}_d .*

Proof. Let us split the left-hand side of (3.6) in two terms. Let $\alpha > 0$, recalling that $\underline{\mu}^{-1}$ satisfies assumption (2.1), and using the notations of proposition 2.1, we introduce two sesquilinear forms, namely $a : (\mathbf{u}, \mathbf{v}) \mapsto (\underline{\mu}^{-1} \mathbf{curl} \mathbf{u} | \mathbf{curl} \mathbf{v})_{0,\Omega} + \alpha e^{i\theta\mu} (\mathbf{u} | \mathbf{v})_{0,\Omega}$, and $b : (\mathbf{u}, \mathbf{v}) \mapsto (\underline{\varepsilon}' \mathbf{u} | \mathbf{v})_{0,\Omega}$ with $\underline{\varepsilon}' := -\omega^2 \underline{\varepsilon} - \alpha e^{i\theta\mu} \mathbf{1} \in \underline{\mathbf{L}}^\infty(\Omega)$.

We claim that the form a is coercive on $\mathbf{K}_N(\underline{\varepsilon}; \Omega)$. Indeed:

$$\begin{aligned} |a(\mathbf{v}, \mathbf{v})| &= |(\underline{\mu}^{-1} \mathbf{curl} \mathbf{v} | \mathbf{curl} \mathbf{v})_{0,\Omega} + \alpha e^{i\theta\mu} (\mathbf{v} | \mathbf{v})_{0,\Omega}| \\ &\geq \Re \left[e^{-i\theta\mu} (\underline{\mu}^{-1} \mathbf{curl} \mathbf{v} | \mathbf{curl} \mathbf{v})_{0,\Omega} + \alpha (\mathbf{v} | \mathbf{v})_{0,\Omega} \right] \\ &\geq (\mu^{-1})_- \|\mathbf{curl} \mathbf{v}\|_{0,\Omega}^2 + \alpha \|\mathbf{v}\|_{0,\Omega}^2 \\ &\gtrsim \|\mathbf{v}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}^2. \end{aligned}$$

Besides, $|b(\mathbf{u}, \mathbf{v})| \leq \|\underline{\varepsilon}'\|_{\underline{\mathbf{L}}^\infty(\Omega)} \|\mathbf{u}\|_{0,\Omega} \|\mathbf{v}\|_{0,\Omega} \lesssim \|\mathbf{u}\|_{0,\Omega} \|\mathbf{v}\|_{\mathbf{K}_N(\underline{\varepsilon}; \Omega)}$, so the form b is continuous on $\mathbf{L}^2(\Omega) \times \mathbf{K}_N(\underline{\varepsilon}; \Omega)$.

As a consequence, the embedding of $\mathbf{K}_N(\underline{\varepsilon}; \Omega)$ into $\mathbf{L}^2(\Omega)$ is compact by theorem 2.5. Hence problem (3.6) enters the coercive + compact framework.

Regarding finally the bound, one uses simply the bounds on $\|p\|_{1,\Omega}$ (see lemma 3.2) and on $\|\ell_{D,0}\|_{(\mathbf{H}_0(\mathbf{curl}; \Omega))'}$ (see (3.3)). \square

We are now in a position to solve formulation (3.2) by regrouping the previous results.

Theorem 3.1. *The formulation (3.2) with unknown \mathbf{E}_0 fits the coercive + compact framework:*

- either the problem (3.2) admits a unique solution \mathbf{E}_0 in $\mathbf{H}_0(\mathbf{curl}; \Omega)$, which depends continuously on the data \mathbf{f} and \mathbf{E}_d :

$$\|\mathbf{E}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{E}_d\|_{\mathbf{H}(\mathbf{curl}; \Omega)};$$

- or, the problem (3.2) has solutions if, and only if, \mathbf{f} and \mathbf{E}_d satisfy a finite number of compatibility conditions; in this case, the space of solutions is an affine space of finite dimension. Additionally, the component of the solution which is orthogonal (in the sense of the $\mathbf{H}_0(\mathbf{curl}; \Omega)$ scalar product) to the corresponding linear vector space, depends continuously on the data \mathbf{f} and \mathbf{E}_d .

Moreover, each alternative occurs simultaneously for formulation (3.6) and formulation (3.2).

Assumption. In the rest of the manuscript, we assume the problem (3.6) has a unique solution, such that the problem (3.2) is well-posed.

3.2. Extra-regularity of the solution

The next two subsections aim at determining the extra-regularity of the solution and the solution's curl, depending on the extra-regularity of the data. We make the following extra-regularity assumptions for the next two subsections. We assume that $\partial\Omega$ is of class \mathcal{C}^2 and that $\underline{\boldsymbol{\mu}}, \underline{\boldsymbol{\varepsilon}} \in \mathbf{W}^{1, \infty}(\Omega) \cap \mathbf{C}^1(\Omega)$. Regarding the extra-regularity of \mathbf{f} and \mathbf{E}_d , we assume $\operatorname{div} \mathbf{f} \in H^{\mathbf{s}-1}(\Omega) = (H_0^{1-\mathbf{s}}(\Omega))'$, $\mathbf{E}_d \in \mathbf{H}^{\mathbf{r}}(\Omega)$ and $\mathbf{curl} \mathbf{E}_d \in \mathbf{H}^{\mathbf{r}' }(\Omega)$ for given $\mathbf{s}, \mathbf{r}, \mathbf{r}'$ in $[0, 1] \setminus \{\frac{1}{2}\}$.

Let us recall a result on the continuous splitting of fields of $\mathbf{H}_0(\mathbf{curl}; \Omega)$, cf. Lemma 2.4 in Ref. 16, or Theorem 3.6.7 in Ref. 2:

Theorem 3.2. *Let Ω be a domain. For all \mathbf{u} in $\mathbf{H}_0(\mathbf{curl}; \Omega)$, there exist \mathbf{u}^{reg} in $\mathbf{H}^1(\Omega)$ and ϕ in $H_0^1(\Omega)$, such that*

$$\mathbf{u} = \mathbf{u}^{\text{reg}} + \nabla\phi \text{ in } \Omega, \text{ with } \|\mathbf{u}^{\text{reg}}\|_{1, \Omega} + \|\phi\|_{1, \Omega} \lesssim \|\mathbf{u}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}.$$

With theorem 3.2 at hand, we introduce the splitting of \mathbf{E}_0 :

$$\begin{cases} \mathbf{E}_0 = \mathbf{E}^{\text{reg}} + \nabla\phi_E, \text{ with } \mathbf{E}^{\text{reg}} \in \mathbf{H}^1(\Omega), \phi_E \in H_0^1(\Omega), \text{ and} \\ \|\mathbf{E}^{\text{reg}}\|_{1, \Omega} + \|\phi_E\|_{1, \Omega} \lesssim \|\mathbf{E}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)}. \end{cases} \quad (3.7)$$

Taking $\mathbf{F} = \nabla q$ in (3.2) for any $q \in H_0^1(\Omega)$, it holds that $-\omega^2 (\underline{\boldsymbol{\varepsilon}} \mathbf{E}_0 | \nabla q)_{0, \Omega} = (\mathbf{f} + \omega^2 \underline{\boldsymbol{\varepsilon}} \mathbf{E}_d | \nabla q)_{0, \Omega}$. As a consequence, ϕ_E is governed by the Dirichlet problem

$$\begin{cases} \text{Find } \phi_E \in H_0^1(\Omega) \text{ such that} \\ \omega^2 (\underline{\boldsymbol{\varepsilon}} \nabla \phi_E | \nabla q)_{0, \Omega} = (\operatorname{div} \mathbf{f} + \omega^2 \operatorname{div} \underline{\boldsymbol{\varepsilon}} \mathbf{E}_d + \omega^2 \operatorname{div} \underline{\boldsymbol{\varepsilon}} \mathbf{E}^{\text{reg}} | q)_{0, \Omega}, \forall q \in H_0^1(\Omega). \end{cases} \quad (3.8)$$

Let us recall the fundamental regularity result for solutions of the Dirichlet problem, see Theorem 3.4.5 in Ref. 9:

Theorem 3.3 (Shift theorem). *Let Ω be a bounded domain of boundary $\partial\Omega$, ℓ in $(H_0^1(\Omega))'$, and p governed by*

$$\begin{cases} \text{Find } p \in H_0^1(\Omega) \text{ such that} \\ (\underline{\boldsymbol{\xi}} \nabla p | \nabla q)_{0, \Omega} = \ell(q), \forall q \in H_0^1(\Omega). \end{cases} \quad (3.9)$$

If the tensor coefficient $\underline{\xi}$ fulfills the ellipticity condition, then the problem (3.9) is well-posed; if additionally $\underline{\xi} \in \underline{\mathbf{C}}^1(\Omega)$ and $\partial\Omega$ is of class \mathcal{C}^2 , then, $\forall \sigma \in [1, 2] \setminus \{\frac{3}{2}\}$,

$$\ell \in (H_0^{2-\sigma}(\Omega))' \implies p \in H^\sigma(\Omega); \quad (3.10)$$

additionally,

$$\exists C_\sigma > 0, \forall \ell \in (H_0^{2-\sigma}(\Omega))', \|p\|_{H^\sigma(\Omega)} \leq C_\sigma \|\ell\|_{(H_0^{2-\sigma}(\Omega))'}. \quad (3.11)$$

Consequently, we have the following regularity result for \mathbf{E} :

Theorem 3.4. *Let \mathbf{E} , governed by (3.1), be split as $\mathbf{E} = \mathbf{E}_0 + \mathbf{E}_d$ with $\mathbf{E}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$. If $\partial\Omega$ is of class \mathcal{C}^2 , if $\underline{\varepsilon} \in \underline{\mathbf{W}}^{1,\infty}(\Omega) \cap \underline{\mathbf{C}}^1(\Omega)$, if $\mathbf{f} \in \mathbf{L}^2(\Omega)$ is such that $\operatorname{div} \mathbf{f} \in H^{s-1}(\Omega)$ with \mathbf{s} in $[0, 1] \setminus \{\frac{1}{2}\}$, and if $\mathbf{E}_d \in \mathbf{H}^{\mathbf{r}}(\Omega)$ with \mathbf{r} in $[0, 1] \setminus \{\frac{1}{2}\}$, then*

$$\begin{aligned} \mathbf{E} &\in \mathbf{H}^{\min(\mathbf{s}, \mathbf{r})}(\Omega) \text{ and} \\ \|\mathbf{E}\|_{\min(\mathbf{s}, \mathbf{r}), \Omega} &\lesssim \|\mathbf{f}\|_{0, \Omega} + \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{0, \Omega}. \end{aligned} \quad (3.12)$$

Remark 3.3. No regularity assumption on $\underline{\mu}$ is required here.

Proof. We split \mathbf{E}_0 as in (3.7), and we apply theorem 3.3 to the problem (3.8) governing ϕ_E . Let us introduce the right-hand side of (3.8), $\ell \in (H_0^1(\Omega))'$, and defined by

$$\ell : q \mapsto (\operatorname{div} \mathbf{f} + \omega^2 \operatorname{div} \underline{\varepsilon} \mathbf{E}_d + \omega^2 \operatorname{div} \underline{\varepsilon} \mathbf{E}^{\text{reg}} |q)_{0, \Omega}.$$

Consider each term: one has $\operatorname{div} \mathbf{f} \in H^{s-1}(\Omega) = (H_0^{1-s}(\Omega))'$; as $\underline{\varepsilon} \in \underline{\mathbf{W}}^{1,\infty}(\Omega)$, there holds $\underline{\varepsilon} \mathbf{E}^{\text{reg}} \in \mathbf{H}^1(\Omega)$ and $\operatorname{div} \underline{\varepsilon} \mathbf{E}^{\text{reg}} \in L^2(\Omega)$; similarly, $\underline{\varepsilon} \mathbf{E}_d \in \mathbf{H}^{\mathbf{r}}(\Omega)$, so $\operatorname{div} \underline{\varepsilon} \mathbf{E}_d \in H^{\mathbf{r}-1}(\Omega) = (H_0^{1-\mathbf{r}}(\Omega))'$ as soon as $\mathbf{r} \neq \frac{1}{2}$. It follows that $\ell \in H^{\min(\mathbf{s}, \mathbf{r})-1}(\Omega) = (H_0^{1-\min(\mathbf{s}, \mathbf{r})}(\Omega))'$. In addition, one has the bound

$$\begin{aligned} \|\ell\|_{\min(\mathbf{s}, \mathbf{r})-1, \Omega} &\lesssim \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\operatorname{div} \underline{\varepsilon} \mathbf{E}^{\text{reg}}\|_{0, \Omega} + \|\operatorname{div} \underline{\varepsilon} \mathbf{E}_d\|_{\mathbf{r}-1, \Omega} \\ &\lesssim \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\mathbf{E}^{\text{reg}}\|_{1, \Omega} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega} \\ &\lesssim \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\mathbf{E}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega}, \end{aligned}$$

where we used (3.7) to reach the third line. We conclude by the shift theorem 3.3 that $\phi_E \in H^{\min(\mathbf{s}, \mathbf{r})+1}(\Omega)$, and $\|\phi_E\|_{\min(\mathbf{s}, \mathbf{r})+1, \Omega} \lesssim \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\mathbf{E}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega}$. Hence $\mathbf{E} = \mathbf{E}^{\text{reg}} + \nabla \phi_E + \mathbf{E}_d$ belongs to $\mathbf{H}^{\min(\mathbf{s}, \mathbf{r})}(\Omega)$, with the bound

$$\begin{aligned} \|\mathbf{E}\|_{\min(\mathbf{s}, \mathbf{r}), \Omega} &\lesssim \|\mathbf{E}^{\text{reg}}\|_{1, \Omega} + \|\nabla \phi_E\|_{\min(\mathbf{s}, \mathbf{r}), \Omega} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega} \\ &\lesssim \|\mathbf{E}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\nabla \phi_E\|_{\min(\mathbf{s}, \mathbf{r}), \Omega} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega} \\ &\lesssim \|\mathbf{E}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega}, \\ &\lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{0, \Omega} + \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega}, \end{aligned}$$

where we used successively (3.7), the bound on $\|\phi_E\|_{\min(\mathbf{s}, \mathbf{r})+1, \Omega}$ and finally theorem 3.1 to conclude. \square

Corollary 3.1. *Let the assumptions of theorem 3.4 hold. When $\mathbf{f} \in \mathbf{H}(\operatorname{div}; \Omega)$, there holds $\mathbf{E} \in \mathbf{H}^{\mathbf{r}}(\Omega)$, and $\|\mathbf{E}\|_{\mathbf{r}, \Omega} \lesssim \|\mathbf{f}\|_{\mathbf{H}(\operatorname{div}; \Omega)} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{0, \Omega}$.*

3.3. Extra-regularity of the solution's curl

We proceed similarly with the solution's curl, with the same assumptions as in subsection 3.2. One has a theorem analogous to theorem 3.2, on the continuous splitting of fields of $\mathbf{H}(\mathbf{curl}; \Omega)$, cf. Theorem 3.6.7 in Ref. 2 (see also Lemma 2.4 in Ref. 16 for a similar result):

Theorem 3.5. *Let Ω be a domain of the \mathfrak{A} -type (see Definition A.2). For all \mathbf{u} in $\mathbf{H}(\mathbf{curl}; \Omega)$, there exist \mathbf{u}^{reg} in $\mathbf{H}^1(\Omega)$ and ϕ in $H_{\text{zmv}}^1(\Omega)$, such that $\mathbf{u}^{\text{reg}} \cdot \mathbf{n}_{|\partial\Omega} = 0$ and*

$$\mathbf{u} = \mathbf{u}^{\text{reg}} + \nabla\phi \text{ in } \Omega, \text{ with } \|\mathbf{u}^{\text{reg}}\|_{1,\Omega} + \|\phi\|_{1,\Omega} \lesssim \|\mathbf{u}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}.$$

Let $\mathbf{C} := \underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E} \in \mathbf{L}^2(\Omega)$, one has $\mathbf{curl} \mathbf{C} = \mathbf{f} + \omega^2 \underline{\boldsymbol{\epsilon}} \mathbf{E} \in \mathbf{L}^2(\Omega)$, hence $\mathbf{C} \in \mathbf{H}(\mathbf{curl}; \Omega)$ with the bound

$$\|\mathbf{C}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \lesssim \|\mathbf{E}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{f}\|_{0,\Omega}, \quad (3.13)$$

according to proposition 2.1. Observing that $\mathbf{E} = \mathbf{E}_0 + \mathbf{E}_d$ with $\mathbf{E}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$, for any q in $H_{\text{zmv}}^1(\Omega)$, it holds that $(\underline{\boldsymbol{\mu}} \mathbf{C} | \nabla q)_{0,\Omega} = (\mathbf{curl} \mathbf{E} | \nabla q)_{0,\Omega} = (\mathbf{curl} \mathbf{E}_d | \nabla q)_{0,\Omega}$. Next, we introduce the splitting of \mathbf{C} by theorem 3.5:

$$\begin{cases} \mathbf{C} = \mathbf{C}^{\text{reg}} + \nabla\phi_C, \text{ with } \mathbf{C}^{\text{reg}} \in \mathbf{H}^1(\Omega), \phi_C \in H_{\text{zmv}}^1(\Omega), \mathbf{C}^{\text{reg}} \cdot \mathbf{n}_{|\partial\Omega} = 0 \text{ and} \\ \|\mathbf{C}^{\text{reg}}\|_{1,\Omega} + \|\phi_C\|_{1,\Omega} \lesssim \|\mathbf{C}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}. \end{cases} \quad (3.14)$$

Thus ϕ_C is governed by the Neumann problem

$$\begin{cases} \text{Find } \phi_C \in H_{\text{zmv}}^1(\Omega) \text{ such that} \\ (\underline{\boldsymbol{\mu}} \nabla \phi_C | \nabla q)_{0,\Omega} = (\mathbf{curl} \mathbf{E}_d - \underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}} | \nabla q)_{0,\Omega}, \forall q \in H_{\text{zmv}}^1(\Omega). \end{cases} \quad (3.15)$$

Hence, one may use a regularity result for solutions of the Neumann problem, see Theorem 3.4.5 in Ref. 9, to estimate the regularity of ϕ_C , to infer the regularity of \mathbf{C} and finally the regularity of $\mathbf{curl} \mathbf{E}$.

Theorem 3.6 (Shift theorem). *Let Ω be a bounded domain of boundary $\partial\Omega$, ℓ in $(H_{\text{zmv}}^1(\Omega))'$, and p governed by*

$$\begin{cases} \text{Find } p \in H_{\text{zmv}}^1(\Omega) \text{ such that} \\ (\underline{\boldsymbol{\xi}} \nabla p | \nabla q)_{0,\Omega} = \ell(q), \forall q \in H_{\text{zmv}}^1(\Omega). \end{cases} \quad (3.16)$$

If the tensor coefficient $\underline{\boldsymbol{\xi}}$ fulfills the ellipticity condition, then the problem (3.16) is well-posed; assume in addition that $\underline{\boldsymbol{\xi}} \in \mathbf{C}^1(\Omega)$ and $\partial\Omega$ is of class \mathcal{C}^2 .

(i) *Then, $\forall \sigma \in [1, \frac{3}{2}[$,*

$$\ell \in (H_{\text{zmv}}^{2-\sigma}(\Omega))' \implies p \in H^\sigma(\Omega); \quad (3.17)$$

and,

$$\exists C_\sigma > 0, \forall \ell \in (H_{\text{zmv}}^{2-\sigma}(\Omega))', \|p\|_{H^\sigma(\Omega)} \leq C_\sigma \|\ell\|_{(H_{\text{zmv}}^{2-\sigma}(\Omega))'}. \quad (3.18)$$

(ii) *If there exists $\sigma \in]\frac{3}{2}, 2]$ such that ℓ writes $\ell(q) = (f|q)_{0,\Omega} + \langle g, q \rangle_{H^{1/2}(\partial\Omega)}$ with $f \in L^2(\Omega)$ and $g \in H^{\sigma-3/2}(\partial\Omega)$, then $p \in H^\sigma(\Omega)$. Moreover,*

$$\exists C_\sigma > 0, \forall (f, g) \in L^2(\Omega) \times H^{\sigma-3/2}(\partial\Omega), \|p\|_{H^\sigma(\Omega)} \leq C_\sigma \left(\|f\|_{0,\Omega} + \|g\|_{H^{\sigma-3/2}(\partial\Omega)} \right). \quad (3.19)$$

Remark 3.4. In the case (i), the theorem can be understood in a variational manner, just as in theorem 3.3 for the Dirichlet problem. On the other hand, in the case (ii), the proof relies on local analysis arguments; see⁹ for details.

Applying this result to problem (3.15), one finds the regularity of ϕ_C , then of $\mathbf{curl} \mathbf{E}$.

Theorem 3.7. *Let \mathbf{E} , governed by (3.1), be split as $\mathbf{E} = \mathbf{E}_0 + \mathbf{E}_d$ with $\mathbf{E}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$. If $\partial\Omega$ is of class \mathcal{C}^2 , if $\underline{\boldsymbol{\mu}} \in \underline{\mathbf{W}}^{1,\infty}(\Omega) \cap \underline{\mathbf{C}}^1(\Omega)$, and if $\mathbf{curl} \mathbf{E}_d \in \mathbf{H}^{\mathbf{r}' }(\Omega)$ with \mathbf{r}' in $[0, 1] \setminus \{\frac{1}{2}\}$, then*

$$\begin{aligned} \mathbf{curl} \mathbf{E} &\in \mathbf{H}^{\mathbf{r}' }(\Omega) \text{ and} \\ \|\mathbf{curl} \mathbf{E}\|_{\mathbf{r}', \Omega} &\lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{E}_d\|_{0, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega}. \end{aligned} \quad (3.20)$$

Remark 3.5. No regularity assumption on $\underline{\boldsymbol{\varepsilon}}$ (other than $\underline{\boldsymbol{\varepsilon}} \in \underline{\mathbf{L}}^\infty(\Omega)$) is required here. As Ω is a domain with boundary of class \mathcal{C}^2 , it is automatically of the \mathfrak{A} -type.

Proof. We would like to apply theorem 3.6 to the problem (3.15) governing ϕ_C . Let us introduce the right-hand side of (3.15), $\ell \in (H_{\text{zmv}}^1(\Omega))'$, and defined by

$$\ell : q \mapsto (\mathbf{curl} \mathbf{E}_d - \underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}} |\nabla q)_{0, \Omega}.$$

To determine the regularity of ϕ_C , one wants to determine whether the form ℓ belongs to $(H_{\text{zmv}}^{2-\sigma}(\Omega))'$ for $\sigma \in [1, 2] \setminus \frac{3}{2}$ as large as possible.

If $\mathbf{r}' < \frac{1}{2}$, then $\mathbf{H}^{\mathbf{r}' }(\Omega)$ identifies with $\mathbf{H}_0^{\mathbf{r}' }(\Omega)$, the dual space of $\mathbf{H}^{-\mathbf{r}' }(\Omega)$. Hence the product $(\mathbf{curl} \mathbf{E}_d |\nabla q)_{0, \Omega}$ is meaningful as soon as $q \in H_{\text{zmv}}^{1-\mathbf{r}' }(\Omega)$, because $\nabla q \in \mathbf{H}^{-\mathbf{r}' }(\Omega)$ in this case; the same holds for the term $(\underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}} |\nabla q)_{0, \Omega}$. This means that ℓ belongs to $(H_{\text{zmv}}^{1-\mathbf{r}' }(\Omega))'$, and the shift theorem 3.6 (i), with $\sigma = 1 + \mathbf{r}'$, ensures that $\phi_C \in H^{1+\mathbf{r}' }(\Omega)$, with the bound

$$\|\phi_C\|_{1+\mathbf{r}', \Omega} \lesssim \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega} + \|\underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}}\|_{1, \Omega} \lesssim \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega} + \|\mathbf{C}^{\text{reg}}\|_{1, \Omega}.$$

On the other hand, if $\mathbf{r}' > \frac{1}{2}$, then $\mathbf{H}_0^{\mathbf{r}' }(\Omega)$ does not identify with $\mathbf{H}^{\mathbf{r}' }(\Omega)$, and, as soon as $\mathbf{curl} \mathbf{E}_d \cdot \mathbf{n}|_{\partial\Omega} \neq 0$, the product $(\mathbf{curl} \mathbf{E}_d |\nabla q)_{0, \Omega}$ can be meaningless if one has only $q \in H_{\text{zmv}}^{1-\mathbf{r}' }(\Omega)$. However, since $\mathbf{r}' > \frac{1}{2}$, $\mathbf{curl} \mathbf{E}_d \cdot \mathbf{n}|_{\partial\Omega}$ makes sense in $H^{\mathbf{r}' - 1/2}(\partial\Omega)$, and, as $\underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}} \cdot \mathbf{n}|_{\partial\Omega} = 0$, $\ell(q)$ rewrites by integrations by parts $\ell(q) = (\text{div} \underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}} |q)_{0, \Omega} + \langle \mathbf{curl} \mathbf{E}_d \cdot \mathbf{n}, q \rangle_{H^{1/2}(\partial\Omega)}$. As $\text{div} \underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}} \in L^2(\Omega)$, ℓ satisfies the assumptions of the shift theorem 3.6 (ii) with $\sigma = \mathbf{r}' + 1$, and we conclude that $\phi_C \in H^{1+\mathbf{r}' }(\Omega)$, with the bound

$$\begin{aligned} \|\phi_C\|_{1+\mathbf{r}', \Omega} &\lesssim \|\mathbf{curl} \mathbf{E}_d \cdot \mathbf{n}\|_{H^{\mathbf{r}' - 1/2}(\partial\Omega)} + \|\text{div} \underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}}\|_{0, \Omega} \\ &\lesssim \|\mathbf{curl} \mathbf{E}_d\|_{H^{\mathbf{r}' - 1/2}(\partial\Omega)} + \|\text{div} \underline{\boldsymbol{\mu}} \mathbf{C}^{\text{reg}}\|_{0, \Omega} \lesssim \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega} + \|\mathbf{C}^{\text{reg}}\|_{1, \Omega}. \end{aligned}$$

So, $\phi_C \in H^{1+\mathbf{r}' }(\Omega)$ in all cases (with the same upper bound), so that $\mathbf{C} = \mathbf{C}^{\text{reg}} + \nabla \phi_C \in \mathbf{H}^{\mathbf{r}' }(\Omega)$,

$$\begin{aligned} \|\mathbf{C}\|_{\mathbf{r}', \Omega} &\lesssim \|\mathbf{C}^{\text{reg}}\|_{1, \Omega} + \|\nabla \phi_C\|_{\mathbf{r}', \Omega} \lesssim \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega} + \|\mathbf{C}^{\text{reg}}\|_{1, \Omega} \\ \text{(see (3.14))} &\lesssim \|\mathbf{C}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega}. \end{aligned}$$

As $\mathbf{curl} \mathbf{E} = \underline{\boldsymbol{\mu}} \mathbf{C}$ and $\underline{\boldsymbol{\mu}} \in \underline{\mathbf{W}}^{1,\infty}(\Omega)$, it finally holds that $\mathbf{curl} \mathbf{E} \in \mathbf{H}^{\mathbf{r}' }(\Omega)$, with the bound $\|\mathbf{curl} \mathbf{E}\|_{\mathbf{r}', \Omega} \lesssim \|\mathbf{C}\|_{\mathbf{r}', \Omega}$. Finally, with the help of (3.13) and of theorem 3.1, one concludes that

$$\begin{aligned} \|\mathbf{curl} \mathbf{E}\|_{\mathbf{r}', \Omega} &\lesssim \|\mathbf{C}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega} \\ &\lesssim \|\mathbf{E}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega} \\ &\lesssim \|\mathbf{E}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{E}_d\|_{0, \Omega} + \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega} \\ &\lesssim \|\mathbf{E}_d\|_{0, \Omega} + \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega}. \end{aligned} \quad \square$$

The last theorem sums up the regularity results of this section.

Theorem 3.8. *Let \mathbf{E} , governed by (3.1), be split as $\mathbf{E} = \mathbf{E}_0 + \mathbf{E}_d$ with $\mathbf{E}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$. If $\partial\Omega$ is of class \mathcal{C}^2 , if $\underline{\varepsilon}, \underline{\mu} \in \mathbf{W}^{1,\infty}(\Omega) \cap \mathbf{C}^1(\Omega)$, if $\mathbf{f} \in \mathbf{L}^2(\Omega)$ and is such that $\operatorname{div} \mathbf{f} \in H^{s-1}(\Omega)$ with s in $[0, 1] \setminus \{\frac{1}{2}\}$, and if $\mathbf{E}_d \in \mathbf{H}^s(\Omega)$ and is such that $\mathbf{curl} \mathbf{E}_d \in \mathbf{H}^{s'}(\Omega)$ with \mathbf{r}, \mathbf{r}' in $[0, 1] \setminus \{\frac{1}{2}\}$, then*

$$\begin{cases} \mathbf{E} \in \mathbf{H}^{\min(s, \mathbf{r})}(\Omega), \mathbf{curl} \mathbf{E} \in \mathbf{H}^{s'}(\Omega), \text{ and} \\ \|\mathbf{E}\|_{\min(s, \mathbf{r}, \Omega)} \lesssim \|\mathbf{f}\|_{0, \Omega} + \|\operatorname{div} \mathbf{f}\|_{s-1, \Omega} + \|\mathbf{E}_d\|_{\mathbf{r}, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{0, \Omega}, \\ \|\mathbf{curl} \mathbf{E}\|_{\mathbf{r}', \Omega} \lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{E}_d\|_{0, \Omega} + \|\mathbf{curl} \mathbf{E}_d\|_{\mathbf{r}', \Omega}. \end{cases} \quad (3.21)$$

4. Analysis of the Neumann problem

In this section, the time-harmonic Maxwell's equation is supplemented by a Neumann boundary condition:

$$\begin{cases} \text{Find } \mathbf{E} \in \mathbf{H}(\mathbf{curl}; \Omega) \text{ such that} \\ \mathbf{curl}(\underline{\mu}^{-1} \mathbf{curl} \mathbf{E}) - \omega^2 \underline{\varepsilon} \mathbf{E} = \mathbf{f} \text{ in } \Omega, \\ \underline{\mu}^{-1} \mathbf{curl} \mathbf{E} \times \mathbf{n} = \mathbf{j} \text{ on } \partial\Omega, \end{cases} \quad (4.1)$$

where \mathbf{j} is a boundary data which can be interpreted as a surface current.

We assume that the tensors $\underline{\varepsilon}, \underline{\mu} \in \mathbf{L}^\infty(\Omega)$ are elliptic. We also assume that $\mathbf{f} \in \mathbf{L}^2(\Omega)$ and that \mathbf{j} is the tangential trace of a field \mathbf{B}_d defined on Ω , i.e. $\mathbf{j} = \mathbf{B}_d \times \mathbf{n}|_{\partial\Omega}$, with $\mathbf{B}_d \in \mathbf{H}(\mathbf{curl}; \Omega)$.

4.1. Variational formulation and well-posedness

The equivalent variational formulation of the problem is obtained by the integration by parts formula (1.1). It writes:

$$\begin{cases} \text{Find } \mathbf{E} \in \mathbf{H}(\mathbf{curl}; \Omega) \text{ such that} \\ (\underline{\mu}^{-1} \mathbf{curl} \mathbf{E} | \mathbf{curl} \mathbf{F})_{0, \Omega} - \omega^2 (\underline{\varepsilon} \mathbf{E} | \mathbf{F})_{0, \Omega} = \ell_N(\mathbf{F}), \quad \forall \mathbf{F} \in \mathbf{H}(\mathbf{curl}; \Omega), \end{cases} \quad (4.2)$$

where $\ell_N : \mathbf{F} \mapsto (\mathbf{f} | \mathbf{F})_{0, \Omega} - \gamma \langle \mathbf{j}, \pi^T \mathbf{F} \rangle_\pi$ belongs to $(\mathbf{H}(\mathbf{curl}; \Omega))'$, and

$$\|\ell_N\|_{(\mathbf{H}(\mathbf{curl}; \Omega))'} \lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{j}\|_\gamma \lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{B}_d\|_{\mathbf{H}(\mathbf{curl}; \Omega)}. \quad (4.3)$$

The analysis follows the same reasoning as in the section 3. For this reason, some proofs are just outlined. For the Neumann problem, our analysis relies on a second-kind Helmholtz decomposition (see theorem 2.3).

Lemma 4.1. *The formulation (4.2) can be equivalently recast as: set $\mathbf{E} = \tilde{\mathbf{E}} + \nabla p$, with $p \in H_{\text{zmv}}^1(\Omega)$ and $\tilde{\mathbf{E}} \in \mathbf{K}_T(\underline{\varepsilon}; \Omega)$, respectively governed by*

$$\begin{cases} \text{Find } p \in H_{\text{zmv}}^1(\Omega) \text{ such that} \\ -\omega^2 (\underline{\varepsilon} \nabla p | \nabla q)_{0, \Omega} = \ell_N(\nabla q), \quad \forall q \in H_{\text{zmv}}^1(\Omega), \end{cases} \quad (4.4)$$

and

$$\begin{cases} \text{Find } \tilde{\mathbf{E}} \in \mathbf{K}_T(\underline{\varepsilon}; \Omega) \text{ such that} \\ (\underline{\mu}^{-1} \mathbf{curl} \tilde{\mathbf{E}} | \mathbf{curl} \tilde{\mathbf{F}})_{0, \Omega} - \omega^2 (\underline{\varepsilon} \tilde{\mathbf{E}} | \tilde{\mathbf{F}})_{0, \Omega} = \omega^2 (\underline{\varepsilon} \nabla p | \tilde{\mathbf{F}})_{0, \Omega} + \ell_N(\tilde{\mathbf{F}}), \quad \forall \tilde{\mathbf{F}} \in \mathbf{K}_T(\underline{\varepsilon}; \Omega). \end{cases} \quad (4.5)$$

Proof. Direct. Let us introduce the second-kind Helmholtz decomposition (2.8) of \mathbf{E} : $\mathbf{E} = \tilde{\mathbf{E}} + \nabla p$, with $p \in H_{\text{zmv}}^1(\Omega)$ and $\tilde{\mathbf{E}} \in \mathbf{K}_T(\underline{\varepsilon}; \Omega)$. Taking $\mathbf{F} = \nabla q$ in (4.2) for any $q \in H_{\text{zmv}}^1(\Omega)$ yields

$$-\omega^2 (\underline{\varepsilon} (\tilde{\mathbf{E}} + \nabla p) | \nabla q)_{0, \Omega} = \ell_N(\nabla q).$$

Hence, as $\tilde{\mathbf{E}}$ belongs to $\mathbf{H}_0(\operatorname{div} \underline{\varepsilon}; \Omega)$, p is governed by (4.4).

On the other hand, there holds

$$(\underline{\mu}^{-1} \operatorname{curl} \tilde{\mathbf{E}} | \operatorname{curl} \mathbf{F})_{0,\Omega} - \omega^2 (\underline{\varepsilon} \tilde{\mathbf{E}} | \mathbf{F})_{0,\Omega} = \omega^2 (\underline{\varepsilon} \nabla p | \mathbf{F})_{0,\Omega} + \ell_N(\mathbf{F})$$

for any $\mathbf{F} \in \mathbf{H}(\operatorname{curl}; \Omega)$, hence for any $\tilde{\mathbf{F}} \in \mathbf{K}_T(\underline{\varepsilon}; \Omega)$: $\tilde{\mathbf{E}}$ is governed by (4.5).

Reverse. Summing equations (4.4) and (4.5) and introducing $\mathbf{E} := \tilde{\mathbf{E}} + \nabla p \in \mathbf{H}(\operatorname{curl}; \Omega)$, one gets, after rearrangements:

$$(\underline{\mu}^{-1} \operatorname{curl} \mathbf{E} | \operatorname{curl}(\tilde{\mathbf{F}} + \nabla q))_{0,\Omega} - \omega^2 (\underline{\varepsilon} \mathbf{E} | (\tilde{\mathbf{F}} + \nabla q))_{0,\Omega} = \ell_N(\tilde{\mathbf{F}} + \nabla q).$$

As q and $\tilde{\mathbf{F}}$ span respectively $H_{\operatorname{zmv}}^1(\Omega)$ and $\mathbf{K}_T(\underline{\varepsilon}; \Omega)$, we know that the sum $\tilde{\mathbf{F}} + \nabla q$ spans the whole $\mathbf{H}(\operatorname{curl}; \Omega)$ thanks to (2.8); hence \mathbf{E} is governed by (4.2). \square

Here, $\mathbf{K}_T(\underline{\varepsilon}; \Omega)$ is equipped with the norm $\|\cdot\|_{\mathbf{H}(\operatorname{curl}; \Omega)}$. The rest of the analysis proceeds as for the Dirichlet problem.

Theorem 4.1. *The formulation (4.2) fits the coercive + compact framework:*

- either the problem (4.2) admits a unique solution \mathbf{E} in $\mathbf{H}(\operatorname{curl}; \Omega)$, which depends continuously on the data \mathbf{f} and \mathbf{B}_d :

$$\|\mathbf{E}\|_{\mathbf{H}(\operatorname{curl}; \Omega)} \lesssim \|\mathbf{f}\|_{0,\Omega} + \|\mathbf{B}_d\|_{\mathbf{H}(\operatorname{curl}; \Omega)};$$

- or, the problem (4.2) has solutions if, and only if, \mathbf{f} and \mathbf{B}_d satisfy a finite number of compatibility conditions; in this case, the space of solutions is an affine space of finite dimension. Additionally, the component of the solution which is orthogonal (in the sense of the $\mathbf{H}(\operatorname{curl}; \Omega)$ scalar product) to the corresponding linear vector space, depends continuously on the data \mathbf{f} and \mathbf{B}_d .

Proof. The Neumann problem (4.4) is well-posed according to theorem 2.1. In fact, the form ℓ_N is continuous on $\mathbf{H}(\operatorname{curl}; \Omega)$ and the mapping ∇ is continuous from $H_{\operatorname{zmv}}^1(\Omega)$ to $\mathbf{H}(\operatorname{curl}; \Omega)$, so one has $\ell_N \circ \nabla \in (H_{\operatorname{zmv}}^1(\Omega))'$, with the bound $\|p\|_{1,\Omega} \lesssim \|\mathbf{f}\|_{0,\Omega} + \|\mathbf{B}_d\|_{\mathbf{H}(\operatorname{curl}; \Omega)}$.

Besides, the formulation (4.5) fits the coercive + compact framework. Indeed, as in the proof of lemma 3.3, one can split the left-hand side of (4.5) in two terms. Let $\alpha > 0$, the term $(\underline{\mu}^{-1} \operatorname{curl} \tilde{\mathbf{E}} | \operatorname{curl} \tilde{\mathbf{F}})_{0,\Omega} + \alpha e^{i\theta_\mu} (\tilde{\mathbf{E}} | \tilde{\mathbf{F}})_{0,\Omega}$ is coercive on $\mathbf{H}(\operatorname{curl}; \Omega)$ (therefore on $\mathbf{K}_T(\underline{\varepsilon}; \Omega)$), as $\underline{\mu}^{-1}$ satisfies the ellipticity condition (see proposition 2.1). The remaining term $-\omega^2 (\underline{\varepsilon} \tilde{\mathbf{E}} | \tilde{\mathbf{F}})_{0,\Omega} - \alpha e^{i\theta_\mu} (\tilde{\mathbf{E}} | \tilde{\mathbf{F}})_{0,\Omega}$ is continuous on $\mathbf{L}^2(\Omega) \times \mathbf{K}_T(\underline{\varepsilon}; \Omega)$, and the embedding of $\mathbf{K}_T(\underline{\varepsilon}; \Omega)$ into $\mathbf{L}^2(\Omega)$ is compact by theorem 2.7. Hence formulation (4.5) enters the coercive + compact framework, and so does formulation (4.2). The bound $\|\tilde{\mathbf{E}}\|_{\mathbf{H}(\operatorname{curl}; \Omega)} \lesssim \|\nabla p\|_{0,\Omega} + \|\mathbf{f}\|_{0,\Omega} + \|\mathbf{B}_d\|_{\mathbf{H}(\operatorname{curl}; \Omega)}$ follows from (4.3), and the bound on $\|\mathbf{E}\|_{\mathbf{H}(\operatorname{curl}; \Omega)}$ is a consequence of the triangle inequality. \square

Assumption. In the rest of the manuscript, we assume that the problem (4.2) is well-posed.

4.2. Extra-regularity of the solution's curl

As previously, we are now willing to estimate the regularity of the solution and its curl in the Neumann case, depending on the extra-regularity of the data. We make the following extra-regularity assumptions for the next two subsections. We assume that $\partial\Omega$ is of class \mathcal{C}^2 , that $\underline{\varepsilon}, \underline{\mu} \in \mathbf{W}^{1,\infty}(\Omega)$, and additionally that $\underline{\mu} \in \mathbf{C}^1(\Omega)$. Regarding the extra-regularity of \mathbf{f} and \mathbf{B}_d ,

we assume that $\mathbf{f} \in \mathbf{H}(\mathbf{curl}; \Omega) \cap \mathbf{H}^{\mathbf{s}}(\Omega)$ for a given $\mathbf{s} \in [0, 1]$, and that $\mathbf{B}_d \in \mathbf{H}^{\mathbf{r}'}$ and $\mathbf{curl} \mathbf{B}_d \in \mathbf{H}^{\mathbf{r}}$ for given $\mathbf{r}', \mathbf{r} \in [0, 1] \setminus \{\frac{1}{2}\}$. We begin with the estimation on the curl. The analysis relies on the same arguments as in subsection 3.2.

Let us introduce $\mathbf{B} := \underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E} \in \mathbf{L}^2(\Omega)$. There holds $\mathbf{curl} \mathbf{B} = \mathbf{f} + \omega^2 \underline{\boldsymbol{\varepsilon}} \mathbf{E} \in \mathbf{L}^2(\Omega)$, hence $\mathbf{B} \in \mathbf{H}(\mathbf{curl}; \Omega)$, with the bound $\|\mathbf{B}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \lesssim \|\mathbf{E}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{f}\|_{0, \Omega}$, according to proposition 2.1. Moreover, $\mathbf{B} \times \mathbf{n} = \mathbf{B}_d \times \mathbf{n}$ on $\partial\Omega$. Letting $\mathbf{B}_0 := \mathbf{B} - \mathbf{B}_d \in \mathbf{H}_0(\mathbf{curl}; \Omega)$, we introduce the splitting of \mathbf{B}_0 by theorem 3.2:

$$\begin{cases} \mathbf{B}_0 = \mathbf{B}^{\text{reg}} + \nabla \phi_B, \text{ with } \mathbf{B}^{\text{reg}} \in \mathbf{H}^1(\Omega), \phi_B \in H_0^1(\Omega), \text{ and} \\ \|\mathbf{B}^{\text{reg}}\|_{1, \Omega} + \|\phi_B\|_{1, \Omega} \lesssim \|\mathbf{B}_0\|_{\mathbf{H}(\mathbf{curl}; \Omega)}. \end{cases} \quad (4.6)$$

As $\text{div} \underline{\boldsymbol{\mu}} \mathbf{B} = 0$, ϕ_B is governed by the Dirichlet problem

$$\begin{cases} \text{Find } \phi_B \in H_0^1(\Omega) \text{ such that} \\ (\underline{\boldsymbol{\mu}} \nabla \phi_B | \nabla q)_{0, \Omega} = (\text{div} \underline{\boldsymbol{\mu}} \mathbf{B}_d + \text{div} \underline{\boldsymbol{\mu}} \mathbf{B}^{\text{reg}} | q)_{0, \Omega}, \forall q \in H_0^1(\Omega). \end{cases} \quad (4.7)$$

As in subsection 3.2, one can apply the shift theorem 3.3 to get the regularity estimate on ϕ_B and $\mathbf{curl} \mathbf{E}$.

Theorem 4.2. *Let \mathbf{E} be governed by (4.1), and $\mathbf{B} = \underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E}$ be split as $\mathbf{B} = \mathbf{B}_0 + \mathbf{B}_d$ with $\mathbf{B}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$. If $\partial\Omega$ is of class \mathcal{C}^2 , if $\underline{\boldsymbol{\mu}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega) \cap \underline{\mathbf{C}}^1(\Omega)$, and if $\mathbf{B}_d \in \mathbf{H}^{\mathbf{r}'}$ with $\mathbf{r}' \in [0, 1] \setminus \{\frac{1}{2}\}$, then*

$$\begin{aligned} \mathbf{curl} \mathbf{E} &\in \mathbf{H}^{\mathbf{r}'}(\Omega) \text{ and} \\ \|\mathbf{curl} \mathbf{E}\|_{\mathbf{r}', \Omega} &\lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{curl} \mathbf{B}_d\|_{0, \Omega} + \|\mathbf{B}_d\|_{\mathbf{r}', \Omega}. \end{aligned} \quad (4.8)$$

Remark 4.1. No regularity assumption on $\underline{\boldsymbol{\varepsilon}}$ (other than $\underline{\boldsymbol{\varepsilon}} \in \underline{\mathbf{L}}^\infty(\Omega)$) is required here.

Proof. The proof proceeds as in theorem 3.4. We apply the shift theorem 3.3 to the Dirichlet problem (4.7) governing ϕ_B . Let us introduce the right-hand side of (4.7), $\ell \in (H_0^1(\Omega))'$, and defined by

$$\ell : q \mapsto (\text{div} \underline{\boldsymbol{\mu}} \mathbf{B}_d + \text{div} \underline{\boldsymbol{\mu}} \mathbf{B}^{\text{reg}} | q)_{0, \Omega}.$$

We observe that there holds, as $\underline{\boldsymbol{\mu}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega)$, that $\text{div} \underline{\boldsymbol{\mu}} \mathbf{B}^{\text{reg}} \in L^2(\Omega)$ and $\text{div} \underline{\boldsymbol{\mu}} \mathbf{B}_d \in H^{\mathbf{r}'-1}(\Omega)$ (as soon as $\mathbf{r}' \neq \frac{1}{2}$ for the latter). Then $\ell \in H^{\mathbf{r}'-1}(\Omega) = (H_0^{1-\mathbf{r}'}(\Omega))'$. Hence, by the shift theorem 3.3, we have that $\phi_B \in H^{1+\mathbf{r}'}(\Omega)$, with the bound

$$\|\phi_B\|_{1+\mathbf{r}', \Omega} \lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{curl} \mathbf{B}_d\|_{0, \Omega} + \|\mathbf{B}_d\|_{\mathbf{r}', \Omega}.$$

So, $\mathbf{B} = \mathbf{B}^{\text{reg}} + \nabla \phi_B + \mathbf{B}_d \in \mathbf{H}^{\mathbf{r}'}$ and we conclude that (4.8) holds, because $\underline{\boldsymbol{\mu}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega)$ \square

4.3. Extra-regularity of the solution

To estimate the regularity of the solution itself, we follow the same approach as in subsection 3.3, with the same assumptions as in subsection 4.2. To that end, let us introduce $\mathbf{G} := \mathbf{curl} \mathbf{B} = \mathbf{f} + \omega^2 \underline{\boldsymbol{\varepsilon}} \mathbf{E} \in \mathbf{L}^2(\Omega)$. As $\underline{\boldsymbol{\varepsilon}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega)$ and $\mathbf{f} \in \mathbf{H}(\mathbf{curl}; \Omega)$, one finds that $\mathbf{curl} \mathbf{G} = \mathbf{curl} \mathbf{f} +$

$\omega^2 \mathbf{curl} \underline{\boldsymbol{\varepsilon}} \mathbf{E} \in \mathbf{L}^2(\Omega)$, hence $\mathbf{G} \in \mathbf{H}(\mathbf{curl}; \Omega)$, with the bound $\|\mathbf{G}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \lesssim \|\mathbf{E}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{f}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}$. By theorem 3.5, one can introduce the splitting of \mathbf{G} :

$$\begin{cases} \mathbf{G} = \mathbf{G}^{\text{reg}} + \nabla \phi_G, \text{ with } \mathbf{G}^{\text{reg}} \in \mathbf{H}^1(\Omega), \phi_G \in H_{\text{zmv}}^1(\Omega), \mathbf{G}^{\text{reg}} \cdot \mathbf{n}|_{\partial\Omega} = 0 \text{ and} \\ \|\mathbf{G}^{\text{reg}}\|_{1, \Omega} + \|\phi_G\|_{1, \Omega} \lesssim \|\mathbf{G}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}. \end{cases} \quad (4.9)$$

Therefore, for any $q \in H_{\text{zmv}}^1(\Omega)$, there holds $(\mathbf{G}|\nabla q)_{0, \Omega} = (\mathbf{curl} \mathbf{B}|\nabla q)_{0, \Omega} = (\mathbf{curl} \mathbf{B}_d|\nabla q)_{0, \Omega}$, as the part with $\mathbf{B}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$ vanishes by integration by parts. Then ϕ_G is governed by the Neumann problem

$$\begin{cases} \text{Find } \phi_G \in H_{\text{zmv}}^1(\Omega) \text{ such that} \\ (\nabla \phi_G|\nabla q)_{0, \Omega} = (\mathbf{curl} \mathbf{B}_d - \mathbf{G}^{\text{reg}}|\nabla q)_{0, \Omega}, \forall q \in H_{\text{zmv}}^1(\Omega). \end{cases} \quad (4.10)$$

As in subsection 3.3, the regularity of ϕ_G , and \mathbf{E} , is then given by the shift theorem 3.6.

Theorem 4.3. *Let \mathbf{E} be governed by (4.1), and $\mathbf{B} = \underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E}$ be split as $\mathbf{B} = \mathbf{B}_0 + \mathbf{B}_d$ with $\mathbf{B}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$. If $\partial\Omega$ is of class \mathcal{C}^2 , if $\underline{\boldsymbol{\varepsilon}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega)$, if $\mathbf{f} \in \mathbf{H}(\mathbf{curl}; \Omega) \cap \mathbf{H}^s(\Omega)$ with $s \in [0, 1]$, and if $\mathbf{curl} \mathbf{B}_d \in \mathbf{H}^r(\Omega)$ with $r \in [0, 1] \setminus \{\frac{1}{2}\}$, then*

$$\begin{aligned} \mathbf{E} &\in \mathbf{H}^{\min(r, s)}(\Omega) \text{ and} \\ \|\mathbf{E}\|_{\min(r, s), \Omega} &\lesssim \|\mathbf{curl} \mathbf{f}\|_{0, \Omega} + \|\mathbf{f}\|_{s, \Omega} + \|\mathbf{B}_d\|_{0, \Omega} + \|\mathbf{curl} \mathbf{B}_d\|_{r, \Omega}. \end{aligned} \quad (4.11)$$

Remark 4.2. No regularity assumption on $\underline{\boldsymbol{\mu}}$ is required here.

Proof. The proof is as in theorem 3.7: we want to apply the shift theorem 3.6 to the problem (4.10) governing ϕ_G . Let us introduce the right-hand side of (4.10), $\ell \in (H_{\text{zmv}}^1(\Omega))'$, and defined by

$$\ell : q \mapsto (\mathbf{curl} \mathbf{B}_d - \mathbf{G}^{\text{reg}}|\nabla q)_{0, \Omega}.$$

To determine the regularity of ϕ_G , one wants to determine whether the form ℓ belongs to $(H_{\text{zmv}}^{2-\sigma}(\Omega))'$ for $\sigma \in [1, 2] \setminus \frac{3}{2}$ as large as possible.

If $r < \frac{1}{2}$, then $\mathbf{H}^r(\Omega)$ identifies with $\mathbf{H}_0^r(\Omega)$, the dual space of $\mathbf{H}^{-r}(\Omega)$. Hence the product $(\mathbf{curl} \mathbf{B}_d - \mathbf{G}^{\text{reg}}|\nabla q)$ is meaningful as soon as $q \in H_{\text{zmv}}^{1-r}(\Omega)$, because $\nabla q \in \mathbf{H}^{-r}(\Omega)$ in this case. This means that ℓ belongs to $(H_{\text{zmv}}^{1-r}(\Omega))'$, and the shift theorem 3.6 (i), with $\sigma = 1 + r$, ensures that $\phi_G \in H^{1+r}(\Omega)$, with the bound

$$\|\phi_G\|_{1+r, \Omega} \lesssim \|\mathbf{f}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} + \|\mathbf{curl} \mathbf{B}_d\|_{r, \Omega} + \|\mathbf{B}_d\|_{0, \Omega}.$$

On the other hand, if $r > \frac{1}{2}$, the previous argument is not valid anymore. However, $\mathbf{curl} \mathbf{B}_d \cdot \mathbf{n}|_{\partial\Omega}$ makes sense in $H^{r-1/2}(\partial\Omega)$, and, as $\mathbf{G}^{\text{reg}} \cdot \mathbf{n}|_{\partial\Omega} = 0$, $\ell(q)$ rewrites by integrations by parts $\ell(q) = (\text{div} \mathbf{G}^{\text{reg}}|q)_{0, \Omega} + \langle \mathbf{curl} \mathbf{B}_d \cdot \mathbf{n}, q \rangle_{H^{1/2}(\partial\Omega)}$. As $\text{div} \mathbf{G}^{\text{reg}} \in L^2(\Omega)$, ℓ satisfies the assumptions of the shift theorem 3.6 (ii), with $\sigma = 1 + r$, and we conclude that $\phi_G \in H^{1+r}(\Omega)$, with the same bound as above.

Finally, $\phi_G \in H^{1+r}(\Omega)$ in all cases, and $\mathbf{G} = \mathbf{G}^{\text{reg}} + \nabla \phi_G \in \mathbf{H}^r(\Omega)$. In passing, we note that $\underline{\boldsymbol{\varepsilon}}^{-1} \in \underline{\mathbf{W}}^{1, \infty}(\Omega)$, because $\underline{\boldsymbol{\varepsilon}}^{-1} \in \underline{\mathbf{L}}^\infty(\Omega)$ (Proposition 2.1) and $\underline{\boldsymbol{\varepsilon}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega)$. Recalling that $\mathbf{E} = \omega^{-2} \underline{\boldsymbol{\varepsilon}}^{-1} (\mathbf{G} - \mathbf{f})$, with $\mathbf{f} \in \mathbf{H}^s(\Omega)$, we conclude that (4.11) holds. \square

To conclude, we sum up the regularity results of this section.

Theorem 4.4. *Let \mathbf{E} be governed by problem (4.1), and $\mathbf{B} = \underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{E}$ be split as $\mathbf{B} = \mathbf{B}_0 + \mathbf{B}_d$ with $\mathbf{B}_0 \in \mathbf{H}_0(\mathbf{curl}; \Omega)$. If $\partial\Omega$ is of class \mathcal{C}^2 , if $\underline{\boldsymbol{\mu}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega) \cap \underline{\mathbf{C}}^1(\Omega)$, if $\underline{\boldsymbol{\varepsilon}} \in \underline{\mathbf{W}}^{1, \infty}(\Omega)$,*

if $\mathbf{f} \in \mathbf{H}(\mathbf{curl}; \Omega) \cap \mathbf{H}^{\mathbf{s}}(\Omega)$ with \mathbf{s} in $[0, 1]$, and if $\mathbf{B}_d \in \mathbf{H}^{\mathbf{x}'}(\Omega)$ and is such that $\mathbf{curl} \mathbf{B}_d \in \mathbf{H}^{\mathbf{x}}(\Omega)$ with \mathbf{x}', \mathbf{x} in $[0, 1] \setminus \frac{1}{2}$, then

$$\begin{cases} \mathbf{E} \in \mathbf{H}^{\min(\mathbf{x}, \mathbf{s})}(\Omega), & \mathbf{curl} \mathbf{E} \in \mathbf{H}^{\mathbf{x}'}(\Omega), \text{ and} \\ \|\mathbf{E}\|_{\min(\mathbf{x}, \mathbf{s}), \Omega} \lesssim \|\mathbf{f}\|_{\mathbf{s}, \Omega} + \|\mathbf{curl} \mathbf{f}\|_{0, \Omega} + \|\mathbf{B}_d\|_{0, \Omega} + \|\mathbf{curl} \mathbf{B}_d\|_{\mathbf{x}, \Omega}, \\ \|\mathbf{curl} \mathbf{E}\|_{\mathbf{x}', \Omega} \lesssim \|\mathbf{f}\|_{0, \Omega} + \|\mathbf{B}_d\|_{\mathbf{x}', \Omega} + \|\mathbf{curl} \mathbf{B}_d\|_{0, \Omega}. \end{cases} \quad (4.12)$$

5. $\mathbf{H}(\mathbf{curl})$ -conforming finite element discretization

Edge finite element methods are natural candidates for the numerical solution of electromagnetic problems. Since these methods lead to $\mathbf{H}(\mathbf{curl})$ -conforming approximations, some features of the numerical solutions can be rather easily studied by leveraging the results obtained for the exact problems. While the comprehensive numerical analysis of the approximate problems is out of the scope of this paper, this section aims at giving a few numerical results for the considered problems. After introducing a standard edge finite element discretization and basic results, we derive an *a priori* error estimate, which is obtained by using the regularity estimates. Elementary numerical results are then proposed to illustrate the expected convergence rate of the method.

5.1. Discretization and a priori error estimate

We consider a shape regular family of meshes $(\mathcal{T}_h)_h$ for the domain Ω . For the sake of simplicity, we assume that the domain Ω is a Lipschitz polyhedron. Each mesh \mathcal{T}_h is made up of closed non-overlapping tetrahedra, generically denoted by K , and is indexed by $h := \max_K h_K$, where h_K is the diameter of K . Denoting by ρ_K the diameter of the largest ball inscribed in K , we assume that there exists a shape regularity parameter $\varsigma > 0$ such that for all h , for all $K \in \mathcal{T}_h$, it holds $h_K \leq \varsigma \rho_K$.

Finite dimensional subspaces $(\mathbf{V}_h)_h$ of $\mathbf{H}(\mathbf{curl}; \Omega)$ are defined by using the so-called Nédélec's first family of edge finite elements. Elements of degree 1 are considered. One has

$$\mathbf{V}_h := \{\mathbf{v}_h \in \mathbf{H}(\mathbf{curl}; \Omega), \mathbf{v}_h|_K \in \mathcal{R}_1(K), \forall K \in \mathcal{T}_h\},$$

where $\mathcal{R}_1(K)$ is the six-dimensional vector space of polynomials on K

$$\mathcal{R}_1(K) := \{\mathbf{v} \in \mathbf{P}_1(K) : \mathbf{v}(\mathbf{x}) = \mathbf{a} + \mathbf{b} \times \mathbf{x}, \mathbf{a}, \mathbf{b} \in \mathbb{R}^3\}.$$

The subspaces verify the approximability property (see *e.g.* Lemma 7.10 in Ref. 17)

$$\lim_{h \rightarrow 0} \left(\inf_{\mathbf{v}_h \in \mathbf{V}_h} \|\mathbf{v} - \mathbf{v}_h\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \right) = 0, \forall \mathbf{v} \in \mathbf{H}(\mathbf{curl}; \Omega). \quad (5.1)$$

We also introduce the closed subspaces $(\mathbf{V}_h^0)_h$ with $\mathbf{V}_h^0 := \mathbf{V}_h \cap \mathbf{H}_0(\mathbf{curl}; \Omega)$, which also verify the approximability property in $\mathbf{H}_0(\mathbf{curl}; \Omega)$.

Using the standard Galerkin approach, the variational formulation of the approximate problem is obtained by seeking the solution in \mathbf{V}_h with test functions in \mathbf{V}_h^0 or \mathbf{V}_h for the Dirichlet and Neumann cases, respectively. Therefore, the discrete Dirichlet problem reads

$$\begin{cases} \text{Find } \mathbf{E}_h \in \mathbf{V}_h \text{ such that} \\ a(\mathbf{E}_h, \mathbf{F}_h) = \ell_D(\mathbf{F}_h), \forall \mathbf{F}_h \in \mathbf{V}_h^0, \\ \mathbf{E}_h \times \mathbf{n} = \mathbf{g}_h \text{ on } \partial\Omega \end{cases} \quad (5.2)$$

and the discrete Neumann problem reads

$$\begin{cases} \text{Find } \mathbf{E}_h \in \mathbf{V}_h \text{ such that} \\ a(\mathbf{E}_h, \mathbf{F}_h) = \ell_N(\mathbf{F}_h), \forall \mathbf{F}_h \in \mathbf{V}_h, \end{cases} \quad (5.3)$$

with the sesquilinear form

$$a : (\mathbf{u}, \mathbf{v}) \mapsto (\underline{\boldsymbol{\mu}}^{-1} \mathbf{curl} \mathbf{u} | \mathbf{curl} \mathbf{v})_{0,\Omega} - \omega^2 (\underline{\boldsymbol{\varepsilon}} \mathbf{u} | \mathbf{v})_{0,\Omega} \quad (5.4)$$

defined on $\mathbf{H}(\mathbf{curl}; \Omega)$. The linear forms ℓ_D and ℓ_N are defined in sections 3.1 and 4.1, respectively. For simplicity, in the remaining, we assume that the integrals are computed exactly.

If the form a is coercive, an *a priori* error estimate for the numerical solution is obtained thanks to Céa's lemma. The lemma reads as follows.

Theorem 5.1. *If the form a is coercive, it holds that*

$$\exists C > 0, \forall h, \|\mathbf{E} - \mathbf{E}_h\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \leq C \inf_{\mathbf{v}_h \in \mathbf{V}_h} \|\mathbf{E} - \mathbf{v}_h\|_{\mathbf{H}(\mathbf{curl}; \Omega)}. \quad (5.5)$$

The term appearing in the right-hand side of (5.5) is the well-known projection error. Combining this result with the approximability property (*i.e.* equation (5.1)) immediately gives

$$\lim_{h \rightarrow 0} \|\mathbf{E} - \mathbf{E}_h\|_{\mathbf{H}(\mathbf{curl}; \Omega)} = 0,$$

which expresses that the error goes to 0 as h goes to 0.

To obtain a sharper error estimate, we recall an interpolation error result on the finite element spaces. Let π_h denote the classical interpolation operator from $\mathbf{H}_0(\mathbf{curl}; \Omega)$ onto \mathbf{V}_h^0 , resp. from $\mathbf{H}(\mathbf{curl}; \Omega)$ onto \mathbf{V}_h . One has the the following interpolation error estimate³.

Theorem 5.2. *Let $\mathfrak{t} \in (1/2, 1]$ and $\mathfrak{t}' \in (0, 1]$. For all $\mathbf{v} \in \{\mathbf{v} \in \mathbf{H}^{\mathfrak{t}}(\Omega), \mathbf{curl} \mathbf{v} \in \mathbf{H}^{\mathfrak{t}'}(\Omega)\}$, it holds that*

$$\|\mathbf{v} - \pi_h \mathbf{v}\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \lesssim h^{\min(\mathfrak{t}, \mathfrak{t}')} (\|\mathbf{v}\|_{\mathfrak{t}, \Omega} + \|\mathbf{curl} \mathbf{v}\|_{\mathfrak{t}', \Omega}). \quad (5.6)$$

In this result, $\mathfrak{t} > 1/2$ is assumed for simplicity. A similar result can be obtained for $\mathfrak{t} \in (0, 1]$ with the help of the combined interpolation operator (see Section 4.2 in Ref. 6), but this result is more involved. Indeed, the norm of the gradient part of the decomposition of \mathbf{v} (given in theorems 3.2 or 3.5) then appears in the right-hand side of (5.6), in addition to both terms already there. Nevertheless, since the gradient part is bounded by the norm of the data (see again theorems 3.2 or 3.5), the same conclusion stands in this general case.

In order to derive a sharper *a priori* error estimate for both problems, we would like to combine the interpolation error estimate with the regularity results, for which we have assumed a boundary of class \mathcal{C}^2 . To that aim, we make the supplementary assumption that the domain Ω is convex, and we build a family of convex polyhedral meshes $(\Omega_h)_h$ to approximate Ω . We consider the framework introduced by Dello Russo and Alonso¹⁰. More precisely, Lemma 8.1 in Ref. 10 is a *generalised* Céa's lemma, with the projection error and additional terms in the right-hand side, the latter coming from the discrepancies between the exact problem, set in Ω , and the discrete problems, set in Ω_h ; and Lemma 8.4 in Ref. 10 gives the final *a priori* error estimate.

First, observe that one can replace the field \mathbf{v} with \mathbf{E} in equation (5.6). Using theorems 3.8 and 4.4, the norms $\|\mathbf{E}\|_{\mathfrak{t}, \Omega}$ and $\|\mathbf{curl} \mathbf{E}\|_{\mathfrak{t}', \Omega}$ are bounded by the norms on the data, and the

exponents become $\mathbf{t} = \min(\mathbf{s}, \mathbf{r})$ and $\mathbf{t}' = \mathbf{r}'$, where $\mathbf{s}, \mathbf{r}, \mathbf{r}'$ are the extra-regularity exponents for the data. We then have

$$\|\mathbf{E} - \pi_h \mathbf{E}\|_{\mathbf{H}(\mathbf{curl}; \Omega_h)} \lesssim h^{\min(\mathbf{s}, \mathbf{r}, \mathbf{r}')}, \quad (5.7)$$

where the bounds on the exponents are defined in theorems 3.8 and 4.4 for the Dirichlet and Neumann cases, respectively. Then, we combine the interpolation error estimate (5.7) with the generalised Céa's lemma. As in Section 8 in Ref. 10, one can check that all additional terms yield errors in the order of $O(h)$. Since the interpolation error is always greater or equal to the projection error, we finally have the sharper *a priori* error estimate

$$\|\mathbf{E} - \mathbf{E}_h\|_{\mathbf{H}(\mathbf{curl}; \Omega)} \lesssim h^{\min(\mathbf{s}, \mathbf{r}, \mathbf{r}')}, \quad (5.8)$$

where the exponents depend only on the regularity of the data.

5.2. Numerical results with a manufactured benchmark

To illustrate the expected convergence rate with a numerical case, we consider a simple benchmark with a manufactured solution. Let a spherical domain of unit radius centred at the origin, $\Omega = \{\mathbf{x} \in \mathbb{R}^3, \|\mathbf{x}\| < 1\}$, the angular frequency $\omega = 1$ and the material tensors

$$\underline{\boldsymbol{\mu}} = \text{diag}(1, 1, 1), \quad \underline{\boldsymbol{\varepsilon}} = \text{diag}(1 + 10^{-1}i, 1 + 10^{-1}i, -2 + 10^{-1}i), \quad (5.9)$$

which fulfill assumption (2.1). According to Lemma 2.3 in Ref. 18, the sesquilinear form a is coercive for these material tensors. A non-homogeneous Dirichlet or Neumann boundary condition is prescribed on the boundary of the domain. We consider the manufactured solution

$$\mathbf{E}_{\text{ref}} = [-1, 1, 1]^\top \exp(i\pi \mathbf{k} \cdot \mathbf{x}), \quad \text{with } \mathbf{k} = \frac{1}{\sqrt{14}} [3, 2, 1]^\top. \quad (5.10)$$

The volume source term is chosen accordingly, *i.e.* $\mathbf{f} = \mathbf{curl} \mathbf{curl} \mathbf{E}_{\text{ref}} - \omega^2 \underline{\boldsymbol{\varepsilon}} \mathbf{E}_{\text{ref}}$, as well as the right-hand-side term of the boundary conditions.

Numerical simulations are performed with FreeFem¹⁵ using six unstructured meshes made of tetrahedra and first-degree edge finite elements. Because the boundary of the meshes (which are polyhedral) does not exactly match the curved border of the spherical domain, the boundary data used in the numerical simulation are evaluated on the sphere and then projected on the surface mesh. It has been proven that this geometric approximation introduces a geometric error of the order $\mathcal{O}(h)^{10}$.

The relative numerical error in $\mathbf{H}(\mathbf{curl})$ -norm is plotted as a function of the mesh size h on Fig. 1 for both Dirichlet and Neumann cases. As a reference, the relative error corresponding to the projection of the reference solution on the discrete solution space, which corresponds to the best approximation error according to Céa's lemma, is plotted as well. As the solution \mathbf{E}_{ref} is smooth, it belongs to $\mathbf{H}^1(\Omega)$ as well as its curl, and one has $\mathbf{t} = \mathbf{t}' = 1$ in (5.8). Therefore, one expects the error to evolve linearly with the mesh size h . The results reported on Fig. 1 show that the convergence behaves effectively like $O(h)$ for both problems.

6. Conclusion and extensions

We have addressed the mathematical analysis of time-harmonic electromagnetic boundary value problems in complex anisotropic material tensors, which fulfill a general ellipticity condition. After having developed a functional framework suited to these complex material tensors (*i.e.* functional spaces, Helmholtz decompositions, Weber inequalities and compact embedding results),




Fig. 1: Relative error in $\mathbf{H}(\mathbf{curl})$ -norm obtained with six meshes for the manufactured benchmark with Dirichlet and Neumann boundary conditions. The relative projection error in $\mathbf{H}(\mathbf{curl})$ -norm is plotted in both cases. The dashed lines correspond to the linear scale $O(h)$.

we have analysed the well-posedness of the Dirichlet and Neumann problems (see Theorems 3.1 and 4.1, respectively), as well as the regularity of the solution (*i.e.* the electric field) and the solution's curl. The regularity results are summarised in Theorems 3.8 and 4.4. A preliminary numerical analysis with a $\mathbf{H}(\mathbf{curl})$ -conforming finite element discretization has been proposed.

Among possible extensions, one could consider a domain Ω with a non-smooth boundary (*e.g.* polyhedral), possibly non-convex. The shift theorems then have to be revisited. To our knowledge, results are available for piecewise smooth, hermitian and elliptic material tensors (see *e.g.* ^{8,14,4,7} for shift theorems in settings with a non-smooth boundary and hermitian tensors). The theory developed in this work could be extended thanks to these results. One could also consider settings with mixed boundary conditions, *i.e.* when a Dirichlet condition is prescribed on one part of the boundary, and a Neumann condition is prescribed on the rest of the boundary. To carry out the theory and fit the problem within the coercive+compact framework, we refer to ¹². For shift theorems, we refer to the works of Disser *et al.*¹¹. Finally, one could consider variational formulations with the magnetic field as the unknown, which poses no extra difficulty.

Appendix A. Additional definitions

Definition A.1. From a topological point of view, a domain Ω verifies the hypothesis $(\mathbf{Top})_I$ if one of the following conditions holds:

- (1) For all curl-free vector field $\mathbf{v} \in \mathbf{C}^1(\Omega)$, there exists $p \in C^0(\Omega)$ such that $\mathbf{v} = \nabla p$ in Ω .
- (2) There exist $I > 0$ non-intersecting, piecewise plane manifolds, $(\Sigma_j)_{j=1,\dots,I}$, with boundaries $\partial\Sigma_i \subset \partial\Omega$, such that, if we let $\dot{\Omega} = \Omega \setminus \bigcup_{i=1}^I \Sigma_i$, for all curl-free vector field \mathbf{v} , there exists $p \in C^0(\dot{\Omega})$ such that $\mathbf{v} = \nabla p$ in $\dot{\Omega}$.

If the first condition holds, Ω is topologically trivial, and we set $I = 0$ and $\dot{\Omega} = \Omega$. If the second condition holds, Ω is topologically non-trivial. See ¹³ for further details.

The set $\hat{\Omega}$ has pseudo-Lipschitz boundary in the sense of ¹. The continuation operator from $L^2(\hat{\Omega})$ to $L^2(\Omega)$ is denoted by $\tilde{\cdot}$, whereas the jump across Σ_i is denoted by $[\cdot]_{\Sigma_i}$, for $i = 1, \dots, I$. The definition of the jump depends on the (fixed) orientation of the normal vector field to Σ_i . For all i , we let $\langle \cdot, \cdot \rangle_{\Sigma_i}$ denote the duality pairing between $H^{1/2}(\Sigma_i)$ and $(H^{1/2}(\Sigma_i))'$. One has the integration by parts formula ¹:

$$(\mathbf{v}|\nabla\hat{q})_{0,\hat{\Omega}} + (\operatorname{div}\mathbf{v}|\hat{q})_{0,\hat{\Omega}} = \sum_{1 \leq j \leq I} \langle \mathbf{v} \cdot \mathbf{n}, [\hat{q}]_{\Sigma_j} \rangle_{\Sigma_j}, \quad \forall \mathbf{v} \in \mathbf{H}_0(\operatorname{div}; \Omega), \quad \forall \hat{q} \in H^1(\hat{\Omega}). \quad (\text{A.1})$$

For the sake of completeness, we recall Definition 3.6.3 of Ref. 2.

Definition A.2. A domain Ω is said *of the \mathfrak{A} -type* if, for any $\mathbf{x} \in \partial\Omega$, there exists a neighbourhood \mathcal{V} of \mathbf{x} in \mathbb{R}^3 , and a C^2 diffeomorphism that transforms $\Omega \cap \mathcal{V}$ into one of the following types, where (x_1, x_2, x_3) denote the cartesian coordinates and $(\rho, \omega) \in \mathbb{R} \times \mathbb{S}^2$ the spherical coordinates in \mathbb{R}^3 :

- (1) $[x_1 > 0]$, *i.e.* \mathbf{x} is a regular point;
- (2) $[x_1 > 0, x_2 > 0]$, *i.e.* \mathbf{x} is a point on a salient (outward) edge;
- (3) $\mathbb{R}^3 \setminus [x_1 \geq 0, x_2 \geq 0]$, *i.e.* \mathbf{x} is a point on a reentrant (inward) edge;
- (4) $[\rho > 0, \omega \in \tilde{\Omega}]$, where $\tilde{\Omega} \subset \mathbb{S}^2$ is a topologically trivial domain. In particular, if $\partial\tilde{\Omega}$ is smooth, \mathbf{x} is a conical vertex; if $\partial\tilde{\Omega}$ is made of arcs of great circles, \mathbf{x} is a polyhedral vertex.

Acknowledgement

The authors thank DGA (*Direction Générale de l'Armement*) for the doctoral scholarship of the first author.

References

1. C. Amrouche, C. Bernardi, M. Dauge, and V. Girault. Vector potentials in three-dimensional non-smooth domains. *Math. Meth. Appl. Sci.*, 21:823–864, 1998.
2. F. Assous, P. Ciarlet Jr., and S. Labrunie. *Mathematical foundations of computational electromagnetism*, volume 198 of *Applied Mathematical Sciences*. Springer, 2018.
3. A. Bermúdez, R. Rodríguez, and P. Salgado. Numerical treatment of realistic boundary conditions for the eddy current problem in an electrode via Lagrange multipliers. *Mathematics of computation*, 74(249):123–151, 2005.
4. A. Bonito, J.-L. Guermond, and F. Luddens. Regularity of the Maxwell equations in heterogeneous media and Lipschitz domains. *J. Math. Anal. Appl.*, 408:498–512, 2013.
5. A. Buffa, M. Costabel, and D. Sheen. On traces for $\mathbf{H}(\operatorname{curl}; \Omega)$ in Lipschitz domains. *J. Math. Anal. Appl.*, 276:845–867, 2002.
6. P. Ciarlet, Jr. On the approximation of electromagnetic fields by edge finite elements. Part 1: Sharp interpolation results for low-regularity fields. *Computers Math. Applic.*, 71:85–104, 2016.
7. P. Ciarlet Jr. On the approximation of electromagnetic fields by edge finite elements. Part 3: Sensitivity to coefficients. *SIAM Journal on Mathematical Analysis*, 2020.
8. M. Costabel, M. Dauge, and S. Nicaise. Singularities of Maxwell interface problems. *Math. Mod. Num. Anal.*, 33:627–649, 1999.
9. M. Costabel, M. Dauge, and S. Nicaise. Corner singularities and analytic regularity for linear elliptic systems. Part I: smooth domains. 2010.
10. A. Dello Russo and A. Alonso. Finite element approximation of Maxwell eigenproblems on curved Lipschitz polyhedral domains. *Applied numerical mathematics*, 59(8):1796–1822, 2009.
11. K. Disser, H.-C. Kaiser, and J. Rehberg. Optimal Sobolev regularity for linear second-order divergence elliptic operators occurring in real-world problems. *SIAM J. Math. Anal.*, 47:1719–1746, 2015.

12. P. Fernandes and G. Gilardi. Magnetostatic and electrostatic problems in inhomogeneous anisotropic media with irregular boundary and mixed boundary conditions. *Math. Models Meth. App. Sci.*, 7:957–991, 1997.
13. P.W. Gross and P.R. Kotiuga. *Electromagnetic theory and computation: a topological approach*. MSRI Publications Series, **48**. Cambridge University Press, 2004.
14. R. Haller-Dintelmann, H.-C. Kaiser, and J. Rehberg. Direct computation of elliptic singularities across anisotropic, multi-material edges. *J. Math. Sci.*, 172:589–622, 2011.
15. F. Hecht. New development in FreeFem++. *J. Numer. Math.*, 20:251–265, 2012.
16. R. Hiptmair. Finite elements in computational electromagnetics. *Acta Numerica*, pages 237–339, 2002.
17. P. Monk. *Finite element methods for Maxwell's equations*. Oxford University Press, 2003.
18. E. Sébelin, Y. Peysson, X. Litaudon, D. Moreau, J.C. Miellou, and O. Lafitte. Uniqueness and existence result around Lax-Milgram lemma: application to electromagnetic waves propagation in tokamak plasmas. Technical report, Association Euratom-CEA, 1997. online: http://www.iaea.org/inis/collection/NCLCollectionStore/_Public/30/017/30017036.pdf.