



Spermatozoa motility in bivalves: Signaling, flagellar beating behavior, and energetics

Myrina Boulais, Marina Demoy-Schneider, Sayyed Mohammad Hadi Alavi,
Jacky Cosson

► To cite this version:

Myrina Boulais, Marina Demoy-Schneider, Sayyed Mohammad Hadi Alavi, Jacky Cosson. Spermatozoa motility in bivalves: Signaling, flagellar beating behavior, and energetics. *Theriogenology*, 2019, 136, pp.15-27. 10.1016/j.theriogenology.2019.06.025 . hal-02749005

HAL Id: hal-02749005

<https://hal.science/hal-02749005>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Revised

Spermatozoa motility in bivalves: signaling, flagellar beating behavior, and energetics

Myrina Boulais ^{a,*}, Marina Demoy-Schneider ^b, Sayyed Mohammad Hadi Alavi ^c, Jacky Cosson ^d

^a University of Brest, CNRS, IRD, Ifremer, LEMAR, rue Dumont d'Urville, F-29280 Plouzané, France

^b University of French Polynesia, UMR 241 EIO, BP 6570, 98702 Faa'a Aéroport, Tahiti, French Polynesia

^c School of Biology, College of Science, University of Tehran, P.O. Box 14155-6455, Tehran, Iran

^d South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Faculty of Fisheries and Protection of Waters, University of South Bohemia in České Budějovice, Vodňany 389 25, Czech Republic

Short title: Spermatozoa motility in bivalves

* All correspondence to:

M. Boulais

Tel.: 02 98 49 86 70, Fax: 02 98 49 86 45

E-mail: myrina.boulais@gmail.com or myrina.boulais@univ-brest.fr

24	Abstract
25	1. Introduction
26	2. An overview of sperm biology in the most studied bivalve species
27	2.1. Pacific oyster, <i>Crassostrea gigas</i>
28	2.2. Black-lip pearl oyster, <i>Pinctada margaritifera</i>
29	2.3. European flat oyster, <i>Ostrea edulis</i>
30	2.4. Manila clam, <i>Ruditapes philippinarum</i>
31	2.5. Great scallop, <i>Pecten maximus</i> , and Japanese scallop, <i>Patinopecten yessoensis</i>
32	2.6. Seminal plasma characteristics
33	3. Spermatozoa motility in bivalves
34	3.1. Spermatozoa motility in seawater
35	3.2. Spermatozoa hyper-motility in bivalves
36	3.3. Serotonin-induced spermatozoa hyper-motility via 5-HT receptor(s)
37	3.4. Spermatozoa motility signaling in bivalves
38	3.4.1. Osmolality-independent activation of spermatozoa motility
39	3.4.2. pH regulation of spermatozoa motility initiation
40	3.4.3. K ⁺ regulation of spermatozoa motility initiation
41	3.4.4. Na ⁺ regulation of spermatozoa motility initiation
42	3.4.5. Ca ²⁺ regulation of spermatozoa motility initiation
43	3.4.6. Cyclic nucleotide regulation of spermatozoa motility initiation
44	3.4.7. Signaling pathway controlling spermatozoa motility initiation in bivalves
45	4. Flagellar beating in spermatozoa of bivalves
46	5. Energy metabolism in spermatozoa of bivalves
47	5.1. Mitochondrial respiration
48	5.2. Intracellular ATP concentration
49	5.3. Metabolic pathways synthesizing ATP
50	6. Conclusion and future researches
51	Acknowledgements
52	Conflict of interest
53	References
54	
55	Abstract

Though bivalve mollusks are keystone species and major species groups in aquaculture production worldwide, gamete biology is still largely unknown. This review aims to provide a synthesis of current knowledge in the field of sperm biology, including spermatozoa motility, flagellar beating, and energy metabolism; and to illustrate cellular signaling controlling spermatozoa motility initiation in bivalves. Serotonin (5-HT) induces hypermotility in spermatozoa via a 5-HT receptor, suggesting a serotonergic system in the male reproductive tract that might regulate sperm physiology. Acidic pH and high concentration of K^+ are inhibitory factors of spermatozoa motility in the testis. Motility is initiated at spawning by a Na^+ -dependent alkalization of intracellular pH mediated by a Na^+/H^+ exchanger. Increase of 5-HT in the testis and decrease of extracellular K^+ when sperm is released in seawater induce hyperpolarization of spermatozoa membrane potential mediated by K^+ efflux and associated with an increase in intracellular Ca^{2+} via opening of voltage-dependent Ca^{2+} channels under alkaline conditions. These events activate dynein ATPases and Ca^{2+} /calmodulin-dependent proteins resulting in flagellar beating. It may be possible that 5-HT is also involved in intracellular cAMP rise controlling cAMP-dependent protein kinase phosphorylation in the flagellum. Once motility is triggered, flagellum beats in asymmetric wave pattern leading to circular trajectories of spermatozoa. Three different flagellar wave characteristics are reported, including “full”, “twitching”, and “declining” propagation of wave, which are described and illustrated in the present review. Mitochondrial respiration, ATP content, and metabolic pathways producing ATP in bivalve spermatozoa are discussed. Energy metabolism of Pacific oyster spermatozoa differs from previously studied marine species since oxidative phosphorylation synthesizes a stable level of ATP throughout 24-h motility period and the end of movement is not explained by a low intracellular ATP content, revealing different strategy to improve oocyte fertilization success. Finally, our review

highlights physiological mechanisms that require further researches and points out some advantages of bivalve spermatozoa to extend knowledge on mechanisms of motility.

Keywords: Adenosine triphosphate (ATP); Ion signaling; pH; Serotonin (5-HT); Spermatozoa motility; Spermatozoa velocity

1. Introduction

Human consumption of seafood has been rising during the last half century supported by an increase from 39.2 million tons in 1961 to 170.9 million tons in 2016 in the global production of fish, crustaceans, mollusks, and other aquatic organisms [1]. Recently, mollusks have become a major species group in aquaculture and their global production worldwide increased from 3.6 to 17.1 million tons between 1990 and 2016 [1]. Control of artificial seed production is important to continue to grow aquaculture production (e.g. controlled crosses, selective breeding). However, reproductive biology is still largely unknown in mollusks, which are the second largest animal phylum with almost 100 000 species, among which 15 000 species of bivalves [2].

In bivalves, spermatogenesis is seasonal and controlled by environmental factors including temperature, photoperiod, and salinity [3-7]. Development of germ cells is controlled by hormones of the central nervous system and of the testis [8-10]. During spermatogenesis, central nervous system of bivalves synthesizes monoamines including serotonin (5-HT) [11]. In 1957, Welsh [12] identified 5-HT neurons in the nervous system of bivalves. Former studies suggested that 5-HT acts as a neurotransmitter in the nervous system of mollusks [13-15]. Our current knowledge shows that fibers of 5-HT producing neurons innervate the gonad [11]. Matsutani and Nomura [16] reported that 5-HT injection induces

spermatozoa and oocyte release in various bivalve species. Further studies have shown that changes in 5-HT content in the nervous system is involved in bivalve spawning through a receptor mechanism [11,17]. In addition to 5-HT, spermatogenesis is also controlled by neuropeptides and sex steroids [8-10]. A GnRH-like peptide, produced in the central nervous system, acts on testis to induce steroidogenesis in bivalves. The 17 β -estradiol triggers spermatogonial proliferation and spermatogonia will then differentiate into spermatocytes I [9,10]. Each primary spermatocyte undergoes meiosis creating four haploid spermatids. Finally, spermatids undergo morphological transformations producing spermatozoa with an acrosome and a flagellum. Once spermatogenesis is completed, spermatozoa are stored in the gonad (**Fig. 1**) or released to the sperm duct where they are immotile and unable to fertilize.

Insert Fig. 1.

In bivalves, the male gamete consists of a head, a midpiece, and a flagellum [3,5,18] (**Fig. 2**). The head contains the nucleus with DNA material and an acrosome located at the anterior head region. The acrosome is a Golgi-derived secretory vesicle that contains enzymes involved in lysing the oocyte membrane during fertilization. Mitochondria and centrioles are located in the sperm midpiece. Mitochondria produce energy required for the motility apparatus of spermatozoa, also called axoneme. The distal centriole forms the basal body of the axoneme with a structure of 9+2 microtubules surrounded by a plasma membrane as described in Bondarenko and Cosson [19, this issue]. Dynein arms and radial spokes are involved in spermatozoa motility. Fueled by ATP hydrolysis, dynein arms generate force and movement on microtubules, regulating flagellar waveform and velocity [20-23]. It is worth to note that axonemal structure is highly conserved through evolution [24-26]. Spermatozoa motility is initiated at spawning after their release from the reproductive tract into the aquatic

environment. Various chemical signals, including pH, ions, and cyclic nucleotides, are involved in controlling sperm motility in bivalves. The present review aims at emphasizing the originality of spermatozoa motility in bivalves, including signaling for motility activation, flagellar beating, and energy metabolism of spermatozoa; and highlights physiological mechanisms that require further researches. The present work focus on ecologically and economically important bivalve species, including the Pacific oyster *Crassostrea gigas*, black-lip pearl oyster *Pinctada margaritifera*, European flat oyster *Ostrea edulis*, Manila clam *Ruditapes philippinarum*, great scallop *Pecten maximus*, and Japanese scallop *Patinopecten yessoensis*.

2. An overview of sperm biology in the most studied bivalve species

The present section reviews sperm biology in bivalves before delving into spermatozoa motility signaling processes. Morphology and motility kinetics of spermatozoa, and biochemical characteristics of seminal and blood plasma of most studied bivalve species are summarized in [Tables 1 and 2](#).

2.1. Pacific oyster, *Crassostrea gigas*

The Pacific oyster is a keystone species and one of the major bivalve species in aquaculture. This species inhabits estuaries and tolerates variations in salinity from 10 to 38 PSU with an optimum salinity range between 20 and 35 PSU [27]. Spawning occurs around 20°C and external fertilization happens in the water column. Successful fertilization requires contact between spermatozoa and oocytes for at least 10 min at a ratio of 500 spermatozoa per oocyte [27-29]. The head of Pacific oyster spermatozoa is spherical with a length and

width of 2-2.6 and 2.3-2.5 μm , respectively. The height and width of the acrosome are 0.7 and 1.2 μm , respectively. There are four mitochondria (0.8 μm diameter) in the midpiece. The length of flagellum is 37-41 μm [22,23].

2.2. Black-lip pearl oyster, *Pinctada margaritifera*

The head of the black-lip pearl oyster spermatozoon is spherical with an acrosome at the front of the head. There are four mitochondria located in the midpiece. Both proximal and distal centrioles are located in the midpiece, the latter being at the base of the flagellum. The length of flagellum is 47 μm [30,31].

2.3. European flat oyster, *Ostrea edulis*

The European flat oyster is a native species that has been over-exploited in the past and is nowadays classified as an endangered species in Europe. In this brooding species, oocytes are released in the inhalant chamber where fertilization takes place after sperm uptake by females. Spermatozoa are clustered in aggregates called “spermatozeugmata”, an acellular structure with a diameter of 25-80 (average: 60 μm) [32,33] in which spermatozoa heads are embedded. At spawning in seawater, spermatozoa are released from spermatozeugmata [33]. Therefore, biological characteristics of European flat oyster spermatozoa differ substantially from those observed in the Pacific and black-lip pearl oyster, with respect to the reproductive behavior.

2.4. Manila clam, *Ruditapes philippinarum*

The elongated pyramid-like structure of the head of Manila clam spermatozoa is about 6.2 μm long (Fig. 2). The nucleus is cone shaped. The acrosome is located at the tip of the head with a length of about 2.4 μm . There are four mitochondria in the midpiece as well as a pair of centrioles orthogonal to each other. The length of the flagellum is about 42-45 μm [34,35].

Insert Fig. 2.

2.5. Great scallop, *Pecten maximus*, and Japanese scallop, *Patinopecten yessoensis*

Spermatozoa of the great scallop have a bullet-shaped head with a short conical acrosome. There are four mitochondria with a pair of centrioles in the midpiece. In the Japanese scallop, the head is 3.9 μm long. The acrosome measures 0.5 μm both in length and diameter. According to micrographs, there are at least four mitochondria in the midpiece. The length of flagellum is about 50 μm [36].

2.6. Seminal plasma characteristics

Ionic composition and pH of seminal fluid have been studied in a few species of bivalves (Table 2). Osmolality of the testicular fluid and body fluid is very similar to that of seawater [37,38]. High osmolality of bivalve testicular fluid is due to its high Na^+ concentration. Interestingly, K^+ concentration is higher in bivalve testicular fluid than in seawater. Finally, pH of testicular fluid is lower than that of seawater in all studied species.

Insert Table 1

3. Spermatozoa motility in bivalves

Bivalve spermatozoa are immotile in the testis and do not possess fertilizing ability [39-41]. Spermatozoa ability to initiate motility after dilution in seawater varies among bivalve species and some movement characteristics of activated spermatozoa are reported in **Table 1**. Motility parameters of spermatozoa can be measured using a computer-assisted sperm analyzer (CASA) plug-in developed for Image J software [42], which has already been adapted to the Pacific oyster and black-lip pearl oyster spermatozoa [31,43].

3.1. Spermatozoa motility in seawater

In the Pacific oyster, motility of 57% [39] to 100% [44] spermatozoa are initiated in seawater, and the duration of motility lasts for several hours [43,44]. Spermatozoa of the Pacific oyster are motile for a long period, > 24 h [43,44] and up to 72 h post-activation (Boulais, unpublished results), which is among the longest of all marine bivalves studied until present. Studies report a time-dependent reduction of percentage of motile spermatozoa, decreasing down to less than 10% at 30 min [39] or at 24 h [43,44] post-activation in seawater. In this species, Dong et al. [45] reported a high variability in spermatozoa motility initiation (39-82%) after 120 s post dilution in seawater. Riesco et al. [46] reported that 45 to 85% of spermatozoa of the Portuguese oyster, *Crassostrea angulata*, are motile after activation in seawater, with movement lasting three days without any significant decrease in the percentage of motile cells. In the European flat oyster, spermatozoa released from spermatogonia are motile for about 10 min [33]. Lyons et al. [47] reported that motility of 75% of black-lip pearl oyster spermatozoa is triggered in seawater. However, Acosta-Salmón

et al. [48] observed that spermatozoa motility is not fully initiated in seawater, and reported a high inter-individual variation in the initiation of spermatozoa motility (20-53%). A later study showed that spermatozoa motility was not immediately initiated in seawater [30], confirming Acosta-Salmón et al. results [48]. Motility of 50-85% of spermatozoa was triggered after dilution in a saline solution called “DCSB4”, which is composed of 19.5 g L⁻¹ NaCl, 6.25 g L⁻¹ glycine, 0.15 g L⁻¹ CaCl₂, 0.19 g L⁻¹ MgSO₄; 2.42 g L⁻¹ Tris-aminomethane, pH 8.2 [30,31].

In the surf clam, *Spisula solidissima*, 6% of spermatozoa are motile when movement is triggered in seawater [49]. In the Manila clam, Alavi et al. [39] reported that spermatozoa motility lasts for seven days. Motility is triggered in 43% of spermatozoa at 1 min post dilution in seawater and this percentage increases up to 72% at 1 h post dilution in seawater.

Motility of great scallop and Japanese scallop spermatozoa is initiated in about 75% of cells following dilution in seawater [39,40]. This percentage decreases down to 45% within 30 min post-activation in the great scallop and to < 10% at 1-day post-dilution in the Japanese scallop [39]. However, Faure et al. [41] reported that great scallop spermatozoa motility is fully triggered after dilution in seawater with motility duration of 6 h after dilution.

Insert Table 2

3.2. Spermatozoa hyper-motility in bivalves

About eight years after a report by Matsutani and Nomura [16] showing the involvement of 5-HT in spawning regulation in the Japanese scallop, Kadam and Koide [49] reported that 5-HT, but not its metabolite 5-hydroxyindoleacetic acid (5-HIAA), triggers

spermatozoa motility in the surf clam. A recent study shows that 5-HT is capable of inducing spermatozoa hyper-motility in various bivalve species including the Manila clam, Japanese scallop, and Pacific oyster [39]. Percentage of motile spermatozoa in these species increases after dilution of sperm in seawater containing 5-HT. In the Manila clam, 90% of spermatozoa become motile within 3-5 min post-activation in the presence of 5-HT [39]. In the Japanese scallop, 85% of spermatozoa are motile at 2 h post-activation and decreases down to 40% within one day post-activation. Ability for 5-HT to induce hyper-motility differs in concentration among species $\geq 10^{-3}$, 5×10^{-6} , $\geq 10^{-6}$, and $\geq 10^{-7}$ M in Japanese scallop, surf clam, Manila clam, and Pacific oyster spermatozoa, respectively. Furthermore, it was observed that 5-HT-induced spermatozoa hyper-motility is time-dependent, suggesting that the stimulatory effect of 5-HT on spermatozoa hyper-motility may be mediated by a 5-HT receptor. It is worth to note that 5-HT-induced hyper-motility has also been reported in mammals including hamster and human spermatozoa [50,51] and this process is commonly known in mammals as 'hyper-activation'. Furthermore, our unpublished data shows that 5-HT hyperpolarizes membrane potential in 5-HT-induced hyper-motile spermatozoa.

Spermatozoa velocity of the Japanese scallop and Manila clam is higher after motility activation with 5-HT than seawater [39]. This might be due to stimulation of flagellar beating by 5-HT as it was reported in spermatozoa of the blue mussel and surf clam [20].

3.3. Serotonin-induced spermatozoa hyper-motility via 5-HT receptor(s)

So far, 14 structurally and pharmacologically distinct mammalian 5-HT receptors, which are classified into seven groups from 5-HT₁ to 5-HT₇, have been identified. Except for the 5-HT₃ receptor that is a ligand-gated ion channel [52,53], the 5-HT₁, 5-HT₂, 5-HT₄, 5-HT₅, 5-HT₆, and 5-HT₇ belong to the G protein-coupled receptor (GPCR) superfamily [54-

55]. In bivalves, 5-HT receptors were cloned in the Japanese Scallop, 5-HT_{py} [56], and the pearl oyster, *Pinctada fucata*, 5-HT_{pf} [57], and is also predicted for the Pacific oyster [58]. In invertebrates, pharmacological properties of 5-HT receptors do not allow to classify them in any of the mammalian categories, although some signal transduction characteristics are similar [59]. The 5-HT_{py} and 5-HT_{pf} are expressed in most of the organs, including ovary, testis, mantle, adductor muscle, gill, nervous system (cerebral-pedal ganglia and visceral ganglion), digestive gland, and kidney. *In situ* hybridization has demonstrated that 5-HT_{py} mRNA are localized in the oocytes, spermatids, and gonoduct epithelium [56]. Pharmacological characteristics of spermatozoa 5-HT receptor have only been studied in the surf clam [60]. Results reveal that 1 μ M ICS 205930 (antagonist of 5-HT₃ receptor), 2-methyl-5-HT (agonist of 5-HT₃ receptor), 8-OH-DPAT (agonist of 5-HT₁ receptor), BMY 7378 (antagonist of 5-HT₁ receptor), 5-HT, 5-CT (a non-selective agonist of 5-HT₁, 5 and 7 receptors), mianserin (antagonist of 5-HT₂ receptor), methysergide (antagonist of 5-HT₂), α -methyl-5-HT (a non-selective agonist of 5-HT₁ receptor), PBG (agonist of 5-HT₂ receptor), and ketanserin (antagonist of 5-HT₂ receptor) inhibits 45, 43, 37, 32, 31, 31, 30, 26, 13, 4, and 1% of [³H]5-HT binding to sperm plasma membrane, respectively. Considering pharmacological characterization of 5-HT receptors, analogs of 5-HT₃, 5-HT₁, and 5-HT₂ receptors are the most potent to compete with [³H]5-HT binding to the sperm plasma membrane, suggesting that a single or mixed 5-HT₁, 5-HT₂, and 5-HT₃ receptor exists in spermatozoa of bivalves [11,17]. Studies on various bivalve species clearly show the existence of 5-HT fibers in the gonad [61-64]. The 5-HT content has been frequently measured in bivalves and results highlight seasonal variations of 5-HT content in the nervous system and in the gonad with inter-sex differences [63-71].

The situation in vertebrates is similar to bivalves; metergoline (a non-selective antagonist of 5-HT receptors with affinity for 5-HT_{1B} and 5-HT_{1D} receptors) inhibits

spermatozoa motility initiation and fertilization [72]. Meizel and Turner [73] reported the involvement of 5-HT in spermatozoa capacitation in the golden hamster, *Mesocricetus auratus*. These authors reported that spermatozoa capacitated *in vitro* for 4.5 h undergo acrosome reaction within 15 min after addition of 50 μ M 5-HT or addition of 5 μ M 5-methoxytryptamine (5-MT, agonist of the 5-HT₁, 2, 4, 6, and 7 receptors with no affinity for the 5-HT₃ receptor). Furthermore, quipazine (agonist of 5-HT₃ and 5-HT_{2A} receptors) or cyproheptadine (antagonist of the 5-HT₂ receptor) inhibits the acrosome reaction. In rabbit, Young and Laing [74] reported 5-HT binding sites in the head and flagellum of spermatozoa. Jiménez-Trejo et al. [51] also observed that rat spermatozoa display immunoreactivity to the goat polyclonal anti-5-HT_{2A} and 3 receptors.

To better understand signaling pathways of 5-HT-induced spermatozoa motility, Kadam and Koide [49] studied the effects of 5-HT analogs on spermatozoa motility initiation in the surf clam. They observed that motility is initiated in 80, 70, and 76% of spermatozoa in the presence of 5 μ M of 8-OH-DPAT (agonist of 5-HT₁ receptor), 5-MT (agonist of the 5-HT₄ receptor), and 2-methyl-5-HT (a non-selective agonist of 5-HT₁ receptor), respectively; and that motility is not triggered when applying RU24969 (agonist of the 5-HT_{1B} receptor). However, treatment of sperm with mianserin (antagonist of 5-HT₂ receptor), ketanserin (antagonist of 5-HT₂ receptor), or metergoline (antagonist of 5-HT₁ receptor) does not interfere with 5-HT-induced spermatozoa motility initiation.

Overall, these studies demonstrate the involvement of 5-HT receptors in spermatozoa motility signaling and suggest the presence of a serotonergic system in the reproductive tract of male that might regulate sperm physiology and biochemistry. It is likely that 5-HT-induced hyper-motility in bivalve spermatozoa includes intracellular regulatory signals for axonemal beating and is mediated by 5-HT₁ or 5-HT₃ receptors.

3.4. Spermatozoa motility signaling in bivalves

3.4.1. Osmolality-independent activation of spermatozoa motility

Alavi et al. [39] investigated the potency of environmental osmolality to trigger spermatozoa motility in the Japanese scallop, Manila clam, and Pacific oyster by diluting spermatozoa in non-ionic media (1000 mM sucrose, 20 mM Tris, pH 8.0 ± 0.1 containing 5-HT or 10 mM Ca^{2+}) with an osmolality similar to that of seawater. In all studied species, spermatozoa motility was not triggered when activated in these media. Boulais et al. [37] reported that osmolality of Pacific oyster testicular fluid is 1061 ± 6 mOsmol L^{-1} , which is similar to the one of seawater. These results indicate that spermatozoa motility initiation in bivalves is an osmolality-independent mechanism. Dong et al. [45] observed in the Pacific oyster that seawater osmolality below 900 and above 1100 mOsmol kg^{-1} negatively affect spermatozoa motility initiation.

3.4.2. pH regulation of spermatozoa motility initiation

In bivalves, pH of the testicular fluid is lower than that of seawater, inhibiting spermatozoa motility activation [30,33,37,39,40] (Table 2). Spermatozoa of bivalves require an alkaline medium to become motile. Gonadal pH of the Pacific oyster was measured at 5.82 and spermatozoa motility is not triggered if seawater pH value is below 6.0 [37]. Percentages of motile spermatozoa were low ($< 10\%$) in pH values between 6.0 and 7.0, and the optimal range of pH values for spermatozoa motility ranged from pH 7.5 to 9.5 [37]. It was observed that percentage of motile spermatozoa is higher in the presence of 5-HT compared to 5-HT-free seawater for the same pH value. In Pacific oyster spermatozoa, motility is triggered

within 15 min post-activation in artificial seawater at pH 7.0 containing 5-HT [39]. Spermatozoa velocity (VAP: velocity of the average path) also increased in a pH-dependent manner from pH 6.5 to pH 7.5 and did not change from pH 7.5 to pH 9.5 [37].

In the black-lip pearl oyster, pH values of the male gonad and gonadal tract were 6.6 and 7.4, respectively [30]. DCSB4 (pH 8.2) behaves as a motility activator for black-lip pearl oyster spermatozoa. After transfer into alkaline DCSB4 medium, it takes a few minutes to spermatozoa to reach fully activated motility after which, spermatozoa remain motile for 10 min or more. Initiation of spermatozoa motility at alkaline pH is reversed when pH is shifted back to more acidic values. Furthermore, black-lip pearl oyster spermatozoa stored in regular seawater (pH 7.8) at 4°C retain their ability to be fully activated for several days. Indeed, spermatozoa motility can be initiated in an alkaline medium up to 13 days after collection and conservation at 4°C. During storage period, they saved their potential motility behavior and characteristics, even if the percentage of motile spermatozoa progressively declined to 50% [30,31].

In the great scallop, spermatozoa motility is not initiated in seawater pH value below 5.0, and fully triggered at pH above 8.0 [40]. In the Japanese scallop, spermatozoa motility is not triggered in seawater at pH 4, but fully activated at pH above 6.0 [39]. Movement of Manila clam spermatozoa is fully triggered in seawater at pH value above eight [39]. Inhibition of spermatozoa motility at low external pH ($[pH]_e$) is reversible since increasing the activation medium pH or washing spermatozoa with alkaline seawater results in motility triggering [40].

Alavi et al. [39] and Boulais et al. [37] observed time-course pH effects on spermatozoa motility initiation in bivalves. Indeed, spermatozoa become motile within a shorter period of time after dilution in seawater at high pH values (≥ 9) compared to lower pH values (< 9). As an example, Manila clam spermatozoa are fully motile at pH 8 and pH 9

within 15- and 3-min post-activation in seawater, respectively. Similarly, percentage of motile spermatozoa is higher at 5 min post-activation compared to 1 min post-activation in seawater at pH value < 8.5 in the Pacific oyster [37].

Since $[pH]_e$ mediates intracellular pH ($[pH]_i$) [75], some studies have been conducted to modify $[pH]_i$ using NH_3 or NH_4Cl , which rapidly alkalize $[pH]_i$ of spermatozoa. Results show that motility of Japanese pearl oyster and Manila clam spermatozoa is immediately and fully triggered in artificial seawater and 5-HT containing 2 mM NH_3 [39,76]. Similarly, NH_4Cl increases three times the percentage of motile spermatozoa compared to NH_4Cl -free seawater in the Pacific oyster [37]. Taken together, inhibition and stimulation of spermatozoa motility by decreasing and increasing seawater $[pH]_e$, respectively, are consistent with the hypothesis that an increase in $[pH]_i$ is required for optimal dynein ATPase activity and axonemal beating [75,77,78]. It has been reported that optimal pH values for the activity of dynein ATPase range from 7.4 to 8.6, and this activity is inhibited at pH < 7.2 [79]. These results suggest that acidic pH of testicular fluid is a factor that contributes to maintain spermatozoa in the quiescent state while they are in the testis.

3.4.3. K^+ regulation of spermatozoa motility initiation

It has been reported that K^+ ions concentration in testicular fluid of the Pacific oyster and Japanese scallop is higher than that of seawater [37,39] (Table 2). Increasing extracellular K^+ concentration ($[K^+]_e$) to the value of K^+ in the testicular fluid inhibits Japanese scallop spermatozoa motility associated with a decrease in spermatozoa velocity in seawater with or without 5-HT [39]. However, percentage of motile spermatozoa is higher in seawater containing 10 mM K^+ compared to seawater at 0, 40, 100, and 150 mM K^+ , suggesting that the concentration of 10 mM K^+ in seawater creates an optimal condition for

initiation and maintenance of sperm motility in bivalves. Inhibited spermatozoa motility due to $[K^+]_e$ is higher in seawater at pH 7.0 than seawater at pH 8.2. The opposite effect is observed in the presence of 5-HT.

To clarify the role of K^+ ions in spermatozoa motility initiation following sperm release into seawater, percentage of motile spermatozoa were measured in K^+ -free seawater with or without 5-HT [37,39]. Motility of Japanese scallop spermatozoa is triggered in K^+ -free seawater, but percentage of motile spermatozoa is lower than in seawater. Similarly, Boulais et al. [37] reported that spermatozoa of the Pacific oyster become fully motile after dilution in K^+ -free seawater, revealing that K^+ influx is not required to trigger spermatozoa motility. A voltage-dependent K^+ channel inhibitor, 4-AP, reduces Japanese scallop spermatozoa motility at 1 min post-activation in seawater and seawater containing 5-HT (both at pH 8.2) at 5 mM and 10 mM, respectively. Spermatozoa velocity also decreases in seawater and seawater containing 5-HT (both at pH 8.2) at 5 mM and 1 mM 4-AP, respectively. These results indicate that K^+ efflux through a voltage-dependent K^+ channel is required to trigger spermatozoa motility and suggest the involvement of K^+ ions in maintaining spermatozoa in the quiescent state in the testis. It is worth to note that K^+ -inhibited spermatozoa motility initiation is reported for salmonid and sturgeon fishes so far [80-83].

3.4.4. Na^+ regulation of spermatozoa motility initiation

Sodium is one of the main constituents of testicular fluid in bivalves [37,39] (Table 2). In the Japanese scallop, Manila clam, and Pacific oyster, spermatozoa motility is not triggered in Na^+ -rich activating medium (500 mM NaCl, 20 mM Tris, pH 8.0 ± 0.1) containing 5-HT or 10 mM Ca^{2+} . Moreover, adding 20 μ M A23187 (a Ca^{2+} ionophore) into

Na⁺-rich activating medium at 10 mM Ca²⁺ does not activate spermatozoa motility [39]. These results indicate that a Na⁺-rich medium, at an osmolality identical to that of seawater, does not allow initiation of spermatozoa motility, hence containing Ca²⁺ ions.

Even if a high concentration of Na⁺ does not trigger spermatozoa motility, extracellular Na⁺ ions ([Na⁺]_e) are needed for motility. Indeed, it was demonstrated that spermatozoa motility is not initiated in Na⁺-free artificial seawater in the great scallop, Japanese scallop, Manila clam, and Pacific oyster [37,39,40]. Faure et al. [40] reported that seawater containing > 150 mM Na⁺ is required to trigger spermatozoa motility in the great scallop. Inhibition of spermatozoa motility in Na⁺-free seawater is reversible since washing sperm with artificial seawater containing Na⁺ results in triggering spermatozoa motility [40].

In seawater containing 5-HT, [Na⁺]_e requirement for spermatozoa motility initiation seems to be species specific. In the Japanese scallop, spermatozoa motility is not stimulated in Na⁺-free 5-HT, but spermatozoa of the Pacific oyster become motile in Na⁺-free 5-HT although percentage of motility is lower than that of spermatozoa diluted in seawater containing 5-HT. These observations highlight that 5-HT overcomes inhibition of spermatozoa motility initiation in Na⁺-free seawater in the Pacific oyster. Since 5-HT triggers motility of Pacific oyster spermatozoa in Na⁺-free 5-HT, the role of Ca²⁺ ion was also investigated [39]. Chelating extracellular Ca²⁺ ions ([Ca²⁺]_e) results in full suppression of spermatozoa motility activation in Na⁺-free 5-HT. Further experiments using flunarizine, a Na⁺/Ca²⁺ exchanger (NCX) blocker [84,85], resulted in inhibition of spermatozoa motility associated with a decrease in spermatozoa velocity [39]. Overall, [Na⁺]_e is essential to spermatozoa motility initiation, which requires Na⁺ influx mediated by NCX and by regulation of [Ca²⁺]_i.

3.4.5. Ca²⁺ regulation of spermatozoa motility initiation

In the testicular fluid, Ca^{2+} concentration is similar to that of seawater [37,39]. Current knowledge on the physiological roles of Ca^{2+} ions in bivalve spermatozoa motility initiation reveals some differences among species and studies. In the Manila clam, chelating $[\text{Ca}^{2+}]_e$ by 5 mM ethylene glycol-bis(β -aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA) totally suppresses spermatozoa movement activation in seawater with or without 5-HT. In the Japanese scallop, spermatozoa motility is decreased in seawater and 5-HT seawater in the presence of 2.5–5 mM EGTA and movement is totally suppressed at 10 mM EGTA. However, 5-HT seawater containing 1 mM EGTA has no effects on spermatozoa motility initiation in this species compared to 5-HT seawater. In the Pacific oyster, EGTA (range from 1 to 10 mM) reduces the percentage of motile spermatozoa in a dose-dependent manner and spermatozoa motility is not initiated at ≥ 5 mM EGTA [39]. In contrast to this observation, Boulais et al. [37] reported that seawater at 1 mM EGTA increases the percentage of motile spermatozoa compared to seawater in the Pacific oyster [37]. Similarly, other authors reported that Ca^{2+} -free seawater increases motility of Pacific oyster spermatozoa [45].

At 1 mM EGTA, spermatozoa motility of the Japanese scallop is higher in seawater containing 5-HT compared to seawater without 5-HT. In the Pacific oyster, once spermatozoa hyper-motility is induced in 5-HT seawater, higher concentration of EGTA is required to suppress initiation of spermatozoa motility compared to seawater [39]. These studies suggest that spermatozoa may require lower $[\text{Ca}^{2+}]_e$ to trigger initiation of motility in the presence of 5-HT compared to seawater.

Regarding the role of Ca^{2+} in spermatozoa motility activation, it was reported that Ca^{2+} influx is essential for initiation of spermatozoa motility, however Ca^{2+} itself is not the primary factor triggering spermatozoa motility, because the addition of Ca^{2+} ionophore to sucrose- or to Na^+ -activating medium containing 10 mM Ca^{2+} did not trigger sperm motility

[39]. Plasma membrane NCX are one way in which cells regulate Ca^{2+} [86], and Boulais et al. [37] suggested that the higher percentage of motile spermatozoa in Ca^{2+} -free artificial seawater compared to seawater is associated with changes in $[\text{Na}^+]_i$ concentration. Investigating the roles of NCX in controlling spermatozoa motility initiation would be of interest for future studies. It might be possible that differences among studies in Ca^{2+} -dependent initiation of spermatozoa motility might be related to intracellular Ca^{2+} ($[\text{Ca}^{2+}]_i$) store; however the role of $[\text{Ca}^{2+}]_i$ store is still unclear. In mammalian spermatozoa, intracellular Ca^{2+} stores are located in the acrosome, neck, and midpiece [87,88]. In mammals, similar to somatic cells, mobilization of stored Ca^{2+} in sperm may occur in response to generation of inositol-1,4,5-triphosphate (IP_3) by activity of phospholipase C and by Ca^{2+} release at inositol-1,4,5-triphosphate-(IP_3)-sensitive Ca^{2+} channel receptor or the ryanodine receptor following Ca^{2+} influx [89-91].

It was observed that spermatozoa motility in the Japanese scallop decreases following activation in seawater containing Ca^{2+} channel blockers including 50 μM mibefradil, 200 μM verapamil, and 200 μM nifedipine at 1, 120, and 15 min post-activation, respectively [39]. In seawater containing 5-HT, spermatozoa motility is decreased in a shorter period of time compared to seawater by mibefradil, verapamil, and nifedipine at 1, 1, and 5 min post-activation, respectively. Spermatozoa velocity in the Japanese scallop also decreases in seawater with or without 5-HT containing aforementioned Ca^{2+} channel blockers at 1 min post-activation. The observed decrease in both spermatozoa motility and velocity following activation in seawater or 5-HT containing Ca^{2+} channel blockers suggests that Ca^{2+} influx is mediated by voltage-dependent Ca^{2+} channels.

To better understand role of Ca^{2+} in spermatozoa motility initiation, changes in $[\text{Ca}^{2+}]_i$ were investigated in Manila clam spermatozoa following activation in seawater with or without 5-HT containing 10 mM EGTA or 50 μM mibefradil [39]. In this species, 5-HT

triggers spermatozoa motility in a time-dependent manner. Spermatozoa motility initiation is associated with time-dependent increase and oscillation in $[Ca^{2+}]_i$. In seawater, $[Ca^{2+}]_i$ increases and Ca^{2+} oscillations are similar to those observed in 5-HT. In the presence of EGTA, no $[Ca^{2+}]_i$ increase and oscillations are observed in seawater with or without 5-HT. In the presence of mibefradil, Ca^{2+} oscillations are observed, but $[Ca^{2+}]_i$ decrease in 5-HT activated spermatozoa throughout the activation period. These results confirm that $[Ca^{2+}]_e$ play an important role in spermatozoa motility activation. Further experiments have shown that spermatozoa motility in the Manila clam and Pacific oyster is suppressed in artificial seawater and 5-HT containing 100 or 200 μ M W-7 (an inhibitor for Ca^{2+} -calmodulin (CaM) dependent phosphodiesterase), which is explained by $[Ca^{2+}]_e$ requirement for Ca^{2+} /CaM-dependent flagellar beating [39]. Taken together, these results indicate that $[Ca^{2+}]_e$ is required for spermatozoa motility initiation in bivalves. It is suggested that Ca^{2+} influx, mediated by voltage-dependent Ca^{2+} channels, is required to activate Ca^{2+} /CaM proteins in the axoneme leading to flagellar beating. Investigating the roles of NCX, $[Ca^{2+}]_i$ store, and their physiological interactions in controlling spermatozoa motility initiation would be of interest for future studies. Moreover, considering the presence of Ca^{2+} in the testicular fluid, Ca^{2+} is not an inhibitory factor maintaining sperm in the quiescent state in the testis.

3.4.6. Cyclic nucleotide regulation of spermatozoa motility initiation

Our current knowledge about contribution of cyclic nucleotide to regulation of spermatozoa motility initiation in bivalves is limited. Various cation channels modulate cAMP [92] and cGMP levels [93] in spermatozoa of marine organisms. These messengers are essential for the activation of spermatozoa motility via cAMP-dependent phosphorylation of axonemal proteins. In the case of Pacific oyster spermatozoa, cAMP activation is not

related with control neither by K^+ nor by Ca^{2+} , and regulation of spermatozoa cAMP is still unclear in this species. It may be possible that 5-HT is involved in controlling intracellular cAMP rise in bivalve spermatozoa.

In the Asian clam, *Corbicula fluminea*, spermatozoa are biflagellate with equal size, each has a typical “9+2” microtubule axoneme structure, and both flagella are beating [94-96]. The Asian clam is a freshwater species and simultaneous hermaphrodite capable of self-fertilization occurring internally in the gill chamber [97]. Howard et al. [94] observed that spermatozoa of the Asian clam are immotile in the gonad and in freshwater. cGMP triggered spermatozoa motility initiation, however percentage of motile spermatozoa was lowered than those activated by cAMP. cAMP analogs (1 mM dbcAMP and 2 mM 8-Br-cAMP) as well as a phosphodiesterase inhibitor (2 mM IBMX) trigger spermatozoa motility initiation. Percentage of motile spermatozoa is similar when they are treated with dbcAMP or 8-Br-cAMP alone or in combination with IBMX, suggesting that an increase in intracellular cAMP is able to trigger spermatozoa motility initiation. As IBMX is able to inhibit cGMP phosphodiesterases resulting in an increase in cGMP, even more than cAMP in flagella of sea urchin spermatozoa [98], Howard et al. [94] examined the effects of cGMP analogs on sperm motility initiation. They observed that 2 mM 8-Br-cGMP alone or in combination with 8-Br-cAMP triggers sperm motility initiation, suggesting that cAMP and cGMP activate the same population of spermatozoa. In further experiments, these authors [94] used a selective cAMP-dependent protein kinase (PKA) inhibitor (H-89) to clarify cAMP signaling in spermatozoa upon motility initiation. Spermatozoa motility was inhibited by 1 μ M H-89 when activated by 8-Br-cAMP but not when activated by 8-Br-cGMP. In presence of H-89, without 8-Br-cAMP or 8-Br-cGMP, spermatozoa motility is not initiated. To study the contribution of cGMP-dependent protein kinase (PKG) in spermatozoa motility initiation, Howard et al. [94] used a PKG inhibitor (Rp-8-pCPT-cGMPS). In the presence of 5 μ M Rp-8-pCPT-cGMPS,

spermatozoa motility is not initiated, and cGMP-induced spermatozoa motility was inhibited. The Rp-8-pCPT-cGMPS was without effects on cAMP-induced spermatozoa motility initiation. These results indicate that involvement of PKA in spermatozoa motility initiation is downstream of cAMP, but not of cGMP. The PKG is also involved in spermatozoa motility initiation, and it is downstream of cGMP, but not cAMP.

3.4.7. Signaling pathway controlling spermatozoa motility initiation in bivalves

Investigations of ionic signaling in spermatozoa motility initiation highlight that high concentration of K^+ and acidic pH of testicular fluid are the main inhibitory factors of spermatozoa motility in testis of bivalves. Spermatozoa motility and hyper-motility initiation is an osmolality-independent mechanism. Since dynein ATPase activity is inhibited at pH < 7.2, alkalization of spermatozoa $[pH]_i$ is mandatory to activate flagellar beating. During spawning, $[pH]_e$ and $[Na^+]_e$ play major physiological role in spermatozoa motility signaling, since motility is triggered by a Na^+ -dependent $[pH]_i$ alkalization probably mediated by a Na^+/H^+ exchanger (Fig. 3). Consequently, $[pH]_i$ but not Na^+ is the primary factor controlling spermatozoa motility activation. 5-HT rise in the testis contributes to spermatozoa motility initiation by inducing hyperpolarization of spermatozoa membrane potential (Fig. 3). Membrane hyperpolarization is also mediated by K^+ efflux via voltage-dependent K^+ channels and associated with opening of voltage-dependent Ca^{2+} channels under alkaline conditions. Release of Ca^{2+} from intracellular stores may also contribute to increase $[Ca^{2+}]_i$. Subsequent increase in $[Ca^{2+}]_i$ triggers Ca^{2+}/CaM -dependent flagellar beating. $[Na^+]_e$ probably regulates Ca^{2+} exchange via NCX. Furthermore, it may be possible that 5-HT is involved in controlling intracellular cAMP rise in bivalve spermatozoa, leading to cAMP-dependent phosphorylation of axonemal proteins.

Insert Fig. 3

4. Flagellar beating in spermatozoa of bivalves

Once spermatozoa motility in bivalves is triggered in seawater with or without 5-HT, flagellum beats in asymmetric wave pattern (**Fig. 4; Video 1**). The asymmetric beating of the flagellum displays a circular movement of spermatozoa (**Fig. 5**). However, it is unclear which axonemal protein(s) is involved in propagation of asymmetric wave.

Insert Fig. 4 and Fig. 5

Demoy-Schneider and al. [30,31] and Faure [99] described flagellar beating in activated spermatozoa. Three different flagellar wave characteristics were reported, including “full”, “twitching” and “declining” propagation of wave. In spermatozoa with fully propagated wave, flagellar waves are developed along flagella, and all spermatozoa have a similar flagellar shape described as flagellar sinusoidal and planar waves from the spermatozoon head to distal tip (**Fig. 6 and Fig. 7**). Wave amplitude (5 to 6 μm) is nearly constant along the flagellum length. Sperm swimming tracks are almost mostly circular with head trajectories describing arcs interspersed with short linear segments. Indeed, sperm tracks draw large arcs alternating with tighter ones, drawing a polygonal shape (**Fig. 5**). In addition, sperm are drawing tracks according to a clockwise direction for spermatozoa swimming close to the cover slip surface and counter-clockwise for cells swimming in the vicinity of glass slide surface. Sperm heads alternate both sides of the mean track.

Insert Fig. 6 and Fig. 7

Spermatozoa showing only one local bending point along the flagellum, and thus no flagellar sinusoidal waves, do not show an efficient movement, which is described as “twitching” behavior (Fig. 8). This occurs mainly during a short period of time at the beginning of movement activation for spermatozoa collected into seawater. Before reaching the behavior of fully activated, previously described, spermatozoa will go through “half activated” (around 50% of motile spermatozoa) and alternatively twitching and stopping several times (Fig. 8). High viscosity swimming medium alters wave shape of spermatozoa. Flagellar wave progresses along flagellum, but wave beat frequency is about five times lower than in seawater leading to reduced forward progression in viscous seawater (Fig. 9).

Insert Fig. 8 and Fig. 9

5. Energy metabolism in spermatozoa of bivalves

5.1. Mitochondrial respiration

In contrast to fish spermatozoa which present a short period of motility, from 30 s to 20 min maximum, respiration rate of oyster spermatozoa can be easily measured because of the long duration of their motility phase. Mitochondrial respiration is closely bound to the activation of motility since spermatozoa oxygen consumption is strongly increased during motility activation. In the black-lip pearl oyster, oxygen consumption is ranging from 0 prior to motility activation to 29 nmol O₂ min⁻¹ per 10⁹ spermatozoa after activation in an alkaline solution (seawater at pH 10.5) [30]. In Pacific oyster spermatozoa activated in seawater,

oxygen consumption is ranging from 56 to 65 nmol O₂ min⁻¹ per 10⁹ spermatozoa at 10 and 120 min after motility initiation, respectively. Maximum respiration rate value is obtained by dissipation of the proton gradient across the inner mitochondrial membrane using an oxidative phosphorylation (OXPHOS) uncoupler such as CCCP (Carbonyl cyanide m-chlorophenyl hydrazine). Maximal rate of mitochondrial oxygen consumption is 151.8 ± 6.5 nmol O₂ min⁻¹ per 10⁹ spermatozoa in the Pacific oyster (Boulais, unpublished results), highlighting that electron transport activity of the mitochondrial respiratory chain can be greatly increased in Pacific oyster spermatozoa.

5.2. Intracellular ATP concentration

Adenosine triphosphate (ATP) is required to fuel spermatozoa movement phase because flagellar movement occurs through a reaction catalyzed by dynein-ATPases located in the flagellum [100]. In most marine species, including fish and sea urchins, intracellular ATP content controls the duration of sperm movement phase. Indeed, ATP content declines within tens of seconds to few minutes depending on species leading to arrest of spermatozoa movement [101]. Changes in ATP content during the movement phase of spermatozoa have been little documented in bivalves. In the great scallop, the halt in spermatozoa movement after a 10 h motility phase is due to the exhaustion of intracellular ATP content [102]. Two min after motility activation, median intracellular ATP concentration of black-lip pearl oyster spermatozoa is 700 nmol per 10⁹ spermatozoa after activation in seawater and 543 nmol per 10⁹ spermatozoa when activated in the alkaline solution DCSB4 [30]. In the Pacific oyster, Boulais et al. [103] reported an intracellular ATP content of 157.2 ± 36.5 nmol 10⁻⁹ spermatozoa at 1 h after activation in seawater. In this species, a stable level of ATP throughout 24-h motility period of spermatozoa is observed, with ATP concentrations of

185.6 \pm 15.7 nmol per 10⁹ spermatozoa and 140.1 \pm 7.8 nmol per 10⁹ spermatozoa at 90 min and 9 h after activation, respectively, remaining at this level for the rest of the movement phase. A similar pattern was previously reported in spermatozoa of this species with an ATP content remaining constant until the end of motility [44]. Overall, energy metabolism of Pacific oyster spermatozoa seems to differ from previously studied marine species.

In the Pacific oyster, ending of spermatozoa movement is not explained by a low intracellular ATP content (75-94% of the initial content) [43,44] nor by the death of spermatozoa since percentage of live cells remain high (> 94%) at the end of the motility phase (24 h post activation) [43]. One hypothesis to explain the halt in sperm movement could be the disruption of ATP transport from mitochondria to the flagellum, which would suggest the involvement of phosphagens (see section 5.3).

5.3. Metabolic pathways synthesizing ATP

There are two metabolic pathways producing ATP: OXPHOS and glycolysis. OXPHOS is the most efficient way to produce ATP, because it produces 19 times more ATP per molecule of glucose than glycolysis and takes place in the mitochondria. These pathways have been little documented in marine invertebrate spermatozoa. In sea urchins, spermatozoa do not undergo glycolysis and rely entirely on the oxidation of endogenous substrates [104-107]. The long movement phase of Pacific oyster spermatozoa compared to other marine invertebrates [44] led to a study to investigate the involvement of OXPHOS in sustaining Pacific oyster sperm movement [43]. Using CCCP, it was demonstrated that OXPHOS produces a stable level of ATP throughout 24-h motility period of Pacific oyster spermatozoa. Boulais et al. [43] suggested that the steady decrease of intracellular ATP content during the

first 9 h of the movement phase stimulated ATP synthesis via OXPHOS, which compensates for ATP hydrolysis from 9 h post activation to the end of the movement phase.

Interestingly, when mitochondrial ATP synthesis is inhibited by 1 μ M CCCP, Pacific oyster spermatozoa are motile during the first 2.5 h of movement [43]. A decrease in the percentage of motile spermatozoa is observed from 2.5 h and totally stopped after 4 h of incubation with CCCP whereas spermatozoa are still motile in the control after 4 h of activation. These results reveal that ATP-sustaining flagellar movement can originate from alternative metabolic processes during the first 2 h of motility, including glycolysis, mobilization of stored ATP or phosphagens. Phosphagens are high-energy compounds, such as arginine- or creatine-phosphate, that serve as intracellular ATP transport from mitochondria to ATPases and as ATP storage for subsequent use during periods of high-energy need [108].

Overall, both the low long motility phase and energy metabolism, including ability to produce ATP during several hours with or without OXPHOS and to increase respiration, of Pacific oyster spermatozoa reveal a specific strategy developed by this species, resulting in a potentially long total distance covered by spermatozoa during their movement phase. Such a strategy may reflect the adaptation of Pacific oyster to sessile life, improving oocyte fertilization success [109]. Furthermore, this movement strategy may be one of the biological traits contributing to the dispersion of Pacific oyster in the wild [110].

6. Conclusion and future researches

Current knowledge highlights that pH and ions control spermatozoa motility in bivalves. Acidic pH and high concentration of K^+ in the seminal fluid of bivalves maintain spermatozoa in the quiescent state. In contrast to fish spermatozoa, osmolality is not involved

in spermatozoa movement in bivalves. During spawning, Na^+ influx allows alkalization of intracellular pH. Decrease of extracellular K^+ causes K^+ efflux leading to membrane hyperpolarization and increase of intracellular Ca^{2+} . These events activate dynein ATPases and Ca^{2+} /CaM-dependent proteins resulting in ATP-dependent flagellum beating. In contrast to the short motility phase of fish spermatozoa, from a few seconds to minutes, bivalve spermatozoa are motile for a long period of time, up to several days in the Pacific oyster. This is achieved by spermatozoa capacity to partly compensate for ATP hydrolysis during movement phase. Motility duration and energy metabolism of spermatozoa highlight an adaptive strategy of bivalves to sedentary life and absence of breeding mates, in contrast to mating behavior in fish. This results in a potentially long total distance covered by spermatozoa during their movement phase and improving oocyte fertilization success.

The long motility period of bivalve spermatozoa is an advantage to investigate mechanisms of spermatozoa motility. Identification and characterization of ion channels that regulate ion fluxes remain to be studied. Proteins activated by cAMP and Ca^{2+} and leading to axonemal beating are still unknown. Furthermore, asymmetrical beating of bivalve spermatozoa flagella is an asset to study evolutionary characterization of axonemal proteins that control spermatozoa motility behavior. The mechanism through which 5-HT induces spermatozoa hyper-motility is still unclear. As molecular identifications reveal the presence of 5-HT receptor(s) in the testis and spermatozoa, further research effort might also be directed towards producing bivalves without 5-HT receptor using genome manipulation. Mechanisms of action of 5-HT need to be studied. It might be possible that 5-HT induces adenylyl cyclase via G_{as} to produce cAMP for activation of PKA and/or induces phospholipase C via $G_{aq/11}$ to produce inositol (1,4,5) trisphosphate (IP_3) and release of Ca^{2+} from intracellular stores. Contribution of Ca^{2+} released from intracellular stores would be of interest to address in the future. In the end, control of spermatozoa motility initiation, for

instance via 5-HT- or pH-induced spermatozoa motility, would help to manage artificial reproduction in bivalve aquaculture.

Acknowledgements

We are grateful to Profs. M. Osada, K. Inaba, M. Morisawa; Associate Profs. K.G. Takahashi, N. Itoh, M. Yoshida, and Assistant Profs. K. Shiba, K. Nagasawa (Tohoku University, University of Tsukuba, University of Tokyo, Japan), and M. Suquet (IFREMER, France) for their supports, advises, and encouragements during research on the biology of sperm in bivalve mollusks. We also thank M. Suquet for his critical reading of the manuscript draft. MB research was supported by CRB-anim (ANR-11-INBS-003) and Reproseed (FP7-KBBE-2009-3). MDS work was supported by the French *Ministère des Outremer*s and the research program “La cryoconservation des spermatozoïdes de l’huître perlière *Pinctada margaritifera*, un atout pour une perliculture durable et la sauvegarde de sa biodiversité”. SMHA was supported by the University of Tehran and by the National Elites Foundation of Iran. JC is thankful to the agencies that supported our work, among which grants from Czech Republic: CENAKVA LM2018099, CZ.02.1.01./0.0/0.0/16_025/0007370 Reproductive and genetic procedures for preserving fish biodiversity and aquaculture, and by the Grant Agency of the University of South Bohemia in Ceske Budejovice (125/2016/Z).

Conflict of interest

The authors declare that they have no conflict and financial interests.

References

- [1] Food and Agriculture Organization of the United Nations. Fishery and Aquaculture Statistics 2016. <http://www.fao.org/fishery/statistics/global-production/query/en> [accessed 11 March 2019].
- [2] Barnes RSK, Calow P, Olive PJW, Golding DW. The invertebrates: A synthesis. Cambridge: Blackwell Scientific Ltd; 1993. ISBN 0-632-04761-5.
- [3] Sastry AN. Pelecypoda (excluding Ostreidae). In: Giese AC, Pearse JS, editors. Reproduction of marine invertebrates, Vol. V: Molluscs: Pelecypods and lesser classes, New York: Academic Press; 1979, p. 113-292.
- [4] Dohmen MR. Gametogenesis. In: Verdonk NH, van den Biggelaar JAM, Tompa AS, editors. The Mollusca, Vol. 3: Development, New York: Academic Press Inc; 1983, p. 1-48.
- [5] Mackie GL. Bivalves. In: Tompa AS, Verdonk NH, van den Biggelaar JAM, editors. The Mollusca: Vol. 7: Reproduction, New York: Academic Press Inc; 1984, p. 351-418.
- [6] Gosling E. Bivalve molluscs: Biology, ecology and culture. Oxford: Blackwell Publishing; 2003, p. 1-456. ISBN: 978-0-852-38234-9
- [7] Fabioux C, Huvet A, Le Souchu P, Le Pennec M, Pouvreau S. Temperature and photoperiod drive *Crassostrea gigas* reproductive internal clock. Aquaculture 2005;250:458-470. <https://doi.org/10.1016/j.aquaculture.2005.02.038>.
- [8] Croll RP, Wang C. Possible roles of sex steroids in the control of reproduction in bivalve molluscs. Aquaculture 2007;272:76-86. <https://doi.org/10.1016/j.aquaculture.2007.06.031>.
- [9] Osada M, Treen N. Molluscan GnRH Associated with Reproduction. Gen Comp Endocrinol 2013;181:254-258. <https://doi.org/10.1016/j.ygcen.2012.09.002>.
- [10] Osada M, Matsumoto T. Endocrine Control of Gametogenesis and Spawning in Bivalves. In: Saleuddin, S, Mukai S, editors. Physiology of molluscs: A collection of selected reviews, Vol. 2, Oakville, ON, Canada: Apple Academic Press; 2019, p. 179-219.
- [11] Alavi SMH, Nagasawa K, Takahashi KG, Osada M. Structure-function of serotonin in bivalve molluscs. In: Kaneez FS, editor. Serotonin - A chemical messenger between all types of living cells, London: InTechOpen; 2017, p. 33-63. <https://doi.org/10.5772/intechopen.69165>.
- [12] Welsh JH. Serotonin as a possible neurohumoral agent: Evidence obtained in lower animals. Ann NY Acad Sci 1957;66:618-630. <https://doi.org/10.1111/j.1749-6632.1957.tb40752.x>.
- [13] Welsh JH. Excitation of the heart of *Venus mercenaria*. Arch Exp Pathol Pharmacol 1953;219:23-29.
- [14] Twarog BM. Responses of a molluscan smooth muscle to acetylcholine and 5-hydroxytryptamine. J Cell Comp Physiol 1954;44:141-163. <https://doi.org/10.1002/jcp.1030440112>.
- [15] Hoyle G, Lowy J. The paradox of *Mytilus* muscle: A new interpretation. J Exp Biol 1956;33:295-310.
- [16] Matsutani T, Nomura T. Induction of spawning by serotonin in the scallop *Patinopecten yessoensis*. Marine Biol Lett 1982;3:353-358.
- [17] Alavi SMH, Nagasawa K, Takahashi KG, Osada M. Pharmacology and molecular identity of serotonin receptor in bivalve mollusks. In: Kaneez FS, editor. Serotonin - A chemical messenger between all types of living cells, London: InTechOpen; 2017, p. 7-31. <https://doi.org/10.5772/intechopen.69680>.
- [18] Morse MP, Zardus JD. Bivalvia. In: Harrison FW, Kohn AJ, editors. Microscopic anatomy of invertebrates, Vol. 6A: Mollusca II, New York: John Wiley & Sons Inc; 1997, p. 7-118.

- [19] Bondarenko V, Cosson J. Structure and beating behavior of the sperm motility apparatus of aquatic animals. *Theriogenology* 2019, this issue.
- [20] Stephens RE, Prior G. Dynein from serotonin-activated cilia and flagella: extraction characteristics and distinct sites for cAMP-dependent protein phosphorylation. *J. Cell Sci* 1992;103:999-1012.
- [21] Gwo JC, Yang WT, Shen YT, Cheng HY. Spermatozoan morphology of four species of bivalve (Heterodonta, Veneridae) from Taiwan. *Tissue Cell* 2002;34:39-43. <https://doi.org/10.1054/tice.2002.0222>.
- [22] Dong Q, Huang C, Tiersch TR. Spermatozoal ultrastructure of diploid and tetraploid Pacific oysters. *Aquaculture* 2005;249:487-496. <https://doi.org/10.1016/j.aquaculture.2005.03.009>.
- [23] Drozdov AL, Sharina SN, Tyurin SA. Sperm ultrastructure in representatives of six bivalve families from Peter the Great Bay, Sea of Japan. *Russ J Mar Biol* 2009;35:236-241.
- [24] Witman GB. Axonemal dyneins. *Curr Opin Cell Biol* 1992;4:74-79. [https://doi.org/10.1016/0955-0674\(92\)90061-G](https://doi.org/10.1016/0955-0674(92)90061-G).
- [25] Inaba K. Molecular basis of sperm flagellar axonemes: structural and evolutionary aspects. *Ann NY Acad Sci* 2007;1101:506-526. <https://doi.org/10.1196/annals.1389.017>.
- [26] Inaba K. Calcium sensors of ciliary outer arm dynein: functions and phylogenetic considerations for eukaryotic evolution. *Cilia* 2015;4:6. <https://doi.org/10.1186/s13630-015-0015-z>.
- [27] Kinnby A. Effects of reduced salinity on fertilization and larval development in the Pacific Oyster, *Crassostrea gigas* (Thunberg, 1789). Master of Science Thesis, Department of Biological and Environmental Sciences, University of Gothenburg, 2015.
- [28] FAO Fisheries and Aquaculture Department. Species fact sheet - *Crassostrea gigas*, http://www.fao.org/fishery/culturedspecies/Crassostrea_gigas/en; 2007 [accessed 11 March 2019].
- [29] Song YP, Suquet M, Quéau I, Lebrun L. Setting of a procedure for experimental fertilisation of Pacific oyster (*Crassostrea gigas*) oocytes. *Aquaculture* 2009;287:311-314. <https://doi.org/10.1016/j.aquaculture.2008.10.018>
- [30] Demoy-Schneider M, Levêque A, Schmitt N, Le Pennec M, Cosson J. Motility activation and metabolism characteristics of spermatozoa of the black-lip-pearl oyster *Pinctada margaritifera* var: *cumingii* (Jameson 1901). *Theriogenology* 2012;77:53-64. <https://doi.org/10.1016/j.theriogenology.2011.07.014>.
- [31] Demoy-Schneider M, Schmitt N, Le Pennec G, Suquet M, Cosson J. Quality assessment of cryopreserved black-lip pearl oyster *Pinctada margaritifera* spermatozoa. *Aquaculture* 2018;497:278-286. <https://doi.org/10.1016/j.aquaculture.2018.07.067>.
- [32] Foighil DO. Role of spermatozeugmata in the spawning ecology of the brooding oyster *Ostrea edulis*. *Gamete Res* 1989;24:219-28. <https://doi.org/10.1002/mrd.1120240209>.
- [33] Suquet M, Pouvreau S, Queau I, Boulais M, Le Grand J, Ratiskol D, Cosson J. Biological characteristics of sperm in European flat oyster (*Ostrea edulis*) *Aquat Living Resour* 2018;31:20. <https://doi.org/10.1051/alr/2018008>.
- [34] Dai PF, Rao XZ, Chen YS. Ultrastructural studies on spermatozoon and spermatogenesis of *Ruditapes philippinarum*. *Chin J Zool* 2004;39:26-32
- [35] Kim JH, Chung JS, Lee K-Y. Ultrastructural Characteristics of the Testis, Spermatogenesis and Taxonomic Values of Sperm Morphology in Male *Ruditapes philippinarum* in Western Korea. *Dev Reprod* 2013;17:121-132. <https://doi.org/10.12717/DR.2013.17.2.121>.

- [36] Li Q, Osada M, Kashihara M, Hirohashi K, Kijima A. Effects of ultraviolet irradiation on genetical inactivation and morphological structure of sperm of the Japanese scallop, *Patinopecten yessoensis*. *Aquaculture* 2000;186:233-242. [https://doi.org/10.1016/S0044-8486\(99\)00384-1](https://doi.org/10.1016/S0044-8486(99)00384-1)
- [37] Boulais M, Suquet M, Arsenault-Pernet EJ, Malo F, Queau I, Pignet P, Ratiskol D, Le Grand J, Huber M, Cosson J. pH controls spermatozoa motility in the Pacific oyster (*Crassostrea gigas*). *Biol Open* 2018;7:bio031427. <https://doi.org/10.1242/bio.031427>.
- [38] Melo FFAP. Development and standardization of a protocol for sperm cryopreservation of two important commercial oyster species. Thesis for Master of Science Degree, Faculty of Sciences and Technology, University of Algarve, 2015, 65 pages. https://sapientia.ualg.pt/bitstream/10400.1/7863/1/TESE_Francisca_Final_2.pdf [accessed 11 March 2019].
- [39] Alavi SMH, Matsumura N, Shiba K, Itoh N, Takahashi KG, Inaba K, Osada M. Roles of extracellular ions and pH in 5-HT-induced sperm motility in marine bivalve. *Reproduction* 2014;147:331-345. <https://doi.org/10.1530/REP-13-0418>.
- [40] Faure C, Devauchelle N, Girard JP. Ionic factors affecting motility, respiration and fertilization of the sperm of the bivalve *Pecten maximus* (L.). *J Comp Physiol Part A* 1994;164:444-450. <https://doi.org/10.1007/BF00714581>.
- [41] Faure C, Devauchelle N, Girard J-P, Cosson J. The quality of *Pecten maximus* sperm. *Proceedings of the 9th International Pectinid Workshop, J Fish Res Board Can.* 1993.
- [42] Wilson-Leedy JG, Ingermann RL. Development of a novel CASA system based on open source software for characterization of zebrafish sperm motility parameters. *Theriogenology* 2007;67:661–672. <https://doi.org/10.1016/j.theriogenology.2006.10.003>
- [43] Boulais M., Soudant P, Le Goïc N, Quéré C, Boudry P, Suquet M. Involvement of mitochondrial activity and OXPHOS in ATP synthesis during the motility phase of spermatozoa in the Pacific oyster, *Crassostrea gigas*. *Biol Reprod* 2015;93:1-7. <https://doi.org/10.1095/biolreprod.115.128538>
- [44] Suquet M, Labbe C, Brizard R, Donval A, Le Coz JR, Quere C, Haffray P. Changes in motility, ATP content, morphology and fertilisation capacity during the movement phase of tetraploid Pacific oyster (*Crassostrea gigas*) sperm. *Theriogenology* 2010;74:111–117. <https://doi.org/10.1016/j.theriogenology.2010.01.021>.
- [45] Dong Q, Eudeline B, Allen SK Jr, Tiersch TR. Factors affecting sperm motility of tetraploid Pacific oysters. *J Shellfish Res* 2002;21:719-723.
- [46] Riesco MF, Félix F, Matias D, Joaquim S, Suquet M, Cabrita E. First study in cryopreserved *Crassostrea angulata* sperm. *Gen Comp Endocrinol* 2017;245:108-115. <https://doi.org/10.1016/j.ygcen.2016.05.003>.
- [47] Lyons L, Jerry DR, Southgate PC. Cryopreservation of black-lip pearl oyster (*Pinctada margaritifera*, l.) spermatozoa: effects of cryoprotectants on spermatozoa motility. *J Shellfish Res* 2005;24:1187-1190. [https://doi.org/10.2983/0730-8000\(2005\)24\[1187:COBPOP\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2005)24[1187:COBPOP]2.0.CO;2)
- [48] Acosta-Salmón H, Jerry DR, Southgate PC. Effects of cryoprotectant agents and freezing protocol on motility of black-lip pearl oyster (*Pinctada margaritifera* L.) spermatozoa. *Cryobiology* 2007;54:13-18. <https://doi.org/10.1016/j.cryobiol.2006.10.189>.
- [49] Kadam AL, Koide SS. Stimulation of *Spisula* sperm motility by 5-hydroxytryptamine analogs. *Invertebrate Reprod Dev* 1990;17:33-37. <https://doi.org/10.1080/07924259.1990.9672085>
- [50] Fujinoki M. Serotonin-enhanced hyperactivation of hamster sperm. *Reproduction* 2011;142:255–266. <https://doi.org/10.1530/REP-11-0074>.

- [51] Jiménez-Trejo F, Tapia-Rodríguez M, Cerbon M, Kuhn DM, Manjarrez-Gutiérrez G, Mendoza-Rodríguez CA, Picazo O. Evidence of 5-HT components in human sperm: implications for protein tyrosine phosphorylation and the physiology of motility. *Reproduction* 2012;144:677–685. <https://doi.org/10.1530/REP-12-0145>.
- [52] Derkach V, Surprenant A, North RA. 5-HT₃ receptors are membrane ion channels. *Nature* 1989;339:706-709. <https://doi.org/10.1038/339706a0>.
- [53] Maricq AV, Peterson AS, Brake AJ, Myers RM, Julius D. Primary structure and functional expression of the 5HT₃ receptor, a serotonin-gated ion channel. *Science* 1991;254:432-437. <https://doi.org/10.1126/science.1718042>
- [54] Hoyer D, Hannon, JP, Martin GR. Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacol Biochem Behav* 2002;71:533-554. [https://doi.org/10.1016/S0091-3057\(01\)00746-8](https://doi.org/10.1016/S0091-3057(01)00746-8).
- [55] Bockaert J, Claeyssen S, Dumuis A, Marin P. Classification and signaling characteristics of 5-HT receptors. In: Müller CP, Jacobs BL, editors. *Handbook of the behavioral neurobiology of serotonin, Handbook of behavioral neuroscience, Vol 21*. Amsterdam: Elsevier; 2010. p. 103-121.
- [56] Tanabe T, Yuan Y, Nakamura S, Itoh N, Takahashi KG, Osada M. The role in spawning of a putative serotonin receptor isolated from the germ and ciliary cells of the gonoduct in the gonad of the Japanese scallop, *Patinopecten yessoensis*. *Gen Comp Endocrinol* 2010;166:620-627. <https://doi.org/10.1016/j.ygcen.2010.01.014>.
- [57] Wang Q, He M. Molecular characterization and analysis of a putative 5-HT receptor involved in reproduction process of the pearl oyster *Pinctada fucata*. *Gen Comp Endocrinol* 2014;204:71-79. <https://doi.org/10.1016/j.ygcen.2014.05.010>.
- [58] Zhang G, Fang X, Guo X, Li L, Luo R, Xu F, Yang P, Zhang L, Wang X, Qi H, Xiong Z, Que H, Xie Y, Holland PW, Paps J, Zhu Y, Wu F, Chen Y, Wang J, Peng C, Meng J, Yang L, Liu J, Wen B, Zhang N, Huang Z, Zhu Q, Feng Y, Mount A, Hedgecock D, Xu Z, Liu Y, Domazet-Lošo T, Du Y, Sun X, Zhang S, Liu B, Cheng P, Jiang X, Li J, Fan D, Wang W, Fu W, Wang T, Wang B, Zhang J, Peng Z, Li Y, Li N, Wang J, Chen M, He Y, Tan F, Song X, Zheng Q, Huang R, Yang H, Du X, Chen L, Yang M, Gaffney PM, Wang S, Luo L, She Z, Ming Y, Huang W, Zhang S, Huang B, Zhang Y, Qu T, Ni P, Miao G, Wang J, Wang Q, Steinberg CE, Wang H, Li N, Qian L, Zhang G, Li Y, Yang H, Liu X, Wang J, Yin Y, Wang J. The oyster genome reveals stress adaptation and complexity of shell formation. *Nature* 2012;490:49-54. <https://doi.org/10.1038/nature11413>.
- [59] Tierney AJ. Structure and function of invertebrate 5-HT receptors: a review. *Comp Biochem Physiol A* 2001;128:791-804. [https://doi.org/10.1016/S1095-6433\(00\)00320-2](https://doi.org/10.1016/S1095-6433(00)00320-2).
- [60] Bandivdekar AH, Segal SJ, Koide SS. Binding of 5-hydroxytryptamine analogs by isolated *Spisula* sperm membrane. *Invertebr Reprod Dev* 1992;21:43-46. <https://doi.org/10.1080/07924259.1992.9672218>.
- [61] Matsutani T, Nomura T. Serotonin-like immunoreactivity in the central nervous system and gonad of the scallop, *Patinopecten yessoensis*. *Cell Tissue Res* 1986;244:515-517.
- [62] Garnerot F, Pellerin J, Blaise C, Mathieu M. Immunohistochemical localization of serotonin (5-hydroxytryptamine) in the gonad and digestive gland of *Mya arenaria* (Mollusca: Bivalvia). *Gen Comp Endocrinol* 2006;149:278-284. <https://doi.org/10.1016/j.ygcen.2006.06.008>.
- [63] Masseau I, Bannon P, Anctil M, Dubé F. Localization and quantification of gonad serotonin during gametogenesis of the Surf clam, *Spisula solidissima*. *Biol Bull* 2002;202:23-33. <https://doi.org/10.2307/1543219>.

- 953 [64] Croll RP, Too CKL, Pani AK, Nason J. Distribution of serotonin in the sea scallop
 954 *Placopecten magellanicus*. Invertebr Reprod Dev 1995;28:125-135.
 955 <https://doi.org/10.1080/07924259.1995.9672473>.
- 956 [65] Paulet YM, Donval A, Bekhadra F. Monoamines and reproduction in *Pecten maximus*, a
 957 preliminary approach. Invertebrate Reproduction and Development. 1993;23:89-94.
 958 <https://doi.org/10.1080/07924259.1993.9672299>.
- 959 [66] Klouche MS, De Deurwaerdère P, Dellu-Hagedorn F, Lakhdar-Ghazal N, Benomar S.
 960 Monoamine content during the reproductivte cycle of *Perna perna* depends on site of
 961 origin on the Atlantic Coast of Morocco. Sci Rep 2015;5:13715.
 962 <https://doi.org/10.1038/srep13715>.
- 963 [67] López-Sánchez JA, Maeda-Martínez AN, Croll RP, Acosta-Salmón H. Monoamine
 964 fluctuations during the reproductive cycle of the Pacific lion's paw scallop *Nodipecten*
 965 *subnodosus*. Comp Biochem Physiol A 2009;154:425-428.
 966 <https://doi.org/10.1016/j.cbpa.2009.07.021>.
- 967 [68] Martínez G, Saleh F, Mettifogo L, Campos E, Inestrosa N. Monoamines and the release
 968 of gametes by the scallop *Argopecten purpuratus*. J Exp Zool 1996;274:365-372.
 969 [https://doi.org/10.1002/\(SICI\)1097-010X\(19960415\)274:6<365::AID-JEZ5>3.0.CO;2-](https://doi.org/10.1002/(SICI)1097-010X(19960415)274:6<365::AID-JEZ5>3.0.CO;2-M)
 970 M.
- 971 [69] Pani AK, Croll RP. Distribution of catecholamines, indoleamines, and their precursors
 972 and metabolites in the scallop, *Placopecten magellanicus* (Bivalvia, Pectinidae). Cell
 973 Mol Neurobiol 1995;15:371-386. <https://doi.org/10.1007/BF02089947>.
- 974 [70] Smith JR. A survey of endogenous dopamine and serotonin in ciliated and nervous
 975 tissues of ve species of marine bivalves, with evidence for speci c, high-a nity dopa-
 976 mine receptors in ciliated tissue of *Mytilus californianus*. Comp Biochem Physiol C.
 977 1982;71:57-61. [https://doi.org/10.1016/0306-4492\(82\)90010-7](https://doi.org/10.1016/0306-4492(82)90010-7).
- 978 [71] Stefano GB, Catapane J, Aiello E. Dopaminergic agents: Infuence on serotonin in the
 979 molluscan nervous system. Science 1976;194:539-541
- 980 [72] Parisi E, De Prisco P, Capasso A, del Prete M. Serotonin and sperm motility. Cell Biol
 981 Int Rep 1984;8:95. [https://doi.org/10.1016/0309-1651\(84\)90075-4](https://doi.org/10.1016/0309-1651(84)90075-4).
- 982 [73] Meizel S, Turner KO. Serotonin or its agonist 5-methoxytryptamine can stimulate
 983 hamster sperm acrosome reactions in a more direct manner than catecholamines. J Exp
 984 Zool 1983;226:171-174. <https://doi.org/10.1002/jez.1402260120>.
- 985 [74] Young RJ, Laing JC. Biogenic amine binding sites in rabbit spermatozoa. Biochem Int
 986 1990;21:781-787.
- 987 [75] Christen RW, Schackmann RW, Shapiro BM. Elevation of intracellular pH activates
 988 respiration and motility of sperm of the sea urchin *Strongylocentrotus purpuratus*. J
 989 Biol Chem 1982;257:14881-14890.
- 990 [76] Ohta H, Kawamoto T, Isowa K, Aoki H, Hayashi M, Narita T, Komaru A. Motility of
 991 spermatozoa obtained from testes of Japanese pearl oyster *Pinctada fucata martensii*.
 992 Fish Sci 2007;73:107-111. <https://doi.org/10.1111/j.1444-2906.2007.01308.x>.
- 993 [77] Boitano S, Omoto CK. Membrane hyperpolarization activates trout sperm without an
 994 increase in intracellular pH. J Cell Sci 1991;98:343-349.
- 995 [78] Nakajima A, Morita M, Takemura A, Kamimura S, Okuno M. Increase in intracellular
 996 pH induces phosphorylation of axonemal proteins for flagellar motility activation in
 997 starfish sperm. J Exp Biol 2005;208:4411-4418. <https://doi.org/10.1242/jeb.01906>.
- 998 [79] Woolsey J, Ingermann RL. Acquisition of the potential for sperm motility in steelhead
 999 (*Oncorhynchus mykiss*): Effect of pH on dynein ATPase. Fish Physiol Biochem
 1000 2003;19:47-56. <https://doi.org/10.1023/B:FISH.0000035896.84954.f6>.

- [80] Cosson J, Billard R, Cibert C, Dreanno C, Suquet M. Ionic factors regulating the motility of fish sperm. In: Gagnon C, editor. The male gamete: from basic to clinical applications. Vienna, IL: Cache Rive Press; 1999, p. 161-86.
- [81] Alavi SMH, Cosson J. Sperm motility in fishes: (II) Effects of ions and osmotic pressure. *Cell Biol Int* 2006;30:1-14. <https://doi.org/10.1016/j.cellbi.2005.06.004>.
- [82] Morisawa M. Adaptation and strategy for fertilization in the sperm of teleost fishes. *J Appl Ichthyol* 2008;24:362-370. <https://doi.org/10.1111/j.1439-0426.2008.01126.x>.
- [83] Alavi SMH, Cosson J, Bondarenko O, Linhart O. Sperm motility in fishes: (III) Diversity of regulatory signals from membrane to the axoneme. *Theriogenology* 2019, In press.
- [84] Vines CA, Yoshida K, Griffin FJ, Pillai MC, Morisawa M, Yanagimachi R, Cherr GN. Motility initiation in herring sperm is regulated by reverse sodium–calcium exchange. *PNAS* 2002;99:2026-2031. <https://doi.org/10.1073/pnas.042700899>.
- [85] Shiba K, Marian T, Krasznai Z, Baba SA, Morisawa M, Yoshida M. Na⁺/Ca²⁺ exchanger modulates the flagellar wave pattern for the regulation of motility activation and chemotaxis in the ascidian spermatozoa. *Cell Motil Cytoskeleton* 2006;63:623-632. <https://doi.org/10.1002/cm.20149>.
- [86] Blaustein MP, Lederer WJ. Sodium/calcium exchange: its physiological implications. *Physiol Rev* 1999;79:763-854. <https://doi.org/10.1152/physrev.1999.79.3.763>.
- [87] Costello S, Michelangeli F, Nash K, Lefievre L, Morris J, Machado-Oliveira G, Barratt C, Kirkman-Brown J, Publicover S. Ca²⁺-stores in sperm: their identities and functions. *Reproduction* 2009;138:425-37. <https://doi.org/10.1530/REP-09-0134>.
- [88] Correia J, Michelangeli F, Publicover S. Regulation and roles of Ca²⁺ stores in human sperm. *Reproduction* 2015;150:65-76. <https://doi.org/10.1530/REP-15-0102>.
- [89] Kuroda Y, Kaneko S, Yoshimura Y, Nozawa S, Mikoshiba K. Are there inositol 1,4,5-triphosphate (IP3) receptors in human sperm? *Life Sciences* 1999;65:135–143. [https://doi.org/10.1016/S0024-3205\(99\)00230-1](https://doi.org/10.1016/S0024-3205(99)00230-1).
- [90] Ho HC, Suarez SS. Characterization of the intracellular calcium store at the base of the sperm flagellum that regulates hyperactivated motility. *Biol Reprod* 2003;68:1590-1596. <https://doi.org/10.1095/biolreprod.102.011320>.
- [91] Naaby-Hansen S, Wolkowicz MJ, Klotz K, Bush LA, Westbrook VA, Shibahara H, Shetty J, Coonrod SA, Reddi PP, Shannon J et al. Co-localization of the inositol 1,4,5-trisphosphate receptor and calreticulin in the equatorial segment and in membrane bounded vesicles in the cytoplasmic droplet of human spermatozoa. *Mol Human Reprod* 2001;7:923–933. <https://doi.org/10.1093/molehr/7.10.923>.
- [92] Sanchez D, Labarca P, Darszon A. Sea urchin sperm cation-selective channels directly modulated by cAMP. *FEBS Lett* 2001;503:111-115. [https://doi.org/10.1016/S0014-5793\(01\)02713-2](https://doi.org/10.1016/S0014-5793(01)02713-2).
- [93] Galindo BEC, Beltran C, Cragoe Jr EJ, Darszon A. Participation of a K⁺ channel modulated directly by cGMP in the speract-induced signaling cascade of *Strongylocentrotus purpuratus* sea urchin sperm. *Dev Biol* 200;221:285-294. <https://doi.org/10.1006/dbio.2000.9678>.
- [94] Howard DR, Trantow CM, Thaler CD. Motility of a biflagellate sperm: waveform analysis and cyclic nucleotide activation. *Cell Motil Cytoskeleton* 2004;259:120-130. <https://doi.org/10.1002/cm.20027>.
- [95] Komaru A, Konishi K. Ultrastructure of biflagellate spermatozoa in the freshwater clam, *Corbicula leana* (Prime). *Invert Reprod Dev* 1996;29:193–197. <https://doi.org/10.1080/07924259.1996.9672513>.

- [96] Konishi K, Kawamura K, Furuita H, Komaru A. Spermatogenesis of the freshwater clam *Corbicula aff. fluminea* Mueller (Bivalvia: Corbiculidae). J Shellfish Res 1998;17:185-189.
- [97] Kraemer LR, Swanson C, Galloway M, Kraemer R. Biological basis of behavior in *Corbicula fluminea*. II. Functional morphology of reproduction and development and review of evidence for self-fertilization. In: Proceedings of the Second International *Corbicula* Symposium. Am Malacol Bull 1986;193-201.
- [98] Cook SP, Babcock DF. Selective modulation by cGMP of the K channel activated by speract. J Biol Chem 1993;268:22402–22407.
- [99] Faure C. Paramètres physiologiques de l'émission et de l'activation des gamètes mâles de deux mollusques bivalves: la coquille Saint Jacques *Pecten maximus* (L.) et l'huître creuse *Crassostrea gigas* (Thunberg). PhD thesis, University of Paris 7, 1996.
- [100] Cosson J. A moving image of flagella: news and views on the mechanisms involved in axonemal beating. Cell Biol Int 1996;20:83–94. <http://dx.doi.org/10.1006/cbir.1996.0012>.
- [101] Cosson J, Dreanno C, Fauvel C, Groison AL, Suquet M, Billard R. Marine fish spermatozoa: racing ephemeral swimmers. Reproduction 2008;136:277-294. <https://doi.org/10.1530/REP-07-0522>.
- [102] Suquet M, Quere C, Mingant C, Lebrun L, Ratiskol D, Miner P, Cosson J. Effect of sampling location, release technique and time after activation on the movement characteristics of scallop (*Pecten maximus*) sperm. Aquat Living Resour 2013;26:215–220. <https://doi.org/10.1051/alr/2013048>.
- [103] Boulais M, Soudant P, Le Goïc N, Quéré C, Boudry P, Suquet M. ATP content and viability of spermatozoa drive variability of fertilization success in the Pacific oyster (*Crassostrea gigas*). Aquaculture 2017;479:114-119. <https://doi.org/10.1016/j.aquaculture.2017.05.035>.
- [104] Mita M, Nakamura M. Phosphatidylcholine is an endogenous substrate for energy metabolism in spermatozoa of sea urchins of the order Echinoidea. Zoolog Sci 1993;10:73–83.
- [105] Mita M, Nakamura M. Energy metabolism of sea urchin spermatozoa: an approach based on echinoid phylogeny. Zoolog Sci 1998;15:1–10. <https://doi.org/10.2108/zsj.15.1>.
- [106] Mita M, Oguchi A, Kikuyama S, Desantis R, Nakamura M. Ultrastructural study of endogenous energy substrates in spermatozoa of the sea urchin urchins *Arbacia lixula* and *Paracentrotus lividus*. Zoolog Sci 1994;11: 701–705.
- [107] Mohri H. Endogenous substrates of respiration in sea-urchin spermatozoa. J Fac Sci Univ Tokyo, sec IV 1957;8:51–63.
- [108] Ellington W. Evolution and physiological roles of phosphagen systems. Annu Rev Physiol 2001;63:289–325. <https://doi.org/10.1146/annurev.physiol.63.1.289>.
- [109] Suquet M, Cosson J, Donval A, Labbé C, Bernard I, Fauvel C. Marathon versus sprint racers: an adaptation of sperm characteristics to the reproductive strategy of Pacific oyster, turbot and seabass. J Appl Ichthyol 2012;28:956–960. <https://doi.org/10.1111/jai.12061>.
- [110] Troost K. Causes and effects of a highly successful marine invasion: case study of the introduced Pacific oyster *Crassostrea gigas* in continental NW European estuaries. J Sea Res 2010;64:145–165. <https://doi.org/10.1016/j.seares.2010.02.004>.
- [111] Suquet M, Malo F, Quéau I, Ratiskol D, Quéré C, Le Grand J, Fauvel C. Seasonal variation of sperm quality in Pacific oyster (*Crassostrea gigas*). Aquaculture 2016;464:638-641. <https://doi.org/10.1016/j.aquaculture.2016.07.016>.

1098 **Table 1.** Morphology and motility parameters of spermatozoa in most studied bivalve
 1099 species.

Species	Black-lip pearl oyster	Pacific oyster	Japanese scallop	Manila clam
Reference	30, 31	22, 23, 39, 43, 44, 103, 111	39, 36	39, 34, 35
Head length (μm)		2-3	4	6
Head width (μm)		2-3		
Number of mitochondria	4	4	4	4
Flagellar length (μm)	47	37-41	50	42-45
Motility in seawater (%)	0	55-73	78	43
Motility in DCSB ₄ (%)	85	77	85	95
VCL - SW ($\mu\text{m s}^{-1}$)			255	100
VSL - SW ($\mu\text{m s}^{-1}$)	67			
VAP - SW ($\mu\text{m s}^{-1}$)	166	60-70		
Waves velocity ($\mu\text{m s}^{-1}$)	403			
Beat frequency (Hz)	49			
Wave amplitude (μm)	5			
Wave length (μm)	16			
Local curvature or bend angle	From -69 to +65			
Number of curvatures	3			
Diameter of circular head tracks (μm)	347			
ATP content (nmol 10^{-8} spz)	4-70	5-19		

1100
 1101 DCSB₄ – saline solution composed of 19.5 g L⁻¹ NaCl, 6.25 g L⁻¹ glycine, 0.15 g L⁻¹ CaCl₂,
 1102 0.19 g L⁻¹ MgSO₄, 2.42 g L⁻¹ Tris-aminomethane, pH 8.2, SW – seawater, VAP – velocity of
 1103 the average path, VCL – curvilinear velocity, VSL – straight-line velocity. Black-lip pearl
 1104 oyster, *Pinctada margaritifera*; Japanese scallop, *Patinopecten yessoensis*; Manila clam,
 1105 *Ruditapes philippinarum*; Pacific oyster, *Crassostrea gigas*

1106

1107 **Table 2.** pH and ionic constituents in testicular fluid (TF) and blood plasma (BP) of bivalves
 1108 compared to seawater.

		Na ⁺ mM	K ⁺ mM	Ca ²⁺ mM	pH	Osmolality mOsmol kg ⁻¹	Reference
Artificial seawater		469	11	10	8.2	1000	39
Pacific oyster	TF	360	18	11	5.5 - 6.5		39
	BP	418	20	14	ND		39
	TF	427	32	9	5.82	1061	37
Japanese scallop	TF	351	41	7	7.0		39
	BP	523	11	9	ND		39
European flat oyster	TF				6.31		33
Great scallop	TF				6.5		40
Black-lip pearl oyster	TF				6.6		30

1109 ND – not determined. Black-lip pearl oyster, *Pinctada margaritifera*; European flat oyster,
 1110 *Ostrea edulis*; great scallop, *Pecten maximus*; Japanese scallop, *Patinopecten yessoensis*;
 1111 Pacific oyster, *Crassostrea gigas*

Figure legends

Fig. 1. Morphology of sexually ripe Pacific oyster, *Crassostrea gigas*, after upper shell removal. The gonad (G) is the white part of the animal and sperm (s) is released through the gonopore (g).

Fig. 2. Morphology of spermatozoon in bivalves. High speed video frame of a motile spermatozoon of the Pacific oyster, *Crassostrea gigas*, in artificial seawater containing 10^{-5} M of 5-Hydroxytryptamine creatinine sulfate (5-HT), a – acrosome, h – head, m – midpiece, f – flagellum.

Fig. 3. Current known signaling pathway of spermatozoa motility in bivalves. During spawning, spermatozoa motility is triggered by a Na^+ -dependent alkalization of internal pH mediated by a Na^+/H^+ exchanger. In addition, decrease of extracellular K^+ or presence of extracellular 5-hydroxytryptamine creatinine sulfate (5-HT) induces hyperpolarization of spermatozoa membrane potential (E_m), mediated by K^+ efflux via voltage-dependent K^+ channels and 5-HT influx, associated with opening of voltage-dependent Ca^{2+} channels under alkaline condition. Subsequent increase in intracellular Ca^{2+} content ($[\text{Ca}^{2+}]_i$) triggers Ca^{2+} /calmodulin (CaM)-dependent flagellar beating. Release of Ca^{2+} from intracellular stores may also contribute to increase $[\text{Ca}^{2+}]_i$. Na^+ influx probably regulates Ca^{2+} exchange via $\text{Na}^+/\text{Ca}^{2+}$ exchanger. The mechanism through which 5-HT induces spermatozoa hypermotility is unclear and it may be possible that 5-HT is involved in controlling intracellular cAMP rise in bivalve spermatozoa, leading to cAMP-dependent protein kinase (PKA) phosphorylation in the flagellum.

Fig. 4. Successive frames of flagellar beating of a Manila clam, *Ruditapes philippinarum*, spermatozoon. Motility of spermatozoa was recorded under a phase contrast microscope (Olympus BX51, 20X magnification) using a high-speed CDD camera (HAS-220; Ditect) with an acquisition rate of 200 frames per second. Some parts of the flagellum are in focus while others are slightly out of focus illustrating the 3D distortion of the flagellar plane of beating. To activate motility, sperm was diluted 2000 folds in artificial seawater containing 10^{-5} M 5-hydroxytryptamine creatinine sulfate (5-HT) and BSA (0.1% w : v) was added to prevent sperm adhering to the glass slide, no cover slip was used.

Fig. 5. Head trajectories in motile spermatozoa of the Manila clam, *Ruditapes philippinarum*. Motility of spermatozoa was activated in artificial seawater or artificial seawater containing 10^{-5} M 5-hydroxytryptamine creatinine sulfate (5-HT) at 5 min, 30, 96, and 144 h post-activation, and recorded under a phase-contrast microscope (Olympus BX51) connected to a digital CCD camera (Hamamatsu Photonics) with an acquisition rate of 50 frames per second. Spermatozoa movement trajectories were analyzed using semen motility analysis system (SMAS). To activate spermatozoa motility, sperm was diluted 500 to 1000 folds in artificial seawater with or without 5-HT. BSA (0.1% w : v) was added to prevent sperm adhering to the glass slide and no cover slip was used.

Fig. 6. Successive frames of flagellar beating of a Pacific oyster, *Crassostrea gigas*, spermatozoon in seawater. Spermatozoa motility was activated in seawater and observed using a phase contrast microscope (dark-field, Olympus BX51), connected to a high-speed camera at 1000 images s^{-1} (interval between two frames is 1 millisecond). Some parts of the flagellum are in focus while others are slightly out of focus, illustrating the 3D distortion of the flagellar plane of beating. To activate spermatozoa motility, sperm was diluted 500 folds

in seawater and pluronic acid (0.1% w : v) was added to prevent sperm adhering to the glass slide, no cover slip was used.

Fig. 7. Successive frames of flagellar beating of a Pacific oyster, *Crassostrea gigas*, spermatozoon in seawater at 1 mM 5-HT. Spermatozoa movement was observed using a phase contrast microscope (dark-field, Olympus BX51) supplied by stroboscopic illumination, and connected to a video camera at 50 frames s⁻¹ at 150 Hz (interval between two frames is 20 millisecond). A: Each frame shows one position of the same spermatozoon B: Each frame includes three successive positions of the same spermatozoon. To activate spermatozoa motility, sperm was diluted 500 folds in seawater and pluronic acid (0.1% w : v) was added to prevent sperm adhering to the glass slide, no cover slip was used.


Fig. 8. Successive frames of “twitching” behavior in a Pacific oyster, *Crassostrea gigas*, spermatozoon. Spermatozoon shows only one local bending point along the flagellum, and thus no flagellar sinusoidal waves. Spermatozoa motility was observed at the beginning of movement activation, prior to full activation of spermatozoa motility in seawater, using a phase contrast microscope (dark-field, Olympus BX51) supplied by stroboscopic illumination, and connected to a video camera at 50 frames s⁻¹ (interval between two frames is 20 millisecond) at 150 Hz. Each frame includes three successive positions of the same spermatozoon. To activate spermatozoa motility, sperm was diluted 500 folds in seawater and pluronic acid (0.1% w : v) was added to prevent sperm adhering to the glass slide, no cover slip was used.


Fig. 9. Successive frames of flagellar beating of a Pacific oyster, *Crassostrea gigas*, spermatozoon in viscous seawater. Spermatozoa motility was activated in seawater


1187 containing methyl cellulose (2% w : v). Waves smoothly develop and progress from the head
1188 junction to the tip of the flagellum, but wave beat frequency is about five times lower than in
1189 seawater and wave shape is highly affected. Progressiveness of spermatozoon is highly
1190 reduced in viscous seawater. Spermatozoa motility was observed using a phase contrast
1191 microscope (dark-field, Olympus BX51) supplied by stroboscopic illumination, and
1192 connected to a video camera at 50 frames s⁻¹ (interval between two frames is 20 millisecond)
1193 at 150 Hz. To activate spermatozoa motility, sperm was diluted 500 folds in seawater and
1194 pluronic acid (0.1% w : v) was added to prevent sperm adhering to the glass slide, no cover
1195 slip was used.


1196

1197 **Video 1.** Available at <https://vimeo.com/312367037>. Once motility is triggered, flagellum
1198 beats in asymmetric wave pattern leading to circular trajectories of spermatozoa.
1199 Spermatozoa movement of several mollusk species was recorded while swimming. In oyster
1200 spermatozoa, flagellar waves are not fully co-planar but rather 3D, while in scallops,
1201 flagellum remains in a single plan.

















Seawater




Serotonin (5-HT)





A



B

