



HAL
open science

Variability of key biological parameters of round sardinella *Sardinella aurita* and the effects of environmental changes

Bocar S. Balde, Fambaye N. Sow, Kamarel Ba, Werner Ekau, Patrice Brehmer, Justin Kantoussan, Massal Fall, Malick Diouf

► To cite this version:

Bocar S. Balde, Fambaye N. Sow, Kamarel Ba, Werner Ekau, Patrice Brehmer, et al.. Variability of key biological parameters of round sardinella *Sardinella aurita* and the effects of environmental changes. *Journal of Fish Biology*, 2019, 94 (3), pp.391-401. 10.1111/jfb.13903 . hal-02749018

HAL Id: hal-02749018

<https://hal.science/hal-02749018v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Variability of key biological parameters of round sardinella *Sardinella aurita* and the effects of environmental changes

Baldé Bocar Sabaly ^{1,2,3,4,*}, Sow Fambaye N. ¹, Ba Kamarel ^{1,4}, Ekau Werner ², Brehmer Patrice ^{1,3}, Kantoussan Justin ⁵, Fall Massal ¹, Diouf Malick ⁴

¹ ISRA, CRODT, PRH, Dakar, Senegal.

² Leibniz Ctr Trop Marine Ecol ZMT, Dept Ecol, Bremen, Germany.

³ Ifremer, Inst Rech Dev France IRD, UMR Lemar, CNRS,UBO,IRD, Dakar, Senegal.

⁴ Univ Cheikh Anta Diop, IUPA, UCAD II, Dakar, Senegal.

⁵ Univ Gaston Berger, Unite Format & Rech Sci Agronom Aquaculture & Tec, St Louis, Senegal.

* Corresponding author : Bocar Sabaly Baldé, email address : bocarbalde2005@hotmail.com

Abstract :

We examined growth rates and reproductive characteristics of *Sardinella aurita* off Senegal and other coastal areas over a 20 year period (1995-2014) to determine how they relate to variations in environmental characteristics of coastal waters. Based on fish length-frequency data and a coastal upwelling index, we found that *S. aurita* recruitment tends to occur during the periods of most intensive upwelling (March-April off Senegal). Peak reproduction corresponds to periods of low sea-surface temperature (in February or March). The sex ratio was remarkably consistent during the 30 year study period and so was not affected by environmental changes. We hypothesise that *S. aurita* takes advantage of the higher zooplankton productivity that occurs in coastal waters when upwelling brings nutrient-rich water to the surface (i.e., it increases its growth rate and accumulates energy reserves for spawning). Growth performance appears to be strongly dependent on environmental conditions. The timing of spawning seems to occur when food (zooplankton) is most available for supplying the energy requirements needed by adults for spawning and early development of larvae. Environmental changes seem to have a significant effect on *S. aurita* growth and reproduction, which endorses their high phenotypic plasticity.

Keywords : data poor fisheries, environmental variability, size structure, small pelagic, spawning period, west Africa

31

1. INTRODUCTION

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

The Canary Current off the West Africa coast initiates an upwelling zone that modifies the trophic structure of coastal ecosystems profoundly (Auger et al., 2016). The coastal upwelling region from 10° N to 35° N has been identified as one of many large marine ecosystems (“LMEs”) of the world, with the southern part of this LME represented by the Senegal-Mauritanian upwelling system (Auger et al., 2016). In the more southern portion of this system off Senegal (between 12° N and 17° N, upwelling occurs only during the winter-spring period. This upwelling is a response to the southern extension of the trade wind, which is defined by the position of the Intertropical Convergence Zone (Auger et al., 2016). The wind-forced upwelling of deep, nutrient-rich water is responsible for high primary productivity (of phytoplankton) in coastal waters, which provides the energy (food) for higher trophic levels. In the Senegal-Mauritanian region (10°–19° N), the geographic expansion of offshore phytoplankton production that occurs during winter and spring is followed by an abrupt decline in summer production (Auger et al., 2016). Bakun (1990) hypothesized that climate change may lead to an intensification of coastal upwelling in response to increasing favourable upwelling wind, and Barton et al. (2013) has predicted there is an increase in SST in general. In either case, changes in the upwelling intensity of the Senegal-Mauritanian system will influence the abundance of small pelagic fish, including *Sardinella aurita* (round Sardinella) and *Sardinella maderensis* (flat Sardinella) (Thiaw et al., 2017; Diankha et al., 2018). Thiaw et al. (2017) and Diankha et al. (2018) report that abundance and recruitment success of *S. aurita* are associated with distinct ranges of sea surface temperature, upwelling intensity, wind-induced turbulence, concentration of chlorophyll-a and north Atlantic oscillation index in Senegal.

The species *S. aurita* inhabits tropical and subtropical regions of the eastern Atlantic Ocean and occurs in high concentrations along the west coast of Africa in three main areas (Roy et al., 1989): between southern Morocco (Western Sahara) and Guinea (10°–26° N), between Ivory Coast and Ghana (7° – 5 °N) and further south between Gabon and southern Angola (0°–18° S) (Froese & Pauly, 2016). *S. aurita* also occurs in the western Atlantic Ocean from Cape Cod (USA) to Argentina, including the Caribbean, West Indies, and Gulf of Mexico (Felder, 2009). It also inhabits the Mediterranean Sea (Boltachev & Karpova, 2014). In all regions, the species preferentially inhabits continental shelves where it prefers saline (between 34 ppt and 36.5 ppt), and warm sea temperatures (between 17° C and 30 °C) (Boëly, 1979). In Senegal, massive reproduction events occur south of the Cap-Vert peninsula (Dakar) to the delta of Saloum at the border with The Gambia (Conand, 1977; Boëly,

59 1982). In terms of exploitation, *S. aurita* has already been intensively exploited by artisanal fleets in Senegal
60 (Thiao, 2009) and in Mauritania by foreign fishing fleets under fisheries agreements (e.g. Union European and
61 Russian Federation) and Morocco with a small artisanal fleet fishing near the coast (Isselmou & Baye, 2009). Due
62 to the increasing demand for *S. aurita* for human consumption, the fishing effort of artisanal fleets in Senegal and
63 Mauritania has steadily increased in recent years.

64 Growth characteristics and breeding season of *S. aurita* populations vary among areas. For example, in the eastern
65 Atlantic (e.g., off Senegal), *S. aurita* possesses an asymptotic length of 36.5 cm total length (TL) (Boëly et al.,
66 1982) and a breeding season that extends from February to May and from October to November (Boëly, 1982). In
67 western Atlantic Ocean (off Venezuela), the species possesses an asymptotic length of 32.6 cm (TL) and a
68 breeding period from January to March and from October to December (Gassman et al., 2008). In the
69 Mediterranean (off Greece), *S. aurita* reaches an asymptotic length of 21.4 cm (TL) (Tsikliras et al., 2005) and a
70 rather limited breeding period between May and June (Tsikliras & Antonopoulou, 2006). Biological
71 characteristics of *S. aurita*, such as growth and reproduction, have been shown to be strongly influenced by
72 environmental conditions (e.g., upwelling intensity and sea surface temperature) in both the Mediterranean
73 (Tsikliras et al., 2005; Tsikliras & Antonopoulou, 2006) and the West Atlantic (Fréon et al. 1997). Such findings
74 suggest that growth and reproductive periods of *S. aurita* in the eastern Atlantic (off Senegal) might also be
75 influenced by upwelling intensity and sea surface temperature.

76 To determine how environmental conditions might be affecting *S. aurita* populations and their biology, we
77 analysed landing data available for coastal Senegal from 1995 to 2014. The data collections included reproductive
78 condition (age of sexual maturity, length-frequency distributions, and condition factor) and growth parameters.
79 The results of this study were compared and discussed in relation to relevant past studies in various north tropical
80 Atlantic (Eastern and Western) and Mediterranean Sea areas.

2. MATERIAL AND METHODS

2.1. Biological data

Data on the growth of *S. aurita* were obtained from fish that landed at the seven main artisanal fishing harbours along the Senegal coast from July 2013 to June 2014 (Fig. 1). Three harbours (Kayar, Saint-Louis and Yoff) are located in the northern part of the coast (from 16° 04' N to 14° 36' N) where the continental shelf is narrow. The other harbours (Hann, Mbour, Joal and Kafountine) are located along the southern coast (from 14° 36' N to 13° 36' N and from 13° 04' N to 12 °20' N) where the continental shelf is wider than the northern part. Length frequency data were collected randomly at about 5 days per week. This study sampled only specimens caught using purse seine nets to have the same fishing gear selectivity for monthly comparisons of the length distribution.

Total length (TL in cm) of fish sampled (n = 32 346 individuals) was measured to the nearest 1 mm, while the body weight (W) was determined to the nearest 0.01 g. Sampled individuals were grouped into 1-cm-interval size classes to calculate size-frequency distributions and to estimate growth parameters.

Length-weight relationship (Le Cren, 1951) was determined for both males and females:

$$W = a \times TL^b \quad (1)$$

Where W is the weight and TL is the Total Length, and a and b allometric coefficients.

The data we used for our study of *S. aurita* reproduction were collected by other surveys of the artisanal fisheries along the southern Senegal coast from 1995 to 2011, including surveys by the Fishery Research Vessel (FRV) Fridtjof Nansen (Charouki et al., 2011) and by the Centre de Recherches Océanographiques de Dakar-Thiaroye (CRODT in French) covering Senegal (i.e., Senegal Fisheries Research Centre). These fish surveys were conducted regularly from November through December with a few surveys covering the June-to-July period. In Senegal, massive reproduction events occur south of the Cap-Vert peninsula (Dakar) to the delta of Saloum at the border with The Gambia (Conand, 1977; Boëly, 1982) (Fig.1). A total of 4397 fish samples were recorded during the surveys (Table I). The FRV Fridtjof Nansen stopped collecting samples in 2006, at the end of its research project, but technicians involved in the surveys from the CRODT continued to collect seasonal samples until 2011. Weight of gonads (0.1 g) and sex were determined for 4397 fish. Stages of maturity were determined macroscopically using the Fontana (1969) maturity scale (Table II).

108 The von Bertalanffy (1938) growth parameters [coefficient of growth rate (K), asymptotic length (L_{∞}), time “t”
 109 when the fish length is zero (t_0)] were estimated from monthly length-frequency data using the ‘TropFishR’
 110 (Tropical Fisheries Analysis with R) package (Mildenberger et al., 2017), while natural mortality was estimated
 111 using the McQuinn et al. (1990) method:

$$112 \quad M = 3/t_{max} \quad (2)$$

113 Where t_{max} is the longevity of the species that is by definition the age corresponding to the size average maximum
 114 ($L_{max} = 95\% L_{\infty}$) that a fish can reach and estimated by the following equation of Pauly (1983):

$$115 \quad t_{max} = (3/K) + t_0 \quad (3)$$

116 The growth performance index (ϕ') was used to compare growth rates among *S. aurita* and evaluate growth
 117 performance potential under different environmental constraints (Tsikliras et al., 2005). Growth parameters of *S.*
 118 *aurita* from the different Mediterranean, Central Atlantic stocks were used to construct an auximetric double
 119 logarithmic plot of an individual’s coefficient of growth rate (K) against an estimate of its asymptotic length (L_{∞}).
 120 The spawning period of *S. aurita* was determined using two approaches: (1) a qualitative approach based on
 121 monitoring monthly fluctuations in the percentage of different stages of gonad development and (2) a quantitative
 122 approach based on the monthly change in the gonado-somatic index (GSI; in %). GSI was calculated for each
 123 sampled individual based on Le Bec, (1983):

$$124 \quad GSI = \frac{W_g}{W_{ev}} \times 100 \quad (4)$$

125 Where W_g is gonad weight and W_{ev} is eviscerated fish weight.

126 We used the equation proposed by Le Cren (1951) to calculate each individual’s condition factor (Kn):

$$127 \quad Kn = \frac{W_T}{W_{th}} \quad (5)$$

128 W_T : total individual fish weight and W_{th} : the theoretical total weight calculated from Eq. 1. This relationship (Eq.
 129 5) indicates that the bigger the fish, the better the environment. The study of this factor makes it possible to see
 130 which cycle, seasonal and annual, obeys weight gain of individuals.

131 Mean length at first sexual maturity (L_{50} ; in cm) was estimated using only fish collected during reproductive
 132 periods (February to June and October to November) (Boëly, 1982; Cury & Fontana, 1988) from 1995 to 2011. A
 133 logistic function with an inflection point at 0.5 ($L_{50\%}$) was used to model the relationship between the percent of
 134 mature females ($M \geq$ Stage III) and total length TL. The following equation was used, where b is a constant of the
 135 model:

$$136 \quad M = \frac{1}{1+e^{-b(TL_t-L_{50\%})}} \times 100 \quad (6)$$

137 For comparison with the results of this study, growth and reproductive parameters from various north tropical
138 Atlantic (Eastern and Western) and Mediterranean areas are shown in Appendix, as well as the data sources.

139 **2.2. Landing data**

140 The data set was total landing of *S. aurita* by the Senegal artisanal fishery from 1995 to 2014 by the type of
141 fishing gear used (purse seine) and landing port (Kayar, Saint-Louis and Yoff, Hann, Mbour, Joal, and
142 Kafountine). These data set was extracted from the CRODT database (Figure 1). The number of trips per fishing
143 gear was recorded on a daily basis, whereas landing data were collected randomly at about 5 days per week. After
144 sorting the data by port, gear type and period (fortnightly intervals), total landings per port were estimated by
145 multiplying the mean number of fish landed for the sampled trips by the total number of fishing trips made
146 (Chaboud et al., 2015). The landings were summed by month for each landing port (Chaboud et al., 2015).
147 Growth (July 2013 to June 2014) and reproduction (1995 to 2011) data by size, sex, maturity stages and landing
148 sites were extrapolated with landing data from 1995 to 2014..

149 **2.3. Environmental data**

150 We used local environmental indices [Coastal Upwelling Index (CUI) and coastal sea surface temperature (SST)]
151 for the period from 1995 to 2011 to analyse the effect of the environmental conditions on seasonal and inter-
152 annual variability of key biological parameters (mainly reproduction) of *S. aurita*. Both data sets were obtained
153 from the Southern Senegal coast (Fig. 1). The SST (°C) data (averaged on a monthly basis) were obtained from
154 advanced, very-high-resolution radiometer (AVHRR) satellite images (5 km resolution) from Southern Senegal
155 coast to the 200 m isobath, where upwelling-induced SST anomalies are generally maximal (Thiaw et al., 2017).
156 The CUI ($\text{m}^3 \text{s}^{-1} \text{m}^{-1}$) was deduced from wind speed data obtained from the U.S. NOAA Environmental Research
157 Division website (ERD, Upwelling and Environmental Index Products, [http://www.pfeg.noaa.gov: Upwelling and
158 Environmental Index Products, last visited: 20/11/2017](http://www.pfeg.noaa.gov: Upwelling and Environmental Index Products, last visited: 20/11/2017)). In order to calculate the CUI in the Southern Senegal,
159 we used the method of Ekman's theory (Ekman, 1905) of the transportation of masses of surface water by wind in
160 the north or north-east direction, coupled with the rotation of the earth:

$$161 \quad CUI = \frac{\tau}{\rho \omega f} \quad (7)$$

162 Where τ is the along shore component of wind stress within 270 km of the coastline (positive southward), ρ is the
163 seawater density (1025 kg m^{-3}) and f is the Coriolis parameter ($= 2 \Omega \sin(\theta)$, with Ω and θ equal to the Earth's
164 angular velocity and latitude, respectively).

165 **2.4. Statistical analysis**

166 All fork length (FL) measurements were converted to total length (TL) according to the relationship (Sylla et al.,
167 2016):

$$168 \quad FL \text{ (mm)} = 10.8264 + 0.8066TL \text{ (mm)} \quad \leftrightarrow \quad TL = (FL - 10.8264)/0.8066 \quad (8)$$

169 All statistical analyses were performed using the “stats”, “mgcv” and “Hmisc” R packages (De Mendiburu, 2014;
170 Harrell Jr, 2017; Wood & Wood, 2017), with a significance level of $\alpha < 0.05$.

171 Condition factor (Kn) was assessed over annual and interannual temporal scales to test whether environmental
172 parameters affected *S. aurita* reproduction. The Generalized Additive Model (GAM) (Hastie & Tibshirani, 1986;
173 Diankha et al., 2018) was used: (i) to quantify the contribution of the environmental variables (CUI, SST); and (ii)
174 to investigate ranges of each environmental variable associated with condition factor. This model does not require
175 any prior assumption on the functional link between each variable and the condition factor. The explanatory
176 variables included in the model were: SST, CUI. We used the smooth function "s", estimated from the data. This
177 function is used to define smooth terms in the GAM model formulas. The function does not evaluate a smooth
178 (spline), but exists only to help configure a model using spline-based smoothing (Wood & Wood, 2017). Multi-
179 collinearity of explanatory variables was tested using the variance inflation factor (VIF) on SST, CUI (Tiedemann
180 & Brehmer, 2017). The variance inflation factors (VIFs) were calculated for all environmental variables to detect
181 possible high dimensional collinearities (Zuur et al., 2010; Tiedemann & Brehmer, 2017; Diankha et al., 2018). A
182 VIF < 2 for all tested explanatory variables was considered adequate (Zuur et al., 2010). Because recruitment was
183 estimated by season [to focus on species biology (the seasonal influence on fecundity)] (Boëly, 1982), season was
184 the temporal unit used to evaluate the effect of environmental variables. The seasonal and interannual means of
185 these three variables were also computed.

$$186 \quad \text{Model} < - \text{gam}(\text{Kn} \sim s(\text{CUI}) + s(\text{SST}), \text{data} = \text{Input data}) \quad (9)$$

187 The χ^2 -test was used to compare sex ratios and proportion of individual in each stage, which made it possible to
188 determine whether the variables observed for the Petite côte and the Northern part of *S. aurita* were independent.

189 The Petite côte (from $14^\circ 36' \text{ N}$ to $13^\circ 36' \text{ N}$) extends from the south of the peninsula of Cap Vert (Dakar) to the

190 delta of Saloum at the border with The Gambia, and includes a 130 km long and 10 km wide continental shelf
191 (Fig. 1). We used analyses of variance (ANOVAs) to compare reproductive parameters between males and
192 females and the growth performance index (ϕ') of different oceanic regions (Mediterranean, Western and Eastern
193 Central Atlantic), followed by Kruskal-Wallis tests. The growth performance index (ϕ') was used to compare
194 growth rates and to assess the potential for growth under various environmental stresses.

195 **2.5. Ethics statement**

196 No specific authorization was required for any activities undertaken during this work at any of the sites visited.
197 The study was conducted in the waters of the Senegalese State. The activities carried out were in collaboration
198 with the Local Fishing Committee (CLP in French) and the local fishermen. Threatened or protected species have
199 not been involved in any part of the field studies.. No approval was required from the Institutional Animal Care
200 and Use Committee, and no field license was required during any part of the experiment.

201

3. RESULTS

3.1. Size spectra and length-weight relationships

The length-frequency distributions between coastal areas (northern and southern part) showed that *S. aurita* off the northern part are larger (mode: 32 ± 2 cm) than fish caught off the southern coast (two modes: 13 ± 1 and 24 ± 3 cm) (Fig. 2). The size distribution of the southern population exhibited a higher peak at a smaller size (24 cm) relative to the northern individuals (peak at 32 cm).

The relationship between the total weight and total length was allometric [$a = 0.008 \pm 0.001$; $b = 3.02 \pm 0.01$, and $r = 0.93$ ($n = 4397$)]. Fig. 3 indicates fish growth in length to be faster than gaining weight when it is young and the older the fish the less the fish is growing in length but is gaining in weight. Only in a very short period, it seems that the fish is relatively seen gaining weight and length at the same rate.

3.2. Temporal variability of condition factor and environmental parameters

The condition factor (Kn) varied seasonally ($X^2 = 22.01$, $df = 11$, $p = 0.001$) with maximum values in May and September (Fig. 4a). The increase in Kn from February to May corresponds to a period of high upwelling intensity (CUI) and low sea surface temperature (SST) along the “Petite côte” (Fig.1) in March–April. Thereafter, Kn declined until August as the CUI declined. A maximum Kn in September was again followed by a decline in CUI and an accelerated decline in SST. The seasonal cycle of declining SST coincides with an increase in CUI because strong offshore, wind-induced upwelling in winter brings cold water to the surface. The relationship between Kn and CUI and SST showed wider amplitude on an annual basis than did seasonal variations (0.88 to 1.14, Fig. 4b). The condition factor for *S. aurita* remained fairly stable over the study 17-year period, except for two peaks, one in 1998 (1.2) and the other in 2003 (1.1). Over the period 1995–2011, SST and CUI varied inter-annually. Mean sea surface temperatures increased in 1998, 2001, 2005, 2008 and 2010, but decreased in 1999 and 2009. The most striking feature during this period was the slight drop in the upwelling index in 1997 at $0.84 \text{ m}^3 \text{ s}^{-1} \text{ m}^{-1}$. However, as a result of this slight weakening of the CUI, the CUI showed a rapid and considerable increase in 2004 and 2009. Although both condition factor and environmental phenomena varied seasonally, the variation in Kn was related to CUI (F-stat = 6.186; $p = 0.0143$) and not for the SST (F-stat = 3.238; $p = 0.0745$). The model explains 5.19 % of the observed variability with R-sq. (adj) = 0.03. During the study period (1995–2011), the condition factor reached its maximum at $\text{CUI} \approx 1 \text{ m}^3 \text{ s}^{-1} \text{ m}^{-1}$ in 1995, 2003 and 2006 (Fig. 5).

229 3.3. Growth parameters

230 Based on the length frequency data, L_{∞} , K and ϕ' for July 2014 to June 2015 were estimated to be 35.4 cm, 0.26
231 year⁻¹ and 2.51 respectively. *Sardinella aurita* grows quickly during its first year, while subsequent growth rates
232 decline gradually with age. No statistical differences between the growth performance index (ϕ') of historical data
233 collected off the Senegal coast and data collected from other areas of the Eastern Central Atlantic was found
234 (ANOVA: F value = 2.371, $p = 0.139$) (Appendix, Table I). However, we did find a significant difference
235 between ϕ' data from the Senegal coast and other regions (West Atlantic, East Atlantic, and Mediterranean)
236 (ANOVA: F value = 10.81, $p = 0.001$) (Appendix, Table I). The Eastern Central Atlantic and Western Central
237 Atlantic did not differ statistically (Tukey: $df = 0.15$, $p = 0.171$), while the Eastern Central Atlantic and the
238 Mediterranean differed significantly (Tukey: $df = 0.33$, $p = 0.001$). The growth parameters (L_{∞} , K and ϕ') of *S.*
239 *aurita* in the Eastern Central Atlantic were higher than Western Central Atlantic and Mediterranean Sea (Fig. 6)
240 and decline over time (from 1932 to the present study), while in the West Atlantic, there was an increasing of L_{∞} .
241 Asymptotic length (L_{∞}) was lower in the Western Central Atlantic than in the Eastern Central Atlantic (Tukey: df
242 = 0.15, $p = 0.001$). *Sardinella aurita* inhabiting these two coastal systems (Eastern and Western Central Atlantic)
243 are characterized by higher lifespans (t_{max}), higher maximum lengths (L_{max}) and longer asymptotic lengths (L_{∞})
244 than those inhabiting the Mediterranean (Tukey: $df = 0.15$, $p = 0.001$) (Appendix, Table I). However, growth
245 parameters seem to have remained constant in the Mediterranean over the period examined (from 1932 to 2012;
246 Fig. 6).

247 3.4. Maturity

248 Sex ratio (SR) for *S. aurita* was in balance with around 51% females. The cumulative frequency curve of
249 individuals that had reached or passed maturity Stage 3 indicated that 50% (L_{50}) of males were mature at $24.8 \pm$
250 1.0 cm, while 50 % of females were mature at 25.7 ± 1.8 cm. We found significant differences between maturity
251 stages (proportions of fish in each stage) for the period investigated (1995–2011) (Table III). This sentence is
252 linked with the results showed in the Fig. 7 and 8.

253 Two main reproductive periods were determined from our analysis of sexual maturity stages and GSI. For this
254 study, the reproductive period was defined as the period between gonad maturation (Stage ≥ 3) to spawning. The
255 first reproductive period occurred from February to May off Senegal. Within this period, gonad maturation
256 occurred in February, April and May. Our analysis of sexual maturity showed that individuals in Stages V were

257 present in January, March, and April and from October to December after the first reproductive period. A second
258 reproductive period, with continuous spawning (more intense than during the first period), occurred from October
259 to the end of December.

260 The monthly record of sexual maturity stages showed that females in stage VI and VII were more abundant from
261 January to February and from July to December. The proportion of females at stages > IV peaked in January to
262 April, and September to December (Fig. 7a). Stage III females were mainly present in June. Stage V males were
263 present in February-March and from August to December. Stage III and IV males were found throughout the year
264 mostly from May to November (Fig. 7b).

265 Monthly changes in GSI followed sexual maturity Stages IV and V (Fig. 8). There were two spawning peaks in
266 April and October. In October, there was a higher presence of Stage IV and V individuals and higher GSI values
267 relative to September and November for males and females. A resting period between spawning occurred in June
268 and August.

4. DISCUSSION

4.1. Growth strategy of *Sardinella aurita*

Three main size classes exist among *S. aurita* inhabiting Senegal coastal waters, suggesting that the adults mainly inhabit more northern coastal waters, while juveniles and young breeders (<25 cm) are concentrated off the southern coast. The highest concentration of juveniles and young breeders off the “Petite côte” of Senegal has been linked to a nursery area located there (Roy et al., 1989; Tiedemann & Brehmer, 2017). Roy et al. (1989) assumed that the concentration of juveniles and young breeders in Senegal’s southern coastal waters may be due to differential functioning, whereas the northern waters, northward the Cap Vert peninsula (Dakar), where physical properties differ significantly, i.e., mesoscale activity, Ekman coastal divergence and upwelling patterns (Ndoye et al., 2017), which play a role in system productivity and retention processes mean (e.g., larval retention). Indeed, the “Petite côte” of Senegal is characterised by high phytoplankton biomass (Auger et al., 2016) and high chlorophyll content during upwelling events (Auger et al., 2016), which are conditions optimal for providing a suitable nursery (Tiedemann & Brehmer, 2017). Tiedemann & Brehmer (2017) suggested that the retention of ichthyoplankton over the shelf was related to food availability, and is important for the reproduction of many pelagic species during the planktonic stages. Indeed, they are able to detect variations in temperature and can follow masses of water whose behavior or quality (nutritive, calorific) suits them (Chikhi, 1995). Thus, temperature and food availability is generally the most important condition for habitat selection.

Growth parameters (L_{∞} , K and ϕ') for *S. aurita* in the Eastern Central Atlantic differ from patterns in the Mediterranean Sea, but no significant differences were found between the Eastern and Western Central Atlantic. Differences in *S. aurita* growth between the central Atlantic and Mediterranean may be related to differences in food abundance or environmental conditions (Wootton, 1998). The growth parameter and asymptotic length for a same short life species are higher in the most productive area (Fig. 6). The availability of food and temperature in Mediterranean is much lower than the Eastern Central Atlantic (Stergiou, 2000). Indeed, the instability of primary production and hydrographic conditions (Puerta et al., 2015) could make the Mediterranean less suitable for *S. aurita* populations. The complex mesoscale eddy complexes of the western Mediterranean resulting from the influx of Atlantic waters have different implications for feeding, transport and larval survival. This has important consequences for the productivity and functioning of the Mediterranean pelagic ecosystem (Puerta et al., 2015). According to Bakun (2013), some mid-scale vorticity features provide improved feeding and retention conditions

297 for pelagic larvae. While in the Californian Current, a significant delay in the spring transition (Barth et al., 2007)
298 might change the ecosystem, affecting primary production (Thomas & Brickley, 2006), zooplankton (Mackas et
299 al., 2006), and fish (Brodeur et al., 2018). For example, the growth of Atlantic herring *Clupea harengus*
300 *harengus* related in part to the temperature of summer water (Anthony & Fogarty, 1985) while in Pacific herring
301 *Clupea harengus pallasii*, Haist and Stocker (1985) found that SST of 12 ° - 13 ° C was optimal for juvenile
302 growth. In the Eastern Central Atlantic, the seasonal variability of wind forcing is the main driver of seasonal
303 variability in surface chlorophyll (Auger et al., 2016) and affects the distribution and abundance of fish
304 populations, and their associated fisheries, over a large range of timescales (Thiaw et al., 2017). The
305 environmental processes responsible for changes in growth rates between different regions are not always clearly
306 identified, but in most of the examples given, they seem to be related to temperature and abundance of food.

307 In addition to environmental factors, high rates of fishing mortality can alter population structure and growth
308 rates. For example, in the Mediterranean Sea (Greece), it increased from 32 cm in 1952 (Ananiades, 1952) to 21
309 cm in 2005 (Tsikliras et al., 2005) ($\Delta = - 11$ cm). On the contrary, there is an increase in asymptotic length in the
310 western central Atlantic of Venezuela (Mendoza et al., 1994) estimated at 26 cm (in 1994) and recently, the study
311 of Barrios et al. (Barrios et al., 2010) showed a slight increase in asymptotic length to 27 cm ($\Delta = + 1$ cm). In
312 Atlantic Eastern Central, we observe a decrease of asymptotic length L_{∞} of *S. aurita*, particularly in Senegal, from
313 42 cm in 1950 (Postel, 1955) to 35.4 cm in this study (2013-2014) ($\Delta = - 7$ cm). In Greece, the commercial
314 exploitation of *S. aurita* has steadily increased since the early 1990's (National Statistical Service of Hellas, 1990-
315 2002) due to a high fish market demand drive by the canning industry as bait for the profitable tuna and swordfish
316 fisheries. Total landings increased from 69 t in 1990 to 2 733 t in 2002 (Tsikliras, 2004). At the same time, in the
317 west-central Atlantic area, *S. aurita* has not yet reached the state of overexploitation. In addition, a drastic
318 reduction in catches was reported in 2009 (WECAFC, 2014). *Sardinella aurita* has been overexploited off West
319 Africa for a decade (FAO, 2016) and Thiaw et al. (2017) assumed that increased fishing presence could have a
320 strong influence on the decline in *S. aurita* growth in Senegal. As concluding remark, the environmental
321 characteristics and the level of fishing pressure on a small pelagic fish species had a balanced impact on their
322 growth parameters and asymptotic length at least for *S. aurita*, and thus should be considered as phenotypic
323 parameters.

4.2. Reproductive strategy of *Sardinella aurita*

325 A sex ratio biased slightly toward females (51.1%) agrees with results of several authors studying the *S. aurita*
326 population off Senegal (Conand, 1977; Boëly, 1982), Mauritania (Wague & Mbodj, 2002) and the Mediterranean.
327 Boëly (1979) noted there was a slight predominance of females (55%) among younger and older individuals off
328 the Senegal coast. This slight difference in sex ratio could be attributed to differences in reproductive growth
329 [(more females are produced) (Boëly, 1982)], differential mortality or differences in the energy cost of
330 reproduction between sexes (Tsikliras & Antonopoulou, 2006). Size at first sexual maturity for males and females
331 was 24.8 and 25.7 cm, respectively in our present study. Conand (1977), Boëly (1982), and Fréon (1988) found
332 smaller size at first sexual maturity than present study while Goudiaby et al. (2008) and Samba (2011) found
333 larger sizes at first sexual maturity than present study (see Appendix, Table II) for both males and females.
334 According to Boëly (1982), size of *S. aurita* at first sexual maturity may vary from one year to another or within
335 the same year. In the Mediterranean, the size at first maturity of *S. aurita* varies from 15 cm to 16.8 cm for
336 females and from 14 cm to 15.8 cm for males, whereas in the Western Central Atlantic (where most studies have
337 focused on females), the size at first sexual maturity ranged from 18 cm to 20 cm. We can therefore assume that
338 the size at first sexual maturity for *S. aurita* also varies throughout its distribution. Size at first maturity depends
339 on environmental and genetic factors (Wootton, 1998), but may also be influenced by other factors, such as long-
340 term fishing pressure (Thiaw et al., 2017). Adaptive phenotypic plasticity is often the only way for populations to
341 react quickly to environmental change and thus ensure their survival. A change in size at first maturity for an *S.*
342 *aurita* population is probably one of the best indicators of its response to changes and/or stress in its environment.
343 The length-weight relationship parameters calculated for *S. aurita* caught in the Eastern Central Atlantic
344 (Mauritania and Senegal) are similar in values to those calculated for fish in the Western Central Atlantic (off
345 Venezuela). However, lower values have been calculated for fish caught in the Mediterranean (Koutrakis &
346 Tsikliras, 2003). Many factors (*e.g.*, morphology, proportion of fatty acids, sex, sexual stage, egg density,
347 condition of the digestive tract, etc.) can lead to fluctuations in the weight of an individual fish (Wootton, 1998).
348 Weight is determined by the condition factor (Kn). Variations in Kn suggest that males and females adopt a
349 similar energy intake strategy during gonad maturation and spawning in that Kn reaches its maximum value the
350 month before the beginning of spawning (May or September). According to Boëly (1982) and Fréon (1988), the
351 weight of *S. aurita* seems to be intimately related to upwelling intensity and agrees with the results obtained in
352 present study..
353

354 Studies of *S. aurita* populations sampled off the coast of Senegal have shown that sexual activity occurs over
355 most of the year and that number and magnitude of spawning peaks change from year to year and from one area
356 to another (see Appendix, Table II). The spawning pattern we found in our study corresponds to the patterns
357 identified by Cury and Fontana (1988) in Senegal. However, the pattern differed from those identified by Conand
358 (1977), Boëly (1982), Goudiaby et al (2008) and Samba (2011) for Senegal and by Wague and Bodj (2002) for
359 Mauritania. Spawning seems to be correlate with periods of high zooplankton abundance. In West Africa,
360 breeding strategies are associated with local environmental conditions (Fréon, 1988; Roy et al., 1989). In places
361 where spawning occurs in twice annually, *S. aurita* is better able to exploit its environment and maintain its
362 population. A single peak of reproduction over a two-month (Tsikliras & Antonopoulou, 2006), three-month
363 (Gaamour et al., 2001), or four-month period (Ananiades, 1952) occurs in the Mediterranean, whereas a more
364 extended breeding season occurs in the Western Central Atlantic (October to April), which supports our results
365 for those coastal areas. We observed in present study that peak spawning corresponds to periods of low sea
366 surface temperature (February or March, depending on the year). This may be due to temperature having a direct
367 effect on the flow of energy through an ecosystem and population dynamics (Thiaw et al., 2017; Diankha et al.,
368 2018). Species-specific differences in fish tolerance to abiotic factors (e.g., temperature, food availability, and
369 other abiotic factors) show species-specific changes in population fecundity. In patients, embryonic stage duration
370 and larval survival are treated with physical processes, at water temperature and food availability, which affect
371 recruitment in upwelling systems (Tiedemann & Brehmer, 2017). Repetitive breeding tends to have a longer
372 spawning season than recruited spawners; thus increasing the likelihood that their offspring will encounter
373 favourable environmental conditions for growth and survival. The many environments studied, it is generally
374 accepted that changes in SST alter ecological stability. Ben-Tuvia (1960) found a positive relationship between
375 the SST in March (pre-spawning period) and the strength of the year-round sardinella age classes in Israel
376 (Mediterranean). In Southern California (Eastern Central Atlantic), warm temperatures allow larval development
377 and successful breeding of sardines (MacCall, 2009). Thus, water temperatures and food availability can affect
378 abundance (Thiaw et al., 2017), recruitment (Diankha et al., 2018), stages of egg and larvae (Tiedemann &
379 Brehmer, 2017), distribution (Sarré et al., 2018) and productivity (Auger et al., 2016) pelagic fish. Any
380 modification at the level of temperature and upwelling might cause a mismatch between *S. aurita* reproduction
381 and occurrence of their food and other organisms, leading to recruitment failures.

382 *Sardinella aurita* adopts a specific growth and reproductive strategy that is a reaction to specific environmental
383 cues. We observed significant differences in the Eastern Atlantic Ocean vs the Mediterranean Sea. The primary
384 spawning period for *S. aurita* occurs in times of high zooplankton abundance (during upwelling) and in places
385 that is generally the most favourable for early life stage development. This strategy allows *S. aurita* to adapt its
386 abundance to fluctuations in food availability. Surprisingly, even under the effect of multiple stressors (mainly
387 environmental changes), the size at spawning for *S. aurita* does not vary in the Canary Current LME. Such result
388 should be considered in management purposes particularly in the context of climate change, and suggest that such
389 parameters must be regularly updated for accurate fisheries management.

390 **ACKNOWLEDGEMENTS**

391 We would like to thank the editor and two anonymous reviewers for their insightful comments on the manuscript
392 and their very useful suggestions, which significantly improved its quality. Data analysis was done inside the
393 AWA project funded by IRD and the BMBF (grant 01DG12073E), www.awa.ird.fr (SRFC: Sub Regional
394 Fisheries Commission), the Preface project funded by the European Commission's Seventh Framework Program
395 (2007-2013) under Grant Agreement number 603521, <https://preface.b.uib.no/> and the DAAD program under
396 Grant Agreement number 57314022. The data for this study were collected through the CRODT (Senegal)
397 fisheries information system from 1996–2013 and the ComFish project from 2014-2015 (funded by USAID). We
398 are thankful to USAID Comfish project for allowing the use of their database and all CRODT technicians for
399 their professionalism during the data collection on the field all the year around.

400 **AUTHOR CONTRIBUTIONS**

401 Conceptualization: BSB PB WE.

402 Data curation: BSB FNS.

403 Formal analysis: BSB KB JK.

404 Methodology: BSB KB.

405 Project administration: BSB PB.

406 Resources: FNS.

407 Software: BSB KB.

408 Validation: PB, WE.

409 Supervision: MF MD.

410 Visualization: BSB.

411 Writing – original draft: BSB.

412 Writing – review & editing: PB WE JK KB.

413 All authors contributed critically to the drafts and gave final approval for publication.

414 **COMPETING INTERESTS**

415 The authors have declared that no competing interests exist.

- 417 Ananiades, C. (1952) Quelques Considérations Sur L'allache Biométriques (*Sardinella aurita* CV). *Praktika*
 418 *Hellenic Hydrobiological Institute* **5**, 5–45.
- 419 Anthony, V. C. & Fogarty, M. J. (1985) Environmental Effects on Recruitment, Growth, and Vulnerability of
 420 Atlantic Herring (*Clupea Harengus Harengus*) in the Gulf of Maine Region. *Canadian Journal of Fisheries and*
 421 *Aquatic Sciences* **42**, s158–s173.
- 422 Auger, P.-A., Gorgues, T., Machu, E., Aumont, O. & Brehmer, P. (2016) What Drives the Spatial Variability of
 423 Primary Productivity and Matter Fluxes in the North-West African Upwelling System? A Modelling Approach
 424 and Box Analysis. *Biogeosciences* **13**, 6419–6440.
- 425 Bakun, A. (1990) Global Climate Change and Intensification of Coastal Ocean Upwelling. *Science* **247**, 198–201.
- 426 Bakun, A. (2013) Ocean Eddies, Predator Pits and Bluefin Tuna: Implications of an Inferred 'low Risk-limited
 427 Payoff' reproductive Scheme of a (Former) Archetypical Top Predator. *Fish and Fisheries* **14**, 424–438.
- 428 Barrios, A., Guzmán, R., Mendoza, J., Gómez, G. & Vizcaino, G. (2010) Parámetros de Crecimiento Y
 429 Mortalidad Total de La Sardina *Sardinella aurita* (Valenciennes, 1810) En El Nororiente Venezolano Durante El
 430 Período 1990 a 2008. *Zootecnia Tropical* **28**, 457–466.
- 431 Barth, J. A., Menge, B. A., Lubchenco, J., Chan, F., Bane, J. M., Kirincich, A. R., McManus, M. A., Nielsen, K.
 432 J., Pierce, S. D. & Washburn, L. (2007) Delayed Upwelling Alters Nearshore Coastal Ocean Ecosystems in the
 433 Northern California Current. *Proceedings of the National Academy of Sciences* **104**, 3719–3724.
- 434 Barton, E. D., Field, D. B. & Roy, C. (2013) Canary Current Upwelling: More or Less? *Progress in*
 435 *Oceanography* **116**, 167–178.
- 436 Ben-Tuvia, A. (1960) Synopsis of Biological Data on *Sardinella Aurita* of the Mediterranean Sea and Other
 437 Waters. *Proceedings of the World. Scientific Meeting on the Biology* **14**, 287–312.
- 438 von Bertalanffy, L. (1938) A Quantitative Theory of Organic Growth (Inquiries on Growth Laws. II). *Human*
 439 *biology* **10**, 181–213.
- 440 Boëly, T. (1979) Biologie de deux espèces de sardinelles *Sardinella aurita* Valenciennes, 1847 et *Sardinella*
 441 *maderensis* Lowe, 1841 des côtes sénégalaises. Thèse de doctorat, Paris: Université Paris VI.
- 442 Boëly, T. (1982) Etude Du Cycle Sexuel de La Sardinelle Ronde (*Sardinella aurita* Val. 1847) Au Sénégal.
 443 *Océanographie tropicale* **17**, 3–13.
- 444 Boëly, T., Fréon, P. & Stéquert, B. (1982) La Croissance de *Sardinella aurita* (Val. 1847) Au Sénégal.
 445 *Océanographie Tropicale* **17**, 103–119.
- 446 Boltachev, A. R. & Karpova, E. P. (2014) Faunistic Revision of Alien Fish Species in the Black Sea. *Russian*
 447 *journal of biological invasions* **5**, 225–241.
- 448 Brodeur, R. D., Hunsicker, M. E., Hann, A. & Miller, T. W. (2018) Effects of Warming Ocean Conditions on
 449 Feeding Ecology of Small Pelagic Fishes in a Coastal Upwelling Ecosystem: A Shift to Gelatinous Food Sources.
 450 *Marine Ecology Progress Series*.
- 451 Chaboud, C., Fall, M., Ferraris, J., Fontana, A., Fonteneau, A., Laloë, F., Samba, A. & Thiao, D. (2015)
 452 Comment on 'Fisheries Catch Misreporting and Its Implications: The Case of Senegal'. *Fisheries Research* **164**,
 453 322–324.
- 454 Charouki, N., Sarré, A., Serghini, M., Ebaye, O. S. & Krakstad, J. O. (2011) Contribution to Acoustic Targets
 455 Identification in Northwest Africa by Statistical Schools Properties. In *Science and Management of Small*
 456 *Pelagics. Symposium on Science and the Challenge of Managing Small Pelagic Fisheries on Shared Stocks in*
 457 *Northwest Africa* pp. 285–294 Rome: FAO.
- 458 Chikhi, L. (1995) Différenciation génétique chez *Sardinella aurita* et *S. maderensis* Allozymes et ADN
 459 mitochondrial. Thèse nouveau doctorat, France: Université de Paris 06.
- 460 Conand, C. (1977) Contribution À L'étude Du Cycle Sexuel et de La Fécondité de La Sardinelle Ronde,
 461 *Sardinella aurita*: Pêche Sardinière Dakaroise En 1975 et Premier Semestre 1976. *Cahiers ORSTOM. Série*
 462 *Océanographie* **15**, 301–312.
- 463 Craney, T. A. & Surlles, J. G. (2002) Model-Dependent Variance Inflation Factor Cutoff Values. *Quality*
 464 *Engineering* **14**, 391–403.
- 465 Cury, P. & Fontana, A. (1988) Compétition et Stratégies Démographiques Comparées de Deux Espèces de
 466 Sardinelles (*Sardinella aurita* et *Sardinella maderensis*) Des Côtes Ouest-Africaines. *Aquatic living resources* **1**,
 467 165–180.
- 468 De Mendiburu, F. (2014) *Agricolae: Statistical Procedures for Agricultural Research. R package version 1*.

469 Diankha, O., Ba, A., Brehmer, P., Brochier, T., Sow, B. A., Thiaw, M., Gaye, A. T., Ngom, F. & Demarcq, H.
470 (2018) Contrasted Optimal Environmental Windows for Both *Sardinella* Species in Senegalese Waters. *Fisheries*
471 *Oceanography* **27**, 351–365.

472 Ekman, V. W. (1905) On the Influence of the Earth's Rotation on Ocean-Currents. *Arkiv Foer Matematik,*
473 *Astronomi och Fysik* **2**, 1–52.

474 FAO. (2016) *Report of the FAO working group on the assessment of small pelagic fish off Northwest Africa*
475 *Casablanca, Morocco, 20–25 July 2015/ Rapport du groupe de travail de la FAO sur l'évaluation des petits*
476 *pélagiques au large de l'Afrique Nord-Occidentale Casablanca, Maroc, 20-25 juillet 2015 (Casablanca, Maroc:*
477 *FAO). N°1122. Casablanca, Maroc: FAO. 243 p.*

478 Felder, D. L. (2009) *Gulf of Mexico Origin, Waters, and Biota: Biodiversity*, College Station. Biodiversity. Texas:
479 Texas A&M University Press. Vol. 1.

480 Fontana, A. (1969) Etude de La Maturité Sexuelle Des Sardinelles *Sardinella eba* (Val) et *Sardinella aurita* C. et
481 V. de La Région de Pointe-Noire. *Cahiers ORSTOM, serie Oceanographique* **VII**, 111–114.

482 Fréon, P. (1988) Réponses et adaptations des stocks de Clupéidés d'Afrique de l'Ouest à la variabilité du milieu
483 et de l'exploitation: analyse et réflexion à partir de l'exemple du Sénégal. Thèse de doctorat, Paris: Université
484 d'Aix Marseille II.

485 Froese, R. & Pauly, D. (2016) FishBase: World Wide Web Electronic Publication Version.

486 Gaamour, A., Missaoui, H., Ben-Abdallah, L. & El Ahmed, A. (2001) Parametres Biologiques de La Sardinelle
487 Ronde (*Sardinella aurita* Valenciennes, 1847) Dans La Région Du Cap Bon (Canal Siculo-Tunisien). *GFCM,*
488 *Kavala. Greece 2001.*

489 Gassman, J., Eslava, N. & González, L. W. (2008) Reproducción de La Sardina, *Sardinella aurita* (Cupleiformes:
490 Clupeidae) Del Sureste de La Isla de Margarita, Venezuela. *Revista de biología tropical* **56**, 1813–1824.

491 Goudiaby, K. D., Samb, B. & Sylla, M. (2008) Contribution À La Connaissance de La Biologie Des Sardinelles
492 (*Sardinella aurita* et *Sardinella maderensis*) Du Littoral Sénégalais. In *Science et aménagement des petits*
493 *pélagiques. Symposium sur la science et le défi de l'aménagement des pêcheries de petits pélagiques sur les*
494 *stocks partagés en Afrique nord-occidentale* pp. 39–56 Rome: Food and Agriculture Organization of the United
495 Nations.

496 Haist, V. & Stocker, M. (1985) Growth and Maturation of Pacific Herring (*Clupea Harengus Pallasii*) in the Strait
497 of Georgia. *Canadian Journal of Fisheries and Aquatic Sciences* **42**, s138–s146.

498 Harrell Jr, F. E. (2017) *With Contributions from Charles Dupont and Many Others: Hmisc: Harrell*
499 *Miscellaneous. R Package Version 3.17-2.*

500 Hastie, T. & Tibshirani, R. (1986) Generalized Additive Models. *Statistical Science* **1**, 297–318.

501 Isselmou, O. & Baye, C. (2009) Evaluation et cartographie de l'abondance du stock de la sardinelle ronde dans la
502 zone Nord-Ouest Africaine. Thèse de maîtrise, Pôle Halieutique (Agrocamus Ouest) et CRH de Sète.

503 Koutrakis, E. T. & Tsikliras, A. C. (2003) Length–weight Relationships of Fishes from Three Northern Aegean
504 Estuarine Systems (Greece). *Journal of Applied Ichthyology* **19**, 258–260.

505 Le Bec, C. (1983) Cycle Sexuel et Fécondité de La Sole *Solea Vulgaris* (Quensel, 1806) Du Golfe de Gascogne.
506 *Revue des travaux de l'Institut des pêches maritimes* **47**, 179–189.

507 Le Cren, E. D. (1951) The Length-Weight Relationship and Seasonal Cycle in Gonad Weight and Condition in
508 the Perch (*Perca Fluviatilis*). *The Journal of Animal Ecology* **20**, 201–219.

509 MacCall, A. D. (2009) Mechanisms of Low-Frequency Fluctuations in Sardine and Anchovy Populations. In
510 *Climate Change and Small Pelagic Fish* pp. 285–299 Cambridge: Cambridge University Press.

511 Mackas, D. L., Peterson, W. T., Ohman, M. D. & Lavaniegos, B. E. (2006) Zooplankton Anomalies in the
512 California Current System before and during the Warm Ocean Conditions of 2005. *Geophysical Research Letters*
513 **33**, S07.

514 McQuinn, I. H., Cleary, L. & O'Boyle, R. N. (1990) Estimation de La Mortalité Naturelle. In *Méthodes*
515 *d'évaluation des stocks halieutiques* pp. 441– 464 Nouvelle- Ecosse, Canada: Projet CIEO-860060, Centre
516 International d'Exploitation des Oceans, Halifax.

517 Mendoza, J. J., Fréon, P. & Guzmán, R. (1994) VPA Estimates of Fishing Mortality and Exploited Biomass from
518 *Sardinella Aurita* Catch-at-Length Data in Eastern Venezuela. *NAGA, the ICLARM quarterly* **17**, 32–36.

519 Mildenberger, T. K., Taylor, M. H. & Wolff, M. (2017) Extending ELEFAN in R. *Figshare*.

520 Ndoye, S., Capet, X., Estrade, P., Sow, B., Machu, E., Brochier, T., Döring, J. & Brehmer, P. (2017) Dynamics of
521 a 'low-enrichment High-retention' Upwelling Center over the Southern Senegal Shelf. *Geophysical Research*
522 *Letters* **44**.

523 Pauly, D. (1983) *Some Simple Methods for the Assessment of Tropical Fish Stocks*. Fisheries Report 234. Food
524 and Agriculture Organization of the United Nations. 52.

525 Peck, M. A., Reglero, P., Takahashi, M. & Catalán, I. A. (2013) Life Cycle Ecophysiology of Small Pelagic Fish
526 and Climate-Driven Changes in Populations. *Progress in Oceanography* **116**, 220–245.

527 Postel, E. (1955) Résumé Des Connaissances Acquisées Sur Les Clupéidés de l'Ouest Africain. *Rapp. Rapport*
528 *Conseil d'Exploration de La Mer* **137**, 14–17.

529 Puerta, P., Hunsicker, M. E., Quetglas, A., Álvarez-Berastegui, D., Esteban, A., González, M. & Hidalgo, M.
530 (2015) Spatially Explicit Modeling Reveals Cephalopod Distributions Match Contrasting Trophic Pathways in the
531 Western Mediterranean Sea. *PloS one* **10**, e0133439.

532 Roy, C., Cury, P., Fontana, A. & Belvèze, H. (1989) Stratégies Spatio-Temporelles de La Reproduction Des
533 Clupéidés Des Zones D'upwelling d'Afrique de l'Ouest. *Aquatic Living Resources* **2**, 21–29.

534 Samba, O. (2011) Nouvelle évaluation des caractéristiques biologiques de *Sardinella aurita* (valenciennes, 1847)
535 et *sardinella maderensis* (lowe, 1841). Mémoire DEA, Senegal: Université Cheikh Anata Diop de Dakar (UCAD).

536 Sarré, A., Krakstad, J.-O., Brehmer, P. & Mbye, E. M. (2018) Spatial Distribution of Main Clupeid Species in
537 Relation to Acoustic Assessment Surveys in the Continental Shelves of Senegal and The Gambia. *Aquatic Living*
538 *Resources* **31**, 9.

539 Stergiou, K. I. (2000) Life-History Patterns of Fishes in the Hellenic Seas. *Web ecology* **1**, 1–10.

540 Sylla, M., Brochier, T., Pecquerie, L., Ndiaye, O., Richaume, M. & Brehmer, P. (2016) Senegalese *Sardinella*
541 *aurita* from 1995 to 2011: Review of Size Spectra, Sex Ratio, Gonadal Somatic and Condition Indexes from
542 Small Scale Main National Landing Sites. In *Brehmer Patrice (Ed.), Ba B. (Ed.), Kraus G. (Ed.)* pp. 51–51
543 Dakar, Senegal: SRFC/CSRP; IRD.

544 Thiao, D. (2009) Un système d'indicateurs de durabilité des pêcheries côtières comme outil de gestion intégrée
545 des ressources halieutiques sénégalaises. Ph.D. Thesis, France: Université de Versailles Saint-Quentin-en-
546 Yvelines.

547 Thiaw, M., Auger, P.-A., Sow, F. N., Brochier, T., FAYE, S., Diankha, O. & Brehmer, P. (2017) Effect of
548 Environmental Conditions on the Seasonal and Inter-Annual Variability of Small Pelagic Fish Abundance off
549 North-West Africa: The Case of Both Senegalese *Sardinella*. *Fisheries Oceanography* **26**, 583–601.

550 Thomas, A. C. & Brickley, P. (2006) Satellite Measurements of Chlorophyll Distribution during Spring 2005 in
551 the California Current. *Geophysical Research Letters* **33**.

552 Tiedemann, M. & Brehmer, P. (2017) Larval Fish Assemblages across an Upwelling Front: Indication for Active
553 and Passive Retention. *Estuarine, Coastal and Shelf Science* **187**, 118–133.

554 Tsikliras, A. C. (2004) Biology and Population Dynamics of Round *Sardinella* (*Sardinella aurita* Valenciennes,
555 1847) in Kavala Gulf (Northern Aegean Sea, Greece). Ph.D. Thesis, Greece: Aristotle University of Thessaloniki.

556 Tsikliras, A. C. & Antonopoulou, E. (2006) Reproductive Biology of Round *Sardinella* (*Sardinella aurita*) in
557 North-Eastern Mediterranean. *Scientia Marina* **70**, 281–290.

558 Tsikliras, A. C., Koutrakis, E. T. & Stergiou, K. I. (2005) Age and Growth of Round *Sardinella* (*Sardinella*
559 *aurita*) in the Northeastern Mediterranean. *Scientia marina* **69**, 231–240.

560 Wague, A. & Mbodj, O. B. (2002) Etude de Quelques Aspects de La Reproduction Chez La Sardinelle Ronde
561 *Sardinella aurita* (Valenciennes, 1847) Pechee Le Long Des Côtes Mauritanienes. *Bulletin Scientifique de*
562 *l'Institut Mauritanien de Recherches Océanographiques et des Pêches* 13–18.

563 WECAFC, W. C. A. F. (2014) *Status of the stocks of the Western Central Atlantic-FAO statistical area 31. Port of*
564 *Spain, Trinidad and Tobago, 26-28 March 2014*. Fifteenth session. Port of Spain, Trinidad and Tobago: FAO. 9.

565 Wood, S. & Wood, M. S. (2017) Package 'mgcv'. *R package version* 1–7.

566 Wootton, R. J. (1998) *Ecology of Teleost Fishes*, 2nd ed. Dordrecht: Kuwer Academic Publishers.

567 Zuur, A. F., Ieno, E. N. & Elphick, C. S. (2010) A Protocol for Data Exploration to Avoid Common Statistical
568 Problems. *Methods in ecology and evolution* **1**, 3–14.

569

570 **Figures captions**

571 FIG. 1: Map of the study area with the localization of sampling stations corresponding to the main
572 landing ports along the Senegal coast. The northern section includes Saint-Louis, Kayar, and Yoff; the
573 southern section includes the “Petite côte” (Hann, Mbour, and Joal) and Casamance (Kafountine).
574 Gambian EEZ was not taking into account in the study, neither were Mauritanian EEZ and Bissau
575 Guinean EEZ.

576 FIG. 2: Size frequency distributions for *Sardinella aurita* caught off the Senegal coast from July 2013 to
577 June 2014. Total length was used. Three modes exist in the frequency data: one for fish located off the
578 northern coast (black) (32 cm, n = 10.109) and two for fish caught off the southern coast (grey) (13 and
579 24 cm, n = 22.237).

580 FIG. 3: Length-weight relationships for *Sardinella aurita* caught off the coast of Senegal (data: 1995 to
581 2011; n = 4397). The black line is the exponential regression curve fitted to all measurements.

582 FIG. 4: *Sardinella aurita* condition factor (K_n) data (black line), combined to seasonal environmental
583 components for Senegal coastal waters (1995 to 2011): coastal Sea Surface Temperature (dashed
584 line) (Modis AVHRR) and Coastal Upwelling Index (grey line). (a) Monthly and (b) annual mean data.


585 FIG. 5: Condition factor (K_n) of *Sardinella aurita* as a function of Coastal Upwelling Index (CUI)
586 throughout the study (1995-2011).


587 FIG. 6: Auximetric plot (double logarithmic plot of parameters K and L_∞) for *Sardinella aurita*
588 populations for four coastal areas. Data from this study in Senegal (square) and from a literature review
589 of Eastern Central Atlantic (circles), Mediterranean Sea (triangles) and Western Central Atlantic (plus
590 signs). Abbreviations: L_∞ : (asymptotic length, in cm) and K : [coefficient of growth rate (by year)].

591 FIG. 7: Monthly sexual maturity stages (*sensu* Fontana 1969) for *Sardinella aurita* off the Senegal
592 coast (1995 to 2011). (a) female (n = 2131) and (b) male (n = 2266)

593 FIG. 8: Gonado-somatic index (GSI) for *Sardinella aurita* in Senegal coastal waters (1995 to 2011). (a)
594 female (n = 2131) and (b) male (n = 2266). Means are represented by the horizontal line inside the box
595 plots. Standard errors are represented by circles.

596






601

602 FIG. 2


603



604


605 FIG. 3

606



607

608 a)




609

610 b)

611 FIG. 4

612


613



614

615 FIG. 5

616




617

618 FIG. 6

619


620



621

622

a)



623


624

b)

625


FIG. 7

626



627

628 a)



629

630 b)

631 FIG. 8

632

633 TABLE I: *Sardinella aurita*. Number of monthly observations for all variables except weight, by sex
 634 and maturity stages.

	Month													
		1	2	3	4	5	6	7	8	9	10	11	12	Total
Sex	F	219	135	170	276	229	401	113	112	86	113	244	168	2266
	M	209	115	167	227	199	362	119	93	91	140	255	154	2131
	Total	428	250	337	503	428	763	232	205	177	253	499	322	4397
Stages	I	22	60	18	56	110	160	98	56	6	27	56	4	673
	II	33	27	45	66	105	184	38	51	9	12	56	21	647
	III	28	31	34	36	39	135	17	11	1	7	35	13	387
	IV	91	58	73	75	40	114	39	33	22	29	122	78	774
	V	175	38	97	173	64	68	20	36	107	139	132	119	1168
	VI	65	18	33	68	51	77	14	12	31	25	77	67	538
	VII	14	18	37	29	19	25	6	6	1	14	21	20	210
	Total	428	250	337	503	428	763	232	205	177	253	499	322	4397

635

636 TABLE II: A brief description of the maturity stages of *Sardinella aurita* used in this study according to
 637 Fontana (1969).

Stage	State	Description
I	Immature	Female gonad small and firm, transparent or light pink, invisible oocytes. Male gonad white or slightly translucent, very fine.
II	Rest	Characteristics of female and male gonads substantially identical to Stage I.
III	Maturation start	Female gonad, firm and of color varying from pale pink to light orange, some oocytes are sometimes visible through the ovarian membrane. Male gonad whitish firm, no liquid will flow if an incision is made.
IV	Pre-egg / pre-release	Female gonad bigger and less firm usually light orange then dark. Oocytes visible through the ovarian membrane make the surface of the ovary granular Male gonad softer and white. A whitish liquid flows as soon as an incision is made.
V	Sperm laying / Emission	Female gonad very large occupying the entire abdominal cavity. Very fine ovarian membrane. Hyaline and large-sized ovules are perfectly visible and are expelled at the slightest pressure on the abdomen. Large and soft male gonad. The sperm flows at the slightest pressure exerted on the abdomen.
VI	End of spawning / Emission and recovery before subsequent spawning	The female ovary is flask very vascularized. Its color varies from pink salmon to red. Through the ovarian membrane the oocytes of 460 micrometers are perfectly visible. Numerous hyaline spaces / Flaccid male gonad with a very fine vascularization especially in the posterior part.
VII	End of the last laying / Emission and involution of the ovary	Female ovary completely collapsed very flaccid of red color due to a very strong vascularization. At this stage, the ovary has the characteristic appearance of an empty bag. Male gonad very flaccid, exhausted and strongly vascularized.

638

639 TABLE III: Proportion of *Sardinella aurita* male and female at the different sexual maturity stages (report to table I) in Senegal (data: 1995 to 2011). χ^2 : Chi-square
640 test; + : < 0.05 .

Stages	Female (%)	Male (%)	X^2	p -value
I	56.8	43.2	2.658E+09	+
II	66.6	33.4	1.0519e+10	+
III	12.6	87.4	2.188e+10	+
IV	20.8	79.2	2.2785e+10	+
V	34.9	65.1	6.424E+09	+
VI	93.2	6.8	2.9401e+10	+
VII	98.9	1.1	1.3085e+10	+

641
642

643 **Appendix**

644 TABLE I: Growth parameters and natural mortality of *Sardinella aurita* estimated in Senegal (present study and historical review) and from Eastern and Western
645 Central Atlantic as well as the Mediterranean Sea. The growth parameters have been estimated according to three different methods, *i.e.*, scale, otoliths or length
646 frequencies as in the present study (size expressed in cm). TL: Total Length, FL: Fork Length, L_{∞} : L_t value when the growth rate is zero; K: coefficient of growth
647 rate, t_0 : the time “t” when the fish length is zero, ϕ' : growth performance index, M: natural mortality, t_{max} : the longevity of the species. NR: not reported.

Country	Method	L_{∞}	L_{∞}	K	t_0	ϕ'	t_{max}	M	References
		FL	TL						
Western Central Atlantic									
Brazil	Scales	20.7	24.4	0.44	-0.39	2.42	6.42	0.46	Matsuura (1971)
Brazil	Otoliths	19.6	23.0	0.72	-0.25	2.58	3.91	0.76	Saccardo et al. (1988)
Brazil	Otoliths	23.2	27.5	0.55	-0.30	2.62	5.15	0.58	Cergole et al. (2002)
Venezuela	NR	20.5	24.1	0.32	-0.55	2.27	8.82	0.34	Heald & Griffiths (1967)
Venezuela	Length frequencies	22.5	26.6	1.26	-0.13	2.95	2.25	1.33	Mendoza et al. (1994)
Venezuela	Length frequencies	27.3	32.6	0.66	-0.05	2.85	4.48	0.66	Barrios et al. (2010)
Mediterranean									
Algeria	Length frequencies	29.3	35.0	0.24	-0.70	2.47	12.01	0.24	Bouaziz et al. (2001)
Croatia	Length frequencies	28.6	34.2	0.36	-0.94	2.62	7.39	0.40	Mustać and Sinovčić (2012a)
Egypt	NR	22.0	26.0	0.53	0.34	2.55	5.00	0.60	El-Maghraby et al. (1970)
Greece	Scales	27.3	32.6	0.28	-0.57	2.47	10.14	0.29	Ananiades (1952)
Greece	Scales	18.3	21.4	0.32	-0.55	2.17	8.82	0.34	Tsikliras et al. (2005)
Israel	Length frequencies	26.1	31.1	0.25	-1.80	2.38	10.20	0.29	Ben Tuvia (1956)
Spain	Length frequencies	23.2	27.5	0.45	-0.38	2.53	6.28	0.47	Navarro (1932)
Tunisia	Otolith	26.3	31.3	0.24	-2.50	2.37	10.00	0.30	Gaamour et al. (2001)
Eastern Central Atlantic									
Canary Islands	Length frequencies	34.6	41.5	0.31	-0.63	2.71	11.34	0.26	Navarro (1932)
Congo	Length frequencies	31.0	37.0	0.69	-0.56	2.96	5.82	0.51	Rosignol (1955)
Mauritania	Scales	34.6	41.6	0.26	-0.87	2.65	13.11	0.22	Chesheva (1998)
Mauritania	Otoliths	31.0	37.0	0.39	1.46	2.72	9.15	0.32	Pascual-Alayón et al. (2008)
Mauritania	Scales and Otoliths	32.9	39.4	0.32	-1.78	2.69	10.89	0.27	Pascual-Alayón et al. (2008)
Mauritania	Otoliths	30.4	36.3	0.41	-0.94	2.72	6.38	0.47	Iglesias et al. (2009)
Mauritania - Senegal	Scales	34.0	40.7	0.33	-0.60	2.74	10.81	0.27	Thuoc & Szypula (1973)
Mauritania - Senegal	Scales	38.1	45.9	0.36	-0.69	2.87	9.91	0.30	Maxim and Maxim (1988)
Mauritania - Senegal	Length frequencies and Scales	31.2	37.3	0.97	-0.21	3.12	4.58	0.65	Boëly et al. (1979)
Senegal	Length frequencies	35.0	42.0	0.60	-0.24	3.02	4.75	0.63	Postel (1955)
Senegal	Scales	35.0	42.1	0.28	-0.99	2.70	12.22	0.24	Krzepowski (1982)
Senegal	Length frequencies and Scales	30.6	36.5	1.21	-0.06	3.20	3.97	0.75	Boëly et al. (1982)
Senegal	Length frequencies	29.4	35.4	0.26	-0.76	2.51	10.77	0.27	Present study

648 TABLE II: Reproductive parameters in Western Central Atlantic (*i.e.* Brazil and Venezuela), in Mediterranean Sea (*i.e.* Greece, Croatia and Tunisia) and in Eastern
649 Central Atlantic (*i.e.* Senegal and Mauritanian) for *Sardinella aurita*. TL: Total Length (cm); FL: Fork Length (cm); N: Number of individual; SR (%): Sex Ratio;
650 L₅₀ (cm) represents the size at which 50% of individuals in a given population reach the maturity; ‘a’ and ‘b’: allometric coefficient; (*) Reproduction period; NR:
651 not reported.

Country	Method	N	Month												Male	Female	a	b	SR	References
			1	2	3	4	5	6	7	8	9	10	11	12	L ₅₀	L ₅₀				
Western Central Atlantic Ocean																				
Brazil	TL	250	*	*	*	*							*	*	NR	18.60	NR	NR	NR	Petermann and Schwingel (2016)
Venezuela	TL	92834	*	*	*								*	*	NR	19.70	1.0377	3.399	NR	Fréon et al. (1997)
Venezuela	TL	3736	*	*	*							*	*	NR	20.00	NR	NR	55.5	Gassman et al. (2008)	
Mediterranean																				
Greece	TL	NR					*	*	*	*					14.00	15.00	NR	NR	NR	Ananiades (1952)
Croatia	TL	2033						*	*	*					15.80	16.60	NR	NR	NR	Mustać and Sinovčić (2012b)
Greece	TL	19	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	0.0087	2.950	NR	Koutrakis and Tsikliras (2003)
Greece	TL	7942					*	*							15.50	16.80	NR	NR	50.5	Tsikliras and Antonopoulou (2006)
Tunisia	TL	918							*	*	*				14.10	15.20	NR	NR	NR	Gaamour et al. (2001)
Eastern Central Atlantic Ocean																				
Mauritania	FL	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	0.007940	3.227	NR	Lawal and Mylnikov (1988)
Mauritania	TL	3073						*	*	*	*				34.61	34.61	NR	NR	55.0	Wague and Mbodj (2002)
Senegal	TL	1334					*	*				*	*		NR	23.45	NR	NR	52.0	Conand (1977)
Senegal	FL	2232	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	0.000005	3.194	NR	Boëly (1982a)
Senegal	TL	9645		*	*	*	*					*	*		NR	21.59	NR	NR	55.0	Boëly (1982b)
Senegal	FL	2433	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	0.006392	3.274	NR	Fréon (1988)
Senegal	NR	NR		*	*	*	*	*			*	*	*		NR	NR	NR	NR	NR	Cury and Fontana (1988)
Senegal	TL	3947				*	*	*				*	*		27.66	30.98	0.000038	3.162	50.5	Goudiaby et al. (2008)
Senegal	TL	NR										*	*		21.10	26.55	0.000007	3.040	NR	Samba (2011)
Senegal	TL	4397		*	*	*	*	*			*	*	*		24.82	25.72	0.008	3.02	51.1	Present study

652

653