

An Application of the Universality Theorem for Tverberg Partitions

Imre Bárány*

Nabil H. Mustafa†

Abstract

We show that, as a consequence of a new result of Pór on universal Tverberg partitions, any large-enough set P of points in \mathbb{R}^d has a $(d+2)$ -sized subset whose Radon point has half-space depth at least $c_d \cdot |P|$, where $c_d \in (0, 1)$ depends only on d . We then give an application of this result to computing weak ϵ -nets by random sampling. We further show that given any set P of points in \mathbb{R}^d and a parameter $\epsilon > 0$, there exists a set of $O\left(\frac{1}{\epsilon^{\lfloor \frac{d}{2} \rfloor + 1}}\right) \lfloor \frac{d}{2} \rfloor$ -dimensional simplices (ignoring polylogarithmic factors) spanned by points of P such that they form a transversal for all convex objects containing at least $\epsilon \cdot |P|$ points of P .

Keywords: Tverberg's theorem, Radon's lemma, weak ϵ -nets, half-space depth, transversals.

1 Introduction

Radon's lemma states that, given any set Q of $(d+2)$ points in \mathbb{R}^d , there always exists a partition of Q into two sets, say Q_1 and Q_2 , such that $\text{conv } Q_1 \cap \text{conv } Q_2 \neq \emptyset$. Further, if Q is in general position, then a dimension argument implies that such a partition $\{Q_1, Q_2\}$ —called a Radon partition of Q —is unique and $\text{conv } Q_1 \cap \text{conv } Q_2$ consists of a single point, called the *Radon point* of Q and denoted by $\text{Radon } Q$.

In this paper we present an application of the following statement, which is one consequence of a recent theorem of Pór (see [2]).

Lemma 1 (Proof in Section 2). *For every $d \in \mathbb{N}$ there is $f(d) \in \mathbb{N}$ such that every set $P \subset \mathbb{R}^d$ of $f(d)$ points in general position contains two disjoint sets $A, B \subset P$ with $|A| = d+2$, $|B| = d+1$ and the Radon point of A is contained in $\text{conv } B$. Furthermore, the Radon partition of A consists of two sets of sizes $\lfloor \frac{d}{2} \rfloor + 1$ and $\lceil \frac{d}{2} \rceil + 1$.*

We use Lemma 1 to prove the following theorem. Given a set P of points in \mathbb{R}^d , the *half-space depth* of a point $q \in \mathbb{R}^d$ with respect to P is defined to be the minimum number of points of P contained in any half-space containing q .

*Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences 13 Reáltanoda Street Budapest 1053 Hungary and Department of Mathematics University College London Gower Street, London, WC1E 6BT, UK. barany.imre@renyi.mta.hu.

†Université Paris-Est, Laboratoire d'Informatique Gaspard-Monge, Equipe A3SI, ESIEE Paris. mustafan@esiee.fr.

Theorem 2 (Proof in Section 2). *For every $d \in \mathbb{N}$ there is $h(d) \in \mathbb{N}$ such that every set P of at least $h(d)$ points in \mathbb{R}^d in general position contains a set $P' \subseteq P$ of size $(d+2)$ with Radon P' being contained in at least $\frac{|P|}{h(d)}$ vertex-disjoint simplices spanned by the points of $P \setminus P'$. In particular, Radon P' has half-space depth at least $\frac{|P|}{h(d)}$.*

We expect that Theorem 2 will find further applications in discrete and combinatorial geometry. Here we give an application to the computation of a weak ϵ -net for a given set P of points in \mathbb{R}^d in general position.

Definition 3. *Given a set P of points in \mathbb{R}^d and a parameter $\epsilon > 0$, a set $N \subseteq \mathbb{R}^d$ is a weak ϵ -net with respect to convex sets for P if for every convex set K with $|K \cap P| \geq \epsilon \cdot |P|$, we have $K \cap N \neq \emptyset$.*

Consider the following simple algorithm to compute a weak ϵ -net for a given set P of points in \mathbb{R}^d .

Algorithm Weak-Nets (Input: a set of points P , parameter $\epsilon > 0$)

Let $R \subseteq P$ be a uniform random sample of size $\frac{g(d)}{\epsilon} \log \frac{1}{\epsilon}$, for a constant $g(d)$ depending only on d .

$$Q = \left\{ \text{Radon } R' : R' \in \binom{R}{d+2} \right\}.$$

return $Q \cup R$.

Our application of Theorem 2 is the following.

Theorem 4 (Proof in Section 3). *Let P be a set of points in \mathbb{R}^d in general position and $\epsilon > 0$ a given parameter. Then there is a $g(d) \in \mathbb{N}$ such that a uniform random sample $R \subseteq P$ of size $\frac{g(d)}{\epsilon} \log \frac{1}{\epsilon}$ satisfies the following properties with probability at least $\frac{9}{10}$.*

1. $Q \cup R$ is a weak ϵ -net for P , where Q is the set of Radon points of all $(d+2)$ -sized subsets of R . That is, [Algorithm Weak-Nets](#) returns a weak ϵ -net for P of size $O\left(\frac{1}{\epsilon^{d+2}}\right)$, and
2. each convex object containing at least $\epsilon|P|$ points of P intersects the convex hull of at least one $\left(\left\lfloor \frac{d}{2} \right\rfloor + 1\right)$ -sized subset of R .

Remark. The first part of Theorem 4 gives a bound on the size of the ϵ -net that is weaker than the current best bound due to Matoušek and Wagner [6], which is of the order of $O\left(\frac{1}{\epsilon^d}\right)$ (ignoring polylogarithmic factors). Yet our construction of a weak ϵ -net is novel and interesting as it uses certain Radon points of the underlying set P .

The constant $\frac{9}{10}$ can be increased by picking a larger random sample R .

2 Proof of Lemma 1 and Theorem 2.

We need some definitions. We set $m = (r-1)(d+1) + 1$, and for $k \in [d+1]$ the block B_k is the set of integers $\{(r-1)(k-1) + 1, (r-1)(k-1) + 2, \dots, (r-1)k + 1\}$. The blocks are of size r each

and they almost form a partition of $[m]$, only neighboring blocks have a common element, namely $(r-1)k+1 \in B_k \cap B_{k+1}$ for all $k \in [d]$. Call an r -partition $\{I_1, \dots, I_r\}$ of $[m]$ *special* if $|I_j \cap B_k| = 1$ for every $j \in [r]$ and every $k \in [d+1]$.

Pór's result is about sequences $S = (a_1, \dots, a_N)$ of vectors in \mathbb{R}^d . A sequence (b_1, \dots, b_t) is a subsequence of length t of S if $b_j = a_{i_j}$ for all $j \in [t]$ where $1 \leq i_1 < i_2 < \dots < i_t \leq N$. Given a sequence $S = (a_1, \dots, a_N)$, $a_i \in \mathbb{R}^d$, an r -partition $\{S_1, S_2, \dots, S_r\}$ of S is in one-to-one correspondence with an r -partition $\{I_1, \dots, I_r\}$ of $[m]$ via $a_i \in S_j$ if and only if $i \in I_j$. An r -partition of S is called *special* if the corresponding r -partition of $[m]$ is special.

Tverberg's theorem states that given a set P of $(r-1)(d+1)+1$ points in \mathbb{R}^d , there exists a partition of P into r sets whose convex-hulls contain a common point.

We can now state Pór's result [10].

Theorem A (Universality theorem for Tverberg partitions). *Assume $d, r, t \in \mathbb{N}$, $r \geq 2$, and $m = (r-1)(d+1)+1 \leq t$. Then there exists $N = N(d, r, t) \in \mathbb{N}$ such that every sequence $S = (a_1, \dots, a_N)$ of vectors (in general position) in \mathbb{R}^d contains a subsequence $S' = (b_1, \dots, b_t)$ (of length t) such that the Tverberg partitions of every subsequence of length m of S' are exactly the special partitions.*

Note that when the points of S (or P) come from the moment curve $\Gamma(x) = \{\gamma(x) : x \in \mathbb{R}^+\}$ where $\gamma(x) = (x, x^2, \dots, x^d)$, then there is a natural ordering $S = (\gamma(x_1), \dots, \gamma(x_n))$ with $x_1 < x_2 < \dots < x_n$. Now let $0 < x_1 < \dots < x_n$ a *rapidly increasing* sequence of real numbers, meaning that, for every $h \in [n-1]$, x_{h+1}/x_h is at least as large as some (large) constant $c_{d,r,h}$ depending only on d, r, h . It is not hard to check that in this case all Tverberg partitions of all $m = (r-1)(d+1)+1$ long subsequences of S are the special ones. This (and other examples as well) show that no other set of partitions can be universal, i.e., that exist as Tverberg partitions in a large-enough point set.

We are going to apply the universality theorem in the special case $r = 3$ and $t = m = (r-1)(d+1)+1 = 2d+3$. In this case $N(d, r, t)$ depends on d only and thus we can set $f(d) = N(d, r, t) = N(d, 3, 2d+3)$.

Proof of Lemma 1. Order the elements of P arbitrarily to obtain a sequence $S = (p_1, \dots, p_{f(d)})$. Apply Theorem A to S with $r = 3$, $t = m = 2d+3$. We get a subsequence S' of length m all of whose Tverberg 3-partitions are exactly the special ones. Define $I_1 = \{z \in [m] : z \equiv 1 \pmod{4}\}$ and $I_2 = \{z \in [m] : z \equiv 3 \pmod{4}\}$ and $I_3 = \{z \in [m] : z \text{ is even}\}$. Note that $|I_1| = \lceil \frac{d}{2} \rceil + 1$, $|I_2| = \lfloor \frac{d}{2} \rfloor + 1$ and $|I_3| = d+1$.

It is easy to see that $\{I_1, I_2, I_3\}$ is a special partition of $[m]$: every block contains exactly one element of I_1, I_2, I_3 . Let the corresponding partition of S' be $\{S_1, S_2, S_3\}$. So $\bigcap_{i=1}^3 \text{conv } S_i \neq \emptyset$. Set $A = S_1 \cup S_2$ and $B = S_3$. Then the Radon point of A , which is $\text{conv } S_1 \cap \text{conv } S_2$, is contained in $\text{conv } B$. \square

Proof of Theorem 2. Consider a $(2d+3)$ -uniform hypergraph $\mathcal{H} = (P, E)$ on the vertex set P , where $e \in E$ if and only if the $(2d+3)$ points of e can be partitioned into two sets $e = e_1 \cup e_2$ such that $|e_1| = d+2$, and Radon $e_1 \in \text{conv } e_2$. We will call the set e_1 the *Radon-base* of the edge e . By the result of de Caen [4], any r -uniform hypergraph on n vertices and m edges contains an

independent set of size at least

$$\frac{r-1}{r^{\frac{r}{r-1}}} \cdot \frac{n^{\frac{r}{r-1}}}{m^{\frac{1}{r-1}}}.$$

On the other hand, Lemma 1 implies that any set Q of $f(d)$ points of P must contain two disjoint sets— $A_Q \subseteq Q$ of size $(d+2)$ and $B_Q \subseteq Q$ of size $(d+1)$ —such that $\text{Radon } A_Q \in \text{conv } B_Q$. Then the $(2d+3)$ points $A_Q \cup B_Q$ form an edge in \mathcal{H} . This implies that no subset of P of size $f(d)$ can be independent in \mathcal{H} . Thus, with $r = 2d+3$, we have

$$\frac{2d+2}{(2d+3)^{\frac{2d+3}{2d+2}}} \cdot \frac{|P|^{\frac{2d+3}{2d+2}}}{|E|^{\frac{1}{2d+2}}} \leq \text{size of max. ind. set in } \mathcal{H} < f(d) \implies |E| \geq \frac{|P|^{2d+3}}{2(2d+3)f(d)^{2d+2}}.$$

By the pigeonhole principle, there exists a $(d+2)$ -sized set $P' \subseteq P$ that is the Radon-base of a set E' of edges of E , where

$$|E'| \geq \frac{|E|}{\binom{|P|}{d+2}} \geq \frac{\frac{|P|^{2d+3}}{2(2d+3)f(d)^{2d+2}}}{\binom{|P|}{d+2}}.$$

The $(d+1)$ -uniform hypergraph consisting of the sets $E'' = \{e' \setminus P' : e' \in E'\}$ has the property that the convex hull of the elements of each set contains $\text{Radon } P'$. It suffices to show that it contains a matching of size $\Omega(|P|)$ —and this follows from known lower-bounds on matchings in uniform hypergraphs (see [1]). For simplicity, we instead present a direct argument, though with worse constants.

Iteratively construct a matching by adding a $(d+1)$ -sized set from E'' to the matching, and deleting all sets from E'' whose intersection with this added set is non-empty. Each set added to the matching can cause the deletion of at most $(d+1) \cdot \binom{|P|}{d}$ sets of E'' , as a vertex of $P \setminus P'$ can belong to at most $\binom{n}{d}$ sets of E'' (each set in E'' has size $(d+1)$). The size of the final matching is the number of iterations, which, by the above discussion, is lower-bounded by

$$\frac{\frac{|P|^{2d+3}}{2(2d+3)f(d)^{2d+2}}}{\binom{|P|}{d+2}} \bigg/ \binom{|P|}{d} (d+1).$$

A calculation then shows that

$$\frac{\frac{|P|^{2d+3}}{2(2d+3)f(d)^{2d+2}}}{\binom{|P|}{d+2}} \bigg/ \binom{|P|}{d} (d+1) \geq \frac{|P|}{h(d)}, \quad \text{where } h(d) = \frac{2(2d+3)(d+1)f(d)^{2d+2}}{(d+2)!d!}.$$

□

3 Proof of Theorem 4

Definition 5. Given positive integers d, p, q with $p \geq q > \lfloor \frac{d}{2} \rfloor$, let $\text{CHS}(d, p, q)$ denote the smallest integer such that the following holds. For any compact convex object $K \subseteq \mathbb{R}^d$ and any set $P \subseteq \mathbb{R}^d \setminus K$ of points, if every subset of P of size p has a q -sized subset whose convex hull is disjoint from K , then P can be separated from K with $\text{CHS}(d, p, q)$ half-spaces (that is, there exists a set \mathcal{H} of $\text{CHS}(d, p, q)$ half-spaces such that $K \subseteq \bigcap_{h \in \mathcal{H}} h$ and $(\bigcap_{h \in \mathcal{H}} h) \cap P = \emptyset$).

Then the key statement is the following.

Lemma 6. *Let P be a set of n points in \mathbb{R}^d and $\epsilon \in [0, \frac{1}{2}]$ a given parameter. Further let $q > \lfloor \frac{d}{2} \rfloor$ be an integer and define $p = q \cdot h(d)$, where $h(d)$ is the function from Theorem 2. Assume $\text{CHS}(d, p, q)$ is finite. Let R be a uniform random sample of P of size*

$$\frac{c_2 \cdot d \cdot \text{CHS}(d, p, q) \cdot \log \text{CHS}(d, p, q)}{\epsilon} \log \frac{1}{\epsilon},$$

where c_2 is a large-enough constant independent of d, ϵ and q . Then with probability at least $\frac{9}{10}$,

1. $R \cup Q$ is a weak ϵ -net for P , where Q is the set of Radon points of all $(d+2)$ -sized subsets of R , and
2. each convex object containing at least $\epsilon|P|$ points of P intersects the convex hull of at least one $(\lfloor \frac{d}{2} \rfloor + 1)$ -sized subset of R .

Proof. The proof follows the method of Mustafa and Ray [7]; however they assumed a more restrictive case, so we present a proof modified appropriately to give a more general bound.

Claim 7. *With probability at least $\frac{9}{10}$, R is an ϵ -net for the set system induced on P by the intersection of $\text{CHS}(d, p, q)$ half-spaces in \mathbb{R}^d .*

Proof. The set system induced by the intersection of k half-spaces in \mathbb{R}^d has VC-dimension $\Theta(dk \log k)$ [3]. Thus by the ϵ -net theorem, a uniform random sample of size

$$\Theta\left(\frac{dk \log k}{\epsilon} \log \frac{1}{\epsilon}\right) = \frac{c_2 \cdot d \cdot \text{CHS}(d, p, q) \cdot \log \text{CHS}(d, p, q)}{\epsilon} \log \frac{1}{\epsilon}$$

is an ϵ -net with probability at least $\frac{9}{10}$ (see [9]), where c_2 is a large-enough constant independent of d, ϵ and q . \square

Assume that R is such an ϵ -net. Let K be any convex object containing at least $\epsilon|P|$ points of P .

Claim 8. *There exists $R_K \subseteq R$ of size p such that the convex hull of every subset of R_K of size q intersects K .*

Proof. If for every subset of R of size p there exists a q -sized subset whose convex hull is *disjoint* from K , then by the definition of $\text{CHS}(d, p, q)$, all points of R can be separated from K by a set \mathcal{H} of $\text{CHS}(d, p, q)$ half-spaces. The common intersection of these half-spaces contains K and hence at least $\epsilon|P|$ points of P and no point of R , a contradiction to Claim 7. \square

By Theorem 2, R_K has a $(d+2)$ -sized subset, say R'_K , such that Radon $R'_K \in Q$ is contained in at least $\frac{|R_K|}{h(d)}$ vertex-disjoint simplices spanned by points of $R_K \setminus R'_K$. Now Radon R'_K must lie inside K : otherwise the half-space separating it from K must contain at least one point from each simplex containing Radon R'_K —namely it must contain at least $\frac{|R_K|}{h(d)} = \frac{p}{h(d)} = q$ points of R_K . But then the convex hull of these q points does not intersect K , a contradiction to Claim 8. Thus $R \cup Q$ is a weak ϵ -net for P , and further, the Radon partition of R'_K of size $\lfloor \frac{d}{2} \rfloor + 1$ must intersect K . This completes the proof. \square

Proof of Theorem 4. It is known that $\text{CHS}(d, p, q)$ is finite for large-enough values of q —this together with Lemma 6 implies the proof. In particular,

1. ([5]) For $p \geq q = d + 1$ we have

$$\text{CHS}(d, p, q) = O\left(p^{d^2} \log^{c' d^3 \log d} p\right),$$

where c' is an absolute constant.

2. ([8]) For any real $\beta > 0$ and $p \geq q = (1 + \beta) \cdot \lfloor \frac{d}{2} \rfloor$ we have

$$\text{CHS}(d, p, q) = O\left(q^2 p^{1+\frac{1}{\beta}} \log p\right).$$

The second bound is stronger, but both of them imply, by setting $q = (d + 1)$, $p = q \cdot h(d)$ and applying Lemma 6, the existence of a function

$$g(d) = O\left(d \cdot \text{CHS}(d, (d + 1) \cdot h(d), (d + 1)) \cdot \log \text{CHS}(d, (d + 1) \cdot h(d), (d + 1))\right).$$

□

Acknowledgements. The first author was supported by the Hungarian National Research, Development and Innovation Office NKFIH Grants K 111827 and K 116769. The second author was supported by the grant ANR grant ADDS (ANR-19-CE48-0005).

References

- [1] N. Alon, P. Frankl, H. Huang, V. Rödl, A. Ruciński, and B. Sudakov. Large matchings in uniform hypergraphs and the conjectures of Erdős and Samuels. *Journal of Combinatorial Theory, Series A*, 119(6):1200–1215, 2012.
- [2] I. Bárány and P. Soberón. Tverberg’s theorem is 50 years old: a survey. *Bull. Amer. Math. Soc.*, 55:459–492, 2018.
- [3] M. Csikós, N. H. Mustafa, and A. Kupavskii. Tight lower bounds on the VC-dimension of geometric set systems. *Journal of Machine Learning Research*, 20(81):1–8, 2019.
- [4] D. de Caen. Extension of a theorem of Moon and Moser on complete subgraphs. *Ars Combin.*, 16:5–10, 1983.
- [5] C. Keller, S. Smorodinsky, and G. Tardos. Improved bounds on the Hadwiger–Debrunner numbers. *Israel Journal of Mathematics*, 225(2):925–945, Apr 2018.
- [6] J. Matoušek and U. Wagner. New constructions of weak epsilon-nets. *Discrete & Computational Geometry*, 32(2):195–206, 2004.
- [7] N. H. Mustafa and S. Ray. Weak ϵ -nets have a basis of size $O(1/\epsilon \log 1/\epsilon)$. *Comp. Geom.: Theory and Appl.*, 40(1):84–91, 2008.

- [8] N. H. Mustafa and S. Ray. On a problem of Danzer. *Combinatorics, Probability and Computing*, page 1–10, 2018.
- [9] N. H. Mustafa and K. Varadarajan. Epsilon-approximations and epsilon-nets. In J. E. Goodman, J. O’Rourke, and C. D. Tóth, editors, *Handbook of Discrete and Computational Geometry*, pages 1241–1268. CRC Press LLC, 2017.
- [10] A. Pór. Universality of vector sequences and universality of Tverberg partitions. *arXiv:1805.07197*, 2018.